

human T

[125825]in a midwestern neural tube defect population. *Teratology*
59: 331–341,
[125826]1999.

[125827]

[125828]

[125829]10910.

[125830]Bellomo, D.; Headrick, J. P.; Silins, G. U.; Paterson, C. A.;
Thomas,

[125831]P. S.; Gartside, M.; Mould, A.; Cahill, M. M.; Tonks, I. D.;
Grimmond,

[125832]S. M.; Townson, S.; Wells, C.; Little, M.; Cummings, M. C.;
Hayward,

[125833]N. K.; Kay, G. F.: Mice lacking the vascular endothelial
growth factor–B

[125834]gene (*Vegfb*) have smaller hearts, dysfunctional coronary
vasculature,

[125835]and impaired recovery from cardiac ischemia. *Circ. Res.*
86: e29–e35,

[125836]2000.

[125837]

[125838]

[125839]10911.

[125840]Gerace, L; Cirenei, N.; Cappelletti, M.; Petraroli, R.; Sebastiani,

[125841]F.; Marziliano, N.: Assignment of the mouse Vegfb gene to mouse chromosome

[125842]19B by in situ hybridization. Cytogenet. Cell Genet. 95: 242–243,

[125843]2001.

[125844]

[125845]

[125846]10912.

[125847]Grimmond, S.; Lagercrantz, J.; Drinkwater, C.; Silins, G.; Townson,

[125848]S.; Pollock, P.; Gotley, D.; Carson, E.; Rakar, S.; Nordenskjold,

[125849]M.; Ward, L.; Hayward, N.; Weber, G.: Cloning and characterization

[125850]of a novel human gene related to vascular endothelial growth factor. Genome

[125851]Res. 6: 124–131, 1996.

[125852]

[125853]

[125854]10913.

[125855]Olofsson, B.; Pajusola, K.; Kaipainen, A.; von Euler, G.;
Joukov,

[125856]V.; Saksela, O.; Orpana, A.; Pettersson, R. F.; Alitalo, K.;
Eriksson,

[125857]U.: Vascular endothelial growth factor B, a novel growth
factor for

[125858]endothelial cells. Proc. Nat. Acad. Sci. 93: 2576–2581,
1996.

[125859]

[125860]

[125861]10914.

[125862]Olofsson, B.; Pajusola, K.; von Euler, G.; Chilov, D.; Alitalo,

[125863]K.; Eriksson, U.: Genomic organization of the mouse and
human genes

[125864]for vascular endothelial growth factor B (VEGF-B) and
characterization

[125865]of a second splice isoform. J. Biol. Chem. 271:
19310–19317, 1996.

[125866]

[125867]

[125868]10915.

[125869]Gunduz, M.; Ouchida, M.; Fukushima, K.; Hanafusa, H.;
Etani, T.;

[125870]Nishioka, S.; Nishizaki, K.; Shimizu, K.: Genomic structure
of the

[125871]human ING1 gene and tumor-specific mutations detected
in head and

[125872]neck squamous cell carcinomas. Cancer Res. 60:
3143–3146, 2000.

[125873]

[125874]

[125875]10916.

[125876]Hasegawa, H.; Kiyokawa, E.; Tanaka, S.; Nagashima, K.;
Gotoh, N.;

[125877]Shibuya, M.; Kurata, T.; Matsuda, M.: DOCK180, a major
CRK-binding

[125878]protein, alters cell morphology upon translocation to the
membrane. Molec.

[125879]Cell Biol. 16: 1770–176, 1996.

[125880]

[125881]

[125882]10917.

[125883]Savill, J.: Phagocytic docking without shocking. Nature
442–443,

[125884]1998.

[125885]

[125886]

[125887]10918.

[125888]Takai, S.; Hasegawa, H.; Kiyokawa, E.; Yamada, K.; Kurata,
T.;

[125889]Matsuda, M.: Chromosomal mapping of the gene encoding DOCK180, a

[125890]major Crk-binding protein, to 10q26.13–q26.3 by fluorescence in situ

[125891]hybridization. Genomics 35: 403–404, 1996.

[125892]

[125893]

[125894]10919.

[125895]Wu, Y.-C.; Horvitz, H. R.: C. elegans phagocytosis and cell-migration

[125896]protein CED-5 is similar to human DOCK180. Nature 392: 501-504,

[125897]1998.

[125898]

[125899]

[125900]10920.

[125901]Alexander, J. M.; Bikkal, H. A.; Zervas, N. T.; Laws, E. R., Jr.;

[125902]Klibanski, A.: Tumor-specific expression and alternate splicing of

[125903]messenger ribonucleic acid encoding activin/transforming growth factor-beta

[125904]receptors in human pituitary adenomas. J. Clin. Endocr. Metab. 81:

[125905]783-790, 1996.

[125906]

[125907]

[125908]10921.

[125909]Su, G. H.; Bansal, R.; Murphy, K. M.; Montgomery, E.; Yeo, C. J.;

[125910]Hruban, R. H.; Kern, S. E.: ACVR1B (ALK4, activin receptor type 1B)

[125911]gene mutations in pancreatic carcinoma. Proc. Nat. Acad. Sci. 98:

[125912]3254–3257, 2001.

[125913]

[125914]

[125915]10922.

[125916]Xu, J.; Matsuzaki, K.; McKeegan, K.; Wang, F.; Kan, M.; McKeegan,

[125917]W. L.: Genomic structure and cloned cDNAs predict that four variants

[125918]in the kinase domain of serine/threonine kinase receptors arise by

[125919]alternative splicing and poly(A) addition. Proc. Nat. Acad. Sci. 91:

[125920]7957–7961, 1994.

[125921]

[125922]

[125923]10923.

[125924]Zhou, Y.; Sun, H.; Danila, D. C.; Johnson, S. R.; Sigai, D. P.;

[125925]Zhang, X.; Klibanski, A.: Truncated activin type I receptor
Alk4

[125926]isoforms are dominant negative receptors inhibiting ac-
tivin signaling. Molec.

[125927]Endocr. 14: 2066–2075, 2000.

[125928]

[125929]

[125930]10924.

[125931]Mann, S. S.; Pettenati, M. J.; von Kap-herr, C.; Hart, T. C.:

[125932]Reassignment of peptidyl prolyl isomerase-like 1 gene
(PPIL1) to human

[125933]chromosome region 6p21.1 by radiation hybrid mapping
and fluorescence

[125934]in situ hybridization. Cytogenet. Cell Genet. 83: 228–229,
1998.

[125935]

[125936]

[125937]10925.

[125938]Ozaki, K.; Fujiwara, T.; Kawai, A.; Shimizu, F.; Takami, S.;
Okuno,

[125939]S.; Takeda, S.; Shimada, Y.; Nagata, M.; Watanabe, T.;
Takaichi, A.;

[125940]Takahashi, E.; Nakamura, Y.; Shin, S.: Cloning, expression
and chromosomal

[125941]mapping of a novel cyclophilin-related gene (PPIL1) from
human fetal

[125942]brain. Cytogenet. Cell Genet. 72: 242–245, 1996.

[125943]

[125944]

[125945]10926.

[125946]Graef, I. A.; Chen, F.; Chen, L.; Kuo, A.; Crabtree, G. R.:
Signals

[125947]transduced by Ca(2+)/calcineurin and NFATc3/c4 pattern
the developing

[125948]vasculature. Cell 105: 863–875, 2001.

[125949]

[125950]

[125951]10927.

[125952]Wang, M. G.; Yi, H.; Guerini, D.; Klee, C. B.; McBride, O. W.:

[125953]Calcineurin A alpha (PPP3CA), calcineurin A beta (PPP3CB)
and calcineurin

[125954]B (PPP3R1) are located on human chromosomes 4,
10q21–q22 and 2p16–p15,

[125955]respectively. Cytogenet. Cell Genet. 72: 236–241, 1996.

[125956]

[125957]

[125958]10928.

[125959]Zeng, H.; Chattarji, S.; Barbarosie, M.; Rondi-Reig, L.;
Philpot,

[125960]B. D.; Miyakawa, T.; Bear, M. F.; Tonegawa, S.: Forebrain–
specific

[125961]calcineurin knockout selectively impairs bidirectional
synaptic plasticity

[125962]and working/episodic–like memory. Cell 107: 617–629,
2001.

[125963]

[125964]

[125965]10929.

[125966]Kolodrubetz, D.; Burgum, A.: Sequence and genetic analysis of

[125967]NHP2: a moderately abundant high mobility group-like nuclear protein

[125968]with an essential function in *Saccharomyces cerevisiae*.

Yeast 7:

[125969]79–90, 1991.

[125970]

[125971]

[125972]10930.

[125973]Saito, H.; Fujiwara, T.; Shin, S.; Okui, K.; Nakamura, Y.: Cloning

[125974]and mapping of a human novel cDNA (NHP2L1) that encodes a protein

[125975]highly homologous to yeast nuclear protein NHP2. Cytogenet. Cell

[125976]Genet. 72: 191–193, 1996.

[125977]

[125978]

[125979]10931.

[125980]Mooseker, M. S.; Cheney, R. E.: Unconventional myosins.
Annu.

[125981]Rev. Cell Dev. Biol. 11: 633–675, 1995.

[125982]

[125983]

[125984]10932.

[125985]Bement, W. M.; Wirth, J. A.; Mooseker, M. S.: Cloning and
mRNA

[125986]expression of human unconventional myosin-IC: a homo-
logue of amoeboid

[125987]myosins-I with a single IQ motif and an SH3 domain. J.
Molec. Biol. 243:

[125988]356–363, 1994.

[125989]

[125990]

[125991]10933.

[125992]Bement, W. M.; Hasson, T.; Wirth, J. A.; Cheney, R. E.;
Mooseker,

[125993]M. S.: Identification and overlapping expression of multi-
ple unconventional

[125994]myosin genes in vertebrate cell types. Proc. Nat. Acad. Sci.

91:

[125995]6549–6553, 1994. Erratum: Proc. Nat. Acad. Sci. 91:

11767, 1994.

[125996]

[125997]

[125998]10934.

[125999]Crozet, F.; Amraoui, A. E.; Blanchard, S.; Lenoir, M.; Ripoll,

[126000]C.; Vago, P.; Hamel, C.; Fizames, C.; Levi-Acobas, F.; Depetris, D.;

[126001]Mattei, M.–G.; Weil, D.; Pujol, R.; Petit, C.: Cloning of the genes

[126002]encoding two murine and human cochlear unconventional type I myosins. Genomics 40:

[126003]332–341, 1997.

[126004]

[126005]

[126006]10935.

[126007]Goppelt, A.; Stelzer, G.; Lottspeich, F.; Meisterernst, M.: A

[126008]mechanism for repression of class II gene transcription

through specific

[126009]binding of NC2 to TBP-promoter complexes via heterodimeric histone

[126010]fold domains. EMBO J. 15: 3105–3116, 1996.

[126011]

[126012]

[126013]10936.

[126014]Inostroza, J. A.; Mermeistein, F. H.; Ha, I.; Lane, W. S.; Reinberg,

[126015]D.: Dr1, a TATA-binding protein-associated phosphoprotein and inhibitor

[126016]of class II gene transcription. Cell 70: 477–489, 1992.

[126017]

[126018]

[126019]10937.

[126020]Kamada, K.; Shu, F.; Chen, H.; Malik, S.; Stelzer, G.; Roeder,

[126021]R. G.; Meisterernst, M.; Burley, S. K.: Crystal structure of negative

[126022]cofactor 2 recognizing the TBP-DNA transcription com-

plex. Cell 106:

[126023]71–81, 2001.

[126024]

[126025]

[126026]10938.

[126027]Mermelstein, F.; Yeung, K.; Cao, J.; Inostroza, J. A.; Erdju-
ment–Bromage,

[126028]H.; Eagelson, K.; Landsman, D.; Levitt, P.; Tempst, P.;
Reinberg,

[126029]D.: Requirement of a corepressor for Dr1–mediated re-
pression of transcription. Genes

[126030]Dev. 10: 1033–1048, 1996.

[126031]

[126032]

[126033]10939.

[126034]Purello, M.; Di Pietro, C.; Rapisarda, A.; Viola, A.; Corsaro,

[126035]C.; Motta, S.; Grzeschik, K.–H.; Sichel, G.: Genomic local-
ization

[126036]of the human gene encoding Dr1, a negative modulator of
transcription

[126037]of class II and class III genes. Cytogenet. Cell Genet. 75:
186–189,

[126038]1996.

[126039]

[126040]

[126041]10940.

[126042]Rozet, J.–M.; Gerber, S.; Perrault, I.; Camuzat, A.; Calvas,
P.;

[126043]Viegas–Pequignot, E.; Molina–Gomes, D.; Le Paslier, D.;
Chumakov,

[126044]I.; Munnich, A.; Kaplan, J.: Structure and physical mapping
of DR1,

[126045]a TATA–binding protein–associated phosphoprotein gene,
to chromosome

[126046]1p22.1 and its exclusion in Stargardt disease (STGD). Ge-
nomics 36:

[126047]554–556, 1996.

[126048]

[126049]

[126050]10941.

[126051]Willy, P. J.; Kobayashi, R.; Kadonaga, J. T.: A basal tran-
scription

[126052]factor that activates or represses transcription. Science
290: 982–984,

[126053]2000.

[126054]

[126055]

[126056]10942.

[126057]Yeung, K.; Kim, S.; Reinberg, D.: Functional dissection of a
human

[126058]Dr1–DRAP1 repressor complex. Molec. Cell. Biol. 17:
36–45, 1997.

[126059]

[126060]

[126061]10943.

[126062]Dong, L. Q.; Du, H.; Porter, S. G.; Kolakowski, L. F., Jr.;
Lee,

[126063]A. V.; Mandarino, J.; Fan, J.; Yee, D.; Liu, F.: Cloning, chro-
mosome

[126064]localization, expression, and characterization of an Src
homology

[126065]2 and pleckstrin homology domain-containing insulin receptor binding

[126066]protein hGrb10-gamma. J. Biol. Chem. 272: 29104-29112, 1997.

[126067]

[126068]

[126069]10944.

[126070]Margolis, B.; Silvennoinen, O.; Comoglio, F.; Roonprapunt, C.;

[126071]Skolnik, E.; Ullrich, A.; Schlessinger, J.: High-efficiency expression/cloning

[126072]of epidermal growth factor-receptor-binding proteins with Src homology

[126073]2 domains. Proc. Nat. Acad. Sci. 89: 8894-8898, 1992.

[126074]

[126075]

[126076]10945.

[126077]Skolnik, E. Y.; Margolis, B.; Mohammadi, M.; Lowenstein, E.; Fischer,

[126078]R.; Drepps, A.; Ullrich, A.; Schlessinger, J.: Cloning of PI3 kinase-

[126079]associated p85 utilizing a novel method for expression/
cloning of

[126080]target proteins for receptor tyrosine kinases. Cell 65:
83–90, 1991.

[126081]

[126082]

[126083]10946.

[126084]Tanaka, S.; Mori, M.; Akiyoshi, T.; Tanaka, Y.; Mafune, K.;
Wands,

[126085]J. R.; Sugimachi, K.: A novel variant of human Grb7 is as-
sociated

[126086]with invasive esophageal carcinoma. J. Clin. Invest. 102:
821–827,

[126087]1998.

[126088]

[126089]

[126090]10947.

[126091]Blagitko, N.; Mergenthaler, S.; Schulz, U.; Wollmann, H. A.;
Craig,

[126092]W.; Eggermann, T.; Ropers, H.–H.; Kalscheuer, V. M.: Hu-
man GRB10

[126093]is imprinted and expressed from the paternal and maternal allele in

[126094]a highly tissue- and isoform-specific fashion. Hum.

Molec. Genet. 9:

[126095]1587-1595, 2000.

[126096]

[126097]

[126098]10948.

[126099]Frantz, J. D.; Giorgetti-Peraldi, S.; Ottinger, E. A.; Shoelson, son,

[126100]S. E.: Human GRB-IR-beta/GRB10: splice variants of an insulin and

[126101]growth factor receptor-binding protein with PH and SH2 domains. J.

[126102]Biol. Chem. 272: 2659-2667, 1997.

[126103]

[126104]

[126105]10949.

[126106]Hannula, K.; Lipsanen-Nyman, M.; Kontiokari, T.; Kere, J.:
A narrow

[126107]segment of maternal uniparental disomy of chromosome
7q31-qter in

[126108]Silver-Russell syndrome delimits a candidate gene region.
Am. J.

[126109]Hum. Genet. 68: 247-253, 2001.

[126110]

[126111]

[126112]10950.

[126113]Jerome, C. A.; Scherer, S. W.; Tsui, L.-C.; Gietz, R. D.;
Triggs-Raine,

[126114]B.: Assignment of growth factor receptor-bound protein
10 (GRB10)

[126115]to human chromosome 7p11.2-p12. Genomics 40:
215-216, 1997.

[126116]

[126117]

[126118]10951.

[126119]Liu, F.; Roth, R. A.: Grb-IR: a SH2-domain-containing pro-
tein

[126120]that binds to the insulin receptor and inhibits its function.
Proc.

[126121]Nat. Acad. Sci. 92: 10287–10291, 1995.

[126122]

[126123]

[126124]10952.

[126125]Ooi, J.; Yajnik, V.; Immanuel, D.; Gordon, M.; Moskow, J. J.;

[126126]Buchberg, A. M.; Margolis, B.: The cloning of Grb10 re-
veals a new

[126127]family of SH2 domain proteins. Oncogene 10: 1621–1630,
1995.

[126128]

[126129]

[126130]10953.

[126131]Daly, R. J.; Sanderson, G. M.; Janes, P. W.; Sutherland, R. L.

[126132]: Cloning and characterization of GRB14, a novel member
of the GRB7

[126133]gene family. J. Biol. Chem. 271: 12502–12510, 1996.

[126134]

[126135]

[126136]10954.

[126137]Hakala, B. E.; White, C.; Recklies, A. D.: Human cartilage
gp-39,

[126138]a major secretory product of articular chondrocytes and
synovial cells,

[126139]is a mammalian member of a chitinase protein family. J.
Biol. Chem. 268:

[126140]25803-25810, 1993.

[126141]

[126142]

[126143]10955.

[126144]Rehli, M.; Krause, S. W.; Andreesen, R.: Molecular charac-
terization

[126145]of the gene for human cartilage gp-39 (CHI3L1), a mem-
ber of the chitinase

[126146]protein family and marker for late stages of macrophage
differentiation. Genomics 43:

[126147]221-225, 1997.

[126148]

[126149]

[126150]10956.

[126151]Dry, K.; Kenwrick, S.; Rosenthal, A.; Platzner, M.: The complete

[126152]sequence of the human locus for NgCAM-related cell adhesion molecule

[126153]reveals a novel alternative exon in chick and man and conserved genomic

[126154]organization for the L1 subfamily. Gene 273: 115–122, 2001.

[126155]

[126156]

[126157]10957.

[126158]Grumet, M.; Mauro, V.; Burgoon, M. P.; Edelman, G. M.; Cunningham,

[126159]B. A.: Structure of a new nervous system glycoprotein, Nr-CAM, and

[126160]its relationship to subgroups of neural cell adhesion molecules. J.

[126161]Cell. Biol. 113: 1399–1412, 1991.

[126162]

[126163]

[126164]10958.

[126165]Kayyem, J. F.; Roman, J. M.; de la Rosa, E. J.; Schwarz, U.;
Dreyer,

[126166]W. J.: Bravo/Nr-CAM is closely related to the cell adhesion
molecules

[126167]L1 and Ng-CAM and has a similar heterodimer structure.
J. Cell. Biol. 118:

[126168]1259–1270, 1992.

[126169]

[126170]

[126171]10959.

[126172]Lane, R. P.; Chen, X.-N.; Yamakawa, K.; Vielmetter, J.; Ko-
renberg,

[126173]J. R.; Dreyer, W. J.: Characterization of a highly conserved
human

[126174]homolog to the chicken neural cell surface protein Bravo/
Nr-CAM that

[126175]maps to chromosome band 7q31. Genomics 35: 456–465,
1996.

[126176]

[126177]

[126178]10960.

[126179]Wang, B.; Williams, H.; Du, J.-S.; Terrett, J.; Kenwrick, S.:

[126180]Alternative splicing of human NrCAM in neural and non-neural tissues. *Molec.*

[126181]*Cell. Neurosci.* 10: 287–295, 1998.

[126182]

[126183]

[126184]10961.

[126185]Damen, J. E.; Liu, L.; Rosten, P.; Humphries, R. K.; Jefferson, son,

[126186]A. B.; Majerus, P. W.; Krystal, G.: The 145-kDa protein induced to

[126187]associate with Shc by multiple cytokines is an inositol tetraphosphate

[126188]and phosphatidylinositol 3,4,5-triphosphate 5-phosphatase. *Proc.*

[126189]*Nat. Acad. Sci.* 93: 1689–1693, 1996.

[126190]

[126191]

[126192]10962.

[126193]Drayer, A. L.; Pesesse, X.; De Smedt, F.; Woscholski, R.;

Parker,

[126194]P.; Erneux, C.: Cloning and expression of a human placenta inositol

[126195]1,3,4,5-tetrakisphosphate and phosphatidylinositol 3,4,5-triphosphate

[126196]5-phosphatase. Biochem. Biophys. Res. Commun. 225: 243-249, 1996.

[126197]

[126198]

[126199]10963.

[126200]Helgason, C. D.; Damen, J. E.; Rosten, P.; Grewal, R.; Sorensen,

[126201]P.; Chappel, S. M.; Borowski, A.; Jirik, F.; Krystal, G.; Humphries,

[126202]R. K.: Targeted disruption of SHIP leads to hemopoietic perturbations,

[126203]lung pathology, and a shortened life span. Genes Dev. 12: 1610-1620,

[126204]1998.

[126205]

[126206]

[126207]10964.

[126208]Huber, M.; Helgason, C. D.; Damen, J. E.; Liu, L.;
Humphries, R.

[126209]K.; Krystal, G.: The src homology 2-containing inositol
phosphatase

[126210](SHIP) is the gatekeeper of mast cell degranulation. Proc.
Nat. Acad.

[126211]Sci. 95: 11330–11335, 1998.

[126212]

[126213]

[126214]10965.

[126215]Kavanaugh, W. M.; Pot, D. A.; Chin, S. M.; Deuter-Rein-
hard, M.;

[126216]Jefferson, A. B.; Norris, F. A.; Masiarz, F. R.; Cousens, L. S.;
Majerus,

[126217]P. W.; Williams, L. T.: Multiple forms of an inositol
polyphosphate

[126218]5-phosphatase form signaling complexes with Shc and
Grb2. Curr.

[126219]Biol. 6: 438–445, 1996.

[126220]

[126221]

[126222]10966.

[126223]Lioubin, M. N.; Algate, P. A.; Tsai, S.; Carlberg, K.; Aebersold,

[126224]R.; Rohrschneider, L. R.: p150(Ship), a signal transduction molecule

[126225]with inositol polyphosphate-5-phosphatase activity.

Genes Dev. 10:

[126226]1084-1095, 1996.

[126227]

[126228]

[126229]10967.

[126230]Liu, Q.; Amgen EST Program; Dumont, D. J.: Molecular cloning

[126231]and chromosomal localization in human and mouse of the SH2-containing

[126232]inositol phosphatase, INPP5D (SHIP). Genomics 39: 109-112, 1997.

[126233]

[126234]

[126235]10968.

[126236]Liu, Q.; Shalaby, F.; Jones, J.; Bouchard, D.; Dumont, D. J.:

[126237]The SH2-containing inositol polyphosphate

5-phosphatase, Ship, is

[126238]expressed during hematopoiesis and spermatogenesis.

Blood 91: 2753-2759,

[126239]1998.

[126240]

[126241]

[126242]10969.

[126243]Takeshita, S.; Namba, N.; Zhao, J. J.; Jiang, Y.; Genant, H.

K.;

[126244]Silva, M. J.; Brodt, M. D.; Helgason, C. D.; Kalesnikoff, J.;

Rauh,

[126245]M. J.; Humphries, R. K.; Krystal, G.; Teitelbaum, S. L.; Ross,

F.

[126246]P.: SHIP-deficient mice are severely osteoporotic due to

increased

[126247]numbers of hyper-resorptive osteoclasts. Nature Med. 8:

943-949,

[126248]2002.

[126249]

[126250]

[126251]10970.

[126252]Wang, J.-W.; Howson, J. M.; Ghansah, T.; Desponts, C.; Ninos,
nos,

[126253]J. M.; May, S. L.; Nguyen, K. H. T.; Toyama-Sorimachi, N.;
Kerr, W.

[126254]G.: Influence of SHIP on the NK repertoire and allogeneic
bone marrow

[126255]transplantation. Science 295: 2094–2097, 2002.

[126256]

[126257]

[126258]10971.

[126259]Ware, M. D.; Rosten, P.; Damen, J. E.; Liu, L.; Humphries,
R.

[126260]K.; Krystal, G.: Cloning and characterization of human
SHIP, the

[126261]145-kD inositol 5-phosphatase that associates with SHC
after cytokine

[126262]stimulation. Blood 88: 2833–2840, 1996.

[126263]

[126264]

[126265]10972.

[126266]Cates, C. A.; Michael, R. L.; Staybrook, K. R.; Harvey, K. A.;
Burke,

[126267]Y. D.; Randall, S. K.; Crowell, P. L.; Crowell, D. N.: Prenyla-
tion

[126268]of oncogenic human PTP(CAAX) protein tyrosine phos-
phatases. Cancer

[126269]Lett. 110: 49–55, 1996.

[126270]

[126271]

[126272]10973.

[126273]Montagna, M.; Serova, O.; Sylla, B. S.; Feunteun, J.; Lenoir,
G.

[126274]M.: A 100–kb physical and transcriptional map around the
EDH17B2

[126275]gene: identification of three novel genes and a pseudo-
gene of a human

[126276]homologue of the rat PRL–1 tyrosine phosphatase. Hum.
Genet. 96:

[126277]532–538, 1995.

[126278]

[126279]

[126280]10974.

[126281]Zeng, Q.; Hong, W.; Tan, Y. H.: Mouse PRL–2 and PRL–3,
two potentially

[126282]prenylated protein tyrosine phosphatases homologous to
PRL–1. Biochem.

[126283]Biophys. Res. Commun. 244: 421–427, 1998.

[126284]

[126285]

[126286]10975.

[126287]Muneer, S.; Ramalingam, V.; Wyatt, R.; Schultz, R. A.;
Minna, J.

[126288]D.; Kamibayashi, C.: Genomic organization and mapping
of the gene

[126289]encoding the PP2A B56–gamma regulatory subunit. Ge-
nomics 79: 344–348,

[126290]2002.

[126291]

[126292]

[126293]10976.

[126294]Deveraux, Q.; Jensen, C.; Rechsteiner, M.: Molecular
cloning and

[126295]expression of a 26 S protease subunit enriched in
dileucine repeats. J.

[126296]Biol. Chem. 270: 23726–23729, 1995.

[126297]

[126298]

[126299]10977.

[126300]Deveraux, Q.; Ustrell, V.; Pickart, C.; Rechsteiner, M.: A 26

[126301]S protease subunit that binds ubiquitin conjugates. J. Biol.
Chem. 269:

[126302]7059–7061, 1994.

[126303]

[126304]

[126305]10978.

[126306]Hu, R.-J.; Lee, M. P.; Johnson, L. A.; Feinberg, A. P.: A
novel

[126307]human homologue of yeast nucleosome assembly protein,
65 kb centromeric

[126308]to the p57(KIP2) gene, is biallelically expressed in fetal
and adult

[126309]tissues. Hum. Molec. Genet. 5: 1743–1748, 1996.

[126310]

[126311]

[126312]10979.

[126313]Abderrahim, H.; Jaramillo–Babb, V. L.; Zhou, Z.; Vollrath,
D.:

[126314]Characterization of the murine TIGR/myocilin gene. Mam–
malian Genome 9:

[126315]673–675, 1998.

[126316]

[126317]

[126318]10980.

[126319]Adam, M. F.; Belmouden, A.; Binisti, P.; Brezin, A. P.; Val–
tot,

[126320]F.; Bechetoille, A.; Dascotte, J.–C.; Copin, B.; Gomez, L.;
Chaventre,

[126321]A.; Bach, J.-F.; Garchon, H.-J.: Recurrent mutations in a single

[126322]exon encoding the evolutionarily conserved olfactomedin-homology domain

[126323]of TIGR in familial open-angle glaucoma. Hum. Molec. Genet. 6: 2091-2097,

[126324]1997.

[126325]

[126326]

[126327]10981.

[126328]Baird, P. N.; Dickinson, J.; Craig, J. E.; Mackey, D. A.: The

[126329]Taa1 restriction enzyme provides a simple means to identify the Q368STOP

[126330]mutation of the myocilin gene in primary open angle glaucoma. Am.

[126331]J. Ophthalm. 131: 510-511, 2001.

[126332]

[126333]

[126334]10982.

[126335]Rodriguez, P.; Munroe, D.; Prawitt, D.; Chu, L. L.; Bric, E.;

Kim,

[126336]J.; Reid, L. H.; Davies, C.; Nakagama, H.; Loebbert, R.;

Winterpacht,

[126337]A.; Petruzzi, M.-J.; Higgins, M. J.; Nowak, N.; Evans, G.;

Shows,

[126338]T.; Weissman, B. E.; Zabel, B.; Housman, D. E.; Pelletier, J.:

Functional

[126339]characterization of human nucleosome assembly protein–

2 (NAP1L4) suggests

[126340]a role as a histone chaperone. Genomics 44: 253–265,

1997.

[126341]

[126342]

[126343]10983.

[126344]Stegmaier, K.; Pendse, S.; Barker, G. F.; Bray–Ward, P.;

Ward,

[126345]D. C.; Montgomery, K. T.; Krauter, K. S.; Reynolds, C.;

Sklar, J.;

[126346]Donnelly, M.; Bohlander, S. K.; Rowley, J. D.; Sallan, S. E.;

Gilliland,

[126347]D. G.; Golub, T. R.: Frequent loss of heterozygosity at the

TEL gene

[126348] locus in acute lymphoblastic leukemia of childhood. Blood
86: 38–44,
[126349] 1995.

[126350]

[126351]

[126352] 10984.

[126353] Arakawa, H.; Nagase, H.; Hayashi, N.; Fujiwara, T.; Ogawa,
M.;

[126354] Shin, S.; Nakamura, Y.: Molecular cloning and expression
of a novel

[126355] human gene that is highly homologous to human
FK506–binding protein

[126356] 12kDa (hFKBP–12) and characterization of two alterna-
tively spliced

[126357] transcripts. Biochem. Biophys. Res. Commun. 200:
836–843, 1994.

[126358]

[126359]

[126360] 10985.

[126361] Xin, H.–B.; Senbonmatsu, T.; Cheng, D.–S.; Wang, Y.–X.
Copello,

[126362]J. A.; Ji, G.-J.; Collier, M. L.; Deng, K.-Y.; Jeyakumar, L. H.;
Magnuson,

[126363]M. A.; Inagami, T.; Kotlikoff, M. I.; Fleischer, S.: Oestrogen
protects

[126364]FKBP12.6 null mice from cardiac hypertrophy. Nature 416:
334–337,

[126365]2002.

[126366]

[126367]

[126368]10986.

[126369]Stein, R.; Mori, N.; Matthews, K.; Lo, L.-C.; Anderson, D. J.:

[126370]The NGF-inducible SCG10 mRNA encodes a novel mem-
brane-bound protein

[126371]present in growth cones and abundant in developing neu-
rons. Neuron 1:

[126372]463–476, 1988.

[126373]

[126374]

[126375]10987.

[126376]Dong, J.-T.; Isaacs, W. B.; Barrett, J. C.; Isaacs, J. T.: Ge-
nomic

[126377]organization of the human KAI1 metastasis-suppressor
gene. Genomics 41:

[126378]25-32, 1997.

[126379]

[126380]

[126381]10988.

[126382]Dong, J.-T.; Lamb, P. W.; Rinker-Schaeffer, C. W.;
Vukanovic, J.;

[126383]Ichikawa, T.; Isaacs, J. T.; Barrett, J. C.: KAI1, a metastasis
suppressor

[126384]gene for prostate cancer on human chromosome 11p11.2.
Science 268:

[126385]884-886, 1995.

[126386]

[126387]

[126388]10989.

[126389]Guo, X.-Z.; Friess, H.; Di Mola, F. F.; Heinicke, J.-M.;

Abou-Shady,

[126390]M.; Graber, H. U.; Baer, H. U.; Zimmermann, A.; Korc, M.;

Buchler,

[126391]M. W.: KAI1, a new metastasis suppressor gene, is reduced

in metastatic

[126392]hepatocellular carcinoma. Hepatology 28: 1481–1488,

1998.

[126393]

[126394]

[126395]10990.

[126396]Mashimo, T.; Watabe, M.; Hirota, S.; Hosobe, S.; Miura, K.;

Tegtmeyer,

[126397]P. J.; Rinker-Shaeffer, C. W.; Watabe, K.: The expression of

the

[126398]KAI1 gene, a tumor metastasis suppressor, is directly acti-

vated by

[126399]p53. Proc. Nat. Acad. Sci. 95: 11307–11311, 1998.

[126400]

[126401]

[126402]10991.

[126403]Miyazaki, T.; Kato, H.; Shitara, Y.; Yoshikawa, M.; Tajima,

K.;

[126404]Masuda, N.; Shouji, H.; Tsukada, K.; Nakajima, T.; Kuwano, H.: Mutation

[126405]and expression of the metastasis suppressor gene KAI1 in esophageal

[126406]squamous cell carcinoma. Cancer 89: 955–962, 2000.

[126407]

[126408]

[126409]10992.

[126410]Gupta, P.; Soyombo, A. A.; Atashband, A.; Wisniewski, K. E.; Shelton,

[126411]J. M.; Richardson, J. A.; Hammer, R. E.; Hofmann, S. L.: Disruption

[126412]of PPT1 or PPT2 causes neuronal ceroid lipofuscinosis in knockout

[126413]mice. Proc. Nat. Acad. Sci. 98: 13566–13571, 2001.

[126414]

[126415]

[126416]10993.

[126417]Ardell, M. D.; Aragon, I.; Oliveira, L.; Porche, G. E.; Burke,

[126418]E.; Pittler, S. J.: The beta subunit of human rod photoreceptor cGMP-gated

[126419]cation channel is generated from a complex transcription unit. FEBS

[126420]Lett. 389: 213–218, 1996.

[126421]

[126422]

[126423]10994.

[126424]Ardell, M. D.; Makhija, A. K.; Oliveira, L.; Miniou, P.; Viegas-Pequignot,

[126425]E.; Pittler, S. J.: cDNA, gene structure, and chromosomal localization

[126426]of human GAR1 (CNCG3L), a homolog of the third subunit of bovine photoreceptor

[126427]cGMP-gated channel. Genomics 28: 32–38, 1995.

[126428]

[126429]

[126430]10995.

[126431]Bareil, C.; Hamel, C. P.; Delague, V.; Arnaud, B.; Demaille, J.;

[126432]Claustres, M.: Segregation of a mutation in CNGB1 encoding the beta-subunit

[126433]of the rod cGMP-gated channel in a family with autosomal recessive

[126434]retinitis pigmentosa. Hum. Genet. 108: 328–334, 2001.

[126435]

[126436]

[126437]10996.

[126438]Bird, A. C.: Retinal photoreceptor dystrophies. LI. Edward Jackson

[126439]Memorial Lecture. Am. J. Ophthal. 119: 543–562, 1995.

[126440]

[126441]

[126442]10997.

[126443]Chen, T.-Y.; Illing, M.; Molday, L. L.; Hsu, Y.-T.; Yau, K.-W.;

[126444]Molday, R. S.: Subunit 2 (or beta) of retinal rod cGMP-gated cation

[126445]channel is a component of the 240-kDa channel-associated protein and

- [126446]mediates Ca(2+)-calmodulin modulation. Proc. Nat. Acad. Sci. 91: 11757-11761, 1994.
- [126448]
- [126449]
- [126450]10998.
- [126451]Chen, T. Y.; Peng, Y. W.; Dhallan, R. S.; Ahamed, B.; Reed, R.; Yau, K. W.: A new subunit of the cyclic nucleotide-gated cation channel in retinal rods. Nature 362: 764-767, 1993.
- [126454]
- [126455]
- [126456]10999.
- [126457]Korschen, H. G.; Beyermann, M.; Muller, F.; Heck, M.; Vantler, M.; Koch, K.-W.; Kellner, R.; Wolfrum, U.; Bode, C.; Hofmann, K. P.; Kaupp, U. B.: Interaction of glutamic-acid-rich proteins with the

[126460]cGMP signalling pathway in rod photoreceptors. Nature
400: 761–766,

[126461]1999.

[126462]

[126463]

[126464]11000.

[126465]Kauppi, P.; Laitinen, T.; Ollikainen, V.; Mannila, H.; Laiti-
nen,

[126466]L. A.; Kere, J.: The IL9R region contribution in asthma is
supported

[126467]by genetic association in an isolated population. Europ. J.
Hum.

[126468]Genet. 8: 788–792, 2000.

[126469]

[126470]

[126471]11001.

[126472]Kermouni, A.; Van Roost, E.; Arden, K. C.; Vermeesch, J.
R.; Weiss,

[126473]S.; Godelaine, D.; Flint, J.; Lurquin, C.; Szikora, J.–P.;
Higgs,

[126474]D. R.; Marynen, P.; Renauld, J.-C.: The IL-9 receptor gene (IL9R):

[126475]genomic structure, chromosomal localization in the pseudoautosomal

[126476]region of the long arm of the sex chromosomes, and identification

[126477]of IL9R pseudogenes at 9qter, 10pter, 16pter, and 18pter.

Genomics 29:

[126478]371-382, 1995.

[126479]

[126480]

[126481]11002.

[126482]Kvaloy, K.; Galvagni, F.; Brown, W. R. A.: The sequence organization

[126483]of the long arm pseudoautosomal region of the human sex chromosomes. Hum.

[126484]Molec. Genet. 3: 771-778, 1994.

[126485]

[126486]

[126487]11003.

[126488]Renauld, J.-C.; Druez, C.; Kermouni, A.; Houssiau, F.; Uyt-
tenhove,

[126489]C.; Van Roost, E.; Van Snick, J.: Expression cloning of the
murine

[126490]and human interleukin 9 receptor cDNAs. Proc. Nat. Acad.
Sci. 89:

[126491]5690–5694, 1992.

[126492]

[126493]

[126494]11004.

[126495]Vermeesch, J. R.; Petit, P.; Kermouni, A.; Renauld, J.-C.;
Van

[126496]Den Berghe, H.; Marynen, P.: The IL-9 receptor gene, lo-
cated in the

[126497]Xq/Yq pseudoautosomal region, has an autosomal origin,
escapes X inactivation

[126498]and is expressed from the Y. Hum. Molec. Genet. 6: 1–8,
1997.

[126499]

[126500]

[126501]11005.

[126502]Blair, H. J.; Ho, M.; Monaco, A. P.; Fisher, S.; Craig, I. W.;

[126503]Boyd, Y.: High-resolution comparative mapping of the
proximal region

[126504]of the mouse X chromosome. Genomics 28: 305–310,
1995.

[126505]

[126506]

[126507]11006.

[126508]Cox, J. P. D.; Yamamoto, K.; Christie, P. T.; Wooding, C.;
Feest,

[126509]T.; Flinter, F. A.; Goodyer, P. R.; Leumann, E.; Neuhaus, T.;
Reid,

[126510]C.; Williams, P. F.; Wrong, O.; Thakker, R. V.: Renal chlo-
ride channel,

[126511]CLCN5, mutations in Dent's disease. J. Bone Min. Res. 14:
1536–1542,

[126512]1999.

[126513]

[126514]

[126515]11007.

[126516]Craig, I.: Personal Communication. Oxford, England
1/31/1995.

[126517]

[126518]

[126519]11008.

[126520]Devuyst, O.; Christie, P. T.; Courtoy, P. J.; Beauwens, R.;
Thakker,

[126521]R. V.: Intra-renal and subcellular distribution of the hu-
man chloride

[126522]channel, CLC-5, reveals a pathophysiological basis for
Dent's disease. Hum.

[126523]Molec. Genet. 8: 247-257, 1999.

[126524]

[126525]

[126526]11009.

[126527]Dutzler, R.; Campbell, E. B.; Cadene, M.; Chait, B. T.;
MacKinnon,

[126528]R.: X-ray structure of a CIC chloride channel at 3.0
angstrom reveals

[126529]the molecular basis of anion selectivity. Nature 415:
287-294, 2002.

[126530]

[126531]

[126532]11010.

[126533]Fisher, S. E.; Black, G. C. M.; Lloyd, S. E.; Hatchwell, E.;
Wrong,

[126534]O.; Thakker, R. V.; Craig, I. W.: Isolation and partial characterization

[126535]of a chloride channel gene which is expressed in kidney
and is a candidate

[126536]for Dent's disease (an X-linked hereditary nephrolithiasis).
Hum.

[126537]Molec. Genet. 3: 2053–2059, 1994.

[126538]

[126539]

[126540]11011.

[126541]Fisher, S. E.; Van Bakel, I.; Lloyd, S. E.; Pearce, S. H. S.;
Thakker,

[126542]R. V.; Craig, I. W.: Cloning and characterization of CLCN5,
the human

[126543]kidney chloride channel gene implicated in Dent disease
(an X-linked

[126544]hereditary nephrolithiasis). Genomics 29: 598–606, 1995.

[126545]

[126546]

[126547]11012.

[126548]Gunther, W.; Luchow, A.; Cluzeaud, F.; Vandewalle, A.;
Jentsch,

[126549]T. J.: CIC–5, the chloride channel mutated in Dent's dis-
ease, colocalizes

[126550]with the proton pump in endocytotically active kidney
cells. Proc.

[126551]Nat. Acad. Sci. 95: 8075–8080, 1998.

[126552]

[126553]

[126554]11013.

[126555]Igarashi, T.; Hayakawa, H.; Shiraga, H.; Kawato, H.; Yan,
K.; Kawaguchi,

[126556]H.; Yamanake, T.; Tsuchida, S.; Akagi, K.: Hypercalciuria
and nephrocalcinosis

[126557]in patients with idiopathic low molecular weight protein-
uria in Japan:

[126558]is the disease identical to Dent's disease in the United Kingdom? Nephron. 69:

[126559]242–247, 1995.

[126560]

[126561]

[126562]11014.

[126563]Lloyd, S. E.; Gunther, W.; Pearce, S. H. S.; Thomson, A.; Bianchi,

[126564]M. L.; Bosio, M.; Craig, I. W.; Fisher, S. E.; Scheinman, S. J.; Wrong,

[126565]O.; Jentsch, T. J.; Thakker, R. V.: Characterisation of renal chloride

[126566]channel, CLCN5, mutations in hypercalciuric nephrolithiasis (kidney

[126567]stones) disorders. Hum. Molec. Genet. 6: 1233–1239, 1997.

[126568]

[126569]

[126570]11015.

[126571]Lloyd, S. E.; Pearce, S. H. S.; Fisher, S. E.; Steinmeyer, K.;

[126572]Schwappach, B.; Schelnman, S. J.; Harding, B.; Bolino, A.;
Devoto,

[126573]M.; Goodyer, P.; Rigden, S. P. A.; Wrong, O.; Jentsch, T. J.;
Craig,

[126574]I. W.; Thakker, R. V.: A common molecular basis for three
inherited

[126575]kidney stone diseases. Nature 370: 445–449, 1996.

[126576]

[126577]

[126578]11016.

[126579]Lloyd, S. E.; Pearce, S. H. S.; Gunther, W.; Kawaguchi, H.;
Igarashi,

[126580]T.; Jentsch, T. J.; Thakker, R. V.: Idiopathic low molecular
weight

[126581]proteinuria associated with hypercalciuric nephrocalci-
nosis in Japanese

[126582]children is due to mutations of the renal chloride channel
(CLCN5). J.

[126583]Clin. Invest. 99: 967–974, 1997.

[126584]

[126585]

[126586]11017.

[126587]Oudet, C.; Martin-Coignard, D.; Pannetier, S.; Praud, E.;
Champion,

[126588]G.; Hanauer, A.: A second family with XLRH displays the
mutation

[126589]S244L in the CLCN5 gene. Hum. Genet. 99: 781–784,
1997.

[126590]

[126591]

[126592]11018.

[126593]Piwon, N.; Gunther, W.; Schwake, M.; Bosl, M. R.; Jentsch,
T.

[126594]J.: CIC-5 Cl(–)-channel disruption impairs endocytosis in a
mouse

[126595]model for Dent's disease. Nature 408: 369–373, 2000.

[126596]

[126597]

[126598]11019.

[126599]Schurman, S. J.; Norden, A. G. W.; Scheinman, S. J.: X-
linked

[126600]recessive nephrolithiasis: presentation and diagnosis in children. J.

[126601]Pediat. 132: 859–862, 1998.

[126602]

[126603]

[126604]11020.

[126605]Cabeza–Arvelaiz, Y.; Shih, L.–C. N.; Hardman, N.; Asselbergs, F.;

[126606]Bilbe, G.; Schmitz, A.; White, B.; Siciliano, M. J.; Lachman, L. B.

[126607]: Cloning and genetic organization of the human kinesin light-chain

[126608](KLC) gene. DNA Cell Biol. 12: 881–892, 1993.

[126609]

[126610]

[126611]11021.

[126612]Pearce, S. H. S.; Cheetham, T.; Imrie, H.; Vaidya, B.; Barnes,

[126613]N. D.; Bilous, R. W.; Carr, D.; Meeran, K.; Shaw, N. J.; Smith, C.

[126614]S.; Toft, A. D.; Williams, G.; Kendall-Taylor, P.: A common and recurrent

[126615]13-bp deletion in the autoimmune regulator gene in British kindreds

[126616]with autoimmune polyendocrinopathy type 1. Am. J. Hum. Genet. 63:

[126617]1675-1684, 1998.

[126618]

[126619]

[126620]11022.

[126621]Mitnick, M.; Reichlin, S.: Enzymatic synthesis of thyrotropin-releasing

[126622]hormone (TRH) by hypothalamic 'TRH synthetase.'. Endocrinology 91:

[126623]1145-1153, 1972.

[126624]

[126625]

[126626]11023.

[126627]Battini, R.; Ferrari, S.; Kaczmarek, L.; Calabretta, B.; Chen,

[126628]S.-T.; Baserga, R.: Molecular cloning of a cDNA for a hu-

man ADP/ATP

[126629]carrier which is growth-regulated. J. Biol. Chem. 262:
4355-4359,

[126630]1987.

[126631]

[126632]

[126633]11024.

[126634]Chen, S.-T.; Chang, C.-D.; Huebner, K.; Ku, D.-H.; McFar-
land, M.;

[126635]DeRiel, J. K.; Baserga, R.; Wurzel, J.: A human ADP/ATP
translocase

[126636]gene has seven pseudogenes and localizes to chromo-
some X. Somat.

[126637]Cell Molec. Genet. 16: 143-149, 1990.

[126638]

[126639]

[126640]11025.

[126641]Ku, D.-H.; Kagan, J.; Chen, S.-T.; Chang, C.-D.; Baserga,
R.; Wurzel,

[126642]J.: The human fibroblast adenine nucleotide translocator

gene: molecular

[126643]cloning and sequence. J. Biol. Chem. 265: 16060–16063,
1990.

[126644]

[126645]

[126646]11026.

[126647]Schiebel, K.; Mertz, A.; Winkelmann, M.; Nagaraja, R.; Rap-
pold,

[126648]G.: Localization of the adenine nucleotide translocase
gene ANT2

[126649]to chromosome Xq24–q25 with tight linkage to DXS425.
Genomics 24:

[126650]605–606, 1994.

[126651]

[126652]

[126653]11027.

[126654]Lu, Q. R.; Sun, T.; Zhu, Z.; Ma, N.; Garcia, M.; Stiles, C. D.;

[126655]Rowitch, D. H.: Common developmental requirement for
Olig function

[126656]indicates a motor neuron/oligodendrocyte connection.

Cell 109: 75–86,
[126657]2002.

[126658]

[126659]

[126660]11028.

[126661]Lu, Q. R.; Yuk, D.; Alberta, J. A.; Zhu, Z.; Pawlitzky, I.;
Chan,

[126662]J.; McMahon, A. P.; Stiles, C. D.; Rowitch, D. H.: Sonic
hedgehog-regulated

[126663]oligodendrocyte lineage genes encoding bHLH proteins in
the mammalian

[126664]central nervous system. Neuron 25: 317–329, 2000.

[126665]

[126666]

[126667]11029.

[126668]Raff, M. C.; Miller, R. H.; Noble, M.: A glial progenitor cell

[126669]that develops in vitro into an astrocyte or an oligodendro-
cyte depending

[126670]on culture medium. Nature 303: 390–396, 1983.

[126671]

[126672]

[126673]11030.

[126674]Zhou, Q.; Anderson, D. J.: The bHLH transcription factors
OLIG2

[126675]and OLIG1 couple neuronal and glial subtype specifica-
tion. Cell 109:

[126676]61–73, 2002.

[126677]

[126678]

[126679]11031.

[126680]Zhou, Q.; Wang, S.; Anderson, D. J.: Identification of a
novel

[126681]family of oligodendrocyte lineage-specific basic helix-
loop-helix

[126682]transcription factors. Neuron 25: 331–343, 2000.

[126683]

[126684]

[126685]11032.

[126686]Guimaraes, M. J.; Peterson, D.; Vicari, A.; Cocks, B. G.;
Copeland,

[126687]N. G.; Gilbert, D. J.; Jenkins, N. A.; Ferrick, D. A.; Kastelein,

[126688]R. A.; Bazan, J. F.; Zlotnik, A.: Identification of a novel selD

[126689]homolog from eukaryotes, bacteria, and archaea: is there
an autoregulatory

[126690]mechanism in selenocysteine metabolism? Proc. Nat.
Acad. Sci. 93:

[126691]15086–15091, 1996.

[126692]

[126693]

[126694]11033.

[126695]Bates, E. E. M.; Dieu, M.–C.; Ravel, O.; Zurawski, S. M.; Pa-
tel,

[126696]S.; Bridon, J.–M.; Ait–Yahia, S.; Vega, F., Jr.; Banchereau, J.;
Lebecque,

[126697]S.: CD40L activation of dendritic cells down–regulates
DORA, a novel

[126698]member of the immunoglobulin superfamily. Molec. Im-
mun. 35: 513–524,

[126699]1998.

[126700]

[126701]

[126702]11034.

[126703]Bates, E. E. M.; Kissenpfennig, A.; Peronne, C.; Mattei, M.-G.;

[126704]Fossiez, F.; Malissen, B.; Lebecque, S.: The mouse and human IGSF6

[126705](DORA) genes map to the inflammatory bowel disease 1 locus and are

[126706]embedded in an intron of a gene of unknown function.

Immunogenetics 52:

[126707]112–120, 2000.

[126708]

[126709]

[126710]11035.

[126711]Boldin, M. P.; Mett, I. L.; Wallach, D.: A protein related to

[126712]a proteasomal subunit binds to the intracellular domain of the p55

[126713]TNF receptor upstream to its 'death domain.' FEBS Lett. 367: 39–44,

[126714]1995.

[126715]

[126716]

[126717]11036.

[126718]Dunbar, J. D.; Song, H. Y.; Guo, D.; Wu, L.-W.; Donner, D. B.:

[126719]Two-hybrid cloning of a gene encoding TNF receptor-associated protein

[126720]2, a protein that interacts with the intracellular domain of the type

[126721]1 TNF receptor: identity with subunit 2 of the 26S protease. J. Immun. 158:

[126722]4252-4259, 1997.

[126723]

[126724]

[126725]11037.

[126726]Tsurumi, C.; Shimizu, Y.; Saeki, M.; Kato, S.; Demartino, G. N.;

[126727]Slaughter, C. A.; Fujimuro, M.; Yokosawa, H.; Yamasaki, M.; Hendil,

[126728]K. B.; Toh-E, A.; Tanahashi, N.; Tanaka, K.: cDNA cloning
and functional

[126729]analysis of the p97 subunit of the 26S proteasome, a
polypeptide identical

[126730]to the type-1 tumor-necrosis-factor-receptor-associated
protein-2/55.11. Europ.

[126731]J. Biochem. 239: 912-921, 1996.

[126732]

[126733]

[126734]11038.

[126735]Koesters, R.; Adams, V.; Betts, D.; Moos, R.; Schmid, M.;
Siermann,

[126736]A.; Hassam, S.; Weitz, S.; Lichter, P.; Heitz, P. U.; von
Knebel Doeberitz,

[126737]M.; Briner, J.: Human eukaryotic initiation factor EIF2C1
gene: cDNA

[126738]sequence, genomic organization, localization to chromo-
somal bands

[126739]1q34-p35, and expression. Genomics 61: 210-218, 1999.

[126740]

[126741]

[126742]11039.

[126743]Martinez, J.; Patkaniowska, A.; Urlaub, H.; Luhrmann, R.;

Tuschi,

[126744]T.: Single-stranded antisense siRNAs guide target RNA
cleavage in

[126745]RNAi. Cell 110: 563–574, 2002.

[126746]

[126747]

[126748]11040.

[126749]Suetsugu, S.; Miki, H.; Takenawa, T.: Identification of two
human

[126750]WAVE/SCAR homologues as general actin regulatory
molecules which associate

[126751]with the Arp2/3 complex. Biochem. Biophys. Res. Com-
mun. 260: 296–302,

[126752]1999.

[126753]

[126754]

[126755]11041.

[126756]Myers, L. C.; Gustafsson, c. M.; Bushnell, D. A.; Lui, M.;

Erdjument-Bromage,

[126757]H.; Tempst, P.; Kornberg, R. D.: The Med proteins of yeast
and their

[126758]function through the RNA polymerase II carboxy-terminal
domain. Genes

[126759]Dev. 12: 45–54, 1998.

[126760]

[126761]

[126762]11042.

[126763]Hanaoka, E.; Ozaki, T.; Ohira, M.; Nakamura, Y.; Suzuki,
M.; Takahashi,

[126764]E.; Moriya, H.; Nakagawara, A.; Sakiyama, S.: Molecular
cloning and

[126765]expression analysis of the human DA41 gene and its
mapping to chromosome

[126766]9q21.2–q21.3. J. Hum. Genet. 45: 188–191, 2000.

[126767]

[126768]

[126769]11043.

[126770]Ozaki, T.; Hishiki, T.; Toyama, Y.; Yuasa, S.; Nakagawara,

A.;

[126771]Sakiyama, S.: Identification of a new cellular protein that
can interact

[126772]specifically with DAN. DNA Cell Biol. 16: 985–991, 1997.

[126773]

[126774]

[126775]11044.

[126776]Venkateswarlu, K.; Gunn–Moore, F.; Oatey, P. B.; Tavaré, J.
M.;

[126777]Cullen, P. J.: Nerve growth factor– and epidermal growth
factor–stimulated

[126778]translocation of the ADP–ribosylation factor–exchange
factor GRP1

[126779]to the plasma membrane of PC12 cells requires activation
of phosphatidylinositol

[126780]3–kinase and the GRP1 pleckstrin homology domain.
Biochem. J. 335:

[126781]139–146, 1998.

[126782]

[126783]

[126784]11045.

[126785]Dammann, R.; Li, C.; Yoon, J.-H.; Chin, P. L.; Bates, S.;

Pfeifer,

[126786]G. P.: Epigenetic inactivation of a RAS association domain family

[126787]protein from the lung tumour suppressor locus 3p21.3.

Nature Genet. 25:

[126788]315–319, 2000.

[126789]

[126790]

[126791]11046.

[126792]Dreijerink, K.; Braga, E.; Kuzmin, I.; Geil, L.; Duh, F.-M.;

Angeloni,

[126793]D.; Zbar, B.; Lerman, M. I.; Stanbridge, E. J.; Minna, J. D.;

Protopopov,

[126794]A.; Li, J.; Kashuba, V.; Klein, G.; Zabarovsky, E. R.: The

candidate

[126795]tumor suppressor gene, RASSF1A, from human chromo-

some 3p21.3 is involved

[126796]in kidney tumorigenesis. Proc. Nat. Acad. Sci. 98:

7504–7509, 2001.

[126797]

[126798]

[126799]11047.

[126800]Harada, K.; Toyooka, S.; Maitra, A.; Maruyama, R.; Toyooka, K.

[126801]O.; Timmons, C. F.; Tomlinson, G. E.; Mastrangelo, D.; Hay, R. J.;

[126802]Minna, J. D.; Gazdar, A. F.: Aberrant promoter methylation and silencing

[126803]of the RASSF1A gene in pediatric tumors and cell lines.

Oncogene 21:

[126804]4345–4349, 2002.

[126805]

[126806]

[126807]11048.

[126808]Sekido, Y.; Ahmadian, M.; Wistuba, I. I.; Latif, F.; Bader, S.;

[126809]Wei, M.–H.; Duh, F.–M.; Gazdar, A. F.; Lerman, M. I.;

Minna, J. D.

[126810]: Cloning of a breast cancer homozygous deletion junction narrows

[126811]the region of search for a 3p21.3 tumor suppressor gene.

Oncogene 16:

[126812]3151–3157, 1998.

[126813]

[126814]

[126815]11049.

[126816]Dann, C. E.; Hsieh, J.–C.; Rattner, A.; Sharma, D.; Nathans, J.;

[126817]Leahy, D. J.: Insights into Wnt binding and signalling from the structures

[126818]of two Frizzled cysteine–rich domains. Nature 412: 86–90, 2001.

[126819]

[126820]

[126821]11050.

[126822]Hoang, B.; Moos, M., Jr.; Vukicevic, S.; Luyten, F. P.: Primary

[126823]structure and tissue distribution of FRZB, a novel protein related

[126824]to Drosophila frizzled, suggest a role in skeletal morphogenesis. J.

[126825]Biol. Chem. 271: 26131–26137, 1996.

[126826]

[126827]

[126828]11051.

[126829]Leyns, L.; Bouwmeester, T.; Kim, S.-H.; Piccolo, S.; De Robertis,

[126830]E. M.: Frzb-1 is a secreted antagonist of Wnt signaling expressed

[126831]in the Spemann organizer. Cell 88: 747-756, 1997.

[126832]

[126833]

[126834]11052.

[126835]Schumann, H.; Holtz, J.; Zerkowski, H.-R.; Hatzfeld, M.: Expression

[126836]of secreted frizzled related proteins 3 and 4 in human ventricular

[126837]myocardium correlates with apoptosis related gene expression. Cardiovasc.

[126838]Res. 45: 720-728, 2000.

[126839]

[126840]

[126841]11053.

[126842]Bouchon, A.; Dietrich, J.; Colonna, M.: Cutting edge: inflammatory

[126843]responses can be triggered by TREM-1, a novel receptor expressed on

[126844]neutrophils and monocytes. J. Immun. 164: 4991-4995, 2000.

[126845]

[126846]

[126847]11054.

[126848]Bouchon, A.; Facchetti, F.; Weigand, M. A.; Colonna, M.: TREM-1

[126849]amplifies inflammation and is a crucial mediator of septic shock. Nature 410:

[126850]1103-1107, 2001.

[126851]

[126852]

[126853]11055.

[126854]Benghezal, M.; Benachour, A.; Rusconi, S.; Aebi, M.; Conzelmann,

[126855]A.: Yeast Gpi8p is essential for GPI anchor attachment onto proteins. EMBO

[126856]J. 15: 6575–6583, 1996.

[126857]

[126858]

[126859]11056.

[126860]Yu, J.; Nagarajan, S.; Knez, J. J.; Udenfriend, S.; Chen, R.; Medof,

[126861]M. E.: The affected gene underlying the class K glycosylphosphatidylinositol

[126862](GPI) surface protein defect codes for the GPI transamidase. Proc.

[126863]Nat. Acad. Sci. 94: 12580–12585, 1997.

[126864]

[126865]

[126866]11057.

[126867]Yu, Y.; Xu, F.; Peng, H.; Fang, X.; Zhao, S.; Li, Y.; Cuevas, B.;

[126868]Kuo, W.–L.; Gray, J. W.; Siciliano, M.; Mills, G. B.; Bast, R. C.,

[126869]Jr.: NOEY2 (ARHI), an imprinted putative tumor suppressor gene in

[126870]ovarian and breast carcinomas. Proc. Nat. Acad. Sci. 96: 214–219,

[126871]1999.

[126872]

[126873]

[126874]11058.

[126875]Walowsky, C.; Fitzhugh, D. J.; Castano, I. B.; Ju, J. Y.; Levin,

[126876]N. A.; Christman, M. F.: The topoisomerase–related function gene

[126877]TRF4 affects cellular sensitivity to the antitumor agent camptothecin. J.

[126878]Biol. Chem. 274: 7302–7308, 1999.

[126879]

[126880]

[126881]11059.

[126882]Wang, Z.; Castano, I. B.; De Las Penas, A.; Adams, C.; Christman,

[126883]M. F.: Pol kappa: a DNA polymerase required for sister

chromatid

[126884]cohesion. Science 289: 774–779, 2000.

[126885]

[126886]

[126887]11060.

[126888]Clark, J.; Lu, Y.-J.; Sidhar, S. K.; Parker, C.; Gill, S.; Smedley,

[126889]D.; Hamoudi, R.; Linehan, W. M.; Shipley, J.; Cooper, C. S.:
Fusion

[126890]of splicing factor genes PSF and NonO (p54-nrb) to the
TFE3 gene in

[126891]papillary renal cell carcinoma. Oncogene 15: 2233–2239,
1997.

[126892]

[126893]

[126894]11061.

[126895]Patton, J. G.; Porro, E. B.; Galceran, J.; Tempst, P.; Nadal-Ginard,

[126896]B.: Cloning and characterization of PSF, a novel pre-mRNA
splicing

[126897]factor. Genes Dev. 7: 393–406, 1993.

[126898]

[126899]

[126900]11062.

[126901]Piquemal, D.; Joulia, D.; Balaguer, P.; Basset, A.; Marti, J.;

[126902]Commes, T.: Differential expression of the RTP/

Drg1/Ndr1 gene product

[126903]in proliferating and growth arrested cells. Biochim. Bio-
phys. Acta 1450:

[126904]364-373, 1999.

[126905]

[126906]

[126907]11063.

[126908]LeClerc, S.; Palaniswami, R.; Xie, B.; Govdan, M. V.: Molec-
ular

[126909]cloning and characterization of a factor that binds the hu-
man glucocorticoid

[126910]receptor gene and represses its expression. J. Biol. Chem.
266:

[126911]17333-17340, 1991.

[126912]

[126913]

[126914]11064.

[126915]Tikoo, A.; Czekay, S.; Viars, C.; White, S.; Heath, J. K.; Arden,

[126916]K.; Maruta, H.: p190-A, a human tumor suppressor gene, maps to the

[126917]chromosomal region 19q13.3 that is reportedly deleted in some gliomas. Gene 257:

[126918]23-31, 2000.

[126919]

[126920]

[126921]11065.

[126922]Boccaccio, I.; Glatt-Deeley, H.; Watrin, F.; Roeckel, N.; Lande,

[126923]M.; Muscatelli, F.: The human MAGEL2 gene and its mouse homologue

[126924]are paternally expressed and mapped to the Prader-Willi region. Hum.

[126925]Molec. Genet. 8: 2497-2505, 1999.

[126926]

[126927]

[126928]11066.

[126929]Lee, S.; Kozlov, S.; Hernandez, L.; Chamberlain, S. J.; Bran-
nan,

[126930]C. I.; Stewart, C. L.; Wevrick, R.: Expression and imprinting
of

[126931]MAGEL2 suggest a role in Prader-Willi syndrome and the
homologous

[126932]murine imprinting phenotype. Hum. Molec. Genet. 9:
1813-1819, 2000.

[126933]

[126934]

[126935]11067.

[126936]Wang, A. H.; Bertos, N. R.; Vezmar, M.; Pelletier, N.;
Crosato,

[126937]M.; Heng, H. H.; Th'ng, J.; Han, J.; Yang, X.-J.: HDAC4, a
human

[126938]histone deacetylase related to yeast HDA1, is a transcrip-
tional corepressor. Molec.

[126939]Cell. Biol. 19: 7816-7827, 1999.

[126940]

[126941]

[126942]11068.

[126943]Nishimura, Y.; Hayashi, M.; Inada, H.; Tanaka, T.: Molecular cloning

[126944]and characterization of mammalian homologues of vesicle-associated

[126945]membrane protein-associated (VAMP-associated) proteins. Biochem.

[126946]Biophys. Res. Commun. 254: 21–26, 1999.

[126947]

[126948]

[126949]11069.

[126950]Weir, M. L.; Klip, A.; Trimble, W. S.: Identification of a human

[126951]homologue of the vesicle-associated membrane protein (VAMP)-associated

[126952]protein of 33 kDa (VAP-33): a broadly expressed protein that binds

[126953]to VAMP. Biochem. J. 333: 247–251, 1998.

[126954]

[126955]

[126956]11070.

[126957]Perez Jurado, L. A.; Wang, Y.-K.; Francke, U.; Cruces, J.:

TBL2,

[126958]a novel transducin family member in the WBS deletion:

characterization

[126959]of the complete sequence, genomic structure, transcrip-

tional variants

[126960]and the mouse ortholog. Cytogenet. Cell Genet. 86:

277-284, 1999.

[126961]

[126962]

[126963]11071.

[126964]Bates, E. E. M.; Ravel, O.; Dieu, M.-C.; Ho, S.; Guret, C.;

Bridon,

[126965]J.-M.; Ait-Yahia, S.; Briere, F.; Caux, C.; Banchereau, J.;

Lebecque,

[126966]S.: Identification and analysis of a novel member of the

ubiquitin

[126967]family expressed in dendritic cells and mature B cells. Eu-

rop. J.

[126968]Immun. 27: 2471–2477, 1997.

[126969]

[126970]

[126971]11072.

[126972]Akhmanova, A.; Hoogenraad, C. C.; Drabek, K.; Stepanova,
T.; Dortland,

[126973]B.; Verkerk, T.; Vermeulen, W.; Burgering, B. M.; De
Zeeuw, C. I.;

[126974]Grosveld, F.; Galjart, N.: CLASPs are CLIP–115 and –170
associating

[126975]proteins involved in the regional regulation of microtubule
dynamics

[126976]in motile fibroblasts. Cell 104: 923–935, 2001.

[126977]

[126978]

[126979]11073.

[126980]Cattanach, B. M.; Barr, J. A.; Beechey, C. V.; Martin, J.;
Noebels,

[126981]J.; Jones, J.: A candidate model for Angelman syndrome in
the mouse. Mammalian

[126982]Genome 8: 472–478, 1997.

[126983]

[126984]

[126985]11074.

[126986]Dhar, M.; Webb, L. S.; Smith, L.; Hauser, L.; Johnson, D.;
West,

[126987]D. B.: A novel ATPase on mouse chromosome 7 is a candi-
date gene for

[126988]increased body fat. Physiol. Genomics 4: 93–100, 2000.

[126989]

[126990]

[126991]11075.

[126992]Halleck, M. S.; Lawler, J. F., Jr.; Blackshaw, S.; Gao, L.; Na-
garajan,

[126993]P.; Hacker, C.; Pyle, S.; Newman, J. T.; Nakanishi, Y.; Ando,
H.;

[126994]Weinstock, D.; Williamson, P.; Schlegel, R. A.: Differential
expression

[126995]of putative transbilayer amphipath transporters. Physiol.
Genomics 1:

[126996]139–150, 1999.

[126997]

[126998]

[126999]11076.

[127000]Fitzgibbon, J.; Neat, M. J.; Foot, N.; Hill, A. S.; Lister, T.

[127001]A.; Gupta, R. K.: Assignment of brain acid-soluble protein
1 (BASP1)

[127002]to human chromosome 5p15.1–p14, differential expres-
sion in human cancer

[127003]cell lines as a result of alterations in gene dosage. Cyto-
genet.

[127004]Cell Genet. 89: 147–149, 2000.

[127005]

[127006]

[127007]11077.

[127008]Mosevitsky, M. I.; Capony, J. P.; Skladchikova, G. Y.; Novit-
skaya,

[127009]V. A.; Plekhanov, A. Y.; Zakharov, V. V.: The BASP1 family
of myristoylated

[127010]proteins abundant in axonal termini: primary structure

analysis and

[127011]physico-chemical properties. Biochimie 79: 373-384,
1997.

[127012]

[127013]

[127014]11078.

[127015]Park, S.; Kim, Y.-I.; Kim, B.; Seong, C.; Oh, Y.; Baek, K.;
Yoon,

[127016]J.: Characterization of bovine and human cDNAs encoding
NAP-22 (22

[127017]kDa neuronal tissue-enriched acidic protein) homologs.
Molec. Cells 8:

[127018]471-477, 1998.

[127019]

[127020]

[127021]11079.

[127022]Shichijo, S.; Nakao, M.; Imai, Y.; Takasu, H.; Kawamoto,
M.; Niiya,

[127023]F.; Yang, D.; Toh, Y.; Yamana, H.; Itoh, K.: A gene encod-
ing antigenic

[127024]peptides of human squamous cell carcinoma recognized

by cytotoxic

[127025]T lymphocytes. J. Exp. Med. 187: 277–288, 1998.

[127026]

[127027]

[127028]11080.

[127029]Valenta, R.; Natter, S.; Seiberler, S.; Wichlas, S.; Maurer, D.;

[127030]Hess, M.; Pavelka, M.; Grote, M.; Ferreira, F.; Szepefalusi,
Z.; Valent,

[127031]P.; Stingl, G.: Molecular characterization of an autoaller-
gen, Hom

[127032]s 1, identified by serum IgE from atopic dermatitis pa-
tients. J.

[127033]Invest. Derm. 111: 1178–1183, 1998.

[127034]

[127035]

[127036]11081.

[127037]Loftus, B. J.; Kim, U.-J.; Sneddon, V. P.; Kalush, F.; Bran-
don,

[127038]R.; Fuhrmann, J.; Mason, T.; Crosby, M. L.; Barnstead, M.;
Cronin,

[127039]L.; Mays, A. D.; Cao, Y.; Xu, R. X.; Kang, H.-L.; Mitchell, S.;
Eichler,
[127040]E. E.; Harris, P. C.; Venter, J. C.; Adams, M. D.: Genome
duplications
[127041]and other features in 12 Mb of DNA sequence from hu-
man chromosome
[127042]16p and 16q. Genomics 60: 295–308, 1999.
[127043]
[127044]
[127045]11082.
[127046]Brauner–Osborne, H.; Krogsgaard–Larsen, P.: Sequence
and expression
[127047]pattern of a novel human orphan G–protein–coupled re-
ceptor, GPRC5B,
[127048]a family C receptor with a short amino–terminal domain.
Genomics 65:
[127049]121–128, 2000.
[127050]
[127051]
[127052]11083.

[127053]Robbins, M. J.; Michalovich, D.; Hill, J.; Calver, A. R.; Medhurst,

[127054]A. D.; Gloger, I.; Sims, M.; Middlemiss, D. N.; Pangalos, M. N.:

[127055]Molecular cloning and characterization of two novel retinoic acid-inducible

[127056]orphan G-protein-coupled receptors (GPRC5B and GPRC5C). Genomics 67:

[127057]8–18, 2000.

[127058]

[127059]

[127060]11084.

[127061]Fan, W.; Cai, W.; Parimoo, S.; Lennon, G. G.; Schwarz, D. C.; Weissman,

[127062]S. M.: Identification of seven new human MHC class I region genes

[127063]around the HLA-F locus. Immunogenetics 44: 97–103, 1996.

[127064]

[127065]

[127066]11085.

[127067]Autieri, M. V.; Feuerstein, G. Z.; Yue, T.-L.; Ohlstein, E. H.;

[127068]Douglas, S. A.: Use of differential display to identify differentially

[127069]expressed mRNAs induced by rat carotid artery balloon angioplasty. Lab.

[127070]Invest. 72: 656–661, 1995.

[127071]

[127072]

[127073]11086.

[127074]Autieri, M. V.; Haines, D. S.; Romanic, A. M.; Ohlstein, E. H.

[127075]: Expression of 14–3–3–gamma in injured arteries and growth factor–

[127076]and cytokine–stimulated human vascular smooth muscle cells. Cell

[127077]Growth Diff. 7: 1453–1460, 1996.

[127078]

[127079]

[127080]11087.

[127081]Horie, M.; Suzuki, M.; Takahashi, E.; Tanigami, A.:

Cloning, expression,

[127082]and chromosomal mapping of the human 14-3-3gamma gene (YWHAG) to 7q11.23. Genomics 60:

[127083]241-243, 1999.

[127084]

[127085]

[127086]11088.

[127087]Morrison, D.: 14-3-3: modulators of signaling proteins?
Science 266:

[127088]56-57, 1994.

[127089]

[127090]

[127091]11089.

[127092]Brazas, R.; Ganem, D.: A cellular homolog of hepatitis
delta antigen:

[127093]implications for viral replication and evolution. Science
274: 90-94,

[127094]1996.

[127095]

[127096]

[127097]11090.

[127098]Long, M.; de Souza, S. J.; Gilbert, W.: Delta-interacting
protein

[127099]A and the origin of hepatitis delta antigen. Science 276:
824–825,

[127100]1997.

[127101]

[127102]

[127103]11091.

[127104]Robertson, H. D.: How did replicating and coding RNAs
first get

[127105]together? Science 274: 66–67, 1996.

[127106]

[127107]

[127108]11092.

[127109]Asada, H.; Kawamura, Y.; Maruyama, K.; Kume, H.; Ding,
R.–G.; Kanbara,

[127110]N.; Kuzume, H.; Sanbo, M.; Yagi, T.; Obata, K.: Cleft palate

and

[127111]decreased brain gamma-aminobutyric acid in mice lacking
the 67-kDa

[127112]isoform of glutamic acid decarboxylase. Proc. Nat. Acad.
Sci. 94:

[127113]6496–6499, 1997.

[127114]

[127115]

[127116]11093.

[127117]Brilliant, M. H.; Szabo, G.; Katarova, Z.; Kozak, C. A.;
Glaser,

[127118]T. M.; Greenspan, R. J.; Housman, D. E.: Sequences ho-
mologous to

[127119]glutamic acid decarboxylase cDNA are present on mouse
chromosomes

[127120]2 and 10. Genomics 6: 115–122, 1990.

[127121]

[127122]

[127123]11094.

[127124]Bu, D.-F.; Tobin, A. J.: The exon-intron organization of

the genes

[127125](GAD1 and GAD2) encoding two human glutamate decarboxylases (GAD-67

[127126]and GAD-65) suggests that they derive from a common ancestral GAD. Genomics 21:

[127127]222–228, 1994.

[127128]

[127129]

[127130]11095.

[127131]Condie, B. G.; Bain, G.; Gottlieb, D. I.; Capecchi, M. R.:
Cleft

[127132]palate in mice with a targeted mutation in the gamma-aminobutyric

[127133]acid-producing enzyme glutamic acid decarboxylase 67.
Proc. Nat.

[127134]Acad. Sci. 94: 11451–11455, 1997.

[127135]

[127136]

[127137]11096.

[127138]Erlander, M. G.; Tillakaratne, N. J. K.; Feldblum, S.; Patel,

N.;

[127139]Tobin, A. J.: Two genes encode distinct glutamate decarboxylases. Neuron 7:

[127140]91–100, 1991.

[127141]

[127142]

[127143]11097.

[127144]Kelly, C. D.; Edwards, Y.; Johnstone, A. P.; Harfst, E.; No-gradi,

[127145]A.; Nussey, S. S.; Povey, S.; Carter, N. D.: Nucleotide sequence

[127146]and chromosomal assignment of a cDNA encoding the large isoform of

[127147]human glutamate decarboxylase. Ann. Hum. Genet. 56: 255–265, 1992.

[127148]

[127149]

[127150]11098.

[127151]Krishnamoorthy, K. S.: Pyridoxine-dependency seizure: report

[127152]of a rare presentation. Ann. Neurol. 13: 103–104, 1983.

[127153]

[127154]

[127155]11099.

[127156]Sparkes, R. S.; Kaufman, D. L.; Heinzmann, C.; Tobin, A. J.;
Mohandas,

[127157]T.: Brain glutamate decarboxylase (GAD) gene assigned to
human chromosome

[127158]2 by somatic cell hybrid analysis. (Abstract) Cytogenet.
Cell Genet. 46:

[127159]696 only, 1987.

[127160]

[127161]

[127162]11100.

[127163]Kedra, D.; Seroussi, E.; Fransson, I.; Trifunovic, J.; Clark,
M.;

[127164]Lagercrantz, J.; Blennow, E.; Mehlin, H.; Dumanski, J.: The
germinal

[127165]center kinase gene and a novel CDC25-like gene are lo-
cated in the

[127166]vicinity of the PYGM gene on 11q13. Hum. Genet. 100:
611-619, 1997.

[127167]

[127168]

[127169]11101.

[127170]Hirsch, D. S.; Pirone, D. M.; Burbelo, P. D.: A new family of

[127171]Cdc42 effector proteins, CEPs, function in fibroblast and
epithelial

[127172]cell shape changes. J. Biol. Chem. 276: 875–883, 2001.

[127173]

[127174]

[127175]11102.

[127176]Joberty, G.; Perlungher, R. R.; Macara, I. G.: The Borgs, a
new

[127177]family of Cdc42 and TC10 GTPase–interacting proteins.
Molec. Cell.

[127178]Biol. 19: 6585–6597, 1999.

[127179]

[127180]

[127181]11103.

[127182]Yang, Z.–Q.; Imoto, I.; Fukuda, Y.; Pimkhaokham, A.; Shi–

mada, Y.;

[127183]Imamura, M.; Sugano, S.; Nakamura, Y.; Inazawa, J.: Identification

[127184]of a novel gene, GASC1, within an amplicon at 9p23–24 frequently detected

[127185]in esophageal cancer cell lines. Cancer Res. 60: 4735–4739, 2000.

[127186]

[127187]

[127188]11104.

[127189]Naar, A. M.; Beaurang, P. A.; Zhou, S.; Abraham, S.; Solomon, W.;

[127190]Tjian, R.: Composite co-activator ARC mediates chromatin-directed

[127191]transcriptional activation. Nature 398: 828–832, 1999.

[127192]

[127193]

[127194]11105.

[127195]Rachez, C.; Lemon, B. D.; Suldan, Z.; Bromleigh, V.; Gamble, M.;

[127196]Naar, A. M.; Erdjument-Bromage, H.; Tempst, P.; Freed-

man, L. P.:

[127197]Ligand-dependent transcription activation by nuclear receptors requires

[127198]the DRIP complex. Nature 398: 824–828, 1999.

[127199]

[127200]

[127201]11106.

[127202]Scott, A. F.: Personal Communication. Baltimore, Md.
3/7/2001.

[127203]

[127204]

[127205]11107.

[127206]LeCouter, J.; Kowalski, J.; Foster, J.; Hass, P.; Zhang, Z.;
Dillard-Telm,

[127207]L.; Frantz, G.; Rangell, L.; DeGuzman, L.; Keller, G.–A.;
Peale, F.;

[127208]Gurney, P.; Hillan, K. J.; Ferrara, N.: Identification of an angiogenic

[127209]mitogen selective for endocrine gland endothelium. Nature 412: 877–884,

[127210]2001.

[127211]

[127212]

[127213]11108.

[127214]Li, M.; Bullock, C. M.; Knauer, D. J.; Ehlert, F. J.; Zhou, Q.

[127215]Y.: Identification of two prokineticin cDNAs: recombinant proteins

[127216]potently contract gastrointestinal smooth muscle. Molec. Pharm. 59:

[127217]692–698, 2001.

[127218]

[127219]

[127220]11109.

[127221]Sun, Q.–A.; Wu, Y.; Zappacosta, F.; Jeang, K.–T.; Lee, B. J.; Hatfield,

[127222]D. L.; Gladyshev, V. N.: Redox regulation of cell signaling by selenocysteine

[127223]in mammalian thioredoxin reductases. J. Biol. Chem. 274: 24522–24530,

[127224]1999.

[127225]

[127226]

[127227]11110.

[127228]Nagase, T.; Nakayama, M.; Nakajima, D.; Kikuno, R.;
Ohara, O.:

[127229]Prediction of the coding sequences of unidentified human
genes. XX.

[127230]The complete sequences of 100 new cDNA clones from
brain which code

[127231]for large proteins in vitro. DNA Res. 8: 85–95, 2001.

[127232]

[127233]

[127234]11111.

[127235]Telford, E. A. R.; Moynihan, L. M.; Markham, A. F.; Lench,
N. J.

[127236]: Isolation and characterisation of a cDNA encoding the
precursor

[127237]for a novel member of the acyl–CoA dehydrogenase gene
family. Biochim.

[127238]Biophys. Acta 1446: 371–376, 1999.

[127239]

[127240]

[127241]11112.

[127242]Jaquemar, D.; Schenker, T.; Trueb, B.: An ankyrin-like protein

[127243]with transmembrane domains is specifically lost after oncogenic transformation

[127244]of human fibroblasts. J. Biol. Chem. 274: 7325–7333, 1999.

[127245]

[127246]

[127247]11113.

[127248]Schenker, T.; Trueb, B.: Down-regulated proteins of mesenchymal

[127249]tumor cells. Exp. Cell Res. 239: 161–168, 1998.

[127250]

[127251]

[127252]11114.

[127253]Cerretti, D. P.; DuBose, R. F.; Black, R. A.; Nelson, N.: Isolation

[127254]of two novel metalloproteinase-disintegrin (ADAM) cDNAs

that show

[127255]testis-specific gene expression. Biochem. Biophys. Res. Commun. 263:

[127256]810–815, 1999.

[127257]

[127258]

[127259]11115.

[127260]Xu, R.; Cai, J.; Xu, T.; Zhou, W.; Ying, B.; Deng, K.; Zhao, S.;

[127261]Li, C.: Molecular cloning and mapping of a novel ADAM gene (ADAM29)

[127262]to human chromosome 4. Genomics 62: 537–539, 1999.

[127263]

[127264]

[127265]11116.

[127266]Jacobs, S.; Schilf, C.; Fliegert, F.; Kolling, S.; Weber, Y.; Schurmann,

[127267]A.; Joost, H.–G.: ADP-ribosylation factor (ARF)-like 4, 6, and 7

[127268]represent a subgroup of the ARF family characterized by rapid nucleotide

[127269]exchange and a nuclear localization signal. FEBS Lett. 456:
384–388,
[127270]1999.

[127271]

[127272]

[127273]11117.

[127274]Lako, M.; Lindsay, S.; Bullen, P.; Wilson, D. I.; Robson, S.
C.;

[127275]Strachan, T.: A novel mammalian Wnt gene, WNT8B, shows
brain–restricted

[127276]expression in early development, with sharply delimited
expression

[127277]boundaries in the developing forebrain. Hum. Molec.
Genet. 7: 813–822,

[127278]1998.

[127279]

[127280]

[127281]11118.

[127282]Lako, M.; Strachan, T.; Curtis, A. R. J.; Lindsay, S.: Isolation

[127283]and characterization of WNT8B, a novel human Wnt gene

that maps to

[127284]10q24. Genomics 35: 386–388, 1996.

[127285]

[127286]

[127287]11119.

[127288]Toda, T.; Iida, A.; Miwa, T.; Nakamura, Y.; Imai, T.: Isolation
tion

[127289]and characterization of a novel gene encoding nuclear
protein at a

[127290]locus (D11S636) tightly linked to multiple endocrine neoplasia type
plasia type

[127291]1 (MEN1). Hum. Molec. Genet. 3: 465–470, 1994.

[127292]

[127293]

[127294]11120.

[127295]Nusse, R.; Varmus, H. E.: Many tumors induced by the
mouse mammary

[127296]tumor virus contain a provirus integrated in the same region of the
gion of the

[127297]host genome. Cell 31: 99–109, 1982.

[127298]

[127299]

[127300]11121.

[127301]Scholler, J. K.; Kanner, S. B.: The human p167 gene encodes a

[127302]unique structural protein that contains centrosomin A homology and

[127303]associates with a multicomponent complex. DNA Cell Biol. 16: 515–531,

[127304]1997.

[127305]

[127306]

[127307]11122.

[127308]Abdulkadir, S. A.; Magee, J. A.; Peters, T. J.; Kaleem, Z.; Naughton,

[127309]C. K.; Humphrey, P. A.; Milbrandt, J.: Conditional loss of Nkx3.1

[127310]in adult mice induces prostatic intraepithelial neoplasia. Molec.

[127311]Cell. Biol. 22: 1495–1503, 2002.

[127312]

[127313]

[127314]11123.

[127315]Bieberich, C. J.; Fujita, K.; He, W. W.; Jay, G.: Prostate-specific
cific

[127316]and androgen-dependent expression of a novel home-
obox gene. J. Biol.

[127317]Chem. 271: 31779–31782, 1996.

[127318]

[127319]

[127320]11124.

[127321]Korkmaz, K. S.; Korkmaz, C. G; Ragnhildstveit, E.;
Kizildag, S.;

[127322]Pretlow, T. G.; Saatcioglu, F.: Full-length cDNA sequence
and genomic

[127323]organization of human NKX3A--alternative forms and
regulation by both

[127324]androgens and estrogens. Gene 260: 25–36, 2000.

[127325]

[127326]

[127327]11125.

[127328]He, W. W.; Sciavolino, P. J.; Wing, J.; Augustus, M.; Hudson, P.;

[127329]Meissner, P. S.; Curtis, R. T.; Shell, B. K.; Bostwick, D. G.; Tindall,

[127330]D. J.; Gelmann, E. P.; Abate-Shen, C.; Carter, K. C.: A novel human

[127331]prostate-specific, androgen-regulated homeobox gene (NKX3.1) that

[127332]maps to 8p21, a region frequently deleted in prostate cancer. Genomics 43:

[127333]69–77, 1997.

[127334]

[127335]

[127336]11126.

[127337]Argyropoulos, G.; Brown, A. M.; Willi, S. M.; Zhu, J.; He, Y.;

[127338]Reitman, M.; Geva, S. M.; Spruill, I.; Garvey, W. T.: Effects of

[127339]mutations in the human uncoupling protein 3 gene on the respiratory

[127340]quotient and fat oxidation in severe obesity and type 2 di-

abetes. J.

[127341]Clin. Invest. 102: 1345–1351, 1998.

[127342]

[127343]

[127344]11127.

[127345]Boss, O.; Giacobino, J.–P.; Muzzin, P.: Genomic structure
of uncoupling

[127346]protein–3 (UCP3) and its assignment to chromosome
11q13. Genomics 47:

[127347]425–426, 1998.

[127348]

[127349]

[127350]11128.

[127351]Boss, O.; Samec, S.; Paoloni–Giacobino, A.; Rossier, C.;
Dulloo,

[127352]A.; Seydoux, J.; Muzzin, P.; Giacobino, J.–P.: Uncoupling
protein–3:

[127353]a new member of the mitochondrial carrier family with
tissue–specific

[127354]expression. FEBS Lett. 408: 39–42, 1997.

[127355]

[127356]

[127357]11129.

[127358]Brown, A. M.; Dolan, J. W.; Willi, S. M.; Garvey, W. T.; Argypoulos,

[127359]G.: Endogenous mutations in human uncoupling protein 3 alter its

[127360]functional properties. FEBS Lett. 464: 189–193, 1999.

[127361]

[127362]

[127363]11130.

[127364]Bilaud, T.; Brun, C.; Ancelin, K.; Koering, C. E.; Laroche, T.;

[127365]Gilson, E.: Telomeric localization of TRF2, a novel human telobox

[127366]protein. Nature Genet. 17: 236–239, 1997.

[127367]

[127368]

[127369]11131.

[127370]Broccoli, D.; Smogorzewska, A.; Chong, L.; de Lange, T.:

Human

[127371]telomeres contain two distinct Myb-related proteins, TRF1 and TRF2. Nature

[127372]Genet. 17: 231–235, 1997.

[127373]

[127374]

[127375]11132.

[127376]Karlseder, J.; Smogorzewska, A.; de Lange, T.: Senescence induced

[127377]by altered telomere state, not telomere loss. Science 295: 2446–2449,

[127378]2002.

[127379]

[127380]

[127381]11133.

[127382]Sakaguchi, A. Y.; Padalecki, S. S.; Mattern, V.; Rodriguez, A.;

[127383]Leach, R. J.; McGill, J. R.; Chavez, M.; Giambernardi, T. A.: Chromosomal

[127384]sublocalization of the transcribed human telomere repeat

binding factor

[127385]2 gene and comparative mapping in the mouse. Somat.

Cell Molec. Genet. 24:

[127386]157–163, 1998.

[127387]

[127388]

[127389]11134.

[127390]van Steensel, B.; Smogorzewska, A.; de Lange, T.: TRF2
protects

[127391]human telomeres from end-to-end fusions. Cell 92:
401–413, 1998.

[127392]

[127393]

[127394]11135.

[127395]Parr, B. A.; McMahon, A. P.: Sexually dimorphic development of

[127396]the mammalian reproductive tract requires Wnt-7a. Nature 395: 707-710,

[127397]1998.

[127398]

[127399]

[127400]11136.

[127401]Barron-Casella, E. A.; Torres, M. A.; Scherer, S. W.; Heng, H.

[127402]H. Q.; Tsui, L.-C.; Casella, J. F.: Sequence analysis and chromosomal

[127403]localization of human Cap Z: conserved residues within the actin-binding

[127404]domain may link Cap Z to gelsolin/severin and profilin protein families. J.

[127405]Biol. Chem. 270: 21472-21479, 1995.

[127406]

[127407]

[127408]11137.

[127409]Hart, M. C.; Korshunova, Y. O.; Cooper, J. A.: Mapping of
the

[127410]mouse actin capping protein alpha subunit genes and
pseudogenes. Genomics 39:

[127411]264–270, 1997.

[127412]

[127413]

[127414]11138.

[127415]Cooper, J. A.: Personal Communication. St. Louis, Mo.
3/8/1999.

[127416]

[127417]

[127418]11139.

[127419]Cardoso, C.; Mignon, C.; Hetet, G.; Grandchamps, B.;
Fontes, M.;

[127420]Colleaux, L.: The human EZH2 gene: genomic organisation
and revised

[127421]mapping in 7q35 within the critical region for malignant
myeloid disorders. Europ.

[127422]J. Hum. Genet. 8: 174–180, 2000.

[127423]

[127424]

[127425]11140.

[127426]Chen, H.; Rossier, C.; Antonarakis, S. E.: Cloning of a human
man

[127427]homolog of the Drosophila enhancer of zeste gene (EZH2)
that maps

[127428]to chromosome 21q22.2. Genomics 38: 30–37, 1996.

[127429]

[127430]

[127431]11141.

[127432]Dohner, K.; Brown, J.; Hehmann, U.; Hetzel, C.; Stewart, J.;
Lowther,

[127433]G.; Scholl, C.; Frohling, S.; Cuneo, A.; Tsui, L. C.; Lichter,
P.;

[127434]Scherer, S. W.; Dohner, H.: Molecular cytogenetic charac-
terization

[127435]of a critical region in bands 7q35–q36 commonly deleted
in malignant

[127436]myeloid disorders. Blood 92: 4031–4035, 1998.

[127437]

[127438]

[127439]11142.

[127440]Laible, G.; Haynes, A. R.; Lebersorger, A.; O'Carroll, D.;
Mattei,

[127441]M.–G.; Denny, P.; Brown, S. D. M.; Jenuwein, T.: The
murine polycomb–group

[127442]genes Ezh1 and Ezh2 map close to Hox gene clusters on
mouse chromosomes

[127443]11 and 6. Mammalian Genome 10: 311–314, 1999.

[127444]

[127445]

[127446]11143.

[127447]Varambally, S.; Dhanasekaran, S. M.; Zhou, M.; Barrette, T.
R.;

[127448]Kumar–Sinha, C.; Sanda, M. G.; Ghosh, D.; Pienta, K. J.;
Sewalt, R.

[127449]G. A. B.; Otte, A. P.; Rubin, M. A.; Chinnaiyan, A. M.: The
polycomb

[127450]group protein EZH2 is involved in progression of prostate cancer. Nature 419:

[127451]624–629, 2002.

[127452]

[127453]

[127454]11144.

[127455]Morohoshi, F.; Arai, K.; Takahashi, E.; Tanigami, A.; Ohki, M.

[127456]: Cloning and mapping of a human RBP56 gene encoding a putative RNA

[127457]binding protein similar to FUS/TLS and EWS proteins. Genomics 38:

[127458]51–57, 1996.

[127459]

[127460]

[127461]11145.

[127462]Batt, J.; Asa, S.; Fladd, C.; Rotin, D.: Pituitary, pancreatic

[127463]and gut neuroendocrine defects in protein tyrosine phosphatase-sigma-deficient

[127464]mice. Molec. Endocr. 16: 155–169, 2002.

[127465]

[127466]

[127467]11146.

[127468]Elchebly, M.; Wagner, J.; Kennedy, T. E.; Lanctot, C.;
Michaliszyn,

[127469]E.; Itie, A.; Drouin, J.; Tremblay, M. L.: Neuroendocrine
dysplasia

[127470]in mice lacking protein tyrosine phosphatase sigma. Na-
ture Genet. 21:

[127471]330–333, 1999.

[127472]

[127473]

[127474]11147.

[127475]Pulido, R.; Serra–Pages, C.; Tang, M.; Streuli, M.: The LAR/
PTP

[127476]delta/PTP sigma subfamily of transmembrane protein–ty-
rosine–phosphatases:

[127477]multiple human LAR, PTP delta, and PTP sigma isoforms
are expressed

[127478]in a tissue–specific manner and associate with the LAR–
interacting

[127479]protein LIP.1. Proc. Nat. Acad. Sci. 92: 11686–11690,
1995.

[127480]

[127481]

[127482]11148.

[127483]Wagner, J.; Gordon, L. A.; Heng, H. H. Q.; Tremblay, M. L.;
Olsen,

[127484]A. S.: Physical mapping of receptor type protein tyrosine
phosphatase

[127485]sigma (PTPRS) to human chromosome 19p13.3. Genomics
38: 76–78, 1996.

[127486]

[127487]

[127488]11149.

[127489]Wallace, M. J.; Batt, J.; Fladd, C. A.; Henderson, J. T.;
Skarnes,

[127490]W.; Rotin, D.: Neuronal defects and posterior pituitary hy-
poplasia

[127491]in mice lacking the receptor tyrosine phosphatase PTP-
sigma. Nature

[127492]Genet. 21: 334–338, 1999.

[127493]

[127494]

[127495]11150.

[127496]Bruyns, E.; Mincheva, A.; Bruyns, R. M.; Kirchgessner, H.;
Weitz,

[127497]S.; Lichter, P.; Meuer, S.; Schraven, B.: Sequence, genomic
organization,

[127498]and chromosomal localization of the human LPAP
(PTPRCAP) and mouse

[127499]CD45-AP/LSM-1 genes. Genomics 38: 79-83, 1996.

[127500]

[127501]

[127502]11151.

[127503]Schraven, B.; Schoenhaut, D.; Bruyns, E.; Koretzky, G.;
Eckerskorn,

[127504]C.; Wallich, R.; Kirchgessner, H.; Sakorafas, P.; Labkovsky,
B.; Ratnofsky,

[127505]S.; Meuer, S.: LPAP, a novel 32-kDa phosphoprotein that
interacts

[127506]with CD45 in human lymphocytes. J. Biol. Chem. 269:
29102-29111,

[127507]1994.

[127508]

[127509]

[127510]11152.

[127511]Bates, S.; Rowan, S.; Vousden, K. H.: Characterisation of
human

[127512]cyclin G1 and G2: DNA damage inducible genes. Onco-
gene 13: 1103-1109,

[127513]1996.

[127514]

[127515]

[127516]11153.

[127517]Endo, Y.; Fujita, T.; Tamura, K.; Tsuruga, H.; Nojima, H.:
Structure

[127518]and chromosomal assignment of the human cyclin G
gene. Genomics 38:

[127519]92-95, 1996.

[127520]

[127521]

[127522]11154.

[127523]Horne, M. C.; Goolsby, G. L.; Donaldson, K. L.; Tran, D.;
Neubauer,

[127524]M.; Wahl, A. F.: Cyclin G1 and cyclin G2 comprise a new
family of

[127525]cyclins with contrasting tissue-specific and cell cycle-
regulated

[127526]expression. J. Biol. Chem. 271: 6050-6061, 1996.

[127527]

[127528]

[127529]11155.

[127530]Jensen, M. R.; Factor, V. M.; Zimonjic, D. B.; Miller, M. J.;
Keck,

[127531]C. L.; Thorgeirsson, S. S.: Chromosome localization and
structure

[127532]of the murine cyclin G1 gene promoter sequence. Ge-
nomics 45: 297-303,

[127533]1997.

[127534]

[127535]

[127536]11156.

[127537]Spicer, A. P.; Augustine, M. L.; McDonald, J. A.: Molecular cloning

[127538]and characterization of a putative mouse hyaluronan synthase. J.

[127539]Biol. Chem. 271: 23400–23406, 1996.

[127540]

[127541]

[127542]11157.

[127543]Skalhegg, B. S.; Huang, Y.; Su, T.; Idzerda, R. L.; McKnight, G.

[127544]S.; Burton, K. A.: Mutation of the C-alpha subunit of PKA leads to

[127545]growth retardation and sperm dysfunction. Molec. Endocr. 16: 630–639,

[127546]2002.

[127547]

[127548]

[127549]11158.

[127550]Tasken, K.; Solberg, R.; Zhao, Y.; Hansson, V.; Jahnsen, T.;

Siciliano,

[127551]M. J.: The gene encoding the catalytic subunit C- α of
cAMP-dependent

[127552]protein kinase (locus PRKACA) localizes to human chro-
mosome region

[127553]19p13.1. Genomics 36: 535-538, 1996.

[127554]

[127555]

[127556]11159.

[127557]Kim, J.; Uyemura, K.; Van Dyke, M. K.; Legaspi, A. J.; Rea,
T.

[127558]H.; Shuai, K.; Modlin, R. L.: A role for IL-12 receptor ex-
pression

[127559]and signal transduction in host defense in leprosy. J. Im-
mun. 167:

[127560]779-786, 2001.

[127561]

[127562]

[127563]11160.

[127564]Morton, S. M.; Bocaccio, I.; Depetris, D.; Mattei, M.; Des-

sein,

[127565]A.: Assignment of IL12RB2 to human chromosome
1p31.3–p31.2 between

[127566]D1S230 and D1S198. Cytogenet. Cell Genet. 79: 282–283,
1997.

[127567]

[127568]

[127569]11161.

[127570]McCright, B.; Brothman, A. R.; Virshup, D. M.: Assignment
of human

[127571]protein phosphatase 2A regulatory subunit genes
B56–alpha, B56–beta,

[127572]B56–gamma, B56–delta, and B56–epsilon
(PPP2R5A–PPP2R5E), highly expressed

[127573]in muscle and brain, to chromosome regions 1q41,
11q12, 3p21, 6p21.1,

[127574]and 7p11.2–to–p12. Genomics 36: 168–170, 1996.

[127575]

[127576]

[127577]11162.

[127578]McCright, B.; Virshup, D. M.: Identification of a new family

of

[127579]protein phosphatase 2A regulatory subunits. J. Biol. Chem. 270:

[127580]26123–26128, 1995.

[127581]

[127582]

[127583]11163.

[127584]Horikawa, S.; Tsukada, K.: Molecular cloning and developmental

[127585]expression of a human kidney S-adenosylmethionine synthetase. FEBS

[127586]Lett. 312: 37–41, 1992.

[127587]

[127588]

[127589]11164.

[127590]Mao, Z.; Liu, S.; Cai, J.; Huang, Z.-Z.; Lu, S. C.: Cloning and

[127591]functional characterization of the 5-prime-flanking region of human

[127592]methionine adenosyltransferase 2A gene. Biochem. Biophys. Res. Commun. 248:

[127593]479–484, 1998.

[127594]

[127595]

[127596]11165.

[127597]Clark, J. A.; Bonner, T. I.; Kim, A. S.; Usdin, T. B.: Multiple

[127598]regions of ligand discrimination revealed by analysis of
chimeric

[127599]parathyroid hormone 2 (PTH2) and PTH/PTH-related pep-
tide (PTHrP) receptors. Molec.

[127600]Endocr. 12: 193–206, 1998.

[127601]

[127602]

[127603]11166.

[127604]Usdin, T. B.; Gruber, C.; Bonner, T. I.: Identification and
functional

[127605]expression of a receptor selectively recognizing parathy-
roid hormone,

[127606]the PTH2 receptor. J. Biol. Chem. 270: 15455–15458,
1995.

[127607]

[127608]

[127609]11167.

[127610]Usdin, T. B.; Modi, W.; Bonner, T. I.: Assignment of the human

[127611]PTH2 receptor gene (PTHR2) to chromosome 2q33 by fluorescence in situ

[127612]hybridization. Genomics 37: 140–141, 1996.

[127613]

[127614]

[127615]11168.

[127616]Faure, S.; Meyer, L.; Costagliola, D.; Vaneensberghe, C.; Genin,

[127617]E.; Autran, B.; French ALT and IMMUNOCO Study Groups; Delfraisay,

[127618]J.-F.; SEROCO Study Group; McDermott, D. H.; Murphy, P. M.; Debre,

[127619]P.; Theodorou, I.; Cambadiere, C.: Rapid progression to AIDS in HIV+

[127620]individuals with a structural variant of the chemokine receptor CX(3)CR1. Science 287:

[127621]2274–2277, 2000.

[127622]

[127623]

[127624]11169.

[127625]Imai, T.; Hieshima, K.; Haskell, C.; Baba, M.; Nagira, M.;
Nishimura,

[127626]M.; Kakizaki, M.; Takagi, S.; Nomiyama, H.; Schall, T. J.;
Yoshie,

[127627]O.: Identification and molecular characterization of
fractalkine

[127628]receptor CX3CR1, which mediates both leukocyte migra-
tion and adhesion. Cell 91:

[127629]521–530, 1997.

[127630]

[127631]

[127632]11170.

[127633]Moatti, D.; Faure, S.; Fumeron, F.; Amara, M. E. W.; Sek-
nadj,

[127634]P.; McDermott, D. H.; Debre, P.; Aumont, M. C.; Murphy, P.
M.; de

[127635]Prost, D.; Combadiere, C.: Polymorphism in the fractalkine
receptor

[127636]CX3CR1 as a genetic risk factor for coronary artery disease. Blood 97:

[127637]1925–1928, 2001.

[127638]

[127639]

[127640]11171.

[127641]Raport, C. J.; Schweickart, V. L.; Eddy, R. L., Jr.; Shows, T.

[127642]B.; Gray, P. W.: The orphan G-protein-coupled receptor-encoding gene

[127643]V28 is closely related to genes for chemokine receptors and is expressed

[127644]in lymphoid and neural tissues. Gene 163: 295–299, 1995.

[127645]

[127646]

[127647]11172.

[127648]Tripp, R. A.; Jones, L. P.; Haynes, L. M.; Zheng, H.; Murphy, P.

[127649]M.; Anderson, L. J.: CX3C chemokine mimicry by respiratory syncytial

[127650]virus G glycoprotein. Nature Immun. 2: 732–738, 2001.

[127651]

[127652]

[127653]11173.

[127654]Lafreniere, R. G.; Rochefort, D. L.; Chretien, N.; Neville, C.

[127655]E.; Korneluk, R. G.; Zuo, L.; Wei, Y.; Lichter, J.; Rouleau, G.

A.

[127656]: Isolation and genomic structure of a human homolog of
the yeast

[127657]periodic tryptophan protein 2 (PWP2) gene mapping to
21q22.3. Genome

[127658]Res. 6: 1216–1226, 1996.

[127659]

[127660]

[127661]11174.

[127662]Lalioi, M. D.; Chen, H.; Rossier, C.; Shafaatian, R.; Reid, J.

[127663]D.; Antonarakis, S. E.: Cloning the cDNA of human PWP2,
which encodes

[127664]a protein with WD repeats and maps to 21q22.3. Ge-

nomics 35: 321–327,

[127665]1996.

[127666]

[127667]

[127668]11175.

[127669]Yamakawa, K.; Gao, D.–Q.; Korenberg, J. R.: A periodic
tryptophan

[127670]protein 2 gene homologue (PWP2H) in the candidate re-
gion of progressive

[127671]myoclonus epilepsy on 21q22.3. Cytogenet. Cell Genet.
74: 140–145,

[127672]1996.

[127673]

[127674]

[127675]11176.

[127676]Ben–David, Y.; Letwin, K.; Tannock, L.; Bernstein, A.; Paw-
son,

[127677]T.: A mammalian protein kinase with potential for serine/
threonine

[127678]and tyrosine phosphorylation is related to cell cycle regu-

lators. EMBO

[127679]J. 10: 317–325, 1991.

[127680]

[127681]

[127682]11177.

[127683]Colwill, K.; Feng, L. L.; Yeakley, J. M.; Gish, G. D.; Caceres,

[127684]J. F.; Pawson, T.; Fu, X.–D.: SRPK1 and Clk/Sty protein kinases show

[127685]distinct substrate specificities for serine/arginine-rich splicing

[127686]factors. J. Biol. Chem. 271: 24569–24575, 1996.

[127687]

[127688]

[127689]11178.

[127690]Hanes, J.; von der Kammer, H.; Klaudiny, J.; Scheit, K. H.: Characterization

[127691]by cDNA cloning of two new human protein kinases: evidence by sequence

[127692]comparison of a new family of mammalian protein kinases. J. Molec.

[127693]Biol. 244: 665–672, 1994.

[127694]

[127695]

[127696]11179.

[127697]Howell, B. W.; Afar, D. E. H.; Lew, J.; Douville, E. M. J.; Icely,

[127698]P. L. E.; Gray, D. A.; Bell, J. C.: STY, a tyrosine–
phosphorylating

[127699]enzyme with sequence homology to serine/threonine ki–
nases. Molec.

[127700]Cell Biol. 11: 568–572, 1991.

[127701]

[127702]

[127703]11180.

[127704]Johnson, K. W.; Smith, K. A.: Molecular cloning of a novel
human

[127705]cdc2/CDC28–like protein kinase. J. Biol. Chem. 266:
3402–3407, 1991.

[127706]

[127707]

[127708]11181.

[127709]Talmadge, C. B.; Finkernagel, S.; Sumegi, J.; Sciorra, L.;
Rabinow,

[127710]L.: Chromosomal mapping of three human LAMMER pro-
tein-kinase-encoding

[127711]genes. Hum. Genet. 103: 523-524, 1998.

[127712]

[127713]

[127714]11182.

[127715]Andersen, G.; Poterszman, A.; Egly, J. M.; Moras, D.;
Thierry,

[127716]J.-C: The crystal structure of human cyclin H. FEBS Lett.
397:

[127717]65-69, 1996.

[127718]

[127719]

[127720]11183.

[127721]Eki, T.; Okumura, K.; Abe, M.; Kagotani, K.; Taguchi, H.;
Murakami,

[127722]Y.; Pan, Z.-Q.; Hanaoka, F.: Mapping of the human genes

encoding

[127723]cyclin H (CCNH) and the CDK-activating kinase (CAK) assembly factor

[127724]MAT1 (MNAT1) to chromosome bands 5q13.3-q14 and 14q23, respectively. Genomics 47:

[127725]115-120, 1998.

[127726]

[127727]

[127728]11184.

[127729]Fisher, R. P.; Morgan, D. O.: A novel cyclin associates with MO15/CDK7

[127730]to form the CDK-activating kinase. Cell 78: 713-724, 1994.

[127731]

[127732]

[127733]11185.

[127734]Shiekhattar, R.; Mermelstein, F.; Fisher, R. P.; Drapkin, R.; Dynlacht,

[127735]B.; Wessling, H. C.; Morgan, D. O.; Reinberg: Cdk-activating kinase

[127736]complex is a component of human transcription factor

TFIIH. Nature 374:

[127737]283–287, 1995.

[127738]

[127739]

[127740]11186.

[127741]Alders, M.; Hodges, M.; Hadjantonakis, A.–K.; Postmus, J.;
van

[127742]Wijk, I.; Blik, J.; de Meulemeester, M.; Westerveld, A.;
Guillemot,

[127743]F.; Oudejans, C.; Little, P.; Mannens, M.: The human
Achaete–Scute

[127744]homologue 2 (ASCL2,HASH2) maps to chromosome
11p15.5, close to IGF2

[127745]and is expressed in extravillous trophoblasts. Hum. Molec.
Genet. 6:

[127746]859–867, 1997.

[127747]

[127748]

[127749]11187.

[127750]Guillemot, F.; Nagy, A.; Auerbach, A.; Rossant, J.; Joyner,

A.

[127751]L.: Essential role of Mash-2 in extraembryonic development. Nature 371:

[127752]333-336, 1994.

[127753]

[127754]

[127755]11188.

[127756]Banga, S. S.; Ozer, H. L.; Park, S.-K.; Lee, S.-T.: Assignment

[127757]of PTK7 encoding a receptor protein tyrosine kinase-like molecule

[127758]to human chromosome 6p21.1-p12.2 by fluorescence in situ hybridization. Cytogenet.

[127759]Cell Genet. 76: 43-44, 1997.

[127760]

[127761]

[127762]11189.

[127763]Lee, S.-T.; Strunk, K. M.; Spritz, R. A.: A survey of protein

[127764]tyrosine kinase mRNAs expressed in normal human melanocytes. Oncogene 8:

[127765]3403–3410, 1993.

[127766]

[127767]

[127768]11190.

[127769]Mossie, K.; Jallal, B.; Alves, F.; Sures, I.; Plowman, G. D.;
Ullrich,

[127770]A.: Colon carcinoma kinase–4 defines a new subclass of
the receptor

[127771]tyrosine kinase family. *Oncogene* 11: 2179–2184, 1995.

[127772]

[127773]

[127774]11191.

[127775]Park, S.–K.; Lee, H.–S.; Lee, S.–T.: Characterization of the
human

[127776]full–length PTK7 cDNA encoding a receptor protein tyro–
sine kinase–like

[127777]molecule closely related to chick KLG. *J. Biochem.* 119:
235–239,

[127778]1996.

[127779]

[127780]

[127781]11192.

[127782]Sotiropoulou, G.; Anisowicz, A.; Sager, R.: Identification,
cloning,

[127783]and characterization of cystatin M, a novel cysteine pro-
teinase inhibitor,

[127784]down-regulated in breast cancer. J. Biol. Chem. 272:
903-910, 1997.

[127785]

[127786]

[127787]11193.

[127788]Stenman, G.; Astrom, A.-K.; Roijer, E.; Sotiropoulou, G.;
Zhang,

[127789]M.; Sager, R.: Assignment of a novel cysteine proteinase
inhibitor

[127790](CST6) to 11q13 by fluorescence in situ hybridization. Cy-
togenet.

[127791]Cell Genet. 76: 45-46, 1997.

[127792]

[127793]

[127794]11194.

[127795]Kohler, M.; Ansieau, S.; Prehn, S.; Leutz, A.; Haller, H.;
Hartmann,

[127796]E.: Cloning of two novel human importin- α subunits
and analysis

[127797]of the expression pattern of the importin- α protein
family. FEBS

[127798]Lett. 417: 104–108, 1997.

[127799]

[127800]

[127801]11195.

[127802]Moore, M. S.; Blobel, G.: The two steps of nuclear import,
targeting

[127803]to the nuclear envelope and translocation through the nu-
clear pore,

[127804]require different cytosolic factors. Cell 69: 939–950,
1992.

[127805]

[127806]

[127807]11196.

[127808]Takeda, S.; Fujiwara, T.; Shimizu, F.; Kawai, A.; Shinomiya, K.;

[127809]Okuno, S.; Ozaki, K.; Katagiri, T.; Shimada, Y.; Nagata, M.; Watanabe,

[127810]T.; Takaichi, A.; Kuga, Y.; Suzuki, M.; Hishigaki, H.; Taka-
hashi,

[127811]E.; Shin, S.; Nakamura, Y.; Hirai, Y.: Isolation and mapping
of karyopherin

[127812]alpha-3 (KPNA3), a human gene that is highly homolo-
gous to genes encoding

[127813]Xenopus importin, yeast SRP1 and human RCH1. Cyto-
genet. Cell Genet. 76:

[127814]87-93, 1997.

[127815]

[127816]

[127817]11197.

[127818]Zhang, C.-Y.; Baffy, G.; Perret, P.; Krauss, S.; Peroni, O.;
Grujic,

[127819]D.; Hagen, T.; Vidal-Puig, A.; Boss, O.; Kim, Y.-B.; Zheng,
X. X.;

[127820]Wheeler, M. B.; Shulman, G. I.; Chan, C. B.; Lowell, B. B.:
Uncoupling

[127821]protein-2 negatively regulates insulin secretion and is a major link

[127822]between obesity, beta cell dysfunction, and type 2 diabetes. Cell 105:

[127823]745–755, 2001.

[127824]

[127825]

[127826]11198.

[127827]Edlund, A.; Johansson, T.; Leidvik, B.; Hansson, L.: Structure

[127828]of the human kappa-casein gene. Gene 174: 65–69, 1996.

[127829]

[127830]

[127831]11199.

[127832]Huber, R.; Carrell, R. W.: Implications of the three-dimensional

[127833]structure of alpha 1-antitrypsin for structure and function of serpins. Biochemistry 28:

[127834]8951–8966, 1989.

[127835]

[127836]

[127837]11200.

[127838]Sprecher, C. A.; Morgenstern, K. A.; Mathewes, S.; Dahlen,
J. R.;

[127839]Schrader, S. K.; Foster, D. C.; Kisiel, W.: Molecular cloning,
expression,

[127840]and partial characterization of two novel members of the
ovalbumin

[127841]family of serine proteinase inhibitors. J. Biol. Chem. 270:
29854–29861,

[127842]1995.

[127843]

[127844]

[127845]11201.

[127846]Chevalier, D.; Cauffiez, C.; Bernard, C.; Lo-Guidice, J.–M.;
Allorge,

[127847]D.; Fazio, F.; Ferrari, N.; Libersa, C.; Lhermitte, M.;
D'Halluin,

[127848]J.–C.; Broly, F.: Characterization of new mutations in the
coding

[127849]sequence and 5-prime-untranslated region of the human prostacyclin

[127850]synthase gene (CYP8A1). Hum. Genet. 108: 148–155, 2001.

[127851]

[127852]

[127853]11202.

[127854]Barrett, A. J.; Rawlings, N. D.: Evolutionary lines of cysteine

[127855]peptidases. Biol. Chem. 382: 727–733, 2001.

[127856]

[127857]

[127858]11203.

[127859]Fernandes-Alnemri, T.; Takahashi, A.; Armstrong, R.; Krebs, J.;

[127860]Fritz, L.; Tomaselli, K. J.; Wang, L.; Yu, Z.; Croce, C. M.; Salveson,

[127861]G.; Earnshaw, W. C.; Litwack, G.; Alnemri, E. S.: Mch3, a novel human

[127862]apoptotic cysteine protease highly related to CPP32. Cancer Res. 55:

[127863]6045–6052, 1995.

[127864]

[127865]

[127866]11204.

[127867]Riedl, S. J.; Fuentes–Prior, P.; Renatus, M.; Kairies, N.;
Krapp,

[127868]S.; Huber, R.; Salvesen, G. S.; Bode, W.: Structural basis for
the

[127869]activation of human procaspase–7. Proc. Nat. Acad. Sci.
98: 14790–14795,

[127870]2001.

[127871]

[127872]

[127873]11205.

[127874]Park, W. S.; Lee, J. H.; Shin, M. S.; Park, J. Y.; Kim, H. S.;

[127875]Lee, J. H.; Kim, Y. S.; Lee, S. N.; Xiao, W.; Park, C. H.; Lee,
S.

[127876]H.; Yoo, N. J.; Lee, J. Y.: Inactivating mutations of the cas–
pase–10

[127877]gene in gastric cancer. Oncogene 21: 2919–2925, 2002.

[127878]

[127879]

[127880]11206.

[127881]Shin, M. S.; Kim, H. S.; Kang, C. S.; Park, W. S.; Kim, S. Y.;

[127882]Lee, S. N.; Lee, J. H.; Park, J. Y.; Jang, J. J.; Kim, C. W.; Kim,

[127883]S. H.; Lee, J. Y.; Yoo, N. J.; Lee, S. H.: Inactivating mutations
tions

[127884]of CASP10 gene in non-Hodgkin lymphomas. Blood 99:
4094–4099, 2002.

[127885]

[127886]

[127887]11207.

[127888]Vincenz, C.; Dixit, V. M.: Fas-associated death domain
protein

[127889]interleukin-1-beta-converting enzyme 2 (FLICE2), an ICE/
Ced-3 homologue,

[127890]is proximally involved in CD95- and p55-mediated death
signaling. J.

[127891]Biol. Chem. 272: 6578–6583, 1997.

[127892]

[127893]

[127894]11208.

[127895]Wang, J.; Chun, H. J.; Wong, W.; Spencer, D. M.; Lenardo, M. J.

[127896]: Caspase-10 is an initiator caspase in death receptor signaling. Proc.

[127897]Nat. Acad. Sci. 98: 13884-13888, 2001.

[127898]

[127899]

[127900]11209.

[127901]Wang, J.; Zheng, L.; Lobito, A.; Chan, F. K.; Dale, J.; Sneller,

[127902]M.; Yao, X.; Puck, J. M.; Straus, S. E.; Lenardo, M. J.: Inherited

[127903]human caspase 10 mutations underlie defective lymphocyte and dendritic

[127904]cell apoptosis in autoimmune lymphoproliferative syndrome type II. Cell 98:

[127905]47-58, 1999.

[127906]

[127907]

[127908]11210.

[127909]Chun, H. J.; Zheng, L.; Ahmad, M.; Wang, J.; Speirs, C. K.;
Siegel,

[127910]R. M.; Dale, J. K.; Puck, J.; Davis, J.; Hall, C. G.; Skoda–
Smith,

[127911]S.; Atkinson, T. P.; Straus, S. E.; Lenardo, M. J.: Pleiotropic
defects

[127912]in lymphocyte activation caused by caspase–8 mutations
lead to human

[127913]immunodeficiency. Nature 419: 395–399, 2002.

[127914]

[127915]

[127916]11211.

[127917]Grenet, J.; Teitz, T.; Wei, T.; Valentine, V.; Kidd, V. J.:
Structure

[127918]and chromosome localization of the human CASP8 gene.
Gene 226: 225–232,

[127919]1999.

[127920]

[127921]

[127922]11212.

[127923]Kischkel, F. C.; Kioschis, P.; Weitz, S.; Poustka, A.; Lichter,

[127924]P.; Krammer, P. H.: Assignment of CASP8 to human chromosome band

[127925]2q33–q34 and Casp8 to the murine syntenic region on chromosome 1B–proximal

[127926]C by in situ hybridization. Cytogenet. Cell Genet. 82: 95–96, 1998.

[127927]

[127928]

[127929]11213.

[127930]Sanchez, I.; Xu, C.–J.; Juo, P.; Kakizaka, A.; Blenis, J.; Yuan,

[127931]J.: Caspase–8 is required for cell death induced by expanded polyglutamine

[127932]repeats. Neuron 22: 623–633, 1999.

[127933]

[127934]

[127935]11214.

[127936]Berkovic, S. F.; Kennerson, M. L.; Howell, R. A.; Scheffer, I.

[127937]E.; Hwang, P. A.; Nicholson, G. A.: Phenotypic expression
of benign

[127938]familial neonatal convulsions linked to chromosome 20.
Arch. Neurol. 51:

[127939]1125–1128, 1994.

[127940]

[127941]

[127942]11215.

[127943]Varfolomeev, E. E.; Schuchmann, M.; Luria, V.; Chian-
nilkulchai,

[127944]N.; Beckmann, J. S.; Mett, I. L.; Rebrikov, D.; Brodianski, V.
M.;

[127945]Kemper, O. C.; Kollet, O.; Lapidot, T.; Soffer, D.; Sobe, T.;
Avraham,

[127946]K. B.; Goncharov, T.; Holtmann, H.; Lonai, P.; Wallach, D.:
Targeted

[127947]disruption of the mouse Caspase 8 gene ablates cell death
induction

[127948]by the TNF receptors, Fas/Apo1, and DR3 and is lethal
prenatally. Immunity 9:

[127949]267–276, 1998.

[127950]

[127951]

[127952]11216.

[127953]Kalchman, M. A.; Koide, H. B.; McCutcheon, K.; Graham, R.
K.; Nichol,

[127954]K.; Nishiyama, K.; Kazemi-Esfarjani, P.; Lynn, F. C.;
Wellington,

[127955]C.; Metzler, M.; Goldberg, Y. P.; Kanazawa, I.; Gietz, R. D.;
Hayden,

[127956]M. R.: HIP1, a human homologue of *S. cerevisiae* Slap2,
interacts

[127957]with membrane-associated huntingtin in the brain. *Nature*
Genet. 16:

[127958]44–53, 1997.

[127959]

[127960]

[127961]11217.

[127962]Wanker, E. E.; Rovira, C.; Scherzinger, E.; Hasenbank, R.;
Walter,

[127963]S.; Tait, D.; Colicelli, J.; Lehrach, H.: HIP-I: a huntingtin in-
teracting

[127964]protein isolated by the yeast two-hybrid system. Hum.

Molec. Genet. 6:

[127965]487-495, 1997.

[127966]

[127967]

[127968]11218.

[127969]Eki, T; Okumura, K.; Shiratori, A.; Abe, M.; Nogami, M.;

Taguchi,

[127970]H.; Shibata, T.; Murakami, Y.; Hanaoka, F.: Assignment of

the closest

[127971]human homologue (DNA2L; KIAA0083) of the yeast Dna2

helicase gene

[127972]to chromosome band 10q21.3-q22.1 Genomics 37:

408-410, 1996.

[127973]

[127974]

[127975]11219.

[127976]Nagase, T; Miyajima, N; Tanaka, A.; Sazuka, T.; Seki, N.;

Sato,

[127977]S.; Tabata, S.; Ishikawa, K.; Kawarabayashi, Y.; Kotani, H.;

Nomura,

[127978]N.: Prediction of the coding sequences of unidentified human genes.

[127979]III. The coding sequences of 40 new genes (KIAA0081–KIAA0120) deduced

[127980]by analysis of cDNA clones from human cell line KG-1.
DNA Res. 2:

[127981]37–43, 1995.

[127982]

[127983]

[127984]11220.

[127985]Kim, J. W.; Lee, Y.; Lee, I. A.; Kang, H. B.; Choe, Y. K.;
Choe,

[127986]I. S.: Cloning and expression of human cDNA encoding
Na(+),K(+)-ATPase

[127987]gamma-subunit. Biochim. Biophys. Acta 1350: 133–135,
1997.

[127988]

[127989]

[127990]11221.

[127991]Langkopf, A.; Hammarback, J. A.; Muller, R.; Vallee, R. B.;
Garner,

[127992]C. C.: Microtubule-associated proteins 1A and LC2: Two proteins encoded

[127993]in one messenger RNA. J. Biol. Chem. 267: 16561–16566, 1992.

[127994]

[127995]

[127996]11222.

[127997]Sun, X.-H.; Copeland, N. G.; Jenkins, N. A.; Baltimore, D.: Id

[127998]proteins Id1 and Id2 selectively inhibit DNA binding by one class

[127999]of helix-loop-helix proteins. Molec. Cell. Biol. 11: 5603–5611,

[128000]1991.

[128001]

[128002]

[128003]11223.

[128004]Kordeli, E.; Bennett, V.: Distinct ankyrin isoforms at neuron

[128005]cell bodies and nodes of Ranvier resolved using erythrocyte ankyrin-deficient

[128006]mice. J. Cell Biol. 114: 1243–1259, 1991.

[128007]

[128008]

[128009]11224.

[128010]Reche, P. A.; Soumelis, V.; Gorman, D. M.; Clifford, T.; Liu, M.;

[128011]Travis, M.; Zurawski, S. M.; Johnston, J.; Liu, Y.-J.; Spits, H.;

[128012]de Waal Malefyt, R.; Kastelein, R. A.; Bazan, J. F.: Human thymic

[128013]stromal lymphopoietin preferentially stimulates myeloid cells. J.

[128014]Immun. 167: 336–343, 2001.

[128015]

[128016]

[128017]11225.

[128018]Nagase, T.; Kikuno, R.; Nakayama, M.; Hirosawa, M.; Ohara, O.:

[128019]Prediction of the coding sequences of unidentified human genes. XVIII.

[128020]The complete sequences of 100 new cDNA clones from
brain which code

[128021]for large proteins in vitro. DNA Res. 7: 273–281, 2000.

[128022]

[128023]

[128024]11226.

[128025]Mukai, J.; Hachiya, T.; Shoji–Hoshino, S.; Kimura, M. T.;
Nadano,

[128026]D.; Suvanto, P.; Hanaoka, T.; Li, Y.; Irie, S.; Greene, L. A.;
Sato,

[128027]T.–A.: NADE, a p75NTR–associated cell death executor, is
involved

[128028]in signal transduction mediated by the common neu-
rotrophin receptor

[128029]p75NTR. J. Biol. Chem. 275: 17566–17570, 2000.

[128030]

[128031]

[128032]11227.

[128033]Rapp, G.; Freudenstein, J.; Klaudiny, J.; Mucha, J.; Wempe,
F.;

[128034]Zimmer, M.; Scheit, K. H.: Characterization of three abundant mRNAs

[128035]from human ovarian granulosa cells. DNA Cell Biol. 9: 479–485, 1990.

[128036]

[128037]

[128038]11228.

[128039]Scott, A. F.: Personal Communication. Baltimore, Md. 10/8/2001.

[128040]

[128041]

[128042]11229.

[128043]Kurochkin, I. V.; Yonemitsu, N.; Funahashi, S.; Nomura, H.: ALEX1,

[128044]a novel human armadillo repeat protein that is expressed differentially

[128045]in normal tissues and carcinomas. Biochem. Biophys. Res. Commun. 280:

[128046]340–347, 2001.

[128047]

[128048]

[128049]11230.

[128050]Nagase, T.; Ishikawa, K.; Miyajima, N.; Tanaka, A.; Kotani, H.;

[128051]Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified

[128052]human genes. IX. The complete sequences of 100 new cDNA clones from

[128053]brain which can code for large proteins in vitro. DNA Res. 5: 31–39,

[128054]1998.

[128055]

[128056]

[128057]11231.

[128058]Chuang, T.–H.; Ulevitch, R. J.: Cloning and characterization of

[128059]a sub-family of human Toll-like receptors: hTLR7, hTLR8 and hTLR9. Europ.

[128060]Cytokine Netw. 11: 372–378, 2000.

[128061]

[128062]

[128063]11232.

[128064]Du, X.; Poltorak, A.; Wei, Y.; Beutler, B.: Three novel mammalian

[128065]toll-like receptors: gene structure, expression, and evolution. Europ.

[128066]Cytokine Netw. 11: 362–371, 2000.

[128067]

[128068]

[128069]11233.

[128070]Kadowaki, N.; Ho, S.; Antonenko, S.; de Waal Malefyt, R.; Kastelein,

[128071]R. A.; Bazan, F.; Liu, Y.–J.: Subsets of human dendritic cell precursors

[128072]express different Toll-like receptors and respond to different microbial

[128073]antigens. J. Exp. Med. 194: 863–869, 2001.

[128074]

[128075]

[128076]11234.

[128077]Wirth, B.; Schmidt, T.; Hahnen, E.; Rudnik-Schoneborn, S.; Krawczak,

[128078]M.; Muller-Myhsok, B.; Schonling, J.; Zerres, K.: De novo rearrangements

[128079]found in 2% of index patients with spinal muscular atrophy: mutational

[128080]mechanisms, parental origin, mutation rate, and implications for genetic

[128081]counseling. Am. J. Hum. Genet. 61: 1102-1111, 1997.

[128082]

[128083]

[128084]11235.

[128085]Frattini, A.; Orchard, P. J.; Sobacchi, C.; Giliani, S.; Abinun,

[128086]M.; Mattsson, J. P.; Keeling, D. J.; Andersson, A.-K.; Wallbrandt,

[128087]P.; Zecca, L.; Notarangelo, L. D.; Vezzoni, P.; Villa, A.: Defects

[128088]in TCIRG1 subunit of the vacuolar proton pump are responsible for

[128089]a subset of human autosomal recessive osteopetrosis. Nature Genet. 25:

[128090]343–346, 2000.

[128091]

[128092]

[128093]11236.

[128094]Afzal, A. R.; Rajab, A.; Fenske, C. D.; Oldridge, M.; Elanko, N.;

[128095]Ternes–Pereira, E.; Tuysuz, B.; Murday, V. A.; Patton, M. A.; Wilkie,

[128096]A. O. M.; Jeffery, S.: Recessive Robinow syndrome, allelic to dominant

[128097]brachydactyly type B, is caused by mutation of ROR2. Nature Genet. 25:

[128098]419–422, 2000.

[128099]

[128100]

[128101]11237.

[128102]van Bokhoven, H.; Celli, J.; Kayserili, H.; van Beusekom, E.;

[128103]Balci, S.; Brussel, W.; Skovby, F.; Kerr, B.; Percin, E. F.; Akarsu,

[128104]N.; Brunner, H. G.: Mutation of the gene encoding the

ROR2 tyrosine

[128105]kinase causes autosomal recessive Robinow syndrome.

Nature Genet. 25:

[128106]423–426, 2000.

[128107]

[128108]

[128109]11238.

[128110]den Dunnen, J. T.; Grootsholten, P. M.; Bakker, E.; Blonden,

[128111]L. A. J.; Ginjaar, H. B.; Wapenaar, M. C.; van Paassen, H. M. B.;

[128112]van Broeckhoven, C.; Pearson, P. L.; van Ommen, G. J. B.:

Topography

[128113]of the Duchenne muscular dystrophy (DMD) gene: FIGE and cDNA analysis

[128114]of 194 cases reveals 115 deletions and 13 duplications.

Am. J. Hum.

[128115]Genet. 45: 835–847, 1989.

[128116]

[128117]

[128118]11239.

[128119] Claudio, J. O.; Liew, C.-C.; Dempsey, A. A.; Cukerman, E.;
Stewart,

[128120] A. K.; Na, E.; Atkins, H. L.; Iscove, N. N.; Hawley, R. G.:
Identification

[128121] of sequence-tagged transcripts differentially expressed
within the

[128122] human hematopoietic hierarchy. Genomics 50: 44–52,
1998.

[128123]

[128124]

[128125] 11240.

[128126] Claudio, J. O.; Liew, C.-C.; Ma, J.; Heng, H. H. Q.; Stewart,
A.

[128127] K.; Hawley, R. G.: Cloning and expression analysis of a
novel WD

[128128] repeat gene, WDR3, mapping to 1p12–p13. Genomics 59:
85–89, 1999.

[128129]

[128130]

[128131] 11241.

[128132] Eng, B. H.; Guerinot, M. L.; Eide, D.; Saier, M. H., Jr.: Se-

quence

[128133]analyses and phylogenetic characterization of the ZIP
family of metal

[128134]ion transport proteins. J. Membr. Biol. 166: 1–7, 1998.

[128135]

[128136]

[128137]11242.

[128138]Lioumi, M.; Ferguson, C. A.; Sharpe, P. T.; Freeman, T.;
Marenholz,

[128139]I.; Mischke, D.; Heizmann, C.; Ragoussis, J.: Isolation and
characterization

[128140]of human and mouse ZIRTL, a member of the IRT1 family
of transporters,

[128141]mapping within the epidermal differentiation complex.
Genomics 62:

[128142]272–280, 1999.

[128143]

[128144]

[128145]11243.

[128146]Lioumi, M.; Olavesen, M. G.; Nizetic, D.; Ragoussis, J.:

High-resolution

[128147]YAC fragmentation map of 1q21. Genomics 49: 200–208, 1998.

[128148]

[128149]

[128150]11244.

[128151]Leiper, J. M.; Santa Maria, J.; Chubb, A.; MacAllister, R. J.;

[128152]Charles, I. G.; Whitley, G. S.; Vallance, P.: Identification of two

[128153]human dimethylarginine dimethylaminohydrolases with distinct tissue

[128154]distributions and homology with microbial arginine deiminases. Biochem.

[128155]J. 343: 209–214, 1999.

[128156]

[128157]

[128158]11245.

[128159]Tran, C. T. L.; Fox, M. F.; Vallance, P.; Leiper, J. M.: Chromosomal

[128160]localization, gene structure, and expression pattern of

DDAH1: comparison

[128161]with DDAH2 and implications for evolutionary origins. Genomics 68:

[128162]101–105, 2000.

[128163]

[128164]

[128165]11246.

[128166]Vanhalst, K.; Kools, P.; Eynde, E. V.; van Roy, F.: The human

[128167]and murine protocadherin-beta one-exon gene families show high evolutionary

[128168]conservation, despite the difference in gene number. FEBS Lett. 495:

[128169]120–125, 2001.

[128170]

[128171]

[128172]11247.

[128173]Wu, Q.; Maniatis, T.: A striking organization of a large family

[128174]of human neural cadherin like cell adhesion genes. Cell

97: 779–790,

[128175]1999.

[128176]

[128177]

[128178]11248.

[128179]Katayama, H.; Ota, T.; Morita, K.; Terada, Y.; Suzuki, F.;
Katoh,

[128180]O.; Tatsuka, M.: Human AIM–1: cDNA cloning and reduced
expression

[128181]during endomitosis in megakaryocyte–lineage cells. Gene
224: 1–7,

[128182]1998.

[128183]

[128184]

[128185]11249.

[128186]Kimura, M.; Matsuda, Y.; Yoshioka, T.; Sumi, N.; Okano,
Y.: Identification

[128187]and characterization of STK12/Aik2: a human gene re–
lated to aurora

[128188]of Drosophila and yeast IPL1. Cytogenet. Cell Genet. 82:

147–152,

[128189]1998.

[128190]

[128191]

[128192]11250.

[128193]Tatsuka, M.; Katayama, H.; Ota, T.; Tanaka, T.; Odashima, S.; Suzuki,

[128194]F.; Terada, Y.: Multinuclearity and increased ploidy caused by overexpression

[128195]of the aurora- and lpl1-like midbody-associated protein mitotic kinase

[128196]in human cancer cells. Cancer Res. 58: 4811–4816, 1998.

[128197]

[128198]

[128199]11251.

[128200]Chen, C. M.; Kraut, N.; Groudine, M.; Weintraub, H.: I-mf, a novel

[128201]myogenic repressor, interacts with members of the MyoD family. Cell 86:

[128202]731–741, 1996.

[128203]

[128204]

[128205]11252.

[128206]Kraut, N.: The gene encoding I-mf (Mdfi) maps to human chromosome

[128207]6p21 and mouse chromosome 17. Mammalian Genome 8: 618-619, 1997.

[128208]

[128209]

[128210]11253.

[128211]Kraut, N.; Snider, L.; Chen, C.-M. A.; Tapscott, S. J.; Grou-dine,

[128212]M.: Requirement of the mouse I-mfa gene for placental development

[128213]and skeletal patterning. EMBO J. 17: 6276-6288, 1998.

[128214]

[128215]

[128216]11254.

[128217]Boynton, S.; Tully, T.: Latheo, a new gene involved in as-

sociative

[128218]learning and memory in *Drosophila melanogaster*, identified from P

[128219]element mutagenesis. *Genetics* 131: 655–672, 1992.

[128220]

[128221]

[128222]11255.

[128223]Pinto, S.; Quintana, D. G.; Smith, P.; Mihalek, R. M.; Hou, Z.-H.;

[128224]Boynton, S.; Jones, C. J.; Hendricks, M.; Velinzon, K.; Wohlschlegel,

[128225]J. A.; Austin, R. J.; Lane, W. S.; Tully, T.; Dutta, A.: Latheo encodes

[128226]a subunit of the origin recognition complex and disrupts neuronal

[128227]proliferation and adult olfactory memory when mutant. Neuron 23:

[128228]45–54, 1999.

[128229]

[128230]

[128231]11256.

[128232]Springer, J.; Nanda, I.; Hoehn, K.; Schmid, M.; Grummt, F.:
Identification

[128233]and chromosomal localization of murine ORC3, a new
member of the mouse

[128234]origin recognition complex. Cytogenet. Cell Genet. 87:
245–251,

[128235]1999.

[128236]

[128237]

[128238]11257.

[128239]Tugal, T.; Zou–Yang, X. H.; Gavin, K.; Pappin, D.; Canas,
B.; Kobayashi,

[128240]R.; Hunt, T.; Stillman, B.: The Orc4p and Orc5p subunits of
the Xenopus

[128241]and human origin recognition complex are related to
Orc1p and Cdc6p. J.

[128242]Biol. Chem. 273: 32421–32429, 1998.

[128243]

[128244]

[128245]11258.

[128246]Butcher, S.; Arney, K. L.; Cook, G. P.: MAFA-L, an ITIM-containing

[128247]receptor encoded by the human NK cell gene complex and expressed by

[128248]basophils and NK cells. *Europ. J. Immun.* 28: 3755–3762, 1998.

[128249]

[128250]

[128251]11259.

[128252]Hanke, T.; Corral, L.; Vance, R. E.; Raulet, D. H.: 2F1 antigen,

[128253]the mouse homolog of the rat 'mast cell function-associated antigen',

[128254]is a lectin-like type II transmembrane receptor expressed by natural

[128255]killer cells. *Europ. J. Immun.* 28: 4409–4417, 1998.

[128256]

[128257]

[128258]11260.

[128259]Lamers, M. B. A. C.; Lamont, A. G.; Williams, D. H.: Human MAFA

[128260]has alternatively spliced variants. Biochim. Biophys. Acta
1399:

[128261]209–212, 1998.

[128262]

[128263]

[128264]11261.

[128265]Voehringer, D.; Kaufmann, M.; Pircher, H.: Genomic structure,
ture,

[128266]alternative splicing, and physical mapping of the killer cell
lectin-like

[128267]receptor G1 gene (KLRG1), the mouse homologue of
MAFA. Immunogenetics 52:

[128268]206–211, 2001.

[128269]

[128270]

[128271]11262.

[128272]Bahler, M.: Myosins on the move to signal transduction.
Curr.

[128273]Opin. Cell Biol. 8: 18–22, 1996.

[128274]

[128275]

[128276]11263.

[128277]Gorman, S. W.; Haider, N. B.; Grieshammer, U.; Swiderski, R. E.;

[128278]Kim E; Welch, J. W.; Searby, C.; Leng, S.; Carmi, R.; Sheffield, V.

[128279]C.; Duhl, D. M.: The cloning and developmental expression of unconventional

[128280]myosin IXA (MYO9A) a gene in the Bardet–Biedl syndrome (BBS4) region

[128281]at chromosome 15q22–q23. Genomics 59: 150–160, 1999.

[128282]

[128283]

[128284]11264.

[128285]Dunham, I.; Shimizu, N.; Roe, B. A.; Chissoe, S.; Hunt, A. R.;

[128286]Collins, J. E.; Bruskiewich, R.; Beare, D. M.; Clamp, M.; Smink, L.

[128287]J.; Ainscough, R.; Almeida, J. P.; and 205 others: The DNA sequence

[128288]of human chromosome 22. Nature 402: 489–495, 1999.

[128289]

[128290]

[128291]11265.

[128292]Kimura, T.; Ivell, R.; Rust, W.; Mizumoto, Y.; Ogita, K.;
Kusui,

[128293]C.; Matsumura, Y.; Azuma, C.; Murata, Y.: Molecular
cloning of a

[128294]human MafF homologue, which specifically binds to the
oxytocin receptor

[128295]gene in term myometrium. Biochem. Biophys. Res. Com-
mun. 264: 86–92,

[128296]1999.

[128297]

[128298]

[128299]11266.

[128300]Hiki, K.; D'Andrea, R. J.; Furze, J.; Crawford, J.; Woollatt, E.;

[128301]Sutherland, G. R.; Vadas, M. A.; Gamble, J. R.: Cloning,
characterization,

[128302]and chromosomal location of a novel human K(+)-Cl(-)

cotransporter. J.

[128303]Biol. Chem. 274: 10661–10667, 1999.

[128304]

[128305]

[128306]11267.

[128307]Mount, D. B.: Personal Communication. Nashville, Tenn.
3/28/2000.

[128308]

[128309]

[128310]11268.

[128311]Mount, D. B.; Mercado, A.; Song, L.; Xu, J.; George, A. L.,
Jr.;

[128312]Delpire, E.; Gamba, G.: Cloning and characterization of
KCC3 and

[128313]KCC4, new members of the cation–chloride cotransporter
gene family. J.

[128314]Biol. Chem. 274: 16355–16362, 1999.

[128315]

[128316]

[128317]11269.

[128318]Herzing, L. B. K.; Kim, S.-J.; Cook, E. H., Jr.; Ledbetter, D.

[128319]H.: The human aminophospholipid-transporting ATPase
gene ATP10C maps

[128320]adjacent to UBE3A and exhibits similar imprinted expres-
sion. Am.

[128321]J. Hum. Genet. 68: 1501-1505, 2001.

[128322]

[128323]

[128324]11270.

[128325]Dode, C.; Le Du, N.; Cuisset, L.; Letourneur, F.; Berthelot,
J.-M.;

[128326]Vaudour, G.; Meyrier, A.; Watts, R. A.; Scott, D. G. I.;
Nicholls,

[128327]A.; Granel, B.; Frances, C.; Garcier, F.; Edery, P.;
Boulinguez, S.;

[128328]Domergues, J.-P.; Delpech, M.; Grateau, G.: New muta-
tions of CIAS1

[128329]that are responsible for Muckle-Wells syndrome and fa-
milial cold urticaria:

[128330]a novel mutation underlies both syndromes. Am. J. Hum.

Genet. 70:

[128331]1498–1506, 2002.

[128332]

[128333]

[128334]11271.

[128335]Feldmann, J.; Prieur, A.–M.; Quartier, P.; Berquin, P.; Certain,
tain,

[128336]S.; Cortis, E.; Teillac–Hamel, D.; Fischer, A.; de Saint
Basile, G.

[128337]: Chronic infantile neurological cutaneous and articular
syndrome

[128338]is caused by mutations in CIAS1, a gene highly expressed
in polymorphonuclear

[128339]cells and chondrocytes. Am. J. Hum. Genet. 71: 198–203,
2002.

[128340]

[128341]

[128342]11272.

[128343]Greeve, I.; Hermans–Borgmeyer, I.; Brellinger, C.; Kasper,
D.;

[128344]Gomez–Isla, T.; Behl, C.; Levkau, B.; Nitsch, R. M.: The hu–

man DIMINUTO/DWARF1

[128345]homolog seladin-1 confers resistance to Alzheimer's disease-associated

[128346]neurodegeneration and oxidative stress. J. Neurosci. 20: 7345-7352,

[128347]2000.

[128348]

[128349]

[128350]11273.

[128351]Sarkar, D.; Imai, T.; Kambe, F.; Shibata, A.; Ohmori, S.; Siddiq,

[128352]A.; Hayasaka, S.; Funahashi, H.; Seo, H.: The human homolog of Diminuto/Dwarf1

[128353]gene (hDiminuto): a novel ACTH-responsive gene overexpressed in benign

[128354]cortisol-producing adrenocortical adenomas. J. Clin. Endocr. Metab. 86:

[128355]5130-5137, 2001.

[128356]

[128357]

[128358]11274.

[128359]Hattori, M.; Fujiyama, A.; Taylor, T. D.; Watanabe, H.;
Yada, T.;

[128360]Park, H.-S.; Toyoda, A.; Ishii, K.; Totoki, Y.; Choi, D.-K.;
Groner,

[128361]Y.; Soeda, E.; and 52 others: The DNA sequence of human
chromosome

[128362]21. Nature 405: 311–319, 2000. Note: Erratum: Nature:
407: 110 only,

[128363]2000.

[128364]

[128365]

[128366]11275.

[128367]Wilcox, E. R.; Burton, Q. L.; Naz, S.; Riazuddin, S.; Smith,
T.

[128368]N.; Ploplis, B.; Belyatseva, I.; Ben-Yosef, T.; Liburd, N. A.;
Morell,

[128369]R. J.; Kachar, B.; Wu, D. K.; Griffith, A. J.; Riazuddin, S.;
Friedman,

[128370]T. B.: Mutations in the gene encoding tight junction
claudin-14 cause

[128371]recessive deafness DFNB29. Cell 104: 165–172, 2001.

[128372]

[128373]

[128374]11276.

[128375]Dunlevy, J. R.; Berryhill, B. L.; Vergnes, J.-P.; SundarRaj, N.;

[128376]Hassell, J. R.: Cloning, chromosomal localization, and
characterization

[128377]of cDNA from a novel gene, SH3BP4, expressed by human
corneal fibroblasts. Genomics 62:

[128378]519–524, 1999.

[128379]

[128380]

[128381]11277.

[128382]Wong, W. T.; Schumacher, C.; Salcini, A. E.; Romano, A.;
Castagnino,

[128383]P.; Pelicci, P. G.; DiFiore, P. P.: A protein–binding domain,
EH,

[128384]identified in the receptor tyrosine kinase substrate Eps15
and conserved

[128385]in evolution. Proc. Nat. Acad. Sci. 92: 9530–9534, 1995.

[128386]

[128387]

[128388]11278.

[128389]Yamadori, T.; Baba, Y.; Matsushita, M.; Hashimoto, S.;
Kurosaki,

[128390]M.; Kurosaki, T.; Kishimoto, T.; Tsukada, S.: Bruton's tyro-
sine kinase

[128391]activity is negatively regulated by Sab, the Btk-SH3 do-
main-binding

[128392]protein. Proc. Nat. Acad. Sci. 96: 6341-6346, 1999.

[128393]

[128394]

[128395]11279.

[128396]Engqvist-Goldstein, A. E. Y.; Kessels, M. M.; Chopra, V. S.;
Hayden,

[128397]M. R.; Drubin, D. G.: An actin-binding protein of the
Sla2/Huntingtin

[128398]interacting protein 1 family is a novel component of
clathrin-coated

[128399]pits and vesicles. J. Cell Biol. 147: 1503-1518, 1999.

[128400]

[128401]

[128402]11280.

[128403]Baba, Y.; Matsushita, M.; Matsuda, Y.; Inazawa, J.; Yamadori, T.;

[128404]Hashimoto, S.; Kishimoto, T.; Tsukada, S.: Assignment of SH3BP5/Sh3bp5

[128405]encoding Sab, an SH3 domain-binding protein which preferentially associates

[128406]with Bruton's tyrosine kinase, to human chromosome 1q43 and mouse

[128407]chromosome 14B by in situ hybridization. Cytogenet. Cell Genet. 87:

[128408]221-222, 1999.

[128409]

[128410]

[128411]11281.

[128412]Matsushita, M.; Yamadori, T.; Kato, S.; Takemoto, Y.; Inazawa,

[128413]J.; Baba, Y.; Hashimoto, S.; Sekine, S.; Arai, S.; Kunikata, T.; Kurimoto,

[128414]M.; Kishimoto, T.; Tsukada, S.: Identification and characterization

[128415]of a novel SH3-domain binding protein, Sab, which preferentially associates

[128416]with Bruton's tyrosine kinase (Btk). Biochem. Biophys. Res. Commun. 245:

[128417]337–343, 1998.

[128418]

[128419]

[128420]11282.

[128421]Itoh, T.; Koshiba, S.; Kigawa, T.; Kikuchi, A.; Yokoyama, S.; Takenawa,

[128422]T.: Role of the ENTH domain in phosphatidylinositol-4,5-bisphosphate

[128423]binding and endocytosis. Science 291: 1047–1051, 2001.

[128424]

[128425]

[128426]11283.

[128427]Seki, N.; Muramatsu, M.; Sugano, S.; Suzuki, Y.; Nakagawara, A.;

[128428]Ohhira, M.; Hayashi, A.; Hori, T.; Saito, T.: Cloning, expression

[128429]analysis, and chromosomal localization of HIP1R, an isolog of huntingtin

[128430]interacting protein (HIP1). J. Hum. Genet. 43: 268–271, 1998.

[128431]

[128432]

[128433]11284.

[128434]Marton, M. J.; Vazquez de Aldana, C. R.; Qiu, H.; Chakraborty,

[128435]K.; Hinnebusch, A. G.: Evidence that GCN1 and GCN20, translational

[128436]regulators of GCN4, function on elongating ribosomes in activation

[128437]of eIF2- α kinase GCN2. Molec. Cell. Biol. 17: 4474–4489, 1997.

[128438]

[128439]

[128440]11285.

[128441]Nash, S. R.; Giros, B.; Kingsmore, S. F.; Kim, K. M.; El-Mestikawy,

[128442]S.; Dong, Q.; Fumagalli, F.; Seldin, M. F.; Caron, M. G.: Cloning,

[128443]gene structure and genomic localization of an orphan transporter from

[128444]mouse kidney with six alternatively-spliced isoforms. Receptors Channels 6:

[128445]113–128, 1998.

[128446]

[128447]

[128448]11286.

[128449]Scott, A. F.: Personal Communication. Baltimore, Md. 2/5/2001.

[128450]

[128451]

[128452]11287.

[128453]Meguro, M.; Kashiwagi, A.; Mitsuya, K.; Nakao, M.; Kondo, I.; Saitoh,

[128454]S.; Oshimura, M.: A novel maternally expressed gene,
ATP10C, encodes

[128455]a putative aminophospholipid translocase associated with
Angelman

[128456]syndrome. Nature Genet. 28: 19–20, 2001.

[128457]

[128458]

[128459]11288.

[128460]Nakakura, E. K.; Watkins, D. N.; Schuebel, K. E.; Sriuran-
pong,

[128461]V.; Borges, M. W.; Nelkin, B. D.; Ball, D. W.: Mammalian
Scratch:

[128462]a neural-specific Snail family transcriptional repressor.
Proc. Nat.

[128463]Acad. Sci. 98: 4010–4015, 2001.

[128464]

[128465]

[128466]11289.

[128467]Scott, A. F.: Personal Communication. Baltimore, Md.
6/21/2001.

[128468]

[128469]

[128470]11290.

[128471]Hoatlin, M. E.; Zhi, Y.; Ball, H.; Silvery, K.; Melnick, A.;
Stone,

[128472]S.; Arai, S.; Hawe, N.; Owen, G.; Zelent, A.; Licht, J. D.: A
novel

[128473]BTB/POZ transcriptional repressor protein interacts with
the Fanconi

[128474]anemia group C protein and PLZF. Blood 94: 3737–3747,
1999.

[128475]

[128476]

[128477]11291.

[128478]Lin, W.; Lai, C.-H.; Tang, C.-J. C.; Huang, C.-J.; Tang, T. K.

[128479]: Identification and gene structure of a novel human PLZF-
related

[128480]transcription factor gene, TZFP. Biochem. Biophys. Res.
Commun. 264:

[128481]789–795, 1999.

[128482]

[128483]

[128484]11292.

[128485]Yokoyama–Kobayashi, M.; Sugano, S.; Kato, T.; Kato, S.: A
signal

[128486]sequence detection system using secreted protease activ-
ity as an indicator. Gene 163:

[128487]193–196, 1995.

[128488]

[128489]

[128490]11293.

[128491]Yokoyama–Kobayashi, M.; Yamaguchi, T.; Sekine, S.; Kato,
S.: Selection

[128492]of cDNAs encoding putative type II membrane proteins on
the cell surface

[128493]from a human full-length cDNA bank. Gene 228:
161–167, 1999.

[128494]

[128495]

[128496]11294.

[128497]Olivier, E.; Soury, E.; Risler, J. L.; Smith, F.; Schneider, K.;

[128498]Lochner, K.; Jouzeau, J. Y.; Fey, G. H.; Salier, J. P.: A novel
set

[128499]of hepatic mRNAs preferentially expressed during an
acute inflammation

[128500]in rat represents mostly intracellular proteins. Genomics
57: 352–364,

[128501]1999.

[128502]

[128503]

[128504]11295.

[128505]Olivier, E.; Soury, E.; Ruminy, P.; Husson, A.; Parmentier,
F.;

[128506]Daveau, M.; Salier, J.–P.: Fetuin–B, a second member of
the fetuin

[128507]family in mammals. Biochem. J. 350: 589–597, 2000.

[128508]

[128509]

[128510]11296.

[128511]Hampe, J.; Grebe, J.; Nikolaus, S.; Solberg, C.; Croucher, P.

J.

[128512]P.; Mascheretti, S.; Jahnsen, J.; Moum, B.; Klump, B.;
Krawczak, M.;

[128513]Mirza, M. M.; Foelsch, U. R.; Vatn, M.; Schreiber, S.: Asso-
ciation

[128514]of NOD2 (CARD 15) genotype with clinical course of
Crohn's disease:

[128515]a cohort study. Lancet 359: 1661–1665, 2002.

[128516]

[128517]

[128518]11297.

[128519]Hugot, J.–P.; Chamaillard, M.; Zouali, H.; Lesage, S.;
Cezard,

[128520]J.–P.; Belaiche, J.; Almer, S.; Tysk, C.; O'Morain, C. A.; Gas-
sull,

[128521]M.; Binder, V.; Finkel, Y.; and 8 others: Association of
NOD2 leucine-rich

[128522]repeat variants with susceptibility to Crohn's disease. Na-
ture 411:

[128523]599–603, 2001.

[128524]

[128525]

[128526]11298.

[128527]Murillo, L.; Crusius, J. B. A.; van Bodegraven, A. A.; Al-izadeh,

[128528]B. Z.; Pena, A. S.: CARD15 gene and the classification of Crohn's

[128529]disease. Immunogenetics 54: 59–61, 2002.

[128530]

[128531]

[128532]11299.

[128533]Ogura, Y.; Bonen, D. K.; Inohara, N.; Nicolae, D. L.; Chen, F.

[128534]F.; Ramos, R.; Britton, H.; Moran, T.; Karaliuskas, R.; Duerr, R.

[128535]H.; Achkar, J.–P.; Brant, S. R.; Bayless, T. M.; Kirschner, B. S.;

[128536]Hanauer, S. B.; Nunez, G.; Cho, J. H.: A frameshift mutation in Nod2

[128537]associated with susceptibility to Crohn's disease. Nature 411: 603–606,

[128538]2001.

[128539]

[128540]

[128541]11300.

[128542]Ogura, Y.; Inohara, N.; Benito, A.; Chen, F. F.; Yamaoka, S.;
Nunez,

[128543]G.: Nod2, a Nod1/Apaf-1 family member that is restricted
to monocytes

[128544]and activates NF-kappa-B. J. Biol. Chem. 276: 4812-4818,
2001.

[128545]

[128546]

[128547]11301.

[128548]Lesage, S.; Zouali, H.; Cezard, J.-P.; EPWG-IBD Group;
Colombel,

[128549]J.-F.; EPIMAD Group; Belaiche, J.; GETAID Group; Almer, S.;
Tysk,

[128550]C.; O'Morain, C.; Gassull, M.; Binder, V.; Finkel, Y.;
Modigliani,

[128551]R.; Gower-Rousseau, C.; Macry, J.; Merlin, F.; Chamaillard,
M.; Jannot,

[128552]A.-S.; Thomas, G.; Hugot, J.-P.: CARD15/NOD2 mutational

analysis

[128553]and genotype–phenotype correlation in 612 patients with
inflammatory

[128554]bowel disease. Am. J. Hum. Genet. 70: 845–857, 2002.

[128555]

[128556]

[128557]11302.

[128558]van Heel, D. A.; McGovern, D. P. B.; Cardon, L. R.;
Dechairo, B.

[128559]M.; Lench, N. J.; Carey, A. H.; Jewell, D. P.: Fine mapping of
the

[128560]IBD1 locus did not identify Crohn disease–associated
NOD2 variants:

[128561]implications for complex disease genetics. Am. J. Med.
Genet. 111:

[128562]253–259, 2002.

[128563]

[128564]

[128565]11303.

[128566]Vermeire, S.; Wild, G.; Kocher, K.; Cousineau, J.; Dufresne,

L.;

[128567]Bitton, A.; Langelier, D.; Pare, P.; Lapointe, G.; Cohen, A.;
Daly,

[128568]M. J.; Rioux, J. D.: CARD15 genetic variation in a Quebec
population:

[128569]prevalence, genotype–phenotype relationship, and haplo-
type structure. Am.

[128570]J. Hum. Genet. 71: 74–83, 2002.

[128571]

[128572]

[128573]11304.

[128574]Yamazaki, K.; Takazoe, M.; Tanaka, T.; Kazumori, T.;
Nakamura,

[128575]Y.: Absence of mutation in the NOD2/CARD15 gene
among 483 Japanese

[128576]patients with Crohn's disease. J. Hum. Genet. 47:
469–472, 2002.

[128577]

[128578]

[128579]11305.

[128580]Fliss, M. S.; Hinkle, P. M.; Bancroft, C.: Expression cloning

[128581]and characterization of PREB (prolactin regulatory element binding),

[128582]a novel WD motif DNA-binding protein with a capacity to regulate prolactin

[128583]promoter activity. Molec. Endocr. 13: 644–657, 1999.

[128584]

[128585]

[128586]11306.

[128587]Taylor Clelland, C. L.; Levy, B.; McKie, J. M.; Duncan, A. M. V.;

[128588]Hirschhorn, K.; Bancroft, C.: Cloning and characterization of human

[128589]PREB; a gene that maps to a genomic region associated with trisomy

[128590]2p syndrome. Mammalian Genome 11: 675–681, 2000.

[128591]

[128592]

[128593]11307.

[128594]Li, B.; Oestreich, S.; de Lange, T.: Identification of human Rap1:

[128595]implications for telomere evolution. Cell 101: 471–483,
2000.

[128596]

[128597]

[128598]11308.

[128599]Lieb, J. D.; Liu, X.; Botstein, D.; Brown, P. O.: Promoter-spe-
cific

[128600]binding of Rap1 revealed by genome-wide maps of pro-
tein–DNA association. Nature

[128601]Genet. 28: 327–334, 2001.

[128602]

[128603]

[128604]11309.

[128605]Honore, B.; Leffers, H.; Madsen, P.; Rasmussen, H. H.;
Vandekerckhove,

[128606]J.; Celis, J. E.: Molecular cloning and expression of a trans-
formation-sensitive

[128607]human protein containing the TPR motif and sharing iden-
tity to the

[128608]stress-inducible yeast protein STI1. J. Biol. Chem. 267:
8485–8491,

[128609]1992.

[128610]

[128611]

[128612]11310.

[128613]Scheufler, C.; Brinker, A.; Bourenkov, G.; Pegoraro, S.; Mo-
roder,

[128614]L.; Bartunik, H.; Hartl, F. U.; Moarefi, I.: Structure of TPR
domain-peptide

[128615]complexes: critical elements in the assembly of the
Hsp70-Hsp90 multichaperone

[128616]machine. Cell 101: 199-210, 2000.

[128617]

[128618]

[128619]11311.

[128620]Chong, S. S.; Tanigami, A.; Roschke, A. V.; Ledbetter, D.
H.:

[128621]14-3-3-epsilon has no homology to LIS1 and lies telom-
eric to it on

[128622]chromosome 17p13.3 outside the Miller-Dieker syndrome
chromosome region. Genome

[128623]Res. 6: 735–741, 1996.

[128624]

[128625]

[128626]11312.

[128627]Conklin, D. S.; Galaktionov, K.; Beach, D.: 14–3–3 proteins
associate

[128628]with cdc25 phosphatases. Proc. Nat. Acad. Sci. 92:
7892–7896, 1995.

[128629]

[128630]

[128631]11313.

[128632]Jin, D.–Y.; Lyu, M. S.; Kozak, C. A.; Jeang, K.–T.: Function
of

[128633]14–3–3 proteins. Nature 382: 308 only, 1996.

[128634]

[128635]

[128636]11314.

[128637]Luk, S. C. W.; Garcia-Barcelo, M.; Tsui, S. K. W.; Fung, K.
P.;

[128638]Lee, C. Y.; Waye, M. M. Y.: Assignment of the human
14-3-3 epsilon

[128639]isoform (YWHAE) to human chromosome 17p13 by in situ
hybridization. Cytogenet.

[128640]Cell Genet. 78: 105-106, 1997.

[128641]

[128642]

[128643]11315.

[128644]Slentz-Kesler, K.; Moore, J. T.; Lombard, M.; Zhang, J.;
Hollingsworth,

[128645]R.; Weiner, M. P.: Identification of the human Mnk2 gene
(MKNK2)

[128646]through protein interaction with estrogen receptor beta.
Genomics 69:

[128647]63-71, 2000.

[128648]

[128649]

[128650]11316.

[128651]Li, Y.; He, X.; Schembri-King, J.; Jakes, S.; Hayashi, J.:
Cloning

[128652]and characterization of human Lnk, an adaptor protein
with pleckstrin

[128653]homology and Src homology 2 domains that can inhibit T
cell activation. J.

[128654]Immun. 164: 5199–5206, 2000.

[128655]

[128656]

[128657]11317.

[128658]Takaki, S.; Sauer, K.; Iritani, B. M.; Chien, S.; Ebihara, Y.;

[128659]Tsuji, K.; Takatsu, K.; Perlmutter, R. M.: Control of B cell
production

[128660]by the adaptor protein Lnk: definition of a conserved fam-
ily of signal-modulating

[128661]proteins. Immunity 13: 599–609, 2000.

[128662]

[128663]

[128664]11318.

[128665]Fiscella, M.; Zhang, H.; Fan, S.; Sakaguchi, K.; Shen, S.;
Mercer,

[128666]W. E.; Vande Woude, G. F.; O'Connor, P. M.; Appella, E.:

Wip1, a

[128667]novel human protein phosphatase that is induced in response to ionizing

[128668]radiation in a p53-dependent manner. Proc. Nat. Acad. Sci. 94: 6048–6053,

[128669]1997.

[128670]

[128671]

[128672]11319.

[128673]Li, J.; Yang, Y.; Peng, Y.; Austin, R. J.; van Eyndhoven, W. G.;

[128674]Nguyen, K. C. Q.; Gabriele, T.; McCurrach, M. E.; Marks, J. R.; Hoey,

[128675]T.; Lowe, S. W.; Powers, S.: Oncogenic properties of PPM1D located

[128676]within a breast cancer amplification epicenter at 17q23. Nature Genet. 31:

[128677]133–134, 2002.

[128678]

[128679]

[128680]11320.

[128681]Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayasi, Y.;

[128682]1. Fiscella, M.; Zhang, H.; Fan, S.; Sakaguchi, K.; Shen, S.; Mercer,

[128683]W. E.; Vande Woude, G. F.; O'Connor, P. M.; Appella, E.: Wip1, a

[128684]novel human protein phosphatase that is induced in response to ionizing

[128685]radiation in a p53-dependent manner. Proc. Nat. Acad. Sci. 94: 6048–6053,

[128686]1997.

[128687]

[128688]

[128689]11321.

[128690]Stover, C. M.; Schwaeble, W. J.; Lynch, N. J.; Thiel, S.; Speicher,

[128691]M. R.: Assignment of the gene encoding mannan-binding lectin-associated

[128692]serine protease 2 (MASP2) to human chromosome 1p36.3–p36.2 by in situ

[128693]hybridization and somatic cell hybrid analysis. Cytogenet. Cell Genet. 84:

[128694]148–149, 1999.

[128695]

[128696]

[128697]11322.

[128698]Stover, C. M.; Thiel, S.; Thelen, M.; Lynch, N. J.; Vorup-Jensen,

[128699]T.; Jensenius, J. C.; Schwaebble, W. J.: Two constituents of the initiation

[128700]complex of the mannan-binding lectin activation pathway of complement

[128701]are encoded by a single structural gene. J. Immun. 162: 3481–3490,

[128702]1999.

[128703]

[128704]

[128705]11323.

[128706]Takahashi, M.; Endo, Y.; Fujita, T.; Matsushita, M.: A truncated

[128707]form of mannose-binding lectin-associated serine protease (MASP)–2

[128708]expressed by alternative polyadenylation is a component
of the lectin

[128709]complement pathway. *Int. Immun.* 11: 859–863, 1999.

[128710]

[128711]

[128712]11324.

[128713]Thiel, S.; Vorup-Jensen, T.; Stover, C. M.; Schwaebble, W.;
Laursen,

[128714]S. B.; Poulsen, K.; Willis, A. C.; Eggleton, P.; Hansen, S.;
Holmskov,

[128715]U.; Reid, K. B. M.; Jensenius, J. C.: A second serine pro-
tease associated

[128716]with mannan-binding lectin that activates complement.
Nature 386:

[128717]506–510, 1997.

[128718]

[128719]

[128720]11325.

[128721]Ferguson, K. M.; Kavran, J. M.; Sankaran, V. G.; Fournier,
E.;

[128722]Isakoff, S. J.; Skolnik, E. Y.; Lemmon, M. A.: Structural basis for

[128723]discrimination of 3-phosphoinositides by pleckstrin homology domains. Molec.

[128724]Cell 6: 373–384, 2000.

[128725]

[128726]

[128727]11326.

[128728]Franco, M.; Boretto, J.; Robineau, S.; Monier, S.; Goud, B.; Chardin,

[128729]P.; Chavrier, P.: ARNO3, a Sec7-domain guanine nucleotide exchange

[128730]factor for ADP ribosylation factor 1, is involved in the control of

[128731]Golgi structure and function. Proc. Nat. Acad. Sci. 95: 9926–9931,

[128732]1998.

[128733]

[128734]

[128735]11327.

[128736]Klarlund, J. K.; Guilherme, A.; Holik, J. J.; Virbasius, J. V.;

[128737]Chawla, A.; Czech, M. P.: Signaling by phosphoinositide-
3,4,5-triphosphate

[128738]through proteins containing pleckstrin and Sec7 homol-
ogy domains. Science 275:

[128739]1927-1930, 1997.

[128740]

[128741]

[128742]11328.

[128743]Drewes, T.; Senkel, S.; Holewa, B.; Ryffel, G. U.: Human
hepatocyte

[128744]nuclear factor 4 isoforms are encoded by distinct and dif-
ferentially

[128745]expressed genes. Molec. Cell. Biol. 16: 925-931, 1996.

[128746]

[128747]

[128748]11329.

[128749]Taraviras, S.; Mantamadiotis, T.; Dong-Si, T.; Mincheva,
A.; Lichter,

[128750]P.; Drewes, T.; Ryffel, G. U.; Monaghan, A. P.; Schutz, G.:

Primary

[128751]structure, chromosomal mapping, expression and transcriptional activity

[128752]of murine hepatocyte nuclear factor 4-gamma. Biochim. Biophys. Acta 1490:

[128753]21–32, 2000.

[128754]

[128755]

[128756]11330.

[128757]Buck, J.; Sinclair, M. L.; Schapal, L.; Cann, M. J.; Levin, L.

[128758]R.: Cytosolic adenylyl cyclase defines a unique signaling molecule

[128759]in mammals. Proc. Nat. Acad. Sci. 96: 79–84, 1999.

[128760]

[128761]

[128762]11331.

[128763]Chen, Y.; Cann, M. J.; Litvin, T. N.; Iourgenko, V.; Sinclair,

[128764]M. L.; Levin, L. R.; Buck, J.: Soluble adenylyl cyclase as an evolutionarily

[128765]conserved bicarbonate sensor. Science 289: 625–628,
2000.

[128766]

[128767]

[128768]11332.

[128769]Reed, B. Y.; Gitomer, W. L.; Heller, H. J.; Hsu, M. C.; Lemke,

[128770]M.; Padalino, P.; Pak, C. Y. C.: Identification and character-
ization

[128771]of a gene with base substitutions associated with the ab-
sorptive hypercalciuria

[128772]phenotype and low spinal bone density. J. Clin. Endocr.
Metab. 87:

[128773]1476–1485, 2002.

[128774]

[128775]

[128776]11333.

[128777]Reed, B. Y.; Heller, H. J.; Gitomer, W. L.; Pak, C. Y. C.:

Mapping

[128778]a gene defect in absorptive hypercalciuria to chromosome
1q23.3–q24. J.

[128779]Clin. Endocr. Metab. 84: 3907–3913, 1999.

[128780]

[128781]

[128782]11334.

[128783]Gauss, R.; Seifert, R.; Kaupp, U. B.: Molecular identification

[128784]of a hyperpolarization–activated channel in sea urchin
sperm. Nature 393:

[128785]583–587, 1998.

[128786]

[128787]

[128788]11335.

[128789]Seifert, R.; Scholten, A.; Gauss, R.; Mincheva, A.; Lichter,
P.;

[128790]Kaupp, U. B.: Molecular characterization of a slowly gating
human

[128791]hyperpolarization–activated channel predominantly ex-
pressed in thalamus,

[128792]heart, and testis. Proc. Nat. Acad. Sci. 96: 9391–9396,
1999.

[128793]

[128794]

[128795]11336.

[128796]Aihara, Y.; Mashima, H.; Onda, H.; Hisano, S.; Kasuya, H.;
Hori,

[128797]T.; Yamada, S.; Tomura, H.; Yamada, Y.; Inoue, I.; Kojima,
I.; Takeda,

[128798]J.: Molecular cloning of a novel brain-type
Na(+)-dependent inorganic

[128799]phosphate cotransporter. J. Neurochem. 74: 2622–2625,
2000.

[128800]

[128801]

[128802]11337.

[128803]Bellocchio, E. E.; Reimer, R. J.; Fremeau, R. T., Jr.; Edwards,

[128804]R. H.: Uptake of glutamate into synaptic vesicles by an in-
organic

[128805]phosphate transporter. Science 289: 957–960, 2000.

[128806]

[128807]

[128808]11338.

[128809]Ni, B.; Du, Y.; Wu, X.; DeHoff, B. S.; Rosteck, P. R., Jr.; Paul,

[128810]S. M.: Molecular cloning, expression, and chromosomal
localization

[128811]of a human brain-specific Na(+)-dependent inorganic
phosphate cotransporter. J.

[128812]Neurochem. 66: 2227–2238, 1996.

[128813]

[128814]

[128815]11339.

[128816]Takamori, S.; Rhee, J. S.; Rosenmund, C.; Jahn, R.: Identifi-
cation

[128817]of a vesicular glutamate transporter that defines a gluta-
matergic

[128818]phenotype in neurons. Nature 407: 189–194, 2000.

[128819]

[128820]

[128821]11340.

[128822]Blackwood, D.: P300, a state and a trait marker in
schizophrenia. Lancet 355:

[128823]771–772, 2000.

[128824]

[128825]

[128826]11341.

[128827]Blackwood, D. H. R.; Fordyce, A.; Walker, M. T.; St. Clair,
D.

[128828]M.; Porteous, D. J.; Muir, W. J.: Schizophrenia and affective
disorders--cosegregation

[128829]with a translocation at chromosome 1q42 that directly
disrupts brain-expressed

[128830]genes: clinical and P300 findings in a family. Am. J. Hum.
Genet. 69:

[128831]428-433, 2001.

[128832]

[128833]

[128834]11342.

[128835]Ekelund, J.; Hovatta, I.; Parker, A.; Paunio, T.; Varilo, T.;
Martin,

[128836]R.; Suhonen, J.; Ellonen, P.; Chan, G.; Sinsheimer, J. S.; So-
bel,

[128837]E.; Juvonen, H.; Arajärvi, R.; Partonen, T.; Suvisaari, J.;
Lonnqvist,

[128838]J.; Meyer, J.; Peltonen, L.: Chromosome 1 loci in Finnish schizophrenia

[128839]families. Hum. Molec. Genet. 10: 1611–1617, 2001.

[128840]

[128841]

[128842]1 1343.

[128843]Millar, J. K.; Wilson–Annan, J. C.; Anderson, S.; Christie, S.;

[128844]Taylor, M. S.; Semple, C. A. M.; Devon, R. S.; St. Clair, D.
M.; Muir,

[128845]W. J.; Blackwood, D. H. R.; Porteous, D. J.: Disruption of
two novel

[128846]genes by a translocation co–segregating with schizophrenia. Hum.

[128847]Molec. Genet. 9: 1415–1423, 2000.

[128848]

[128849]

[128850]1 1344.

[128851]St. Clair, D.; Blackwood, D.; Muir, W.; Carothers, A.;
Walker,

[128852]M.; Spowart, G.; Gosden, C.; Evans, H. J.: Association

within a family

[128853]of a balanced autosomal translocation with major mental illness. Lancet 336:

[128854]13–16, 1990.

[128855]

[128856]

[128857]11345.

[128858]Bulfone, A.; Menguzzato, E.; Broccoli, V.; Marchitello, A.; Gattuso,

[128859]C.; Mariani, M.; Consalez, G. G.; Martinez, S.; Ballabio, A.; Banfi,

[128860]S.: Barhl1, a gene belonging to a new subfamily of mammalian homeobox

[128861]genes, is expressed in migrating neurons of the CNS. Hum. Molec.

[128862]Genet. 9: 1443–1452, 2000.

[128863]

[128864]

[128865]11346.

[128866]Alessi, D. R.; Deak, M.; Casamayor, A.; Caudwell, F. B.;

Morrice,

[128867]N.; Norman, D. G.; Gaffney, P.; Reese, C. B.; MacDougall,
C. N.; Harbison,

[128868]D.; Ashworth, A.; Bownes, M.:

3-phosphoinositide-dependent protein

[128869]kinase-1 (PDK1): structural and functional homology with
the *Drosophila*

[128870]DSTPK61 kinase. *Curr. Biol.* 7: 776-789, 1997.

[128871]

[128872]

[128873]11347.

[128874]Bertin, J.; Nir, W.-J.; Fischer, C. M.; Tayber, O. V.; Errada,

[128875]P. R.; Grant, J. R.; Keilty, J. J.; Gosselin, M. L.; Robison, K.
E.;

[128876]Wong, G. H. W.; Glucksmann, M. A.; DiStefano, P. S.: Hu-
man CARD4

[128877]protein is a novel CED-4/Apaf-1 cell death family member
that activates

[128878]NF-kappa-B. *J. Biol. Chem.* 274: 12955-12958, 1999.

[128879]

[128880]

[128881]11348.

[128882]Cohen, D. E.; Green, R. M.; Wu, M. K.; Beier, D. R.: Cloning,

[128883]tissue-specific expression, gene structure and chromosomal localization

[128884]of human phosphatidylcholine transfer protein. Biochim. Biophys.

[128885]Acta 1447: 265–270, 1999.

[128886]

[128887]

[128888]11349.

[128889]van Helvoort, A.; de Brouwer, A.; Ottenhoff, R.; Brouwers, J. F.

[128890]H. M.; Wijnholds, J.; Beijnen, J. H.; Rijneveld, A.; van der Valk,

[128891]M. A.; Majoor, D.; Voorhout, W.; Wirtz, K. W. A.; Elferink, R. P.

[128892]J. O.; Borst, P.: Mice without phosphatidylcholine transfer protein

[128893]have no defects in the secretion of phosphatidylcholine into bile

[128894]or into lung airspaces. Proc. Nat. Acad. Sci. 96:

11501–11506, 1999.

[128895]

[128896]

[128897]11350.

[128898]de Rooij, J.; Zwartkruis, F. J. T.; Verheijen, M. H. G.; Cool,

[128899]R. H.; Nijman, S. M. B.; Wittinghofer, A.; Bos, J. L.: Epac is a

[128900]Rap1 guanine–nucleotide–exchange factor directly activated by cyclic

[128901]AMP. *Nature* 396: 474–477, 1998.

[128902]

[128903]

[128904]11351.

[128905]Kawasaki, H.; Springett, G. M.; Mochizuki, N.; Toki, S.;

Nakaya,

[128906]M.; Matsuda, M.; Housman, D. E.; Graybiel, A. M.: A family of cAMP–binding

[128907]proteins that directly activate Rap1. *Science* 282: 2275–2279, 1998.

[128908]

[128909]

[128910]11352.

[128911]Hahn, Y.; Lee, J.; Seong, C.; Yoon, J.; Chung, J. H.: Struc-
tural

[128912]analysis of phylogenetically conserved J domain protein
gene. Biochim.

[128913]Biophys. Acta 1447: 325–333, 1999.

[128914]

[128915]

[128916]11353.

[128917]Hollopeter, G.; Jantzen, H.–M.; Vincent, D.; Li, G.; England,
L.;

[128918]Ramakrishnan, V.; Yang, R.–B.; Nurden, P.; Nurden, A.;
Julius, D.;

[128919]Conley, P. B.: Identification of the platelet ADP receptor
targeted

[128920]by antithrombotic drugs. Nature 409: 202–206, 2001.

[128921]

[128922]

[128923]11354.

[128924]Nurden, P.; Savi, P.; Heilmann, E.; Bihour, C.; Herbert, J.-M.;

[128925]Maffrand, J.-P.; Nurden, A.: An inherited bleeding disorder linked

[128926]to a defective interaction between ADP and its receptor on platelets:

[128927]its influence on glycoprotein IIb-IIIa complex function. J. Clin.

[128928]Invest. 95: 1612-1622, 1995.

[128929]

[128930]

[128931]11355.

[128932]Hill, K. E.; Dasouki, M.; Phillips, J. A., III; Burk, R. F.: Human man

[128933]selenoprotein P gene maps to 5q31. Genomics 36: 550-551, 1996.

[128934]

[128935]

[128936]11356.

[128937]Hill, K. E.; Lloyd, R. S.; Burk, R. F.: Conserved nucleotide sequences

[128938]in the open reading frame and 3-prime untranslated region of selenoprotein

[128939]P mRNA. Proc. Nat. Acad. Sci. 90: 537–541, 1993.

[128940]

[128941]

[128942]11357.

[128943]Keshan Disease Research Group of the Chinese Academy of Medical

[128944]Sciences: Observations on effect of sodium selenite in prevention

[128945]of Keshan disease. Chinese Med. J. 92: 471–476, 1979.

[128946]

[128947]

[128948]11358.

[128949]Eberhart, C. G.; Maines, J. Z.; Wasserman, S. A.: Meiotic cell

[128950]cycle requirement for a fly homologue of human Deleted in Azoospermia. Nature 381:

[128951]783–785, 1996.

[128952]

[128953]

[128954]11359.

[128955]Rugglu, M.; Speed, R.; Taggart, M.; McKay, S. J.; Kianowski, F.;

[128956]Saunders, P.; Derin, J.; Cooke, H. J.: The mouse Dazla gene encodes

[128957]a cytoplasmic protein essential for gametogenesis. Nature 389: 73–77,

[128958]1997.

[128959]

[128960]

[128961]11360.

[128962]Shan, Z.; Hirschmann, P.; Seebacher, T.; Edelman, A.; Jauch, A.;

[128963]Morell, J.; Urbitsch, P.; Vogt, P. H.: A SPGY copy homologous to

[128964]the mouse gene Dazla and the Drosophila gene boule is autosomal and

[128965]expressed only in the human male gonad. Hum. Molec. Genet. 5: 2005–2011,

[128966]1996.

[128967]

[128968]

[128969]11361.

[128970]Slee, R.; Grimes, B.; Speed, R. M.; Taggart, M.; Maguire, S.
M.;

[128971]Ross, A.; McGill, N. I.; Saunders, P. T. K.; Cooke, H. J.: A
human

[128972]DAZ transgene confers partial rescue of the mouse Dazl
null phenotype. Proc.

[128973]Nat. Acad. Sci. 96: 8040–8045, 1999.

[128974]

[128975]

[128976]11362.

[128977]Yen, P. H.; Chai, N. N.; Salido, E. C.: The human autosomal
gene

[128978]DAZLA: testis specificity and a candidate for male infertili-
ty. Hum.

[128979]Molec. Genet. 5: 2013–2017, 1996.

[128980]

[128981]

[128982]11363.

[128983]Akiyama, T. E.; Sakai, S.; Lambert, G.; Nicol, C. J.; Matsue,

[128984]K.; Pimprale, S.; Lee, Y.-H.; Ricote, M.; Glass, C. K.; Brewer, H.

[128985]B., Jr.; Gonzalez, F. J.: Conditional disruption of the peroxisome

[128986]proliferator-activated receptor gamma gene in mice results in lowered

[128987]expression of ABCA1, ABCG1, and apoE in macrophages and reduced cholesterol

[128988]efflux. Molec. Cell. Biol. 22: 2607-2619, 2002.

[128989]

[128990]

[128991]11364.

[128992]Ardlie, K. G.; Lunetta, K. L.; Seielstad, M.: Testing for population

[128993]subdivision and association in four case-control studies. Am. J.

[128994]Hum. Genet. 71: 304-311, 2002.

[128995]

[128996]

[128997]11365.

[128998]Barak, Y.; Nelson, M. C.; Ong, E. S.; Jones, Y. Z.; Ruiz-
Lozano,

[128999]P.; Chien, K. R.; Koder, A.; Evans, R. M.: PPAR-gamma is
required

[129000]for placental, cardiac, and adipose tissue development.
Molec. Cell 4:

[129001]585-595, 1999.

[129002]

[129003]

[129004]11366.

[129005]Barroso, I.; Gurnell, M.; Crowley, V. E. F.; Agostini, M.;
Schwabel,

[129006]J. W.; Soos, M. A.; Masien, G. L.; Williams, T. D. M.; Lewis,
H.;

[129007]Schafer, A. J.; Chatterjee, V. K. K.; O'Rahilly, S.: Dominant
negative

[129008]mutations in human PPAR-gamma associated with severe
insulin resistance,

[129009]diabetes mellitus and hypertension. Nature 402: 880-883,

1999.

[129010]

[129011]

[129012]11367.

[129013]Beamer, B. A.; Yen, C.-J.; Andersen, R. E.; Muller, D.; Elahi,

[129014]D.; Cheskin, L. J.; Andres, R.; Roth, J.; Shuldiner, A. R.: Association
sociation

[129015]of the pro12ala variant in the peroxisome proliferator-activated receptor-gamma-2

[129016]gene with obesity in two Caucasian populations. Diabetes
47: 1806-1808,

[129017]1998.

[129018]

[129019]

[129020]11368.

[129021]Beamer, B. A.; Negri, C.; Yen, C.-J.; Gavrilova, O.; Rumberger,

[129022]J. M.; Durcan, M. J.; Yarnall, D. P.; Hawkins, A. L.; Griffin,
C.

[129023]A.; Burns, D. K.; Roth, J.; Reitman, M.; Shuldiner, A. R.:

Chromosomal

[129024] localization and partial genomic structure of the human peroxisome

[129025] proliferator activated receptor- γ (hPPAR- γ) gene. Biochem.

[129026] Biophys. Res. Commun. 233: 756-759, 1997.

[129027]

[129028]

[129029] 11369.

[129030] Chawla, A.; Boisvert, W. A.; Lee, C.-H.; Laffitte, B. A.; Barak,

[129031] Y.; Joseph, S. B.; Liao, D.; Nagy, L.; Edwards, P. A.; Curtiss, L.

[129032] K.; Evans, R. M.; Tontonoz, P.: A PPAR- γ -LXR-ABCA1 pathway in

[129033] macrophages is involved in cholesterol efflux and atherogenesis. Molec.

[129034] Cell 7: 161-171, 2001.

[129035]

[129036]

[129037] 11370.

[129038]Deeb, S. S.; Fajas, L.; Nemoto, M.; Pihlajamaki, J.; Mykka-
nen,

[129039]L.; Kuusisto, J.; Laakso, M.; Fujimoto, W.; Auwerx, J.: A
pro12ala

[129040]substitution in PPAR-gamma-2 associated with decreased
receptor activity,

[129041]lower body mass index and improved insulin sensitivity.
Nature Genet. 20:

[129042]284-287, 1998.

[129043]

[129044]

[129045]11371.

[129046]Elbrecht, A.; Chen, Y.; Cullinan, C. A.; Hayes, N.; Lei-
bowitz,

[129047]M. D.; Moller, D. E.; Berger, J.: Molecular cloning, expres-
sion and

[129048]characterization of human peroxisome proliferator acti-
vated receptors

[129049]gamma-1 and gamma-2. Biochem. Biophys. Res. Com-
mun. 224: 431-437,

[129050]1996.

[129051]

[129052]

[129053]11372.

[129054]Fajas, L.; Auboeuf, D.; Raspe, E.; Schoonjans, K.; Lefebvre, A.

[129055]M.; Saladin, R.; Najib, J.; Laville, M.; Fruchart, J.-C.; Deeb, S.;

[129056]Vidal-Puig, A.; Flier, J.; Briggs, M. R.; Staels, B.; Vidal, H.; Auwerx,

[129057]J.: The organization, promoter analysis, and expression of the human

[129058]PPAR-gamma gene. J. Biol. Chem. 272: 18779-18789, 1997.

[129059]

[129060]

[129061]11373.

[129062]Gampe, R. T., Jr.; Montana, V. G.; Lambert, M. H.; Miller, A.

[129063]B.; Bledsoe, R. K.; Milburn, M. V.; Kliewer, S. A.; Willson, T. M.;

[129064]Xu, H. E.: Asymmetry in the PPAR-gamma/RXR-alpha crystal structure

[129065]reveals the molecular basis of heterodimerization among nuclear receptors. Molec.

[129066]Cell 5: 545–555, 2000.

[129067]

[129068]

[129069]11374.

[129070]Chakraborti, A.; Lippman, D. L.; Loh, H. H.; Kozak, C. A.; Lee,

[129071]N. M.: Genetic mapping of opioid binding protein gene(s) to mouse

[129072]chromosome 9. Mammalian Genome 4: 179–182, 1993.

[129073]

[129074]

[129075]11375.

[129076]Cho, T. M.; Hasegawa, J.-I.; Ge, B.-I.; Loh, H. H.: Purification

[129077]to apparent homogeneity of a mu-type opioid receptor from rat brain. Proc.

[129078]Nat. Acad. Sci. 83: 4138–4142, 1986.

[129079]

[129080]

[129081]11376.

[129082]Shark, K. B.; Lee, N. M.: Cloning, sequencing and localization

[129083]to chromosome 11 of a cDNA encoding a human opioid-binding cell adhesion

[129084]molecule (OBCAM). Gene 155: 213–217, 1995.

[129085]

[129086]

[129087]11377.

[129088]Burmeister, M.; Meyer, G. E.: The trefoil gene maps to mouse chromosome

[129089]17. Mammalian Genome 8: 223–224, 1997.

[129090]

[129091]

[129092]11378.

[129093]Chinery, R.; Williamson, J.; Poulsom, R.: The gene encoding human

[129094]intestinal trefoil factor (TFF3) is located on chromosome

21q22.3

[129095]clustered with other members of the trefoil peptide family.

Genomics 32:

[129096]281–284, 1996.

[129097]

[129098]

[129099]11379.

[129100]Mashimo , H.; Wu, D.–C.; Podolsky, D. K.; Fishman, M. C.:

Impaired

[129101]defense of intestinal mucosa in mice lacking intestinal

trefoil factor. Science 274:

[129102]262–265, 1996.

[129103]

[129104]

[129105]11380.

[129106]Podolsky, D. K.; Lynch–Devaney, K.; Stow, J. L.; Oates, P.;

Murgue,

[129107]B.; DeBeaumont, M.; Sands, B. E.; Mahida, Y. R.: Identifica–

tion of

[129108]human intestinal trefoil factor: goblet cell–specific ex–

pression of

[129109]a peptide targeted for apical secretion. J. Biol. Chem. 268:
6694–6702,

[129110]1993.

[129111]

[129112]

[129113]11381.

[129114]Probst, J. C.; Zetzsche, T.; Weber, M.; Theilemann, P.;
Skutella,

[129115]T.; Landgraf, R.; Jirikowski, G. F.: Human intestinal trefoil
factor

[129116]is expressed in human hypothalamus and pituitary: evi-
dence for a novel

[129117]neuropeptide. FASEB J. 10: 1518–1523, 1996.

[129118]

[129119]

[129120]11382.

[129121]Schmitt, H.; Wundrack, I.; Beck, S.; Gott, P.; Welter, C.;
Shizuya,

[129122]H.; Simon, M. I.; Blin, N.: A third P-domain peptide gene

(TFF3),

[129123]human intestinal trefoil factor, maps to 21q22.3. Cyto-
genet. Cell

[129124]Genet. 72: 299–302, 1996.

[129125]

[129126]

[129127]11383.

[129128]Taupin, D.; Wu, D.–C.; Jeon, W.–K.; Devaney, K.; Wang, T.
C.; Podolsky,

[129129]D. K.: The trefoil gene family are coordinately expressed
immediate–early

[129130]genes: EGF receptor– and MAP kinase–dependent inter-
regulation. J.

[129131]Clin. Invest. 103: R31–R38, 1999.

[129132]

[129133]

[129134]11384.

[129135]Thim, L.: A new family of growth factor–like peptides: 'tre-
foil'

[129136]disulphide loop structures as a common feature in breast

cancer associated

[129137]peptide (pS2), pancreatic spasmolytic polypeptide (PSP),
and frog

[129138]skin peptides (spasmolysins). FEBS Lett. 250: 85–90,
1989.

[129139]

[129140]

[129141]11385.

[129142]Sugimoto, Y.; Yatsunami, K.; Tsujimoto, M.; Khorana, H.
G.; Ichikawa,

[129143]A.: The amino acid sequence of a glutamic acid–rich pro-
tein from

[129144]bovine retina as deduced from the cDNA sequence. Proc.
Nat. Acad.

[129145]Sci. 88: 3116–3119, 1991.

[129146]

[129147]

[129148]11386.

[129149]Marigo, V.; Roberts, D. J.; Lee, S. M. K.; Tsukurov, O.; Levi,

[129150]T.; Gastier, J. M.; Epstein, D. J.; Gilbert, D. J.; Copeland, N.

G.;

[129151]Seidman, C. E.; Jenkins, N. A.; Seidman, J. G.; McMahon, A. P.; Tabin,

[129152]C.: Cloning, expression, and chromosomal location of SHH and IHH:

[129153]two human homologues of the Drosophila segment polarity gene hedgehog. Genomics 28:

[129154]44–51, 1995.

[129155]

[129156]

[129157]11387.

[129158]McIntire, J. J.; Umetsu, S. E.; Akbari, O.; Potter, M.; Kuchroo,

[129159]V. K.; Barsh, G. S.; Freeman, G. J.; Umetsu, D. T.; DeKruyff, R. H.

[129160]: Identification of Tapr (an airway hyperreactivity regulatory locus)

[129161]and the linked Tim gene family. Nature Immun. 2: 1109–1116, 2001.

[129162]

[129163]

[129164]11388.

[129165]Hwang, B. J.; Ford, J. M.; Hanawalt, P. C.; Chu, G.: Expression

[129166]of the p48 xeroderma pigmentosum gene is p53-dependent and is involved

[129167]in global genomic repair. Proc. Nat. Acad. Sci. 96: 424–428, 1999.

[129168]

[129169]

[129170]11389.

[129171]Tahir, S. A.; Yang, G.; Ebara, S.; Timme, T. L.; Satoh, T.; Li,

[129172]L.; Goltsov, A.; Ittmann, M.; Morrisett, J. D.; Thompson, T. C.:

[129173]Secreted caveolin–1 stimulates cell survival/clonal growth and contributes

[129174]to metastasis in androgen–insensitive prostate cancer.

Cancer Res. 61:

[129175]3882–3885, 2001.

[129176]

[129177]

[129178]11390.

[129179]England, S. K.; Uebele, V. N.; Kodali, J.; Bennett, P. B.;

Tamkun,

[129180]M. M.: A novel K⁺ channel beta-subunit (hKv-beta-1.3) is
produced

[129181]via alternative mRNA splicing. J. Biol. Chem. 270:

28531-28534,

[129182]1995.

[129183]

[129184]

[129185]11391.

[129186]Kratz, C. P.; Emerling, B. M.; Bonifas, J.; Wang, W.; Green,
E.

[129187]D.; Le Beau, M. M.; Shannon, K. M.: Genomic structure of
the PIK3CG

[129188]gene on chromosome band 7q22 and evaluation as a can-
didate myeloid

[129189]tumor suppressor. Blood 99: 372-374, 2002.

[129190]

[129191]

[129192]11392.

[129193]Temple, I. K.; Gardner, R. J.; Robinson, D. O.; Kibirige, M. S.;

[129194]Ferguson, A. W.; Baum, J. D.; Barber, J. C. K.; James, R. S.; Shield,

[129195]J. P. H.: Further evidence for an imprinted gene for neonatal diabetes

[129196]localised to chromosome 6q22–q23. Hum. Molec. Genet. 5: 1117–1124,

[129197]1996.

[129198]

[129199]

[129200]11393.

[129201]Ahmad, W.; Li, S.; Chen, H.; Tuck–Muller, C. M.; Pittler, S. J.;

[129202]Aronson, N. N., Jr.: Lysosomal chitobiase (CTB) and the G–protein

[129203]gamma–5 subunit (GNG5) genes co-localize to human chromosome 1p22. Cytogenet.

[129204]Cell Genet. 71: 44–46, 1995.

[129205]

[129206]

[129207]11394.

[129208]Fisher, K. J.; Aronson, N. N.: Characterization of the cDNA
and

[129209]genomic sequence of a G protein gamma subunit
(gamma-5). Molec. Cell

[129210]Biol. 12: 1585-1591, 1992.

[129211]

[129212]

[129213]11395.

[129214]Fiaschi, T.; Marzella, R.; Veggi, D.; Marzocchini, R.;
Raugei,

[129215]G.; Rocchi, M.; Ramponi, G.: Assignment of the human
erythrocyte

[129216]acylphosphatase gene (ACYP1) to chromosome band
14q24.3. Cytogenet.

[129217]Cell Genet. 81: 235-236, 1998.

[129218]

[129219]

[129220]11396.

- [129221]Fiaschi, T.; Raugei, G.; Marzocchini, R.; Chiarugi, P.; Cirri,
[129222]P.; Ramponi, G.: Cloning and expression of the cDNA cod-
ing for the
[129223]erythrocyte isoenzyme of human acylphosphatase. FEBS
Lett. 367:
[129224]145–148, 1995.
- [129225]
- [129226]
- [129227]11397.
- [129228]Goldowitz, D.; Smeyne, R. J.: Tune into the weaver chan-
nel. Nature
[129229]Genet. 11: 107–109, 1995.
- [129230]
- [129231]
- [129232]11398.
- [129233]Hess, E. J.: Identification of the weaver mouse mutation:
the
[129234]end of the beginning. Neuron 16: 1073–1076, 1996.
- [129235]

[129236]

[129237]11399.

[129238]Lane, P. W.: New mutation: Weaver, wv. Mouse News Letter
32–33,

[129239]1964.

[129240]

[129241]

[129242]11400.

[129243]Lesage, F.; Duprat, F.; Fink, M.; Guillemare, E.; Coppola,
T.;

[129244]Lazdunski, M.; Hugnot, J.–P.: Cloning provides evidence
for a family

[129245]of inward rectifier and G–protein coupled K(+) channels in
the brain. FEBS

[129246]Lett. 353: 37–42, 1994.

[129247]

[129248]

[129249]11401.

[129250]Rakic, P.; Sidman, R. L.: Sequence of developmental ab–

normalities

[129251]leading to granule cell deficit in cerebellar cortex of
weaver mutant

[129252]mice. J. Comp. Neurol. 152: 103–132, 1973.

[129253]

[129254]

[129255]11402.

[129256]Sakura, H.; Bond, C.; Warren–Perry, M.; Horsley, S.; Kear–
ney, L.;

[129257]Tucker, S.; Adelman, J.; Turner, R.; Ashcroft, F. M.: Char–
acterization

[129258]and variation of a human inwardly–rectifying K–channel
gene (KCNJ6):

[129259]a putative ATP–sensitive K–channel subunit. FEBS Lett.
367: 193–197,

[129260]1995.

[129261]

[129262]

[129263]11403.

[129264]Tsaur, M.–L.; Menzel, S.; Lai, F.–P.; Espinosa, R., III; Con–

cannon,

[129265]P.; Spielman, R. S.; Hanis, C. L.; Cox, N. J.; Le Beau, M. M.;

German,

[129266]M. S.; Jan, L. Y.; Bell, G. I.; Stoffel, M.: Isolation of a cDNA

[129267]clone encoding a K(ATP) channel-like protein expressed in
insulin-secreting

[129268]cells, localization of the human gene to chromosome
band 21q22.1 and

[129269]linkage studies with NIDDM. Diabetes 44: 592–596, 1995.

[129270]

[129271]

[129272]11404.

[129273]Yasuda, K.; Sakura, H.; Mori, Y.; Iwamoto, K.; Shimokawa,
K.;

[129274]Kadowaki, H.; Hagura, R.; Akanuma, Y.; Adelman, J. P.;
Yazaki, Y.;

[129275]Ashcroft, F. M.; Kadowaki, T.: No evidence for mutations
in a putative

[129276]subunit of the beta-cell ATP-sensitive potassium channel
(K-ATP channel)

[129277]in Japanese NIDDM patients. Biochem. Biophys. Res. Com-

mun. 211:

[129278]1036–1040, 1995.

[129279]

[129280]

[129281]11405.

[129282]Efiok, B. J. S.; Chiorini, J. A.; Safer, B.: A key transcription

[129283]factor for eukaryotic initiation factor–2–alpha is strongly
homologous

[129284]to developmental transcription factors and may link
metabolic genes

[129285]to cellular growth and development. J. Biol. Chem. 269:
18921–18930,

[129286]1994.

[129287]

[129288]

[129289]11406.

[129290]Gopalakrishnan, L.; Scarpulla, R. C.: Structure, expression,
and

[129291]chromosomal assignment of the human gene encoding
nuclear respiratory

[129292]factor 1. J. Biol. Chem. 270: 18019–18025, 1995.

[129293]

[129294]

[129295]11407.

[129296]Spelbrink, J. N.; Van den Bogert, C.: The pre-mRNA of nuclear
clear

[129297]respiratory factor 1, a regulator of mitochondrial biogenesis, is

[129298]alternatively spliced in human tissues and cell lines. Hum.
Molec.

[129299]Genet. 4: 1591–1596, 1995.

[129300]

[129301]

[129302]11408.

[129303]Tiranti, V.; Rossi, E.; Rocchi, M.; DiDonato, S.; Zuffardi, O.;

[129304]Zeviani, M.: The gene (NFE2L1) for human NRF-1, an activator involved

[129305]in nuclear-mitochondrial interactions, maps to 7q32. Genomics 27:

[129306]555–557, 1995.

[129307]

[129308]

[129309]11409.

[129310]Ohshiro, T.; Yagami, T.; Zhang, C.; Matsuzaki, F.: Role of
cortical

[129311]tumour-suppressor proteins in asymmetric division of
Drosophila neuroblast. Nature 408:

[129312]593–596, 2000.

[129313]

[129314]

[129315]11410.

[129316]Peng, C.–Y.; Manning, L.; Albertson, R.; Doe, C. Q.: The
tumour-suppressor

[129317]genes lgl and dlg regulate basal protein targeting in
Drosophila neuroblasts. Nature 408:

[129318]596–600, 2000.

[129319]

[129320]

[129321]11411.

[129322]Strand, D.; Unger, S.; Corvi, R.; Hartenstein, K.; Schenkel, H.;

[129323]Kalmes, A.; Merdes, G.; Neumann, B.; Krieg-Schneider, F.; Coy, J.

[129324]F.; Poustka, A.; Schwab, M.; Mechler B. M.: A human homologue of

[129325]the Drosophila tumour suppressor gene l(2)gl maps to 17p11.2-12 and

[129326]codes for a cytoskeletal protein that associates with non-muscle myosin

[129327]II heavy chain. Oncogene 11: 291-301, 1995.

[129328]

[129329]

[129330]11412.

[129331]Timchenko, L. T.; Miller, J. W.; Timchenko, N. A.; DeVore, D. R.;

[129332]Datar, K. V.; Lin, L.; Roberts, R.; Caskey, C. T.; Swanson, M. S.

[129333]: Identification of a (CUG)_n triplet repeat RNA-binding protein and

[129334]its expression in myotonic dystrophy. Nucleic Acids Res. 24: 4407-4414,

[129335]1996.

[129336]

[129337]

[129338]11413.

[129339]Timchenko, L. T.; Timchenko, N. A.; Caskey, C. T.;
Roberts, R.

[129340]: Novel proteins with binding specificity for DNA CTG re-
peats and

[129341]RNA CUG repeats: implications for myotonic dystrophy.
Hum. Molec.

[129342]Genet. 5: 115–121, 1996.

[129343]

[129344]

[129345]11414.

[129346]Wary, K. K.; Mariotti, A.; Zurzolo, C.; Giancotti, F. G.: A re-
quirement

[129347]for caveolin–1 and associated kinase Fyn in integrin sig-
naling and

[129348]anchorage–dependent cell growth. Cell 94: 625–634,
1998.

[129349]

[129350]

[129351]11415.

[129352]Yang, G.; Truong, L. D.; Timme, T. L.; Ren, C.; Wheeler, T.
M.;

[129353]Park, S. H.; Nasu, Y.; Bangma, C. H.; Kattan, M. W.;
Scardino, P.

[129354]T.; Thompson, T. C.: Elevated expression of caveolin is
associated

[129355]with prostate and breast cancer. Clin. Cancer Res. 4:
1873–1880,

[129356]1998.

[129357]

[129358]

[129359]11416.

[129360]Meij, I. C.; Koenderink, J. B.; van Bokhoven, H.; Assink, K.
F.

[129361]H.; Groenestege, W. T.; de Pont, J. J. H. H. M.; Bindels, R. J.
M.;

[129362]Monnens, L. A. H.; van den Heuvel, L. P. W. J.; Knoers, N.
V. A. M.

[129363]: Dominant isolated renal magnesium loss is caused by
misrouting of

[129364]the Na⁺,K⁽⁺⁾-ATPase gamma-subunit. Nature Genet. 26:
265–266, 2000.

[129365]Note: Erratum. Nature Genet. 27: 125 only, 2001.

[129366]

[129367]

[129368]11417.

[129369]Sweadner, K. J.; Rael, E.: The FXYD gene family of small
ion transport

[129370]regulators or channels: cDNA sequence, protein signature
sequence,

[129371]and expression. Genomics 68: 41–56, 2000.

[129372]

[129373]

[129374]11418.

[129375]Sweadner, K. J.; Wetzel, R. K.; Arystarkhova, E.: Genomic
organization

[129376]of the human FXYD2 gene encoding the gamma subunit of
the Na,K-ATPase. Biochem.

[129377]Biophys. Res. Commun. 279: 196–201, 2000.

[129378]

[129379]

[129380]11419.

[129381]Blair, E.; Redwood, C.; Ashrafian, H.; Oliveira, M.; Broxholme,

[129382]J.; Kerr, B.; Salmon, A.; Ostman-Smith, I.; Watkins, H.: Mutations

[129383]in the gamma-2 subunit of AMP-activated protein kinase cause familial

[129384]hypertrophic cardiomyopathy: evidence for the central role of energy

[129385]compromise in disease pathogenesis. Hum. Molec. Genet. 10: 1215–1220,

[129386]2001.

[129387]

[129388]

[129389]11420.

[129390]Siderovski, D. P.; Heximer, S. P.; Forsdyke, D. R.: A human gene

[129391]encoding a putative basic helix-loop-helix phosphoprotein whose mRNA

[129392]increases rapidly in cycloheximide-treated blood mononuclear cells. DNA

[129393]Cell Biol. 13: 125-147, 1994.

[129394]

[129395]

[129396]11421.

[129397]Hirai, H.; Roussel, M. F.; Kato, J.-Y.; Ashmun, R. A.; Sherr, C.

[129398]J.: Novel INK4 proteins, p19 and p18, are specific inhibitors of

[129399]cyclin D-dependent kinases CDK4 and CDK6. Molec. Cell. Biol. 15:

[129400]2672-2681, 1995.

[129401]

[129402]

[129403]11422.

[129404]Okuda, T.; Hirai, H.; Valentine, V. A.; Shurtleff, S. A.; Kidd,

[129405]V. J.; Lahti, J. M.; Sherr, C. J.; Downing, J. R.: Molecular

cloning,

[129406]expression pattern, and chromosomal localization of human CDKN2D/INK4d,

[129407]an inhibitor of cyclin D-dependent kinases. Genomics 29: 623–630,

[129408]1995.

[129409]

[129410]

[129411]11423.

[129412]Sago, H.; Kitagawa, M.; Obata, S.; Mori, N.; Taketani, S.; Rochelle,

[129413]J. M.; Seldin, M. F.; Davidson, M.; St. John, T.; Suzuki, S. T.:

[129414]Cloning, expression, and chromosomal localization of a novel cadherin-related

[129415]protein, protocadherin-3. Genomics 29: 631–640, 1995.

[129416]

[129417]

[129418]11424.

[129419]Fiorucci, S.; Mencarelli, A.; Palazzetti, B.; Distrutti, E.; Vergnolle,

- [129420]N.; Hollenberg, M. D.; Wallace, J. L.; Morelli, A.; Cirino, G.:
Proteinase-activated
[129421]receptor 2 is an anti-inflammatory signal for colonic lam-
ina propria
[129422]lymphocytes in a mouse model of colitis. Proc. Nat. Acad.
Sci. 98:
[129423]13936-13941, 2001.
- [129424]
- [129425]
- [129426]11425.
- [129427]Nystedt, S.; Emilsson, K.; Larsson, A.-K.; Strombeck, B.;
Sundelin,
[129428]J.: Molecular cloning and functional expression of the
gene encoding
[129429]the human proteinase activated receptor 2. Europ. J.
Biochem. 232:
[129430]84-89, 1995.
- [129431]

[129432]

[129433]11426.

[129434]Nystedt, S.; Emilsson, K.; Wahlestedt, C.; Sundelin, J.:

Molecular

[129435]cloning of a potential proteinase-activated receptor. Proc.

Nat.

[129436]Acad. Sci. 91: 9208-9212, 1994.

[129437]

[129438]

[129439]11427.

[129440]Devlin, A. M.; Ling, E.; Peerson, J. M.; Fernando, S.; Clarke,

[129441]R.; Smith, A. D.; Halsted, C. H.: Glutamate carboxypepti-

dase II:

[129442]a polymorphism associated with lower levels of serum fo-

late and hyperhomocysteinemia. Hum.

[129443]Molec. Genet. 9: 2837-2844, 2000.

[129444]

[129445]

[129446]11428.

[129447]Israeli, R. S.; Powell, C. T.; Fair, W. R.; Heston, W. D. W.:

[129448]Molecular cloning of a complementary DNA encoding a
prostate-specific

[129449]membrane antigen. Cancer Res. 53: 227–230, 1993.

[129450]

[129451]

[129452]11429.

[129453]Leek, J.; Lench, N.; Maraj, B.; Bailey, A.; Carr, I. M.; Ander-
sen,

[129454]S.; Cross, J.; Whelan, P.; MacLennan, K. A.; Meredith, D. M.;
Markham,

[129455]A. F.: Prostate-specific membrane antigen: evidence for
the existence

[129456]of a second related human gene. Brit. J. Cancer 72:
583–588, 1995.

[129457]

[129458]

[129459]11430.

[129460]Maraj, B. H.; Leek, J. P.; Karayi, M.; Ali, M.; Lench, N. J.;
Markham,

[129461]A. F.: Detailed genetic mapping around a putative prostate-specific

[129462]membrane antigen locus on human chromosome 11p11.2. Cytogenet. Cell

[129463]Genet. 81: 3–9, 1998.

[129464]

[129465]

[129466]11431.

[129467]Ikemoto, M.; Arai, H.; Feng, D.; Tanaka, K.; Aoki, J.; Dohmae,

[129468]N.; Takio, K.; Adachi, H.; Tsujimoto, M.; Inoue, K.: Identification

[129469]of a PDZ-domain-containing protein that interacts with the scavenger

[129470]receptor class B type I. Proc. Nat. Acad. Sci. 97: 6538–6543, 2000.

[129471]

[129472]

[129473]11432.

[129474]O'Keefe, D. S.; Su, S. L.; Bacich, D. J.; Horiguchi, Y.; Luo, Y.;

[129475]Powell, C. T.; Zandvliet, D.; Russell, P. J.; Molloy, P. L.;
Nowak,

[129476]N. J.; Shows, T. B.; Mullins, C.; Vonder Haar, R. A.; Fair, W.
R.;

[129477]Heston, W. D. W.: Mapping, genomic organization and
promoter analysis

[129478]of the human prostate-specific membrane antigen gene.
Biochim. Biophys.

[129479]Acta 1443: 113–127, 1998.

[129480]

[129481]

[129482]11433.

[129483]Rinker-Schaeffer, C. W.; Hawkins, A. L.; Su, S. L.; Israeli, R.

[129484]S.; Griffin, C. A.; Isaacs, J. T.; Heston, W. D. W.: Localiza-
tion

[129485]and physical mapping of the prostate-specific membrane
antigen (PSM)

[129486]gene to human chromosome 11. Genomics 30: 105–108,
1995.

[129487]

[129488]

[129489]11434.

[129490]Beltrame, J. F.; Sasayama, S.; Maseri, A.: Racial heterogeneity

[129491]in coronary artery vasomotor reactivity: differences between Japanese

[129492]and Caucasian patients. J. Am. Coll. Cardiol. 33: 1442–1452, 1999.

[129493]

[129494]

[129495]11435.

[129496]Inagaki, N.; Inazawa, J.; Seino, S.: cDNA sequence, gene structure,

[129497]and chromosomal localization of the human ATP-sensitive potassium

[129498]channel, uK(ATP)–1, gene (KCNJ8). Genomics 30: 102–104, 1995.

[129499]

[129500]

[129501]11436.

[129502]Inagaki, N.; Tsuura, Y.; Namba, N.; Masuda, K.; Gono, T.; Horie,

[129503]M.; Seino, Y.; Mizuta, M.; Seino, S.: Cloning and functional characterization

[129504]of a novel ATP-sensitive potassium channel ubiquitously expressed

[129505]in rat tissues, including pancreatic islets, pituitary, skeletal muscle,

[129506]and heart. J. Biol. Chem. 270: 5691–5694, 1995.

[129507]

[129508]

[129509]11437.

[129510]MacAlpin, R. N.: Cardiac arrest and sudden unexpected death in

[129511]variant angina: complications of coronary spasm that can occur in

[129512]the absence of severe organic coronary stenosis. Am. Heart J. 125:

[129513]1011–1017, 1993.

[129514]

[129515]

[129516]11438.

[129517]Bartsch, J. W.; Mukai, H.; Takahashi, N.; Ronsiek, M.;
Fuchs, S.;

[129518]Jockusch, H.; Ono, Y.: The protein kinase N (PKN) gene
PRKCL1/Prkcl1

[129519]maps to human chromosome 19p12–p13.1 and mouse
chromosome 8 with close

[129520]linkage to the myodystrophy (myd) mutation. Genomics
49: 129–132,

[129521]1998.

[129522]

[129523]

[129524]11439.

[129525]Mukai, H.; Ono, Y.: A novel protein kinase with leucine
zipper–like

[129526]sequences: its catalytic domain is highly homologous to
that of protein

[129527]kinase C. Biochem. Biophys. Res. Commun. 199: 897–904,
1994.

[129528]

[129529]

[129530]11440.

[129531]Palmer, R. H.; Ridden, J.; Parker, P. J.: Identification of multiple,

[129532]novel, protein kinase C-related gene products. FEBS Lett. 356: 5–8,

[129533]1994.

[129534]

[129535]

[129536]11441.

[129537]Palmer, R. H.; Ridden, J.; Parker, P. J.: Cloning and expressions

[129538]patterns of two members of a novel protein-kinase-C-related kinase

[129539]family. Europ. J. Biochem. 227: 344–351, 1995.

[129540]

[129541]

[129542]11442.

[129543]Yu, W.; Liu, J.; Morrice, N. A.; Wettenhall, R. E. H.: Isolation

[129544]and characterization of a structural homologue of human PRK2 from

[129545]rat liver. J. Biol. Chem. 272: 10030–10034, 1997.

[129546]

[129547]

[129548]11443.

[129549]Durkin, M. E.; Loechel, F.; Mattei, M.-G.; Gilpin, B. J.; Albrechtsen,

[129550]R.; Wewer, U. M.: Tissue-specific expression of the human laminin

[129551]alpha-5-chain, and mapping of the gene to human chromosome 20q13.2-13.3

[129552]and to distal mouse chromosome 2 near the locus for the ragged (Ra)

[129553]mutation. FEBS Lett. 411: 296-300, 1997.

[129554]

[129555]

[129556]11444.

[129557]Miner, J. H.; Lewis, R. M.; Sanes, J. R.: Molecular cloning of

[129558]a novel laminin chain, alpha-5, and widespread expression in adult

[129559]mouse tissues. J. Biol. Chem. 270: 28523-28526, 1995.

[129560]

[129561]

[129562]11445.

[129563]Hernandez, A.; Park, J. P.; Lyon, G. J.; Mohandas, T. K.; St.
Germain,

[129564]D. L.: Localization of the type 3 iodothyronine deiodinase
(DIO3)

[129565]gene to human chromosome 14q32 and mouse chromo-
some 12F1. Genomics 53:

[129566]119–121, 1998.

[129567]

[129568]

[129569]11446.

[129570]Huang, S. A.; Tu, H. M.; Harney, J. W.; Venihaki, M.; Butte,
A.

[129571]J.; Kozakewich, H. P. W.; Fishman, S. J.; Larsen, P. R.: Se-
vere hypothyroidism

[129572]caused by type 3 iodothyronine deiodinase in infantile he-
mangiomas. New

[129573]Eng. J. Med. 343: 185–189, 2000.

[129574]

[129575]

[129576]11447.

[129577]Salvatore, D.; Low, S. C.; Berry, M.; Maia, A. L.; Harney, J. W.;

[129578]Croteau, W.; St. German, D. L.; Larsen, P. R.: Type 3
iodothyronine

[129579]deiodinase: cloning, in vitro expression, and functional
analysis

[129580]of the placental selenoenzyme. J. Clin. Invest. 96:
2421–2430, 1995.

[129581]

[129582]

[129583]11448.

[129584]Kolodkin, A. L.; Matthes, D. J.; Goodman, C. S.: The
semaphorin

[129585]genes encode a family of transmembrane and secreted
growth cone guidance

[129586]molecules. Cell 75: 1389–1399, 1993.

[129587]

[129588]

[129589]11449.

[129590]Luo, Y.; Raible, D.; Raper, J. A.: Collapsin: a protein in
brain

[129591]that induces the collapse and paralysis of neuronal growth
cones. Cell 75:

[129592]217–227, 1993.

[129593]

[129594]

[129595]11450.

[129596]Marin, O.; Yaron, A.; Bagri, A.; Tessier–Lavigne, M.;
Rubenstein,

[129597]J. L. R.: Sorting of striatal and cortical interneurons regu-
lated

[129598]by semaphorin–neuropilin interactions. Science 293:
872–875, 2001.

[129599]

[129600]

[129601]11451.

[129602]Sekido, Y.; Bader, S.; Latif, F.; Chen, J.–Y.; Duh, F.–M.; Wei,

[129603]M.–H.; Albanesi, J. P.; Lee, C.–C.; Lerman, M. I.; Minna, J.

D.:

[129604]Human semaphorins A(V) and IV reside in the 3p21.3
small cell lung

[129605]cancer deletion region and demonstrate distinct expres-
sion patterns. Proc.

[129606]Nat. Acad. Sci. 93: 4120–4125, 1996.

[129607]

[129608]

[129609]11452.

[129610]Xiang, R.-H.; Hensel, C. H.; Garcia, D. K.; Carlson, H. C.;
Kok,

[129611]K.; Daly, M. C.; Kerbacher, K.; van den Berg, A.; Veldhuis,
P.; Buys,

[129612]C. H. C. M.; Naylor S. L.: Isolation of the human
semaphorin III/F

[129613]gene (SEMA3F) at chromosome 3p21, a region deleted in
lung cancer. Genomics 32:

[129614]39–48, 1996.

[129615]

[129616]

[129617]11453.

[129618]Ardehali, H.; Tiller, G. E.; Printz, R. L.; Mochizuki, H.; Proc-
hazka,

[129619]M.; Granner, D. K.: A novel (TA)_n polymorphism in the
hexokinase

[129620]II gene: application to noninsulin-dependent diabetes
mellitus in

[129621]the Pima Indians. Hum. Genet. 97: 482–485, 1996.

[129622]

[129623]

[129624]11454.

[129625]Echwald, S. M.; Bjorbaek, C.; Hansen, T.; Clausen, J. O.;
Vestergaard,

[129626]H.; Zierarth, J. R.; Printz, R. L.; Granner, D. K.; Pedersen,
O.:

[129627]Identification of four amino acid substitutions in hexoki-
nase II and

[129628]studies of relationships to NIDDM, glucose effectiveness,
and insulin

[129629]sensitivity. Diabetes 44: 347–353, 1995.

[129630]

[129631]

[129632]11455.

[129633]Heikkinen, S.; Suppola, S.; Malkki, M.; Deeb, S. S.; Janne, J.;

[129634]Laakso, M.: Mouse hexokinase II gene: structure, cDNA, promoter analysis,

[129635]and expression pattern. Mammalian Genome 11: 91–96, 2000.

[129636]

[129637]

[129638]11456.

[129639]Laakso, M.; Malkki, M.; Deeb, S. S.: Amino acid substitutions

[129640]in hexokinase II among patients with NIDDM. Diabetes 44: 330–334,

[129641]1995.

[129642]

[129643]

[129644]11457.

[129645]Lehto, M.; Xiang, K.; Stoffel, M.; Espinosa, R., III; Groop, L.

[129646]C.; Le Beau, M. M.; Bell, G. I.: Human hexokinase II: local–

ization

[129647]of the polymorphic gene to chromosome 2. Diabetologia

36: 1299–1302,

[129648]1993.

[129649]

[129650]

[129651]11458.

[129652]Mathupala, S. P.; Heese, C.; Pedersen, P. L.: Glucose

catabolism

[129653]in cancer cells: the type II hexokinase promoter contains

functionally

[129654]active response elements for the tumor suppressor p53. J.

Biol. Chem. 272:

[129655]22776–22780, 1997.

[129656]

[129657]

[129658]11459.

[129659]Mathupala, S. P.; Rempel, A.; Pedersen, P. L.: Glucose

catabolism

[129660]in cancer cells: isolation, sequence, and activity of the

promoter

[129661]for type II hexokinase. J. Biol. Chem. 270: 16918–16925, 1995.

[129662]

[129663]

[129664]11460.

[129665]Vidal–Puig, A.; Printz, R. L.; Stratton, I. M.; Granner, D. K.;

[129666]Moller, D. E.: Analysis of the hexokinase II gene in subjects with

[129667]insulin resistance and NIDDM and detection of a gln142–to–his substitution. Diabetes 44:

[129668]340–346, 1995.

[129669]

[129670]

[129671]11461.

[129672]Luo, Y.; Roeder, R. G.: Cloning, functional characterization,

[129673]and mechanism of action of the B–cell–specific transcriptional activator

[129674]OCA–B. Molec. Cell. Biol. 15: 4115–4124, 1995.

[129675]

[129676]

[129677]11462.

[129678]Staudt, L. M.; Lenardo, M. J.: Immunoglobulin gene transcription. Ann.

[129679]Rev. Immun. 9: 373–398, 1991.

[129680]

[129681]

[129682]11463.

[129683]Strubin, M.; Newell, J. W.; Matthias, P.: OBF–1, a novel B cell-specific

[129684]coactivator that stimulates immunoglobulin promoter activity through

[129685]association with octamer-binding proteins. Cell 80: 497–506, 1995.

[129686]

[129687]

[129688]11464.

[129689]Endele, S.; Zabel, B.; Winterpacht, A.: Assignment of the

human

[129690]diacylglycerol kinase 4 (DAGK4) gene to chromosome
4p16.3. Genomics 33:

[129691]145–146, 1996.

[129692]

[129693]

[129694]11465.

[129695]Houssa, B.; Schaap, D.; van der Wal, J.; Goto, K.; Kondo,
H.; Yamakawa,

[129696]A.; Shibata, M.; Takenawa, T.; van Blitterswijk, W. J.:
Cloning of

[129697]a novel human diacylglycerol kinase (DGK–theta) contain-
ing three cysteine–rich

[129698]domains, a proline–rich region, and a pleckstrin homology
domain with

[129699]an overlapping Ras–associating domain. J. Biol. Chem.
272: 10422–10428,

[129700]1997.

[129701]

[129702]

[129703]11466.

[129704]Kozak, C. A.; Stephenson, D. A.: Encyclopedia of the
mouse genome

[129705]III. October 1993. Mouse chromosome 5. Mammalian
Genome (Suppl.) 4:

[129706]S72–S87, 1993.

[129707]

[129708]

[129709]11467.

[129710]Hahm, K.; Kim, G.; Turck, C. W.; Smale, S. T.: Isolation of a

[129711]murine gene encoding a nucleic acid–binding protein with
homology

[129712]to hnRNP K. Nucleic Acids Res. 21: 3894 only, 1993.

[129713]

[129714]

[129715]11468.

[129716]Leffers, H.; Dejgaard, K.; Celis, J. E.: Characterisation of
two

[129717]major cellular poly(rC)–binding human proteins, each
containing three

[129718]K–homologous (KH) domains. Europ. J. Biochem. 230:

447–453, 1995.

[129719]

[129720]

[129721]11469.

[129722]Hahm, K.; Kim, G.; Turck, C.; Smale, S. T.: Isolation of a
murine

[129723]gene encoding a nucleic acid-binding protein with homol-
ogy to hnRNP

[129724]K. Nucleic Acids Res. 21: 21–26, 1993.

[129725]

[129726]

[129727]11470.

[129728]Ostareck-Lederer, A.; Ostareck, D. H.; Hentze, M. W.: Cy-
toplasmic

[129729]regulatory functions of the KH-domain proteins hnRNPs K
and E1/E2. Trends

[129730]Biochem. Sci. 23: 409–411, 1998.

[129731]

[129732]

[129733]11471.

[129734]Bouchard, M. J.; Wang, L.-H.; Schneider, R. J.: Calcium signaling

[129735]by HBx protein in hepatitis B virus DNA replication. Science 294:

[129736]2376–2378, 2001.

[129737]

[129738]

[129739]11472.

[129740]Calalb, M. B.; Polte, T. R.; Hanks, S. K.: Tyrosine phosphorylation

[129741]of focal adhesion kinase at sites in the catalytic domain regulates

[129742]kinase activity: a role for Src family kinases. Molec. Cell. Biol. 15:

[129743]954–963, 1995.

[129744]

[129745]

[129746]11473.

[129747]Ganem, D.: The X files—one step closer to closure. Sci–

ence 294:

[129748]2299–2300, 2001.

[129749]

[129750]

[129751]11474.

[129752]Herzog, H.; Nicholl, J.; Hort, Y. J.; Sutherland, G. R.; Shine,

[129753]J.: Molecular cloning and assignment of FAK2, a novel human focal

[129754]adhesion kinase, to 8p11.2–p22 by nonisotopic in situ hybridization. Genomics 32:

[129755]484–486, 1996.

[129756]

[129757]

[129758]11475.

[129759]Lev, S.; Moreno, H.; Martinez, R.; Canoll, P.; Peles, E.;

Musacchio,

[129760]J. M.; Plowman, G. D.; Rudy, B.; Schlessinger, J.: Protein tyrosine

[129761]kinase PYK2 involved in Ca^{2+} –induced regulation of ion channel and

[129762]MAP kinase functions. Nature 376: 737–745, 1995.

[129763]

[129764]

[129765]11476.

[129766]Manser, E.; Leung, T.; Salihuddin, H.; Tan, L.; Lim, L.: A
non-receptor

[129767]tyrosine kinase that inhibits the GTPase activity of
p21(cdc42). Nature 363:

[129768]364–367, 1993.

[129769]

[129770]

[129771]11477.

[129772]Matsuya, M.; Sasaki, H.; Aoto, H.; Mitaka, T.; Nagura, K.;
Ohba,

[129773]T.; Ishino, M.; Takahashi, S.; Suzuki, R.; Sasaki, T.: Cell
adhesion

[129774]kinase beta forms a complex with a new member, Hic-5,
of proteins

[129775]localized at focal adhesions. J. Biol. Chem. 273:
1003–1014, 1998.

[129776]

[129777]

[129778]11478.

[129779]Aubry, F.; Mattei, M.-G.; Barque, J.-P.; Galibert, F.: Chro-
mosomal

[129780]localization and expression pattern of the RNase L in-
hibitor gene. FEBS

[129781]Lett. 381: 135-139, 1996.

[129782]

[129783]

[129784]11479.

[129785]Diriong, S.; Salehzada, T.; Bisbal, C.; Martinand, C.; Tavi-
aux,

[129786]S.: Localization of the ribonuclease L inhibitor gene
(RNS4I), a

[129787]new member of the interferon-regulated 2-5A pathway,
to 4q31 by fluorescence

[129788]in situ hybridization. Genomics 32: 488-490, 1996.

[129789]

[129790]

[129791]11480.

[129792]Luo, Z. G.; Wang, Q.; Zhou, J. Z.; Wang, J.; Lou, Z.; Liu, M.;

[129793]He, X.; Wynshaw-Boris, A.; Xiong, W. C.; Lu, B.; Mei, L.:

Regulation

[129794]of AChR clustering by Dishevelled interacting with MuSK

and PAK1. Neuron 35:

[129795]489–505, 2002.

[129796]

[129797]

[129798]11481.

[129799]Valenzuela, D. M.; Stitt, T. N.; DiStefano, P. S.; Rojas, E.;

Mattsson,

[129800]K.; Compton, D. L.; Nunez, L.; Park, J. S.; Stark, J. L.; Gies,

D.

[129801]R.; Thomas, S.; LeBeau, M. M.; Fernald, A. A.; Copeland, N.

G.; Jenkins,

[129802]N. A.; Burden, S. J.; Glass, D. J.; Yancopoulos, G. D.: Re-

ceptor

[129803]tyrosine kinase specific for the skeletal muscle lineage:

expression

[129804]in embryonic muscle, at the neuromuscular junction, and

after injury. Neuron 15:

[129805]573–584, 1995.

[129806]

[129807]

[129808]11482.

[129809]Meyer, J.; Wirth, J.; Held, M.; Schempp, W.; Scherer, G.:
SOX20,

[129810]a new member of the SOX gene family, is located on chromosome 17p13. Cytogenet.

[129811]Cell Genet. 72: 246–249, 1996.

[129812]

[129813]

[129814]11483.

[129815]Vujic, M.; Rajic, T.; Goodfellow, P. N.; Stevanovic, M.: cDNA

[129816]characterization and high resolution mapping of the human SOX20 gene. Mammalian

[129817]Genome 9: 1059–1061, 1998.

[129818]

[129819]

[129820]11484.

[129821]Ide, H.; Saito-Ohara, F.; Ohnami, S.; Osada, Y.; Ikeuchi, T.;
Yoshida,

[129822]T.; Terada, M.: Assignment of the BMPR1A and BMPR1B
genes to human

[129823]chromosome 10q22.3 and 4q23-q24 by in situ hybridiza-
tion and radiation

[129824]hybrid mapping. Cytogenet. Cell Genet. 81: 285-286,
1998.

[129825]

[129826]

[129827]11485.

[129828]Schuffenhauer, S.; Lichtner, P.; Peykar-Derakhshandeh, P.;
Murken,

[129829]J.; Haas, O. A.; Back, E.; Wolff, G.; Zabel, B.; Barisic, I.;
Rauch,

[129830]A.; Borochowitz, Z.; Dallapiccola, B.; Ross, M.; Meitinger,
T.: Deletion

[129831]mapping on chromosome 10p and definition of a critical
region for

[129832]the second DiGeorge syndrome locus (DGS2). Europ. J.
Hum. Genet. 6:

[129833]213–225, 1998.

[129834]

[129835]

[129836]11486.

[129837]Pizzuti, A.; Amati, F.; Calabrese, G.; Mari, A.; Colosimo, A;
Silani,

[129838]V.; Giardino, L.; Ratti, A.; Penso, D.; Calza, L.; Palka, G.;
Scarlato,

[129839]G.; Novelli, G.; Dallapiccola, B.: cDNA characterization and
chromosomal

[129840]mapping of two human homologs of the *Drosophila* di-
shevelled polarity

[129841]gene. *Hum. Molec. Genet.* 5: 953–958, 1996.

[129842]

[129843]

[129844]11487.

[129845]Semenov, M. V.; Snyder, M.: Human dishevelled genes
constitute

[129846]a DHR-containing multigene family. *Genomics* 42:
302–310, 1997.

[129847]

[129848]

[129849]11488.

[129850]Lee, J.; Hahn, Y.; Yun, J. H.; Mita, K.; Chung, J. H.: Characterization

[129851]of JDP genes, an evolutionarily conserved J domain-only protein family,

[129852]from human and moths. Biochim. Biophys. Acta 1491: 355–363, 2000.

[129853]

[129854]

[129855]11489.

[129856]Qiu, J.; Qian, Y.; Chen, V.; Guan, M.-X.; Shen, B.: Human exonuclease

[129857]1 functionally complements its yeast homologues in DNA recombination,

[129858]RNA primer removal, and mutation avoidance. J. Biol. Chem. 274:

[129859]17893–17900, 1999.

[129860]

[129861]

[129862]11490.

[129863]Schmutte, C.; Marinescu, R. C.; Sadoff, M. M.; Guerrette, S.; Overhauser,

[129864]J.; Fishel, R.: Human exonuclease I interacts with the mismatch repair

[129865]protein hMSH2. Cancer Res. 58: 4537–4542, 1998.

[129866]

[129867]

[129868]11491.

[129869]Tishkoff, D. X.; Amin, N. S.; Viars, C. S.; Arden, K. C.; Kolodner,

[129870]R. D.: Identification of a human gene encoding a homologue of *Saccharomyces*

[129871]*cerevisiae* EXO1, an exonuclease implicated in mismatch repair and

[129872]recombination. Cancer Res. 58: 5027–5031, 1998.

[129873]

[129874]

[129875]11492.

[129876]Wilson, D. M., III; Carney, J. P.; Coleman, M. A.; Adamson, A.

[129877]W.; Christensen, M.; Lamerdin, J. E.: Hex1: a new human Rad2 nuclease

[129878]family member with homology to yeast exonuclease 1. Nucleic Acids

[129879]Res. 26: 3762–3768, 1998.

[129880]

[129881]

[129882]11493.

[129883]Carson–Walter, E. B.; Watkins, D. N.; Nanda, A.; Vogel–stein, B.;

[129884]Kinzler, K. W.; St. Croix, B.: Cell surface tumor endothelial markers

[129885]are conserved in mice and humans. Cancer Res. 61: 6649–6655, 2001.

[129886]

[129887]

[129888]11494.

[129889]Christian, S.; Ahorn, H.; Koehler, A.; Eisenhaber, F.; Rodi, H.–P.;

[129890]Garin-Chesa, P.; Park, J. E.; Rettig, W. J.; Lenter, M. C.:

Molecular

[129891]cloning and characterization of endosialin, a C-type

lectin-like cell

[129892]surface receptor of tumor endothelium. J. Biol. Chem. 276:

7408-7414,

[129893]2001.

[129894]

[129895]

[129896]11495.

[129897]St. Croix, B.; Rago, C.; Velculescu, V.; Traverso, G.; Ro-

mans,

[129898]K. E.; Montgomery, E.; Lal, A.; Riggins, G. J.; Lengauer, C.;

Vogelstein,

[129899]B.; Kinzler, K. W.: Genes expressed in human tumor en-

dothelium. Science 289:

[129900]1197-1202, 2000.

[129901]

[129902]

[129903]11496.

[129904]Gosling, J.; Dairaghi, D. J.; Wang, Y.; Hanley, M.; Talbot, D.;

[129905]Miao, Z.; Schall, T. J.: Cutting edge: identification of a novel

[129906]chemokine receptor that binds dendritic cell- and T cell- active chemokines

[129907]including ELC, SLC, and TECK. J. Immun. 164: 2851-2856, 2000.

[129908]

[129909]

[129910]11497.

[129911]Cohn, R. D.; Campbell, K. P.: Molecular basis of muscular dystrophies. Muscle

[129912]Nerve 23: 1456-1471, 2000.

[129913]

[129914]

[129915]11498.

[129916]Gillespie, C. S.; Lee, M.; Fantes, J. F.; Brophy, P. J.: The gene

[129917]encoding the Schwann cell protein periaxin localizes on mouse chromosome

[129918]7 (Prx). Genomics 41: 297–298, 1997.

[129919]

[129920]

[129921]11499.

[129922]Gillespie, C. S.; Sherman, D. L.; Fleetwood–Walker, S. M.;
Cottrell,

[129923]D. F.; Tait, S.; Garry, E. M.; Wallace, V. C. J.; Ure, J.; Grif-
fiths,

[129924]I. R.; Smith, A. Brophy, P. J.: Peripheral demyelination and
neuropathic

[129925]pain behavior in periaxin–deficient mice. Neuron 26:
523–531, 2000.

[129926]

[129927]

[129928]11500.

[129929]Schuler, G. D.: Sequence mapping by electronic PCR.
Genome Res. 7:

[129930]541–550, 1997.

[129931]

[129932]

[129933]11501.

[129934]Takashima, H.; Boerkoel, C. F.; De Jonghe, P.; Ceuterick, C.; Martin,

[129935]J.-J.; Voit, T.; Schroder, J.-M.; Williams, A.; Brophy, P. J.; Timmerman,

[129936]V.; Lupski, J. R.: Periaxin mutations cause a broad spectrum of demyelinating

[129937]neuropathies. Ann. Neurol. 51: 709–715, 2002.

[129938]

[129939]

[129940]11502.

[129941]Liu, Q.-Y.; Wang, L. F.; Miao, S. Y.; Catterall, J. F.: Expression

[129942]and characterization of a novel human sperm membrane protein. Biol.

[129943]Reprod. 54: 323–330, 1996.

[129944]

[129945]

[129946]11503.

[129947]Miao, S.; Yan, Y.; Li, Y.; Bai, Y.; Wei, S.; Zong, C.; Zhao, M.;

[129948]Zong, S.; Wang, L.: cDNA encoding a human sperm membrane protein

[129949]BS-84. Prog. Natural Sci. 5: 119-122, 1995.

[129950]

[129951]

[129952]11504.

[129953]Wang, H.; Miao, S.; Chen, D.; Wang, L.; Koide, S. S.: Assignment

[129954]of chromosomal locus and evidence for alternatively spliced mRNAs

[129955]of a human sperm membrane protein (hSMP-1). Biochim. Biophys. Acta 1447:

[129956]119-124, 1999.

[129957]

[129958]

[129959]11505.

[129960]Horie, M.; Mitsumoto, Y.; Kyushiki, H.; Kanemoto, N.; Watanabe,

[129961]A.; Taniguchi, Y.; Nishino, N.; Okamoto, T.; Kondo, M.; Mori, T.;

[129962]Noguchi, K.; Nakamura, Y.; Takahashi, E.; Tanigami, A.:

Identification

[129963]and characterization of TMEFF2, a novel survival factor for
hippocampal

[129964]and mesencephalic neurons. Genomics 67: 146–152,
2000.

[129965]

[129966]

[129967]11506.

[129968]Uchida, T.; Wada, K.; Akamatsu, T.; Yonezawa, M.;

Noguchi, H.;

[129969]Mizoguchi, A.; Kasuga, M.; Sakamoto, C.: A novel epider-
mal growth

[129970]factor-like molecule containing two follistatin modules
stimulates

[129971]tyrosine phosphorylation of erbB-4 in MKN28 gastric can-
cer cells. Biochem.

[129972]Biophys. Res. Commun. 266: 593–602, 1999.

[129973]

[129974]

[129975]11507.

[129976]Young, J.; Biden, K. G.; Simms, L. A.; Huggard, P.; Kar-
matic,

[129977]R.; Eyre, H. J.; Sutherland, G. R.; Herath, N.; Barker, M.;
Anderson,

[129978]G. J.; Fitzpatrick, D. R.; Ramm, G. A.; Jass, J. R.; Leggett, B.
A.

[129979]: HPP1: a transmembrane protein–encoding gene com-
monly methylated

[129980]in colorectal polyps and cancers. Proc. Nat. Acad. Sci. 98:
265–270,

[129981]2001.

[129982]

[129983]

[129984]11508.

[129985]Blasina, A.; de Weyer, I. V.; Laus, M. C.; Luyten, W. H.;
Parker,

[129986]A. E.; McGowan, C. H.: A human homologue of the check-
point kinase

[129987]Cds1 directly inhibits Cdc25 phosphatase. Curr. Biol. 14:
1–10,

[129988]1999.

[129989]

[129990]

[129991]11509.

[129992]Brown, A. L.; Lee, C.-H.; Schwarz, J. K.; Mitiku, N.; Piwnica-Worms,

[129993]H.; Chung, J. H.: A human Cds1-related kinase that functions downstream

[129994]of ATM protein in the cellular response to DNA damage. Proc. Nat.

[129995]Acad. Sci. 96: 3745–3750, 1999.

[129996]

[129997]

[129998]11510.

[129999]Chaturvedi, P.; Eng, W. K.; Zhu, Y.; Mattern, M. R.; Mishra, R.;

[130000]Hurle, M. R.; Zhang, X.; Annan, R. S.; Lu, Q.; Faucette, L. F.; Scott,

[130001]G. F.; Li, X.; Carr, S. A.; Johnson, R. K.; Winkler, J. D.; Zhou,

[130002]B. B.: Mammalian Chk2 is a downstream effector of the ATM-dependent

[130003]DNA damage checkpoint pathway. Oncogene 18:
4047–4054, 1999.

[130004]

[130005]

[130006]11511.

[130007]Chehab, N. H.; Malikzay, A.; Appel, M.; Halazonetis, T. D.:
Chk2/hCds1

[130008]functions as a DNA damage checkpoint in G–1 by stabiliz–
ing p53. Genes

[130009]Dev. 14: 278–288, 2000.

[130010]

[130011]

[130012]11512.

[130013]Ino, Y.; Wahrer, D. C. R.; Bell, D. W.; Haber, D. A.; Louis, D.

[130014]N.: Mutation analysis of the hCHK2 gene in primary hu–
man malignant

[130015]gliomas. (Letter) Neurogenetics 3: 45–46, 2000.

[130016]

[130017]

[130018]11513.

[130019]Lee, S. B.; Kim, S. H.; Bell, D. W.; Wahrer, D. C. R.; Schiripo,

[130020]T. A.; Jorczak, M. M.; Sgroi, D. C.; Garber, J. E.; Li, F. P.;

Nichols,

[130021]K. E.; Varley, J. M.; Godwin, A. K.; Shannon, K. M.; Harlow,

E.; Haber,

[130022]D. A.: Destabilization of CHK2 by a missense mutation associated
sociated

[130023]with Li-Fraumeni syndrome. Cancer Res. 61: 8062-8067,
2001.

[130024]

[130025]

[130026]11514.

[130027]Lopes, M.; Cotta-Ramusino, C.; Pelliccioli, A.; Liberi, G.; Plevani,
evani,

[130028]P.; Muzi-Falconi, M. Newlon, C. S.; Foiani, M.: The DNA
replication

[130029]checkpoint response stabilizes stalled replication forks.
Nature 412:

[130030]557-561, 2001.

[130031]

[130032]

[130033]11515.

[130034]Matsuoka, S.; Huang, M.; Elledge, S. J.: Linkage of ATM to
cell

[130035]cycle regulation by the Chk2 protein kinase. Science 282:
1893–1897,

[130036]1998.

[130037]

[130038]

[130039]11516.

[130040]Meijers–Heijboer, H.; van den Ouweland, A.; Klijn, J.;
Wasielewski,

[130041]M.; de Snoo, A.; Oldenburg, R.; Hollestelle, A.; Houben,
M.; Crepin,

[130042]E.; van Veghel–Plandsoen, M.; Elstrodt, F.; van Duijn, C.;
and 29

[130043]others: Low–penetrance susceptibility to breast cancer
due to CHEK2*1100delC

[130044]in noncarriers of BRCA1 or BRCA2 mutations. Nature
Genet. 31: 55–59,

[130045]2002.

[130046]

[130047]

[130048]11517.

[130049]Miller, C. W.; Ikezoe, T.; Krug, U.; Hofmann, W.-K.; Tavor, S.;

[130050]Vegesna, V.; Tsukasaki, K.; Takeuchi, S.; Koeffler, H. P.:
Mutations

[130051]of the CHK2 gene are found in some osteosarcomas, but
are rare in

[130052]breast, lung, and ovarian tumors. Genes Chromosomes
Cancer 33: 17–21,

[130053]2002.

[130054]

[130055]

[130056]11518.

[130057]Sogo, J. M.; Lopes, M.; Foiani, M.: Fork reversal and ssDNA
accumulation

[130058]at stalled replication forks owing to checkpoint defects.
Science 297:

[130059]599–602, 2002.

[130060]

[130061]

[130062]11519.

[130063]Vahteristo, P.; Bartkova, J.; Eerola, H.; Syrjakoski, K.; Ojala,

[130064]S.; Kilpivaara, O.; Tamminen, A.; Kononen, J.; Aittomaki,
K.; Heikkila,

[130065]P.; Holli, K.; Blomqvist, C.; Bartek, J.; Kallioniemi, O.-P.;
Nevanlinna,

[130066]H.: A CHEK2 genetic variant contributing to a substantial
fraction

[130067]of familial breast cancer. Am. J. Hum. Genet. 71: 432–438,
2002.

[130068]

[130069]

[130070]11520.

[130071]Vahteristo, P.; Tamminen, A.; Karvinen, P.; Eerola, H.; Ek-
lund,

[130072]C.; Aaltonen, L. A.; Blomqvist, C.; Aittomaki, K.; Nevan-
linna, H.

[130073]: p53, CHK2, and CHK1 genes in Finnish families with Li-
Fraumeni syndrome:

[130074]further evidence of CHK2 in inherited cancer predisposition. Cancer

[130075]Res. 61: 5718–5722, 2001.

[130076]

[130077]

[130078]11521.

[130079]De, S. K.; Enders, G. C.; Andrews, G. K.: High levels of metallothionein

[130080]messenger RNAs in male germ cells of the adult mouse. Molec. Endocr. 5:

[130081]628–636, 1991.

[130082]

[130083]

[130084]11522.

[130085]Salehi–Ashtiani, K.; Widrow, R. J.; Markert, C. L.; Goldberg, E.

[130086]: Testis–specific expression of a metallothionein I–driven transgene

[130087]correlates with undermethylation of the locus in testicular DNA. Proc.

[130088]Nat. Acad. Sci. 90: 8886–8890, 1993.

[130089]

[130090]

[130091]11523.

[130092]Sugihara, T.; Wadhwa, R.; Kaul, S. C.; Mitsui, Y.: A novel
testis-specific

[130093]metallothionein-like protein, tesmin, is an early marker of
male germ

[130094]cell differentiation. Genomics 57: 130–136, 1999.

[130095]

[130096]

[130097]11524.

[130098]Larrouy, D.; Vidal, H.; Andreelli, F.; Laville, M.; Langin, D.

[130099]: Cloning and mRNA tissue distribution of human PPAR-
gamma coactivator-1. Int.

[130100]J. Obesity 23: 1327–1332, 1999.

[130101]

[130102]

[130103]11525.

[130104]Lin, J.; Wu, H.; Tarr, P. T.; Zhang, C.-Y.; Wu, Z.; Boss, O.;
Michael,

[130105]L. F.; Puigserver, P.; Isotani, E.; Olson, E. N.; Lowell, B. B.;
Bassel-Duby,

[130106]R.; Spiegelman, B. M.: Transcriptional co-activator PGC-
1-alpha drives

[130107]the formation of slow-twitch muscle fibres. Nature 418:
797-801,

[130108]2002.

[130109]

[130110]

[130111]11526.

[130112]Monsalve, M.; Wu, Z.; Adelmant, G.; Puigserver, P.; Fan,
M.; Spiegelman,

[130113]B. M.: Direct coupling of transcription and mRNA process-
ing through

[130114]the thermogenic coactivator PGC-1. Molec. Cell 6:
307-316, 2000.

[130115]

[130116]

[130117]11527.

- [130118]Puigserver, P.; Adelmant, G.; Wu, Z.; Fan, M.; Xu, J.;
O'Malley,
[130119]B.; Spiegelman, B. M.: Activation of PPAR–gamma coacti-
vator–1 through
[130120]transcription factor docking. Science 286: 1368–1371,
1999.
- [130121]
- [130122]
- [130123]11528.
- [130124]Puigserver, P.; Rhee, J.; Lin, J.; Wu, Z.; Yoon, J. C.; Zhang,
[130125]C.–Y.; Krauss, S.; Mootha, V. K.; Lowell, B. B.; Spiegelman,
B. M.
- [130126]: Cytokine stimulation of energy expenditure through p38
MAP kinase
[130127]activation of PPAR–gamma coactivator–1. Molec. Cell 8:
971–982,
[130128]2001.
- [130129]
- [130130]
- [130131]11529.

[130132]Puigserver, P.; Wu, Z.; Park, C. W.; Graves, R.; Wright, M.; Spiegelman,

[130133]B. M.: A cold-inducible coactivator of nuclear receptors linked to

[130134]adaptive thermogenesis. Cell 92: 829–839, 1998.

[130135]

[130136]

[130137]11530.

[130138]Waite, L. L.; Person, E. C.; Zhou, Y.; Lim, K.-H.; Scanlan, T.

[130139]S.; Taylor, R. N.: Placental peroxisome proliferator-activated receptor-gamma

[130140]is up-regulated by pregnancy serum. J. Clin. Endocr. Metab. 85:

[130141]3808–3814, 2000.

[130142]

[130143]

[130144]11531.

[130145]Wu, Z.; Puigserver, P.; Andersson, U.; Zhang, C.; Adelmant, G.;

[130146]Mootha, V.; Troy, A.; Cinti, S.; Lowell, B.; Scarpulla, R. C.;

Spiegelman,

[130147]B. M.: Mechanisms controlling mitochondrial biogenesis
and respiration

[130148]through the thermogenic coactivator PGC-1. Cell 98:
115-124, 1999.

[130149]

[130150]

[130151]11532.

[130152]Yoon, J. C.; Puigserver, P.; Chen, G.; Donovan, J.; Wu, Z.;
Rhee,

[130153]J.; Adelmant, G.; Stafford, J.; Kahn, C. R.; Granner, D. K.;
Newgard,

[130154]C. B.; Spiegelman, B. M.: Control of hepatic gluconeogen-
esis through

[130155]the transcriptional coactivator PGC-1. Nature 413:
131-138, 2001.

[130156]

[130157]

[130158]11533.

[130159]Granger, S. W.; Butrovich, K. D.; Houshmand, P.; Edwards,
W. R.;

[130160]Ware, C. F.: Genomic characterization of LIGHT reveals linkage to

[130161]an immune response locus on chromosome 19p13.3 and distinct isoforms

[130162]generated by alternate splicing or proteolysis. J. Immun. 167: 5122–5128,

[130163]2001.

[130164]

[130165]

[130166]11534.

[130167]Mauri, D. N.; Ebner, R.; Montgomery, R. I.; Kochel, K. D.; Cheung,

[130168]T. C.; Yu, G. L.; Ruben, S.; Murphy, M.; Eisenberg, R. J.; Cohen,

[130169]G. H.; Spear, P. G.; Ware, C. F.: LIGHT, a new member of the TNF

[130170]superfamily, and lymphotoxin alpha are ligands for herpesvirus entry

[130171]mediator. Immunity 8: 21–30, 1998.

[130172]

[130173]

[130174]11535.

[130175]Harrop, J. A.; McDonnell, P. C.; Brigham–Burke, M.; Lyn, S. D.;

[130176]Minton, J.; Tan, K. B.; Dede, K.; Spampanato, J.; Silverman, C.;

[130177]Hensley, P.; DiPrinzio, R.; Emery, J. G.; Deen, K.; Eichman, C.; Chabot–Fletcher,

[130178]M.; Truneh, A.; Young, P. R.: Herpesvirus entry mediator ligand (HVEM–L),

[130179]a novel ligand for HVEM/TR2, stimulates proliferation of T cells and

[130180]inhibits HT29 cell growth. . J. Biol. Chem. 273: 27548–27556, 1998.

[130181]

[130182]

[130183]11536.

[130184]Hadjantonakis, A.–K.; Formstone, C. J.; Little, P. F. R.: mCelsr1

[130185]is an evolutionarily conserved seven–pass transmembrane receptor and

[130186]is expressed during mouse embryonic development. Mech. Dev. 78:

[130187]91–95, 1998.

[130188]

[130189]

[130190]11537.

[130191]Hadjantonakis, A.–K.; Sheward, W. J.; Harmar, A. J.; de
Galan,

[130192]L.; Hoovers, J. M. N.; Little, P. F. R.: Celsr1, a neural-spe-
cific

[130193]gene encoding an unusual seven-pass transmembrane re-
ceptor, maps to

[130194]mouse chromosome 15 and human chromosome 22qter.
Genomics 45: 97–104,

[130195]1997.

[130196]

[130197]

[130198]11538.

[130199]Hewett–Emmett, D.; Tashian, R. E.: Functional diversity,
conservation,

[130200]and convergence in the evolution of the alpha–, beta–,
and gamma–carbonic

[130201]anhydrase gene families. Molec. Phylogenet. Evol. 5:
50–77, 1996.

[130202]

[130203]

[130204]11539.

[130205]Hewett–Emmett, D.; Wiebauer, K.: Personal Communica-
tion. Houston,

[130206]Texas 9/1999.

[130207]

[130208]

[130209]11540.

[130210]Kleiderlein, J. J.; Nisson, P. E.; Jessee, J.; Li, W.–B.; Becker,

[130211]K. G.; Derby, M. L.; Ross, C. A.; Margolis, R. L.: CCG re-
peats in

[130212]cDNAs from human brain. Hum. Genet. 103: 666–673,
1998.

[130213]

[130214]

[130215]11541.

[130216]Lovejoy, D. A.; Hewett–Emmett, D.; Porter, C. A.; Cepoi, D.; Sheffield,

[130217]A.; Vale, W. W.; Tashian, R. E.: Evolutionarily conserved, 'acatalytic'

[130218]carbonic anhydrase–related protein XI contains a sequence motif present

[130219]in the neuropeptide sauvagine: the human CA–RP XI gene (CA11) is embedded

[130220]between the secretor gene cluster and the DBP gene at 19q13.3. Genomics 54:

[130221]484–493, 1998.

[130222]

[130223]

[130224]11542.

[130225]Bellingham, J.; Greogory–Evans, K.; Gregory–Evans, C. Y.: Sequence

[130226]and tissue expression of a human novel carbonic anhydrase–related

[130227]protein, CARP–2, mapping to chromosome 19q13.3. Biochem. Biophys.

[130228]Res. Comm. 253: 364–367, 1998.

[130229]

[130230]

[130231]11543.

[130232]Fujikawa–Adachi, K.; Nishimori, I.; Taguchi, T.; Yuri, K.;
Onishi,

[130233]S.: cDNA sequence, mRNA expression, and chromosomal
localization

[130234]of human carbonic anhydrase–related protein, CA–RP XI.
Biochim. Biophys.

[130235]Acta 1431: 518–524, 1999.

[130236]

[130237]

[130238]11544.

[130239]Lindsay, M. E.; Plafker, K.; Smith, A. E.; Clurman, B. E.;
Macara,

[130240]I. G.: Npap60/Nup50 is a tri–stable switch that stimulates
importin–alpha:beta–mediated

[130241]nuclear protein import. Cell 110: 349–360, 2002.

[130242]

[130243]

[130244]11545.

[130245]Trichet, V.; Shkolny, D.; Dunham, I.; Beare, D.; McDermid, H. E.

[130246]: Mapping and complex expression pattern of the human NPAP60L nucleoporin

[130247]gene. Cytogenet. Cell Genet. 85: 221–226, 1999.

[130248]

[130249]

[130250]11546.

[130251]Martin, L.; Fanarraga, M. L.; Aloria, K.; Zabala, J. C.: Tubulin

[130252]folding cofactor D is a microtubule destabilizing protein. FEBS Lett. 470:

[130253]93–95, 2000.

[130254]

[130255]

[130256]11547.

[130257]de Veer, M. J.; Sim, H.; Whisstock, J. C.; Devenish, R. J.; Ralph,

[130258]S. J.: IFI60/ISG60/IFIT4, a new member of the human

IFI54/IFIT2 family

[130259]of interferon-stimulated genes. Genomics 54: 267-277, 1998.

[130260]

[130261]

[130262]11548.

[130263]Yu, M.; Tong, J.-H.; Mao, M.; Kan, L.-X.; Liu, M.-M.; Sun, Y.-W.;

[130264]Fu, G.; Jing, Y.-K.; Yu, L.; Lepaslier, D.; Lanotte, M.; Wang, Z.-Y.;

[130265]Chen, Z.; Waxman, S.; Wang, Y.-X.; Tan, J.-Z.; Chen, S.-J.: Cloning

[130266]of a gene (RIG-G) associated with retinoic acid-induced differentiation

[130267]of acute promyelocytic leukemia cells and representing a new member

[130268]of a family of interferon-stimulated genes. Proc. Nat. Acad. Sci. 94:

[130269]7406-7411, 1997.

[130270]

[130271]

[130272]11549.

[130273]Zhu, H.; Cong, J.-P.; Shenk, T.: Use of differential display analysis

[130274]to assess the effect of human cytomegalovirus infection on the accumulation

[130275]of cellular RNAs: induction of interferon-responsive RNAs. Proc.

[130276]Nat. Acad. Sci. 94: 13985–13990, 1997.

[130277]

[130278]

[130279]11550.

[130280]Hirai, H.; Tanaka, K.; Takano, S.; Ichimasa, M.; Nakamura, M.;

[130281]Nagata, K.: Cutting edge: agonistic effect of indomethacin on a prostaglandin

[130282]D2 receptor, CRTH2. J. Immun. 168: 981–985, 2002.

[130283]

[130284]

[130285]11551.

[130286]Nagata, K.; Tanaka, K.; Ogawa, K.; Kemmotsu, K.; Imai,

Yoshie,

[130287]O.; Abe, H.; Tada, K.; Nakamura, M.; Sugamura, K.;

Takano, S.: Selective

[130288]expression of a novel surface molecule by human Th2
cells in vivo. J.

[130289]Immun. 162: 1278–1286, 1999.

[130290]

[130291]

[130292]11552.

[130293]Grundemann, D.; Schechinger, B.; Rappold, G. A.;

Schomig, E.:

[130294]Molecular identification of the corticosterone–sensitive
extraneuronal

[130295]catecholamine transporter. Nature Neurosci. 1: 349–351,
1998.

[130296]

[130297]

[130298]11553.

[130299]Boettger, T.; Hubner, C. A.; Maler, H.; Rust, M. B.; Beck, F.
X.;

[130300]Jentsch, T. J.: Deafness and renal tubular acidosis in mice

lacking

[130301]the K-Cl co-transporter Kcc4. Nature 416: 874-878, 2002.

[130302]

[130303]

[130304]11554.

[130305]Kotake, K.; Ozaki, N.; Mizuta, M.; Sekiya, S.; Inagaki, N.; Seino,

[130306]S.: Noc2, a putative zinc finger protein involved in exocytosis in

[130307]endocrine cells. J. Biol. Chem. 272: 29407-29410, 1997.

[130308]

[130309]

[130310]11555.

[130311]Smith, J. S.; Tachibana, I.; Allen, C.; Chiappa, S. A.; Lee, H.

[130312]K.; McIver, B.; Jenkins, R. B.; Raffel, C.: Cloning of a human ortholog

[130313](RPH3AL) of (RNO)Rph3al from a candidate 17p13.3 medulloblastoma tumor

[130314]suppressor locus. Genomics 59: 97-101, 1999.

[130315]

[130316]

[130317]11556.

[130318]Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.;

Kawarabayasi,

[130319]Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.:

Prediction

[130320]of the coding sequences of unidentified human genes. I.

The coding

[130321]sequences of 40 new genes (KIAA0001–KIAA0040) de-

duced by analysis

[130322]of randomly sampled cDNA clones from human immature

myeloid cell line

[130323]KG-1. DNA Res. 1: 27–35, 1994.

[130324]

[130325]

[130326]11557.

[130327]Kimura, M.; Okano, Y.: Identification and assignment of

the human

[130328]NIMA-related protein kinase 7 gene (NEK7) to human

chromosome 1q31.3. Cytogenet.

[130329]Cell Genet. 94: 33–38, 2001.

[130330]

[130331]

[130332]11558.

[130333]Li, M. Z.; Yu, L.; Liu, Q.; Chu, J. Y.; Zhao, S. Y.: Assignment

[130334]of NEK6, a NIMA–related gene, to human chromosome

9q33.3–q34.11 by

[130335]radiation hybrid mapping. Cytogenet. Cell Genet. 87:

271–272, 1999.

[130336]

[130337]

[130338]11559.

[130339]Favier, D.; Gonda, T. J.: Detection of proteins that bind to
the

[130340]leucine zipper motif of c–Myb. Oncogene 9: 305–311,
1994.

[130341]

[130342]

[130343]11560.

[130344]Keough, R.; Woollatt, E.; Crawford, J.; Sutherland, G. R.;
Plummer,

[130345]S.; Casey, G.; Gonda, T. J.: Molecular cloning and chromo-
somal mapping

[130346]of the human homologue of MYB binding protein (P160)
1A (MYBBP1A)

[130347]to 17p13.3. Genomics 62: 483–489, 1999.

[130348]

[130349]

[130350]11561.

[130351]Tavner, F. J.; Simpson, R.; Tashiro, S.; Favier, D.; Jenkins,
N.

[130352]A.; Gilbert, D. J.; Copeland, N. G.; Macmillan, E. M.;
Lutwyche, J.;

[130353]Keough, R. A.; Ishii, S.; Gonda, T. J.: Molecular cloning re-
veals

[130354]that the p160 Myb-binding protein is a novel, predomi-
nantly nucleolar

[130355]protein which may play a role in transactivation by Myb.
Molec. Cell

[130356]Biol. 18: 989–1002, 1998.

[130357]

[130358]

[130359]11562.

[130360]Gilbert, D. J.; Engel, H.; Wang, X.; Grzeschik, K.-H.;
Copeland,

[130361]N. G.; Jenkins, N. A.; Kilimann, M. W.: The neurobeachin
gene (Nbea)

[130362]identifies a new region of homology between mouse cen-
tral chromosome

[130363]3 and human chromosome 13q13. Mamm. Genome 10:
1030-1031, 1999.

[130364]

[130365]

[130366]11563.

[130367]Nagle, D. L.; Karim, M. A.; Woolf, E. A.; Holmgren, L.; Bork,
P.;

[130368]Misumi, D. J.; McGrail, S. H.; Dussault, B. J., Jr.; Perou, C.
M.;

[130369]Boissy, R. E.; Duyk, G. M.; Spritz, R. A.; Moore, K. J.: Iden-
tification

[130370]and mutation analysis of the complete gene for Chediak-

Higashi syndrome. Nature

[130371]Genet. 14: 307–311, 1996.

[130372]

[130373]

[130374]11564.

[130375]Wang, X.; Herberg, F. W.; Laue, M. M.; Wullner, C.; Hu, B.;
Petrascch–Parwez,

[130376]E.; Kilimann, M, W.: Neurobeachin: a protein kinase A–
anchoring,

[130377]beige/Chediak–Higashi protein homolog implicated in
neuronal membrane

[130378]traffic. J. Neurosci. 20: 8551–8565, 2000.

[130379]

[130380]

[130381]11565.

[130382]Connor, F.; Wright, E.; Denny, P.; Koopman, P.; Ashworth,
A.:

[130383]The Sry–related HMG box–containing gene Sox6 is ex–
pressed in the adult

[130384]testis and developing nervous system of the mouse. Nu–

cleic Acids

[130385]Res. 23: 3365–3372, 1995.

[130386]

[130387]

[130388]11566.

[130389]Wunderle, V. M.; Critcher, R.; Ashworth, A.; Goodfellow, P.
N.

[130390]: Cloning and characterization of SOX5, a new member of
the human

[130391]SOX gene family. Genomics 36: 354–358, 1996.

[130392]

[130393]

[130394]11567.

[130395]Gomez–Escobar, N.; Chou, C.–F.; Lin, W.–W.; Hsieh, S.–L.;
Campbell,

[130396]R. D.: The G11 gene located in the major histocompatibil–
ity complex

[130397]encodes a novel nuclear serine/threonine protein kinase.
J. Biol.

[130398]Chem. 273: 30954–30960, 1998.

[130399]

[130400]

[130401]11568.

[130402]Sargent, C. A.; Anderson, M. J.; Hsieh, S.-L.; Kendall, E.;
Gomez-Escobar,

[130403]N.; Campbell, R. D.: Characterisation of the novel gene
G11 lying

[130404]adjacent to the complement C4A gene in the human major
histocompatibility

[130405]complex. Hum. Molec. Genet. 3: 481-488, 1994.

[130406]

[130407]

[130408]11569.

[130409]Shen, L.; Wu, L.; Sanlioglu, S.; Chen, R.; Mendoza, A. R.;
Dangel,

[130410]A. W.; Carroll, M. C.; Zipf, W. B.; Yu, C. Y.: Structure and
genetics

[130411]of the partially duplicated gene RP located immediately
upstream of

[130412]the complement C4A and the C4B genes in the HLA class
III region:

[130413]molecular cloning, exon–intron structure, composite
retroposon, and

[130414]breakpoint gene duplication. J. Biol. Chem. 269:
8466–8476, 1994.

[130415]

[130416]

[130417]11570.

[130418]Bunn, R. C.; Jensen, M. A.; Reed, B. C.: Protein interactions

[130419]with the glucose transporter binding protein GLUT1CBP
that provide

[130420]a link between GLUT1 and the cytoskeleton. Molec. Biol.
Cell 10:

[130421]819–832, 1999.

[130422]

[130423]

[130424]11571.

[130425]De Vries, L.; Lou, X.; Zhao, G.; Zheng, B.; Farquhar, M. G.:
GIPC,

[130426]a PDZ domain containing protein, interacts specifically
with the C

[130427]terminus of RGS-GAIP. Proc. Nat. Acad. Sci. 95:
12340-12345, 1998.

[130428]

[130429]

[130430]11572.

[130431]Von Kap-Herr, C.; Kandala, G.; Mann, S. S.; Hart, T. C.;
Pettenati,

[130432]M. J.; Setaluri, V.: Assignment of PDZ domain-containing
protein

[130433]GIPC gene (C19orf3) to human chromosome band
19p13.1 by in situ hybridization

[130434]and radiation hybrid mapping. Cytogenet. Cell Genet. 89:
234-235,

[130435]2000.

[130436]

[130437]

[130438]11573.

[130439]Alessi, D. R.; James, S. R.; Downes, C. P.; Holmes, A. B.;
Gaffney,

[130440]P. R. J.; Reese, C. B.; Cohen, P.: Characterization of a
3-phosphoinositide-dependent

[130441]protein kinase which phosphorylates and activates protein
kinase B-alpha. Curr.

[130442]Biol. 7: 261-269, 1997.

[130443]

[130444]

[130445]11574.

[130446]Sandrini, F.; Farmakidis, C.; Kirschner, L. S.; Wu, S.-M.;
Tullio-Pelet,

[130447]A.; Lyonnet, S.; Metzger, D. L.; Bourdony, C. J.; Tiosano,
D.; Chan,

[130448]W.-Y.; Stratakis, C. A.: Spectrum of mutations of the AAAS
gene in

[130449]Allgrove syndrome: lack of mutations in six kindreds with
isolated

[130450]resistance to corticotropin. J. Clin. Endocr. Metab. 86:
5433-5437,

[130451]2001.

[130452]

[130453]

[130454]11575.

[130455]Ludwig, D.; Lorenz, J.; Dejana, E.; Bohlen, P.; Hicklin, D. J.;

[130456]Witte, L.; Pytowski, B.: cDNA cloning, chromosomal mapping, and expression

[130457]analysis of human VE-cadherin-2. Mammalian Genome
11: 1030-1033,

[130458]2000.

[130459]

[130460]

[130461]11576.

[130462]Telo, P.; Breviario, F.; Huber, P.; Panzeri, C.; Dejana, E.:
Identification

[130463]of a novel cadherin (vascular endothelial cadherin-2) located at intercellular

[130464]junctions in endothelial cells. J. Biol. Chem. 273:
17565-17572,

[130465]1998.

[130466]

[130467]

[130468]11577.

[130469]Adams, J. C.; Seed, B.; Lawler, J.: Muskelin, a novel intra-cellular

[130470]mediator of cell adhesive and cytoskeletal responses to thrombospondin-1. EMBO

[130471]J. 17: 4964-4974, 1998.

[130472]

[130473]

[130474]11578.

[130475]Adams, J. C.; Zhang, L.: cDNA cloning of human muskelin and localisation

[130476]of the muskelin (MKLN1) gene to human chromosome 7q32 and mouse chromosome

[130477]6B1/B2 by physical mapping and FISH. Cytogenet. Cell Genet. 87:

[130478]19-21, 1999.

[130479]

[130480]

[130481]11579.

[130482]Moynihan, T. P.; Ardley, H. C.; Nuber, U.; Rose, S. A.;

Jones,

[130483]P. F.; Markham, A. F.; Scheffner, M.; Robinson, P. A.: The
ubiquitin-conjugating

[130484]enzymes Ubch7 and Ubch8 interact with RING finger/IBR
motif-containing

[130485]domains of HHARI and H7-AP1. J. Biol. Chem. 274:
30963-30968, 1999.

[130486]

[130487]

[130488]11580.

[130489]Tan, N. G. S.; Ardley, H. C.; Rose, S. A.; Leek, J. P.;

Markham,

[130490]A. F.; Robinson, P. A.: Characterisation of the human and
mouse orthologues

[130491]of the Drosophila ariadne gene. Cytogenet. Cell Genet. 90:
242-245,

[130492]2000.

[130493]

[130494]

[130495]11581.

[130496]Janssen, J. W. G.; Imoto, I.; Inoue, J.; Shimada, Y.; Ueda, M.;

[130497]Imamura, M.; Bartram, C. R.; Inazawa, J.: MYEOV, a gene at 11q13,

[130498]is coamplified with CCND1, but epigenetically inactivated in a subset

[130499]of esophageal squamous cell carcinomas. J. Hum. Genet. 47: 460–464,

[130500]2002.

[130501]

[130502]

[130503]11582.

[130504]Janssen, J. W. G.; Vaandrager, J.–W.; Heuser, T.; Jauch, A.; Kluin,

[130505]P. M.; Geelen, E.; Bergsagel, P. L.; Kuehl, W. M.; Drexler, H. G.;

[130506]Otsuki, T.; Bartram, C. R.; Schuuring, E.: Concurrent activation

[130507]of a novel putative transforming gene, myeov, and cyclin D1 in a subset

[130508]of multiple myeloma cell lines with t(11;14)(q13;q32).

Blood 95:

[130509]2691–2698, 2000.

[130510]

[130511]

[130512]11583.

[130513]Di Cunto, F.; Calautti, E.; Hsiao, J.; Ong, L.; Topley, G.;

Turco,

[130514]E.; Dotto, G. P.: Citron Rho–interacting kinase, a novel tissue-specific

[130515]ser/thr kinase encompassing the Rho–Rac–binding protein citron. J.

[130516]Biol. Chem. 273: 29706–29711, 1998.

[130517]

[130518]

[130519]11584.

[130520]Di Cunto, F.; Imarisio, S.; Hirsch, E.; Broccoli, V.; Bulfone,

[130521]A.; Migheli, A.; Atzori, C.; Turco, E.; Triolo, R.; Dotto, G. P.;

[130522]Silengo, L.; Altruda, F.: Defective neurogenesis in citron

kinase

[130523]knockout mice by altered cytokinesis and massive apoptosis. Neuron 28:

[130524]115–127, 2000.

[130525]

[130526]

[130527]11585.

[130528]Madaule, P.; Furuyashiki, T.; Reid, T.; Ishizaki, T.; Watanabe,

[130529]G.; Morii, N.; Narumiya, S.: A novel partner for the GTP-bound forms

[130530]of rho and rac. FEBS Lett. 377: 243–248, 1995.

[130531]

[130532]

[130533]11586.

[130534]Ishizaki, J.; Suzuki, N.; Higashino, K.; Yokota, Y.; Ono, T.; Kawamoto,

[130535]K.; Fujii, N.; Arita, H.; Hanasaki, K.: Cloning and characterization

[130536]of novel mouse and human secretory phospholipase A(2)s.

J. Biol.

[130537]Chem. 274: 24973–24979, 1999.

[130538]

[130539]

[130540]11587.

[130541]Baudat, F.; Manova, K.; Yuen, J. P.; Jasin, M.; Keeney, S.:

Chromosome

[130542]synapsis defects and sexually dimorphic meiotic progression in mice

[130543]lacking Spo11. Molec. Cell 6: 989–998, 2000.

[130544]

[130545]

[130546]11588.

[130547]Romanienko, P. J.; Camerini–Otero, R. D.: The mouse

Spo11 gene

[130548]is required for meiotic chromosome synapsis. Molec. Cell 6: 975–987,

[130549]2000.

[130550]

[130551]

[130552]11589.

[130553]Romanienko, P. J.; Camerini-Otero, R. D.: Cloning, characterization,

[130554]and localization of mouse and human SPO11. Genomics
61: 156–169,

[130555]1999.

[130556]

[130557]

[130558]11590.

[130559]Shannon, M.; Richardson, L.; Christian, A.; Handel, M. A.;
Thelen,

[130560]M. P.: Differential gene expression of mammalian
SPO11/TOP6A homologs

[130561]during meiosis. FEBS Lett. 462: 329–334, 1999.

[130562]

[130563]

[130564]11591.

[130565]Lopez-Coviella, I.; Berse, B.; Krauss, R.; Thies, R. S.; Blusz-

tajn,

[130566]J. K.: Induction and maintenance of the neuronal cholinergic phenotype

[130567]in the central nervous system by BMP-9. Science 289: 313-316, 2000.

[130568]

[130569]

[130570]11592.

[130571]Burn, T. C.; Connors, T. D.; Van Raay, T. J.; Dackowski, W. R.;

[130572]Millholland, J. M.; Klinger, K. W.; Landes, G. M.: Generation of

[130573]a transcriptional map for a 700-kb region surrounding the polycystic

[130574]kidney disease type 1 (PKD1) and tuberous sclerosis type 2 (TSC2)

[130575]disease genes on human chromosome 16p13.3. Genome Res. 6: 525-537,

[130576]1996.

[130577]

[130578]

[130579]11593.

[130580]Pullen, N.; Dennis, P. B.; Andjelkovic, M.; Dufner, A.;
Kozma,

[130581]S. C.; Hemmings, B. A.; Thomas, G.: Phosphorylation and
activation

[130582]of p70(s6k) by PDK1. Science 279: 707–710, 1998.

[130583]

[130584]

[130585]11594.

[130586]Stephens, L.; Anderson, K.; Stokoe, D.; Erdjument–Bro–
mage, H.;

[130587]Painter, G. F.; Holmes, A. B.; Gaffney, P. R. J.; Reese, C. B.;
McCormick,

[130588]F.; Tempst, P.; Coadwell, J.; Hawkins, P. T.: Protein kinase
B kinases

[130589]that mediate phosphatidylinositol

3,4,5–triphosphate–dependent activation

[130590]of protein kinase B. Science 279: 710–714, 1998.

[130591]

[130592]

[130593]11595.

[130594]Uebele, V. N.; Lagrutta, A.; Wade, T.; Figueroa, D. J.; Liu, Y.;

[130595]McKenna, E.; Austin, C. P.; Bennett, P. B.; Swanson, R.:
Cloning

[130596]and functional expression of two families of beta-
subunits of the

[130597]large conductance calcium-activated K(+) channel. J. Biol.
Chem. 275:

[130598]23211-23218, 2000.

[130599]

[130600]

[130601]11596.

[130602]Kouroku, Y.; Soyama, A.; Fujita, E.; Urase, K.; Tsukahara, T.;

[130603]Momoi, T.: RA70 is a src kinase-associated protein expressed ubiquitously. Biochem.

[130604]Biophys. Res. Comm. 252: 738-742, 1998.

[130605]

[130606]

[130607]11597.

[130608]Liu, J.; Kang, H.; Raab, M.; da Silva, A. J.; Kraeft, S.-K.;
Rudd,

[130609]C. E.: FYB (FYN binding protein) serves as a binding part-
ner for

[130610]lymphoid protein and FYN kinase substrate SKAP55 and a
SKAP55-related

[130611]protein in T cells. Proc. Nat. Acad. Sci. 95: 8779–8784,
1998.

[130612]

[130613]

[130614]11598.

[130615]Marie-Cardine, A.; Verhagen, A. M.; Eckerskorn, C.;
Schraven, B.

[130616]: SKAP-HOM, a novel adaptor protein homologous to the
FYN-associated

[130617]protein SKAP55. FEBS Lett. 435: 55–60, 1998.

[130618]

[130619]

[130620]11599.

[130621]Chai, J.; Du, C.; Wu, J.-W.; Kyin, S.; Wang, X.; Shi, Y.:

Structural

[130622]and biochemical basis of apoptotic activation by Smac/

DIABLO. Nature 406:

[130623]855–862, 2000.

[130624]

[130625]

[130626]11600.

[130627]Du, C.; Fang, M.; Li, Y.; Li, L.; Wang, X.: Smac, a mitochondrial

[130628]protein that promotes cytochrome c-dependent caspase activation by

[130629]eliminating IAP inhibition. Cell 102: 33–42, 2000.

[130630]

[130631]

[130632]11601.

[130633]Okada, H.; Suh, W.-K.; Jin, J.; Woo, M.; Du, C.; Elia, A.;
Duncan,

[130634]G. S.; Wakeham, A.; Itie, A.; Lowe, S. W.; Wang, X.; Mak, T.
W.:

[130635]Generation and characterization of Smac/DIA-
BLO-deficient mice. Molec.

[130636]Cell. Biol. 22: 3509–3517, 2002.

[130637]

[130638]

[130639]11602.

[130640]Scott, A. F.: Personal Communication. Baltimore, Md.
8/18/2000.

[130641]

[130642]

[130643]11603.

[130644]Verhagen, A. M.; Ekert, P. G.; Pakusch, M.; Silke, J.; Con-
nolly,

[130645]L. M.; Reid, G. E.; Moritz, R. L.; Simpson, R. J.; Vaux, D. L.:
Identification

[130646]of DIABLO, a mammalian protein that promotes apoptosis
by binding

[130647]to and antagonizing IAP proteins. Cell 102: 43–53, 2000.

[130648]

[130649]

[130650]11604.

[130651]Behrens, R.; Nolting, A.; Reimann, F.; Schwarz, M.; Wald-
schutz,

[130652]R.; Pongs, O.: hKCNMB3 and hKCNMB4, cloning and char-
acterization

[130653]of two members of the large-conductance calcium-acti-
vated potassium

[130654]channel beta subunit family. FEBS Lett. 474: 99–106,
2000.

[130655]

[130656]

[130657]11605.

[130658]Brenner, R.; Jegla, T. J.; Wickenden, A.; Liu, Y.; Aldrich, R.

[130659]W.: Cloning and functional characterization of novel large
conductance

[130660]calcium-activated potassium channel beta subunits, hKC-
NMB3 and hKCNMB4. J.

[130661]Biol. Chem. 275: 6453–6461, 2000.

[130662]

[130663]

[130664]11606.

[130665]Riazi, M. A.; Brinkman–Mills, P.; Johnson, A.; Naylor, S. L.;

Minoshima,

[130666]S.; Shimizu, N.; Baldini, A.; McDermid, H. E.: Identification
of

[130667]a putative regulatory subunit of a calcium–activated
potassium channel

[130668]in the dup(3q) syndrome region and a related sequence on
22q11.2. Genomics 62:

[130669]90–94, 1999.

[130670]

[130671]

[130672]11607.

[130673]Hemmi, H.; Takeuchi, O.; Kawai, T.; Kaisho, T.; Sato, S.;

Sanjo,

[130674]H.; Matsumoto, M.; Hoshino, K.; Wagner, H.; Takeda, K.;

Akira, S.

[130675]: A toll–like receptor recognizes bacterial DNA. Nature
408: 740–745,

[130676]2000.

[130677]

[130678]

[130679]11608.

[130680]Leadbetter, E. A.; Rifkin, I. R.; Hohlbaum, A. M.; Beaudette,
B.

[130681]C.; Shlomchik, M. J.; Marshak–Rothstein, A.: Chromatin–
IgG complexes

[130682]activate B cells by dual engagement of IgM and Toll–like
receptors. Nature 416:

[130683]603–607, 2002.

[130684]

[130685]

[130686]11609.

[130687]Takeshita, F.; Leifer, C. A.; Gursel, I.; Ishii, K. J.; Takeshita,

[130688]S.; Gursel, M.; Klinman, D. M.: Cutting edge: role of Toll–
like receptor

[130689]9 in CpG DNA–induced activation of human cells. J. Im–
mun. 167: 3555–3558,

[130690]2001.

[130691]

[130692]

[130693]11610.

[130694]Verthelyi, D.; Ishii, K. J.; Gursel, M.; Takeshita, F.; Klinman,

[130695]D. K.: Human peripheral blood cells differentially recognize and

[130696]respond to two distinct CpG motifs. J. Immun. 166: 2372–2377, 2001.

[130697]

[130698]

[130699]11611.

[130700]Bagrodia, S.; Taylor, S. J.; Jordon, K. A.; Van Aelst, L.; Cerione,

[130701]R. A.: A novel regulator of p21-activated kinases. J. Biol. Chem. 273:

[130702]23633–23636, 1998.

[130703]

[130704]

[130705]11612.

[130706]Oh, W. K.; Yoo, J. C.; Jo, D.; Song, Y. H.; Kim, M. G.; Park, D.

[130707]: Cloning of a SH3 domain-containing proline-rich pro-

tein, p85SPR,

[130708]and its localization in focal adhesion. Biochem. Biophys.

Res. Commun. 235:

[130709]794–798, 1997.

[130710]

[130711]

[130712]11613.

[130713]Ehrmann, D. A.; Schwarz, P. E. H.; Hara, M.; Tang, X.;

Horikawa,

[130714]Y.; Imperial, J.; Bell, G. I.; Cox, N. J.: Relationship of cal-

pain-10

[130715]genotype to phenotypic features of polycystic ovary syn-

drome. J.

[130716]Clin. Endocr. Metab. 87: 1669–1673, 2002.

[130717]

[130718]

[130719]11614.

[130720]Elbein, S. C.; Chu, W.; Ren, Q.; Hemphill, C.; Schay, J.; Cox,

[130721]N. J.; Hanis, C. L.; Hasstedt, S. J.: Role of calpain-10 gene

variants

[130722]in familial type 2 diabetes in Caucasians. J. Clin. Endocr.

Metab. 87:

[130723]650–654, 2002.

[130724]

[130725]

[130726]11615.

[130727]Fullerton, S. M.; Bartoszewicz, A.; Ybazeta, G.; Horikawa,
Y.;

[130728]Bell, G. I.; Kidd, K. K.; Cox, N. J.; Hudson, R. R.; Di Rienzo,
A.

[130729]: Geographic and haplotype structure of candidate type 2
diabetes–susceptibility

[130730]variants at the calpain–10 locus. Am. J. Hum. Genet. 70:
1096–1106,

[130731]2002.

[130732]

[130733]

[130734]11616.

[130735]Tsai, H.–J.; Sun, G.; Weeks, D. E.; Kaushal, R.; Wolujewicz,
M.;

[130736]McGarvey, S. T.; Tufa, J.; Viali, S.; Deka, R.: Type 2 diabetes and

[130737]three calpain-10 gene polymorphisms in Samoans: no evidence of association. Am.

[130738]J. Hum. Genet. 69: 1236–1244, 2001.

[130739]

[130740]

[130741]11617.

[130742]Weiss, K. M.; Terwilliger, J. D.: How many diseases does it take

[130743]to map a gene with SNPs? Nature Genet. 26: 151–157, 2000.

[130744]

[130745]

[130746]11618.

[130747]Hanis, C. L.; Boerwinkle, E.; Chakraborty, R.; Ellsworth, D. L.;

[130748]Concannon, P.; Stirling, B.; Morrison, V. A.; Wapelhorst, B.; Spielman,

[130749]R. S.; Gogolin-Ewens, K. J.; Shephard, J. M.; Williams, S. R.; and

[130750]21 others: A genome-wide search for human non-insulin-dependent

[130751](type 2) diabetes genes reveals a major susceptibility locus on chromosome

[130752]2. Nature Genet. 13: 161–166, 1996.

[130753]

[130754]

[130755]11619.

[130756]Weber, A.; Wienker, T. F.; Jung, M.; Easton, D.; Dean, H. J.; Heinrichs,

[130757]C.; Reis, A.; Clark, A. J. L.: Linkage of the gene for the triple

[130758]A syndrome to chromosome 12q13 near the type II keratin gene cluster. Hum.

[130759]Molec. Genet. 5: 2061–2066, 1996.

[130760]

[130761]

[130762]11620.

[130763]Bousquet, O.; Basseville, M.; Vila-Porcile, E.; Billette de Villemeur,

[130764]T.; Hauw, J.-J.; Landrieu, P.; Portier, M.-M.: Aggregation of
a subpopulation

[130765]of vimentin filaments in cultured human skin fibroblasts
derived from

[130766]patients with giant axonal neuropathy. Cell. Motil. Cy-
toskeleton 33:

[130767]115-129, 1996.

[130768]

[130769]

[130770]11621.

[130771]Kuhlenbaumer, G.; Young, P.; Oberwittler, C.; Hunermund,
G.; Schirmacher,

[130772]A.; Domschke, K.; Ringelstein, B.; Stogbauer, F.: Giant ax-
onal neuropathy

[130773](GAN): case report and two novel mutations in the gigax-
onin gene. Neurology 58:

[130774]1273-1276, 2002. Note: Erratum: Neurology 58: 1444,
2002.

[130775]

[130776]

[130777]11622.

[130778]Pena, S. D.: Giant axonal neuropathy: an inborn error of organization

[130779]of intermediate filaments. Muscle Nerve 5: 166–172, 1982.

[130780]

[130781]

[130782]11623.

[130783]Prineas, J. W.; Ouvrier, R. A.; Wright, R. G.; Walsh, J. C.; McLeod,

[130784]J. G.: Giant axonal neuropathy: a generalized disorder of cytoplasmic

[130785]microfilament formation. J. Neuropath. Exp. Neurol. 35: 458–470,

[130786]1976.

[130787]

[130788]

[130789]11624.

[130790]Bowe, A. E.; Finnegan, R.; Jan de Beur, S. M.; Cho, J.; Levine,

[130791]M. A.; Kumar, R.; Schiavi, S. C.: FGF–23 inhibits renal tubular phosphate

[130792]transport and is a PHEX substrate. Biochem. Biophys. Res. Commun. 284: 977–981, 2001.

[130794]

[130795]

[130796]11625.

[130797]Shimada, T.; Mizutani, S.; Muto, T.; Yoneya, T.; Hino, R.; Takeda,

[130798]S.; Takeuchi, Y.; Fujita, T.; Fukumoto, S.; Yamashita, T.: Cloning

[130799]and characterization of FGF23 as a causative factor of tumor-induced

[130800]osteomalacia. Proc. Nat. Acad. Sci. 98: 6500–6505, 2001.

[130801]

[130802]

[130803]11626.

[130804]Strewler, G. J.: FGF23, hypophosphatemia, and rickets: has phosphorylation

[130805]been found? (Commentary) Proc. Nat. Acad. Sci. 98: 5945–5946, 2001.

[130806]

[130807]

[130808]11627.

[130809]White, K. E.; Jonsson, K. B.; Carn, G.; Hampson, G.; Spector, T.

[130810]D.; Mannstadt, M.; Lorenz-Depiereux, B.; Miyauchi, A.; Yang, I. M.;

[130811]Ljunggren, O.; Meitinger, T.; Strom, T. M.; Juppner, H.; Econs, M.

[130812]J.: The autosomal dominant hypophosphatemic rickets (ADHR) gene is

[130813]a secreted polypeptide overexpressed by tumors that cause phosphate

[130814]wasting. J. Clin. Endocr. Metab. 86: 497–500, 2001.

[130815]

[130816]

[130817]11628.

[130818]Yamashita, T.; Yoshioka, M.; Itoh, N.: Identification of a novel

[130819]fibroblast growth factor, FGF–23, preferentially expressed in the

[130820]ventrolateral thalamic nucleus of the brain. Biochem. Biophys. Res.

[130821]Commun. 277: 494–498, 2000.

[130822]

[130823]

[130824]11629.

[130825]Barker, R. L.; Gleich, G. J.; Pease, L. R.: Acidic precursor revealed

[130826]in human eosinophil granule major basic protein cDNA. J. Exp. Med. 168:

[130827]1493–1498, 1988.

[130828]

[130829]

[130830]11630.

[130831]Li, M.–S.; Sun, L.; Satoh, T.; Fisher, L. M.; Spry, C. J. F.:

[130832]Human eosinophil major basic protein, a mediator of allergic inflammation,

[130833]is expressed by alternative splicing from two promoters. Biochem.

[130834]J. 305: 921–927, 1995.

[130835]

[130836]

[130837]11631.

[130838]McGrogan, M.; Simonsen, C.; Scott, R.; Griffith, J.; Ellis, N.;

[130839]Kennedy, J.; Campanelli, D.; Nathan, C.; Gabay, J.: Isolation
of

[130840]a complementary DNA clone encoding a precursor to hu-
man eosinophil

[130841]major basic protein. J. Exp. Med. 168: 2295–2308, 1988.

[130842]

[130843]

[130844]11632.

[130845]Plager, D. A.; Weiler, D. A.; Loegering, D. A.; Johnson, W.
B.;

[130846]Haley, L.; Eddy, R. L.; Shows, T. B.; Gleich, G. J.: Compara-
tive

[130847]structure, proximal promoter elements, and chromosome
location of

[130848]the human eosinophil major basic protein genes. Ge-
nomics 71: 271–281,

[130849]2001.

[130850]

[130851]

[130852]11633.

[130853]Wasmoen, T. L.; Bell, M. P.; Loegering, D. A.; Gleich, G. J.;

Prendergast,

[130854]F. G.; McKean, D. J.: Biochemical and amino acid sequence
analysis

[130855]of human eosinophil granule major basic protein. J. Biol.
Chem. 263:

[130856]12559–12563, 1988.

[130857]

[130858]

[130859]11634.

[130860]Weller, P. F.; Ackerman, S. J.; Smith, J. A.: Eosinophil gran-
ule

[130861]proteins: major basic protein is distinct from the smaller
subunit

[130862]of eosinophil peroxidase. J. Leukoc. Biol. 43: 1–4, 1988.

[130863]

[130864]

[130865]11635.

[130866]Yoshimatsu, K.; Ohya, Y.; Shikata, Y.; Seto, T.; Hasegawa, Y.;

[130867]Tanaka, I.; Kawamura, T.; Kitoh, K.; Toyoshima, S.; Osawa, T.: Purification

[130868]and cDNA cloning of a novel factor produced by a human T-cell hybridoma:

[130869]sequence homology with animal lectins. Molec. Immun. 29: 537–546,

[130870]1992.

[130871]

[130872]

[130873]11636.

[130874]Frey, N.; Richardson, J. A.; Olson, E. N.: Calsarcins, a novel

[130875]family of sarcomeric calcineurin-binding proteins. Proc. Nat. Acad.

[130876]Sci. 97: 14632–14637, 2000.

[130877]

[130878]

[130879]11637.

[130880]Faulkner, G.; Pallavicini, A.; Comelli, A.; Salamon, M.; Bor-
toletto,

[130881]G.; Ievolella, C.; Trevisan, S.; Kojic, S.; Dalla Vecchia, F.;
Laveder,

[130882]P.; Valle, G.; Lanfranchi, G.: FATZ, a filamin-, actinin-, and
telethonin-binding

[130883]protein of the Z-disc of skeletal muscle. J. Biol. Chem.
275: 41234–41242,

[130884]2000.

[130885]

[130886]

[130887]11638.

[130888]Takada, F.; Vander Woude, D. L.; Tong, H.–Q.; Thompson,
T. G.;

[130889]Watkins, S. C.; Kunkel, L. M.; Beggs, A. H.: Myozenin: an
alpha-actinin–

[130890]and gamma-filamin-binding protein of skeletal muscle Z
lines. Proc.

[130891]Nat. Acad. Sci. 98: 1595–1600, 2001.

[130892]

[130893]

[130894]11639.

[130895]Scott, A. F.: Personal Communication. Baltimore, Md.
3/13/2001.

[130896]

[130897]

[130898]11640.

[130899]Drane, P.; Barel, M.; Balbo, M.; Frade, R.: Identification of
[130900]RB18A, a 205 kDa new p53 regulatory protein which
shares antigenic

[130901]and functional properties with p53. Oncogene 15:
3013–3024, 1997.

[130902]

[130903]

[130904]11641.

[130905]Mansharamani, M.; Hewetson, A.; Chilton, B. S.: Cloning
and characterization

[130906]of an atypical type IV P-type ATPase that binds to the
RING motif

[130907]of RUSH transcription factors. J. Biol. Chem. 276:

3641–3649, 2001.

[130908]

[130909]

[130910]11642.

[130911]Nakayama, Y.; Weissman, S. M.; Bothwell, A. L. M.: BX–
MAS1 identifies

[130912]a cluster of homologous genes differentially expressed in
B cells. Biochem.

[130913]Biophys. Res. Commun. 285: 830–837, 2001.

[130914]

[130915]

[130916]11643.

[130917]Bashirova, A. A.; Geijtenbeek, T. B. H.; van Duijnhoven, G.
C.

[130918]F.; van Vliet, S. J.; Eilering, J. B. G.; Martin, M. P.; Wu, L.;
Martin,

[130919]T. D.; Viebig, N.; Knolle, P. A.; KewalRamani, V. N.; van
Kooyk, Y.;

[130920]Carrington, M.: A dendritic cell–specific intercellular adhe–
sion

[130921]molecule 3–grabbing nonintegrin (DC–SIGN)–related pro–

tein is highly

[130922]expressed on human liver sinusoidal endothelial cells and promotes

[130923]HIV-1 infection. J. Exp. Med. 193: 671-678, 2001.

[130924]

[130925]

[130926]11644.

[130927]Desai, R.; Peretz, A.; Idelson, H.; Lazarovici, P.; Attali, B.

[130928]: Ca(2+)-activated K(+) channels in human leukemic Jurkat T cells:

[130929]molecular cloning, biochemical and functional characterization. J.

[130930]Biol. Chem. 275: 39954-39963, 2000.

[130931]

[130932]

[130933]11645.

[130934]Maguchi, M.; Nishida, W.; Kohara, K.; Kuwano, A.; Kondo, I.; Hiwada,

[130935]K.: Molecular cloning and gene mapping of human basic and acidic

[130936]calponins. Biochem. Biophys. Res. Commun. 217:
238–244, 1995.

[130937]

[130938]

[130939]11646.

[130940]Miano, J. M.; Krahe, R.; Garcia, E.; Elliott, J. M.; Olson, E.

[130941]N.: Expression, genomic structure and high resolution
mapping to

[130942]19p13.2 of the human smooth muscle cell calponin gene.
Gene 197:

[130943]215–224, 1997.

[130944]

[130945]

[130946]11647.

[130947]Miano, J. M.; Thomas, S.; Distèche, C. M.: Expression and
chromosomal

[130948]mapping of the mouse smooth muscle calponin gene.
Mammalian Genome 12:

[130949]187–191, 2001.

[130950]

[130951]

[130952]11648.

[130953]Strasser, P.; Gimona, M.; Moessler, H.; Herzog, M.; Small, J. V.

[130954]: Mammalian calponin: identification and expression of genetic variants. FEBS

[130955]Lett. 330: 13–18, 1993.

[130956]

[130957]

[130958]11649.

[130959]Duetsch, G.; Illig, T.; Loesgen, S.; Rohde, K.; Kloop, N.; Herbon,

[130960]N.; Cohlke, H.; Altmueller, J.; Wjst, M.: STAT6 as an asthma candidate

[130961]gene: polymorphism–screening, association and haplo–type analysis in

[130962]a Caucasian sib–pair study. Hum. Molec. Genet. 11: 613–621, 2002.

[130963]

[130964]

[130965]11650.

[130966]Wu, H.-K.; Heng, H. H. Q.; Shi, X.-M.; Forsdyke, D. R.;
Tsui, L.-C.;

[130967]Mak, T. W.; Minden, M. D.; Siderovski, D. P.: Differential
expression

[130968]of a basic helix-loop-helix phosphoprotein gene, G0S8, in
acute leukemia

[130969]and localization to human chromosome 1q31. Leukemia
9: 1291-1298,

[130970]1995.

[130971]

[130972]

[130973]11651.

[130974]Bogerd, H. P.; Fridell, R. A.; Madore, S.; Cullen, B. R.: Iden-
tification

[130975]of a novel cellular cofactor for the Rev/Rex class of retro-
viral regulatory

[130976]proteins. Cell 82: 485-494, 1995.

[130977]

[130978]

[130979]11652.

[130980]Fritz, C. C.; Zapp, M. L.; Green, M. R.: A human nucleoporin-like

[130981]protein that specifically interacts with HIV Rev. Nature
376: 530–533,

[130982]1995.

[130983]

[130984]

[130985]11653.

[130986]Jones, T.; Sheer, D.; Bevec, D.; Kappel, B.; Hauber, J.;
Steinkasserer,

[130987]A.: The human HIV–1 Rev binding–protein hRIP/Rab (HRB)
maps to chromosome

[130988]2q36. Genomics 40: 198–199, 1997.

[130989]

[130990]

[130991]11654.

[130992]Kang–Decker, N.; Mantchev, G. T.; Juneja, S. C.; McNiven,
M. A.;

[130993]van Deursen, J. M. A.: Lack of acrosome formation in Hrb–

deficient

[130994]mice. Science 294: 1531–1533, 2001.

[130995]

[130996]

[130997]11655.

[130998]Salcini, A. E.; Confalonieri, S.; Doria, M.; Santolini, E.;

Tassi,

[130999]E.; Minenkova, O.; Cesareni, G.; Pelicci, P. G.; Di Fiore, P.

P.:

[131000]Binding specificity and in vivo targets of the EH domain, a

novel

[131001]protein–protein interaction module. Genes Dev. 11:

2239–2249, 1997.

[131002]

[131003]

[131004]11656.

[131005]Fish, K. J.; Cegielska, A.; Getman, M. E.; Landes, G. M.;

Virshup,

[131006]D. M.: Isolation and characterization of human casein ki-

nase I–epsilon

[131007](CKI), a novel member of the CKI gene family. J. Biol.

Chem. 270:

[131008]14875–14883, 1995.

[131009]

[131010]

[131011]11657.

[131012]Kloss, B.; Price, J. L.; Saez, L.; Blau, J.; Rothenfluh, A.; Wesley,
ley,

[131013]C. S.; Young, M. W.: The Drosophila clock gene double-
time encodes

[131014]a protein closely-related to human casein kinase I-
epsilon. Cell 94:

[131015]97–107, 1998.

[131016]

[131017]

[131018]11658.

[131019]Lowrey, P. L.; Shimomura, K.; Antoch, M. P.; Yamazaki, S.;
Zamenides,

[131020]P. D.; Ralph, M. R.; Menaker, M.; Takahashi, J. S.: Posi-
tional syntenic

[131021]cloning and functional characterization of the mammalian

circadian

[131022]mutation tau. Science 288: 483–491, 2000.

[131023]

[131024]

[131025]11659.

[131026]Graves, P. R.; Haas, D. W.; Hagedorn, C. H.; DePaoli–
Roach, A.

[131027]A.; Roach, P. J.: Molecular cloning, expression, and char–
acterization

[131028]of a 49–kilodalton casein kinase I isoform from rat testis.
J. Biol.

[131029]Chem. 268: 6394–6401, 1993.

[131030]

[131031]

[131032]11660.

[131033]Kusuda, J.; Hidari, N.; Hirai, M.; Hashimoto, K.: Sequence
analysis

[131034]of the cDNA for the human casein kinase 1–delta
(CSNK1D) gene and

[131035]its chromosomal localization. Genomics 32: 140–143,

1996.

[131036]

[131037]

[131038]11661.

[131039]Kools, P. F. J.; Roebroek, A. J. M.; van de Velde, H. J. K.;

Marynen,

[131040]P.; Bullerdiek, J.; Van de Ven, W. J. M.: Regional mapping

of the

[131041]human NSP gene to chromosome region 14q21–q22 by

fluorescence in situ

[131042]hybridization analysis. Cytogenet. Cell Genet. 66: 48–50,

1994.

[131043]

[131044]

[131045]11662.

[131046]Roebroek, A. J. M.; Ayoubi, T. A. Y.; van de Velde, H. J. K.;

Schoenmakers,

[131047]E. F. P. M.; Pauli, I. G. L.; Van de Ven, W. J. M.: Genomic

organization

[131048]of the human NSP gene, prototype of a novel gene family

encoding reticulons. Genomics 32:

[131049]191–199, 1996.

[131050]

[131051]

[131052]11663.

[131053]Roebroek, A. J. M.; van de Velde, H. J. K.; Van Bokhoven,
A.; Broers,

[131054]J. L. V.; Ramaekers, F. C. S.; Van de Ven, W. J. M.: Cloning
and

[131055]expression of alternative transcripts of a novel neuroen-
docrine-specific

[131056]gene and identification of its 135-kDa translational prod-
uct. J.

[131057]Biol. Chem. 268: 13439–13447, 1993.

[131058]

[131059]

[131060]11664.

[131061]Senden, N. H. M.; van de Velde, H. J. K.; Broers, J. L. V.;
Timmer,

[131062]E. D. J.; Roebroek, A. J. M.; van de Ven, W. J. M.; Ramaek-
ers, F.

[131063]C. S.: Cluster-10 lung-cancer antibodies recognize NSPs,
novel neuro-endocrine

[131064]proteins associated with membranes of the endoplasmic
reticulum. Int.

[131065]J. Cancer (Suppl. 8): 84-88, 1994.

[131066]

[131067]

[131068]11665.

[131069]van de Velde, H. J. K.; Roebroek, A. J. M.; van Leeuwen, F.
W.;

[131070]Van de Ven, W. J. M.: Molecular analysis of expression in
rat brain

[131071]of NSP-A, a novel neuroendocrine-specific protein of the
endoplasmic

[131072]reticulum. Molec. Brain Res. 23: 81-92, 1994.

[131073]

[131074]

[131075]11666.

[131076]van de Velde, H. J. K.; Senden, N. H. M.; Roskams, T. A. D.;
Broers,

[131077]J. L. V.; Ramaekers, F. C. S.; Roebroek, A. J. M.; Van de Ven, W.

[131078]J. M.: NSP-encoded reticulons are neuroendocrine markers of a novel

[131079]category in human lung cancer diagnosis. Cancer Res. 54: 4769–4776,

[131080]1994.

[131081]

[131082]

[131083]11667.

[131084]Baron, B. W.; Anastasi, J.; Thirman, M. J.; Furukawa, Y.; Fears,

[131085]S.; Kim, D. C.; Simone, F.; Birkenbach, M.; Montag, A.; Sadhu, A.;

[131086]Zelevnik–Le, N.; McKeithan, T. W.: The human programmed cell death–2

[131087](PDCD2) gene is a target of BCL6 repression: implications for a role

[131088]of BCL6 in the down-regulation of apoptosis. Proc. Nat. Acad. Sci. 99:

[131089]2860–2865, 2002.

[131090]

[131091]

[131092]11668.

[131093]Cohen, J. J.; Duke, R. C.: Glucocorticoid activation of a calcium-dependent

[131094]endonuclease in thymocyte nuclei leads to cell death. J. Immun. 132:

[131095]38–42, 1984.

[131096]

[131097]

[131098]11669.

[131099]Kawakami, T.; Furukawa, Y.; Sudo, K.; Saito, H.; Takami, S.; Takahashi,

[131100]E.; Nakamura, Y.: Isolation and mapping of a human gene (PDCD2) that

[131101]is highly homologous to Rp8, a rat gene associated with programmed

[131102]cell death. Cytogenet. Cell Genet. 71: 41–43, 1995.

[131103]

[131104]

[131105]11670.

[131106]Owens, G. P.; Cohen, J. J.: Identification of genes involved
in

[131107]programmed cell death. Cancer Metastasis Rev. 11:
149–156, 1992.

[131108]

[131109]

[131110]11671.

[131111]Sellins, K. S.; Cohen, J. J.: Gene induction by gamma-ir-
radiation

[131112]leads to DNA fragmentation in lymphocytes. J. Immun.
139: 3199–3206,

[131113]1987.

[131114]

[131115]

[131116]11672.

[131117]Wyllie, A. H.: Glucocorticoid-induced thymocyte apoptosis
is associated

[131118]with endogenous endonuclease activation. Nature 284:
555–556, 1980.

[131119]

[131120]

[131121]11673.

[131122]Bullrich, F.; Druck, T.; Kunapuli, P.; Gomez, J.; Gripp, K.
W.;

[131123]Schlegelberger, B.; Lasota, J.; Aronson, M.; Cannizzaro, L.
A.; Huebner,

[131124]K.; Benovic, J. L.: Chromosomal mapping of the genes
GPRK5 and GPRK6

[131125]encoding G protein-coupled receptor kinases GRK5 and
GRK6. Cytogenet.

[131126]Cell Genet. 70: 250–254, 1995.

[131127]

[131128]

[131129]11674.

[131130]Haribabu, B.; Snyderman, R.: Identification of additional
members

[131131]of human G-protein-coupled receptor kinase multigene
family. Proc.

[131132]Nat. Acad. Sci. 90: 9398–9402, 1993.

[131133]

[131134]

[131135]11675.

[131136]Miki, T.; Suzuki, M.; Shibasaki, T.; Uemura, H.; Sato, T.;
Yamaguchi,

[131137]K.; Koseki, H.; Iwanaga, T.; Nakaya, H.; Seino, S.: Mouse
model of

[131138]Prinzmetal angina by disruption of the inward rectifier
Kir6.1. Nature

[131139]Med. 8: 466–472, 2002.

[131140]

[131141]

[131142]11676.

[131143]Prinzmetal, M.; Kennamer, R.; Merliss, R.; Wada, T.; Bor,
N.:

[131144]Angina pectoris. 1. A variant form of angina pectoris: pre-
liminary

[131145]report. Am. J. Med. 27: 375–388, 1959.

[131146]

[131147]

[131148]11677.

[131149]Pristipino, C.; Beltrame, J. F.; Finocchiaro, M. L.; Hattori, R.;

[131150]Fujita, M.; Mongiardo, R.; Cianflone, D.; Sanna, T.; Sasayama, S.;

[131151]Maseri, A.: Major racial differences in coronary constrictor response

[131152]between Japanese and Caucasians with recent myocardial infarction. Circulation 101:

[131153]1102–1108, 2000.

[131154]

[131155]

[131156]11678.

[131157]Cherel, M.; Sorel, M.; Apiou, F.; Lebeau, B.; Dubois, S.; Jacques,

[131158]Y.; Minvielle, S.: The human interleukin–11 receptor alpha gene (IL11RA):

[131159]genomic organization and chromosome mapping. Genomics 32: 49–53,

[131160]1996.

[131161]

[131162]

[131163]11679.

[131164]Cherel, M.; Sorel, M.; Lebeau, B.; Dubois, S.; Moreau, J.-F.;
Bataille,

[131165]R.; Minvielle, S.; Jacques, Y.: Molecular cloning to two iso-
forms

[131166]of a receptor for the human hematopoietic cytokine inter-
leukin-11. Blood 86:

[131167]2534-2540, 1995.

[131168]

[131169]

[131170]11680.

[131171]Neuhaus, H.; Bettenhausen, B.; Bilinski, P.; Simon-
Chazottes, D.;

[131172]Guenet, J.-L.; Gossler, A.: Etl2, a novel putative type-I cy-
tokine

[131173]receptor expressed during mouse embryogenesis at high
levels in skin

[131174]and cells with skeletogenic potential. Dev. Biol. 166:
531-542,

[131175]1994.

[131176]

[131177]

[131178]11681.

[131179]Van Leuven, F.; Stas, L.; Hilliker, C.; Miyake, Y.; Bilinski, P.;

[131180]Gossler, A.: Molecular cloning and characterization of the
human

[131181]interleukin-11 receptor alpha-chain gene, IL11RA, located
on chromosome

[131182]9p13. Genomics 31: 65-70, 1996.

[131183]

[131184]

[131185]11682.

[131186]Magrangeas, F.; Pitiot, G.; Dubois, S.; Bragado-Nilsson, E.;
Cherel,

[131187]M.; Jobert, S.; Lebeau, B.; Boisteau, O.; Lethe, B.; Mallet, J.;
Jacques,

[131188]Y.; Minvielle, S.: Cotranscription and intergenic splicing of
human

[131189]galactose-1-phosphate uridylyltransferase and inter-
leukin-11 receptor

[131190]alpha-chain genes generate a fusion mRNA in normal cells: implication

[131191]for the production of multidomain proteins during evolution. J. Biol.

[131192]Chem. 273: 16005–16010, 1998.

[131193]

[131194]

[131195]11683.

[131196]Avraham, K. B.; Hasson, T.; Sobe, T.; Balsara, B.; Testa, J. R.;

[131197]Skvorak, A. B.; Morton, C. C.; Copeland, N. G.; Jenkins, N. A.: Characterization

[131198]of unconventional MYO6, the human homologue of the gene responsible

[131199]for deafness in Snell's waltzer mice. Hum. Molec. Genet. 6: 1225–1231,

[131200]1997.

[131201]

[131202]

[131203]11684.

[131204]Avraham, K. B.; Hasson, T.; Steel, K. P.; Kingsley, D. M.;
Russell,

[131205]L. B.; Mooseker, M. S.; Copeland, N. G.; Jenkins, N. A.: The
mouse

[131206]Snell's waltzer deafness gene encodes an unconventional
myosin required

[131207]for structural integrity of inner ear hair cells. Nature
Genet. 11:

[131208]369–375, 1995.

[131209]

[131210]

[131211]11685.

[131212]Hasson, T.; Mooseker, M. S.: Porcine myosin–VI: charac-
terization

[131213]of a new mammalian unconventional myosin. J. Cell. Biol.
127: 425–440,

[131214]1994.

[131215]

[131216]

[131217]11686.

[131218]Hasson, T.; Skowron, J. F.; Gilbert, D. J.; Avraham, K. B.;
Perry,

[131219]W. L.; Bement, W. M.; Anderson, B. L.; Sherr, E. H.; Chen,
Z.-Y.;

[131220]Greene, L. A.; Ward, D. C.; Corey, D. P.; Mooseker, M. S.;
Copeland,

[131221]N. G.; Jenkins, N. A.: Mapping of unconventional myosins
in mouse

[131222]and human. Genomics 36: 431–439, 1996.

[131223]

[131224]

[131225]11687.

[131226]Rock, R. S.; Rice, S. E.; Wells, A. L.; Purcell, T. J.; Spudich,

[131227]J. A.; Sweeney, H. L.: Myosin VI is a processive motor with
a large

[131228]step size. Proc. Nat. Acad. Sci. 98: 13655–13659, 2001.

[131229]

[131230]

[131231]11688.

[131232]Wells, A. L.; Lin, A. W.; Chen, L.-Q.; Safer, D.; Cain, S. M.;

[131233]Hasson, T.; Carragher, B. O.; Milligan, R. A.; Sweeney, H.
L.: Myosin

[131234]VI is an actin-based motor that moves backwards. Nature
401: 505–508,

[131235]1999.

[131236]

[131237]

[131238]11689.

[131239]Melchionda, S.; Ahituv, N.; Bisceglia, L.; Sobe, T.; Glaser,
F.;

[131240]Rabionet, R.; Arbones, M. L.; Notarangelo, A.; Di Iorio, E.;
Carella,

[131241]M.; Zelante, L.; Estivill, X.; Avraham, K. B.; Gasparini, P.:
MYO6,

[131242]the human homologue of the gene responsible for deaf-
ness in Snell's

[131243]Waltzer mice, is mutated in autosomal dominant nonsyn-
dromic hearing

[131244]loss. Am. J. Hum. Genet. 69: 635–640, 2001.

[131245]

[131246]

[131247]11690.

[131248]Chittum, H. S.; Himeno, S.; Hill, K. E.; Burk, R. F.: Multiple

[131249]forms of selenoprotein P in rat plasma. Arch. Biochem.

Biophys. 325:

[131250]124–128, 1996.

[131251]

[131252]

[131253]11691.

[131254]Bonne, S.; van Hengel, J.; van Roy, F.: Chromosomal map-

ping of

[131255]human armadillo genes belonging to the

p120(ctn)/plakophilin subfamily. Genomics 51:

[131256]452–454, 1998.

[131257]

[131258]

[131259]11692.

[131260]Drab, M.; Verkade, P.; Elger, M.; Kasper, M.; Lohn, M.;

Lauterbach,

[131261]B.; Menne, J.; Lindschau, C.; Mende, F.; Luft, F. C.; Schedl,

A.;

[131262]Haller, H.; Kurzchalia, T. V.: Loss of caveolae, vascular dysfunction,

[131263]and pulmonary defects in caveolin-1 gene-disrupted mice. Science 293:

[131264]2449–2452, 2001.

[131265]

[131266]

[131267]11693.

[131268]Engelman, J. A.; Zhang, X.; Galbiati, F.; Volonte, D.; Sotgia,

[131269]F.; Pestell, R. G.; Minetti, C.; Scherer, P. E.; Okamoto, T.; Lisanti,

[131270]M. P.: Molecular genetics of the caveolin gene family: implications

[131271]for human cancers, diabetes, Alzheimer disease, and muscular dystrophy. Am.

[131272]J. Hum. Genet. 63: 1578–1587, 1998.

[131273]

[131274]

[131275]11694.

[131276]Engelman, J. A.; Zhang, X. L.; Galbiati, F.; Lisanti, M. P.:
Chromosomal

[131277]localization, genomic organization, and developmental
expression of

[131278]the murine caveolin gene family (Cav-1, -2, and -3): Cav-
1 and Cav-2

[131279]genes map to a known tumor suppressor locus
(6-A2/7q31). FEBS Lett. 429:

[131280]330-336, 1998.

[131281]

[131282]

[131283]11695.

[131284]Engelman, J. A.; Zhang, X. L.; Lisanti, M. P.: Genes encod-
ing

[131285]human caveolin-1 and -2 are co-localized to the D7S522
locus (7q31.1),

[131286]a known fragile site (FRA7G) that is frequently deleted in
human cancers. FEBS

[131287]Lett. 436: 403-410, 1998.

[131288]

[131289]

[131290]11696.

[131291]Feron, O.; Dessy, C.; Moniotte, S.; Desager, J.-P.; Balli-
gand,

[131292]J.-L.: Hypercholesterolemia decreases nitric oxide produc-
tion by

[131293]promoting the interaction of caveolin and endothelial ni-
tric oxide

[131294]synthase. J. Clin. Invest. 103: 897–905, 1999.

[131295]

[131296]

[131297]11697.

[131298]Giordano, S.; Ponzetto, C.; Di Renzo, M. F.; Cooper, C. S.;
Comoglio,

[131299]P. M.: Tyrosine kinase receptor indistinguishable from the
c-met

[131300]protein. Nature 339: 155–156, 1989.

[131301]

[131302]

[131303]11698.

[131304]Glenney, J. R., Jr.: The sequence of human caveolin reveals

identity

[131305]with VIP21, a component of transport vesicles. FEBS Lett.

314: 45–48,

[131306]1992.

[131307]

[131308]

[131309]11699.

[131310]Kurzchalia, T. V.; Dupree, P.; Parton, R. G.; Kellner, R.;

Virta,

[131311]H.; Lehnert, M.; Simons, K.: VIP21, a 21-kD membrane

protein is an

[131312]integral component of trans-Golgi-network-derived

transport vesicles. J.

[131313]Cell Biol. 118 1003–1014, 1992.

[131314]

[131315]

[131316]11700.

[131317]Scherer, P. E.; Okamoto, T.; Chun, M.; Nishimoto, I.;

Lodish,

[131318]H. F.; Lisanti, M. P.: Identification, sequence, and expres–

sion of

[131319]caveolin-2 defines a caveolin gene family. Proc. Nat. Acad. Sci. 93:

[131320]131-135, 1996.

[131321]

[131322]

[131323]11701.

[131324]Scherer, P. E.; Tang, Z.; Chun, M.; Sargiacomo, M.; Lodish, H.

[131325]F.; Lisanti, M. P.: Caveolin isoforms differ in their N-terminal

[131326]protein sequence and subcellular distribution: identification and

[131327]epitope mapping of an isoform-specific monoclonal antibody probe. J.

[131328]Biol. Chem. 270 16395-16401, 1995.

[131329]

[131330]

[131331]11702.

[131332]Razani, B.; Engelman, J. A.; Wang, X. B.; Schubert, W.;

Zhang,

[131333]X. L.; Marks, C. B.; Macaluso, F.; Russell, R. G.; Li, M.;

Pestell,

[131334]R. G.; Di Vizio, D.; Hou, H., Jr.; Kneitz, B.; Lagaud, G.;

Christ,

[131335]G. J.; Edelman, W.; Lisanti, M. P.: Caveolin-1 null mice are
viable

[131336]but show evidence of hyperproliferative and vascular ab-
normalities. J.

[131337]Biol. Chem. 276: 38121-38138, 2001.

[131338]

[131339]

[131340]11703.

[131341]Bell, D. R.; Plant, N. J.; Rider, C. G.; Na, L.; Brown, S.;

Ateitalla,

[131342]I.; Acharya, S. K.; Davies, M. H.; Elias, E.; Jenkins, N. A.;

Gilbert,

[131343]D. J.; Copeland, N. G.; Elcombe, C. R.: Species-specific in-
duction

[131344]of cytochrome P-450 4A RNAs: PCR cloning of partial
guinea-pig, human

[131345]and mouse CYP4A cDNAs. Biochem. J. 294: 173-180,

1993.

[131346]

[131347]

[131348]11704.

[131349]Imaoka, S.; Ogawa, H.; Kimura, S.; Gonzalez, F. J.: Complete cDNA

[131350]sequence and cDNA-directed expression of CYP4A11, a fatty acid omega-hydroxylase

[131351]expressed in human kidney. DNA Cell Biol. 12: 893-899, 1993.

[131352]

[131353]

[131354]11705.

[131355]Kawashima, H.; Kusunose, E.; Kikuta, Y.; Kinoshita, H.; Tanaka,

[131356]S.; Yamamoto, S.; Kishimoto, T.; Kusunose, M.: Purification and cDNA

[131357]cloning of human liver CYP4A fatty acid omega-hydroxylase. J. Biochem. 116:

[131358]74-80, 1994.

[131359]

[131360]

[131361]11706.

[131362]Palmer, C. N. A.; Richardson, T. H.; Griffin, K. J.; Hsu, M.-H.;

[131363]Muerhoff, A. S.; Clark, J. E.; Johnson, E. F.: Characterization of

[131364]a cDNA encoding a human kidney, cytochrome P-450 4A fatty acid omega-hydroxylase

[131365]and the cognate enzyme expressed in Escherichia coli. Biochim. Biophys.

[131366]Acta 1172: 161-166, 1993.

[131367]

[131368]

[131369]11707.

[131370]Muller, J.; Ory, S.; Copeland, T.; Piwnicka-Worms, H.; Morrison,

[131371]D. K.: C-TAK1 regulates Ras signaling by phosphorylating the MAPK

[131372]scaffold, KSR1. Molec. Cell 8: 983-993, 2001.

[131373]

[131374]

[131375]11708.

[131376]Rosenthal, R. L.: Haemorrhage in PTA (factor XI) deficiency.

[131377](Abstract) Proc. 10th Int. Cong. Soc. Hemat., Stockholm , 1964.

[131378]

[131379]

[131380]11709.

[131381]Laporte, J.; Hu, L. J.; Kretz, C.; Mandel, J.-L.; Kioschis, P.;

[131382]Coy, J. F.; Klauck, S. M.; Poustka, A.; Dahl, N.: A gene mutated

[131383]in X-linked myotubular myopathy defines a new putative tyrosine phosphatase

[131384]family conserved in yeast. Nature Genet. 13: 175-182, 1996.

[131385]

[131386]

[131387]11710.

[131388]Rosenthal, R. L.; Dreskin, O. H.; Rosenthal, N.: Plasma

thromboplastin

[131389]antecedent (PTA) deficiency: clinical, coagulation, therapeutic and

[131390]hereditary aspects of a new hemophilia-like disease.

Blood 10: 120-131,

[131391]1955.

[131392]

[131393]

[131394]11711.

[131395]Seligsohn, U.: Factor XI (PTA) deficiency.In: Goodman, R.
E.;

[131396]Motulsky, A. G.: Genetic Diseases Among Ashkenazi Jews.
New York:

[131397]Raven Press (pub.) 1979. Pp. 141-148.

[131398]

[131399]

[131400]11712.

[131401]Vinazzer, H.: Partieller familiaerer Faktor-XI-Mangel. Blut
15:

[131402]263-267, 1967.

[131403]

[131404]

[131405]11713.

[131406]Wistinghausen, B.; Reischer, A.; Oddoux, C.; Ostrer, H.;
Nardi,

[131407]M.; Karpatkin, M.: Severe factor XI deficiency in an Arab
family

[131408]associated with a novel mutation in exon 11. Brit. J.
Haemat. 99:

[131409]575–577, 1997.

[131410]

[131411]

[131412]11714.

[131413]Zivelin, A.; Bauduer, F.; Ducout, L.; Peretz, H.; Rosenberg,
N.;

[131414]Yatuv, R.; Seligsohn, U.: Factor XI deficiency in French
Basques

[131415]is caused predominantly by an ancestral cys38-to-arg
mutation in the

[131416]factor XI gene. Blood 99: 2448–2454, 2002.

[131417]

[131418]

[131419]11715.

[131420]Hsueh, Y.-P.; Wang, T.-F.; Yang, F.-C.; Sheng, M.: Nuclear transcription

[131421]and transcription regulation by the membrane-associated guanylate

[131422]kinase CASK/LIN-2. Nature 404: 298–302, 2000.

[131423]

[131424]

[131425]11716.

[131426]De Smet, C.; Lurquin, C.; van der Bruggen, P.; De Plaen, E.; Brasseur,

[131427]F.; Boon, T.: Sequence and expression pattern of the human MAGE2

[131428]gene. Immunogenetics 39: 121–129, 1994.

[131429]

[131430]

[131431]11717.

[131432]Chambost, H.; Van Baren, N.; Brasseur, F.; Godelaine, D.;
Xerri,

[131433]L.; Landi, S. J.; Theate, I.; Plumas, J.; Spagnoli, G. C.;
Michel,

[131434]G.; Coulie, P. G.; Olive, D.: Expression of gene MAGE-A4 in
Reed-Sternberg

[131435]cells. Blood 95: 3530–3533, 2000.

[131436]

[131437]

[131438]11718.

[131439]Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Miyajima,
N.;

[131440]Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction
of the

[131441]coding sequences of unidentified human genes. XI. The
complete sequences

[131442]of 100 new cDNA clones from brain which code for large
proteins in

[131443]vitro. DNA Res. 5: 277–286, 1998.

[131444]

[131445]

[131446]11719.

[131447]Gospe, S. M., Jr.; Lazaro, R. P.; Lava, N. S.; Grootscholten,

[131448]P. M.; Scott, M. O.; Fischbeck, K. H.: Familial X-linked
myalgia

[131449]and cramps: a nonprogressive myopathy associated with a
deletion in

[131450]the dystrophin gene. *Neurology* 39: 1277–1280, 1989.

[131451]

[131452]

[131453]11720.

[131454]Kingston, H. M.; Sarfarazi, M.; Thomas, N. S. T.; Harper, P.
S.

[131455]: Localisation of the Becker muscular dystrophy gene on
the short

[131456]arm of the X chromosome by linkage to cloned DNA se-
quences. *Hum.*

[131457]*Genet.* 67: 6–17, 1984.

[131458]

[131459]

[131460]11721.

- [131461]Kingston, H. M.; Thomas, N. S. T.; Pearson, P. L.; Sarfarazi,
[131462]M.; Harper, P. S.: Genetic linkage between Becker muscular
dystrophy
[131463]and a polymorphic DNA sequence on the short arm of the
X chromosome. J.
[131464]Med. Genet. 20: 255–258, 1983.
- [131465]
- [131466]
- [131467]11722.
- [131468]Reardon, W.; Middleton–Price, H. R.; Sandkuijl, L.; Phelps,
P.;
- [131469]Bellman, S.; Luxon, L.; Pembrey, M. E.; Malcolm, S.: A mul-
tipedigree

[131470]linkage study of X-linked deafness: linkage to Xq13-q21
and evidence

[131471]for genetic heterogeneity. Genomics 11: 885-894, 1991.

[131472]

[131473]

[131474]11723.

[131475]Cardoso, C.; Timsit, S.; Villard, L.; Khrestchatisky, M.;
Fontes,

[131476]M.; Colleaux, L.: Specific interaction between the XNP/
ATR-X gene

[131477]product and the SET domain of the human EZH2 protein.
Hum. Molec.

[131478]Genet. 7: 679-684, 1998.

[131479]

[131480]

[131481]11724.

[131482]Bairoch, A.: Personal Communication. Geneva, Switzerland
5/13/1994.

[131483]

[131484]

[131485]11725.

[131486]Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.-H.; Goldstein, L. S.

[131487]: Axonal transport of amyloid precursor protein is mediated by direct

[131488]binding to the kinesin light chain subunit of kinesin-I.

Neuron 28:

[131489]449–459, 2000.

[131490]

[131491]

[131492]11726.

[131493]Luo, Y.; Bolon, B.; Kahn, S.; Bennett, B. D.; Babu-Khan, S.; Denis,

[131494]P.; Fan, W.; Kha, H.; Zhang, J.; Gong, Y.; Martin, L.; Louis, J.-C.;

[131495]Yan, Q.; Richards, W. G.; Citron, M.; Vassar, R.: Mice deficient

[131496]in BACE1, the Alzheimer's beta-secretase, have normal phenotype and

[131497]abolished beta-amyloid generation. Nature Neurosci. 4: 231–232,

[131498]2001.

[131499]

[131500]

[131501]11727.

[131502]Gross, J. A.; Dillon, S. R.; Mudri, S.; Johnston, J.; Littau, A.;

[131503]Roque, R.; Rixon, M.; Schou, O.; Foley, K. P.; Haugen, H.;
McMillen,

[131504]S.; Waggle, K.; Schreckhise, R. W.; Shoemaker, K.; Vu, T.;
Moore,

[131505]M.; Grossman, A.; Clegg, C. H.: TACI-Ig neutralizes
molecules critical

[131506]for B cell development and autoimmune disease: impaired
B cell maturation

[131507]in mice lacking BLyS. Immunity 15: 289–302, 2001.

[131508]

[131509]

[131510]11728.

[131511]Liu, Y.; Xu, L.; Opalka, N.; Kappler, J.; Shu, H.-B.; Zhang,
G.

[131512]: Crystal structure of sTALL-1 reveals a virus-like assem-

bly of TNF

[131513]family ligands. Cell 108: 383–394, 2002.

[131514]

[131515]

[131516]11729.

[131517]Moore, P. A.; Belvedere, O.; Orr, A.; Pierl, K.; LaFleur, D.
W.;

[131518]Feng, P.; Soppet, D.; Charters, M.; Gentz, R.; Parmelee, D.;
Li, Y.;

[131519]Galperina, O.; and 10 others: BLyS: member of the tumor
necrosis

[131520]factor family and B lymphocyte stimulator. Science 285:
260–263,

[131521]1999.

[131522]

[131523]

[131524]11730.

[131525]Mukhopadhyay, A.; Ni, J.; Zhai, Y.; Yu, G.–L.; Aggarwal, B.
B.

[131526]: Identification and characterization of a novel cytokine,

THANK,

[131527]a TNF homologue that activates apoptosis, nuclear factor-
kappa-B,

[131528]and c-Jun NH2-terminal kinase. J. Biol. Chem. 274:
15978-15981,

[131529]1999.

[131530]

[131531]

[131532]11731.

[131533]Oren, D. A.; Li, Y.; Volovik, Y.; Morris, T. S.; Dharia, C.;
Das,

[131534]K.; Galperina, O.; Gentz, R.; Arnold, E.: Structural basis of
BLyS

[131535]receptor recognition. Nature Struct. Biol. 9: 288-292,
2002.

[131536]

[131537]

[131538]11732.

[131539]Schiemann, B.; Gommerman, J. L.; Vora, K.; Cachero, T. G.;
Shulga-Morskaya,

[131540]S.; Dobles, M.; Frew, E.; Scott, M. L.: An essential role for

BAFF

[131541]in the normal development of B cells through a BCMA-independent pathway. Science 2111–2114,

[131542]2001.

[131543]

[131544]

[131545]11733.

[131546]Schneider, P.; MacKay, F.; Steiner, V.; Hofmann, K.; Bodmer, J.

[131547]L.; Holler, N.; Ambrose, C.; Lawton, P.; Bixler, S.; Acha-Orbea, H.;

[131548]Valmori, D.; Romero, P.; Werner-Favre, C.; Zubler, R. H.; Browning,

[131549]J. L.; Tschopp, J.: BAFF, a novel ligand of the tumor necrosis factor

[131550]family, stimulates B cell growth. J. Exp. Med. 189: 1747–1756, 1999.

[131551]

[131552]

[131553]11734.

[131554]Shu, H.-B.; Hu, W. H.; Johnson, H.: TALL-1 is a novel

member

[131555]of the TNF family that is down-regulated by mitogens. J.

Leukoc.

[131556]Biol. 65: 680-683, 1999.

[131557]

[131558]

[131559]11735.

[131560]Tribouley, C.; Wallroth, M.; Chan, V.; Paliard, X.; Fang, E.;

[131561]Lamson, G.; Pot, D.; Escobedo, J.; Williams, L. T.: Characterization
terization

[131562]of a new member of the TNF family expressed on antigen-presenting
presenting

[131563]cells. Biol. Chem. 380: 1443-1447, 1999.

[131564]

[131565]

[131566]11736.

[131567]Okamoto, N.; Hori, S.; Akazawa, C.; Hayashi, Y.; Shigemoto, R.;

[131568]Mizuno, N.; Nakanishi, S.: Molecular characterization of a
new metabotropic

[131569]glutamate receptor mGluR7 coupled to inhibitory cyclic
AMP signal

[131570]transduction. J. Biol. Chem. 269: 1231–1236, 1994.

[131571]

[131572]

[131573]11737.

[131574]Ango, F.; Prezeau, L.; Muller, T.; Tu, J. C.; Xiao, B.; Worley,

[131575]P. F.; Pin, J. P.; Bockaert, J.; Fagni, L.: Agonist-independent
activation

[131576]of metabotropic glutamate receptors by the intracellular
protein Homer. Nature 411:

[131577]962–965, 2001.

[131578]

[131579]

[131580]11738.

[131581]Offenberg, H. H.; Schalk, J. A. C.; Meuwissen, R. L. J.; van
Aalderen,

[131582]M.; Kester, H. A.; Dietrich, A. J. J.; Heyting, C.: SCP2: a ma-
jor

[131583]protein component of the axial elements of synaptonemal

complexes

[131584]of the rat. Nucleic Acids Res. 26: 2572–2579, 1998.

[131585]

[131586]

[131587]11739.

[131588]Schalk, J. A. C.; Offenberg, H. H.; Peters, E.; Groot, N. P. B.;

[131589]Hoovers, J. M. N.; Heyting, C.: Isolation and characteriza-
tion of

[131590]the human SCP2 cDNA and chromosomal localization of
the gene. Mammalian

[131591]Genome 10: 642–644, 1999.

[131592]

[131593]

[131594]11740.

[131595]Komada, M.; McLean, D. J.; Griswold, M. D.; Russell, L. D.;
Soriano,

[131596]P.: E-MAP-115, encoding a microtubule-associated pro-
tein, is a retinoic

[131597]acid-inducible gene required for spermatogenesis. Genes
Dev. 14:

[131598]1332–1342, 2000.

[131599]

[131600]

[131601]11741.

[131602]Masson, D.; Kreis, T. E.: Identification and molecular characterization

[131603]of E-MAP-115, a novel microtubule-associated protein predominantly

[131604]expressed in epithelial cells. J. Cell Biol. 123: 357–371, 1993.

[131605]

[131606]

[131607]11742.

[131608]Kowalski, P. E.; Freeman, J. D.; Nelson, D. T.; Mager, D. L.:

[131609]Genomic structure and evolution of a novel gene (PLA2L) with duplicated

[131610]phospholipase A2-like domains. Genomics 39: 38–46, 1997.

[131611]

[131612]

[131613]11743.

[131614]Liu, M.; Parker, R. M. C.; Darby, K.; Eyre, H. J.; Copeland, N.

[131615]G.; Crawford, J.; Gilbert, D. J.; Sutherland, G. R.; Jenkins, N. A.;

[131616]Herzog, H.: GPR56, a novel secretin-like human G-protein-coupled

[131617]receptor gene. Genomics 55: 296–305, 1999.

[131618]

[131619]

[131620]11744.

[131621]Zendman, A. J. W.; Cornelissen, I. M. H. A.; Weidle, U. H.; Ruiter,

[131622]D. J.; van Muijen, G. N. P.: TM7XN1, a novel human EGF-TM7-like cDNA,

[131623]detected with mRNA differential display using human melanoma cell

[131624]lines with different metastatic potential. FEBS Lett. 446: 292–298,

[131625]1999.

[131626]

[131627]

[131628]11745.

[131629]Nicolaou, M.; Song, Y.-Q.; Sato, C. A.; Orlacchio, A.;
Kawarai,

[131630]T.; Medeiros, H.; Liang, Y.; Sorbi, S.; Richard, E.; Rogaev, E.
I.;

[131631]Moliaka, Y.; Bruni, A. C.; Jorge, R.; Percy, M.; Duara, R.;
Farrer,

[131632]L. A.; St George-Hyslop, P.; Rogaeva, E. A.: Mutations in
the open

[131633]reading frame of the beta-site APP cleaving enzyme
(BACE) locus are

[131634]not a common cause of Alzheimer's disease. Neurogenet-
ics 3: 203-206,

[131635]2001.

[131636]

[131637]

[131638]11746.

[131639]Saunders, A. J.; Kim, T.-W.; Tanzi, R. E.: BACE maps to
chromosome

[131640]11 and a BACE homolog, BACE2, reside in the obligate

Down syndrome

[131641]region of chromosome 21. Science 286: 1255a, 1999.

Note: Electronic

[131642]Publication.

[131643]

[131644]

[131645]11747.

[131646]Sinha, S.; Anderson, J. P.; Barbour, R.; Basi, G. S.; Caccavello,

[131647]R.; Davis, D.; Doan, M.; Dovey, H. F.; Frigon, N.; Hong, J.; Jacobson-Croak,

[131648]K.; Jewett, N.; and 15 others: Purification and cloning of amyloid

[131649]precursor protein beta-secretase from human brain. Nature 402: 537-540,

[131650]1999.

[131651]

[131652]

[131653]11748.

[131654]Vassar, R.; Bennett, B. D.; Babu-Khan, S.; Kahn, S.; Men-

diaz,

[131655]E. A.; Dents, P.; Taplow, D. B.; Ross, S.; Amaranta, P.;

Loeloff,

[131656]R.; Luo, Y.; Fisher, S.; and 12 others: Beta-secretase cleav-

age

[131657]of Alzheimer's amyloid precursor protein by the trans-

membrane aspartic

[131658]protease BACE. Science 286: 735-741, 1999.

[131659]

[131660]

[131661]11749.

[131662]Yan, R.; Bienkowski, M. J.; Shuck, M. E.; Miao, H.; Tory, M.

C.;

[131663]Pauley, A. M.; Brashler, J. R.; Stratman, N. C.; Mathews, W.

R.; Buhl,

[131664]A. E.; Carter, D. B.; Tomasselli, A. G.; Parodi, L. A.; Hein-

rikson,

[131665]R. L.; Gurney, M. E.: Membrane-anchored aspartyl pro-

tease with Alzheimer's

[131666]disease beta-secretase activity. Nature 402: 533-536,

1999.

[131667]

[131668]

[131669]11750.

[131670]Jagla, K.; Dolle, P.; Mattei, M.-G.; Jagla, T.; Schuhbaur, B.;

[131671]Dretzen, G.; Bellard, F.; Bellard, M.: Mouse Lbx1 and human LBX1

[131672]define a novel mammalian homeobox gene family related to the Drosophila

[131673]lady bird genes. Mech. Dev. 53: 345–356, 1995.

[131674]

[131675]

[131676]11751.

[131677]Mennerich, D.; Schafer, K.; Braun, T.: Pax-3 is necessary but

[131678]not sufficient for lbx1 expression in myogenic precursor cells of

[131679]the limb. Mech. Dev. 73: 147–158, 1998.

[131680]

[131681]

[131682]11752.

[131683]Schafer, K.; Braun, T.: Early specification of limb muscle precursor

[131684]cells by the homeobox gene Lbx1h. Nature Genet. 23: 213–216, 1999.

[131685]

[131686]

[131687]11753.

[131688]Wilson, P. J.; McGlinn, E.; Marsh, A.; Evans, T.; Arnold, J.; Wright,

[131689]K.; Biden, K.; Young, J.; Wainwright, B.; Wicking, C.; Chenevix-Trench,

[131690]G.: Sequence variants of DLC1 in colorectal and ovarian tumours. Hum.

[131691]Mutat. 15: 156–165, 2000.

[131692]

[131693]

[131694]11754.

[131695]Yuan, B.-Z.; Miller, M. J.; Keck, C. L.; Zimonjic, D. B.; Thorgeirsson,

[131696]S. S.; Popescu, N. C.: Cloning, characterization, and chromosomal

[131697]localization of a gene frequently deleted in human liver cancer (DLC-1)

[131698]homologous to rat RhoGAP. Cancer Res. 58: 2196-2199, 1998.

[131699]

[131700]

[131701]11755.

[131702]Yuan, B.-Z.; Yang, Y.; Keck-Waggoner, C. L.; Zimonjic, D. B.; Thorgeirsson,

[131703]S. S.; Popescu, N. C.: Assignment and cloning of mouse Arhgap7 to

[131704]chromosome 8A4-B2, a conserved syntenic region of human chromosome

[131705]8p22-p21. Cytogenet. Cell Genet. 87: 189-190, 1999.

[131706]

[131707]

[131708]11756.

[131709]Arnould, C.; Philippe, C.; Bourdon, V.; Gregoire, M. J.; Berger,

[131710]R.; Jonveaux, P.: The signal transducer and activator of transcription

[131711]STAT5b gene is a new partner of retinoic acid receptor al-
pha in acute

[131712]promyelocytic-like leukaemia. Hum. Molec. Genet. 8:
1741-1749, 1999.

[131713]

[131714]

[131715]11757.

[131716]Lin, J.-X.; Mietz, J.; Modi, W. S.; John, S.; Leonard, W. J.:

[131717]Cloning of human Stat5B: reconstitution of interleukin-
2-induced Stat5A

[131718]and Stat5B DNA binding activity in COS-7 cells. J. Biol.
Chem. 271:

[131719]10738-10744, 1996.

[131720]

[131721]

[131722]11758.

[131723]Shinoura, N.; Shamraj, O. I.; Hugenholtz, H.; Zhu, J. G.;
McBlack,

[131724]P.; Warnick, R.; Tew, J. J.; Wani, M. A.; Menon, A. G.: Iden-
tification

[131725]and partial sequence of a cDNA that is differentially expressed in

[131726]human brain tumors. Cancer Lett. 89: 215–221, 1995.

[131727]

[131728]

[131729]11759.

[131730]Rumsby, G.; Cregeen, D. P.: Identification and expression of a

[131731]cDNA for human hydroxypyruvate/glyoxylate reductase. Biochim. Biophys.

[131732]Acta 1446: 383–388, 1999.

[131733]

[131734]

[131735]11760.

[131736]Webster, K. E.; Ferree, P. M.; Holmes, R. P.; Cramer, S. D.: Identification

[131737]of missense, nonsense, and deletion mutations in the GRHPR gene in

[131738]patients with primary hyperoxaluria type II (PH2). Hum. Genet. 107:

[131739]176–185, 2000.

[131740]

[131741]

[131742]11761.

[131743]Cremona, O.; Di Paolo, G.; Wenk, M. R.; Luthi, A.; Kim, W.
T.;

[131744]Takei, K.; Daniell, L.; Nemoto, Y.; Shears, S. B.; Flavell, R.
A.;

[131745]McCormick, D. A.; De Camilli, P.: Essential role of phos-
phoinositide

[131746]metabolism in synaptic vesicle recycling. Cell 99:
179–188, 1999.

[131747]

[131748]

[131749]11762.

[131750]Cremona, O.; Nimmakayalu, M.; Haffner, C.; Bray–Ward, P.;
Ward,

[131751]D. C.; De Camilli, P.: Assignment of SYNJ1 to human chro-
mosome 21q22.2

[131752]and Synj12 (sic) to the murine homologous region on
chromosome 16C3–4

[131753]by in situ hybridization. Cytogenet. Cell Genet. 88: 89–90,
2000.

[131754]

[131755]

[131756]11763.

[131757]Haffner, C.; Takei, K.; Chen, H.; Ringstad, N.; Hudson, A.;
Butler,

[131758]M. H.; Salcini, A. E.; Di Fiore, P. P.; De Camilli, P.: Synapto-
janin

[131759]1: localization on coated endocytic intermediates in nerve
terminals

[131760]and interaction of its 170 kDa isoform with Eps15. FEBS
Lett. 419:

[131761]175–180, 1997.

[131762]

[131763]

[131764]11764.

[131765]McPherson, P. S.; Garcia, E. P.; Slepnev, V. I.; David, C.;
Zhang,

[131766]X.; Grabs, D.; Sossin, W. S.; Bauerfeind, R.; Nemoto, Y.; De
Camilli,

[131767]P.: A presynaptic inositol-5-phosphatase. Nature 379:
353-357,

[131768]1996.

[131769]

[131770]

[131771]11765.

[131772]Saito, T.; Guan, F.; Papolos, D. F.; Lau, S.; Klein, M.; Fann,

[131773]C. S. J.; Lachman, H. M.: Mutation analysis of SYNJ1: a
possible

[131774]candidate gene for chromosome 21q22-linked bipolar
disorder. Molec.

[131775]Psychiat. 6: 387-395, 2001.

[131776]

[131777]

[131778]11766.

[131779]Aita, V. M.; Liang, X. H.; Murty, V. V. V. S.; Pincus, D. L.;
Yu,

[131780]W.; Cayanis, E.; Kalachikov, S.; Gilliam, T. C.; Levine, B.:
Cloning

[131781]and genomic organization of beclin 1, a candidate tumor

suppressor

[131782]gene on chromosome 17q21. Genomics 59: 59–65, 1999.

[131783]

[131784]

[131785]11767.

[131786]Liang, X. H.; Jackson, S.; Seaman, M.; Brown, K.; Kempkes, B.;

[131787]Hibshoosh, H.; Levine, B.: Induction of autophagy and inhibition

[131788]of tumorigenesis by beclin 1. Nature 402: 672–676, 1999.

[131789]

[131790]

[131791]11768.

[131792]Liang, X. H.; Kleeman, L. K.; Jiang, H. H.; Gordon, G.; Goldman,

[131793]J. E.; Berry, G.; Herman, B.; Levine, B.: Protection against fatal

[131794]Sindbis virus encephalitis by beclin, a novel Bcl-2-interacting protein. J.

[131795]Virol. 72: 8586–8596, 1998.

[131796]

[131797]

[131798]11769.

[131799]Liang, X. H.; Yu, J.; Brown, K.; Levine, B.: Beclin 1 contains

[131800]a leucine-rich nuclear export signal that is required for its
autophagy

[131801]and tumor suppressor function. Cancer Res. 61:
3443–3449, 2001.

[131802]

[131803]

[131804]11770.

[131805]Hatamura, I.; Kanauchi, Y.; Takahara, M.; Fujiwara, M.;
Muragaki,

[131806]Y.; Ooshima, A.; Ogino, T.: A nonsense mutation in TRPS1
in a Japanese

[131807]family with tricho-rhino-phalangeal syndrome type I.
(Letter) Clin.

[131808]Genet. 59: 366–367, 2001.

[131809]

[131810]

[131811]11771.

[131812]Hilton, M. J.; Sawyer, J. M.; Gutierrez, L.; Hogart, A.; Kung,

[131813]T. C.; Wells, D. E.: Analysis of novel and recurrent mutations responsible

[131814]for the tricho-rhino-phalangeal syndromes. J. Hum.

Genet. 47: 103-106,

[131815]2002.

[131816]

[131817]

[131818]11772.

[131819]Patel, N.; Brinkman-Van der Linden, E. C. M.; Altmann, S.

W.; Gish,

[131820]K.; Balasubramanian, S.; Timans, J. C.; Peterson, D.; Bell,

M. P.;

[131821]Bazan, J. F.; Varki, A.; Kastelein, R. A.: OB-BP1/Siglec-6: a

leptin-

[131822]and sialic acid-binding protein of the immunoglobulin su-

perfamily. J.

[131823]Biol. Chem. 274: 22729-22738, 1999. Note: Erratum: J.

Biol. Chem.

[131824]274: 28058 only, 1999.

[131825]

[131826]

[131827]11773.

[131828]Miller, A. F.; Harvey, S. A. K.; Thies, R. S.; Olson, M. S.:
Bone

[131829]morphogenetic protein-9: an autocrine/paracrine cytokine
in the liver. J.

[131830]Biol. Chem. 275: 17937-17945, 2000.

[131831]

[131832]

[131833]11774.

[131834]Scott, A. F.: Personal Communication. Baltimore, Md.
7/13/2000.

[131835]

[131836]

[131837]11775.

[131838]Lin, C.-Y.; Anders, J.; Johnson, M.; Sang, Q. A.; Dickson, R.
B.

[131839]: Molecular cloning of cDNA for matriptase, a matrix-

degrading serine

[131840]protease with trypsin-like activity. J. Biol. Chem. 274:

18231–18236,

[131841]1999.

[131842]

[131843]

[131844]11776.

[131845]Oberst, M.; Anders, J.; Xie, B.; Singh, B.; Ossandon, M.;

Johnson,

[131846]M.; Dickson, R. B.; Lin, C.-Y.: Matriptase and HAI-1 are

expressed

[131847]by normal and malignant epithelial cells in vitro and in

vivo. Am.

[131848]J. Path. 158: 1301–1311, 2001.

[131849]

[131850]

[131851]11777.

[131852]Kandil, E.; Kohda, K.; Ishibashi, T.; Tanaka, K.; Kasahara,

M.

[131853]: PA28 subunits of the mouse proteasome: primary struc-

tures and chromosomal

[131854] localization of the genes. Immunogenetics 46: 337–344, 1997.

[131855]

[131856]

[131857] 11778.

[131858] Nikaido, T.; Shimada, K.; Shibata, M.; Hata, M.; Sakamoto, M.;

[131859] Takasaki, Y.; Sato, C.; Takahashi, T.; Nishida, Y.: Cloning and nucleotide

[131860] sequence of cDNA for Ki antigen, a highly conserved nuclear protein

[131861] detected with sera from patients with systemic lupus erythematosus. Clin.

[131862] Exp. Immun. 79: 209–214, 1990.

[131863]

[131864]

[131865] 11779.

[131866] Bednarek, A. K.; Keck–Waggoner, C. L.; Daniel, R. L.; Laflin, K.

[131867] J.; Bergsagel, P. L.; Kiguchi, K.; Brenner, A. J.; Aldaz, C. M.:

[131868]WWOX, the FRA16D gene, behaves as a suppressor of tumor growth. Cancer

[131869]Res. 61: 8068–8073, 2001.

[131870]

[131871]

[131872]11780.

[131873]Bednarek, A. K.; Laflin, K. J.; Daniel, R. L.; Liao, Q.;
Hawkins,

[131874]K. A.; Aldaz, C. M.: WWOX, a novel WW domain-containing
protein mapping

[131875]to human chromosome 16q23.3–24.1, a region frequently
affected in

[131876]breast cancer. Cancer Res. 60: 2140–2145, 2000.

[131877]

[131878]

[131879]11781.

[131880]Chang, N.-S.; Pratt, N.; Heath, J.; Schultz, L.; Slevin, D.;
Carey,

[131881]G. B.; Zevotek, N.: Hyaluronidase induction of a WW domain-containing

[131882]oxidoreductase that enhances tumor necrosis factor cytotoxicity. J.

[131883]Biol. Chem. 276: 3361–3370, 2001.

[131884]

[131885]

[131886]11782.

[131887]Krummel, K. A.; Denison, S. R.; Calhoun, E.; Phillips, L. A.; Smith,

[131888]D. I.: The common fragile site FRA16D and its associated gene WWOX

[131889]are highly conserved in the mouse at Fra8E1. Genes Chromosomes Cancer 34:

[131890]154–167, 2002.

[131891]

[131892]

[131893]11783.

[131894]Miyake, A.; Mochizuki, S.; Yokoi, H.; Kohda, M.; Furuichi, K.:

[131895]New ether-a-go-go K(+) channel family members localized in human telencephalon. J.

[131896]Biol. Chem. 274: 25018–25025, 1999.

[131897]

[131898]

[131899]11784.

[131900]Acampora, D.; Postiglione, M. P.; Avantaggiato, V.; Di
Bonito,

[131901]M.; Vaccarino, F. M.; Michaud, J.; Simeone, A.: Progressive
impairment

[131902]of developing neuroendocrine cell lineages in the hy-
pothalamus of

[131903]mice lacking the Orthopedia gene. Genes Dev. 13:
2787–2800, 1999.

[131904]

[131905]

[131906]11785.

[131907]Lin, X.; State, M. W.; Vaccarino, F. M.; Greally, J.; Hass, M.;

[131908]Leckman, J. F.: Identification, chromosomal assignment,
and expression

[131909]analysis of the human homeodomain-containing gene Or-
thopedia (OTP). Genomics 60:

[131910]96–104, 1999.

[131911]

[131912]

[131913]11786.

[131914]Pessino, A.; Sivori, S.; Bottino, C.; Malaspina, A.; Morelli, L.;

[131915]Moretta, L.; Biassoni, R.; Moretta, A.: Molecular cloning of NKp46:

[131916]a novel member of the immunoglobulin superfamily involved in triggering

[131917]of natural cytotoxicity. J. Exp. Med. 188: 953–960, 1998.

[131918]

[131919]

[131920]11787.

[131921]Sivori, S.; Vitale, M.; Morelli, L.; Sanseverino, L.; Augugliaro,

[131922]R.; Bottino, C.; Moretta, L.; Moretta, A.: p46, a novel natural killer

[131923]cell-specific surface molecule that mediates cell activation. J.

[131924]Exp. Med. 186: 1129–1136, 1997.

[131925]

[131926]

[131927]11788.

[131928]Vankayalapati, R.; Wizel, B.; Weis, S. E.; Safi, H.; Lakey, D.

[131929]L.; Mandelboim, O.; Samten, B.; Porgador, A.; Barnes, P. F.:

The

[131930]NKp46 receptor contributes to NK cell lysis of mononuclear phagocytes

[131931]infected with an intracellular bacterium. J. Immun. 168: 3451–3457,

[131932]2002.

[131933]

[131934]

[131935]11789.

[131936]Gongora, C.; David, G.; Pintard, L.; Tissot, C.; Hua, T. D.; Dejean,

[131937]A.; Mehti, N.: Molecular cloning of a new interferon-induced PML

[131938]nuclear body-associated protein. J. Biol. Chem. 272:

19457–19463,

[131939]1997.

[131940]

[131941]

[131942]11790.

[131943]Mattei, M. G.; Tissot, C.; Gongora, C.; Mechti, N.: Assign-
ment

[131944]of ISG20 encoding a new interferon–induced PML nuclear
body–associated

[131945]protein, to chromosome 15q26 by in situ hybridization.
Cytogenet.

[131946]Cell Genet. 79: 286–287, 1997.

[131947]

[131948]

[131949]11791.

[131950]Pentecost, B. T.: Expression and estrogen regulation of the
HEM45

[131951]mRNA in human tumor lines and in the rat uterus. J.
Steroid Biochem.

[131952]Molec. Biol. 64: 25–33, 1998.

[131953]

[131954]

[131955]11792.

[131956]Berchtold, S.; Muhl-Zurbes, P.; Heufler, C.; Winklehner, P.;
Schuler,

[131957]G.; Steinkasserer, A.: Cloning, recombinant expression
and biochemical

[131958]characterization of the murine CD83 molecule which is
specifically

[131959]upregulated during dendritic cell maturation. FEBS Lett.
461: 211–216,

[131960]1999.

[131961]

[131962]

[131963]11793.

[131964]Fujimoto, Y.; Tu, L.; Miller, A. S.; Bock, C.; Fujimoto, M.;
Doyle,

[131965]C.; Steeber, D. A.; Tedder, T. F.: CD83 expression influ-
ences CD4+

[131966]T cell development in the thymus. Cell 108: 755–767,
2002.

[131967]

[131968]

[131969]11794.

[131970]Kozlow, E. J.; Wilson, G. L.; Fox, C. H.; Kehrl, J. H.: Sub-
tractive

[131971]cDNA cloning of a novel member of the Ig gene superfam-
ily expressed

[131972]at high levels in activated B lymphocytes. Blood 81:
454-461, 1993.

[131973]

[131974]

[131975]11795.

[131976]Olavesen, M.G.; Bentley, E.; Mason, R. V.; Stephens, R. J.;
Ragoussis,

[131977]J.: Fine mapping of 39 ESTs on human chromosome
6p23-p25. Genomics 46:

[131978]303-306, 1997.

[131979]

[131980]

[131981]11796.

[131982]Scholler, N.; Hayden-Ledbetter, M.; Hellstrom, K.-E.; Hellstrom,

[131983]I.; Ledbetter, J. A.: CD83 is a sialic acid-binding Ig-like lectin

[131984](Siglec) adhesion receptor that binds monocytes and a subset of activated

[131985]CD8(+) T cells. J. Immun. 166: 3865-3872, 2001.

[131986]

[131987]

[131988]11797.

[131989]Schinkmann, K.; Blenis, J.: Cloning and characterization of a

[131990]human STE20-like protein kinase with unusual cofactor requirements. J.

[131991]Biol. Chem. 272: 28695-28703, 1997.

[131992]

[131993]

[131994]11798.

[131995]Zhou, T.-H.; Ling, K.; Guo, J.; Zhou, H.; Wu, Y.-L.; Jing, Q.;

[131996]Ma, L.; Pei, G.: Identification of a human brain-specific

isoform

[131997]of mammalian STE20-like kinase 3 that is regulated by
cAMP-dependent

[131998]protein kinase. J. Biol. Chem. 275: 2513–2519, 2000.

[131999]

[132000]

[132001]11799.

[132002]St-Pierre, M. V.; Hagenbuch, B.; Ugele, B.; Meier, P. J.;
Stallmach,

[132003]T.: Characterization of an organic anion-transporting
polypeptide

[132004](OATP-B) in human placenta. J. Clin. Endocr. Metab. 87:
1856–1863,

[132005]2002.

[132006]

[132007]

[132008]11800.

[132009]Murthy, A.; Gonzalez-Agosti, C.; Cordero, E.; Pinney, D.;
Candia,

[132010]C.; Solomon, F.; Gusella, J.; Ramesh, V.: NHE-RF, a regula-

tory cofactor

[132011]for Na(+)-H(+) exchange, is a common interactor for
merlin and ERM

[132012](MERM) proteins. J. Biol. Chem. 273: 1273–1276, 1998.

[132013]

[132014]

[132015]11801.

[132016]Reczek, D.; Berryman, M.; Bretscher, A.: Identification of
EBP50:

[132017]a PDZ-containing phosphoprotein that associates with
members of the

[132018]ezrin–radixin–moesin family. J. Cell Biol. 139: 169–179,
1997.

[132019]

[132020]

[132021]11802.

[132022]Shenolikar, S.; Voltz, J. W.; Minkoff, C. M.; Wade, J. B.;
Weinman,

[132023]E. J.: Targeted disruption of the mouse NHERF-1 gene
promotes internalization

[132024]of proximal tubule sodium–phosphate cotransporter type

Ila and renal

[132025]phosphate wasting. Proc. Nat. Acad. Sci. 99:
11470–11475, 2002.

[132026]

[132027]

[132028]11803.

[132029]Boucher, C. A.; Winchester, C. L.; Hamilton, G. M.; Winter,
A.

[132030]D.; Johnson, K. J.; Bailey, M. E. S.: Structure, mapping and
expression

[132031]of the human gene encoding the homeodomain protein,
SIX2. Gene 247:

[132032]145–151, 2000.

[132033]

[132034]

[132035]11804.

[132036]Hasan, S.; Stucki, M.; Hassa, P. O.; Imhof, R.; Gehrig, P.;
Hunziker,

[132037]P.; Hubscher, U.; Hottiger, M. O.: Regulation of human flap
endonuclease–1

[132038]activity by acetylation through the transcriptional coacti–

vator p300. Molec.

[132039]Cell 7: 1221–1231, 2001.

[132040]

[132041]

[132042]11805.

[132043]Hiraoka, L. R.; Harrington, J. J.; Gerhard, D. S.; Lieber, M.
R.;

[132044]Hsieh, C.-L.: Sequence of human FEN-1, a structure-spe-
cific endonuclease,

[132045]and chromosomal localization of the gene (FEN1) in
mouse and human. Genomics 25:

[132046]220–225, 1995.

[132047]

[132048]

[132049]11806.

[132050]Hosfield, D. J.; Mol, C. D.; Shen, B.; Tainer, J. A.: Structure

[132051]of the DNA repair and replication endonuclease and ex-
onuclease FEN-1:

[132052]coupling DNA and PCNA binding to FEN-1 activity. Cell
95: 135–146,

[132053]1998.

[132054]

[132055]

[132056]11807.

[132057]Kucherlapati, M.; Yang, K.; Kuraguchi, M.; Zhao, J.; Lia, M.;
Heyer,

[132058]J.; Kane, M. F.; Fan, K.; Russell, R.; Brown, A. M. C.; Kneitz,
B.;

[132059]Edelmann, W.; Kolodner, R. D.; Lipkin, M.; Kucherlapati, R.:
Haploinsufficiency

[132060]of Flap endonuclease (Fen1) leads to rapid tumor progres-
sion. Proc.

[132061]Nat. Acad. Sci. 99: 9924–9929, 2002.

[132062]

[132063]

[132064]11808.

[132065]Otto, C. J.; Almqvist, E.; Hayden, M. R.; Andrew, S. E.: The
'flap'

[132066]endonuclease gene FEN1 is excluded as a candidate gene
implicated

[132067]in the CAG repeat expansion underlying Huntington disease. Clin.

[132068]Genet. 59: 122–127, 2001.

[132069]

[132070]

[132071]11809.

[132072]Spiro, C.; Pelletier, R.; Rolfsmeier, M. L.; Dixon, M. J.;
Lahue,

[132073]R. S.; Gupta, G.; Park, M. S.; Chen, X.; Mariappan, S. V. S.;
McMurray,

[132074]C. T.: Inhibition of FEN–1 processing by DNA secondary
structure

[132075]at trinucleotide repeats. Molec. Cell 4: 1079–1085, 1999.

[132076]

[132077]

[132078]11810.

[132079]Tishkoff, D. X.; Filosi, N.; Gaida, G. M.; Kolodner, R. D.: A

[132080]novel mutation avoidance mechanism dependent on *S.*
cerevisiae RAD27

[132081]is distinct from DNA mismatch repair. Cell 88: 253–263,

1997.

[132082]

[132083]

[132084]11811.

[132085]Vermeesch, J. R.; Mertens, G.; David, G.; Marynen, P.: Assignment
segment

[132086]of the human glypican gene (GPC1) to 2q35-q37 by fluorescence in situ

[132087]hybridization. Genomics 25: 327-329, 1995.

[132088]

[132089]

[132090]11812.

[132091]Ono, Y.; Ohno, M.; Shimura, Y.: Identification of a putative
RNA

[132092]helicase (HRH1), a human homolog of yeast Prp22. Molec.
Cell. Biol. 14:

[132093]7611-7620, 1994.

[132094]

[132095]

[132096]11813.

[132097]Fulop, V.; Bocskei, Z.; Polgar, L.: Prolyl oligopeptidase: an

[132098]unusual beta-propeller domain regulates proteolysis. Cell

94: 161–170,

[132099]1998.

[132100]

[132101]

[132102]11814.

[132103]Goossens, F. J.; Wauters, J. G.; Vanhoof, G. C.; Bossuyt, P.

J.;

[132104]Schatterman, K. A.; Loens, K.; Scharpe, S. L.: Subregional

mapping

[132105]of the human lymphocyte prolyl oligopeptidase gene

(PREP) to human

[132106]chromosome 6q22. Cytogenet. Cell Genet. 74: 99–101,

1996.

[132107]

[132108]

[132109]11815.

[132110]Vanhoof, G.; Goossens, F.; Hendriks, L.; De Meester, I.;

Hendriks,

[132111]D.; Vriend, G.; Van Broeckhoven, C.; Scharpe, S.: Cloning
and sequence

[132112]analysis of the gene encoding human lymphocyte prolyl
endopeptidase. Gene 149:

[132113]363–366, 1994.

[132114]

[132115]

[132116]11816.

[132117]Goldstein, L. A.; Gherzi, G.; Pineiro–Sanchez, M. L.; Salam–
one,

[132118]M.; Yeh, Y.; Flessate, D.; Chen, W.–T.: Molecular cloning of
seprase:

[132119]a serine integral membrane protease from human
melanoma. Biochim.

[132120]Biophys. Acta 1361: 11–19, 1997.

[132121]

[132122]

[132123]11817.

[132124]Mathew, S.; Scanlan, M. J.; Mohan Raj, B. K.; Murty, V. V. V.

S.;

[132125]Garin-Chesa, P.; Old, L. J.; Rettig, W. J.; Chaganti, R. S. K.:

The

[132126]gene for fibroblast activation protein alpha (FAP), a putative cell

[132127]surface-bound serine protease expressed in cancer stroma and wound

[132128]healing, maps to chromosome band 2q23. Genomics 25: 335-337, 1995.

[132129]

[132130]

[132131]11818.

[132132]Pineiro-Sanchez, M. L.; Goldstein, L. A.; Dodt, J.; Howard, L.;

[132133]Yeh, Y.; Tran, H.; Argraves, W. S.; Chen, W.-T.: Identification of

[132134]the 170-kDa melanoma membrane-bound gelatinase (seprase) as a serine

[132135]integral membrane protease. J. Biol. Chem. 272: 7595-7601, 1997.

[132136]

[132137]

[132138]11819.

[132139]Scanlan, M. J.; Raj, B. K. M.; Calvo, B.; Garin-Chesa, P.;
Sanz-Moncasi,

[132140]M. P.; Healey, J. H.; Old, L. J.; Rettig, W. J.: Molecular
cloning

[132141]of fibroblast activation protein alpha, a member of the
serine protease

[132142]family selectively expressed in stromal fibroblasts of ep-
ithelial

[132143]cancers. Proc. Nat. Acad. Sci. 91: 5657-5661, 1994.

[132144]

[132145]

[132146]11820.

[132147]Okumura, K.; Nogami, M.; Taguchi, H.; Dean, F. B.; Chen,
M.; Pan,

[132148]Z.-Q.; Hurwitz, J.; Shiratori, A.; Murakami, Y.; Ozawa, K.;
Eki, T.

[132149]: Assignment of the 36.5-kDa (RFC5), 37-kDa (RFC4),
38-kDa (RFC3),

[132150]and 40-kDa (RFC2) subunit genes of human replication
factor C to chromosome

[132151]bands 12q24.2–q24.3, 3q27, 13q12.3–q13, and 7q11.23.

Genomics 25:

[132152]274–278, 1995.

[132153]

[132154]

[132155]11821.

[132156]Peoples, R.; Perez–Jurado, L.; Wang, Y.–K.; Kaplan, P.;
Francke,

[132157]U.: The gene for replication factor C subunit 2 (RFC2) is
within

[132158]the 7q11.23 Williams syndrome deletion. (Letter) Am. J.
Hum. Genet. 58:

[132159]1370–1373, 1996.

[132160]

[132161]

[132162]11822.

[132163]Wang, Y.; Cortez, D.; Yazdi, P.; Neff, N.; Elledge, S. J.; Qin,

[132164]J.: BASC, a super complex of BRCA1–associated proteins
involved in

[132165]the recognition and repair of aberrant DNA structures.

Genes Dev. 14:

[132166]927–939, 2000.

[132167]

[132168]

[132169]11823.

[132170]Thim, L.; Woldike, H. F.; Nielsen, P. F.; Christensen, M.;
Lynch–Devaney,

[132171]K.; Podolsky, D. K.: Characterization of human and rat in-
testinal

[132172]trefoil factor produced in yeast. Biochemistry 34:
4757–4764, 1995.

[132173]

[132174]

[132175]11824.

[132176]Itoh, T.; Mori, T.; Ohkubo, H.; Yamaizumi, M.: A newly
identified

[132177]patient with clinical xeroderma pigmentosum phenotype
has a non-sense

[132178]mutation in the DDB2 gene and incomplete repair in (6–4)
photoproducts. J.

[132179]Invest. Derm. 113: 251–257, 1999.

[132180]

[132181]

[132182]11825.

[132183]Kimura, S.; Hara, Y.; Pineau, T.; Fernandez-Salguero, P.;
Fox,

[132184]C. H.; Ward, J. M.; Gonzalez, F. J.: The T/ebp null mouse:
thyroid-specific

[132185]enhancer-binding protein is essential for the organogene-
sis of the

[132186]thyroid, lung, ventral forebrain, and pituitary. Genes Dev.
10:

[132187]60-69, 1996.

[132188]

[132189]

[132190]11826.

[132191]Fernandes-Alnemri, T.; Armstrong, R. C.; Krebs, J.; Srini-
vasula,

[132192]S. M.; Wang, L.; Bullrich, F.; Fritz, L. C.; Trapani, J. A.;
Tomaselli,

[132193]K. J.; Litwack, G.; Alnemri, E. S.: In vitro activation of
CPP32

[132194]and Mch3 by Mch4, a novel human apoptotic cysteine protease containing

[132195]two FADD-like domains. Proc. Nat. Acad. Sci. 93: 7464–7469, 1996.

[132196]

[132197]

[132198]11827.

[132199]Fernandes-Alnemri, T.; Litwack, G.; Alnemri, E. S.: CPP32, a novel

[132200]human apoptotic protein with homology to Caenorhabditis elegans cell

[132201]death protein Ced-3 and mammalian interleukin-1 beta-converting enzyme. J.

[132202]Biol. Chem. 269: 30761–30764, 1994.

[132203]

[132204]

[132205]11828.

[132206]Fernando, P.; Kelly, J. F.; Balazsi, K.; Slack, R. S.; Megeney,

[132207]L. A.: Caspase 3 activity is required for skeletal muscle differentiation. Proc.

[132208]Nat. Acad. Sci. 99: 11025–11030, 2002.

[132209]

[132210]

[132211]11829.

[132212]Huang, Y.; Shin, N.–H.; Sun, Y.; Wang, K. K. W.: Molecular cloning

[132213]and characterization of a novel caspase–3 variant that attenuates

[132214]apoptosis induced by proteasome inhibition. Biochem. Biophys. Res.

[132215]Commun. 283: 762–769, 2001.

[132216]

[132217]

[132218]11830.

[132219]Kuida, K.; Zheng, T. S.; Na, S.; Kuan, C.; Yang, D.; Karasuyama,

[132220]H.; Rakio, P.; Flavell, R. A.: Decreased apoptosis in the brain and

[132221]premature lethality in CPP32–deficient mice. Nature 384: 368–372,

[132222]1996.

[132223]

[132224]

[132225]11831.

[132226]Levkau, B.; Koyama, H.; Raines, E. W.; Clurman, B. E.; Herren,

[132227]B.; Orth, K.; Roberts, J. M.; Ross, R.: Cleavage of p21(Cip1/Waf1)

[132228]and p27(Kip1) mediates apoptosis in endothelial cells through activation

[132229]of Cdk2: role of a caspase cascade. Molec. Cell 1: 553–563, 1998.

[132230]

[132231]

[132232]11832.

[132233]Nasir, J.; Theilmann, J. L.; Chopra, V.; Jones, A. M.; Walker,

[132234]D.; Rasper, D. M.; Vaillancourt, J. P.; Hewitt, J. E.; Nicholson,

[132235]D. W.; Hayden, M. R.: Localization of the cell death genes CPP32

[132236]and Mch-2 to human chromosome 4q. Mammalian
Genome 8: 56-59, 1997.

[132237]

[132238]

[132239]11833.

[132240]Tiso, N.; Pallavicini, A.; Muraro, T.; Zimbello, R.; Apolloni,

[132241]E.; Valle, G.; Lanfranchi, G.; Danieli, G. A.: Chromosomal
localization

[132242]of the human genes, CPP32, Mch2, Mch3, and Ich-1, in-
volved in cellular

[132243]apoptosis. Biochem. Biophys. Res. Commun. 225:
983-989, 1996.

[132244]

[132245]

[132246]11834.

[132247]Woo, M.; Hakem, R.; Soengas, M. S.; Duncan, G. S.;
Shahinian,

[132248]A.; Kagi, K.; Hakem, A.; McCurrach, M.; Khoo, W.; Kauf-
man, S. A.;

[132249]Senaldi, G.; Howard, T.; Lowe, S. W.; Mak, T. W.: Essential
contribution

[132250]of caspase 3/CPP32 to apoptosis and its associated nuclear changes. Genes

[132251]Dev. 12: 806–819, 1998.

[132252]

[132253]

[132254]11835.

[132255]Nicholson, D. W.; Ali, A.; Thornberry, N. A.; Vaillancourt, J.

[132256]P.; Ding, C. K.; Gallant, M.; Gareau, Y.; Griffin, P. R.; Labelle,

[132257]M.; Lazebnik, Y. A.; Munday, N. A.; Raju, S. M.; Smulson, M. E.; Yamin,

[132258]T.–T.; Yu, V. L.; Miller, D. K.: Identification and inhibition of

[132259]the ICE/CED–3 protease necessary for mammalian apoptosis. Nature 376:

[132260]37–43, 1995. MEDLINE UID: 95319529

[132261]

[132262]

[132263]11836.

[132264]Bergeron, L.; Perez, G. I.; Macdonald, G.; Shi, L.; Sun, Y.;

Juriscova,

[132265]A.; Varmuza, S.; Latham, K. E.; Flaws, J. A.; Salter, J. C. M.;

Hara,

[132266]H.; Moskowitz, M. A.; Li, E.; Greenberg, A.; Tilly, J. L.;

Yuan, J.

[132267]: Defects in regulation of apoptosis in caspase-2-deficient mice. *Genes*

[132268]*Dev.* 12: 1304-1314, 1998.

[132269]

[132270]

[132271]11837.

[132272]Garcia, C. K.; Goldstein, J. L.; Pathak, R. K.; Anderson, R. G.

[132273]W.; Brown, M. S.: Molecular characterization of a membrane transporter

[132274]for lactate, pyruvate, and other monocarboxylates: implications for

[132275]the Cori cycle. *Cell* 76: 865-873, 1994.

[132276]

[132277]

[132278]11838.

[132279]Garcia, C. K.; Li, X.; Luna, J.; Francke, U.: cDNA cloning of
[132280]the human monocarboxylate transporter 1 and chromosomal localization
[132281]of the SLC16A1 locus to 1p13.2–p12. Genomics 23:
500–503, 1994.

[132282]

[132283]

[132284]11839.

[132285]Kim, C. M.; Goldstein, J. L.; Brown, M. S.: cDNA cloning of
mev,

[132286]a mutant protein that facilitates cellular uptake of mevalonate, and

[132287]identification of the point mutation responsible for its
gain of function. J.

[132288]Biol. Chem. 267: 23113–23121, 1992.

[132289]

[132290]

[132291]11840.

[132292]Kingsmore, S. F.; Souryal, C. A.; Watson, M. L.; Patel, D. D.;

[132293]Seldin, M. F.: Physical and genetic linkage of the genes

encoding

[132294]Ly-9 and CD48 on mouse and human chromosomes 1.

Immunogenetics 42:

[132295]59-62, 1995.

[132296]

[132297]

[132298]11841.

[132299]Kozak, C. A.; Davidson, W. F.; Morse, H. C., III: Genetic and

[132300]functional relationships of the retroviral and lymphocyte

alloantigen

[132301]loci on mouse chromosome 1. Immunogenetics 19:

163-168, 1984.

[132302]

[132303]

[132304]11842.

[132305]Sandrin, M. S.; Gumley, T. P.; Henning, M. M.; Vaughan, H.

A.;

[132306]Gonez, L. J.; Trapani, J. A.; McKenzie, I. F. C.: Isolation and

characterization

[132307]of cDNA clones for mouse Ly-9. J. Immun. 149:

1636–1641, 1992.

[132308]

[132309]

[132310]11843.

[132311]Leek, J. P.; Moynihan, T. P.; Anwar, R.; Bonthron, D. T.;

Markham,

[132312]A. F.; Lench, N. J.: Assignment of Indian hedgehog (IHH) to
human

[132313]chromosome bands 2q33–q35 by in situ hybridization.

Cytogenet. Cell

[132314]Genet. 76: 187–188, 1997.

[132315]

[132316]

[132317]11844.

[132318]Porter, J. A.; Young, K. E.; Beachy, P. A.: Cholesterol modi-
fication

[132319]of hedgehog signaling proteins in animal development.

Science 274:

[132320]255–259, 1996.

[132321]

[132322]

[132323]11845.

[132324]Zhang, X. M.; Ramalho-Santos, M.; McMahon, A. P.:

Smoothened mutants

[132325]reveal redundant roles for Shh and Ihh signaling including regulation

[132326]of L/R asymmetry by the mouse node. Cell 105: 781–792, 2001.

[132327]

[132328]

[132329]11846.

[132330]Geurts, J. M. W.; Schoenmakers, E. F. P. M.; Roijer, E.; Aström,

[132331]A.-K.; Stenman, G.; van de Ven, W. J. M.: Identification of NFIB

[132332]as recurrent translocation partner gene of HMGIC in pleomorphic adenomas. Oncogene 16:

[132333]865–872, 1998.

[132334]

[132335]

[132336]11847.

[132337]Fiedorek, F. T., Jr.; Kay, E. S.: Mapping of the insulin promoter

[132338]factor 1 gene (Ipf1) to distal mouse chromosome 5. Genomics 28:

[132339]581–584, 1995.

[132340]

[132341]

[132342]11848.

[132343]Hani, E. H.; Stoffers, D. A.; Chevre, J.–C.; Durand, E.; Stanojevic,

[132344]V.; Dina, C.; Habener, J. F.; Froguel, P.: Defective mutations in

[132345]the insulin promoter factor–1 (IPF–1) gene in late-onset type 2 diabetes

[132346]mellitus. J. Clin. Invest. 104: R41–R48, 1999.

[132347]

[132348]

[132349]11849.

[132350]Tang, J. Y.; Hwang, B. J.; Ford, J. M.; Hanawalt, P. C.; Chu,

G.

[132351]: Xeroderma pigmentosum p48 gene enhances global genomic repair and

[132352]suppresses UV-induced mutagenesis. Molec. Cell 5:
737–744, 2000.

[132353]

[132354]

[132355]11850.

[132356]Bermingham, J. R., Jr.; Arden, K. C.; Naumova, A. K.;
Sapienza,

[132357]C.; Viars, C. S.; Fu, X.-D.; Khotz, J.; Manley, J. L.; Rosen-
feld,

[132358]M. G.: Chromosomal localization of mouse and human
genes encoding

[132359]the splicing factors ASF/SF2 (SFRS1) and SC-35 (SFRS2).
Genomics 29:

[132360]70–79, 1995.

[132361]

[132362]

[132363]11851.

[132364]Wang, J.; Takagaki, Y.; Manley, J. L.: Targeted disruption of

[132365]an essential vertebrate gene: ASF/SF2 is required for cell viability. Genes

[132366]Dev. 10: 2588–2599, 1996.

[132367]

[132368]

[132369]11852.

[132370]Fu, X.-D.; Maniatis, T.: Isolation of a complementary DNA that

[132371]encodes the mammalian splicing factor SC35. Science
256: 535–538,

[132372]1992.

[132373]

[132374]

[132375]11853.

[132376]Kuhlenbaeumer, G.; Meuleman, J.; Schirmacher, A.; Stoeg-
bauer, F.;

[132377]Ringelstein, E. B.; Wehnert, M.; Hoeltzenbein, M.; Broeck-
hoven, C.

[132378]V.; Timmerman, V.: Mutation analysis of a putative sialyl-
transferase

[132379]gene, the SFRS2 splicing factor gene and the c-myb ET-
locus in two

[132380]families with hereditary neuralgic amyotrophy (HNA). Ann.
Hum. Genet. 62:

[132381]397–400, 1998.

[132382]

[132383]

[132384]11854.

[132385]Wang, H.–Y.; Xu, X.; Ding, J.–H.; Bermingham, J. R., Jr.; Fu,
X.–D.

[132386]: SC35 plays a role in T cell development and alternative
splicing

[132387]of CD45. Molec. Cell 7: 331–342, 2001.

[132388]

[132389]

[132390]11855.

[132391]Hopfner, K.–P.; Karcher, A.; Craig, L.; Woo, T. T.; Carney, J.

[132392]P.; Tainer, J. A.: Structural biochemistry and interaction
architecture

[132393]of the DNA double–strand break repair Mre11 nuclease

and Rad50-ATPase. Cell 105:

[132394]473-485, 2001.

[132395]

[132396]

[132397]11856.

[132398]Stracker, T. H.; Carson, C. T.; Weitzman, M. D.: Adenovirus
oncoproteins

[132399]inactivate the Mre11-Rad50-NBS1 DNA repair complex.

Nature 418:

[132400]348-352, 2002.

[132401]

[132402]

[132403]11857.

[132404]Zhong, Q.; Chen, C.-F.; Li, S.; Chen, Y.; Wang, C.-C.; Xiao,
J.;

[132405]Chen, P.-L.; Sharp, Z. D.; Lee, W.-H.: Association of BRCA1
with

[132406]the hRad50-hMre11-p95 complex and the DNA damage
response. Science 285:

[132407]747-750, 1999.

[132408]

[132409]

[132410]11858.

[132411]Zhu, X.-D.; Kuster, B.; Mann, M.; Petrini, J. H. J.; de Lange,

[132412]T.: Cell-cycle-regulated association of

RAD50/MRE11/NBS1 with TRF2

[132413]and human telomeres. *Nature Genet.* 25: 347–352, 2000.

[132414]

[132415]

[132416]11859.

[132417]Virbasius, C. A.; Virbasius, J. V.; Scarpulla, R. C.: NRF-1,

an

[132418]activator involved in nuclear-mitochondrial interactions,

utilizes

[132419]a new DNA-binding domain conserved in a family of de-

velopmental regulators. *Genes*

[132420]*Dev.* 7: 2431–2445, 1993.

[132421]

[132422]

[132423]11860.

[132424]Choi, S. S.; Park, I.-C.; Yun, J. W.; Sung, Y. C.; Hong, S.-I.;

[132425]Shin, H.-S.: A novel Bcl-2 related gene, Bfl-1, is overex-
pressed

[132426]in stomach cancer and preferentially expressed in bone
marrow. Oncogene 11:

[132427]1693-1698, 1995.

[132428]

[132429]

[132430]11861.

[132431]Choi, S. S.; Park, S. H.; Kim, U.-J.; Shin, H.-S.: Bfl-1, a Bcl-
2-related

[132432]gene, is the human homolog of the murine A1, and maps
to chromosome

[132433]15q24.3. Mammalian Genome 8: 781-782, 1997.

[132434]

[132435]

[132436]11862.

[132437]D'Sa-Eipper, C.; Subramanian, T.; Chinnadurai, G.: Bfl-1, a

bcl-2

[132438]homologue, suppresses p53-induced apoptosis and ex-
hibits potent cooperative

[132439]transforming activity. Cancer Res. 56: 3879-3882, 1996.

[132440]

[132441]

[132442]11863.

[132443]Lin, E. Y.; Kozak, C. A.; Orlofsky, A.; Prystowsky, M. B.:

The

[132444]bcl-2 family member, Bcl2al, maps to mouse chromosome
9 and human

[132445]chromosome 15. Mammalian Genome 8: 293-294, 1997.

[132446]

[132447]

[132448]11864.

[132449]Lin, E. Y.; Orlofsky, A.; Berger, M. S.; Prystowsky, M. B.:

Characterization

[132450]of A1, a novel hemopoietic-specific early-response gene
with sequence

[132451]similarity to bcl-2. J. Immun. 151: 1979-1988, 1993.

[132452]

[132453]

[132454]11865.

[132455]Lin, E. Y.; Orlofsky, A.; Wang, H.-G.; Reed, J. C.; Prys-
towsky,

[132456]M. B.: A1, a Bcl-2 family member, prolongs cell survival
and permits

[132457]myeloid differentiation. Blood 87: 983-992, 1996.

[132458]

[132459]

[132460]11866.

[132461]Xiang, Z.; Ahmed, A. A.; Moller, C.; Nakayama, K.;
Hatakeyama,

[132462]S.; Nilsson, G.: Essential role of the prosurvival bcl-2 ho-
mologue

[132463]A1 in mast cell survival after allergic activation. J. Exp.
Med. 194:

[132464]1561-1569, 2001.

[132465]

[132466]

[132467]11867.

[132468]Bass, B. L.; Weintraub, H.: An unwinding activity that covalently

[132469]modifies its double-stranded RNA substrate. Cell 55:
1089-1098,

[132470]1988.

[132471]

[132472]

[132473]11868.

[132474]Kim, U.; Wang, Y.; Sanford, T.; Zeng, Y.; Nishikura, K.:
Molecular

[132475]cloning of cDNA for double-stranded RNA adenosine
deaminase, a candidate

[132476]enzyme for nuclear RNA editing. Proc. Nat. Acad. Sci. 91:
11457-11461,

[132477]1994.

[132478]

[132479]

[132480]11869.

[132481]O'Connell, M. A.; Krause, S.; Higuchi, M.; Hsuan, J. J.;

Totty,

[132482]N. F.; Jenny, A.; Keller, W.: Cloning of cDNAs encoding
mammalian

[132483]double-stranded RNA-specific adenosine deaminase.

Molec. Cell. Biol. 15:

[132484]1389–1397, 1995.

[132485]

[132486]

[132487]11870.

[132488]Patterson, J. B.; Samuel, C. E.: Expression and regulation
by

[132489]interferon of a double-stranded-RNA-specific adenosine deaminase from

[132490]human cells: evidence for two forms of the deaminase. Molec. Cell.

[132491]Biol. 15: 5376–5388, 1995.

[132492]

[132493]

[132494]11871.

[132495]Wang, Q.; Khillan, J.; Gadue, P.; Nishikura, K.: Requirement of

[132496]the RNA editing deaminase ADAR1 gene for embryonic erythropoiesis. Science 290:

[132497]1765–1768, 2000.

[132498]

[132499]

[132500]11872.

[132501]Wang, Y.; Zeng, Y.; Murray, J. M.; Nishikura, K.: Genomic organization

[132502]and chromosomal location of the human dsRNA adenosine deaminase gene:

[132503]the enzyme for glutamate-activated ion channel RNA editing. J. Molec.

[132504]Biol. 254: 184-195, 1995.

[132505]

[132506]

[132507]11873.

[132508]Weier, H.-U. G.; George, C. X.; Greulich, K. M.; Samuel, C. E.

[132509]: The interferon-inducible, double-stranded RNA-specific adenosine

[132510]deaminase gene (DSRAD) maps to human chromosome 1q21.1-21.2. Genomics 30:

[132511]372-375, 1995.

[132512]

[132513]

[132514]11874.

[132515]Weier, H.-U. G.; George, C. X.; Lersch, R. A.; Breitwieser, S.;

[132516]Cheng, J.-F.; Samuel, C. E.: Assignment of the RNA-specific adenosine

[132517]deaminase gene (Adar) to mouse chromosome 3F2 by in

situ hybridization. Cytogenet.

[132518]Cell Genet. 89: 214–215, 2000.

[132519]

[132520]

[132521]11875.

[132522]Kawagoe, H.; Soma, O.; Goji, J.; Nishimura, N.; Narita, M.;
Inazawa,

[132523]J.; Nakamura, H.; Sano, K.: Molecular cloning and chromo-
somal assignment

[132524]of the human brain-type phosphodiesterase I/nucleotide
pyrophosphatase

[132525]gene (PDNP2). Genomics 30: 380–384, 1995.

[132526]

[132527]

[132528]11876.

[132529]Murata, J.; Lee, H. Y.; Clair, T.; Krutzsch, H. C.; Arestad, A.

[132530]A.; Sobel, M. E.; Liotta, L. A.; Stracke, M. L.: cDNA cloning
of

[132531]human tumor motility-stimulating protein, autotaxin, re-
veals a homology

[132532]with phosphodiesterases. J. Biol. Chem. 269:
30479–30484, 1994.

[132533]

[132534]

[132535]11877.

[132536]Narita, M.; Goji, J.; Nakamura, H.; Sano, K. Molecular
cloning,

[132537]expression, and localization of a brain-specific phospho-
diesterase

[132538]I/nucleotide (PD-I-alpha) from rat brain. J. Biol. Chem.
269: 28235–28242,

[132539]1994.

[132540]

[132541]

[132542]11878.

[132543]Piao, J.-H.; Matsuda, Y.; Nakamura, H.; Sano, K.: Assign-
ment of

[132544]PdnP2, the gene encoding phosphodiesterase I/nucleotide
pyrophosphatase

[132545]2, to mouse chromosome 15D2. Cytogenet. Cell Genet.
87: 172–174,

[132546]1999.

[132547]

[132548]

[132549]11879.

[132550]Tokumura, A.; Majima, E.; Kariya, Y.; Tominaga, K.;
Kogure, K.;

[132551]Yasuda, K.; Fukuzawa, K.: Identification of human plasma
lysophospholipase

[132552]D, a lysophosphatidic acid-producing enzyme, as auto-
taxin, a multifunctional

[132553]phosphodiesterase. J. Biol. Chem. 277: 39436-39442,
2002.

[132554]

[132555]

[132556]11880.

[132557]Lin, Q.; Schwarz, J.; Bucana, C.; Olson, E. N.: Control of
mouse

[132558]cardiac morphogenesis and myogenesis by transcription
factor MEF2C. Science 276:

[132559]1404-1407, 1997.

[132560]

[132561]

[132562]11881.

[132563]Erickson, H. P.: A tenascin knockout with a phenotype.

Nature

[132564]Genet. 17: 5–7, 1997.

[132565]

[132566]

[132567]11882.

[132568]Akama, T. O.; Nakagawa, H.; Sugihara, K.; Narisawa, S.;

Ohyama,

[132569]C.; Nishimura, S.–I.; O'Brien, D. A.; Moremen, K. W.; Millan,

J. L.;

[132570]Fukuda, M. N.: Germ cell survival through carbohydrate-me-

diated interaction

[132571]with Sertoli cells. Science 295: 124–127, 2002.

[132572]

[132573]

[132574]11883.

[132575]Bai, R.-Y.; Koester, C.; Ouyang, T.; Hahn, S. A.; Hammer-
schmidt,

[132576]M.; Peschel, C.; Duyster, J.: SMIF, a Smad4-interacting
protein that

[132577]functions as a co-activator in TGF-beta signalling. Nature
Cell Biol. 4:

[132578]181-190, 2002.

[132579]

[132580]

[132581]11884.

[132582]Akiyama, Y.; Tsubouchi, N.; Yuasa, Y.: Frequent somatic
mutations

[132583]of hMSH3 with reference to microsatellite instability in
hereditary

[132584]nonpolyposis colorectal cancer. Biochem. Biophys. Res.
Commun. 236:

[132585]248-252, 1997.

[132586]

[132587]

[132588]11885.

[132589]Chittenden, T.; Harrington, E. A.; O'Connor, R.; Fleming-
ton, C.;

[132590]Lutz, R. J.; Evan, G. I.; Guild, B. C.: Induction of apoptosis
by

[132591]the Bcl-2 homologue Bak. Nature 374: 733-736, 1995.

[132592]

[132593]

[132594]11886.

[132595]Herberg, J. A.; Phillips, S.; Beck, S.; Jones, T.; Sheer, D.;
Wu,

[132596]J. J.; Prochazka, V.; Barr, P. J.; Kiefer, M. C.; Trowsdale, J.:

[132597]Genomic structure and domain organisation of the human
Bak gene. Gene 211:

[132598]87-94, 1998.

[132599]

[132600]

[132601]11887.

[132602]Kiefer, M. C.; Brauer, M. J.; Powers, V. C.; Wu, J. J.; Uman-
sky,

[132603]S. R.; Tomei, L. D.; Barr, P. J.: Modulation of apoptosis by

the

[132604]widely distributed Bcl-2 homologue Bak. Nature 374:
736-739, 1995.

[132605]

[132606]

[132607]11888.

[132608]Ulrich, E.; Kauffmann-Zeh, A.; Hueber, A.-O.; Williamson,
J.; Chittenden,

[132609]T.; Ma, A.; Evan, G.: Gene structure, cDNA sequence, and
expression

[132610]of murine Bak, a proapoptotic Bcl-2 family member. Ge-
nomics 44:

[132611]195-200, 1997.

[132612]

[132613]

[132614]11889.

[132615]Higuchi, M.; Maas, S.; Single, F. N.; Hartner, J.; Rozov, A.;
Burnashev,

[132616]N.; Feldmeyer, D.; Sprengel, R.; Seeburg, P. H.: Point mu-
tation in

[132617]an AMPA receptor gene rescues lethality in mice deficient

in the RNA-editing

[132618]enzyme ADAR2. Nature 406: 78–81, 2000.

[132619]

[132620]

[132621]11890.

[132622]Lai, F.; Chen, C.-X.; Carter, K. C.; Nishikura, K.: Editing of

[132623]glutamate receptor B subunit ion channel RNAs by four al-
ternatively

[132624]spliced DRADA2 double-stranded RNA adenosine deami-
nases. Molec. Cell.

[132625]Biol. 17: 2413–2424, 1997.

[132626]

[132627]

[132628]11891.

[132629]Melcher, T.; Maas, S.; Herb, A.; Sprengel, R.; Seeburg, P.
H.;

[132630]Higuchi, M.: A mammalian RNA editing enzyme. Nature
379: 460–463,

[132631]1996.

[132632]

[132633]

[132634]11892.

[132635]Mittaz, L.; Scott, H. S.; Rossier, C.; Seeburg, P. H.; Higuchi,

[132636]M.; Antonarakis, S. E.: Cloning of a human RNA editing
deaminase

[132637](ADARB1) of glutamate receptors that maps to chromo-
some 21q22.3. Genomics 41:

[132638]210–217, 1997.

[132639]

[132640]

[132641]11893.

[132642]O'Connell, M. A.; Gerber, A.; Keller, W.: Purification of hu-
man

[132643]double-stranded RNA-specific editase 1 (hRED1) involved
in editing

[132644]of brain glutamate receptor B pre-mRNA. J. Biol. Chem.
272: 473–478,

[132645]1997.

[132646]

[132647]

[132648]11894.

[132649]Villard, L.; Tassone, F.; Haymowicz, M.; Welborn, R.; Gardiner,

[132650]K.: Map location, genomic organization and expression patterns of

[132651]the human RED1 RNA editase. Somat. Cell Molec. Genet. 23: 135–145,

[132652]1997.

[132653]

[132654]

[132655]11895.

[132656]Yang, J.–H.; Sklar, P.; Axel, R.; Maniatis, T.: Purification and

[132657]characterization of a human RNA adenosine deaminase for glutamate

[132658]receptor B pre–mRNA editing. Proc. Nat. Acad. Sci. 94: 4354–4359,

[132659]1997.

[132660]

[132661]

[132662]11896.

[132663]Lu, C.-M.; Han, J.; Rado, T. A.; Brown, G. B.: Differential expression

[132664]of two sodium channel subtypes in human brain. FEBS Letters 303:

[132665]53–58, 1992.

[132666]

[132667]

[132668]11897.

[132669]Zhao, Z.; Lee, C.-C.; Monckton, D. G.; Yazdani, A.; Coolbaugh,

[132670]M. I.; Li, X.; Bailey, J.; Shen, Y.; Caskey, C. T.: Characterization

[132671]and genomic mapping of genes and pseudogenes of a new human protein

[132672]tyrosine phosphatase. Genomics 35: 172–181, 1996.

[132673]

[132674]

[132675]11898.

[132676]Chopra, V. S.; Metzler, M.; Rasper, D. M.; Engqvist-Goldstein,

[132677]A. E. Y.; Singaraja, R.; Gan, L.; Fichter, K. M.; McCutcheon, K.;

[132678]Drubin, D.; Nicholson, D. W.; Hayden, M. R.: HIP12 is a non-proapoptotic

[132679]member of a gene family including HIP1, an interacting protein with

[132680]huntingtin. Mammalian Genome 11: 1006–1015, 2000.

[132681]

[132682]

[132683]11899.

[132684]Borras, T.; Rowlette, L. L. S.; Tamm, E. R.; Gottanka, J.; Epstein,

[132685]D. L.: Effects of elevated intraocular pressure on outflow facility

[132686]and TIGR/MYOC expression in perfused human anterior segments. Invest.

[132687]Ophthal. Vis. Sci. 43: 33–40, 2002.

[132688]

[132689]

[132690]11900.

[132691]Cerretti, D. P.; Nelson, N.: Characterization of the genes
for

[132692]mouse LERK-3/Ephrin-A3 (Epl3), mouse LERK-
4/Ephrin-A4 (Epl4), and

[132693]human LERK-6/Ephrin-A2 (EPLG6): conservation of in-
tron/exon structure. Genomics 47:

[132694]131-135, 1998.

[132695]

[132696]

[132697]11901.

[132698]Ma, T.; Yang, B.; Kuo, W.-L.; Verkman, A. S.: cDNA cloning
and

[132699]gene structure of a novel water channel expressed exclu-
sively in human

[132700]kidney: evidence for a gene cluster of aquaporins at chro-
mosome locus

[132701]12q13. Genomics 35: 543-550, 1996.

[132702]

[132703]

[132704]11902.

[132705]Yasui, M.; Kwon, T.-H.; Knepper, M. A.; Nielsen, S.; Agre, P.:

[132706]Aquaporin-6: an intracellular vesicle water channel protein in renal

[132707]epithelia. Proc. Nat. Acad. Sci. 96: 5808-5813, 1999.

[132708]

[132709]

[132710]11903.

[132711]Shan, X.; Bourdeau, A.; Rhoton, A.; Wells, D. E.; Cohen, E. H.;

[132712]Landgraf, B. E.; Palfree, R. G. E.: Characterization and mapping

[132713]to human chromosome 8q24.3 of Ly-6-related gene 9804 encoding an apparent

[132714]homologue of mouse TSA-1. J. Immun. 160: 197-208, 1998.

[132715]

[132716]

[132717]11904.

[132718]Bova, G. S.; MacGrogan, D.; Levy, A.; Pin, S. S.; Bookstein, R.;

[132719]Isaacs, W. B.: Physical mapping of chromosome 8p22 markers and their

[132720]homozygous deletion in a metastatic prostate cancer. Genomics 35:

[132721]46–54, 1996.

[132722]

[132723]

[132724]11905.

[132725]An, S.; Yang, J.; Xia, M.; Goetzl, E. J.: Cloning and expression

[132726]of the EP2 subtype of human receptors for prostaglandin E2. Biochem.

[132727]Biophys. Res. Comm. 197: 263–270, 1993.

[132728]

[132729]

[132730]11906.

[132731]Bastien, L.; Sawyer, N.; Grygorczyk, R.; Metters, K. M.; Adam,

[132732]M.: Cloning, functional expression and characterization of the human

[132733]prostaglandin E2 receptor EP2 subtype. J. Biol. Chem. 269: 11873–11877,

[132734]1994.

[132735]

[132736]

[132737]11907.

[132738]Coleman, R. A.; Grix, S. P.; Head, S. A.; Louttit, J. B.; Mallett,

[132739]A.; Sheldrick, R. L. G.: A novel inhibitory prostanoid receptor in

[132740]piglet saphenous vein. Prostaglandins 47: 151–168, 1994.

[132741]

[132742]

[132743]11908.

[132744]Foord, S. M.; Marks, B.; Stolz, M.; Bufflier, E.; Fraser, N. J.;

[132745]Lee, M. G.: The structure of the prostaglandin EP4 receptor gene

[132746]and related pseudogenes. Genomics 35: 182–188, 1996.

[132747]

[132748]

[132749]11909.

[132750]Mori, K.; Tanaka, I.; Kotani, M.; Miyaoka, F.; Sando, T.;
Muro,

[132751]S.; Sasaki, Y.; Nakagawa, O.; Ogawa, Y.; Usui, T.; Ozaki, S.;
Ichikawa,

[132752]A.; Narumiya, S.; Nakao, K.: Gene expression of the hu-
man prostaglandin

[132753]E receptor EP4 subtype: differential regulation in monocy-
toid and

[132754]lymphoid lineage cells by phorbol ester. J. Molec. Med. 74:
333-336,

[132755]1996.

[132756]

[132757]

[132758]11910.

[132759]Regan, J. W.; Bailey, T. J.; Pepperl, D. J.; Pierce, K. L.; Boga-
rdus,

[132760]A. M.; Donello, J. E.; Fairbairn, C. E.; Kedzie, K. M.; Wood-
ward,

[132761]D. F.; Gil, D. W.: Cloning of a novel human prostaglandin receptor

[132762]with characteristics of the pharmacologically defined EP2 subtype. Molec.

[132763]Pharm. 46: 213–220, 1994.

[132764]

[132765]

[132766]11911.

[132767]Segi, E.; Sugimoto, Y.; Yamasaki, A.; Aze, Y.; Oida, H.; Nishimura,

[132768]T.; Murata, T.; Matsuoka, T.; Ushikubi, F.; Hirose, M.; Tanaka, T.;

[132769]Yoshida, N.; Narumiya, S.; Ichikawa, A.: Patent ductus arteriosus

[132770]and neonatal death in prostaglandin receptor EP4-deficient mice. Biochem.

[132771]Biophys. Res. Comm. 246: 7–12, 1998.

[132772]

[132773]

[132774]11912.

- [132775]Yoshida, K.; Oida, H.; Kobayashi, T.; Maruyama, T.;
Tanaka, M.;
- [132776]Katayama, T.; Yamaguchi, K.; Segi, E.; Tsuboyama, T.;
Matsushita,
- [132777]M.; Ito, K.; Ito, Y.; Sugimoto, Y.; Ushikubi, F.; Ohuchida, S.;
Kondo,
- [132778]K.; Nakamura, T.; Narumiya, S.: Stimulation of bone for-
mation and
- [132779]prevention of bone loss by prostaglandin E EP4 receptor
activation. Proc.
- [132780]Nat. Acad. Sci. 99: 4580–4585, 2002.
- [132781]
- [132782]
- [132783]11913.
- [132784]Jadayel, D. M.; Osborne, L. R.; Coignet, L. J. A.; Zani, V. J.;
- [132785]Tsui, L.-C.; Scherer, S. W.; Dyer, M. J. S.: The BCL7 gene
family:
- [132786]deletion of BCL7B in Williams syndrome. Gene 224:
35–44, 1998.
- [132787]
- [132788]

[132789]11914.

[132790]Arima, T.; Drewell, R. A.; Arney, K. L.; Inoue, J.; Makita, Y.;

[132791]Hata, A.; Oshimura, M.; Wake, N.; Surani, M. A.: A con-
served imprinting

[132792]control region at the HYMAI/ZAC domain is implicated in
transient

[132793]neonatal diabetes mellitus. Hum. Molec. Genet. 10:
1475–1483, 2001.

[132794]

[132795]

[132796]11915.

[132797]Joung, I.; Strominger, J. L.; Shin, J.: Molecular cloning of a

[132798]phosphotyrosine-independent ligand of the p56-lck SH2
domain. Proc.

[132799]Nat. Acad. Sci. 93: 5991–5995, 1996.

[132800]

[132801]

[132802]11916.

[132803]Laurin, N.; Brown, J. P.; Morissette, J.; Raymond, V.: Recur-

rent

[132804]mutation of the gene encoding sequestosome 1
(SQSTM1/p62) in Paget

[132805]disease of bone. Am. J. Hum. Genet. 70: 1582–1588,
2002.

[132806]

[132807]

[132808]11917.

[132809]Park, I.; Chung, J.; Walsh, C. T.; Yun, Y.; Strominger, J. L.;

[132810]Shin, J.: Phosphotyrosine-independent binding of a
62-kDa protein

[132811]to the src homology 2 (SH2) domain of p56-lck and its
regulation by

[132812]phosphorylation of ser-59 in the lck unique N-terminal
region. Proc.

[132813]Nat. Acad. Sci. 92: 12338–12342, 1995.

[132814]

[132815]

[132816]11918.

[132817]Vadlamudi, R. K.; Joung, I.; Strominger, J. L.; Shin, J.: p62,

[132818]a phosphotyrosine-independent ligand of the SH2 domain
of p56-lck,

[132819]belongs to a new class of ubiquitin-binding proteins. J.
Biol. Chem. 271:

[132820]20235-20237, 1996.

[132821]

[132822]

[132823]11919.

[132824]Akbar, G. K. M.; Dasari, V. R.; Webb, T. E.; Ayyanathan, K.;
Pillarisetti,

[132825]K.; Sandhu, A. K.; Athwal, R. S.; Daniel, J. L.; Ashby, B.;
Barnard,

[132826]E. A.; Kunapuli, S. P.: Molecular cloning of a novel P2
purinoceptor

[132827]from human erythroleukemia cells. J. Biol. Chem. 271:
18363-18367,

[132828]1996.

[132829]

[132830]

[132831]11920.

[132832]Chen, X.-S.; Sheller, J. R.; Johnson, E. N.; Funk, C. D.: Role
[132833]of leukotrienes revealed by targeted disruption of
5-lipoxygenase
[132834]gene. Nature 372: 179–182, 1994.

[132835]

[132836]

[132837]11921.

[132838]Devchand, P. R.; Keller, H.; Peters, J. M.; Vazquez, M.;
Gonzalez,

[132839]F. J.; Wahli, W.: The PPAR alpha-leukotriene B4 pathway to
inflammation

[132840]control. Nature 384: 39–43, 1996.

[132841]

[132842]

[132843]11922.

[132844]Kato, K.; Yokomizo, T.; Izumi, T.; Shimizu, T.: Cell-
specific

[132845]transcriptional regulation of human leukotriene B4 recep-
tor gene. J.

[132846]Exp. Med. 192: 413–420, 2000.

[132847]

[132848]

[132849]11923.

[132850]Owman, C.; Nilsson, C.; Lolait, S. J.: Cloning of cDNA encoding

[132851]a putative chemoattractant receptor. Genomics 37: 187–194, 1996.

[132852]

[132853]

[132854]11924.

[132855]Samuelsson, B.; Dahlen, S. E.; Lindgren, J. A.; Rouzer, C. A.;

[132856]Serhan, C. N.: Leukotrienes and lipoxins: structures, biosynthesis,

[132857]and biological effects. Science 237: 1171–1176, 1987.

[132858]

[132859]

[132860]11925.

[132861]Yokomizo, T.; Izumi, T.; Chang, K.; Takuwa, Y.; Shimizu,

T.: A

[132862]G-protein-coupled receptor for leukotriene B₄ that mediates chemotaxis. *Nature* 387:

[132863]620–624, 1997.

[132864]

[132865]

[132866]11926.

[132867]Fernandes-Alnemri, T.; Litwack, G.; Alnemri, E. S.: Mch2, a new

[132868]member of the apoptotic Ced-3/Ice cysteine protease gene family. *Cancer*

[132869]*Res.* 55: 2737–2742, 1995.

[132870]

[132871]

[132872]11927.

[132873]Orth, K.; Chinnaiyan, A. M.; Garg, M.; Froelich, C. J.; Dixit,

[132874]V. M.: The CED-3/ICE-like protease Mch2 is activated during apoptosis

[132875]and cleaves the death substrate lamin A. *J. Biol. Chem.* 271: 16443–16446,

[132876]1996.

[132877]

[132878]

[132879]11928.

[132880]Verhaegh, G. W. C. T.; Jongmans, W.; Jaspers, N. G. J.;
Natarajan,

[132881]A. T.; Oshimura, M.; Lohman, P. H. M.; Zdzienicka, M. Z.: A
gene

[132882]that regulates DNA replication in response to DNA dam-
age is located

[132883]on human chromosome 4q. Am. J. Hum. Genet. 57:
1095–1103, 1995.

[132884]

[132885]

[132886]11929.

[132887]Kubo, Y.; Reuveny, E.; Slesinger, P. A.; Jan, Y. N.; Jan, L. Y.

[132888]: Primary structure and functional expression of a rat G-
protein-coupled

[132889]muscarinic potassium channel. Nature 364: 802–806,
1993.

[132890]

[132891]

[132892]11930.

[132893]Schoots, O.; Voskoglou, T.; Van Tol, H. H. M.: Genomic organization

[132894]and promoter analysis of the human G-protein-coupled K⁺ channel Kir3.1

[132895](KCNJ3/HGIRK1). Genomics 39: 279–288, 1997.

[132896]

[132897]

[132898]11931.

[132899]Schoots, O.; Yue, K.-T.; MacDonald, J. F.; Hampson, D. R.; Nobrega,

[132900]J. N.; Dixon, L. M.; Van Tol, H. H. M.: Cloning of a G protein-activated

[132901]inwardly rectifying potassium channel from human cerebellum. Molec.

[132902]Brain Res. 39: 23–30, 1996.

[132903]

[132904]

[132905]11932.

[132906]Stoffel, M.; Espinosa, R., III; Powell, K. L.; Philipson, L. H.;

[132907]Le Beau, M. M.; Bell, G. I.: Human G-protein-coupled inwardly rectifying

[132908]potassium channel (GIRK1) gene (KCNJ3): localization to chromosome

[132909]2 and identification of a simple tandem repeat polymorphism. Genomics 21:

[132910]254–256, 1994.

[132911]

[132912]

[132913]11933.

[132914]Phillips, H. A.; Scheffer, I. E.; Berkovic, S. F.; Hollway, G.

[132915]E.; Sutherland, G. R.; Mulley, J. C.: Localization of a gene for

[132916]autosomal dominant nocturnal frontal lobe epilepsy to chromosome 20q13.2. Nature

[132917]Genet. 10: 117–118, 1995.

[132918]

[132919]

[132920]11934.

[132921]Phillips, H. A.; Scheffer, I. E.; Crossland, K. M.; Bhatia, K.

[132922]P.; Fish, D. R.; Marsden, C. D.; Howell, S. J. L.; Stephenson,
J.

[132923]B. P.; Tolmie, J.; Plazzi, G.; Eeg-Olofsson, O.; Singh, R.;
Lopes-Cendes,

[132924]I.; Andermann, E.; Andermann, F.; Berkovic, S. F.; Mulley, J.
C.:

[132925]Autosomal dominant nocturnal frontal-lobe epilepsy: ge-
netic heterogeneity

[132926]and evidence for a second locus at 15q24. Am. J. Hum.
Genet. 63:

[132927]1108–1116, 1998.

[132928]

[132929]

[132930]11935.

[132931]D'Arcangelo, G.: Personal Communication. Nutley, N. J.
6/2/1995.

[132932]

[132933]

[132934]11936.

[132935]D'Arcangelo, G.; Homayouni, R.; Keshvara, L.; Rice, D. S.;
Sheldon,

[132936]M.; Curran, T.: Reelin is a ligand for lipoprotein receptors.
Neuron 24:

[132937]471–479, 1999.

[132938]

[132939]

[132940]11937.

[132941]D'Arcangelo, G.; Miao, G. G.; Chen, S.–C.; Soares, H. D.;
Morgan,

[132942]J. I.; Curran, T.: A protein related to extracellular matrix
proteins

[132943]deleted in the mouse mutant reeler. Nature 374:
719–723, 1995.

[132944]

[132945]

[132946]11938.

[132947]DeSilva, U.; D'Arcangelo, G.; Braden, V. V.; Chen, J.; Miao,
G.

[132948]G.; Curran, T.; Green, E. D.: The human reelin gene: isolation, sequencing,

[132949]and mapping on chromosome 7. *Genome Res.* 7: 157–164, 1997.

[132950]

[132951]

[132952]1 1939.

[132953]Green, M. C.: Catalog of mutant genes and polymorphic loci. In:

[132954]Lyon, M. F.; Searle, A. G.: Genetic Variants and Strains of the Laboratory

[132955]Mouse. Oxford: Oxford Univ. Press (pub.) (2nd ed.): 1989.

[132956]

[132957]

[132958]1 1940.

[132959]Hack, I.; Bancila, M.; Loulier, K.; Carroll, P.; Cremer, H.: Reelin

[132960]is a detachment signal in tangential chain-migration during postnatal

[132961]neurogenesis. *Nature Neurosci.* 5: 939–945, 2002.

[132962]

[132963]

[132964]11941.

[132965]Hirotsune, S.; Takahara, T.; Sasaki, N.; Hirose, K.; Yoshiki, A.;

[132966]Ohashi, T.; Kusakabe, M.; Murakami, Y.; Muramatsu, M.; Watanabe, S.;

[132967]Nakao, K.; Katsuki, M.; Hayashizaki, Y.: The reeler gene encodes

[132968]a protein with an EGF-like motif expressed by pioneer neurons. Nature

[132969]Genet. 10: 77–83, 1995.

[132970]

[132971]

[132972]11942.

[132973]Hong, S. E.; Shugart, Y. Y.; Huang, D. T.; Al Shahwan, S.; Grant,

[132974]P. E.; Hourihane, J. O.; Martin, N. D. T.; Walsh, C. A.: Autosomal

[132975]recessive lissencephaly with cerebellar hypoplasia is associated with

[132976]human RELN mutations. Nature Genet. 26: 93–96, 2000.

Note: Erratum:

[132977]Nature Genet. 27: 225 only, 2001.

[132978]

[132979]

[132980]11943.

[132981]Hourihane, J. O.; Bennett, C. P.; Chaudhuri, R.; Robb, S. A.;

[132982]Martin, N. D. T.: A sibship with a neuronal migration defect, cerebellar

[132983]hypoplasia and congenital lymphedema. Neuropediatrics
24: 43–46,

[132984]1993.

[132985]

[132986]

[132987]11944.

[132988]Impagnatiello, F.; Guidotti, A. R.; Pesold, C.; Dwivedi, Y.;
Caruncho,

[132989]H.; Pisu, M. G.; Uzunov, D. P.; Smalheiser, N. R.; Davis, J.
M.; Pandey,

[132990]G. N.; Pappas, G. D.; Tueting, P.; Sharma, R. P.; Costa, E.:

A decrease

[132991]of reelin expression as a putative vulnerability factor in schizophrenia. Proc.

[132992]Nat. Acad. Sci. 95: 15718–15723, 1998.

[132993]

[132994]

[132995]11945.

[132996]Magdaleno, S.; Keshvara, L.; Curran, T.: Rescue of ataxia and

[132997]preplate splitting by ectopic expression of reelin in reeler mice. Neuron 33:

[132998]573–586, 2002.

[132999]

[133000]

[133001]11946.

[133002]Royaux, I.; Lambert de Rouvroit, C.; D'Arcangelo, G.; Demirov,

[133003]D.; Goffinet, A. M.: Genomic organization of the mouse reelin gene. Genomics 46:

[133004]240–250, 1997.

[133005]

[133006]

[133007]11947.

[133008]Yip, J. W.; Yip, Y. P. L.; Nakajima, K.; Capriotti, C.: Reelin

[133009]controls position of autonomic neurons in the spinal cord.

Proc.

[133010]Nat. Acad. Sci. 97: 8612–8616, 2000.

[133011]

[133012]

[133013]11948.

[133014]Cattaneo, M.; Lecchi, A.; Randi, A. M.; McGregor, J. L.;

Mannucci,

[133015]P. M.: Identification of a new congenital defect of platelet
function

[133016]characterized by severe impairment of platelet responses
to adenosine

[133017]diphosphate. Blood 80: 2787–2796, 1992.

[133018]

[133019]

[133020]11949.

[133021]Pawson, T.; Gish, G. D.: SH2 and SH3 domains: from
structure to

[133022]function. Cell 71: 359–362, 1992.

[133023]

[133024]

[133025]11950.

[133026]Welsh, M.; Mares, J.; Karlsson, T.; Lavergne, C.; Breant, B.;
Claesson-Welsh,

[133027]L.: Shb is a ubiquitously expressed Src homology 2 pro-
tein. Oncogene 9:

[133028]19–27, 1994.

[133029]

[133030]

[133031]11951.

[133032]Yulug, I. G.; Hillermann, R.; Fisher, E. M. C.: The SHB
adaptor

[133033]protein maps to human chromosome 9. Genomics 24:
615–617, 1994.

[133034]

[133035]

[133036]11952.

[133037]Bansal–Pakala, P.; Jember, A. G.–H.; Croft, M.: Signaling through

[133038]OX40 (CD134) breaks peripheral T–cell tolerance. Nature Med. 7:

[133039]907–912, 2001.

[133040]

[133041]

[133042]11953.

[133043]Birkeland, M. L.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[133044]A.; Barclay, A. N.: Gene structure and chromosomal localization of

[133045]the mouse homologue of rat OX40 protein. Europ. J. Immun. 25: 926–930,

[133046]1995.

[133047]

[133048]

[133049]11954.

[133050]Jember, A. G.-H.; Zuberi, R.; Liu, F.-T.; Croft, M.: Development

[133051]of allergic inflammation in a murine model of asthma is dependent

[133052]on the costimulatory receptor OX40. J. Exp. Med. 193: 387-392, 2001.

[133053]

[133054]

[133055]11955.

[133056]Latza, U.; Durkop, H.; Schnittger, S.; Ringeling, J.; Eitelbach,

[133057]F.; Hummel, M.; Fonatsch, C.; Stein, H.: The human OX40 homolog:

[133058]cDNA structure, expression and chromosomal assignment of the ACT35

[133059]antigen. Europ. J. Immun. 24: 677-683, 1994.

[133060]

[133061]

[133062]11956.

[133063]Anderson, D. W.; Probst, F. J.; Belyantseva, I. A.; Fridell, R.

[133064]A.; Beyer, L.; Martin, D. M.; Wu, D.; Kachar, B.; Friedman, T. B.;

[133065]Raphael, Y.; Camper, S. A.: The motor and tail regions of myosin

[133066]XV are critical for normal structure and function of auditory and

[133067]vestibular hair cells. Hum. Molec. Genet. 9: 1729–1738, 2000.

[133068]

[133069]

[133070]11957.

[133071]Probst, F. J.; Fridell, R. A.; Raphael, Y.; Saunders, T. L.; Wang,

[133072]A.; Liang, Y.; Morell, R. J.; Touchman, J. W.; Lyons, R. H.; Noben-Trauth,

[133073]K.; Friedman, T. B.; Camper, S. A.: Correction of deafness in shaker-2

[133074]mice by an unconventional myosin in a BAC transgene. Science 280:

[133075]1444–1447, 1998.

[133076]

[133077]

[133078]11958.

[133079]Wang, A.; Liang, Y.; Fridell, R. A.; Probst, F. J.; Wilcox, E.

[133080]R.; Touchman, J. W.; Morton, C. C.; Morell, R. J.; Noben-
Trauth, K.;

[133081]Camper, S. A.; Friedman, T. B.: Association of unconven-
tional myosin

[133082]MYO15 mutations with human nonsyndromic deafness
DFNB3. Science 280:

[133083]1447–1451, 1998.

[133084]

[133085]

[133086]11959.

[133087]Wilton, S. D.; Lim, L.; Dorosz, S. D.; Gunn, H. C.; Eyre, H.
J.;

[133088]Callen, D. F.; Laing, N. G.: Assignment of the human al-
pha-tropomyosin

[133089]gene TPM4 to band 19p13.1 by fluorescence in situ hy-
bridization. Cytogenet.

[133090]Cell Genet. 72: 294–296, 1996.

[133091]

[133092]

[133093]11960.

[133094]Lazzaro, M. A.; Picketts, D. J.: Cloning and characterization

[133095]of the murine Imitation Switch (ISWI) genes: differential expression

[133096]patterns suggest distinct developmental roles for Snf2h and Snf2l. J.

[133097]Neurochem. 77: 1145–1156, 2001.

[133098]

[133099]

[133100]11961.

[133101]Okabe, I.; Bailey, L. C.; Attree, O.; Srinivasan, S.; Perkel, J.

[133102]M.; Laurent, B. C.; Carlson, M.; Nelson, D. L.; Nussbaum, R. L.:

[133103]Cloning of human and bovine homologs of SNF2/SWI2: a global activator

[133104]of transcription in yeast *S. cerevisiae*. Nucleic Acids Res. 20:

[133105]4649–4655, 1992.

[133106]

[133107]

[133108]11962.

[133109]Stahl, W. L.; Eakin, T. J.; Owens, J. W. M., Jr.; Breininger, J.

[133110]F.; Filuk, P. E.; Anderson, W. R.: Plasma membrane

Ca(2+)-ATPase

[133111]isoforms: distribution of mRNAs in rat brain by in situ hybridization.

[133112]Molec. Brain Res. 16: 223–231, 1992.

[133113]

[133114]

[133115]11963.

[133116]Cavallo, A.: The pineal gland in human beings: relevance to pediatrics. J.

[133117]Pediat. 123: 843–851, 1993.

[133118]

[133119]

[133120]11964.

[133121]Donohue, S. J.; Roseboom, P. H.; Illnerova, H.; Weller, J. L.;

[133122]Klein, D. C.: Human hydroxyindole-O-methyltransferase:
presence of

[133123]LINE-1 fragment in a cDNA clone and pineal mRNA. DNA
Cell Biol. 12:

[133124]715-727, 1993.

[133125]

[133126]

[133127]11965.

[133128]Yi, H.; Donohue, S. J.; Klein, D. C.; McBride, O. W.: Local-
ization

[133129]of the hydroxyindole-O-methyltransferase gene to the
pseudoautosomal

[133130]region: implications for mapping of psychiatric disorders.
Hum. Molec.

[133131]Genet. 2: 127-131, 1993.

[133132]

[133133]

[133134]11966.

[133135]De Plaen, E.; Arden, K.; Traversari, C.; Gaforio, J. J.;

Szikora,

[133136]J.-P.; De Smet, C.; Brasseur, F.; van der Bruggen, P.; Lethe, B.;

[133137]Lurquin, C.; Brasseur, R.; Chomez, P.; De Backer, O.; Cavenee, W.;

[133138]Boon, T.: Structure, chromosomal localization, and expression of

[133139]12 genes of the MAGE family. Immunogenetics 40: 360–369, 1994.

[133140]

[133141]

[133142]11967.

[133143]Rogner, U. C.; Wilke, K.; Steck, E.; Korn, B.; Poustka, A.: The

[133144]melanoma antigen gene (MAGE) family is clustered in the chromosomal

[133145]band Xq28. Genomics 29: 725–731, 1995.

[133146]

[133147]

[133148]11968.

[133149]Chakarova, C.; Wehnert, M. S.; Uhl, K.; Sakthivel, S.; Vos–

berg,

[133150]H.-P.; van der Ven, P. F. M.; Furst, D. O.: Genomic structure and

[133151]fine mapping of the two human filamin gene paralogues FLNB and FLNC

[133152]and comparative analysis of the filamin gene family. Hum. Genet. 107:

[133153]597–611, 2000.

[133154]

[133155]

[133156]11969.

[133157]Fink, J. M.; Dobyns, W. B.; Guerrini, R.; Hirsch, B. A.: Identification

[133158]of a duplication of Xq28 associated with bilateral periventricular

[133159]nodular heterotopia. Am. J. Hum. Genet. 61: 379–387, 1997.

[133160]

[133161]

[133162]11970.

[133163]Fox, J. W.; Lamperti, E. D.; Eksioglu, Y. Z.; Hong, S. E.;

Feng,

[133164]Y.; Graham, D. A.; Scheffer, I. E.; Dobyns, W. B.; Hirsch, B. A.;

[133165]Radtke, R. A.; Berkovic, S. F.; Huttenlocher, P. R.; Walsh, C. A.

[133166]: Mutations in filamin 1 prevent migration of cerebral cortical neurons

[133167]in human periventricular heterotopia. Neuron 21: 1315–1325, 1998.

[133168]

[133169]

[133170]11971.

[133171]Gariboldi, M.; Maestrini, E.; Canzian, F.; Manenti, G.; De Gregorio,

[133172]L.; Rivella, S.; Chatterjee, A.; Herman, G. E.; Archidiacono, N.;

[133173]Antonacci, R.; Pierotti, M. A.; Dragani, T. A.; Toniolo, D.: Comparative

[133174]mapping of the actin-binding protein 280 genes in human and mouse. Genomics 21:

[133175]428–430, 1994.

[133176]

[133177]

[133178]11972.

[133179]Bashir, R.; Britton, S.; Strachan, T.; Keers, S.; Vafiadaki, E.;

[133180]Lako, M.; Richard, I.; Marchand, S.; Bourg, N.; Argov, Z.;
Sadeh,

[133181]M.; Mahjneh, I.; Marconi, G.; Passos-Bueno, M. R.; Moreira,
E. S.;

[133182]Zatz, M.; Beckmann, J. S.; Bushby, K.: A gene related to
Caenorhabditis

[133183]elegans spermatogenesis factor fer-1 is mutated in limb-
girdle muscular

[133184]dystrophy type 2B. Nature Genet. 20: 37-42, 1998.

[133185]

[133186]

[133187]11973.

[133188]Liu, J.; Aoki, M.; Illa, I.; Wu, C.; Fardeau, M.; Angelini, C.;

[133189]Serrano, C.; Urtizberea, J. A.; Hentati, F.; Ben Hamida, M.;
Bohlega,

[133190]S.; Culper, E. J.; Amato, A. A.; Bossie, K.; Oeltjen, J.; Be-
jaoui,

[133191]K.; McKenna-Yasek, D.; Hosler, B. A.; Schurr, E.; Arahata, K.; de

[133192]Jong, P. J.; Brown, R. H., Jr.: Dysferlin, a novel skeletal muscle

[133193]gene, is mutated in Miyoshi myopathy and limb girdle muscular dystrophy. Nature

[133194]Genet. 20: 31–36, 1998.

[133195]

[133196]

[133197]11974.

[133198]Aghai, E.; Yaniv, I.; David, M.: Factor XI deficiency in an Arab

[133199]Moslem family in Israel. Scand. J. Haemat. 32: 327–331, 1984.

[133200]

[133201]

[133202]11975.

[133203]Asakai, R.; Chung, D. W.; Davie, E. W.; Seligsohn, U.: Factor

[133204]XI deficiency in Ashkenazi Jews in Israel. New Eng. J. Med. 325:

[133205]153–158, 1991.

[133206]

[133207]

[133208]11976.

[133209]Asakai, R.; Chung, D. W.; Ratnoff, O. D.; Davie, E. W.: Factor
tor

[133210]XI (plasma thromboplastin antecedent) deficiency in
Ashkenazi Jews

[133211]is a bleeding disorder that can result from three types of
point mutations. Proc.

[133212]Nat. Acad. Sci. 86: 7667–7671, 1989.

[133213]

[133214]

[133215]11977.

[133216]Asakai, R.; Davie, E. W.; Chung, D. W.: Organization of the
gene

[133217]for human factor XI. Biochemistry 26: 7221–7228, 1987.

[133218]

[133219]

[133220]11978.

[133221]Bauduer, F.; Dupreuilh, F.; Ducout, L.; Marti, B.: Factor XI deficiency

[133222]in the French Basque country. Haemophilia 5: 187–190, 1999.

[133223]

[133224]

[133225]11979.

[133226]Bertina, R. M.; Veltkamp, J. J.: A genetic variant of factor XI

[133227]with decreased capacity for Ca ion binding. Brit. J.

Haemat. 42:

[133228]623–635, 1979.

[133229]

[133230]

[133231]11980.

[133232]Biggs, R.; MacFarlane, R. G.: Human Blood Coagulation and Its

[133233]Disorders. Oxford: Blackwell (pub.) (3rd ed.): 1962.

[133234]

[133235]

[133236]11981.

[133237]Bolton–Maggs, P. H. B.: Factor XI deficiency. Bailliere's
Clin.

[133238]Haemat. 9: 355–368, 1996.

[133239]

[133240]

[133241]11982.

[133242]Braunstein, K. M.; Noyes, C. M.; Griffith, M. J.; Lundblad, R.

[133243]L.; Roberts, H. R.: Characterization of the defect in activa-
tion

[133244]of factor IX Chapel Hill by human factor XIa. J. Clin. Invest.
68:

[133245]1420–1426, 1981.

[133246]

[133247]

[133248]11983.

[133249]Buetow, K. H.; Shiang, R.; Yang, P.; Nakamura, Y.; Lathrop,
G.

[133250]M.; White, R.; Wasmuth, J. J.; Wood, S.; Berdahl, L. D.; Ley-
sens,

[133251]N. J.: A detailed multipoint map of human chromosome 4
provides evidence

[133252]for linkage heterogeneity and position-specific recombina-
tion rates. Am.

[133253]J. Hum. Genet. 48: 911-925, 1991.

[133254]

[133255]

[133256]1 1984.

[133257]Clarkson, K.; Rosenfeld, B.; Fair, J.; Klein, A.; Bell, W.: Fac-
tor

[133258]XI deficiency acquired by liver transplantation. Ann. Intern.
Med. 115:

[133259]877-879, 1991.

[133260]

[133261]

[133262]1 1985.

[133263]Dodds, W. J.; Kull, J. E.: Canine factor XI (plasma thrombo-
plastin

[133264]antecedent) deficiency. J. Lab. Clin. Med. 78: 746–752,
1971.

[133265]

[133266]

[133267]11986.

[133268]Dzik, W. H.; Arkin, C. F.; Jenkins, R. L.: Transfer of con-
genital

[133269]factor XI deficiency from a donor to a recipient by liver
transplantation.

[133270](Letter) New Eng. J. Med. 316: 1217–1218, 1987.

[133271]

[133272]

[133273]11987.

[133274]Fujikawa, K.; Chung, D. W.; Hendrickson, L. E.; Davie, E.
W.:

[133275]Amino acid sequence of human factor XI, a blood coagu-
lation factor

[133276]with four tandem repeats that are highly homologous with
plasma prekallikrein. Biochemistry 25:

[133277]2417–2424, 1986.

[133278]

[133279]

[133280]1 1988.

[133281]Imanaka, Y.; Lal, K.; Nishimura, T.; Bolton–Maggs, P. H. B.;
Tuddenham,

[133282]E. G. D.; McVey, J. H.: Identification of two novel mutations
in

[133283]non–Jewish factor XI deficiency. Brit. J. Haemat. 90:
916–920, 1995.

[133284]

[133285]

[133286]1 1989.

[133287]Kato, A.; Asakai, R.; Davie, E. W.; Aoki, N.: Factor XI gene

[133288](F11) is located on the distal end of the long arm of hu–
man chromosome

[133289]4. Cytogenet. Cell Genet. 52: 77–78, 1989.

[133290]

[133291]

[133292]1 1990.

[133293]Litz, C. E.; Swaim, W. R.; Dalmasso, A. P.: Factor XI deficiency:

[133294]genetic and clinical studies of a single kindred. Am. J. Hemat. 28:

[133295]8–12, 1988.

[133296]

[133297]

[133298]11991.

[133299]Mannhalter, C.; Hellstern, P.; Deutsch, E.: Identification of

[133300]a defective factor XI cross-reacting material in a factor XI-deficient

[133301]patient. Blood 70: 31–37, 1987.

[133302]

[133303]

[133304]11992.

[133305]Meijers, J. C. M.; Davie, E. W.; Chung, D. W.: Expression of

[133306]human blood coagulation factor XI: characterization of the defect

[133307]in factor XI type III deficiency. Blood 79: 1435–1440,

1992.

[133308]

[133309]

[133310]1 1993.

[133311]Mitchell, M.; Cutler, J.; Thompson, S.; Moore, G.; Rees, E.
J.;

[133312]Smith, M.; Savidge, G.; Alhaq, A.: Heterozygous factor XI
deficiency

[133313]associated with three novel mutations. Brit. J. Haemat.
107: 763–765,

[133314]1999.

[133315]

[133316]

[133317]1 1994.

[133318]Muir, W. A.; Ratnoff, O. D.: The prevalence of plasma
thromboplastin

[133319]antecedent (PTA factor XI) deficiency. Blood 44: 569–570,
1974.

[133320]

[133321]

[133322]11995.

[133323]Niskanen, E. O.; Saito, H.; Cline, M. J.: Plasma thrombo-
plastin

[133324]antecedent (factor XI) deficiency in a black family. Arch.
Intern.

[133325]Med. 141: 936–941, 1981.

[133326]

[133327]

[133328]11996.

[133329]Peretz, U.; Zivelin, A.; Usher, S.; Eichel, R.; Seligsohn, U.

[133330]: Identification of a new mutation in the factor XI gene of
an Ashkenazi–Jew

[133331]with severe factor XI deficiency. (Abstract) Blood 82
(suppl. 1):

[133332]66a only, 1993.

[133333]

[133334]

[133335]11997.

[133336]Pugh, R. E.; McVey, J. H.; Tuddenham, E. G. D.; Hancock, J.

F.

[133337]: Six point mutations that cause factor XI deficiency. Blood
85:

[133338]1509–1516, 1995.

[133339]

[133340]

[133341]1 1998.

[133342]Ragni, M. V.; Sinha, D.; Seaman, F.; Lewis, J. H.; Spero, J.
A.;

[133343]Walsh, P. N.: Comparison of bleeding tendency, factor XI
coagulant

[133344]activity, and factor XI antigen in 25 factor XI-deficient kin-
dreds. Blood 65:

[133345]719–724, 1985.

[133346]

[133347]

[133348]1 1999.

[133349]Rapaport, S. I.; Proctor, R. R.; Patch, M. J.; Yettra, M.: The

[133350]mode of inheritance of PTA deficiency: evidence for the
existence

[133351]of major PTA deficiency and minor PTA deficiency. Blood
18: 149–165,
[133352]1961.

[133353]

[133354]

[133355]12000.

[133356]Rimon, A.; Schiffman, S.; Feinstein, D. I.; Rapaport, S. I.:

[133357]Factor–XI activity and factor–XI antigen in homozygous
and heterozygous

[133358]factor–XI deficiency. Blood 48: 165–174, 1976.

[133359]

[133360]

[133361]12001.

[133362]Foresti, V.; Ferrari, C.: Central hypothyroidism: isolated
thyrotropin–releasing

[133363]hormone deficiency or resistance of pituitary thyrotropes?.
(Letter) J.

[133364]Endocr. Invest. 8: 577 only, 1985.

[133365]

[133366]

[133367]12002.

[133368]Katakami, H.; Kato, Y.; Inada, M.; Imura, H.: Hypothalamic
hypothyroidism

[133369]due to isolated thyrotropin-releasing hormone (TRH) deficiency. J.

[133370]Endocr. Invest. 7: 231-233, 1984.

[133371]

[133372]

[133373]12003.

[133374]Niimi, H.; Inomata, H.; Sasaki, N.; Nakajima, H.: Congenital
isolated

[133375]thyrotrophin releasing hormone deficiency. Arch. Dis.
Child. 57:

[133376]877-878, 1982.

[133377]

[133378]

[133379]12004.

[133380]Roller, M. L.; Camper, S. A.: Localization of the thy-

rotropin-releasing

[133381]hormone gene, Trh, on mouse chromosome 6. Mam-
malian Genome 6: 443-444,

[133382]1995.

[133383]

[133384]

[133385]12005.

[133386]Yamada, M.; Radovick, S.; Wondisford, F. E.; Nakayama, Y.;
Weintraub,

[133387]B. D.; Wilber, J. F.: Cloning and structure of human ge-
nomic DNA

[133388]and hypothalamic cDNA encoding human preprothy-
rotropin-releasing hormone. Molec.

[133389]Endocr. 4: 551-556, 1990.

[133390]

[133391]

[133392]12006.

[133393]Yamada, M.; Satoh, T.; Monden, T.; Mori, M.: Assignment
of the

[133394]thyrotropin-releasing hormone gene (TRH) to human

chromosome 3q13.3–q21

[133395]by in situ hybridization. Cytogenet. Cell Genet. 87: 275
only, 1999.

[133396]

[133397]

[133398]12007.

[133399]Yamada, M.; Wondisford, F. E.; Radovick, S.; Nakayama, Y.;
Weintraub,

[133400]B. D.; Wilber, J. F.: Assignment of human preprothy-
rotropin–releasing

[133401]hormone (TRH) gene to chromosome 3. Somat. Cell Molec.
Genet. 17:

[133402]97–100, 1991.

[133403]

[133404]

[133405]12008.

[133406]McDermott, J. C.; Cardoso, M. C.; Yu, Y.–T.; Andres, V.;
Leifer,

[133407]D.; Krainc, D.; Lipton, S. A.; Nadal–Ginard, B.: hMEF2C
gene encodes

[133408]skeletal muscle– and brain–specific transcription factors.

Molec.

[133409]Cell. Biol. 13: 2564–2577, 1993.

[133410]

[133411]

[133412]12009.

[133413]Anderson, G. J.; Murphy, T. L.; Cowley, L.; Evans, B. A.;
Halliday,

[133414]J. W.; McLaren, G. D.: Mapping the gene for sex-linked
anemia: an

[133415]inherited defect of intestinal iron absorption in the mouse.

Genomics 48:

[133416]34–39, 1998.

[133417]

[133418]

[133419]12010.

[133420]Bannerman, R. M.: Genetic defects of iron transport. Fed.
Proc. 35:

[133421]2281–2285, 1976.

[133422]

[133423]

[133424]12011.

[133425]Edwards, J. A.; Hoke, J. E.; Mattioli, M.; Reichlin, M.: Fer-
ritin

[133426]distribution and synthesis in sex-linked anemia. J. Lab.
Clin. Med. 90:

[133427]68-76, 1977.

[133428]

[133429]

[133430]12012.

[133431]Falconer, D. S.; Isaacson, J. H.: The genetics of sex-linked
anaemia

[133432]in the mouse. Genet. Res. 3: 248-250, 1962.

[133433]

[133434]

[133435]12013.

[133436]Grewal, M. S.: A sex-linked anaemia in the mouse. Genet.
Res. 3:

[133437]238-247, 1962.

[133438]

[133439]

[133440]12014.

[133441]Lee, G. R.; Nacht, S.; Lukens, J. N.; Cartwright, G. E.: Iron

[133442]metabolism in copper-deficient swine. J. Clin. Invest. 47:
2058–2069,

[133443]1968.

[133444]

[133445]

[133446]12015.

[133447]Vulpe, C. D.; Kuo, Y.-M.; Murphy, T. L.; Cowley, L.;
Askwith, C.;

[133448]Libina, N.; Gitschier, J.; Anderson, G. J.: Hephaestin, a
ceruloplasmin

[133449]homologue implicated in intestinal iron transport, is de-
fective in

[133450]the sla mouse. Nature Genet. 21: 195–199, 1999.

[133451]

[133452]

[133453]12016.

[133454]Karumanchi, S. A.; Jha, V.; Ramchandran, R.; Karihaloo, A.;

Tsiokas,

[133455]L.; Chan, B.; Dhanabal, M.; Hanai, J.; Venkataraman, G.;

Shriver,

[133456]Z.; Keiser, N.; Kalluri, R.; and 9 others: Cell surface glypi-

cans

[133457]are low-affinity endostatin receptors. *Molec. Cell* 7:

811–822, 2001.

[133458]

[133459]

[133460]12017.

[133461]Veugelers, M.; Vermeesch, J.; Watanabe, K.; Yamaguchi,

Y.; Marynen,

[133462]P.; David, G.: GPC4, the gene for human K-glypican, flanks

GPC3 on

[133463]Xq26: deletion of the GPC3–GPC4 gene cluster in one

family with Simpson–Golabi–Behmel

[133464]syndrome. *Genomics* 53: 1–11, 1998.

[133465]

[133466]

[133467]12018.

[133468]Emes, R. D.; Ponting, C. P.: A new sequence motif linking
lissencephaly,

[133469]Treacher Collins and oral-facial-digital type 1 syndromes,
microtubule

[133470]dynamics and cell migration. Hum. Molec. Genet. 10:
2813-2820, 2001.

[133471]

[133472]

[133473]12019.

[133474]Copley, L. M.; Zhao, W. D.; Kopacz, K.; Herman, G. E.;
Kioschis,

[133475]P.; Poustka, A.; Taudien, S.; Platzer, M.: Exclusion of mu-
tations

[133476]in the MTMR1 gene as a frequent cause of X-linked my-
otubular myopathy.

[133477](Letter) Am. J. Med. Genet. 107: 256-258, 2002.

[133478]

[133479]

[133480]12020.

[133481]Kioschis, P.; Wiemann, S.; Heiss, N. S.; Francis, F.; Gotz, C.;

[133482]Poustka, A.; Taudien, S.; Platzer, M.; Wiehe, T.; Beckmann, G.; Weber,

[133483]J.; Nordsiek, G.; Rosenthal, A.: Genomic organization of a 225-kb

[133484]region in Xq28 containing the gene for X-linked myotubular myopathy

[133485](MTM1) and a related gene (MTMR1). Genomics 54: 256–266, 1998.

[133486]

[133487]

[133488]12021.

[133489]Laporte, J.; Blondeau, F.; Buj-Bello, A.; Tentler, D.; Kretz, C.;

[133490]Dahl, N.; Mandel, J.–L.: Characterization of the myotubularin dual

[133491]specificity phosphatase gene family from yeast to human. Hum. Molec.

[133492]Genet. 7: 1703–1712, 1998.

[133493]

[133494]

[133495]12022.

[133496]Gorlin, J. B.; Henske, E.; Warren, S. T.; Kunst, C. B.; D'Urso,

[133497]M.; Palmieri, G.; Hartwig, J. H.; Bruns, G.; Kwiatkowski, D.

J.:

[133498]Actin-binding protein (ABP-280) filamin gene (FLN) maps
telomeric

[133499]to the color vision locus (R/GCP) and centromeric to G6PD
in Xq28. Genomics 17:

[133500]496-498, 1993.

[133501]

[133502]

[133503]12023.

[133504]Jager, R. J.; Anvret, M.; Hall, K.; Scherer, G.: A human XY
female

[133505]with a frame shift mutation in the candidate testis-
determining gene

[133506]SRY. Nature 348: 452-454, 1990.

[133507]

[133508]

[133509]12024.

[133510]Koopman, P.; Gubbay, J.; Vivian, N.; Goodfellow, P.; Lovell–
Badge,

[133511]R.: Male development of chromosomally female mice
transgenic for

[133512]Sry. Nature 351: 117–121, 1991.

[133513]

[133514]

[133515]12025.

[133516]Koopman, P.; Munsterberg, A.; Capel, B.; Vivian, N.;
Lovell–Badge,

[133517]R.: Expression of a candidate sex–determining gene dur–
ing mouse testis

[133518]differentiation. Nature 348: 450–452, 1990.

[133519]

[133520]

[133521]12026.

[133522]McElreavey, K. D.; Vilain, E.; Boucekkine, C.; Vidaud, M.;

Jaubert,

[133523]F.; Richaud, F.; Fellous, M.: XY sex reversal associated with
a nonsense

[133524]mutation in SRY. Genomics 13: 838–840, 1992.

[133525]

[133526]

[133527]12027.

[133528]Page, D. C.; Mosher, R.; Simpson, E. M.; Fisher, E. M. C.;
Mardon,

[133529]G.; Pollack, J.; McGillivray, B.; de la Chapelle, A.; Brown, L.
G.

[133530]: The sex-determining region of the human Y chromo-
some encodes a finger

[133531]protein. Cell 51: 1091–1104, 1987.

[133532]

[133533]

[133534]12028.

[133535]Taylor, H.; Barter, R. H.; Jacobson, C. B.: Neoplasms of
dysgenetic

[133536]gonads. Am. J. Obstet. Gynec. 96: 816–823, 1966.

[133537]

[133538]

[133539]12029.

[133540]Vilain, E.; McElreavey, K.; Jaubert, F.; Raymond, J.-P.;
Richaud,

[133541]F.; Fellous, M.: Familial case with sequence variant in the
testis-determining

[133542]region associated with two sex phenotypes. Am. J. Hum.
Genet. 50:

[133543]1008-1011, 1992.

[133544]

[133545]

[133546]12030.

[133547]Gorlin, J. B.; Yamin, R.; Egan, S.; Stewart, M.; Stossel, T. P.;

[133548]Kwiatkowski, D. J.; Hartwig, J. H.: Human endothelial
actin-binding

[133549]protein (ABP-280, nonmuscle filamin): a molecular leaf
spring. J.

[133550]Cell Biol. 111: 1089-1105, 1990.

[133551]

[133552]

[133553]12031.

[133554]Huttenlocher, P. R.; Taravath, S.; Mojtahedi, S.: Periven-
tricular

[133555]heterotopia and epilepsy. Neurology 44: 51–55, 1994.

[133556]

[133557]

[133558]12032.

[133559]Kunst, C. B.; Henske, E.; Hartwig, J. H.; Kwiatkowski, D. J.;
D'Urso,

[133560]M.; Bruns, G.; Warren, S. T.; Gorlin, J. B.: The dystrophin-
like

[133561]actin binding protein 280 gene maps between DXS52 and
G6PD overlapping

[133562]the Emery–Dreifuss muscular dystrophy locus. (Abstract)
Am. J. Hum.

[133563]Genet. 51: A21, 1992.

[133564]

[133565]

[133566]12033.

[133567]Maestrini, E.; Patrosso, C.; Mancini, M.; Rivella, S.; Rocchi,
[133568]M.; Repetto, M.; Villa, A.; Frattini, A.; Zoppe, M.; Vezzoni,
P.;

[133569]Toniolo, D.: Mapping of two genes encoding isoforms of
the actin

[133570]binding protein ABP-280, a dystrophin like protein, to
Xq28 and to

[133571]chromosome 7. Hum. Molec. Genet. 2: 761-766, 1993.

[133572]

[133573]

[133574]12034.

[133575]Maestrini, E.; Rivella, S.; Tribioli, C.; Purtilo, D.; Rocchi,

[133576]M.; Archidiacono, N.; Toniolo, D.: Probes for CpG islands
on the

[133577]distal long arm of the human X chromosome are clustered
in Xq24 and

[133578]Xq28. Genomics 8: 664-670, 1990.

[133579]

[133580]

[133581]12035.

[133582]Patrosso, M. C.; Repetto, M.; Villa, A.; Milanesi, L.; Frattini,
[133583]A.; Faranda, S.; Mancini, M.; Maestrini, E.; Toniolo, D.;
Vezzoni,

[133584]P.: The exon–intron organization of the human X–linked
gene (FLN1)

[133585]encoding actin–binding protein 280. Genomics 21: 71–76,
1994.

[133586]

[133587]

[133588]12036.

[133589]Sheen, V. L.; Dixon, P. H.; Fox, J. W.; Hong, S. E.; Kinton,
L.;

[133590]Sisodiya, S. M.; Duncan, J. S.; Dubeau, F.; Scheffer, I. E.;
Schachter,

[133591]S. C.; Wilner, A.; Henchy, R.; and 18 others: Mutations in
the

[133592]X–linked filamin 1 gene cause periventricular nodular het-
erotopia

[133593]in males as well as in females. Hum. Molec. Genet. 10:
1775–1783,

[133594]2001.

[133595]

[133596]

[133597]12037.

[133598]Small, K.; Wagener, M.; Warren, S. T.: Isolation and characterization

[133599]of the complete mouse emerin gene. Mammalian Genome
8: 337-341,

[133600]1997.

[133601]

[133602]

[133603]12038.

[133604]Vadlamudi, R. K.; Li, F.; Adam, L.; Nguyen, D.; Ohta, Y.;
Stossel,

[133605]T. P.; Kumar, R.: Filamin is essential in actin cytoskeletal
assembly

[133606]mediated by p21-activated kinase 1. Nature Cell Biol. 4:
681-690,

[133607]2002.

[133608]

[133609]

[133610]12039.

[133611]Faranda, S.; Frattini, A.; Vezzoni, P.: The human genes encoding

[133612]renin-binding protein and host cell factor are closely linked in Xq28

[133613]and transcribed in the same direction. Gene 155: 237-239, 1995.

[133614]

[133615]

[133616]12040.

[133617]Frattini, A.; Chatterjee, A.; Faranda, S.; Sacco, M. G.; Villa,

[133618]A.; Herman, G. E.; Vezzoni, P.: The chromosome localization and the

[133619]HCF repeats of the human host cell factor gene (HCFC1) are conserved

[133620]in the mouse homologue. Genomics 32: 277-280, 1996.

[133621]

[133622]

[133623]12041.

[133624]Frattini, A.; Faranda, S.; Redolfi, E.; Zucchi, I.; Villa, A.;

[133625]Patrosso, M. C.; Strina, D.; Susani, L.; Vezzoni, P.: Genomic organization

[133626]of the human VP16 accessory protein, a housekeeping gene (HCFC1) mapping

[133627]to Xq28. Genomics 23: 30–35, 1994.

[133628]

[133629]

[133630]12042.

[133631]Wilson, A. C.; LaMarco, K.; Peterson, M. G.; Herr, W.: The VP16

[133632]accessory protein HCF is a family of polypeptides processed from a

[133633]large precursor protein. Cell 74: 115–125, 1993.

[133634]

[133635]

[133636]12043.

[133637]Wilson, A. C.; Parrish, J. E.; Massa, H. F.; Nelson, D. L.; Trask,

[133638]B. J.; Herr, W.: The gene encoding the VP16–accessory protein HCF

[133639](HCFC1) resides in human Xq28 and is highly expressed in fetal tissues

[133640]and the adult kidney. Genomics 25: 462–468, 1995.

[133641]

[133642]

[133643]12044.

[133644]Zoppe, M.; Frattini, A.; Faranda, S.; Vezzoni, P.: The complete

[133645]sequence of the host cell factor 1 (HCFC1) gene and its promoter:

[133646]a role for YY1 transcription factor in the regulation of its expression. Genomics 34:

[133647]85–91, 1996.

[133648]

[133649]

[133650]12045.

[133651]Maestrini, E.; Tamagnone, L.; Longati, P.; Cremona, O.;
Gulisano,

[133652]M.; Bione, S.; Tamanini, F.; Neel, B. G.; Toniolo, D.; Co-
moglio, P.

[133653]M.: A family of transmembrane proteins with homology to the MET–hepatocyte

[133654]growth factor receptor. Proc. Nat. Acad. Sci. 93: 674–678, 1996.

[133655]

[133656]

[133657]12046.

[133658]Tamagnone, L.; Artigiani, S.; Chen, H.; He, Z.; Ming, G.; Song,

[133659]H.; Chedotal, A.; Winberg, M. L.; Goodman, C. S.; Poo, M.; Tessier–Lavigne,

[133660]M.; Comoglio, P. M.: Plexins are a large family of receptors for

[133661]transmembrane, secreted, and GPI–anchored semaphorins in vertebrates. Cell 99:

[133662]71–80, 1999.

[133663]

[133664]

[133665]12047.

[133666]Becker, P. E.: Two new families of benign sex–linked recessive

[133667]muscular dystrophy. Rev. Canad. Biol. 21: 551–566, 1962.

[133668]

[133669]

[133670]12048.

[133671]Becker, P. E.: Eine neue X-chromosomale Muskeldystrophie. Acta

[133672]Psychiat. Neurol. Scand. 193: 427, 1955.

[133673]

[133674]

[133675]12049.

[133676]Becker, P. E.: Neue Ergebnisse der Genetik der Muskeldystrophien. Acta

[133677]Genet. Statist. Med. 7: 303–310, 1957.

[133678]

[133679]

[133680]12050.

[133681]Bushby, K. M. D.; Cleghorn, N. J.; Curtis, A.; Haggerty, I. D.;

[133682]Nicholson, L. V. B.; Johnson, M. A.; Harris, J. B.; Bhat-

tacharya,

[133683]S. S.: Identification of a mutation in the promoter region
of the

[133684]dystrophin gene in a patient with atypical Becker muscular
dystrophy. Hum.

[133685]Genet. 88: 195–199, 1991.

[133686]

[133687]

[133688]12051.

[133689]Doriguzzi, C.; Palmucci, L.; Mongini, T.; Chiado-Piat, L.;
Restagno,

[133690]G.; Ferrone, M.: Exercise intolerance and recurrent myo-
globinuria

[133691]as the only expression of Xp21 Becker type muscular dys-
trophy. J.

[133692]Neurol. 240: 269–271, 1993.

[133693]

[133694]

[133695]12052.

[133696]England, S. B.; Nicholson, L. V. B.; Johnson, M. A.; Forrest,

[133697]S. M.; Love, D. R.; Zubrzycka-Gaarn, E. E.; Bulman, D. E.;
Harris,

[133698]J. B.; Davies, K. E.: Very mild muscular dystrophy associ-
ated with

[133699]the deletion of 46% dystrophin. Nature 343: 180–182,
1990.

[133700]

[133701]

[133702]12053.

[133703]Aronsson, F. C.; Magnusson, P.; Andersson, B.; Karsten, S.
L.;

[133704]Shibasaki, Y.; Lendon, C. L.; Goate, A. M.; Brookes, A. J:
The NIK

[133705]protein kinase and C17orf1 genes: chromosomal map-
ping, gene structures

[133706]and mutational screening in frontotemporal dementia and
parkinsonism

[133707]linked to chromosome 17. Hum. Genet. 103: 340–345,
1998.

[133708]

[133709]

[133710]12054.

[133711]Malinin, N. L.; Boldin, M. P.; Kovalenko, A. V.; Wallach, D.:

[133712]MAP3K-related kinase involved in NF-kappaB induction by
TNF, CD95

[133713]and IL-1. Nature 385: 540-544, 1997.

[133714]

[133715]

[133716]12055.

[133717]Shinkura, R.; Kitada, K.; Matsuda, F.; Tashiro, K.; Ikuta, K.;

[133718]Suzuki, M.; Kogishi, K.; Serikawa, T.; Honjo, T.: A lympho-
plasia is

[133719]caused by a point mutation in the mouse gene encoding
Nf-kappa-b-inducing

[133720]kinase. Nature Genet. 22: 74-77, 1999.

[133721]

[133722]

[133723]12056.

[133724]Smith, C.; Andreakos, E.; Crawley, J. B.; Brennan, F. M.;
Feldmann,

[133725]M.; Foxwell, B. M. J.: NF-kappa-B-inducing kinase is dispensable

[133726]for activation of NF-kappa-B in inflammatory settings but essential

[133727]for lymphotoxin beta receptor activation of NF-kappa-B in primary

[133728]human fibroblasts. J. Immun. 167: 5895-5903, 2001.

[133729]

[133730]

[133731]12057.

[133732]Yin, L.; Wu, L.; Wesche, H.; Arthur, C. D.; White, J. M.; Goeddel,

[133733]D. V.; Schreiber, R. D.: Defective lymphotoxin-beta receptor-induced

[133734]NF-kappa-B transcriptional activity in NIK-deficient mice. Science 291:

[133735]2162-2165, 2001.

[133736]

[133737]

[133738]12058.

[133739]Aronsson, F. C.; Magnusson, P.; Andersson, B.; Karsten, S.
L.;

[133740]Shibasaki, Y.; Lendon, C. L.; Goate, A. M.; Brookes, A. J.:
The NIK

[133741]protein kinase and C17orf1 genes: chromosomal map-
ping, gene structures

[133742]and mutational screening in frontotemporal dementia and
parkinsonism

[133743]linked to chromosome 17. Hum. Genet. 103: 340–345,
1998.

[133744]

[133745]

[133746]12059.

[133747]Yayoshi–Yamamoto, S.; Taniuchi, I.; Watanabe, T.: FRL, a
novel

[133748]formin–related protein, binds to Rac and regulates cell
motility and

[133749]survival of macrophages. Molec. Cell. Biol. 20:
6872–6881, 2000.

[133750]

[133751]

[133752]12060.

[133753]Gress, T. M.; Muller-Pillasch, F.; Geng, M.; Zimmerhackl, F.; Zehetner,

[133754]G.; Friess, H.; Buchler, M.; Adler, G.; Lehrach, H.: A pancreatic
creatic

[133755]cancer-specific expression profile. Oncogene 13:
1819-1830, 1996.

[133756]

[133757]

[133758]12061.

[133759]Muller-Pillasch, F.; Wallrapp, C.; Lacher, U.; Friess, H.;
Buchler,

[133760]M.; Adler, G.; Gress, T. M.: Identification of a new tumour-
associated

[133761]antigen TM4SF5 and its expression in human cancer. Gene
208: 25-30,

[133762]1998.

[133763]

[133764]

[133765]12062.

[133766]Borggreffe, T.; Masat, L.; Wabl, M.; Riwar, B.; Cattoretti, G.;

[133767]Jessberger, R.: Cellular, intracellular, and developmental
expression

[133768]patterns of murine SWAP-70. Europ. J. Immun. 29:
1812-1822, 1999.

[133769]

[133770]

[133771]12063.

[133772]Borggreffe, T.; Wabl, M.; Akhmedov, A. T.; Jessberger, R.: A
B-cell-specific

[133773]DNA recombination complex. J. Biol. Chem. 273:
17025-17035, 1998.

[133774]

[133775]

[133776]12064.

[133777]Masat, L.; Caldwell, J.; Armstrong, R.; Khoshnevisan, H.;
Jessberger,

[133778]R.; Herndier, B.; Wabl, M.; Ferrick, D.: Association of
SWAP-70 with

[133779]the B cell antigen receptor complex. Proc. Nat. Acad. Sci.
97: 2180-2184,

[133780]2000.

[133781]

[133782]

[133783]12065.

[133784]Masat, L.; Liddell, R. A.; Mock, B. A.; Kuo, W.-L.; Jess-
berger,

[133785]R.; Wabl, M.; Morse, H. C., III: Mapping of the SWAP70
gene to mouse

[133786]chromosome 7 and human chromosome 11p15. Immuno-
genetics 51: 16-19,

[133787]2000.

[133788]

[133789]

[133790]12066.

[133791]Shinohara, M.; Terada, Y.; Iwamatsu, A.; Shinohara, A.;
Mochizuki,

[133792]N.; Higuchi, M.; Gotoh, Y.; Ihara, S.; Nagata, S.; Itoh, H.;
Fukui,

[133793]Y.; Jessberger, R.: SWAP-70 is a guanine-nu-
cleotide-exchange factor

[133794]that mediates signalling of membrane ruffling. Nature
416: 759–763,
[133795]2002.

[133796]

[133797]

[133798]12067.

[133799]Miura, K.; Suzuki, K.; Tokino, T.; Isomura, M.; Inazawa, J.;
Matsuno,

[133800]S.; Nakamura, Y.: Detailed deletion mapping in squamous
cell carcinomas

[133801]of the esophagus narrows a region containing a putative
tumor suppressor

[133802]gene to about 200 kilobases on distal chromosome 9q.
Cancer Res. 56:

[133803]1629–1634, 1996.

[133804]

[133805]

[133806]12068.

[133807]Nishiwaki, T.; Daigo, Y.; Kawasoe, T.; Nakamura, Y.: Isola-
tion

[133808]and mutational analysis of a novel human cDNA, DEC1
(deleted in esophageal

[133809]cancer 1), derived from the tumor suppressor locus in
9q32. Genes

[133810]Chromosomes Cancer 27: 169–176, 2000.

[133811]

[133812]

[133813]12069.

[133814]Kamimoto, T.; Zama, T.; Aoki, R.; Muro, Y.; Hagiwara, M.:
Identification

[133815]of a novel kinesin-related protein, KRMP1, as a target for
mitotic

[133816]peptidyl-prolyl isomerase Pin1. J. Biol. Chem. 276:
37520–37528,

[133817]2001.

[133818]

[133819]

[133820]12070.

[133821]Akagi, T.; Motegi, M.; Tamura, A.; Suzuki, R.; Hosokawa,
Y.; Suzuki,

[133822]H.; Ota, H.; Nakamura, S.; Morishima, Y.; Taniwaki, M.;
Seto, M.:

[133823]A novel gene, MALT1 at 18q21, is involved in t(11;18)
(q21;q21) found

[133824]in low-grade B-cell lymphoma of mucosa-associated lym-
phoid tissue. Oncogene 18:

[133825]5785–5794, 1999.

[133826]

[133827]

[133828]12071.

[133829]Dierlamm, J.; Baens, M.; Wlodarska, I.; Stefanova-
Ouzounova, M.;

[133830]Hernandez, J. M.; Hossfeld, D. K.; De Wolf-Peeters, C.;
Hagemeijer,

[133831]A.; Van den Berghe, H.; Marynen, P.: The apoptosis in-
hibitor gene

[133832]API2 and a novel 18q gene, MLT, are recurrently rear-
ranged in the

[133833]t(11;18)(q21;q21) associated with mucosa-associated
lymphoid tissue

[133834]lymphomas. Blood 93: 3601–3609, 1999.

[133835]

[133836]

[133837]12072.

[133838]Ott, G.; Katzenberger, T.; Greiner, A.; Kalla, J.; Rosenwald, A.;

[133839]Heinrich, U.; Ott, M. M.; Muller-Hermelink, H. K.: The t(11;18)(q21;q21)

[133840]chromosome translocation is a frequent and specific aberration in

[133841]low-grade but not high-grade malignant non-Hodgkin's lymphomas of

[133842]the mucosa-associated lymphoid tissue (MALT-) type. Cancer Res. 57:

[133843]3944-3948, 1997.

[133844]

[133845]

[133846]12073.

[133847]Sato, Y.; Akiyama, Y.; Tanizawa, T.; Shibata, T.; Saito, K.; Mori,

[133848]S.; Kamiyama, R.; Yuasa, Y.: Molecular characterization of the genomic

[133849]breakpoint junction in the t(11;18)(q21;q21) translocation
of a gastric

[133850]MALT lymphoma. Biochem. Biophys. Res. Commun. 280:
301–306, 2001.

[133851]

[133852]

[133853]12074.

[133854]Valladeau, J.; Clair–Moninot, V.; Dezutter–Dambuyant, C.;
Pin,

[133855]J.–J.; Kissenpfennig, A.; Mattei, M.–G.; Ait–Yahia, S.; Bates,
E.

[133856]E. M.; Malissen, B.; Koch, F.; Fossiez, F.; Romani, N.;
Lebecque,

[133857]S.; Saeland, S.: Identification of mouse langerin/CD207 in
Langerhans

[133858]cells and some dendritic cells of lymphoid tissues. J. Im–
mun. 168:

[133859]782–792, 2002.

[133860]

[133861]

[133862]12075.

- [133863]Valladeau, J.; Ravel, O.; Dezutter–Dambuyant, C.; Moore, K.; Kleijmeer, M.; Liu, Y.; Duvert–Frances, V.; Vincent, C.; Schmitt, D.; Davoust, J.; Caux, C.; Lebecque, S.; Saeland, S.: Langerin, a novel C-type lectin specific to Langerhans cells, is an endocytic receptor that induces the formation of Birbeck granules. *Immunity* 12: 71–81, 2000.
- [133868]
- [133869]
- [133870]12076.
- [133871]Reichenberger, E.; Tiziani, V.; Watanabe, S.; Park, L.; Ueki, Y.;
- [133872]Santanna, C.; Baur, S. T.; Shiang, R.; Grange, D. K.; Beighton, P.;
- [133873]Gardner, J.; Hamersma, H.; Sellars, S.; Ramesar, R.; Lidral, A. C.;
- [133874]Sommer, A.; Raposo do Amaral, C. M.; Gorlin, R. J.; Mulliken, J. B.;
- [133875]Olsen, B. R.: Autosomal dominant craniometaphyseal dys–

plasia is caused

[133876]by mutations in the transmembrane protein ANK. Am. J.

Hum. Genet. 68:

[133877]1321–1326, 2001.

[133878]

[133879]

[133880]12077.

[133881]Ansel, K. M.; Harris, R. B. S.; Cyster, J. G.: CXCL13 is re-
quired

[133882]for B1 cell homing, natural antibody production, and body
cavity immunity. Immunity 16:

[133883]67–76, 2002.

[133884]

[133885]

[133886]12078.

[133887]Gunn, M. D.; Ngo, V. N.; Ansel, K. M.; Ekland, E. H.; Cyster,
J.

[133888]G.; Williams, L. T.: A B–cell–homing chemokine made in
lymphoid follicles

[133889]activates Burkitt's lymphoma receptor–1. Nature 391:

799–803, 1998.

[133890]

[133891]

[133892]12079.

[133893]Legler, D. F.; Loetscher, M.; Roos, R. S.; Clark–Lewis, I.;
Baggiolini,

[133894]M.; Moser, B.: B cell–attracting chemokine 1, a human CXC
chemokine

[133895]expressed in lymphoid tissues, selectively attracts B lym-
phocytes

[133896]via BLR1/CXCR5. J. Exp. Med. 187: 655–660, 1998.

[133897]

[133898]

[133899]12080.

[133900]Derst, C.; Engel, H.; Grzeschik, K.–H.; Daut, J.: Genomic
structure

[133901]and chromosome mapping of human and mouse RAMP
genes. Cytogenet.

[133902]Cell Genet. 90: 115–118, 2000.

[133903]

[133904]

[133905]12081.

[133906]Amler, L. C.; Bauer, A.; Corvi, R.; Dihlmann, S.; Praml, C.;
Cavenee,

[133907]W. K.; Schwab, M.; Hampton, G. M.: Identification and
characterization

[133908]of novel genes located at the t(1;15)(p36.2;q24) translo-
cation breakpoint

[133909]in the neuroblastoma cell line NGP. Genomics 64:
195–202, 2000.

[133910]

[133911]

[133912]12082.

[133913]Yanagisawa, H.; Bundo, M.; Miyashita, T.; Okamura–Oho,
Y.; Tadokoro,

[133914]K.; Tokunaga, K.; Yamada, M.: Protein binding of a DRPLA
family through

[133915]arginine–glutamic acid dipeptide repeats is enhanced by
extended polyglutamine. Hum.

[133916]Molec. Genet. 9: 1433–1442, 2000.

[133917]

[133918]

[133919]12083.

[133920]Oh, J.; Takahashi, R.; Kondo, S.; Mizoguchi, A.; Adachi, E.;
Sasahara,

[133921]R. M.; Nishimura, S.; Imamura, Y.; Kitayama, H.; Alexan-
der, D. B.;

[133922]Ide, C.; Horan, T. P.; Arakawa, T.; Yoshida, H.; Nishikawa,
S.; Itoh,

[133923]Y.; Seiki, M.; Itohara, S.; Takahashi, C.; Noda, M.: The
membrane-anchored

[133924]MMP inhibitor RECK is a key regulator of extracellular ma-
trix integrity

[133925]and angiogenesis. Cell 107: 789-800, 2001.

[133926]

[133927]

[133928]12084.

[133929]Takahashi, C.; Sheng, Z.; Horan, T. P.; Kitayama, H.; Maki,
M.;

[133930]Hitomi, K.; Kitaura, Y.; Takai, S.; Sasahara, R. M.; Hori-
moto, A.;

[133931]Ikawa, Y.; Ratzkin, B. J.; Arakawa, T.; Noda, M.: Regulation

of matrix

[133932]metalloproteinase-9 and inhibition of tumor invasion by
the membrane-anchored

[133933]glycoprotein RECK. Proc. Nat. Acad. Sci. 95:
13221-13226, 1998.

[133934]

[133935]

[133936]12085.

[133937]Nagase, T.; Ishikawa, I.; Nakajima, D.; Ohira, M.; Seki, N.;
Miyajima,

[133938]N.; Tanaka, A.; Kotani, H.; Nomura, N.; O'Hara, O.: Predic-
tion of

[133939]the coding sequences of unidentified human genes. VII.
The complete

[133940]sequences of 100 new cDNA clones from brain which can
code for large

[133941]proteins in vitro. DNA Res. 4: 141-150, 1997.

[133942]

[133943]

[133944]12086.

[133945]Del Punta, K.; Leinders-Zufall, T.; Rodriguez, I.; Jukam, D.;

Wysocki,

[133946]C. J.; Ogawa, S.; Zufall, F.; Mombaerts, P.: Deficient
pheromone

[133947]responses in mice lacking a cluster of vomeronasal recep-
tor genes. Nature 419:

[133948]70–74, 2002.

[133949]

[133950]

[133951]12087.

[133952]Pantages, E.; Dulac, C.: A novel family of candidate
pheromone

[133953]receptors in mammals. Neuron 28: 835–845, 2000.

[133954]

[133955]

[133956]12088.

[133957]Rodriguez, I.; Feinstein, P.; Mombaerts, P.: Variable pat-
terns

[133958]of axonal projections of sensory neurons in the mouse
vomeronasal

[133959]system. Cell 97: 199–208, 1999.

[133960]

[133961]

[133962]12089.

[133963]Rodriguez, I.; Greer, C. A.; Mok, M. Y.; Mombaerts, P.: A
putative

[133964]pheromone receptor gene expressed in human olfactory
mucosa. Nature

[133965]Genet. 26: 18–19, 2000.

[133966]

[133967]

[133968]12090.

[133969]Stern, K.; McClintock, M. K.: Regulation of ovulation by
human

[133970]pheromones. Nature 392: 177–179, 1998.

[133971]

[133972]

[133973]12091.

[133974]Koseki, T.; Inohara, N.; Chen, S.; Nunez, G.: ARC, an in-
hibitor

[133975]of apoptosis expressed in skeletal muscle and heart that
interacts

[133976]selectively with caspases. Proc. Nat. Acad. Sci. 95:
5156–5160,

[133977]1998.

[133978]

[133979]

[133980]12092.

[133981]Li, P.–F.; Li, J.; Muller, E.–C.; Otto, A.; Dietz, R.; von Hars–
dorf,

[133982]R.: Phosphorylation by protein kinase CK2: a signaling
switch for

[133983]the caspase–inhibiting protein ARC. Molec. Cell 10:
247–258, 2002.

[133984]

[133985]

[133986]12093.

[133987]Vidal–Taboada, J. M.; Lu, A.; Pique, M.; Pons, G.; Gil, J.;
Oliva,

[133988]R.: Down syndrome critical region gene 2: expression dur–
ing mouse

[133989]development and in human cell lines indicates a function related to

[133990]cell proliferation. Biochem. Biophys. Res. Commun. 272: 156–163,

[133991]2000.

[133992]

[133993]

[133994]12094.

[133995]Muto, A.; Hoshino, H.; Madisen, L.; Yanai, N.; Obinata, M.; Karasuyama,

[133996]H.; Hayashi, N.; Nakauchi, H.; Yamamoto, M.; Groudine, M.; Igarashi,

[133997]K.: Identification of Bach2 as a B–cell–specific partner for small

[133998]Maf proteins that negatively regulate the immunoglobulin heavy chain

[133999]gene 3–prime enhancer. EMBO J. 17: 5734–5743, 1998.

[134000]

[134001]

[134002]12095.

[134003]Sasaki, S.; Ito, E.; Toki, T.; Maekawa, T.; Kanezaki, R.;
Umenai,

[134004]T.; Muto, A.; Nagai, H.; Kinoshita, T.; Yamamoto, M.; In-
azawa, J.;

[134005]Taketo, M. M.; Nakahata, T.; Igarashi, K.; Yokoyama, M.:
Cloning

[134006]and expression of human B cell-specific transcription fac-
tor BACH2

[134007]mapped to chromosome 6q15. Oncogene 19: 3739–3749,
2000.

[134008]

[134009]

[134010]12096.

[134011]Gayle, M. A.; Slack, J. L.; Bonnert, T. P.; Renshaw, B. R.;
Sonoda,

[134012]G.; Taguchi, T.; Testa, J. R.; Dower, S. K.; Sims, J. E.:
Cloning

[134013]of a putative ligand for the T1/ST2 receptor. J. Biol. Chem.
271:

[134014]5784–5789, 1996.

[134015]

[134016]

[134017]12097.

[134018]Shimaoka, T.; Kume, N.; Minami, M.; Hayashida, K.;
Kataoka, H.;

[134019]Kita, T.; Yonehara, S.: Molecular cloning of a novel scav-
enger receptor

[134020]for oxidized low density lipoprotein, SR-PSOX, on
macrophages. J.

[134021]Biol. Chem. 275: 40663–40666, 2000.

[134022]

[134023]

[134024]12098.

[134025]Wilbanks, A.; Zondlo, S. C.; Murphy, K.; Mak, S.; Soler, D.;
Langdon,

[134026]P.; Andrew, D. P.; Wu, L.; Briskin, M.: Expression cloning of
the

[134027]STRL33/BONZO/TYMSTR ligand reveals elements of CC,
CXC, CX3C chemokines. J.

[134028]Immun. 166: 5145–5154, 2001.

[134029]

[134030]

[134031]12099.

[134032]Brill, S.; Li, S.; Lyman, C. W.; Church, D. M.; Wasmuth, J. J.;

[134033]Weissbach, L.; Bernards, A.; Snijders, A. J.: The Ras GT-
Pase-activating-protein-related

[134034]human protein IQGAP2 harbors a potential actin binding
domain and

[134035]interacts with calmodulin and Rho family GTPases. Molec.
Cell. Biol. 16:

[134036]4869-4878, 1996.

[134037]

[134038]

[134039]12100.

[134040]Bitner-Glindzicz, M.; Turnpenny, P.; Hoglund, P.; Kaari-
ainen, H.;

[134041]Sankila, E.-M.; van der Maarel, S. M.; de Kok, Y. J. M.; Rop-
ers, H.-H.;

[134042]Cremers, F. P. M.; Pembrey, M.; Malcolm, S.: Further mu-
tations in

[134043]brain 4 (POU3F4) clarify the phenotype in the X-linked
deafness, DFN3. Hum.

[134044]Molec. Genet. 4: 1467–1469, 1995.

[134045]

[134046]

[134047]12101.

[134048]de Kok, Y. J. M.; Cremers, C. W. R. J.; Ropers, H.–H.; Cremers,

[134049]F. P. M.: The molecular basis of X-linked deafness type 3 (DFN3)

[134050]in two sporadic cases: identification of a somatic mosaicism for a

[134051]POU3F4 missense mutation. Hum. Mutat. 10: 207–211, 1997.

[134052]

[134053]

[134054]12102.

[134055]de Kok, Y. J. M.; Merkx, G. F. M.; van der Maarel, S. M.; Huber,

[134056]I.; Malcolm, S.; Ropers, H.–H.; Cremers, F. P. M.: A duplication/paracentric

[134057]inversion associated with familial X-linked deafness (DFN3) suggests

[134058]the presence of a regulatory element more than 400 kb upstream of

[134059]the POU3F4 gene. Hum. Molec. Genet. 4: 2145–2150, 1995.

[134060]

[134061]

[134062]12103.

[134063]de Kok, Y. J. M.; van der Maarel, S. M.; Bitner–Glindzicz, M.;

[134064]Huber, I.; Monaco, A. P.; Malcolm, S.; Pembrey, M. E.; Ropers, H.–H.;

[134065]Cremers, F. P. M.: Association between X–linked mixed deafness and

[134066]mutations in the POU domain gene POU3F4. Science 267: 685–688, 1995.

[134067]

[134068]

[134069]12104.

[134070]Douville, P. J.; Atanasoski, S.; Tobler, A.; Fontana, A.; Schwab,

[134071]M. E.: The brain–specific POU–box gene Brn4 is a sex–

linked transcription

[134072]factor located on the human and mouse X chromosomes.

Mammalian Genome 5:

[134073]180–182, 1994.

[134074]

[134075]

[134076]12105.

[134077]Friedman, R. A.; Bykhovskaya, Y.; Tu, G.; Talbot, J. M.; Wilson,
son,

[134078]D. F.; Parnes, L. S.; Fischel–Ghodsian, N.: Molecular analysis of
sis of

[134079]the POU3F4 gene in patients with clinical and radiographic
evidence

[134080]of X–linked mixed deafness with perilymphatic gusher.
Ann. Otol.

[134081]Rhinol. Laryng. 106: 320–325, 1997.

[134082]

[134083]

[134084]12106.

[134085]Minowa, O.; Ikeda, K.; Sugitani, Y.; Oshima, T.; Nakai, S.;

Katori,

[134086]Y.; Suzuki, M.; Furukawa, M.; Kawase, T.; Zheng, Y.;

Ogura, M.; Asada,

[134087]Y.; Watanabe, K.; Yamanaka, H.; Gotoh, S.; Nishi-

Takeshima, M.; Sugimoto,

[134088]T.; Kikuchi, T.; Takasaka, T.; Noda, T.: Altered cochlear fi-

brocytes

[134089]in a mouse model of DFN3 nonsyndromic deafness. Sci-

ence 285: 1408-1411,

[134090]1999.

[134091]

[134092]

[134093]12107.

[134094]Phelps, P. D.; Reardon, W.; Pembrey, M.; Bellman, S.;

Luxom, L.

[134095]: X-linked deafness, stapes gushers and a distinctive de-

fect of the

[134096]inner ear. Neuroradiology 33: 326-330, 1991.

[134097]

[134098]

[134099]12108.

[134100]Phippard, D.; Boyd, Y.; Reed, V.; Fisher, G.; Masson, W. K.;
Evans,

[134101]E. P.; Saunders, J. C.; Crenshaw, E. B., III: The sex-linked
fidget

[134102]mutation abolishes Brn4/Pou3f4 gene expression in the
embryonic inner

[134103]ear. Hum. Molec. Genet. 9: 79–85, 2000.

[134104]

[134105]

[134106]12109.

[134107]Brown, C. J.; Miller, A. P.; Carrel, L.; Rupert, J. L.; Davies,

[134108]K. E.; Willard, H. F.: The DXS423E gene in Xp11.21 escapes
X chromosome

[134109]inactivation. Hum. Molec. Genet. 4: 251–255, 1995.

[134110]

[134111]

[134112]12110.

[134113]Sumara, I.; Vorlaufer, E.; Gieffers, C.; Peters, B. H.; Peters,

[134114]J.–M.: Characterization of vertebrate cohesin complexes

and their

[134115]regulation in prophase. J. Cell Biol. 151: 749–761, 2000.

[134116]

[134117]

[134118]12111.

[134119]Lowe, D. G.; Dizhoor, A. M.; Liu, K.; Gu, Q.; Spencer, M.;
Laura,

[134120]R.; Lu, L.; Hurley, J. B.: Cloning and expression of a second
photoreceptor-specific

[134121]membrane retina guanylyl cyclase (RetGC), RetGC-2. Proc.
Nat. Acad.

[134122]Sci. 92: 5535–5539, 1995.

[134123]

[134124]

[134125]12112.

[134126]Teague, P. W.; Aldred, M. A.; Jay, M.; Dempster, M.; Harri-
son,

[134127]C.; Carothers, A. D.; Hardwick, L. J.; Evans, H. J.; Strain, L.;
Brock,

[134128]D. J. H.; Bunday, S.; Jay, B.; Bird, A. C.; Bhattacharya, S. S.;

Wright,

[134129]A. F.: Heterogeneity analysis in 40 X-linked retinitis pigmentosa

[134130]families. Am. J. Hum. Genet. 55: 105–111, 1994.

[134131]

[134132]

[134133]12113.

[134134]Bodrug, S. E.; Ray, P. N.; Gonzalez, I. L.; Schmickel, R. D.;

[134135]Sylvester, J. E.; Worton, R. G.: Molecular analysis of a constitutional

[134136]X-autosome translocation in a female with muscular dystrophy. Science 237:

[134137]1620–1624, 1987.

[134138]

[134139]

[134140]12114.

[134141]Boyce, F. M.; Beggs, A. H.; Feener, C.; Kunkel, L. M.: Dystrophin

[134142]is transcribed in brain from a distant upstream promoter. Proc. Nat.

[134143]Acad. Sci. 88: 1276–1280, 1991.

[134144]

[134145]

[134146]12115.

[134147]Boyd, Y.; Buckle, V. J.: Cytogenetic heterogeneity of
translocations

[134148]associated with Duchenne muscular dystrophy. Clin.
Genet. 29: 108–115,

[134149]1986.

[134150]

[134151]

[134152]12116.

[134153]Bulman, D. E.; Gangopadhyay, S. B.; Bebchuck, K. G.; Wor-
ton, R.

[134154]G.; Ray, P. N.: Point mutation in the human dystrophin
gene: identification

[134155]through Western blot analysis. Genomics 10: 457–460,
1991.

[134156]

[134157]

[134158]12117.

[134159]Burke, J. F.; Mogg, A. E.: Suppression of a nonsense mutation

[134160]in mammalian cells in vivo by the aminoglycoside antibiotics G-418

[134161]and paromomycin. Nucleic Acids Res. 13: 6265-6272, 1985.

[134162]

[134163]

[134164]12118.

[134165]Burnette, W. N.: 'Western blotting': electrophoretic transfer

[134166]of proteins from sodium dodecyl sulfate-polyacrylamide gels to unmodified

[134167]nitrocellulose and radiographic detection with antibody and radioiodinated

[134168]protein A. Anal. Biochem. 112: 195-203, 1981.

[134169]

[134170]

[134171]12119.

[134172]Chamberlain, J. S.; Pearlman, J. A.; Muzny, D. M.; Gibbs, R.

A.;

[134173]Ranier, J. E.; Reeves, A. A.; Caskey, C. T.: Expression of the murine

[134174]Duchenne muscular dystrophy gene in muscle and brain.

Science 239:

[134175]1416–1418, 1988.

[134176]

[134177]

[134178]12120.

[134179]Chelly, J.; Concordet, J.–P.; Kaplan, J.–C.; Kahn, A.: Illegitimate

[134180]transcription: transcription of any gene in any cell type.

Proc.

[134181]Nat. Acad. Sci. 86: 2617–2621, 1989.

[134182]

[134183]

[134184]12121.

[134185]Chelly, J.; Gilgenkrantz, H.; Hugnot, J. P.; Hamard, G.; Lambert,

[134186]M.; Recan, D.; Akli, S.; Cometto, M.; Kahn, A.; Kaplan, J. C.:

Illegitimate

[134187]transcription: application to the analysis of truncated transcripts

[134188]of the dystrophin gene in nonmuscle cultured cells from Duchenne and

[134189]Becker patients. J. Clin. Invest. 88: 1161–1166, 1991.

[134190]

[134191]

[134192]12122.

[134193]Chelly, J.; Hamard, G.; Koulakoff, A.; Kaplan, J.–C.; Kahn, A.;

[134194]Berwald–Netter, Y.: Dystrophin gene transcribed from different promoters

[134195]in neuronal and glial cells. Nature 344: 64–65, 1990.

[134196]

[134197]

[134198]12123.

[134199]Chelly, J.; Kaplan, J.–C.; Maire, P.; Gautron, S.; Kahn, A.:

[134200]Transcription of the dystrophin gene in human muscle and non–muscle

[134201]tissues. Nature 333: 858–860, 1988.

[134202]

[134203]

[134204]12124.

[134205]Clemens, P. R.; Ward, P. A.; Caskey, C. T.; Bulman, D. E.;
Fenwick,

[134206]R. G.: Premature chain termination mutation causing
Duchenne muscular

[134207]dystrophy. Neurology 42: 1775–1782, 1992.

[134208]

[134209]

[134210]12125.

[134211]Cooper, B. J.; Valentine, B. A.; Wilson, S.; Patterson, D. F.;

[134212]Concannon, P. W.: Canine muscular dystrophy: confirma-
tion of X-linked

[134213]inheritance. J. Hered. 79: 405–408, 1988.

[134214]

[134215]

[134216]12126.

[134217]Covone, A. E.; Lerone, M.; Romeo, G.: Genotype–pheno–
type correlation

[134218]and germline mosaicism in DMD/BMD patients with dele–
tions of the dystrophin

[134219]gene. Hum. Genet. 87: 353–360, 1991.

[134220]

[134221]

[134222]12127.

[134223]Cox, G. A.; Cole, N. M.; Matsumura, K.; Phelps, S. F.;
Hauschka,

[134224]S. D.; Campbell, K. P.; Faulkner, J. A.; Chamberlain, J. S.:
Overexpression

[134225]of dystrophin in transgenic mdx mice eliminates dys–
trophic symptoms

[134226]without toxicity. Nature 364: 725–729, 1993.

[134227]

[134228]

[134229]12128.

[134230]Cox, G. A.; Sunada, Y.; Campbell, K. P.; Chamberlain, J. S.:

[134231]Dp71 can restore the dystrophin–associated glycoprotein

complex in

[134232]muscle but fails to prevent dystrophy. Nature Genet. 8:
333–339,

[134233]1994.

[134234]

[134235]

[134236]12129.

[134237]Comi, G. P.; Ciafaloni, E.; de Silva, H. A. R.; Prella, A.; Bar-
doni,

[134238]A.; Rigoletto, C.; Robotti, M.; Bresolin, N.; Moggio, M.;
Fortunato,

[134239]F.; Ciscato, P.; Turconi, A.; Rose, A. D.; Scarlato, G.: A
G(+1)–to–A

[134240]transversion at the 5–prime splice site of intron 69 of the
dystrophin

[134241]gene causing the absence of peripheral nerve Dp116 and
severe clinical

[134242]involvement in a DMD patient. Hum. Molec. Genet. 4:
2171–2174, 1995.

[134243]

[134244]

[134245]12130.

[134246]Crawford, G. E.; Lu, Q. L.; Partridge, T. A.; Chamberlain, J.

[134247]S.: Suppression of revertant fibers in mdx mice by expression of

[134248]a functional dystrophin. Hum. Molec. Genet. 10: 2745–2750, 2001.

[134249]

[134250]

[134251]12131.

[134252]Darras, B. T.; Blattner, P.; Harper, J. F.; Spiro, A. J.; Alter,

[134253]S.; Francke, U.: Intragenic deletions in 21 Duchenne muscular dystrophy

[134254](DMD)/Becker muscular dystrophy (BMD) families studied with the dystrophin

[134255]cDNA: location of breakpoints on HindIII and BglIII exon-containing

[134256]fragment maps, meiotic and mitotic origin of the mutations. Am. J.

[134257]Hum. Genet. 43: 620–629, 1988.

[134258]

[134259]

[134260]12132.

[134261]Darras, B. T.; Francke, U.: Normal human genomic restriction-fragment

[134262]patterns and polymorphisms revealed by hybridization with the entire

[134263]dystrophin cDNA. Am. J. Hum. Genet. 43: 612-619, 1988.

[134264]

[134265]

[134266]12133.

[134267]Darras, B. T.; Francke, U.: A partial deletion of the muscular

[134268]dystrophy gene transmitted twice by an unaffected male. Nature 329:

[134269]556-558, 1987.

[134270]

[134271]

[134272]12134.

[134273]Davies, K. E.; Smith, T. J.; Bunday, S.; Read, A. P.; Flint, T.;

[134274]Bell, M.; Speer, A.: Mild and severe muscular dystrophy associated

[134275]with deletions in Xp21 of the human X chromosome. J. Med. Genet. 25:

[134276]9–13, 1988.

[134277]

[134278]

[134279]12135.

[134280]De Angelis, F. G.; Sthandier, O.; Berarducci, B.; Toso, S.; Galluzzi,

[134281]G.; Ricci, E.; Cossu, G.; Bozzoni, I.: Chimeric snRNA molecules carrying

[134282]antisense sequences against the splice junctions of exon 51 of the

[134283]dystrophin pre-mRNA induce exon skipping and restoration of a dystrophin

[134284]synthesis in delta-48-50 DMD cells. Proc Nat. Acad. Sci. 99: 9456–9461,

[134285]2002.

[134286]

[134287]

[134288]12136.

[134289]den Dunnen, J. T.; Bakker, E.; Klein Breteler, E. G.; Pearson,

[134290]P. L.; van Ommen, G. J. B.: Direct mutation of more than
50% of the

[134291]Duchenne muscular dystrophy mutations by field inver-
sion gels. Nature 329:

[134292]640–642, 1987.

[134293]

[134294]

[134295]12137.

[134296]Dickson, G.; Pizzey, J. A.; Elsom, V. E.; Love, D.; Davies, K.

[134297]E.; Walsh, F. S.: Distinct dystrophin mRNA species are ex-
pressed

[134298]in embryonic and adult mouse skeletal muscle. FEBS Lett.
242: 47–52,

[134299]1988.

[134300]

[134301]

[134302]12138.

[134303]Dominguez–Steglich, M.; Meng, G.; Bettecken, T.; Muller, C. R.;

[134304]Schmid, M.: The dystrophin gene is autosomally located on a microchromosome

[134305]in chicken. Genomics 8: 536–540, 1990.

[134306]

[134307]

[134308]12139.

[134309]Doolittle, R. F.: Similar amino acid sequences: chance or common

[134310]ancestry? Science 214: 149–159, 1981.

[134311]

[134312]

[134313]12140.

[134314]Bassi, M. T.; Bergen, A. A. B.; Bitoun, P.; Charles, S. J.; Clementi,

[134315]M.; Gosselin, R.; Hurst, J.; Lewis, R. A.; Lorenz, B.; Meitinger,

[134316]T.; Messiaen, L.; Ramesar, R. S.; Ballabio, A.; Schiaffino, M. V.

[134317]: Diverse prevalence of large deletions within the OA1
gene in ocular

[134318]albinism type 1 patients from Europe and North America.
Hum. Genet. 108:

[134319]51–54, 2001.

[134320]

[134321]

[134322]12141.

[134323]Bassi, M. T.; Schiaffino, M. V.; Renieri, A.; De Nigris, F.;
Galli,

[134324]L.; Bruttini, M.; Gebbia, M.; Bergen, A. A. B.; Lewis, R. A.;
Ballabio,

[134325]A.: Cloning of the gene for ocular albinism type 1 from the
distal

[134326]short arm of the X chromosome. Nature Genet. 10: 13–19,
1995.

[134327]

[134328]

[134329]12142.

[134330]Bergen, A. A. B.; Samanns, C.; Schuurman, E. J. M.; van
Osch, L.;

[134331]van Dorp, D. B.; Pinckers, A. J. L. G.; Bakker, E.; Gal, A.;
van Ommen,

[134332]G. J. B.; Bleeker–Wagemakers, E. M.: Multipoint linkage
analysis

[134333]in X-linked ocular albinism of the Nettleship–Falls type.
Hum. Genet. 88:

[134334]162–166, 1991.

[134335]

[134336]

[134337]12143.

[134338]Bergen, A. A. B.; Schuurman, E. J. M.; van den Born, L. I.;
Samanns,

[134339]C.; van Dorp, D. B.; Pinckers, A. J. L. G.; Bakker, E.; van
Ommen,

[134340]G. J. B.; Gal, A.; Bleeker–Wagemakers, E. M.: Carrier detec-
tion in

[134341]X-linked ocular albinism of the Nettleship–Falls type by
DNA analysis. Clin.

[134342]Genet. 41: 135–138, 1992.

[134343]

[134344]

[134345]12144.

[134346]Bergen, A. A. B.; Zijp, P.; Schuurman, E. J. M.; Bleeker-
Wagemakers,

[134347]E. M.; Apkarian, P.; van Ommen, G.-J. B.: Refinement of
the localization

[134348]of the X-linked ocular albinism gene. Genomics 16:
272-273, 1993.

[134349]

[134350]

[134351]12145.

[134352]Bouloux, P.-M. G.; Kirk, J.; Munroe, P.; Duke, V.; Meindl,
A.;

[134353]Hilson, A.; Grant, D.; Carter, N.; Betts, D.; Meitinger, T.;
Besser,

[134354]G. M.: Deletion analysis maps ocular albinism proximal to
the steroid

[134355]sulphatase locus. Clin. Genet. 43: 169-173, 1993.

[134356]

[134357]

[134358]12146.

[134359]Charles, S. J.; Green, J. S.; Moore, A. T.; Barton, D. E.;
Yates,

[134360]J. R. W.: Genetic mapping of X-linked ocular albinism:
linkage analysis

[134361]in a large Newfoundland kindred. Genomics 16: 259–261,
1993.

[134362]

[134363]

[134364]12147.

[134365]Charles, S. J.; Moore, A. T.; Yates, J. R. W.: Genetic map-
ping

[134366]of X linked ocular albinism: linkage analysis in British
families. J.

[134367]Med. Genet. 29: 552–554, 1992.

[134368]

[134369]

[134370]12148.

[134371]Cooper, D. N.; Krawczak, M.: Mechanisms of insertional
mutagenesis

[134372]in human genes causing genetic disease. Hum. Genet. 87:
409–415,

[134373]1991.

[134374]

[134375]

[134376]12149.

[134377]Creel, D.; O'Donnell, F. E., Jr.; Witkop, C. J., Jr.: Visual

[134378]system anomalies in human ocular albinos. Science 201:
931-933,

[134379]1978.

[134380]

[134381]

[134382]12150.

[134383]d'Addio, M.; Pizzigoni, A.; Bassi, M. T.; Baschirotto, C.;
Valetti,

[134384]C.; Incerti, B.; Clementi, M.; De Luca, M.; Ballabio, A.; Schi-
affino,

[134385]M. V.: Defective intracellular transport and processing of
OA1 is

[134386]a major cause of ocular albinism type 1. Hum. Molec.
Genet. 9: 3011-3018,

[134387]2000.

[134388]

[134389]

[134390]12151.

[134391]Engelhard, C. F.: Eine Familie mit hereditaerem Nystagmus. Z.

[134392]Ges. Neurol. Psychiat. 28: 319–338, 1915.

[134393]

[134394]

[134395]12152.

[134396]Fialkow, P. J.; Giblett, E. R.; Motulsky, A. G.: Measurable linkage

[134397]between ocular albinism and Xg. Am. J. Hum. Genet. 19: 63–69, 1967.

[134398]

[134399]

[134400]12153.

[134401]Francois, J.; Deweer, J. P.: Albinisme oculaire lie au sexe et

[134402]alterations caracteristiques du fond d'oeil chez les femmes heterozygotes. Ophthalmologica 126:

[134403]209–221, 1953.

[134404]

[134405]

[134406]12154.

[134407]Gillespie, F. D.: Ocular albinism with report of a family
with

[134408]female carriers. Arch. Ophthal. 66: 774–777, 1961.

[134409]

[134410]

[134411]12155.

[134412]Incerti, B.; Cortese, K.; Pizzigoni, A.; Surace, E. M.; Varani,

[134413]S.; Coppola, M.; Jeffery, G.; Seeliger, M.; Jaissle, G.; Ben-
nett,

[134414]D. C.; Marigo, V.; Schiaffino, M. V.; Tacchetti, C.; Ballabio,
A.

[134415]: Oa1 knock-out: new insights on the pathogenesis of oc-
ular albinism

[134416]type 1. Hum. Molec. Genet. 9: 2781–2788, 2000.

[134417]

[134418]

[134419]12156.

[134420]Jaeger, C.; Jay, B.: X-linked ocular albinism: a family containing

[134421]a manifesting heterozygote, and an affected male married to a female

[134422]with autosomal recessive ocular albinism. Hum. Genet. 56: 299–304,

[134423]1981.

[134424]

[134425]

[134426]12157.

[134427]Johnson, G. J.; Gillan, J. G.; Pearce, W. G.: Ocular albinism

[134428]in Newfoundland. Can. J. Ophthal. 6: 237–248, 1971.

[134429]

[134430]

[134431]12158.

[134432]Kidd, J. R.; Castiglione, C. M.; Davies, K. E.; Pakstis, A. J.;

[134433]Gusella, J.; Sparkes, R. S.; Pearson, P.; Willard, H.; Kidd, K.

K.

[134434]: Mapping the locus for X-linked ocular albinism (OA).

(Abstract) Am.

[134435]J. Hum. Genet. 37: A161 only, 1985.

[134436]

[134437]

[134438]12159.

[134439]Krawczak, M.; Reiss, J.; Cooper, D. N.: The mutational
spectrum

[134440]of single base-pair substitutions in mRNA splice junctions
of human

[134441]genes: causes and consequences. Hum. Genet. 90: 41-54,
1992.

[134442]

[134443]

[134444]12160.

[134445]Lein, J. N.; Stewart, C. T.; Moll, F. C.: Sex-linked hereditary

[134446]nystagmus. Pediatrics 18: 214-217, 1956.

[134447]

[134448]

[134449]12161.

[134450]Lyon, M. F.: Sex chromatin and gene action in the mammalian X-chromosome. Am.

[134451]J. Hum. Genet. 14: 135–148, 1962.

[134452]

[134453]

[134454]12162.

[134455]Meindl, A.; Hosenfeld, D.; Bruckl, W.; Schuffenhauer, S.; Jenderny,

[134456]J.; Bacskulin, A.; Oppermann, H.–C.; Swensson, O.; Bouloux, P.; Meitinger,

[134457]T.: Analysis of a terminal Xp22.3 deletion in a patient with six

[134458]monogenic disorders: implications for the mapping of X linked ocular

[134459]albinism. J. Med. Genet. 30: 838–842, 1993.

[134460]

[134461]

[134462]12163.

[134463]Negrelli, B. C.: L'albinisme oculaire lie au sexe dans le
cadre

[134464]du depistage des heterozygotes en ophtalmologie. J.
Genet. Hum. 8:

[134465]108 only, 1959.

[134466]

[134467]

[134468]12164.

[134469]Newton, J. M.; Orlow, S. J.; Barsh, G. S.: Isolation and char-
acterization

[134470]of a mouse homolog of the X-linked ocular albinism (OA1)
gene. Genomics 37:

[134471]219-225, 1996.

[134472]

[134473]

[134474]12165.

[134475]O'Donnell, F. E., Jr.; Green, W. R.; Fleishman, J. A.; Ham-
brick,

[134476]G. W.: X-linked ocular albinism in Blacks: ocular albinism
cum pigmento. Arch.

[134477]Ophthal. 96: 1189–1192, 1978.

[134478]

[134479]

[134480]12166.

[134481]O'Donnell, F. E., Jr.; Hambrick, G. W., Jr.; Green, W. R.; Iliff,

[134482]W. J.; Stone, D. L.: X-linked ocular albinism: an oculocutaneous

[134483]macromelanosomal disorder. Arch. Ophthal. 94:
1883–1892, 1976.

[134484]

[134485]

[134486]12167.

[134487]Oetting, W. S.: New insights into ocular albinism type 1
(OA1):

[134488]mutations and polymorphisms of the OA1 gene. Hum.
Mutat. 19: 85–92,

[134489]2002.

[134490]

[134491]

[134492]12168.

[134493]Pearce, W. G.; Johnson, G. J.; Gillan, J. G.: Nystagmus in a
[134494]female carrier of ocular albinism. J. Med. Genet. 9:
126-128, 1972.

[134495]

[134496]

[134497]12169.

[134498]Pearce, W. G.; Johnson, G. J.; Sanger, R.: Ocular albinism
and

[134499]Xg. (Letter) Lancet I: 1072 only, 1971.

[134500]

[134501]

[134502]12170.

[134503]Pearce, W. G.; Sanger, R.: X mapping in man: evidence
against

[134504]direct measurable linkage between ocular albinism and
deutan colour

[134505]blindness. J. Med. Genet. 13: 319 only, 1976.

[134506]

[134507]

[134508]12171.

[134509]Gaugitsch, H. W.; Prieschl, E. E.; Kalthoff, F.; Huber, N. E.;

[134510]Baumruker, T.: A novel transiently expressed, integral
membrane protein

[134511]linked to cell activation: molecular cloning via the rapid
degradation

[134512]signal AUUUA. J. Biol. Chem 267: 11267–11273, 1992.

[134513]

[134514]

[134515]12172.

[134516]Kanai, Y.; Segawa, H.; Miyamoto, K.; Uchino, H.; Takeda,
H.; Endou,

[134517]H.: Expression cloning and characterization of a trans-
porter for

[134518]large neutral amino acids activated by the heavy chain of
4F2 antigen

[134519](CD98). J. Biol. Chem. 273: 23629–23632, 1998.

[134520]

[134521]

[134522]12173.

[134523]Takahara, K.; Kessler, E.; Biniaminov, L.; Brusel, M.; Eddy, R.

[134524]L.; Jani-Sait, S.; Shows, T. B.; Greenspan, D. S.: Type I pro-collagen

[134525]COOH-terminal proteinase enhancer protein: identification, primary

[134526]structure, and chromosomal localization of the cognate human gene

[134527](PCOLCE). J. Biol. Chem. 269: 26280–26285, 1994.

[134528]

[134529]

[134530]12174.

[134531]Takahara, K.; Osborne, L.; Elliott, R. W.; Tsui, L.-C.;
Scherer,

[134532]S. W.; Greenspan, D. S.: Fine mapping of the human and
mouse genes

[134533]for the type I procollagen COOH-terminal proteinase en-
hancer protein. Genomics 31:

[134534]253–256, 1996.

[134535]

[134536]

[134537]12175.

[134538]Amagai, M.; Wang, Y.; Minoshima, S.; Kawamura, K.;
Green, K. J.;

[134539]Nishikawa, T.; Shimizu, N.: Assignment of the human
genes for desmocollin

[134540]3 (DSC3) and desmocollin 4 (DSC4) to chromosome
18q12. Genomics 25:

[134541]330-332, 1995.

[134542]

[134543]

[134544]12176.

[134545]Kawamura, K.; Watanabe, K.; Suzuki, T.; Yamakawa, T.;
Kamiyama,

[134546]T.; Nakagawa, H.; Tsurufuji, S.: cDNA cloning and expres-
sion of a

[134547]novel human desmocollin. J. Biol. Chem. 269:
26295-26302, 1994.

[134548]

[134549]

[134550]12177.

[134551]King, I. A.; Sullivan, K. H.; Bennett, R., Jr.; Buxton, R. S.:

[134552]The desmocollins of human foreskin epidermis: identifica-
tion and chromosomal

[134553]assignment of a third gene and expression patterns of the
three isoforms. J.

[134554]Invest. Derm. 105: 314-321, 1995.

[134555]

[134556]

[134557]12178.

[134558]Warburg, O.: On the origin of cancer cells. Science 123:
309–314,

[134559]1956.

[134560]

[134561]

[134562]12179.

[134563]Yang, R.–B.; Foster, D. C.; Garbers, D. L.; Fulle, H.–J.: Two
[134564]membrane forms of guanylyl cyclase found in the eye.
Proc. Nat. Acad.

[134565]Sci. 92: 602–606, 1995.

[134566]

[134567]

[134568]12180.

[134569]Dubrovsky, A. L.; Taratuto, A. L.; Sevelever, G.; Schultz, M.;

[134570]Pegoraro, E.; Hoop, R. C.; Hoffman, E. P.: Duchenne mus-
cular dystrophy

[134571]and myotonic dystrophy in the same patient. Am. J. Med.

Genet. 55:

[134572]342–348, 1995.

[134573]

[134574]

[134575]12181.

[134576]Cazzola, M.; Bergamaschi, G.: X-linked Wiskott–Aldrich
syndrome

[134577]in a girl. (Letter) New Eng. J. Med. 338: 1850 only, 1998.

[134578]

[134579]

[134580]12182.

[134581]Emery, A. E. H.: Duchenne Muscular Dystrophy. Oxford,
UK: Oxford

[134582]University Press (pub.) (2nd ed.): 1993.

[134583]

[134584]

[134585]12183.

[134586]Fabb, S. A.; Wells, D. J.; Serpente, P.; Dickson, G.: Adeno-
associated

[134587]virus vector gene transfer and sarcolemmal expression of
a 144 kDa

[134588]micro-dystrophin effectively restores the dystrophin-as-
sociated protein

[134589]complex and inhibits myofibre degeneration in nude/mdx
mice. Hum.

[134590]Molec. Genet. 11: 733–741, 2002.

[134591]

[134592]

[134593]12184.

[134594]Feener, C. A.; Boyce, F. M.; Kunkel, L. M.: Rapid detection
of

[134595]CA polymorphisms in cloned DNA: application to the
5–prime region

[134596]of the dystrophin gene. Am. J. Hum. Genet. 48: 621–627,
1991.

[134597]

[134598]

[134599]12185.

[134600]Ferlini, A.; Galie, N.; Merlini, L.; Sewry, C.; Branzi, A.;
Muntoni,

[134601]F.: A novel Alu-like element rearranged in the dystrophin gene causes

[134602]a splicing mutation in a family with X-linked dilated cardiomyopathy. Am.

[134603]J. Hum. Genet. 63: 436–446, 1998.

[134604]

[134605]

[134606]12186.

[134607]Finnegan, D. J.: Eukaryotic transposable elements and genome

[134608]evolution. Trends Genet. 5: 103–107, 1989.

[134609]

[134610]

[134611]12187.

[134612]Forrest, S. M.; Cross, G. S.; Speer, A.; Gardner-Medwin, D.; Burn,

[134613]J.; Davies, K. E.: Preferential deletion of exons in Duchenne and

[134614]Becker muscular dystrophies. Nature 329: 638–640, 1987.

[134615]

[134616]

[134617]12188.

[134618]Francke, U.; Ochs, H. D.; de Martinville, B.; Giacalone, J.;
Lindgren,

[134619]V.; Distèche, C.; Pagon, R. A.; Hofker, M. H.; van Ommen,
G.-J. B.;

[134620]Pearson, P. L.; Wedgwood, R. J.: Minor Xp21 chromosome
deletion in

[134621]a male associated with expression of Duchenne muscular
dystrophy,

[134622]chronic granulomatous disease, retinitis pigmentosa, and
McLeod syndrome. Am.

[134623]J. Hum. Genet. 37: 250–267, 1985.

[134624]

[134625]

[134626]12189.

[134627]Furst, D.; Nave, R.; Osborn, M.; Weber, K.; Bardosi, A.;
Archidiacono,

[134628]N.; Ferro, M.; Romano, V.; Romeo, G.: Nebulin and titin
expression

[134629]in Duchenne muscular dystrophy appears normal. FEBS

Lett. 224: 49–53,
[134630]1987.

[134631]

[134632]

[134633]12190.

[134634]Giacalone, J. P.; Francke, U.: Common sequence motifs at
the

[134635]rearrangement sites of a constitutional X/autosome
translocation and

[134636]associated deletion. Am. J. Hum. Genet. 50: 725–741,
1992.

[134637]

[134638]

[134639]12191.

[134640]Gillard, E. F.; Chamberlain, J. S.; Murphy, E. G.; Duff, C. L.;

[134641]Smith, B.; Burghes, A. H. M.; Thompson, M. W.; Suther-
land, J.; Oss,

[134642]I.; Bodrug, S. E.; Klamut, H. J.; Ray, P. N.; Worton, R. G.:
Molecular

[134643]and phenotypic analysis of patients with deletions within

the deletion-rich

[134644]region of the Duchenne muscular dystrophy (DMD) gene.

Am. J. Hum.

[134645]Genet. 45: 507–520, 1989.

[134646]

[134647]

[134648]12192.

[134649]Ginjaar, I. B.; Kneppers, A. L. J.; Meulen, J.–D. M.; Ander-
son,

[134650]L. V. B.; Bremmer–Bout, M.; van Deutekom, J. C. T.; Wee-
genaar, J.;

[134651]den Dunnen, J. T.; Bakker, E.: Dystrophin nonsense muta-
tion induces

[134652]different levels of exon 29 skipping and leads to variable
phenotypes

[134653]within one BMD family. Europ. J. Hum. Genet. 8: 793–796,
2000.

[134654]

[134655]

[134656]12193.

[134657]Greenberg, D. S.; Sunada, Y.; Campbell, K. P.; Yaffe, D.;

Nudel,

[134658]U.: Exogenous Dp71 restores the levels of dystrophin associated proteins

[134659]but does not alleviate muscle damage in mdx mice. Nature Genet. 8:

[134660]340–344, 1994.

[134661]

[134662]

[134663]12194.

[134664]Gussoni, E.; Soneoka, Y.; Strickland, C. D.; Buzney, E. A.; Khan,

[134665]M. K.; Flint, A. F.; Kunkel, L. M.; Mulligan, R. C.: Dystrophin expression

[134666]in the mdx mouse restored by stem cell transplantation.

Nature 401:

[134667]390–394, 1999.

[134668]

[134669]

[134670]12195.

[134671]Hagiwara, Y.; Mizuno, Y.; Takemitsu, M.; Matsuzaki, T.;

Nonaka,

[134672]I.; Ozawa, E.: Dystrophin-positive muscle fibers following
C2 myoblast

[134673]transplantation into mdx nude mice. Acta Neuropath. 90:
592–600,

[134674]1995.

[134675]

[134676]

[134677]12196.

[134678]Hagiwara, Y.; Nishio, H.; Kitoh, Y.; Takeshima, Y.; Narita,
N.;

[134679]Wada, H.; Yokoyama, M.; Nakamura, H.; Matsuo, M.: A
novel point mutation

[134680](G(–1) to T) in a 5–prime splice donor site of intron 13 of
the dystrophin

[134681]gene results in exon skipping and is responsible for
Becker muscular

[134682]dystrophy. Am. J. Hum. Genet. 54: 53–61, 1994.

[134683]

[134684]

[134685]12197.

[134686]Hammonds, R. G., Jr.: Protein sequence of DMD gene is related

[134687]to actin-binding domain of alpha-actinin. (Letter) Cell 51: 1, 1987.

[134688]

[134689]

[134690]12198.

[134691]Harper, S. Q.; Hauser, M. A.; DelloRusso, C.; Duan, D.; Crawford,

[134692]R. W.; Phelps, S. F.; Harper, H. A.; Robinson, A. S.; Engelhardt,

[134693]J. F.; Brooks, S. V.; Chamberlain, J. S.: Modular flexibility of

[134694]dystrophin: implications for gene therapy of Duchenne muscular dystrophy. Nature

[134695]Med. 8: 253-261, 2002.

[134696]

[134697]

[134698]12199.

[134699]Hart, K. A.; Hodgson, S.; Walker, A.; Cole, C. G.; Johnson, L.;

[134700]Dubowitz, V.; Bobrow, M.: DNA deletions in mild and severe Becker

[134701]muscular dystrophy. Hum. Genet. 75: 281–285, 1987.

[134702]

[134703]

[134704]12200.

[134705]Hodgson, S. V.; Abbs, S.; Clark, S.; Manzur, A.; Heckmatt, J.

[134706]Z. H.; Dubowitz, V.; Bobrow, M.: Correlation of clinical and deletion

[134707]data in Duchenne and Becker muscular dystrophy, with special reference

[134708]to mental ability. Neuromusc. Disord. 2: 269–276, 1992.

[134709]

[134710]

[134711]12201.

[134712]Hoffman, E. P.; Brown, R. H., Jr.; Kunkel, L. M.: The protein

[134713]product of the Duchenne muscular dystrophy locus. Cell 51: 919–928,

[134714]1987.

[134715]

[134716]

[134717]12202.

[134718]Hoffman, E. P.; Knudson, C. M.; Campbell, K. P.; Kunkel, L. M.

[134719]: Subcellular fractionation of dystrophin to the triads of skeletal

[134720]muscle. Nature 330: 754–758, 1987.

[134721]

[134722]

[134723]12203.

[134724]Hoffman, E. P.; Monaco, A. P.; Feener, C. C.; Kunkel, L. M.:

[134725]Conservation of the Duchenne muscular dystrophy gene in mice and humans. Science 238:

[134726]347–350, 1987.

[134727]

[134728]

[134729]12204.

[134730]Hoop, R. C.; Russo, L. S.; Riconda, D. L.; Schwartz, L. S.;

Hoffman,

[134731]E. P.: Restoration of half the normal dystrophin sequence
in a double-deletion

[134732]Duchenne muscular dystrophy family. Am. J. Med. Genet.
49: 323–327,

[134733]1994.

[134734]

[134735]

[134736]12205.

[134737]Hoffman, E. P.; Fischbeck, K. H.; Brown, R. H.; Johnson, M.;
Medori,

[134738]R.; Loike, J. D.; Harris, J. B.; Waterston, R.; Brooke, M.;
Specht,

[134739]L.; Kupsky, W.; Chamberlain, J.; Caskey, C. T.; Shapiro, F.;
Kunkel,

[134740]L. M.: Characterization of dystrophin in muscle–biopsy
specimens

[134741]from patients with Duchenne's or Becker's muscular dys-
trophy. New

[134742]Eng. J. Med. 318: 1363–1368, 1988.

[134743]

[134744]

[134745]12206.

[134746]Howard, P. L.; Dally, G. Y.; Wong, M. H.; Ho, A.; Weleber, R.

[134747]G.; Pillers, D.-A. M.; Ray, P. N.: Localization of dystrophin isoform

[134748]Dp71 to the inner limiting membrane of the retina suggests a unique

[134749]functional contribution of Dp71 in the retina. Hum. Molec. Genet. 7:

[134750]1385-1391, 1998.

[134751]

[134752]

[134753]12207.

[134754]Hu, X.; Burghes, A. H. M.; Bulman, D. E.; Ray, P. N.; Worton,

[134755]R. G.: Evidence for mutation by unequal sister chromatid exchange

[134756]in the Duchenne muscular dystrophy gene. Am. J. Hum. Genet. 44:

[134757]855-863, 1989.

[134758]

[134759]

[134760]12208.

[134761]Birnbaumer, M.: Vasopressin receptors. TEM 11: 406–410,
2000.

[134762]

[134763]

[134764]12209.

[134765]Kools, P.; Vanhalst, K.; Wan den Eynde, E.; van Roy, F.: The
human

[134766]cadherin–10 gene: complete coding sequence, predomi-
nant expression

[134767]in the brain and mapping on chromosome 5p13–13. FEBS
Lett. 452:

[134768]328–334, 1999.

[134769]

[134770]

[134771]12210.

[134772]Leder, S.; Weber, Y.; Altafaj, X.; Estivill, X.; Joost, H.–G.;

[134773]Becker, W.: Cloning and characterization of DYRK1B, a novel member

[134774]of the DYRK family of protein kinases. Biochem. Biophys. Res. Commun. 254:

[134775]474–479, 1999.

[134776]

[134777]

[134778]12211.

[134779]Lee, K.; Deng, X.; Friedman, E.: Mirk protein kinase is a mitogen-activated

[134780]protein kinase substrate that mediates survival of colon cancer cells. Cancer

[134781]Res. 60: 3631–3637, 2000.

[134782]

[134783]

[134784]12212.

[134785]Coyle, A. J.; Lehar, S.; Lloyd, C.; Tian, J.; Delaney, T.; Manning,

[134786]S.; Nguyen, T.; Burwell, T.; Schneider, H.; Gonzalo, J. A.; Gosselin,

[134787]M.; Owen, L. R.; Rudd, C. E.; Gutierrez-Ramos, J. C.: The
CD28-related

[134788]molecule ICOS is required for effective T cell-dependent
immune responses. Immunity 13:

[134789]95–105, 2000.

[134790]

[134791]

[134792]12213.

[134793]Dong, C.; Juedes, A. E.; Temann, U. A.; Shresta, S.; Allison,
J.

[134794]P.; Ruddle, N. H.; Flavell, R. A.: ICOS co-stimulatory recep-
tor is

[134795]essential for T-cell activation and function. Nature 409:
97–101,

[134796]2001.

[134797]

[134798]

[134799]12214.

[134800]Haimila, K. E.; Partanen, J. A.; Holopainen, P. M.: Genetic
polymorphism

[134801]of the human ICOS gene. Immunogenetics 53:
1028–1032, 2002.

[134802]

[134803]

[134804]12215.

[134805]Hutloff, A.; Dittrich, A. M.; Beier, K. C.; Eljaschewitsch, B.;

[134806]Kraft, R.; Anagnostopoulos, I.; Kroczeck, R. A.: ICOS is an
inducible

[134807]T–cell co–stimulator structurally and functionally related
to CD28. Nature 397:

[134808]263–266, 1999.

[134809]

[134810]

[134811]12216.

[134812]McAdam, A. J.; Greenwald, R. J.; Levin, M. A.; Chernova, T.;
Malenkovich,

[134813]N.; Ling, V.; Freeman, G. J.; Sharpe, A. H.: ICOS is critical
for

[134814]CD40–mediated antibody class switching. Nature 409:
102–105, 2001.

[134815]

[134816]

[134817]12217.

[134818]Pack, S. D.; Pak, E.; Tanigami, A.; Ledbetter, D. H.; Fukuda, M.

[134819]N.: Assignment of the bystin gene BYSL to human chromosome band 6p21.1

[134820]by in situ hybridization. Cytogenet. Cell Genet. 83: 76–77, 1998.

[134821]

[134822]

[134823]12218.

[134824]Medina, M.; Marinescu, R. C.; Overhauser, J.; Kosik, K. S.: Hemizyosity

[134825]of delta-catenin (CTNND2) is associated with severe mental retardation

[134826]in cri-du-chat syndrome. Genomics 63: 157–164, 2000.

[134827]

[134828]

[134829]12219.

[134830]Paffenholz, R.; Franke, W. W.: Identification and localization

[134831]of a neurally expressed member of the plakoglobin/armadillo multigene

[134832]family. Differentiation 61: 293–304, 1997.

[134833]

[134834]

[134835]12220.

[134836]Zhou, J.; Liyanage, U.; Medina, M.; Ho, C.; Simmons, A. D.; Lovett.

[134837]M.; Kosik, K. S.: Presenilin 1 interaction in the brain with a novel

[134838]member of the armadillo family. Neuroreport 8: 2085–2090, 1997.

[134839]

[134840]

[134841]12221.

[134842]Hatzfeld, M.; Nachtsheim, C.: Cloning and characterization of

[134843]a new armadillo family member, p0071, associated with the junctional

[134844]plaque: evidence for a subfamily of closely related proteins. J.

[134845]Cell Sci. 109: 2767–2778, 1996.

[134846]

[134847]

[134848]12222.

[134849]Burger, J.; Fonknechten, N.; Hoeltzenbein, M.; Neumann, L.; Bratanoff,

[134850]E.; Hazan, J.; Reis, A.: Hereditary spastic paraplegia caused by

[134851]mutations in the SPG4 gene. Europ. J. Hum. Genet. 8: 771–776, 2000.

[134852]

[134853]

[134854]12223.

[134855]Burger, J.; Metzke, H.; Paternotte, C.; Schilling, F.; Hazan, J.;

[134856]Reis, A.: Autosomal dominant spastic paraplegia with anticipation

[134857]maps to a 4–cM interval on chromosome 2p21–p24 in a large German family. Hum.

[134858]Genet. 98: 371–375, 1996.

[134859]

[134860]

[134861]12224.

[134862]Errico, A.; Ballabio, A.; Rugarli, E. I.: Spastin, the protein

[134863]mutated in autosomal dominant hereditary spastic paraplegia, is involved

[134864]in microtubule dynamics. Hum. Molec. Genet. 11:
153–163, 2002.

[134865]

[134866]

[134867]12225.

[134868]Lindsey, J. C.; Lusher, M. E.; McDermott, C. J.; White, K. D.;

[134869]Reid, E.; Rubinsztein, D. C.; Bashir, R.; Hazan, J.; Shaw, P.
J.;

[134870]Bushby, K. M. D.: Mutation analysis of the spastin gene
(SPG4) in

[134871]patients with hereditary spastic paraplegia. J. Med. Genet.
37:

[134872]759–765, 2000.

[134873]

[134874]

[134875]12226.

[134876]Takahashi, N.; Tuiki, H.; Saya, H.; Kaibuchi, K.: Localiza-
tion

[134877]of the gene coding for ROCK II/Rho kinase on human
chromosome 2p24. Genomics 55:

[134878]235–237, 1999.

[134879]

[134880]

[134881]12227.

[134882]Gruber, A. D.; Pauli, B. U.: Clustering of the human CLCA
gene

[134883]family on the short arm of chromosome 1 (1p22–31).

Genome 42: 1030–1032,

[134884]1999.

[134885]

[134886]

[134887]12228.

[134888]Kedra, D.; Pan, H.-Q.; Seroussi, E.; Fransson, I.; Guilbaud, C.;

[134889]Collins, J. E.; Dunham, I.; Blennow, E.; Roe, B. A.; Piehl, F.; Dumanski,

[134890]J. P.: Characterization of the human synaptogyrin gene family. Hum.

[134891]Genet. 103: 131–141, 1998.

[134892]

[134893]

[134894]12229.

[134895]Hazan, J.; Davoine, C. S.; Mavel, D.; Fonknechten, N.; Paternotte,

[134896]C.; Fizames, C.; Cruaud, C.; Samson, D.; Muselet, D.; Vega-Czarny,

[134897]N.; Brice, A.; Gyapay, G.; Heilig, R.; Fontaine, B.; Weissenbach,

[134898]J.: A fine integrated map of the SPG4 locus excludes an expanded

[134899]CAG repeat in chromosome 2p-linked autosomal dominant spastic paraplegia. Genomics 60:

[134900]309–319, 1999.

[134901]

[134902]

[134903]12230.

[134904]Hazan, J.; Fonknechten, N.; Mavel, D.; Paternotte, C.; Samson,

[134905]D.; Artiguenave, F.; Davoine, C.-S.; Cruaud, C.; Durr, A.; Wincker,

[134906]P.; Brottier, P.; Cattolico, L.; Barbe, V.; Burgunder, J.-M.; Prud'homme,

[134907]J.-F.; Brice, A.; Fontaine, B.; Heilig, R.; Weissenbach, J.: Spastin,

[134908]a new AAA protein, is altered in the most frequent form of autosomal

[134909]dominant spastic paraplegia. Nature Genet. 23: 296–303, 1999.

[134910]

[134911]

[134912]12231.

[134913]Bedford, F. K.; Ashworth, A.; Enver, T.; Wiedemann, L. M.: HEX:

[134914]a novel homeobox gene expressed during haematopoiesis

and conserved

[134915]between mouse and human. Nucleic Acids Res. 21:
1245–1249, 1993.

[134916]

[134917]

[134918]12232.

[134919]Crompton, M. R.; Bartlett, T. J.; MacGregor, A. D.; Manfio-
letti,

[134920]G.; Buratti, E.; Giancotti, V.; Goodwin, G. H.: Identification
of

[134921]a novel vertebrate homeobox gene expressed in
haematopoietic cells. Nucleic

[134922]Acids Res. 20: 5661–5667, 1992.

[134923]

[134924]

[134925]12233.

[134926]D'Elia, A. V.; Tell, G.; Russo, D.; Arturi, F.; Puglisi, F.; Man-
fioletti,

[134927]G.; Gattei, V.; Mack, D. L.; Cataldi, P.; Filetti, S.; Di Loreto,

[134928]C.; Damante, G.: Expression and localization of the home-

odomain-containing

[134929]protein HEX in human thyroid tumors. J. Clin. Endocr.

Metab. 87:

[134930]1376–1383, 2002.

[134931]

[134932]

[134933]12234.

[134934]Hromas, R.; Radich, J.; Collins, S.: PCR cloning of an orphan

[134935]homeobox gene (PRH) preferentially expressed in myeloid and liver

[134936]cells. Biochem. Biophys. Res. Commun. 195: 976–983, 1993.

[134937]

[134938]

[134939]12235.

[134940]Morgutti, M.; Demori, E.; Pecile, V.; Amoroso, A.; Rustighi, A.;

[134941]Manfioletti, G.: Genomic organization and chromosome mapping of the

[134942]human homeobox gene HHEX. Cytogenet. Cell Genet. 94:

30–32, 2001.

[134943]

[134944]

[134945]12236.

[134946]Tanaka, T.; Inazu, T.; Yamada, K.; Myint, Z.; Keng, V. W;

Inoue,

[134947]Y.; Taniguchi, N.; Noguchi, T.: cDNA cloning and expression of rat

[134948]homeobox gene, Hex, and functional characterization of the protein. Biochem.

[134949]J. 339: 111–117, 1999.

[134950]

[134951]

[134952]12237.

[134953]Miles, M. F.; Barhite, S.; Sganga, M.; Elliott, M.: Phosducin-like

[134954]protein: an ethanol-responsive potential modulator of guanine nucleotide-binding

[134955]protein function. Proc. Nat. Acad. Sci. 90: 10831–10835, 1993.

[134956]

[134957]

[134958]12238.

[134959]Thibault, C.; Wang, J. F.; Charnas, R.; Mirel, D.; Barhite, S.;

[134960]Miles, M. F.: Cloning and characterization of the rat and
human phosducin-like

[134961]protein genes: structure, expression and chromosomal localization. Biochim.

[134962]Biophys. Acta 1444: 346–354, 1999.

[134963]

[134964]

[134965]12239.

[134966]Fusco, C.; Reymond, A.; Zervos, A. S.: Molecular cloning
and characterization

[134967]of a novel retinoblastoma-binding protein. Genomics 51:
351–358,

[134968]1998.

[134969]

[134970]

[134971]12240.

[134972]Schaeper, U.; Subramanian, T.; Lim, L.; Boyd, J. M.; Chin-
nadurai,

[134973]G.: Interaction between a cellular protein that binds to the
C-terminal

[134974]region of adenovirus E1A (CtBP) and a novel cellular pro-
tein is disrupted

[134975]by E1A through a conserved PLDLS motif. J. Biol. Chem.
273: 8549–8552,

[134976]1998.

[134977]

[134978]

[134979]12241.

[134980]Wong, A. K. C.; Ormonde, P. A.; Pero, R.; Chen, Y.; Lian, L.;
Salada,

[134981]G.; Berry, S.; Lawrence, Q.; Dayananth, P.; Ha, P.; Tavti-
gian, S.

[134982]V.; Teng, D. H.-F.; Bartel, P. L.: Characterization of a car-
boxy-terminal

[134983]BRCA1 interacting protein. Oncogene 17: 2279–2285,
1998.

[134984]

[134985]

[134986]12242.

[134987]Yu, X.; Baer, R.: Nuclear localization and cell cycle-specific

[134988]expression of CtIP, a protein that associates with the
BRCA1 tumor

[134989]suppressor. J. Biol. Chem. 275: 18541–18549, 2000.

[134990]

[134991]

[134992]12243.

[134993]Yu, X.; Wu, L. C.; Bowcock, A. M.; Aronheim, A.; Baer, R.:

The

[134994]C-terminal (BRCT) domains of BRCA1 interact in vivo with
CtIP, a protein

[134995]implicated in the CtBP pathway of transcriptional repression. J.

[134996]Biol. Chem. 273: 25388–25392, 1998.

[134997]

[134998]

[134999]12244.

[135000]Pope, R. K.; Pestonjamas, K. N.; Smith, K. P.; Wulfkuhle, J. D.;

[135001]Strassel, C. P.; Lawrence, J. B.; Luna, E. J.: Cloning, characterization,

[135002]and chromosomal localization of human supervillin (SVIL).

Genomics 52:

[135003]342–351, 1998.

[135004]

[135005]

[135006]12245.

[135007]Ting, H.-J.; Yeh, S.; Nishimura, K.; Chang, C.: Supervillin associates

[135008]with androgen receptor and modulates its transcriptional activity. Proc.

[135009]Nat. Acad. Sci. 99: 661–666, 2002.

[135010]

[135011]

[135012]12246.

[135013]Bianchi, V.; Robles, R.; Alberio, L.; Furlan, M.; Lammle, B.:

[135014]Von Willebrand factor–cleaving protease (ADAMTS13) in

thrombocytopenic

[135015]disorders: a severely deficient activity is specific for
thrombotic

[135016]thrombocytopenic purpura. Blood 100: 710–713, 2002.

[135017]

[135018]

[135019]12247.

[135020]Fujikawa, K.; Suzuki, H.; McMullen, B.; Chung, D.: Purifica-
tion

[135021]of human von Willebrand factor–cleaving protease and its
identification

[135022]as a new member of the metalloproteinase family. Blood
98: 1662–1666,

[135023]2001.

[135024]

[135025]

[135026]12248.

[135027]Furlan, M.; Robles, R.; Solenthaler, M.; Wassmer, M.; San-
doz, Pl;

[135028]Lammle, B.: Deficient activity of von Willebrand factor–

cleaving

[135029]protease in chronic relapsing thrombotic thrombocy-
topenic purpura. Blood 89:

[135030]3097-3103, 1997.

[135031]

[135032]

[135033]12249.

[135034]Gerritsen, H. E.; Robles, R.; Lammle, B.; Furlan, M.: Partial

[135035]amino acid sequence of purified von Willebrand factor-
cleaving protease. Blood 98:

[135036]1654-1661, 2001.

[135037]

[135038]

[135039]12250.

[135040]Kokame, K.; Matsumoto, M.; Soejima, K.; Yagi, H.; Ishiza-
shi, H.;

[135041]Funato, M.; Tamai, H.; Konno, M.; Kamide, K.; Kawano, Y.;
Miyata,

[135042]T.; Fujimura, Y.: Mutations and common polymorphisms
in ADAMTS13

[135043]gene responsible for von Willebrand factor–cleaving protease activity. Proc.

[135044]Nat. Acad. Sci. 99: 11902–11907, 2002.

[135045]

[135046]

[135047]12251.

[135048]Re, F.; Strominger, J. L.: Toll–like receptor 2 (TLR2) and TLR4

[135049]differentially activate human dendritic cells. J. Biol. Chem. 276:

[135050]37692–37699, 2001.

[135051]

[135052]

[135053]12252.

[135054]Thoma–Uszynski, S.; Stenger, S.; Takeuchi, O.; Ochoa, M. T.; Engele,

[135055]M.; Sieling, P. A.; Barnes, P. F.; Rollinghoff, M.; Bolcskei, P. L.;

[135056]Wagner, M.; Akira, S.; Norgard, M. V.; Belisle, J. T.; Godowski, P.

[135057]J.; Bloom, B. R.; Modlin, R. L.: Induction of direct antimicrobial

[135058]activity through mammalian Toll-like receptors. Science 291: 1544–1547,

[135059]2001.

[135060]

[135061]

[135062]12253.

[135063]Underhill, D. M.; Ozinsky, A.; Hajjar, A. M.; Stevens, A.; Wilson,

[135064]C. B.; Bassetti, M.; Aderem, A.: The Toll-like receptor 2 is recruited

[135065]to macrophage phagosomes and discriminates between pathogens. Nature 401:

[135066]811–815, 1999.

[135067]

[135068]

[135069]12254.

[135070]Wolfs, T. G. A. M.; Buurman, W. A.; van Schadewijk, A.; de Vries,

[135071]B.; Daemen, M. A. R. C.; Hiemstra, P. S.; van't Veer, C.: In vivo

[135072]expression of Toll-like receptor 2 and 4 by renal epithelial cells:

[135073]IFN-gamma and TNF-alpha mediated up-regulation during inflammation. J.

[135074]Immun. 168: 1286-1293, 2002.

[135075]

[135076]

[135077]12255.

[135078]Alexopoulou, L.; Holt, A. C.; Medzhitov, R.; and Flavell, R. A.

[135079]: Recognition of double stranded RNA and activation of NFkB by Toll-like

[135080]receptor 3. Nature 413: 732-738, 2001.

[135081]

[135082]

[135083]12256.

[135084]Advani, R. J.; Bae, H.-R.; Bock, J. B.; Chao, D. S.; Doung, Y.-C.;

[135085]Prekeris, R.; Yoo, J.-S.; Scheller, R. H.: Seven novel mammalian

[135086]SNARE proteins localize to distinct membrane compartments. J. Biol.

[135087]Chem. 273: 10317–10324, 1998.

[135088]

[135089]

[135090]12257.

[135091]Bock, J. B.; Scheller, R. H.: A fusion of new ideas. Nature 387:

[135092]133–135, 1997.

[135093]

[135094]

[135095]12258.

[135096]Bui, T. D.; Wong, S. H.; Lu, L.; Hong, W.: Endobrevin maps to

[135097]chromosome 2 in human and chromosome 6 in mouse.

Genomics 54: 579–580,

[135098]1998.

[135099]

[135100]

[135101]12259.

[135102]Wong, S. H.; Zhang, T.; Xu, Y.; Subramaniam, V. N.; Griffiths,

[135103]G.; Hong, W.: Endobrevin, a novel synaptobrevin/VAMP-like protein

[135104]preferentially associated with the early endosome. *Molec. Biol. Cell* 9:

[135105]1549–1563, 1998.

[135106]

[135107]

[135108]12260.

[135109]Hewett–Emmett, D.; Tashian, R. E.: Functional diversity, conservation,

[135110]and convergence in the evolution of the alpha-, beta- and gamma- carbonic

[135111]anhydrase gene families. *Molec. Phylogenet. Evol.* 5: 50–77, 1996.

[135112]

[135113]

[135114]12261.

[135115]Nakagawa, Y.; Uemura, H.; Hirao, Y.; Yoshida, K.; Saga, S.;
Yoshikawa,

[135116]K.: Radiation hybrid mapping of the human MN/CA9 locus
to chromosome

[135117]band 9p12–p13. Genomics 53: 118–119, 1998.

[135118]

[135119]

[135120]12262.

[135121]Opavsky, R.; Pastorekova, S.; Zelnik, V.; Gibadulinova, A.;
Stanbridge,

[135122]E. J.; Zavada, J.; Kettmann, R.; Pastorek, J.: Human MN/
CA9 gene,

[135123]a novel member of the carbonic anhydrase family: struc-
ture and exon

[135124]to protein domain relationships. Genomics 33: 480–487,
1996.

[135125]

[135126]

[135127]12263.

[135128]Soilleux, E. J.; Barten, R.; Trowsdale, J.: Cutting edge: DC-SIGN;

[135129]a related gene, DC-SIGNR; and CD23 form a cluster on 19p13. J. Immun. 165:

[135130]2937-2942, 2000.

[135131]

[135132]

[135133]12264.

[135134]Arnold, G. L.; Bixler, D.; Girod, D.: Probable autosomal recessive

[135135]inheritance of polysplenia, situs inversus and cardiac defects in

[135136]an Amish family. Am. J. Med. Genet. 16: 35-42, 1983.

[135137]

[135138]

[135139]12265.

[135140]Brueckner, M.; D'Eustachio, P.; Horwich, A. L.: Linkage mapping

[135141]of a mouse gene, iv, that controls left-right asymmetry of the heart

[135142]and viscera. Proc. Nat. Acad. Sci. 86: 5035–5038, 1989.

[135143]

[135144]

[135145]12266.

[135146]Hanzlik, A. J.; Binder, M.; Layton, W. M.; Rowe, L.; Layton, M.;

[135147]Taylor, B. A.; Osemlak, M. M.; Richards, J. E.; Kurnit, D. M.; Stewart,

[135148]G. D.: The murine situs inversus viscerum (iv) gene responsible

[135149]for visceral asymmetry is linked tightly to the Igh–C cluster on chromosome

[135150]12. Genomics 7: 389–393, 1990.

[135151]

[135152]

[135153]12267.

[135154]McGrath, J.; Horwich, A. L.; Brueckner, M.: Duplication/deficiency

[135155]mapping of situs inversus viscerum (iv), a gene that determines left–right

[135156]asymmetry in the mouse. Genomics 14: 643–648, 1992.

[135157]

[135158]

[135159]12268.

[135160]Okada, Y.; et al.; et al: Abnormal nodal flow precedes situs
inversus

[135161]in iv and inv mice. Molec. Cell 4: 459–468, 1999.

[135162]

[135163]

[135164]12269.

[135165]Supp, D.; et al.; et al: Targeted deletion of the ATP binding

[135166]domain of left–right dynein confirms its role in specifying
development

[135167]of left–right asymmetries. Development 126: 5495–5504,
1999.

[135168]

[135169]

[135170]12270.

[135171]Supp, D. M.; Witte, D. P.; Potter, S. S.; Brueckner, M.: Mu–

tation

[135172]of an axonemal dynein affects left–right asymmetry in in–
versus viscerum

[135173]mice. Nature 389: 963–966, 1997.

[135174]

[135175]

[135176]12271.

[135177]Matoskova, B.; Wong, W. T.; Nomura, N.; Robbins, K. C.; Di
Fiore,

[135178]P. P.: RN–tre specifically binds to the SH3 domain of eps8
with high

[135179]affinity and confers growth advantage to NIH3T3 upon
carboxy–terminal

[135180]truncation. Oncogene 12: 2679–2688, 1996.

[135181]

[135182]

[135183]12272.

[135184]Matoskova, B.; Wong, W. T.; Seki, N.; Nagase, T.; Nomura,
N.; Robbins,

[135185]K. C.; Di Fiore, P. P.: RN–tre identifies a family of tre–

related

[135186]proteins displaying a novel potential protein binding domain. Oncogene 12:

[135187]2563–2571, 1996.

[135188]

[135189]

[135190]12273.

[135191]Lee, M. P.; Brandenburg, S.; Landes, G. M.; Adams, M.; Miller,

[135192]G.; Feinberg, A. P.: Two novel genes in the center of the 11p15 imprinted

[135193]domain escape genomic imprinting. Hum. Molec. Genet. 8: 683–690,

[135194]1999.

[135195]

[135196]

[135197]12274.

[135198]Nakamura, M.; Masuda, H.; Horii, J.; Kuma, K.; Yokoyama, N.; Ohba,

[135199]T.; Nishitani, H.; Miyata, T.; Tanaka, M.; Nishimoto, T.:

When overexpressed,

[135200]a novel centrosomal protein, RanBPM, causes ectopic microtubule nucleation

[135201]similar to gamma-tubulin. J. Cell Biol. 143: 1041–1052, 1998.

[135202]

[135203]

[135204]12275.

[135205]Nishitani, H.; Hirose, E.; Uchimura, Y.; Nakamura, M.; Umeda,

[135206]M.; Nishii, K.; Mori, N.; Nishimoto, T.: Full-sized RanBPM cDNA encodes

[135207]a protein possessing a long stretch of proline and glutamine within

[135208]the N-terminal region, comprising a large protein complex. Gene 272:

[135209]25–33, 2001.

[135210]

[135211]

[135212]12276.

[135213]Miyamoto, T.; Kanazawa, N.; Kato, S.; Kawakami, M.; In-

oue, Y.;

[135214]Kuhara, T.; Inoue, T.; Takeshita, K.; Tsujino, S.: Diagnosis
of Japanese

[135215]patients with HHH syndrome by molecular genetic analysis: a common

[135216]mutation, R179X. J. Hum. Genet. 46: 260–262, 2001.

[135217]

[135218]

[135219]12277.

[135220]Nakajima, M.; Ishii, S.; Mito, T.; Takeshita, K.; Takashima,
S.;

[135221]Takakura, H.; Inoue, I.; Saheki, T.; Akiyoshi, H.; Ichihara,
K.:

[135222]Clinical, biochemical and ultrastructural study on the
pathogenesis

[135223]of hyperornithinemia–hyperammonemia–homocitrullinuria
syndrome. Brain

[135224]Dev. 10: 181–185, 1988.

[135225]

[135226]

[135227]12278.

[135228]Tsujino, S.; Kanazawa, N.; Ohashi, T.; Eto, Y.; Saito, T.;
Kira,

[135229]J.; Yamada, T.: Three novel mutations (G27E, insAAC,
R179X) in the

[135230]ORNT1 gene of Japanese patients with hyperornithinemia,
hyerammonemia,

[135231]and homocitrullinuria syndrome. Ann. Neurol. 47:
625–631, 2000.

[135232]

[135233]

[135234]12279.

[135235]Grand, R. J. A.; Milner, A. E.; Mustoe, T.; Johnson, G. D.;
Owen,

[135236]D.; Grant, M. L.; Gregory, C. D.: A novel protein expressed
in mammalian

[135237]cells undergoing apoptosis. Exp. Cell Res. 218: 439–451,
1995.

[135238]

[135239]

[135240]12280.

[135241]Twist, C. J.; Beier, D. R.; Disteché, C. M.; Edelhoff, S.; Ted–

der,

[135242]T. F.: The mouse Cd83 gene: structure, domain organization, and chromosome

[135243]localization. Immunogenetics 48: 383–393, 1998.

[135244]

[135245]

[135246]12281.

[135247]Lee, S. B.; Rao, A. K.; Lee, K.-H.; Yang, X.; Bae, Y. S.; Rhee,

[135248]S. G.: Decreased expression of phospholipase C- β 2 isozyme in human

[135249]platelets with impaired function. Blood 88: 1684–1691, 1996.

[135250]

[135251]

[135252]12282.

[135253]Mao, G. F.; Vaidyula, V. R.; Kunapuli, S. P.; Rao, A. K.: Lineage-specific

[135254]defect in gene expression in human platelet phospholipase C- β 2

[135255]deficiency. Blood 99: 905–911, 2002.

[135256]

[135257]

[135258]12283.

[135259]Park, D.; Jhon, D.-Y.; Kriz, R.; Knopf, J.; Rhee, S. G.:

Cloning,

[135260]sequencing, expression, and Gq-independent activation
of phospholipase

[135261]C-beta-2. J. Biol. Chem. 267: 16048-16055, 1992.

[135262]

[135263]

[135264]12284.

[135265]Park, S. H.; Ryu, S. H.; Suh, P. G.; Kim, H.: Assignment of
human

[135266]PLCB2 encoding PLC-beta-2 to human chromosome
15q15 by fluorescence

[135267]in situ hybridization. Cytogenet. Cell Genet. 83: 48-49,
1998.

[135268]

[135269]

[135270]12285.

[135271]Rao, A. K.; Kowalska, M. A.; Disa, J.: Impaired cytoplasmic
ionized

[135272]calcium mobilization in inherited platelet secretion de-
fects. Blood 74:

[135273]664–672, 1989.

[135274]

[135275]

[135276]12286.

[135277]Yang, X.; Sun, L.; Ghosh, S.; Rao, A. K.: Human platelet
signaling

[135278]defect characterized by impaired production of 1,4,5 in-
ositol triphosphate

[135279]and phosphatic acid, and diminished pleckstrin phospho-
rylation. Evidence

[135280]for defective phospholipase C activation. Blood 88:
1676–1683, 1996.

[135281]

[135282]

[135283]12287.

[135284]Gillen, C. M.; Brill, S.; Payne, J. A.; Forbush, B., III: Molecu-
lar

[135285]cloning and functional expression of the K-Cl cotransporter from rabbit,

[135286]rat, and human: a new member of the cation-chloride cotransporter

[135287]family. J. Biol. Chem. 271: 16237-16244, 1996.

[135288]

[135289]

[135290]12288.

[135291]Larsen, F.; Solheim, J.; Kristensen, T.; Kolsto, A. B.; Prydz,

[135292]H.: A tight cluster of five unrelated human genes on chromosome 16q22.1. Hum.

[135293]Molec. Genet. 2: 1589-1595, 1993.

[135294]

[135295]

[135296]12289.

[135297]Pellegrino, C. M.; Rybicki, A. C.; Musto, S.; Nagel, R. L.; Schwartz,

[135298]R. S.: Molecular identification and expression of erythroid K:Cl

[135299]cotransporter in human and mouse erythroleukemic cells.

Blood Cells

[135300]Molec. Dis. 24: 31–40, 1998.

[135301]

[135302]

[135303]12290.

[135304]Hammond, E. M.; Brunet, C. L.; Johnson, G. D.; Parkhill, J.;
Milner,

[135305]A. E.; Brady, G.; Gregory, C. D.; Grand, R. J. A.: Homology
between

[135306]a human apoptosis specific protein and the product of
APG5, a gene

[135307]involved in autophagy in yeast. FEBS Lett. 425: 391–395,
1998.

[135308]

[135309]

[135310]12291.

[135311]Schmeiser, K.; Armstrong, S.; Hammond, E. M.; Grand, R.
J. A.:

[135312]Assignment of the yeast APG5 human homologue APG5L
to chromosome band

[135313]6q21 by fluorescence in situ hybridisation. Cytogenet. Cell

Genet. 87:

[135314]213–214, 1999.

[135315]

[135316]

[135317]12292.

[135318]Nakayama, M.; Nakajima, D.; Nagase, T.; Nomura, N.; Seki, N.; Ohara,

[135319]O.: Identification of high-molecular-weight proteins with multiple

[135320]EGF-like motifs by motif-trap screening. Genomics 51: 27–34, 1998.

[135321]

[135322]

[135323]12293.

[135324]Wu, Q.; Maniatis, T.: Large exons encoding multiple ectodomains

[135325]are a characteristic feature of protocadherin genes. Proc. Nat. Acad.

[135326]Sci. 97: 3124–3129, 2000.

[135327]

[135328]

[135329]12294.

[135330]Becker, R. M.; Darrow, C.; Zimonjic, D. B.; Popescu, N. C.;
Watson,

[135331]M. A.; Fleming, T. P.: Identification of mammaglobin B, a
novel member

[135332]of the uteroglobin gene family. Genomics 54: 70–78,
1998.

[135333]

[135334]

[135335]12295.

[135336]Molloy, M. P.; Bolis, S.; Herbert, B. R.; Ou, K.; Tyler, M. I.;

[135337]van Dyk, D. D.; Willcox, M. D.; Gooley, A. A.; Williams, K.
L.; Morris,

[135338]C. A.; Walsh, B. J.: Establishment of the human reflex tear
two-dimensional

[135339]polyacrylamide gel electrophoresis reference map: new
proteins of

[135340]potential diagnostic value. Electrophoresis 18:
2811–2815, 1997.

[135341]

[135342]

[135343]12296.

[135344]Paine–Saunders, S.; Viviano, B. L.; Saunders, S.: GPC6, a novel

[135345]member of the glypican gene family, encodes a product structurally

[135346]related to GPC4 and is colocalized with GPC5 on human chromosome 13. Genomics 57:

[135347]455–458, 1999.

[135348]

[135349]

[135350]12297.

[135351]Veugelers, M.; De Cat, B.; Ceulemans, H.; Bruystens, A. M.; Coomans,

[135352]C.; Durr, J.; Vermeesch, J.; Marynen, P.; David, G.: Glypican–6,

[135353]a new member of the glypican family of cell surface heparan sulfate

[135354]proteoglycans. J. Biol. Chem. 274: 26968–26977, 1999.

[135355]

[135356]

[135357]12298.

[135358]Takei, Y.; Sasaki, S.; Fujiwara, T.; Takahashi, E.; Muto, T.;
Nakamura,

[135359]Y.: Molecular cloning of a novel gene similar to myeloid
antigen

[135360]CD33 and its specific expression in placenta. Cytogenet.
Cell Genet. 78:

[135361]295–300, 1997.

[135362]

[135363]

[135364]12299.

[135365]Jacobs, S.; Schurmann, A.; Becker, W.; Bockers, T. M.;
Copeland,

[135366]N. G.; Jenkins, N. A.; Joost, H.–G.: The mouse ADP–
ribosylation factor–like

[135367]4 gene: two separate promoters direct specific transcrip–
tion in tissues

[135368]and testicular germ cell. Biochem. J. 335: 259–265, 1998.

[135369]

[135370]

[135371]12300.

[135372]Bootcov, M. R.; Bauskin, A. R.; Valenzuela, S. M.; Moore, A. G.;

[135373]Bansal, M.; He, X. Y.; Zhang, H. P.; Donnellan, M.; Mahler, S.; Pryor,

[135374]K.; Walsh, B. J.; Nicholson, R. C.; Fairlie, W. D.; Por, S. B.; Robbins,

[135375]J. M.; Breit, S. N.: MIC-1, a novel macrophage inhibitory cytokine,

[135376]is a divergent member of the TGF-beta superfamily. Proc. Nat. Acad.

[135377]Sci. 94: 11514-11519, 1997.

[135378]

[135379]

[135380]12301.

[135381]Bottner, M.; Laaff, M.; Schechinger, B.; Rappold, G.; Unsicker,

[135382]K.; Suter-Crazzolara, C.: Characterization of the rat, mouse, and

[135383]human genes of growth/differentiation factor-

15/macrophage inhibiting

[135384]cytokine-1 (GDF-15/MIC-1). Gene 237: 105-111, 1999.

[135385]

[135386]

[135387]12302.

[135388]Brown, D. A.; Breit, S. N.; Buring, J.; Fairlie, W. D.; Bauskin,

[135389]A. R.; Liu, T.; Ridker, P. M.: Concentration in plasma of
macrophage

[135390]inhibitory cytokine-1 and risk of cardiovascular events in
women:

[135391]a nested case-control study. Lancet 359: 2159-2163,
2002.

[135392]

[135393]

[135394]12303.

[135395]Fairlie, W. D.; Russell, P. K.; Wu, W. M.; Moore, A. G.;
Zhang,

[135396]H.-P.; Brown, P. K.; Bauskin, A. R.; Breit, S. N.: Epitope
mapping

[135397]of the transforming growth factor-beta superfamily pro-

tein, macrophage

[135398]inhibitory cytokine-1 (MIC-1): identification of at least five distinct

[135399]epitope specificities. Biochemistry 40: 65-73, 2001.

[135400]

[135401]

[135402]12304.

[135403]Hromas, R.; Hufford, M.; Sutton, J.; Xu, D.; Li, Y.; Lu, L.:
PLAB,

[135404]a novel placental bone morphogenetic protein. Biochim.
Biophys. Acta 1354:

[135405]40-44, 1997.

[135406]

[135407]

[135408]12305.

[135409]Moore, A. G.; Brown, D. A.; Fairlie, W. D.; Bauskin, A. R.;
Brown,

[135410]P. K.; Munier, M. L. C.; Russell, P. K.; Salamonsen, L. A.;
Wallace,

[135411]E. M.; Breit, S. N.: The transforming growth factor-beta

superfamily

[135412]cytokine macrophage inhibitory cytokine-1 is present in
high concentrations

[135413]in the serum of pregnant women. J. Clin. Endocr. Metab.
85: 4781-4788,

[135414]2000.

[135415]

[135416]

[135417]12306.

[135418]Paralkar, V. M.; Vail, A. L.; Grasser, W. A.; Brown, T. A.; Xu,

[135419]H.; Vukicevic, S.; Ke, H. Z.; Qi, H.; Owen, T. A.; Thompson,
D. D.

[135420]: Cloning and characterization of a novel member of the
transforming

[135421]growth factor-beta/bone morphogenetic protein family. J.
Biol. Chem. 273:

[135422]13760-13767, 1998.

[135423]

[135424]

[135425]12307.

[135426]Yokoyama–Kobayashi, M.; Saeki, M.; Sekine, S.; Kato, S.:

Human

[135427]cDNA encoding a novel TGF–beta superfamily protein
highly expressed

[135428]in placenta. J. Biochem. 122: 622–626, 1997.

[135429]

[135430]

[135431]12308.

[135432]Conklin, D. C.; Rixon, M. W.; Kuestner, R. E.; Maurer, M. F.;
Whitmore,

[135433]T. E.; Millar, R. P.: Cloning and gene expression of a novel
human

[135434]ribonucleoprotein. Biochim. Biophys. Acta 1492: 465–469,
2000.

[135435]

[135436]

[135437]12309.

[135438]Kataoka, N.; Yong, J.; Kim, V. N.; Velazquez, F.; Perkinson,
R.

[135439]A.; Wang, F.; Dreyfuss, G.: Pre–mRNA splicing imprints
mRNA in the

[135440]nucleus with a novel RNA-binding protein that persists in the cytoplasm. Molec.

[135441]Cell 6: 673–682, 2000.

[135442]

[135443]

[135444]12310.

[135445]Salicioni, A. M.; Xi. M.; Vanderveer, L. A.; Balsara, B.; Testa,

[135446]J. R.; Dunbrack, R. L., Jr.; Godwin, A. K.: Identification and structural

[135447]analysis of human RBM8A and RBM8B: two highly conserved RNA-binding

[135448]motif proteins that interact with OVCA1, a candidate tumor suppressor. Genomics 69:

[135449]54–62, 2000.

[135450]

[135451]

[135452]12311.

[135453]Zhao, X.–F.; Nowak, N. J.; Shows, T. B.; Aplan, P. D.: MAGOH interacts

[135454]with a novel RNA-binding protein. Genomics 63:
145–148, 2000.

[135455]

[135456]

[135457]12312.

[135458]Fischle, W.; Emiliani, S.; Hendzel, M. J.; Nagase, T.; No-
mura,

[135459]N.; Voelter, W.; Verdin, E.: A new family of human histone
deacetylases

[135460]related to *Saccharomyces cerevisiae* HDA1p. J. Biol. Chem.
274: 11713–11720,

[135461]1999.

[135462]

[135463]

[135464]12313.

[135465]Pazin, M. J.; Kadonaga, J. T.: What's up and down with his-
tone

[135466]deacetylation and transcription? Cell 89: 325–328, 1997.

[135467]

[135468]

[135469]12314.

[135470]Zhou, L.-J.; Tedder, T. F.: Human blood dendritic cells selectively

[135471]express CD83, a member of the immunoglobulin superfamily. J. Immun. 154:

[135472]3821–3835, 1995.

[135473]

[135474]

[135475]12315.

[135476]Zhou, L. J.; Schwarting, R.; Smith, H. M.; Tedder, T. F.: A novel

[135477]cell–surface molecule expressed by human interdigitating reticulum

[135478]cells, Langerhans cells, and activated lymphocytes is a new member

[135479]of the Ig superfamily. J. Immun. 149: 735–742, 1992.

[135480]

[135481]

[135482]12316.

[135483]Bost–Usinger, L.; Chen, R. J.; Hillman, D.; Park, P.; Burn–

side,

[135484]B.: Multiple kinesin family members expressed in teleost
retina and

[135485]RPE include a novel C-terminal kinesin. Exp. Eye Res. 64:
781–794,

[135486]1997.

[135487]

[135488]

[135489]12317.

[135490]Hoang, E. H.; Whitehead, J. L.; Dose, A. C.; Burnside, B.:
Cloning

[135491]of a novel C-terminal kinesin (KIFC3) that maps to human
chromosome

[135492]16q13–q21 and thus is a candidate gene for Bardet–Biedl
syndrome. Genomics 52:

[135493]219–222, 1998.

[135494]

[135495]

[135496]12318.

[135497]Norman, D. A. M.; Barton, P. J. R.: Isolation, sequence, and

chromosomal

[135498]localisation of the human I(kappa)BR gene. Ann. Hum.

Genet. 64:

[135499]15–23, 2000.

[135500]

[135501]

[135502]12319.

[135503]Ray, P.; Zhang, D.–H.; Elias, J. A.; Ray, A.: Cloning of a differentially

[135504]expressed I–kappa–B–related protein. J. Biol. Chem. 270:
10680–10685,

[135505]1995.

[135506]

[135507]

[135508]12320.

[135509]Uchiumi, T.; Hinoshita, E.; Haga, S.; Nakamura, T.; Tanaka,
T.;

[135510]Toh, S.; Furukawa, M.; Kawabe, T.; Wada, M.; Kagotani, K.;
Okumura,

[135511]K.; Kohno, K.; Akiyama, S.; Kuwano, M.: Isolation of a

novel human

[135512]canalicular multispecific organic anion transporter,
cMOAT2/MRP3,

[135513]and its expression in cisplatin-resistant cancer cells with
decreased

[135514]ATP-dependent drug transport. Biochem. Biophys. Res.
Commun. 252:

[135515]103–110, 1998.

[135516]

[135517]

[135518]12321.

[135519]Bruhn, S. L.; Pil, P. M.; Essigmann, J. M.; Housman, D. E.;
Lippard,

[135520]S. J.: Isolation and characterization of human cDNA clones
encoding

[135521]a high mobility group box protein that recognizes struc-
tural distortions

[135522]to DNA caused by binding of the anticancer agent cis-
platin. Proc.

[135523]Nat. Acad. Sci. 89: 2307–2311, 1992.

[135524]

[135525]

[135526]12322.

[135527]Orphanides, G.; LeRoy, G.; Chang, C.-H.; Luse, D. S.; Reinberg,

[135528]D.: FACT, a factor that facilitates transcript elongation through

[135529]nucleosomes. Cell 92: 105–116, 1998.

[135530]

[135531]

[135532]12323.

[135533]Acker, J.; Wintzerith, M.; Vigneron, M.; Kedinger, C.: A
14.4

[135534]KDa acidic subunit of human RNA polymerase II with a putative leucine-zipper. DNA

[135535]Seq. 4: 329–331, 1994.

[135536]

[135537]

[135538]12324.

[135539]Pusch, C.; Wang, Z.; Roe, B.; Blin, N.: Genomic structure of

the

[135540]RNA polymerase II small subunit (hRPB14.4) locus (POLRF)
(sic) and

[135541]mapping to 22q13.1 by sequence identity. Genomics 34:
440–442, 1996.

[135542]

[135543]

[135544]12325.

[135545]Caira, F.; Antonson, P.; Pelto–Huikko, M.; Treuter, E.;
Gustafsson,

[135546]J.-A.: Cloning and characterization of RAP250, a novel nuclear receptor

[135547]coactivator. J. Biol. Chem. 275: 5308–5317, 2000.

[135548]

[135549]

[135550]12326.

[135551]Guan, X. Y.; Xu, J.; Anzick, S. L.; Zhang, H.; Trent, J. M.; Meltzer,

[135552]P. S.: Hybrid selection of transcribed sequences from microdissected

[135553]DNA: isolation of genes within amplified region at 20q11–q13.2 in

[135554]breast cancer. Cancer Res. 56: 3446–3450, 1996.

[135555]

[135556]

[135557]12327.

[135558]Ko, L.; Cardona, G. R.; Chin, W. W.: Thyroid hormone receptor-binding

[135559]protein, an LXXLL motif-containing protein, functions as a general

[135560]coactivator. Proc. Nat. Acad. Sci. 97: 6212–6217, 2000.

[135561]

[135562]

[135563]12328.

[135564]Lee, S.–K.; Anzick, S. L.; Choi, J.–E.; Bubendorf, L.; Guan,
X.–Y.;

[135565]Jung, Y.–K.; Kallioniemi, O. P.; Kononen, J.; Trent, J. M.;
Azorsa,

[135566]D.; Jhun, B.–H.; Cheong, J. H.; Lee, Y. C.; Meltzer, P. S.;
Lee, J.

[135567]W.: A nuclear factor, ASC–2, as a cancer–amplified tran-
scriptional

[135568]coactivator essential for ligand–dependent transactivation
by nuclear

[135569]receptors in vivo. J. Biol. Chem. 274: 34283–34293, 1999.

[135570]

[135571]

[135572]12329.

[135573]Mahajan, M. A.; Samuels, H. H: A new family of nuclear re-
ceptor

[135574]coregulators that integrate nuclear receptor signaling
through CREB-binding

[135575]protein. *Molec. Cell. Biol.* 20: 5048–5063, 2000.

[135576]

[135577]

[135578]12330.

[135579]Zhu, Y.; Kan, L.; Qi, C.; Kanwar, Y. S.; Yeldandi, A. V.; Rao,

[135580]M. S.; Reddy, J. K.: Isolation and characterization of per-
oxisome

[135581]proliferator-activated receptor (PPAR) interacting protein
(PRIP)

[135582]as a coactivator for PPAR. *J. Biol. Chem.* 275:
13510–13516, 2000.

[135583]

[135584]

[135585]12331.

[135586]Still, I. H.; Hamilton, M.; Vince, P.; Wolfman, A.; Cowell, J.

[135587]K.: Cloning of TACC1, an embryonically expressed, poten-
tially transforming

[135588]coiled coil containing gene, from the 8p11 breast cancer

amplicon. Oncogene 18:

[135589]4032–4038, 1999.

[135590]

[135591]

[135592]12332.

[135593]Still, I. H.; Vince, P.; Cowell, J. K.: The third member of the

[135594]transforming acidic coiled coil–containing gene family,

TACC3, maps

[135595]in 4p16, close to translocation breakpoints in multiple

myeloma, and

[135596]is upregulated in various cancer cell lines. Genomics 58:

165–170,

[135597]1999.

[135598]

[135599]

[135600]12333.

[135601]Burmester, T.; Weich, B.; Reinhardt, S.; Hankeln, T.: A ver-

tebrate

[135602]globin expressed in the brain. Nature 407: 520–523,

2000.

[135603]

[135604]

[135605]12334.

[135606]Moens, L.; Dewilde, S.: Globins in the brain. Nature 407:
461–462,

[135607]2000.

[135608]

[135609]

[135610]12335.

[135611]Scott, A. F.: Personal Communication. Baltimore, Md.
9/28/2000.

[135612]

[135613]

[135614]12336.

[135615]Vilches, C.; Gardiner, C. M.; Parham, P.: Gene structure
and promoter

[135616]variation of expressed and nonexpressed variants of the
KIR2DL5 gene. J.

[135617]Immun. 165: 6416–6421, 2000.

[135618]

[135619]

[135620]12337.

[135621]Vilches, C.; Rajalingam, R.; Uhrberg, M.; Gardiner, C. M.;
Young,

[135622]N. T.; Parham, P.: KIR2DL5, a novel killer-cell receptor
with a D0-D2

[135623]configuration of Ig-like domains. J. Immun. 164:
5797-5804, 2000.

[135624]

[135625]

[135626]12338.

[135627]Bates, E. E. M.; Fournier, N.; Garcia, E.; Valladeau, J.; Du-
rand,

[135628]I.; Pin, J.-J.; Zurawski, S. M.; Patel, S.; Abrams, J. S.;
Lebecque,

[135629]S.; Garrone, P.; Saeland, S.: APCs express DCIR, a novel C-
type lectin

[135630]surface receptor containing an immunoreceptor tyrosine-
based inhibitory

[135631]motif. J. Immun. 163: 1973-1983, 1999.

[135632]

[135633]

[135634]12339.

[135635]Huang, X.; Yuan, Z.; Chen, G.; Zhang, M.; Zhang, W.; Yu, Y.; Cao,

[135636]X.: Cloning and characterization of a novel ITIM containing lectin-like

[135637]immunoreceptor LLIR and its two transmembrane region deletion variants. Biochem.

[135638]Biophys. Res. Commun. 281: 131–140, 2001.

[135639]

[135640]

[135641]12340.

[135642]Seeger, M.; Kraft, R.; Ferrell, K.; Bech-Otschir, D.; Dumdey, R.;

[135643]Schade, R.; Gordon, C.; Naumann, M.; Dubiel, W.: A novel protein

[135644]complex involved in signal transduction possessing similarities to

[135645]26S proteasome subunits. FASEB J. 12: 469–478, 1998.

[135646]

[135647]

[135648]12341.

[135649]Ann, K.; Kowalchyk, J. A.; Loyet, K. M.; Martin, T. F. J.:
Novel

[135650]Ca(2+)-binding protein (CAPS) related to UNC-31 required
for Ca(2+)-activated

[135651]exocytosis. J. Biol. Chem. 272: 19637-19640, 1997.

[135652]

[135653]

[135654]12342.

[135655]Berwin, B.; Floor, E.; Martin, T. F. J.: CAPS (mammalian
UNC-31)

[135656]protein localizes to membranes involved in dense-core
vesicle exocytosis. Neuron 21:

[135657]137-145, 1998.

[135658]

[135659]

[135660]12343.

[135661]Hirosawa, M.; Nagase, T.; Ishikawa, K.; Kikuno, R.; Nomura, N.;

[135662]Ohara, O.: Characterization of cDNA clones selected by the GeneMark

[135663]analysis from size-fractionated cDNA libraries from human brain. DNA

[135664]Res. 6: 329–336, 1999.

[135665]

[135666]

[135667]12344.

[135668]Walent, J. H.; Porter, B. W.; Martin, T. F. J.: A novel 145 kd

[135669]brain cytosolic protein reconstitutes Ca^{2+} -regulated secretion in

[135670]permeable neuroendocrine cells. Cell 70: 765–775, 1992.

[135671]

[135672]

[135673]12345.

[135674]Veldhuisen, B.; Spruit, L.; Dauwerse, H. G.; Breuning, M. H.; Peters,

[135675]D. J.: Genes homologous to the autosomal dominant poly-

cystic kidney

[135676]disease genes (PKD1 and PKD2). *Europ. J. Hum. Genet.* 7:
860–872,

[135677]1999.

[135678]

[135679]

[135680]12346.

[135681]Hughes, J.; Ward, C. J.; Aspinwall, R.; Butler, R.; Harris, P.

[135682]C.: Identification of a human homologue of the sea urchin
receptor

[135683]for egg jelly: a polycystic kidney disease–like protein.
Hum. Molec.

[135684]Genet. 8: 543–549, 1999.

[135685]

[135686]

[135687]12347.

[135688]Curtis, B. M.; Scharnowske, S.; Watson, A. J.: Sequence and
expression

[135689]of a membrane–associated C–type lectin that exhibits
CD4–independent

[135690]binding of human immunodeficiency virus envelope glycoprotein gp120. Proc.

[135691]Nat. Acad. Sci. 89: 8356–8360, 1992.

[135692]

[135693]

[135694]12348.

[135695]Feinberg, H.; Mitchell, D. A.; Drickamer, K.; Weis, W. I.:
Structural

[135696]basis for selective recognition of oligosaccharides by DC-SIGN and

[135697]DC-SIGNR. Science 294: 2163–2166, 2001.

[135698]

[135699]

[135700]12349.

[135701]Geijtenbeek, T. B. H.; Kwon, D. S.; Torensma, R.; van Vliet, S.

[135702]J.; van Duijnhoven, G. C. F.; Middel, J.; Cornelissen, I. L. M. H.

[135703]A.; Nottet, H. S. L. M.; KewalRamani, V. N.; Littman, D. R.; Figdor,

[135704]C. G.; van Kooyk, Y.: DC-SIGN, a dendritic cell-specific
HIV-1-binding

[135705]protein that enhances trans-infection of T cells. Cell 100:
587-597,

[135706]2000.

[135707]

[135708]

[135709]12350.

[135710]Geijtenbeek, T. B. H.; Torensma, R.; van Vliet, S. J.; van
Duijnhoven,

[135711]G. C. F.; Adema, G. J.; van Kooyk, Y.; Figdor, C. G.: Identi-
fication

[135712]of DC-SIGN, a novel dendritic cell-specific ICAM-3 recep-
tor that supports

[135713]primary immune responses. Cell 100: 575-585, 2000.

[135714]

[135715]

[135716]12351.

[135717]Battini, J.-L.; Rasko, J. E. J.; Miller, A. D.: A human cell-
surface

[135718]receptor for xenotropic and polytropic murine leukemia
viruses: possible

[135719]role in G protein-coupled signal transduction. Proc. Nat.
Acad. Sci. 96:

[135720]1385-1390, 1999.

[135721]

[135722]

[135723]12352.

[135724]Levy, J. A.: Xenotropism: the elusive viral receptor finally
uncovered. Proc.

[135725]Nat. Acad. Sci. 96: 802-804, 1999.

[135726]

[135727]

[135728]12353.

[135729]Tailor, C. S.; Nouri, A.; Lee, C. G.; Kozak, C.; Kabat, D.:
Cloning

[135730]and characterization of a cell surface receptor for
xenotropic and

[135731]polytropic murine leukemia viruses. Proc. Nat. Acad. Sci.
96: 927-932,

[135732]1999.

[135733]

[135734]

[135735]12354.

[135736]Yang, Y.-L.; Guo, L.; Xu, S.; Holland, C. A.; Kitamura, T.;
Hunter,

[135737]K.; Cunningham, J. M.: Receptors for polytropic and
xenotropic mouse

[135738]leukaemia viruses encoded by a single gene at Rmc1. Na-
ture Genet. 21:

[135739]216-219, 1999.

[135740]

[135741]

[135742]12355.

[135743]Aksoy, S.; Raftogianis, R.; Weinshilboum, R.: Human his-
tamine

[135744]N-methyltransferase gene: structural characterization and
chromosomal

[135745]localization. Biochem. Biophys. Res. Commun. 219:
548-554, 1996.

[135746]

[135747]

[135748]12356.

[135749]Preuss, C. V.; Wood, T. C.; Szumlanski, C. L.; Raftogianis, R.

[135750]B.; Otterness, D. M.; Girard, B.; Scott, M. C.; Weinshilboum, R. M.

[135751]: Human histamine N-methyltransferase pharmacogenetics: common genetic

[135752]polymorphisms that alter activity. Molec. Pharm. 53: 708–717, 1998.

[135753]

[135754]

[135755]12357.

[135756]Price, R. A.; Scott, M. C.; Weinshilboum, R. M.: Genetic segregation

[135757]analysis of red blood cell (RBC) histamine N-methyltransferase (HNMT)

[135758]activity. Genet. Epidemiol. 10: 123–131, 1993.

[135759]

[135760]

[135761]12358.

[135762]Scott, M. C.; Van Loon, J. A.; Weinshilboum, R. M.: Pharmacogenetics

[135763]of N-methylation: heritability of human erythrocyte histamine N-methyltransferase

[135764]activity. Clin. Pharm. Ther. 43: 256-262, 1988.

[135765]

[135766]

[135767]12359.

[135768]Yamauchi, K.; Sekizawa, K.; Suzuki, H.; Nakazawa, H.; Ohkawara,

[135769]Y.; Katayose, D.; Ohtsu, H.; Tamura, G.; Shibahara, S.; Takemura,

[135770]M.; Maeyama, K.; Watanabe, T.; Sasaki, H.; Shirato, K.; Takishima,

[135771]T.: Structure and function of human histamine N-methyltransferase:

[135772]critical enzyme in histamine metabolism in airway. Am. J. Physiol. 267:

[135773]L342-L349, 1994.

[135774]

[135775]

[135776]12360.

[135777]Yan, L.; Galinsky, R. E.; Bernstein, J. A.; Liggett, S. B.;
Weinshilboum,

[135778]R. M.: Histamine N-methyltransferase pharmacogenetics:
association

[135779]of a common functional polymorphism with asthma. Phar-
macogenetics 10:

[135780]261-266, 2000.

[135781]

[135782]

[135783]12361.

[135784]Bartoloni, L.; Blouin, J.-L. C.; Sainsbury, A. J.; Gos, A.;
Morris,

[135785]M. A.; Affara, N. A.; DeLozier-Blanchet, C. D.; Antonarakis,
S. E.

[135786]: Assignment of the human dynein heavy chain gene
DNAH17L to human

[135787]chromosome 17p12 by in situ hybridization and radiation
hybrid mapping. Cytogenet.

[135788]Cell Genet. 84: 188–189, 1999.

[135789]

[135790]

[135791]12362.

[135792]Bartoloni, L.; Blouin, J.–L.; Maiti, A. K.; Sainsbury, A.;
Rossier,

[135793]C.; Gehrig, C.; She, J.–X.; Marron, M. P.; Lander, E. S.;
Meeks, M.;

[135794]Chung, E.; Armengot, M.; Jorissen, M.; Scott, H. S.; De–
lozier–Blanchet,

[135795]C. D.; Gardiner, R. M.; Antonarakis, S. E.: Axonemal beta
heavy chain

[135796]dynein DNAH9: cDNA sequence, genomic structure, and
investigation

[135797]of its role in primary ciliary dyskinesia. Genomics 72:
21–33, 2001.

[135798]

[135799]

[135800]12363.

[135801]Maiti, A. K.; Mattei, M.–G.; Jorissen, M.; Volz, A.; Zeigler,
A.;

[135802]Bouvagnet, P.: Identification, tissue specific expression,
and chromosomal

[135803]localisation of several human dynein heavy chain genes.
Europ. J.

[135804]Hum. Genet. 8: 923–932, 2000.

[135805]

[135806]

[135807]12364.

[135808]Milisav, I.; Jones, M. H.; Affara, N. A.: Characterization of

[135809]a novel human dynein–related gene that is specifically ex-
pressed in

[135810]testis. Mammalian Genome 7: 667–672, 1996.

[135811]

[135812]

[135813]12365.

[135814]Chapelin, C.; Duriez, N.; Magnino, F.; Goossens, M.; Es-
cudier,

[135815]E.; Amselem, S.: Isolation of several human axonemal
dynein heavy

[135816]chain genes: genomic structure of the catalytic site, phy-

logenetic

[135817]analysis and chromosomal assignment. FEBS Lett. 412:
325–330, 1997.

[135818]

[135819]

[135820]12366.

[135821]Hart, M. J.; Callow, M. G.; Souza, B.; Polakis, P.: IQGAP1, a

[135822]calmodulin-binding protein with a rasGAP-related do-
main, is a potential

[135823]effector for cdc42Hs. EMBO J. 15: 2997–3005, 1996.

[135824]

[135825]

[135826]12367.

[135827]Sugimoto, N.; Imoto, I.; Fukuda, Y.; Kurihara, N.; Kuroda,
S.;

[135828]Tanigami, A.; Kaibuchi, K.; Kamiyama, R.; Inazawa, J.: IQ-
GAP1, a

[135829]negative regulator of cell–cell adhesion, is upregulated by
gene amplification

[135830]at 15q26 in gastric cancer cell lines HSC39 and 40A. J.

Hum. Genet. 46:

[135831]21–25, 2001.

[135832]

[135833]

[135834]12368.

[135835]Weissbach, L.; Settleman, J.; Kalady, M. F.; Snijders, A. J.;
Murthy,

[135836]A. E.; Yan, Y.–X.; Bernards, A.: Identification of a human
RasGAP–related

[135837]protein containing calmodulin–binding motifs. J. Biol.
Chem. 269:

[135838]20517–20521, 1994.

[135839]

[135840]

[135841]12369.

[135842]Harrington, J. J.; Lieber, M. R.: Functional domains within
FEN–1

[135843]and RAD2 define a family of structure–specific endonucle–
ases: implications

[135844]for nucleotide excision repair. Genes Dev. 8: 1344–1355,

1994.

[135845]

[135846]

[135847]12370.

[135848]Kato, H.; Torigoe, T.: Radioimmunoassay for tumor anti-
gen of human

[135849]cervical squamous cell carcinoma. Cancer 40: 1621-1628,
1977.

[135850]

[135851]

[135852]12371.

[135853]Kuwano, A.; Kondo, I.; Kishi, F.; Suminami, Y.; Kato, H.:
Assignment

[135854]of the squamous cell carcinoma antigen locus (SCC) to
18q21 by in

[135855]situ hybridization. Genomics 30: 626, 1995.

[135856]

[135857]

[135858]12372.

[135859]Suminami, Y.; Kishi, F.; Sekiguchi, K.; Kato, H.: Squamous cell

[135860]carcinoma antigen is a new member of the serine protease inhibitors. Biochem.

[135861]Biophys. Res. Commun. 181: 51–58, 1991.

[135862]

[135863]

[135864]12373.

[135865]Barnes, R. C.; Worrall, D. M.: Identification of a novel human

[135866]serpin gene; cloning sequencing and expression of leupin. FEBS Lett. 373:

[135867]61–65, 1995.

[135868]

[135869]

[135870]12374.

[135871]Schick, C.; Kamachi, Y.; Bartuski, A. J.; Cataltepe, S.; Schechter,

[135872]N. M.; Pemberton, P. A.; Silverman, G. A.: Squamous cell carcinoma

[135873]antigen 2 is a novel serpin that inhibits the chymotrypsin-
like proteinases

[135874]cathepsin G and mast cell chymase. J. Biol. Chem. 272:
1849–1855,

[135875]1997.

[135876]

[135877]

[135878]12375.

[135879]Dahl, M. R.; Thiel, S.; Matsushita, M.; Fujita, T.; Willis, A.

[135880]C.; Christensen, T.; Vorup-Jensen, T.; Jensenius, J. C.:
MASP-3 and

[135881]its association with distinct complexes of the mannan-
binding lectin

[135882]complement activation pathway. Immunity 15: 127–135,
2001.

[135883]

[135884]

[135885]12376.

[135886]Endo, Y.; Sato, T.; Matsushita, M.; Fujita, T.: Exon struc-
ture

[135887]of the gene encoding the human mannose-binding protein-associated

[135888]serine protease light chain: comparison with complement C1r and C1s

[135889]genes. *Int. Immun.* 8: 1355–1358, 1996.

[135890]

[135891]

[135892]12377.

[135893]Matsushita, M.; Thiel, S.; Jensenius, J. C.; Terai, I.; Fujita,

[135894]T.: Proteolytic activities of two types of mannose-binding lectin-associated

[135895]serine protease. *J. Immun.* 165: 2637–2642, 2000.

[135896]

[135897]

[135898]12378.

[135899]Sato, T.; Endo, Y.; Matsushita, M.; Fujita, T.: Molecular characterization

[135900]of a novel serine protease involved in activation of the complement

[135901]system by mannose-binding protein. *Int. Immun.* 6:

665–669, 1994.

[135902]

[135903]

[135904]12379.

[135905]Takada, F.; Seki, N.; Matsuda, Y.; Takayama, Y.; Kawakami, M.:

[135906]Localization of the genes for the 100-kDa complement-activating components

[135907]of Ra-reactive factor (CRARF and Crarf) to human 3q27–q28 and mouse

[135908]16B2–B3. Genomics 25: 757–759, 1995.

[135909]

[135910]

[135911]12380.

[135912]Takada, F.; Takayama, Y.; Hatsuse, H.; Kawakami, M.: A new member

[135913]of the C1s family of complement proteins found in a bactericidal factor,

[135914]Ra-reactive factor, in human serum. Biochem. Biophys. Res. Commun. 196:

[135915]1003–1009, 1993.

[135916]

[135917]

[135918]12381.

[135919]Takayama, Y.; Takada, F.; Nowatari, M.; Kawakami, M.;
Matsu-ura,

[135920]N.: Gene structure of the P100 serine-protease component of the human

[135921]Ra-reactive factor. Molec. Immun. 36: 505-514, 1999.

[135922]

[135923]

[135924]12382.

[135925]Dessen, A.; Tang, J.; Schmidt, H.; Stahl, M.; Clark, J. D.;
Seehra,

[135926]J.; Somers, W. S.: Crystal structure of human cytosolic
phospholipase

[135927]A(2) reveals a novel topology and catalytic mechanism.
Cell 97:

[135928]349-360, 1999.

[135929]

[135930]

[135931]12383.

[135932]Hansen, L.; Urioste, S.; Petersen, H. V.; Jensen, J. N.;
Eiberg,

[135933]H.; Barbetti, F.; Serup, P.; Hansen, T.; Pedersen, O.: Mis-
sense mutations

[135934]in the human insulin promoter factor-1 gene and their re-
lation to

[135935]maturity-onset diabetes of the young and late-onset type
2 diabetes

[135936]mellitus in Caucasians. J. Clin. Endocr. Metab. 85:
1323-1326, 2000.

[135937]

[135938]

[135939]12384.

[135940]Johansson, K. A.; Grapin-Botton, A.: Development and
diseases

[135941]of the pancreas. Clin. Genet. 62: 14-23, 2002.

[135942]

[135943]

[135944]12385.

[135945]Jonsson, J.; Carlsson, L.; Edlund, T.; Edlund, H.: Insulin-promoter-factor

[135946]1 is required for pancreas development in mice. Nature 371: 606–609,

[135947]1994.

[135948]

[135949]

[135950]12386.

[135951]Leonard, J.; Peers, B.; Johnson, T.; Ferreri, I.; Lee, K.; Montminy,

[135952]M. R.: Characterization of somatostatin transactivating factor-1,

[135953]a novel homeobox factor that stimulates somatostatin expression in

[135954]pancreatic islet cells. Molec. Endocr. 7: 1275–1283, 1993.

[135955]

[135956]

[135957]12387.

[135958]Macfarlane, W. M.; Frayling, T. M.; Ellard, S.; Evans, J. C.; Allen,

[135959]L. I. S.; Bulman, M. P.; Ayres, S.; Shepherd, M.; Clark, P.;
Millward,

[135960]A.; Demaine, A.; Wilkin, T.; Docherty, K.; Hattersley, A. T.:
Missense

[135961]mutations in the insulin promoter factor-1 gene predis-
pose to type

[135962]2 diabetes. J. Clin. Invest. 104: R33-R39, 1999.

[135963]

[135964]

[135965]12388.

[135966]Miller, C. P.; McGehee, R. E., Jr.; Habener, J. F.: IDX-1: a

[135967]new homeodomain transcription factor expressed in rat
pancreatic islets

[135968]and duodenum that transactivates the somatostatin gene.
EMBO J. 13:

[135969]1145-1156, 1994.

[135970]

[135971]

[135972]12389.

[135973]Ohlsson, H.; Karlsson, K.; Edlund, T.: IPF1, a home-

odomain-containing

[135974]transactivator of the insulin gene. EMBO J. 12:
4251–4259, 1993.

[135975]

[135976]

[135977]12390.

[135978]Sharma, S.; Jhala, U. S.; Johnson, T.; Ferreri, K.; Leonard, J.;

[135979]Montminy, M.: Hormonal regulation of an islet-specific
enhancer in

[135980]the pancreatic homeobox gene STF-1. Molec. Cell. Biol.
17: 2598–2604,

[135981]1997.

[135982]

[135983]

[135984]12391.

[135985]Stoffel, M.; Stein, R.; Wright, C. V. E.; Espinosa, R., III; Le

[135986]Beau, M. M.; Bell, G. I.: Localization of human home-
odomain transcription

[135987]factor insulin promoter factor 1 (IPF1) to chromosome
band 13q12.1. Genomics 28:

[135988]125–126, 1995.

[135989]

[135990]

[135991]12392.

[135992]Stoffers, D. A.; Ferrer, J.; Clarke, W. L.; Habener, J. F.:

Early-onset

[135993]type-II diabetes mellitus (MODY4) linked to IPF1. (Letter)

Nature

[135994]Genet. 17: 138–141, 1997.

[135995]

[135996]

[135997]12393.

[135998]Stoffers, D. A.; Stanojevic, V.; Habener, J. F.: Insulin promoter

[135999]factor-1 gene mutation linked to early-onset type 2 diabetes mellitus

[136000]directs expression of a dominant negative isoprotein. J. Clin. Invest. 102:

[136001]232–241, 1998.

[136002]

[136003]

[136004]12394.

[136005]Watada, H.; Kajimoto, Y.; Kaneto, H.; Matsuoka, T.; Fujitani,

[136006]Y.; Miyazaki, J.; Yamasaki, Y.: Involvement of the homeodomain-containing

[136007]transcription factor PDX-1 in islet amyloid polypeptide gene transcription. Biochem.

[136008]Biophys. Res. Commun. 229: 746-751, 1996.

[136009]

[136010]

[136011]12395.

[136012]Wright, N. M.; Metzger, D. L.; Borowitz, S. M.; Clarke, W. L.

[136013]: Permanent neonatal diabetes mellitus and pancreatic exocrine insufficiency

[136014]resulting from congenital pancreatic agenesis. Am. J. Dis. Child. 147:

[136015]607-609, 1993.

[136016]

[136017]

[136018]12396.

[136019]Ashford, M. L. J.; Bond, C. T.; Blair, T. A.; Adelman, J. P.:

[136020]Cloning and functional expression of a rat heart KATP
channel. Nature 370:

[136021]456–459, 1994.

[136022]

[136023]

[136024]12397.

[136025]Bond, C. T.; Pessia, M.; Xia, X.–M.; Lagrutta, A.; Ka-
vannah, M.

[136026]P.; Adelman, J. P.: Cloning and expression of a family of
inward

[136027]rectifier potassium channels. Receptors Channels 2:
183–191, 1994.

[136028]

[136029]

[136030]12398.

[136031]Tucker, S. J.; James, M. R.; Adelman, J. P.: Assignment of
K(ATP)–1,

[136032]the cardiac ATP–sensitive potassium channel gene

(KCNJ5), to human

[136033]chromosome 11q24. Genomics 28: 127–128, 1995.

[136034]

[136035]

[136036]12399.

[136037]Wickman, K.; Seldin, M. F.; Gendler, S. J.; Clapham, D. E.:
Partial

[136038]structure, chromosome localization, and expression of the
mouse Girk4

[136039]gene. Genomics 40: 395–401, 1997.

[136040]

[136041]

[136042]12400.

[136043]Jenkins, N. A.: Personal Communication. Frederick, Md.
5/12/1997.

[136044]

[136045]

[136046]12401.

[136047]Khurana, T. S.; Engle, E. C.; Bennett, R. R.; Silverman, G.

A.;

[136048]Selig, S.; Bruns, G. A. P.; Kunkel, L. M.: (CA) repeat polymorphism

[136049]in the chromosome 18 encoded dystrophin-like protein. Hum. Molec.

[136050]Genet. 3: 841 only, 1994.

[136051]

[136052]

[136053]12402.

[136054]Eisenberg, I.; Avidan, N.; Potikha, T.; Hochner, H.; Chen, M.;

[136055]Olender, T.; Barash, M.; Shemesh, M.; Sadeh, M.; Grabov-Nardini, G.;

[136056]Shmlevich, I.; Friedmann, A.; Karpati, G.; Bradley, W. G.; Baumbach,

[136057]L.; Lancet, D.; Ben Asher, E.; Beckmann, J. S.; Argov, Z.; Mitrani-Rosenbaum,

[136058]S.: The UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase

[136059]gene is mutated in recessive hereditary inclusion body myopathy. Nature

[136060]Genet. 29: 83-87, 2001.

[136061]

[136062]

[136063]12403.

[136064]Metzinger, L.; Blake, D. J.; Squier, M. V.; Anderson, L. V. B.;

[136065]Deconinck, A. E.; Nawrotzki, R.; Hilton-Jones, D.; Davies,
K. E.:

[136066]Dystrobrevin deficiency at the sarcolemma of patients
with muscular

[136067]dystrophy. Hum. Molec. Genet. 6: 1185–1191, 1997.

[136068]

[136069]

[136070]12404.

[136071]Sadoulet-Puccio, H. M.; Feener, C. A.; Schaid, D. J.; Thi-
bodeau,

[136072]S. N.; Michels, V. V.; Kunkel, L. M.: The genomic organiza-
tion of

[136073]human dystrobrevin. Neurogenetics 1: 37–42, 1997.

[136074]

[136075]

[136076]12405.

[136077]Sadoulet–Puccio, H. M.; Khurana, T. S.; Cohen, J. B.;
Kunkel, L.

[136078]M.: Cloning and characterization of the human homologue
of a dystrophin

[136079]related phosphoprotein found at the Torpedo electric or-
gan post-synaptic

[136080]membrane. Hum. Molec. Genet. 5: 489–496, 1996.

[136081]

[136082]

[136083]12406.

[136084]Yoshida, M.; Hama, H.; Ishikawa–Sakurai, M.; Imamura, M.;
Mizuno,

[136085]Y.; Araishi, K.; Wakabayashi–Takai, E.; Noguchi, S.;
Sasaoka, T.;

[136086]Ozawa, E.: Biochemical evidence for association of dystro-
brevin with

[136087]the sarcoglycan–sarcospan complex as a basis for under-
standing sarcoglycanopathy. Hum.

[136088]Molec. Genet. 9: 1033–1040, 2000.

[136089]

[136090]

[136091]12407.

[136092]Bauer, W. R.; Hayes, J. J.; White, J. H.; Wolffe, A. P.: Nucleo-
some

[136093]structural changes due to acetylation. J. Molec. Biol. 236:
685-690,

[136094]1994.

[136095]

[136096]

[136097]12408.

[136098]Furukawa, Y.; Kawakami, T.; Sudo, K.; Inazawa, J.; Mat-
sumine, A.;

[136099]Akiyama, T.; Nakamura, Y.: Isolation and mapping of a
human gene

[136100](RPD3L1) that is homologous to RPD3, a transcription fac-
tor in *Saccharomyces*

[136101]*cerevisiae*. Cytogenet. Cell Genet. 73: 130-133, 1996.

[136102]

[136103]

[136104]12409.

[136105]Lee, D. Y.; Hayes, J. J.; Pruss, D.; Wolffe, A. P.: A positive
[136106]role for histone acetylation in transcription factor access
to nucleosomal
[136107]DNA. Cell 72: 73–84, 1993.

[136108]

[136109]

[136110]12410.

[136111]Luo, J.; Su, F.; Chen, D.; Shiloh, A.; Gu, W.: Deacetylation
of

[136112]p53 modulates its effect on cell growth and apoptosis.
Nature 408:

[136113]377–381, 2000.

[136114]

[136115]

[136116]12411.

[136117]Magnaghi–Jaulin, L.; Groisman, R.; Naguibneva, I.; Robin,
P.; Lorain,

[136118]S.; Le Villain, J. P.; Troalen, F.; Trouche, D.; Harel–Bellan,
A.

[136119]: Retinoblastoma protein represses transcription by re–

cruiting a histone

[136120]deacetylase. Nature 391: 601–605, 1998.

[136121]

[136122]

[136123]12412.

[136124]Taunton, J.; Hassig, C. A.; Schreiber, S. L.: A mammalian
histone

[136125]deacetylase related to the yeast transcriptional regulator
Rpd3p. Science 272:

[136126]408–411, 1996.

[136127]

[136128]

[136129]12413.

[136130]Vidal, M.; Gaber, R. F.: RPD3 encodes a second factor re-
quired

[136131]to achieve maximum positive and negative transcriptional
states in

[136132]*Saccharomyces cerevisiae*. Molec. Cell. Biol. 11:
6317–6327, 1991.

[136133]

[136134]

[136135]12414.

[136136]Vidal, M.; Strich, R.; Esposito, R. E.; Gaber, R. F.: RPD1
(SIN3/UME4)

[136137]is required for maximal activation and repression of di-
verse yeast

[136138]genes. Molec. Cell. Biol. 11: 6306–6316, 1991.

[136139]

[136140]

[136141]12415.

[136142]Wade, P. A.: Transcriptional control at regulatory check-
points

[136143]by histone deacetylases: molecular connections between
cancer and

[136144]chromatin. Hum. Molec. Genet. 10: 693–698, 2001.

[136145]

[136146]

[136147]12416.

[136148]Wolffe, A. P.: Histone deacetylase: a regulator of tran-

scription. Science 272:

[136149]371–372, 1996.

[136150]

[136151]

[136152]12417.

[136153]Zhang, Y.; Sun, Z.–W.; Iratni, R.; Erdjument–Bromage, H.;
Tempst,

[136154]P.; Hampsey, M.; Reinberg, D.: SAP30, a novel protein
conserved between

[136155]human and yeast, is a component of a histone deacetylase
complex. Molec.

[136156]Cell 1: 1021–1031, 1998.

[136157]

[136158]

[136159]12418.

[136160]Chappell, T. G.; Welch, W. J.; Schlossman, D. M.; Palter, K.
B.;

[136161]Schlesinger, M. J.; Rothman, J. E.: Uncoating ATPase is a
member

[136162]of the 70 kilodalton family of stress proteins. Cell 45:

3-13, 1986.

[136163]

[136164]

[136165]12419.

[136166]Dworniczak, B.; Mirault, M.-E.: Structure and expression
of a

[136167]human gene coding for a 71 kd heat shock 'cognate' pro-
tein. Nucleic

[136168]Acids Res. 15: 5181-5197, 1987.

[136169]

[136170]

[136171]12420.

[136172]Tavaria, M.; Gabriele, T.; Anderson, R. L.; Mirault, M.-E.;
Baker,

[136173]E.; Sutherland, G.; Kola, I.: Localization of the gene encod-
ing the

[136174]human heat shock cognate protein, HSP73, to chromo-
some 11. Genomics 29:

[136175]266-268, 1995.

[136176]

[136177]

[136178]12421.

[136179]Ungewickell, E.: The 70-kd mammalian heat shock proteins are structurally

[136180]and functionally related to the uncoating protein that releases clathrin

[136181]triskelia from coated vesicles. EMBO J. 4: 3385–3391, 1985.

[136182]

[136183]

[136184]12422.

[136185]Camoretti-Mercado, B.; Forsythe, S. M.; LeBeau, M. M.; Espinosa,

[136186]R., III; Vieira, J. E.; Halayko, A. J.; Willadsen, S.; Kurtz, B.;

[136187]Ober, C.; Evans, G. A.; Thweatt, R.; Shapiro, S.; Niu, Q.; Qin, Y.;

[136188]Padrid, P. A.; Solway, J.: Expression and cytogenetic localization

[136189]of the human SM22 gene (TAGLN). Genomics 49: 452–457, 1998.

[136190]

[136191]

[136192]12423.

[136193]Lawson, D.; Harrison, M.; Shapland, C.: Fibroblast trans-
gelin

[136194]and smooth muscle SM22-alpha are the same protein, the
expression

[136195]of which is down-regulated in many cell lines. Cell Motil.
Cytoskeleton 38:

[136196]250-257, 1997.

[136197]

[136198]

[136199]12424.

[136200]Lees-Miller, J. P.; Heeley, D. H.; Smillie, L. B.; Kay, C. M.:

[136201]Isolation and characterization of an abundant and novel
22-kDa protein

[136202](SM22) from chicken gizzard smooth muscle. J. Biol.
Chem. 262: 2988-2993,

[136203]1987.

[136204]

[136205]

[136206]12425.

[136207]Nishida, W.; Kitami, Y.; Abe, M.; Hiwada, K.: Gene cloning
and

[136208]nucleotide sequence of SM22-alpha from the chicken gizzard smooth

[136209]muscle. Biochem. Int. 23: 663-668, 1991.

[136210]

[136211]

[136212]12426.

[136213]Shapland, C.; Hsuan, J. J.; Totty, N. F.; Lawson, D.: Purification
cation

[136214]and properties of transgelin: a transformation and shape
change sensitive

[136215]actin-gelling protein. J. Cell Biol. 121: 1065-1073, 1993.

[136216]

[136217]

[136218]12427.

[136219]Shapland, C.; Lowings, P.; Lawson, D.: Identification of
new actin-associated

[136220]polypeptides that are modified by viral transformation and

changes

[136221]in cell shape. J. Cell Biol. 107: 153–161, 1988.

[136222]

[136223]

[136224]12428.

[136225]Solway, J.; Seltzer, J.; Samaha, F. F.; Kim, S.; Alger, L. E.;

[136226]Niu, Q.; Morrissey, E. E.; Ip, H. S.; Parmacek, M. S.: Struc-
ture and

[136227]expression of a smooth muscle cell-specific gene,
SM22-alpha. J.

[136228]Biol. Chem. 270: 13460–13469, 1995.

[136229]

[136230]

[136231]12429.

[136232]Stanier, P.; Abu-Hayyeh, S.; Murdoch, J. N.; Eddleston, J.;
Copp,

[136233]A. J.: Paralogous Sm22-alpha (Tagln) genes map to mouse
chromosomes

[136234]1 and 9: further evidence for a paralogous relationship.

Genomics 51:

[136235]144–147, 1998.

[136236]

[136237]

[136238]12430.

[136239]Thweatt, R.; Lumpkin, C. K., Jr.; Goldstein, S.: A novel gene

[136240]encoding a smooth muscle protein is overexpressed in
senescent human

[136241]fibroblasts. Biochem. Biophys. Res. Commun. 187: 1–7,
1992.

[136242]

[136243]

[136244]12431.

[136245]Tamanini, F.; Bontekoe, C.; Bakker, C. E.; van Unen, L.;
Anar,

[136246]B.; Willemsen, R.; Yoshida, M.; Galjaard, H.; Oostra, B. A.;
Hoogeveen,

[136247]A. T.: Different targets for the fragile X–related proteins
revealed

[136248]by their distinct nuclear localizations. Hum. Molec. Genet.
8: 863–869,

[136249]1999.

[136250]

[136251]

[136252]12432.

[136253]Tamanini, F.; Kirkpatrick, L. L.; Schonkeren, J.; van Unen, L.;

[136254]Bontekoe, C.; Bakker, C.; Nelson, D. L.; Galjaard, H.; Oost-
tra, B.

[136255]A.; Hoogeveen, A. T.: The fragile X-related proteins FXR1P
and FXR2P

[136256]contain a functional nucleolar-targeting signal equivalent
to the

[136257]HIV-1 regulatory proteins. Hum. Molec. Genet. 9:
1487-1493, 2000.

[136258]

[136259]

[136260]12433.

[136261]Radice, P.; Pensotti, V.; Jones, C.; Perry, H.; Pierotti, M. A.;

[136262]Tunnacliffe, A.: The human archaia gene, ARCN1, has
highly conserved

[136263]homologs in rice and Drosophila. Genomics 26: 101–106,
1995.

[136264]

[136265]

[136266]12434.

[136267]Tunnacliffe, A.; van de Vrugt, H.; Pensotti, V.; Radice, P.:
The

[136268]coatamer protein delta-COP, encoded by the archain
gene, is conserved

[136269]across diverse eukaryotes. Mammalian Genome 7:
784–786, 1996.

[136270]

[136271]

[136272]12435.

[136273]Hawtin, S. R.; Wesley, V. J.; Parslow, R. A.; Simms, J.; Miles,

[136274]A.; McEwan, K.; Wheatley, M.: A single residue (arg46) lo-
cated within

[136275]the N-terminus of the V(1a) vasopressin receptor is criti-
cal for binding

[136276]vasopressin but not peptide or nonpeptide antagonists.
Molec. Endocr. 16:

[136277]600–609, 2002.

[136278]

[136279]

[136280]12436.

[136281]Morel, A.; O'Carroll, A.–M.; Brownstein, M. J.; Lolait, S. J.:

[136282]Molecular cloning and expression of a rat V1a arginine
vasopressin

[136283]receptor. Nature 356: 523–526, 1992.

[136284]

[136285]

[136286]12437.

[136287]Thibonnier, M.; Auzan, C.; Madhun, Z.; Wilkins, P.; Berti–
Mattera,

[136288]L.; Clauser, E.: Molecular cloning, sequencing, and func–
tional expression

[136289]of a cDNA encoding the human V1a vasopressin receptor.
J. Biol. Chem. 269:

[136290]3304–3310, 1994.

[136291]

[136292]

[136293]12438.

[136294]Thibonnier, M.; Graves, M. K.; Wagner, M. S.; Auzan, C.; Clauser,

[136295]E.; Willard, H. F.: Structure, sequence, expression, and chromosomal

[136296]localization of the human V(1a) vasopressin receptor gene. Genomics 31:

[136297]327–334, 1996.

[136298]

[136299]

[136300]12439.

[136301]Young, L. J.; Nilsen, R.; Waymire, K. G.; MacGregor, G. R.; Insel,

[136302]T. R.: Increased affiliative response to vasopressin in mice expressing

[136303]the V(1A) receptor from a monogamous vole. Nature 400: 766–768,

[136304]1999.

[136305]

[136306]

[136307]12440.

[136308]Langer, L. O., Jr.; Cervenka, J.; Camargo, M.: A severe autosomal

[136309]recessive acromesomelic dysplasia, the Hunter–Thompson type, and comparison

[136310]with the Grebe type. Hum. Genet. 81: 323–328, 1989.

[136311]

[136312]

[136313]12441.

[136314]Robin, N. H.: Personal Communication. Cleveland, Ohio 11/3/1997.

[136315]

[136316]

[136317]12442.

[136318]Runner, M. N.: Linkage of brachypodism: a new member of linkage

[136319]group V of the house mouse. J. Hered. 50: 81–84, 1959.

[136320]

[136321]

[136322]12443.

[136323]Thomas, J. T.; Lin, K.; Nandedkar, M.; Camargo, M.; Cervenka,

[136324]J.; Luyten, F. P.: A human chondrodysplasia due to a mutation in

[136325]a TGF-beta superfamily member. Nature Genet. 12: 315-317, 1996.

[136326]

[136327]

[136328]12444.

[136329]Tsumaki, N.; Tanaka, K.; Arikawa-Hirasawa, E.; Nakase, T.; Kimura,

[136330]T.; Thomas, J. T.; Ochi, T.; Luyten, F. P.; Yamada, Y.: Role of CDMP-1

[136331]in skeletal morphogenesis: promotion of mesenchymal cell recruitment

[136332]and chondrocyte differentiation. J. Cell Biol. 144: 161-173, 1999.

[136333]

[136334]

[136335]12445.

[136336]Huh, T.-L.; Kim, Y.-O.; Oh, I.-U.; Song, B. J.; Inazawa, J.:

Assignment

[136337]of the human mitochondrial NAD(+)-specific isocitrate
dehydrogenase

[136338]alpha subunit (IDH3A) gene to 15q25.1-q25.2 by in situ
hybridization. Genomics 31:

[136339]295-296, 1996.

[136340]

[136341]

[136342]12446.

[136343]Kim, Y.-O.; Oh, I.-U.; Park, H.-S.; Jeng, J.; Song, B. J.; Huh,

[136344]T.-L.: Characterization of a cDNA clone for human
NAD(+)-specific

[136345]isocitrate dehydrogenase alpha-subunit and structural
comparison with

[136346]its isoenzymes from different species. Biochem. J. 308:
63-68, 1995.

[136347]

[136348]

[136349]12447.

[136350]Amann, J.; Kidd, V. J.; Lahti, J. M.: Characterization of putative

[136351]human homologues of the yeast chromosome transmission fidelity gene,

[136352]CHL1. J. Biol. Chem. 272: 3823–3832, 1997.

[136353]

[136354]

[136355]12448.

[136356]Amann, J.; Valentine, M.; Kidd, V. J.; Lahti, J. M.: Localization

[136357]of Chl1-related helicase genes to human chromosome regions 12p11 and

[136358]12p13: similarity between parts of these genes and conserved human

[136359]telomeric-associated DNA. Genomics 32: 260–265, 1996.

[136360]

[136361]

[136362]12449.

[136363]Frank, S.; Werner, S.: The human homologue of the yeast

CHL1 gene

[136364] is a novel keratinocyte growth factor-regulated gene. J.

Biol. Chem. 271:

[136365] 24337-24340, 1996.

[136366]

[136367]

[136368] 12450.

[136369] Barnes, L. D.; Garrison, P. N.; Siprashvili, Z.; Guranowski, A.;

[136370] Robinson, A. K.; Ingram, S. W.; Croce, C. M.; Ohta, M.; Huebner, K.

[136371]: Fhit, a putative tumor suppressor in humans, is a dinucleoside 5-prime,5-triple

[136372] prime-P(1),P(3)-triphosphate hydrolase. Biochemistry 35: 11529-11535,

[136373] 1996.

[136374]

[136375]

[136376] 12451.

[136377] Bernar, J.; Funderburk, S. J.; Sparkes, R. S.: The inducible

fragile

[136378]site on chromosome 3. (Letter) Hum. Genet. 66: 373 only,
1984.

[136379]

[136380]

[136381]12452.

[136382]Jin, C.-J.; Miners, J. O.; Lillywhite, K. J.; Mackenzie, P. I.

[136383]: cDNA cloning and expression of two new members of
the human liver

[136384]UDP-glucuronosyltransferase 2B subfamily. Biochem. Bio-
phys. Res.

[136385]Commun. 194: 496-503, 1993.

[136386]

[136387]

[136388]12453.

[136389]Krasnewich, D.; Kozak, C. A.; Nebert, D. W.; Mackenzie, P.
I.:

[136390]Localization of UDP glucuronosyltransferase gene(s) on
mouse chromosome

[136391]

[136392]12454.

[136393]Somat. Cell Molec. Genet. 13: 179–182, 1987.

[136394]

[136395]5. Monaghan, G.; Clarke, D. J.; Povey, S.; See, C. G.; Boxer, M.;

[136396]Burchell, B.: Isolation of a human YAC contig encompassing a cluster

[136397]of UGT2 genes and its regional localization to chromosome 4q13. Genomics 23:

[136398]496–499, 1994.

[136399]

[136400]

[136401]12455.

[136402]Monaghan, G.; Povey, S.; Burchell, B.; Boxer, M.: Localization

[136403]of a bile acid UDP–glucuronosyltransferase gene (UGT2B) to chromosome

[136404]4 using the polymerase chain reaction. Genomics 13: 908–909, 1992.

[136405]

[136406]

[136407]12456.

[136408]Riedy, M.; Wang, J. Y.; Miller, A. P.; Buckler, A.; Hall, J.;
Guida,

[136409]M.: Genomic organization of the UGT2b gene cluster on
human chromosome

[136410]4q13. Pharmacogenetics 10: 251–260, 2000.

[136411]

[136412]

[136413]12457.

[136414]Monaghan, G.; Clarke, D. J.; Povey, S.; See, C. G.; Boxer,
M.;

[136415]Burchell, B.: Isolation of a human YAC contig encompass–
ing a cluster

[136416]of UGT2 genes and its regional localization to chromo–
some 4q13. Genomics 23:

[136417]496–499, 1994.

[136418]

[136419]

[136420]12458.

[136421]Coffman, B. L.; Tephly, T. R.; Irshaid, Y. M.; Green, M. D.;
Smith,

[136422]C.; Jackson, M. R.; Wooster, R.; Burchell, B.: Characteriza-
tion and

[136423]primary sequence of a human hepatic microsomal estriol
UDP-glucuronosyltransferase. Arch.

[136424]Biochem. Biophys. 281: 170-175, 1990.

[136425]

[136426]

[136427]12459.

[136428]Buc-Caron, M. H.; Condamine, H.; Kerjaschki, D.: Rat Hey-
mann nephritis

[136429]antigen is closely related to brushin, a glycoprotein
present in early

[136430]mouse embryo epithelia. Ann. Inst. Pasteur/Immun. 138:
707-722,

[136431]1987.

[136432]

[136433]

[136434]12460.

[136435]Chatelet, F.; Brianti, E.; Ronco, P.; Roland, J.; Verroust, P.

[136436]: Ultrastructural localization by monoclonal antibodies of
brush border

[136437]antigens expressed by glomeruli: II. Extrarenal distribu-
tion. Am.

[136438]J. Path. 122: 512-519, 1986.

[136439]

[136440]

[136441]12461.

[136442]Chowdhary, B. P.; Lundgren, S.; Johansson, M.; Hjalmar, G.;
Akerstrom,

[136443]G.; Gustavsson, I.; Rask, L.: In situ hybridization mapping
of a

[136444]500-kDa calcium-sensing protein gene (LRP2) to human
chromosome region

[136445]2q31-q32.1 and porcine chromosome region 15q22-q24.
Cytogenet. Cell

[136446]Genet. 71: 120-123, 1995.

[136447]

[136448]

[136449]12462.

[136450]Farquhar, M. G.: The unfolding story of megalin (gp330):
now recognized

[136451]as a drug receptor (Editorial) J. Clin. Invest. 96: 1184 only,
1995.

[136452]

[136453]

[136454]12463.

[136455]Farquhar, M. G.; Saito, A.; Kerjaschki, D.; Orlando, R. A.:
The

[136456]Heymann nephritis antigenic complex: megalin (gp330)
and RAP. J.

[136457]Am. Soc. Nephrol. 6: 35–47, 1995.

[136458]

[136459]

[136460]12464.

[136461]Hjalm, G.; Murray, E.; Crumley, G.; Harazim, W.; Lundgren,
S.;

[136462]Onyango, I.; Bo, E. K.; Larsson, M.; Juhlin, C.; Hellman, P.;
Davis,

[136463]H.; Akerstrom, G.; Rask, L.; Morse, B.: Cloning and sequencing of

[136464]human gp330, a Ca(2+)-binding receptor with potential intracellular

[136465]signaling properties. *Europ. J. Biochem.* 239: 132–137, 1996.

[136466]

[136467]

[136468]12465.

[136469]Arinami, T.; Kondo, I.; Hamaguchi, H.; Nakajima, S.: Multifocal

[136470]meningiomas in a patient with a constitutional ring chromosome 22. *J.*

[136471]*Med. Genet.* 23: 178–180, 1986.

[136472]

[136473]

[136474]12466.

[136475]Petrella, R.; Levine, S.; Wilmot, P. L.; Ashar, K. D.; Casamassima,

[136476]A. C.; Shapiro, L. R.: Multiple meningiomas in a patient with constitutional

[136477]ring chromosome 22. Am. J. Med. Genet. 47: 184–186,
1993.

[136478]

[136479]

[136480]12467.

[136481]Peyrard, M.; Fransson, I.; Xie, Y.–G.; Han, F.–Y.; Rutledge,
M.

[136482]H.; Swahn, S.; Collins, J. E.; Dunham, I.; Collins, V. P.; Du-
manski,

[136483]J. P.: Characterization of a new member of the human
beta–adaptin

[136484]gene family from chromosome 22q12, a candidate menin-
gioma gene. Hum.

[136485]Molec. Genet. 3: 1393–1399, 1994.

[136486]

[136487]

[136488]12468.

[136489]Peyrard, M.; Pan, H.–Q.; Kedra, D.; Fransson, I.; Swahn, S.;
Hartman,

[136490]K.; Clifton, S. W.; Roe, B. A.; Dumanski, J. P.: Structure of
the

[136491]promoter and genomic organization of the human beta-
prime-adaptin

[136492]gene (BAM22) from chromosome 22q12. Genomics 36:
112–117, 1996.

[136493]

[136494]

[136495]12469.

[136496]Zankl, H.; Zang, K. D.: Correlations between clinical and
cytogenetical

[136497]data in 180 human meningiomas. Cancer Genet. Cyto-
genet. 1: 351–356,

[136498]1980.

[136499]

[136500]

[136501]12470.

[136502]Bai, C.; Sen, P.; Hofmann, K.; Ma, L.; Goebel, M.; Harper, J.
W.;

[136503]Elledge, S. J.: SKP1 connects cell cycle regulators to the
ubiquitin

[136504]proteolysis machinery through a novel motif, the F-box.
Cell 86:

[136505]263–274, 1996.

[136506]

[136507]

[136508]12471.

[136509]Kraus, B.; Pohlschmidt, M.; Leung, A. L. S.; Germino, G. G.;
Snarey,

[136510]A.; Schneider, M. C.; Reeders, S. T.; Frischauf, A.–M.: A
novel cyclin

[136511]gene (CCNF) in the region of the polycystic kidney disease
gene (PKD1). Genomics 24:

[136512]27–33, 1994.

[136513]

[136514]

[136515]12472.

[136516]Obermayr, F.; Sutherland, H. F.; Kraus, B.; Frischauf,
A.–M.:

[136517]Mouse cyclin F maps to a conserved linkage group on
mouse chromosome

[136518]17. Mammalian Genome 6: 149–150, 1995.

[136519]

[136520]

[136521]12473.

[136522]Donaldson, S. H.; Hirsh, A.; Li, D. C.; Holloway, G.; Chao, J.;

[136523]Boucher, R. C.; Gabriel, S. E.: Regulation of the epithelial sodium

[136524]channel by serine proteases in human airways. J. Biol. Chem. 277:

[136525]8338–8345, 2002.

[136526]

[136527]

[136528]12474.

[136529]Harvey, K. F.; Dinudom, A.; Cook, D. I.; Kumar, S.: The Nedd4-like

[136530]protein KIAA0439 is a potential regulator of the epithelial sodium

[136531]channel. J. Biol. Chem. 276: 8597–8601, 2001.

[136532]

[136533]

[136534]12475.

[136535]Arriza, J. L.; Kavanaugh, M. P.; Fairman, W. A.; Wu, Y.-N.;
Murdoch,

[136536]G. H.; North, R. A.; Amara, S. G.: Cloning and expression
of a human

[136537]neutral amino acid transporter with structural similarity to
the glutamate

[136538]transporter gene family. J. Biol. Chem. 268:
15329–15332, 1993.

[136539]

[136540]

[136541]12476.

[136542]Hofmann, K.; Duker, M.; Fink, T.; Lichter, P.; Stoffel, W.:
Human

[136543]neutral amino acid transporter ASCT1: structure of the
gene (SLC1A4)

[136544]and localization to chromosome 2p13–p15. Genomics 24:
20–26, 1994.

[136545]

[136546]

[136547]12477.

[136548]Shafqat, S.; Tamarappoo, B. K.; Kilberg, M. S.; Puranam, R.

S.;

[136549]McNamara, J. O.; Guadano-Ferraz, A.; Fremeau, R. T., Jr.:

Cloning

[136550]and expression of a novel Na(+)-dependent neutral amino acid transporter

[136551]structurally related to mammalian Na(+)/glutamate co-transporters. J.

[136552]Biol. Chem. 268: 15351-15355, 1993.

[136553]

[136554]

[136555]12478.

[136556]Zerangue, N.; Kavanaugh, M. P.: ASCT-1 is a neutral amino acid

[136557]exchanger with chloride channel activity. J. Biol. Chem. 271: 27991-27994,

[136558]1996.

[136559]

[136560]

[136561]12479.

[136562]Albertin, G.; Rossi, G. P.; Majone, F.; Tiso, N.; Mattara, A.;

[136563]Danieli, G. A.; Pessina, A. C.; Palu, G.: Fine mapping of the human

[136564]endothelin-converting enzyme gene by fluorescent in situ hybridization

[136565]and radiation hybrids. Biochem. Biophys. Res. Commun.
221: 682–687,

[136566]1996.

[136567]

[136568]

[136569]12480.

[136570]Sharp, J. D.; Pickard, R. T.; Chiou, X. G.; Manetta, J. V.; Ko-
vacevic,

[136571]S.; Miller, J. R.; Varshavsky, A. D.; Roberts, E. F.; Strifler, B.

[136572]A.; Brems, D. N.; Kramer, R. M.: Serine 228 is essential for
catalytic

[136573]activities of 85–kDa cytosolic phospholipase A2. J. Biol.
Chem. 269:

[136574]23250–23254, 1994.

[136575]

[136576]

[136577]12481.

[136578]Tamimi, R.; Steingrimsson, E.; Copeland, N. G.; Dyer-
Montgomery,

[136579]K.; Lee, J. E.; Hernandez, R.; Jenkins, N. A.; Tapscott, S. J.:

The

[136580]NEUROD gene maps to human chromosome 2q32 and mouse chromosome 2. Genomics 34:

[136581]418–421, 1996.

[136582]

[136583]

[136584]12482.

[136585]Newton, J. S.; Deed, R. W.; Mitchell, E. L. D.; Murphy, J. J.;

[136586]Norton, J. D.: A B cell specific immediate early human gene is located

[136587]on chromosome band 1q31 and encodes an alpha helical basic phosphoprotein. Biochim.

[136588]Biophys. Acta 1216: 314–316, 1993.

[136589]

[136590]

[136591]12483.

[136592]Sierra, D. A.; Gilbert, D. J.; Householder, D.; Grishin, N. V.;

[136593]Yu, K.; Ukidwe, P.; Barker, S. A.; He, W.; Wensel, T. G.; Otero, G.;

[136594]Brown, G.; Copeland, N. G.; Jenkins, N. A.; Wilkie, T. M.:

Evolution

[136595]of the regulators of G-protein signaling multigene family
in mouse

[136596]and human. Genomics 79: 177–185, 2002.

[136597]

[136598]

[136599]12484.

[136600]Hofstra, R. M. W.; Valdenaire, O.; Arch, E.; Osinga, J.;

Kroes,

[136601]H.; Loffler, B.–M.; Hamosh, A.; Meijers, C.; Buys, C. H. C.

M.: A

[136602]loss-of-function mutation in the endothelin-converting
enzyme 1 (ECE-1)

[136603]associated with Hirschsprung disease, cardiac defects,
and autonomic

[136604]dysfunction. (Letter) Am. J. Hum. Genet. 64: 304–308,
1999.

[136605]

[136606]

[136607]12485.

[136608]Matsuoka, R.; Sawamura, T.; Yamada, K.; Yoshida, M.; Furutani,

[136609]Y.; Ikura, T.; Shiraki, T.; Hoshikawa, H.; Shimada, K.; Tanzawa, K.;

[136610]Masaki, T.: Human endothelin converting enzyme gene (ECE1) mapped

[136611]to chromosomal region 1p36.1. Cytogenet. Cell Genet. 72: 322–324,

[136612]1996.

[136613]

[136614]

[136615]12486.

[136616]Schmidt, M.; Kroger, B.; Jacob, E.; Seulberger, H.; Subkowski,

[136617]T.; Otter, R.; Meyer, T.; Schmalzing, G.; Hillen, H.: Molecular characterization

[136618]of human and bovine endothelin converting enzyme (ECE–1). FEBS Lett. 356:

[136619]238–243, 1994.

[136620]

[136621]

[136622]12487.

[136623]Shimada, K.; Matsushita, Y.; Wakabayashi, K.; Takahashi, M.; Matsubara,

[136624]A.; Iijima, Y.; Tanzawa, K.: Cloning and functional expression of

[136625]human endothelin-converting enzyme cDNA. Biochem. Biophys. Res. Commun. 207:

[136626]807–812, 1995.

[136627]

[136628]

[136629]12488.

[136630]Valdenaire, O.; Rohrbacher, E.; Mattei, M.–G.: Organization of

[136631]the gene encoding the human endothelin-converting enzyme (ECE–1). J.

[136632]Biol. Chem. 270: 29794–29798, 1995.

[136633]

[136634]

[136635]12489.

[136636]Yorimitsu, K.; Moroi, K.; Inagaki, N.; Saito, T.; Masuda, Y.;

[136637]Masaki, T.; Seino, S.; Kimura, S.: Cloning and sequencing
of a human

[136638]endothelin converting enzyme in renal adenocarcinoma
(ACHN) cells

[136639]producing endothelin-2. Biochem. Biophys. Res. Commun.
208: 721-727,

[136640]1995.

[136641]

[136642]

[136643]12490.

[136644]Ivey-Hoyle, M.; Conroy, R.; Huber, H. E.; Goodhart, P. J.;
Oliff,

[136645]A.; Heimbrook, D. C.: Cloning and characterization of
E2F-2, a novel

[136646]protein with the biochemical properties of transcription
factor E2F. Molec.

[136647]Cell. Biol. 13: 7802-7812, 1993.

[136648]

[136649]

[136650]12491.

[136651]Lees, J. A.; Saito, M.; Vidal, M.; Valentine, M.; Look, T.;
Harlow,

[136652]E.; Dyson, N.; Helin, K.: The retinoblastoma protein binds
to a family

[136653]of E2F transcription factors. Molec. Cell. Biol. 13:
7813–7825,

[136654]1993.

[136655]

[136656]

[136657]12492.

[136658]Cloud, J. E.; Rogers, C.; Reza, T. L.; Ziebold, U.; Stone, J.
R.;

[136659]Picard, M. H.; Caron, A. M.; Bronson, R. T.; Lees, J. A.: Mu-
tant

[136660]mouse models reveal the relative roles of E2F1 and E2F3
in vivo. Molec.

[136661]Cell. Biol. 22: 2663–2672, 2002.

[136662]

[136663]

[136664]12493.

[136665]He, Y.; Armanious, M. K.; Thomas, M. J.; Cress, W. D.:

Identification

[136666]of E2F-3B, an alternative form of E2F-3 lacking a conserved N-terminal

[136667]region. *Oncogene* 19: 3422-3433, 2000.

[136668]

[136669]

[136670]12494.

[136671]Sharp, J. D.; White, D. L.; Chiou, X. G.; Goodson, T.; Gamboa,

[136672]G. C.; McClure, D.; Burgett, S.; Hoskins, J.; Skatrud, P. L.; Sportsman,

[136673]J. R.; Becker, G. W.; Kang, L. H.; Roberts, E. F.; Kramer, R. M.:

[136674]Molecular cloning and expression of human Ca(2+)-sensitive cytosolic

[136675]phospholipase A2. *J. Biol. Chem.* 266: 14850-14853, 1991.

[136676]

[136677]

[136678]12495.

[136679]Sheridan, A. M.; Force, T.; Yoon, H.-J.; O'Leary, E.;

Choukroun,

[136680]G.; Taheri, M. R.; Bonventre, J. V.: PLIP, a novel splice variant

[136681]of Tip60, interacts with group IV cytosolic phospholipase A2, induces

[136682]apoptosis, and potentiates prostaglandin production. Molec. Cell.

[136683]Biol. 21: 4470–4481, 2001.

[136684]

[136685]

[136686]12496.

[136687]Skorecki, K. L.: Personal Communication. Toronto, Canada 4/19/1995.

[136688]

[136689]

[136690]12497.

[136691]Tay, A.; Simon, J. S.; Squire, J.; Hamel, K.; Jacob, H. J.; Skorecki,

[136692]K.: Cytosolic phospholipase A2 gene in human and rat: chromosomal

[136693] localization and polymorphic markers. Genomics 26:
138–141, 1995.

[136694]

[136695]

[136696] 12498.

[136697] Halford, M. M.; Armes, J.; Buchert, M.; Meskenaite, V.;
Grail,

[136698] D.; Hibbs, M. L.; Wilks, A. F.; Farlie, P. G.; Newgreen, D. F.;
Hovens,

[136699] C. M.; Stacker, S. A.: Ryk–deficient mice exhibit craniofa-
cial defects

[136700] associated with perturbed Eph receptor crosstalk. Nature
Genet. 25:

[136701] 414–418, 2000.

[136702]

[136703]

[136704] 12499.

[136705] Nakamura, S.; Stock, D. W.; Wydner, K. L.; Bollekens, J. A.;
Takeshita,

[136706] K.; Nagai, B. M.; Chiba, S.; Kitamura, T.; Freeland, T. M.;
Zhao,

[136707]Z.; Minowada, J.; Lawrence, J. B.; Weiss, K. M.; Ruddle, F.

H.: Genomic

[136708]analysis of a new mammalian distal-less gene: Dlx7. Ge-
nomics 38:

[136709]314–324, 1996.

[136710]

[136711]

[136712]12500.

[136713]Singh, J.; Itahana, Y.; Parrinello, S.; Murata, K.; Desprez,
P.–Y.

[136714]: Molecular cloning and characterization of a zinc finger
protein

[136715]involved in Id–1–stimulated mammary epithelial cell
growth. J. Biol.

[136716]Chem. 276: 11852–11858, 2001.

[136717]

[136718]

[136719]12501.

[136720]Kumar, S.; Kinoshita, M.; Noda, M.; Copeland, N. G.; Jenk-
ins, N.

[136721]A.: Induction of apoptosis by the mouse Nedd2 gene,
which encodes

[136722]a protein similar to the product of the *Caenorhabditis elegans* cell

[136723]death gene *ced-3* and the mammalian IL-
1-beta-converting enzyme. *Genes*

[136724]*Dev.* 8: 1613-1626, 1994.

[136725]

[136726]

[136727]12502.

[136728]Bollag, R. J.; Siegfried, Z.; Cebra-Thomas, J. A.; Garvey, N.;

[136729]Davison, E. M.; Silver, L. M.: An ancient family of embry-
onically

[136730]expressed mouse genes sharing a conserved protein motif
with the T

[136731]locus. *Nature Genet.* 7: 383-389, 1994.

[136732]

[136733]

[136734]12503.

[136735]Campbell, C.; Goodrich, K.; Casey, G.; Beatty, B.: Cloning

and

[136736]mapping of a human gene (TBX2) sharing a highly conserved protein

[136737]motif with the Drosophila omb gene. Genomics 28: 255–260, 1995.

[136738]

[136739]

[136740]12504.

[136741]Campbell, C. E.; Casey, G.; Goodrich, K.: Genomic structure of

[136742]TBX2 indicates conservation with distantly related T-box genes. Mammalian

[136743]Genome 9: 70–73, 1998.

[136744]

[136745]

[136746]12505.

[136747]Jacobs, J. J. L.; Keblusek, P.; Robanus-Maandag, E.; Kristel, P.;

[136748]Lingbeek, M.; Nederlof, P. M.; van Welsem, T.; van de Vijver, M. J.;

[136749]Koh, E. Y.; Daley, G. Q.; van Lohuizen, M.: Senescence by-

pass screen

[136750]identifies TBX2, which represses Cdkn2a(p19ARF) and is amplified in

[136751]a subset of human breast cancers. Nature Genet. 26: 291–299, 2000.

[136752]

[136753]

[136754]12506.

[136755]Law, D. J.; Gebhur, T.; Garvey, N.; Agulnik, S. I.; Silver, L.

[136756]M.: Identification, characterization, and localization to chromosome

[136757]17q21–22 of the human TBX2 homolog, member of a conserved developmental

[136758]gene family. Mammalian Genome 6: 793–797, 1995.

[136759]

[136760]

[136761]12507.

[136762]Pflugfelder, G. O.; Roth, H.; Poeck, B.: A homology domain shared

[136763]between Drosophila optomotor-blind and mouse

Brachyury is involved

[136764]in DNA binding. Biochem. Biophys. Res. Commun. 186:
918–925, 1992.

[136765]

[136766]

[136767]12508.

[136768]Sowden, J. C.; Holt, J. K. L.; Meins, M.; Smith, H. K.; Bhat-
tacharya,

[136769]S. S.: Expression of Drosophila omb-related T-box genes
in the developing

[136770]human and mouse neural retina. Invest. Ophthalm. Vis. Sci.
42: 3095–3102,

[136771]2001.

[136772]

[136773]

[136774]12509.

[136775]Walter, L.; Dirks, B.; Rothermel, E.; Heyens, M.; Szpirer, C.;

[136776]Levan, G.; Gunther, E.: A novel, conserved gene of the rat
that is

[136777]developmentally regulated in the testis. Mammalian

Genome 5: 216–221,

[136778]1994.

[136779]

[136780]

[136781]12510.

[136782]Walter, L.; Marynen, P.; Szpirer, J.; Levan, G.; Gunther, E.:

[136783]Identification of a novel conserved human gene, TEGT.

Genomics 28:

[136784]301–304, 1995.

[136785]

[136786]

[136787]12511.

[136788]Xu, Q.; Reed, J. C.: Bax inhibitor–1, a mammalian apopto–
sis suppressor

[136789]identified by functional screening in yeast. Molec. Cell 1:
337–346,

[136790]1998.

[136791]

[136792]

[136793]12512.

[136794]Hsu, Y.-C.; Perin, M. S.: Human neuronal pentraxin II
(NPTX2):

[136795]conservation, genomic structure, and chromosomal localization. Genomics 28:

[136796]220–227, 1995.

[136797]

[136798]

[136799]12513.

[136800]Schlimgen, A. K.; Helms, J. A.; Vogel, H.; Perin, M. S.: Neuronal

[136801]pentraxin, a secreted protein with homology to acute
phase proteins

[136802]of the immune system. Neuron 14: 519–526, 1995.

[136803]

[136804]

[136805]12514.

[136806]Kumar, S.; White, D. L.; Takai, S.; Turczynowicz, S.; Jut-
tner,

[136807]C. A.; Hughes, T. P.: Apoptosis regulatory gene NEDD2

maps to human

[136808]chromosome segment 7q34–35, a region frequently affected in haematological

[136809]neoplasms. Hum. Genet. 95: 641–644, 1995.

[136810]

[136811]

[136812]12515.

[136813]Lassus, P.; Opitz–Araya, X.; Lazebnik, Y.: Requirement for caspase–2

[136814]in stress–induced apoptosis before mitochondrial permeabilization. Science 297:

[136815]1352–1354, 2002.

[136816]

[136817]

[136818]12516.

[136819]Aksoy, I. A.; Callen, D. F.; Apostolou, S.; Her, C.; Weinshilboum,

[136820]R. M.: Thermolabile phenol sulfotransferase gene (STM) localization

[136821]to human chromosome 16p11.2. Genomics 23: 275–277,

1994.

[136822]

[136823]

[136824]12517.

[136825]Aksoy, I. A.; Weinshilboum, R. M.: Human thermolabile
phenol sulfotransferase

[136826]gene (STM): molecular cloning and structural characteri-
zation. Biochem.

[136827]Biophys. Res. Commun. 208: 786–795, 1995.

[136828]

[136829]

[136830]12518.

[136831]Wood, T. C.; Aksoy, I. A.; Aksoy, S.; Weinshilboum, R. M.:
Human

[136832]liver thermolabile phenol sulfotransferase: cDNA cloning,
expression

[136833]and characterization. Biochem. Biophys. Res. Commun.
198: 1119–1127,

[136834]1994.

[136835]

[136836]

[136837]12519.

[136838]Zhu, X.; Veronese, M. E.; Bernard, C. C. A.; Sansom, L. N.;
McManus,

[136839]M. E.: Identification of two human brain aryl sulfotrans-
ferase cDNAs. Biochem.

[136840]Biophys. Res. Commun. 195: 120–127, 1993.

[136841]

[136842]

[136843]12520.

[136844]Fuhlbrigge, R. C.; Kieffer, J. D.; Armerding, D.; Kupper, T.
S.

[136845]: Cutaneous lymphocyte antigen is a specialized form of
PSGL–1 expressed

[136846]on skin–homing T cells. Nature 389: 978–981, 1997.

[136847]

[136848]

[136849]12521.

[136850]Herron, M. J.; Nelson, C. M.; Larson, J.; Snapp, K. R.;

Kansas,

[136851]G. S.; Goodman, J. L.: Intracellular parasitism by the human granulocytic

[136852]ehrlichiosis bacterium through the P-selectin ligand, PSGL-1. Science 288:

[136853]1653-1656, 2000.

[136854]

[136855]

[136856]12522.

[136857]Veldman, G. M.; Bean, K. M.; Cumming, D. A.; Eddy, R. L.; Sait,

[136858]S. N. J.; Shows, T. B.: Genomic organization and chromosomal localization

[136859]of the gene encoding human P-selectin glycoprotein ligand. J. Biol.

[136860]Chem. 270: 16470-16475, 1995.

[136861]

[136862]

[136863]12523.

[136864]Yang, J.; Galipeau, J.; Kozak, C. A.; Furie, B. C.; Furie, B.:

[136865]Mouse P-selectin glycoprotein ligand-1: molecular cloning, chromosomal

[136866]localization, and expression of a functional P-selectin receptor. Blood 87:

[136867]4176-4186, 1996.

[136868]

[136869]

[136870]12524.

[136871]Brown, C. B.; Boyer, A. S.; Runyan, R. B.; Barnett, J. V.: Requirement

[136872]of type III TGF-beta receptor for endocardial cell transformation

[136873]in the heart. Science 283: 2080-2082, 1999.

[136874]

[136875]

[136876]12525.

[136877]Lewis, K. A.; Gray, P. C.; Blount, A. L.; MacConell, L. A.; Wiater,

[136878]E.; Bilezikjian, L. M.; Vale, W.: Betaglycan binds inhibin and can

[136879]mediate functional antagonism of activin signalling. Nature 404:

[136880]411–414, 2000.

[136881]

[136882]

[136883]12526.

[136884]Hu, Y.-F.; Luscher, B.; Admon, A.; Mermod, N.; Tjian, R.:
Transcription

[136885]factor AP-4 contains multiple dimerization domains that
regulate dimer

[136886]specificity. Genes Dev. 4: 1741–1752, 1990.

[136887]

[136888]

[136889]12527.

[136890]Mermod, N.; Williams, T. J.; Tjian, R.: Enhancer binding
factors

[136891]AP-4 and AP-1 act in concert to activate SV40 late tran-
scription in

[136892]vitro. Nature 332: 557–561, 1988.

[136893]

[136894]

[136895]12528.

[136896]Carr, C. S.; Sharp, P. A.: A helix-loop-helix protein related

[136897]to the immunoglobulin E box-binding proteins. Molec.

Cell. Biol. 10:

[136898]4384-4388, 1990.

[136899]

[136900]

[136901]12529.

[136902]Bonner, C. A.; Loftus, S. K.; Wasmuth, J. J.: Isolation, characterization,

[136903]and precise physical localization of human CDX1, a caudal-type homeobox

[136904]gene. Genomics 28: 206-211, 1995.

[136905]

[136906]

[136907]12530.

[136908]Subramanian, V.; Meyer, B. I.; Gruss, P.: Disruption of the murine

[136909]homeobox gene Cdx1 affects axial skeletal identities by altering the

[136910]mesodermal expression domains of Hox genes. Cell 83: 641–653, 1995.

[136911]

[136912]

[136913]12531.

[136914]Henske, E. P.; Short, M. P.; Jozwiak, S.; Bovey, C. M.; Ram-lakhan,

[136915]S.; Haines, J. L.; Kwiatkowski, D. J.: Identification of VAV2 on
on

[136916]9q34 and its exclusion as the tuberous sclerosis gene
TSC1. Ann.

[136917]Hum. Genet. 59: 25–37, 1995.

[136918]

[136919]

[136920]12532.

[136921]Schwientek, T.; Nomoto, M.; Levery, S. B.; et al: Control of
O-glycan

[136922]branch formation. J. Biol. Chem. 274: 4504–4512, 1999.

[136923]

[136924]

[136925]12533.

[136926]Walczak, H.; Degli-Esposti, M. A.; Johnson, R. S.; Smolak,
P. J.;

[136927]Waugh, J. Y.; Boiani, N.; Timour, M. S.; Gerhart, M. J.;
Schooley,

[136928]K. A.; Smith, C. A.; Goodwin, R. G.; Rauch, C. T.: TRAIL-R2:
a novel

[136929]apoptosis-mediating receptor for TRAIL. EMBO J. 16:
5386-5397, 1997.

[136930]

[136931]

[136932]12534.

[136933]Wu, G. S.; Burns, T. F.; McDonald, E. R., III; Jiang, W.;
Meng,

[136934]R.; Krantz, I. D.; Kao, G.; Gan, D.-D.; Zhou, J.-Y.; Muschel,
R.;

[136935]Hamilton, S. R.; Spinner, N. B.; Markowitz, S.; Wu, G.; El-
Deiry,

[136936]W. S.: KILLER/DR5 is a DNA damage-inducible

p53-regulated death receptor

[136937]gene. Nature Genet. 17: 141–143, 1997.

[136938]

[136939]

[136940]12535.

[136941]Kanaar, R.; Troelstra, C.; Swagemakers, S. M. A.; Essers, J.;
Smit,

[136942]B.; Franssen, J.–H.; Pastink, A.; Bezzubova, O. Y.; Buer–
stedde, J.–M.;

[136943]Clever, B.; Heyer, W.–D.; Hoeijmakers, J. H. J.: Human and
mouse

[136944]homologs of the *Saccharomyces cerevisiae* RAD54 DNA
repair gene: evidence

[136945]for functional conservation. Curr. Biol. 6: 828–838, 1996.

[136946]

[136947]

[136948]12536.

[136949]Matsuda, M.; Miyagawa, K.; Takahashi, M.; Fukuda, T.;
Kataoka,

[136950]T.; Asahara, T.; Inui, H.; Watatani, M.; Yasutomi, M.; Ka–

mada, N.;

[136951]Dohi, K.; Kamiya, K.: Mutations in the RAD54 recombination gene in

[136952]primary cancers. *Oncogene* 18: 3427–3430, 1999.

[136953]

[136954]

[136955]12537.

[136956]Tucker, J. E.; Winkfein, R. J.; Cooper, C. B.; Schnetkamp, P. P.

[136957]: cDNA cloning of the human retinal rod Na–Ca + K exchanger: comparison

[136958]with a revised bovine sequence. *Invest. Ophthal. Vis. Sci.* 39: 435–440,

[136959]1998.

[136960]

[136961]

[136962]12538.

[136963]Tucker, J. E.; Winkfein, R. J.; Murthy, S. K.; Friedman, J. S.;

[136964]Walter, M. A.; Demetrick, D. J.; Schnetkamp, P. P. M.: Chromosomal

[136965]localization and genomic organization of the human retinal rod Na-Ca+K

[136966]exchanger. Hum. Genet. 103: 411-414, 1998.

[136967]

[136968]

[136969]12539.

[136970]Dennis, J. A.; Healy, P. J.; Beaudet, A. L.; O'Brien, W. E.:

[136971]Molecular definition of bovine argininosuccinate synthetase deficiency. Proc.

[136972]Nat. Acad. Sci. 86: 7947-7951, 1989.

[136973]

[136974]

[136975]12540.

[136976]McCarrey, J. R.; Riggs, A. D.: Determinator-inhibitor pairs as

[136977]a mechanism for threshold setting in development: a possible function

[136978]for pseudogenes. Proc. Nat. Acad. Sci. 83: 679-683, 1986.

[136979]

[136980]

[136981]12541.

[136982]Su, T.-S.; Nussbaum, R. L.; Airhart, S.; Ledbetter, D. H.;
Mohandas,

[136983]T.; O'Brien, W. E.; Beaudet, A. L.: Human chromosomal as-
signments

[136984]for 14 argininosuccinate synthetase pseudogenes: cloned
DNAs as reagents

[136985]for cytogenetic analysis. Am. J. Hum. Genet. 36: 954-964,
1984.

[136986]

[136987]

[136988]12542.

[136989]Hara, Y.; Wakamori, M.; Ishii, M.; Maeno, E.; Nishida, M.;
Yoshida,

[136990]T.; Yamada, H.; Shimizu, S.; Mori, E.; Kudoh, J.; Shimizu,
S.; Kurose,

[136991]H.; Okada, Y.; Imoto, K.; Mori, Y.: LTRPC2
Ca(2+)-permeable channel

[136992]activated by changes in redox status confers susceptibility
to cell

[136993]death. Molec. Cell 9: 163–173, 2002.

[136994]

[136995]

[136996]12543.

[136997]Harteneck, C.; Plant, T. D.; Schultz, G.: From worm to
man: three

[136998]subfamilies of TRP channels. Trends Neurosci. 23:
159–166, 2000.

[136999]

[137000]

[137001]12544.

[137002]Kudoh, J.; Nagamine, K.; Asakawa, S.; Abe, I.; Kawasaki,
K.; Maeda,

[137003]H.; Tsujimoto, S.; Minoshima, S.; Ito, F.; Shimizu, N.: Lo-
calization

[137004]of 16 exons to a 450–kb region involved in the autoim-
mune polyglandular

[137005]disease type I (APECED) on human chromosome 21q22.3.
DNA Res. 4:

[137006]45–52, 1997.

[137007]

[137008]

[137009]12545.

[137010]Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.;
Asakawa,

[137011]S.; Ito, F.; Shimizu, N.: Molecular cloning of a novel puta-
tive Ca(2+)

[137012]channel protein (TRPC7) highly expressed in brain. Ge-
nomics 54:

[137013]124–131, 1998.

[137014]

[137015]

[137016]12546.

[137017]Perraud, A.-L.; Fleig, A.; Dunn, C. A.; Bagley, L. A.; Launay,

[137018]P.; Schmitz, C.; Stokes, A. J.; Zhu, Q.; Bessman, M. J.; Pen-
ner, R.;

[137019]Kinet, J.-P.; Scharenberg, A. M.: ADP-ribose gating of the
calcium-permeable

[137020]LTRPC2 channel revealed by Nudix motif homology. Na-
ture 411: 595–599,

[137021]2001.

[137022]

[137023]

[137024]12547.

[137025]Sano, Y.; Inamura, K.; Miyake, A.; Mochizuki, S.; Yokoi, H.;
Matsushime,

[137026]H.; Furuichi, K.: Immunocyte Ca(2+) influx system mediated by LTRPC2. Science 293:

[137027]1327–1330, 2001.

[137028]

[137029]

[137030]12548.

[137031]Koyama, N.; Ishibashi, K.; Kuwahara, M.; Inase, N.; Ichioka,
M.;

[137032]Sasaki, S.; Marumo, F.: Cloning and functional expression
of human

[137033]aquaporin8 cDNA and analysis of its gene. Genomics 54:
169–172,

[137034]1998.

[137035]

[137036]

[137037]12549.

[137038]Viggiano, L.; Rocchi, M.; Svelto, M.; Calamita, G.: Assignment

[137039]of the aquaporin-8 water channel gene (AQP8) to human chromosome 16p12. Cytogenet.

[137040]Cell Genet. 84: 208-210, 1999.

[137041]

[137042]

[137043]12550.

[137044]Graler, M. H.; Bernhardt, G.; Lipp, M.: EDG6, a novel G-protein-coupled

[137045]receptor related to receptors for bioactive lysophospholipids, is

[137046]specifically expressed in lymphoid tissue. Genomics 53: 164-169,

[137047]1998.

[137048]

[137049]

[137050]12551.

[137051]Jedlicka, A. E.; Taylor, E. W.; Meyers, D. A.; Liu, Z.; Levitt,

[137052]R. C.: Localization of the highly polymorphic locus

D19S120 to 19p13.3

[137053]by linkage. Cytogenet. Cell. Genet. 65: 140 only, 1994.

[137054]

[137055]

[137056]12552.

[137057]Koegl, M.; Hoppe, T.; Schlenker, S.; Ulrich, H. D.; Mayer, T.
U.;

[137058]Jentsch, S.: A novel ubiquitination factor, E4, is involved in
multiubiquitin

[137059]chain assembly. Cell 96: 635–644, 1999.

[137060]

[137061]

[137062]12553.

[137063]Nonaka, S.; Tanaka, Y.; Okada, Y.; Takeda, S.; Harada, A.;
Kanai,

[137064]Y.; Kido, M.; Hirokawa, N.: Randomization of left–right
asymmetry

[137065]due to loss of nodal cilia generating leftward flow of ex–

traembryonic

[137066]fluid in mice lacking KIF3B motor protein. Cell 95:
829–837, 1998.

[137067]

[137068]

[137069]12554.

[137070]Yamazaki, H.; Nakata, T.; Okada, Y.; Hirokawa, N.:
KIF3A/B: a

[137071]heterodimeric kinesin superfamily protein that works as a
microtubule

[137072]plus end–directed motor for membrane organelle trans-
port. J. Cell

[137073]Biol. 130: 1387–1399, 1995.

[137074]

[137075]

[137076]12555.

[137077]Allikmets, R.; Schriml, L. M.; Hutchinson, A.; Romano-
Spica, V.;

[137078]Dean, M.: A human placenta–specific ATP–binding cas-
sette gene (ABCP)

[137079]on chromosome 4q22 that is involved in multidrug resis–

tance. Cancer

[137080]Res. 58: 5337–5339, 1998.

[137081]

[137082]

[137083]12556.

[137084]Bailey–Dell, K. J.; Hassel, B.; Doyle, L. A.; Ross, D. D.: Pro-moter

[137085]characterization and genomic organization of the human breast cancer

[137086]resistance protein (ATP–binding cassette transporter G2) gene. Biochim.

[137087]Biophys. Acta 1520: 234–241, 2001.

[137088]

[137089]

[137090]12557.

[137091]Doyle, L. A.; Yang, W.; Abruzzo, L. V.; Krogmann, T.; Gao, Y.;

[137092]Rishi, A. K.; Ross, D. D.: A multidrug resistance trans-porter from

[137093]human MCF–7 breast cancer cells. Proc. Nat. Acad. Sci. 95:

15665–15670,

[137094]1998.

[137095]

[137096]

[137097]12558.

[137098]Suzuki, N.; Zara, J.; Sato, T.; Ong, E.; Bakhiet, N.; Oshima,
R.

[137099]G.; Watson, K. L.; Fukuda, M. N.: A cytoplasmic protein,
bystin,

[137100]interacts with trophinin, tastin, and cytokeratin and may
be involved

[137101]in trophinin-mediated cell adhesion between trophoblast
and endometrial

[137102]epithelial cells. Proc. Nat. Acad. Sci. 95: 5027–5032,
1998.

[137103]

[137104]

[137105]12559.

[137106]Cerutti, A.; Schaffer, A.; Goodwin, R. G.; Shah, S.; Zan, H.;
Ely,

[137107]S.; Casali, P.: Engagement of CD153 (CD30 ligand) by

CD30-positive

[137108]T cells inhibits class switch DNA recombination and anti-body production

[137109]in human IgD-positive IgM-positive B cells. J. Immun. 165: 786-794,

[137110]2000.

[137111]

[137112]

[137113]12560.

[137114]Croager, E. J.; Abraham, L. J.: Characterisation of the human man

[137115]CD30 ligand gene structure. Biochim. Biophys. Acta 1353: 231-235,

[137116]1997.

[137117]

[137118]

[137119]12561.

[137120]Smith, C. A.; Gruss, H.-J.; Davis, T.; Anderson, D.; Farrah, T.;

[137121]Baker, E.; Sutherland, G. R.; Brannan, C. I.; Copeland, N.

G.; Jenkins,

[137122]N. A.; Grabstein, K. H.; Gliniak, B.; and 9 others: CD30 antigen,

[137123]a marker for Hodgkin's lymphoma, is a receptor whose ligand defines

[137124]an emerging family of cytokines with homology to TNF.

Cell 73: 1349–1360,

[137125]1993.

[137126]

[137127]

[137128]12562.

[137129]Hurskainen, T. L.; Hirohata, S.; Seldin, M. F.; Apte, S. S.:

ADAM–TS5,

[137130]ADAM–TS6, and ADAM–TS7, novel members of a new family of zinc metalloproteases:

[137131]general features and genomic distribution of the ADAM–TS family. J.

[137132]Biol. Chem. 274: 25555–25563, 1999.

[137133]

[137134]

[137135]12563.

[137136]Tang, B. L.; Hong, W.: ADAMTS: a novel family of proteases with

[137137]an ADAM protease domain and thrombospondin 1 repeats. FEBS Lett. 445:

[137138]223–225, 1999.

[137139]

[137140]

[137141]12564.

[137142]Tortorella, M. D.; Burn, T. C.; Pratta, M. A.; Abbaszade, I.; Hollis,

[137143]J. M.; Liu, R.; Rosenfeld, S. A.; Copeland, R. A.; Decicco, C. P.;

[137144]Wynn, R.; Rockwell, A.; Yang, F.; and 16 others: Purification and

[137145]cloning of aggrecanase-1: a member of the ADAMTS family of proteins. Science 284:

[137146]1664–1666, 1999.

[137147]

[137148]

[137149]12565.

[137150]Higashide, T.; Inana, G.: Characterization of the gene for
HRG4

[137151](UNC119), a novel photoreceptor synaptic protein homologous to unc-119. Genomics 57:

[137152]446–450, 1999.

[137153]

[137154]

[137155]12566.

[137156]Higashide, T.; McLaren, M. J.; Inana, G.: Localization of
HRG4,

[137157]a photoreceptor protein homologous to unc-119, in ribbon synapse. Invest.

[137158]Ophthal. Vis. Sci. 39: 690–698, 1998.

[137159]

[137160]

[137161]12567.

[137162]Higashide, T.; Murakami, A.; McLaren, M. J.; Inana, G.:
Cloning

[137163]of the cDNA for a novel photoreceptor protein. J. Biol.
Chem. 271:

[137164]1797–1804, 1996.

[137165]

[137166]

[137167]12568.

[137168]Swanson, D. A.; Chang, J. T.; Campochiaro, P. A.; Zack, D.
J.;

[137169]Valle, D.: Mammalian orthologs of *C. elegans* unc-119
highly expressed

[137170]in photoreceptors. Invest. Ophthalm. Vis. Sci. 39:
2085–2094, 1998.

[137171]

[137172]

[137173]12569.

[137174]Gundelfinger, E.: Personal Communication. Madgeburg,
FRG. 1/8/1999.

[137175]

[137176]

[137177]12570.

[137178]Hashida, H.; Goto, J.; Zhao, N.; Takahashi, N.; Hirai, M.;

Kanazawa,

[137179]I.; Sakaki, Y.: Cloning and mapping of ZNF231, a novel
brain-specific

[137180]gene encoding neuronal double zinc finger protein whose
expression

[137181]is enhanced in a neurodegenerative disorder, multiple
system atrophy

[137182](MSA). Genomics 54: 50–58, 1998.

[137183]

[137184]

[137185]12571.

[137186]Santorelli, F. M.; Patrono, C.; Fortini, D.; Tessa, A.; Co-
manducci,

[137187]G.; Bertini, E.; Pierallini, A.; Amabile, G. A.; Casali, C.: In-
trafamilial

[137188]variability in hereditary spastic paraplegia associated with
an SPG4

[137189]gene mutation. Neurology 55: 702–705, 2000.

[137190]

[137191]

[137192]12572.

[137193]Sauter, S.; Mitterski, B.; Klimpe, S.; Bonsch, D.; Schols, L.;

[137194]Visbeck, A.; Papke, T.; Hopf, H. C.; Engel, W.; Deufel, T.;
Epplen,

[137195]J. T.; Neesen, J.: Mutation analysis of the spastin gene
(SPG4) in

[137196]patients in Germany with autosomal dominant hereditary
spastic paraplegia. Hum.

[137197]Mutat. 20: 127–132, 2002.

[137198]

[137199]

[137200]12573.

[137201]Svenson, I. K.; Ashley-Koch, A. E.; Pericak-Vance, M. A.;
Marchuk,

[137202]D. A.: A second leaky splice-site mutation in the spastin
gene. (Letter) Am.

[137203]J. Hum. Genet. 69: 1407–1409, 2001.

[137204]

[137205]

[137206]12574.

[137207]Flannery, C. R.; Hughes, C. E.; Schumacher, B. L.; Tudor,

D.; Aydelotte,

[137208]M. B.; Kuettner, K. E.; Caterson, B.: Articular cartilage superficial

[137209]zone protein (SZP) is homologous to megakaryocyte stimulating factor

[137210]precursor and is a multifunctional proteoglycan with potential growth-promoting,

[137211]cytoprotective, and lubricating properties in cartilage metabolism. Biochem.

[137212]Biophys. Res. Commun. 254: 535–541, 1999.

[137213]

[137214]

[137215]12575.

[137216]Ikegawa, S.; Sano, M.; Koshizuka, Y.; Nakamura, Y.: Isolation,

[137217]characterization and mapping of the mouse and human PRG4 (proteoglycan

[137218]4) genes. Cytogenet. Cell Genet. 90: 291–297, 2000.

[137219]

[137220]

[137221]12576.

- [137222]Merberg, D. M.; Fitz, L. J.; Temple, P.; Giannotti, J.; Murtha,
[137223]P.; Fitzgerald, M.; Scaltreto, H.; Kelleher, K.; Preissner, K.;
Kriz,
[137224]R.; Jacobs, K.; Turner, K.: In: Preissner, K. T.; Rosenblatt,
S.;
[137225]Kost, C.; Wegerhoff, J.; Mosher, D. F. (eds.): Biology of Vit-
ronectins
[137226]and Their Receptors. Elsevier Science, B.V. 1993. Pp.
45–53.
- [137227]
- [137228]
- [137229]12577.
- [137230]Schumacher, B. L.; Block, J. A.; Schmid, T. M.; Aydelotte,
M. B.;
[137231]Kuettner, K. E.: A novel proteoglycan synthesized and se-
creted by
[137232]chondrocytes of the superficial zone of articular cartilage.
Arch.
[137233]Biochem. Biophys. 15: 144–152, 1994.
- [137234]
- [137235]

[137236]12578.

[137237]Pennica, D.; Swanson, T. A.; Welsh, J. W.; Roy, M. A.;

Lawrence,

[137238]D. A.; Lee, J.; Brush, J.; Taneyhill, L. A.; Deuel, B.; Lew, M.;

Watanabe,

[137239]C.; Cohen, R. L.; Melhem, M. F.; Finley, G. G.; Quirke, P.;

Goddard,

[137240]A. D.; Hillan, K. J.; Gurney, A. L.; Botstein, D.; Levine, A. J.:

[137241]WISP genes are members of the connective tissue growth
factor family

[137242]that are up-regulated in Wnt-1-transformed cells and
aberrantly expressed

[137243]in human colon tumors. Proc. Nat. Acad. Sci. 95:

14717-14722, 1998.

[137244]

[137245]

[137246]12579.

[137247]Tanaka, S.; Sugimachi, K.; Saeki, H.; Kinoshita, J.; Ohga,

T.;

[137248]Shimada, M.; Maehara, Y.; Sugimachi, K.: A novel variant
of WISP1

[137249]lacking a von Willebrand type C module overexpressed in
scirrhous

[137250]gastric carcinoma. *Oncogene* 20: 5525–5532, 2001.

[137251]

[137252]

[137253]12580.

[137254]Gee, S.; Krauss, S. W.; Miller, E.; Aoyagi, K.; Arenas, J.;
Conboy,

[137255]J. G.: Cloning of mDEAH9, a putative RNA helicase and
mammalian homologue

[137256]of *Saccharomyces cerevisiae* splicing factor Prp43. *Proc.*
Nat. Acad.

[137257]*Sci.* 94: 11803–11807, 1997.

[137258]

[137259]

[137260]12581.

[137261]Imamura, O.; Sugawara, M.; Furuichi, Y.: Cloning and
characterization

[137262]of a putative human RNA helicase gene of the DEAH-box
protein family. *Biochem.*

[137263]Biophys. Res. Commun. 240: 335–340, 1997.

[137264]

[137265]

[137266]12582.

[137267]Sano, K.; Tanihara, H.; Heimark, R. L.; Obata, S.; Davidson,
M.;

[137268]St. John, T.; Taketani, S.; Suzuki, S.: Protocadherins: a
large family

[137269]of cadherin–related molecules in central nervous system.
EMBO J. 12:

[137270]2249–2256, 1993.

[137271]

[137272]

[137273]12583.

[137274]Frigerio, J.–M.; Berthezene, P.; Garrido, P.; Ortiz, E.;
Barthelley,

[137275]S.; Vasseur, S.; Sastre, B.; Seleznieff, I.; Dagorn, J. C.; Io-
vanna,

[137276]J. L.: Analysis of 2166 clones from a human colorectal
cancer cDNA

[137277]library by partial sequencing. Hum. Molec. Genet. 4:
37–43, 1995.

[137278]

[137279]

[137280]12584.

[137281]Frigerio, J.–M.; Dagorn, J.–C.; Iovanna, J. L.: Cloning, se-
quencing

[137282]and expression of the L5, L21, L27a, L28, S5, S9, S10 and
S29 human

[137283]ribosomal protein mRNAs. Biochim. Biophys. Acta 1262:
64–68, 1995.

[137284]

[137285]

[137286]12585.

[137287]Eisenblatter, T.; Galla, H.–J.: A new multidrug resistance
protein

[137288]at the blood–brain barrier. Biochem. Biophys. Res. Com-
mun. 293:

[137289]1273–1278, 2002.

[137290]

[137291]

[137292]12586.

[137293]Miyake, K.; Mickley, L.; Litman, T.; Zhan, Z.; Robey, R.;
Cristensen,

[137294]B.; Brangi, M.; Greenberger, L.; Dean, M.; Fojo, T.; Bates, S.
E.

[137295]: Molecular cloning of cDNAs which are highly overex-
pressed in mitoxantrone-resistant

[137296]cells: demonstration of homology to ABC transport genes.
Cancer Res. 59:

[137297]8-13, 1999.

[137298]

[137299]

[137300]12587.

[137301]Ozvegy, C.; Litman, T.; Szakacs, G.; Nagy, Z.; Bates, S.;
Varadi,

[137302]A.; Sarkadi, B.: Functional characterization of the human
multidrug

[137303]transporter, ABCG2, expressed in insect cells. Biochem.
Biophys.

[137304]Res. Commun. 285: 111-117, 2001.

[137305]

[137306]

[137307]12588.

[137308]Dominguez, O.; Ashhab, Y.; Sabater, L.; Belloso, E.; Caro, P.;

[137309]Pujol-Borrell, R.: Cloning of ARE-containing genes by AU-motif-directed

[137310]display. Genomics 54: 278–286, 1998.

[137311]

[137312]

[137313]12589.

[137314]Kostrub, C. F.; Knudsen, K.; Subramani, S.; Enoch, T.: Hus1p,

[137315]a conserved fission yeast checkpoint protein, interacts with Rad1p

[137316]and is phosphorylated in response to DNA damage. EMBO J. 17: 2055–2066,

[137317]1998.

[137318]

[137319]

[137320]12590.

[137321]Lieberman, H. B.; Hopkins, K. M.; Nass, M.; Demetrick, D.;
Davey,

[137322]S.: A human homolog of the *Schizosaccharomyces pombe*
rad9+ checkpoint

[137323]control gene. *Proc. Nat. Acad. Sci.* 93: 13890–13895,
1996.

[137324]

[137325]

[137326]12591.

[137327]Katashima, R.; Iwahana, H.; Fujimura, M.; Yamaoka, T.;
Ishizuka,

[137328]T.; Tatibana, M.; Itakura, M.: Molecular cloning of a human
cDNA

[137329]for the 41-kDa phosphoribosylpyrophosphate synthetase–
associated protein. *Biochim.*

[137330]*Biophys. Acta* 1396: 245–250, 1998.

[137331]

[137332]

[137333]12592.

[137334]Katashima, R.; Iwahana, H.; Fujimura, M.; Yamaoka, T.;
Itakura,

[137335]M.: Assignment of the human phosphoribosylpyrophos-
phate synthetase-associated

[137336]protein 41 gene (PRPSAP2) to 17p11.2-p12. Genomics 54:
180-181,

[137337]1998.

[137338]

[137339]

[137340]12593.

[137341]Koike, N.; Hida, A.; Numano, R.; Hirose, M.; Sakaki, Y.; Tei,
H.

[137342]: Identification of the mammalian homologues of the
Drosophila timeless

[137343]gene, timeless1. FEBS Lett. 441: 427-431, 1998.

[137344]

[137345]

[137346]12594.

[137347]Sangoram, A. M.; Saez, L.; Antoch, M. P.; Gekakis, N.;
Staknis,

[137348]D.; Whiteley, A.; Fruechte, E. M.; Vitaterna, M. H.; Shimo-
mura, K.;

[137349]King, D. P.; Young, M. W.; Weitz, C. J.; Takahashi, J. S.:
Mammalian

[137350]circadian autoregulatory loop: a timeless ortholog and
mPer1 interact

[137351]and negatively regulate CLOCK-BMAL1-induced transcrip-
tion. Neuron 21:

[137352]1101-1113, 1998.

[137353]

[137354]

[137355]12595.

[137356]Zylka, M. J.; Shearman, L. P.; Levine, J. D.; Jin, X.; Weaver,

[137357]D. R.; Reppert, S. M.: Molecular analysis of mammalian
timeless. Neuron 21:

[137358]1115-1122, 1998.

[137359]

[137360]

[137361]12596.

[137362]Ardley, H. C.; Rose, S. A.; Tan, N.; Leek, J. P.; Markham, A.

F.;

[137363]Robinson, P. A.: Genomic organization of the human
ubiquitin-conjugating

[137364]enzyme gene, UBE2L6 on chromosome 11q12. Cytogenet.
Cell Genet. 89:

[137365]137–140, 2000.

[137366]

[137367]

[137368]12597.

[137369]Kumar, S.; Kao, W. H.; Howley, P. M.: Physical interaction
between

[137370]specific E2 and Hect E3 enzymes determines functional
cooperativity. J.

[137371]Biol. Chem. 272: 13548–13554, 1997.

[137372]

[137373]

[137374]12598.

[137375]tom Dieck, S.; Sanmarti-Vila, L.; Langnaese, K.; Richter, K.;
Kindler,

[137376]S.; Soyke, A.; Wex, H.; Smalla, K.-H.; Kampf, U.; Franzer,

J.-T.;

[137377]Stumm, M.; Garner, C. C.; Gundelfinger, E. D.: Bassoon, a novel zinc-finger

[137378]CAG/glutamine-repeat protein selectively localized at the active zone

[137379]of presynaptic nerve terminals. J. Cell Biol. 142: 499–509, 1998.

[137380]

[137381]

[137382]12599.

[137383]Winter, c.; tom Dieck, S.; Boeckers, T. M.; Bockmann, J.; Kampf,

[137384]U.; Sanmarti-Vila, L.; Langnaese, K.; Altroch, W.; Stumm, M.; Soyke,

[137385]A.; Wieacker, P.; Garner, C. C.; Gundelfinger, E. D.: The presynaptic

[137386]cytomatrix protein Bassoon: sequence and chromosomal localization

[137387]of the human BSN gene. Genomics 57: 389–397, 1999.

[137388]

[137389]

[137390]12600.

[137391]Bui, T. D.; Levy, E. R.; Subramaniam, V. N.; Lowe, S. L.;
Hong,

[137392]W.: cDNA characterization and chromosomal mapping of
human Golgi

[137393]SNARE GS27 and GS28 to chromosome 17. Genomics 57:
285–288, 1999.

[137394]

[137395]

[137396]12601.

[137397]Lowe, S. L.; Peter, F.; Subramaniam, V. N.; Wong, S. H.;
Hong,

[137398]W.: A SNARE involved in protein transport through the
Golgi apparatus. Nature 389:

[137399]881–884, 1997.

[137400]

[137401]

[137402]12602.

[137403]Nagahama, M.; Orci, L.; Ravazzola, M.; Amherdt, M.; La-
comis, L.;

[137404]Tempst, P.; Rothman, J. E.; Sollner, T. H.: A v-SNARE im-
plicated

[137405]in intra-Golgi transport. J. Cell Biol. 133: 507–516, 1996.

[137406]

[137407]

[137408]12603.

[137409]Subramaniam, V. N.; Peter, F.; Philp, R.; Wong, S. H.; Hong,
W.

[137410]: GS28, a 28-kilodalton Golgi SNARE that participates in
ER-Golgi

[137411]transport. Science 272: 1161–1163, 1996.

[137412]

[137413]

[137414]12604.

[137415]Hay, J. C.; Chao, D. S.; Kuo, C. S.; Scheller, R. H.: Protein

[137416]interactions regulating vesicle transport between the en-
doplasmic

[137417]reticulum and Golgi apparatus in mammalian cells. Cell
89: 149–158,

[137418]1997.

[137419]

[137420]

[137421]12605.

[137422]Hisamatsu, H.; Shimbara, N.; Saito, Y.; Kristensen, P.;
Hendil,

[137423]K. B.; Fujiwara, T.; Takahashi, E.; Tanahashi, N.; Tamura,
T.; Ichihara,

[137424]A.; Tanaka, K.: Newly identified pair of proteasomal sub-
units regulated

[137425]reciprocally by interferon gamma. J. Exp. Med. 183:
1807–1816, 1996.

[137426]

[137427]

[137428]12606.

[137429]Li, J.; Ding, S.-F.; Habib, N. A.; Fermor, B. F.; Wood, C. B.;

[137430]Gilmour, R. S.: Partial characterization of a cDNA for hu-
man stearyl-CoA

[137431]desaturase and changes in its mRNA expression in some
normal and malignant

[137432]tissues. Int. J. Cancer 57: 348–352, 1994.

[137433]

[137434]

[137435]12607.

[137436]Ntambi, J. M.; Miyazaki, M.; Stoehr, J. P.; Lan, H.;

Kendzierski,

[137437]C. M.; Yandell, B. S.; Song, Y.; Cohen, P.; Friedman, J. M.;

Attie,

[137438]A. D.: Loss of stearyl-CoA desaturase-1 function protects mice against

[137439]adiposity. Proc. Nat. Acad. Sci. 99: 11482-11486, 2002.

[137440]

[137441]

[137442]12608.

[137443]Thiede, M. A.; Ozols, J.; Strittmatter, P.: Construction and sequence

[137444]of cDNA for rat liver stearyl coenzyme A desaturase. J.

Biol. Chem. 261:

[137445]13230-13235, 1986.

[137446]

[137447]

[137448]12609.

[137449]Zhang, L.; Ge, L.; Parimoo, S.; Stenn, K.; Prouty, S. M.: Human
man

[137450]stearoyl-CoA desaturase: alternative transcripts generated
from a

[137451]single gene by usage of tandem polyadenylation sites.

Biochem. J. 340:

[137452]255–264, 1999.

[137453]

[137454]

[137455]12610.

[137456]Zheng, Y.; Eilertsen, K. J.; Ge, L.; Zhang, L.; Sundberg, J. P.;

[137457]Prouty, S. M.; Stenn, K. S.; Parimoo, S.: Scd1 is expressed
in sebaceous

[137458]glands and is disrupted in the asebia mouse. (Letter) Na-
ture Genet. 23:

[137459]268–270, 1999.

[137460]

[137461]

[137462]12611.

[137463]Duh, F.-M.; Latif, F.; Weng, Y.; Geil, L.; Modi, W.; Stackhouse,

[137464]T.; Matsumura, F.; Duan, D. R.; Linehan, W. M.; Lerman, M. I.; Gnarra,

[137465]J. R.: cDNA cloning and expression of the human homolog of the sea

[137466]urchin fascin and Drosophila singed genes which encodes an actin-bundling

[137467]protein. DNA Cell Biol. 13: 821-827, 1994.

[137468]

[137469]

[137470]12612.

[137471]Mosialos, G.; Birkenbach, M.; Ayehunie, S.; Matsumura, F.; Pinkus,

[137472]G. S.; Kieff, E.; Langhoff, E.: Circulating human dendritic cells

[137473]differentially express high levels of a 55-kd actin-bundling protein. Am.

[137474]J. Path. 148: 593-600, 1996.

[137475]

[137476]

[137477]12613.

[137478]Ono, S.; Yamakita, Y.; Yamashiro, S.; Matsudaira, P. T.;

Gnarra,

[137479]J. R.; Obinata, T.; Matsumura, F.: Identification of an actin binding

[137480]region and a protein kinase C phosphorylation site on human fascin. J.

[137481]Biol. Chem. 272: 2527–2533, 1997.

[137482]

[137483]

[137484]12614.

[137485]Pinkus, G. S.; Pinkus, J. L.; Langhoff, E.; Matsumura, F.;

Yamashiro,

[137486]S.; Mosialos, G.; Said, J. W.: Fascin, a sensitive new marker for

[137487]Reed–Sternberg cells of Hodgkin's disease: evidence for a dendritic

[137488]or B cell derivation? Am. J. Path. 150: 543–562, 1997.

[137489]

[137490]

[137491]12615.

[137492]Sonderbye, L.; Magerstadt, R.; Blatman, R. N.; Preffer, F. I.;

[137493]Langhoff, E.: Selective expression of human fascin (p55)
by dendritic

[137494]leukocytes. Adv. Exp. Med. Biol. 471: 41–46, 1997.

[137495]

[137496]

[137497]12616.

[137498]Yamakita, Y.; Ono, S.; Matsumura, F.; Yamashiro, S.: Phos-
phorylation

[137499]of human fascin inhibits its actin binding and bundling
activities. J.

[137500]Biol. Chem. 271: 12632–12638, 1996.

[137501]

[137502]

[137503]12617.

[137504]Yamashiro–Matsumura, S.; Matsumura, F.: Intracellular lo-
calization

[137505]of the 55–kD actin–bundling protein in cultured cells:

spatial relationships

[137506]with actin, alpha-actinin, tropomyosin, and fimbrin. J. Cell Biol. 103:

[137507]631–640, 1986.

[137508]

[137509]

[137510]12618.

[137511]Yamashiro–Matsumura, S.; Matsumura, F.: Purification and characterization

[137512]of an F-actin-bundling 55-kilodalton protein from HeLa cells. J.

[137513]Biol. Chem. 260: 5087–5097, 1985.

[137514]

[137515]

[137516]12619.

[137517]Masson, J.–Y.; Stasiak, A. Z.; Stasiak, A.; Benson, F. E.; West,

[137518]S. C.: Complex formation by the human RAD51C and XRCC3 recombination

[137519]repair proteins. Proc. Nat. Acad. Sci. 98: 8440–8446,

2001.

[137520]

[137521]

[137522]12620.

[137523]Selfors, L. M.; Schutzman, J. L.; Borland, C. Z.; Stern, M. J.

[137524]: Soc-2 encodes a leucine-rich repeat protein implicated
in fibroblast

[137525]growth factor receptor signaling. Proc. Nat. Acad. Sci. 95:
6903-6908,

[137526]1998.

[137527]

[137528]

[137529]12621.

[137530]Sieburth, D. S.; Sun, Q.; Han, M.: SUR-8, a conserved Ras-
binding

[137531]protein with leucine-rich repeats, positively regulates
Ras-mediated

[137532]signaling in *C. elegans*. Cell 94: 119-130, 1998.

[137533]

[137534]

[137535]12622.

[137536]Gibbs, P. E. M.; McGregor, W. G.; Maher, V. M.; Nisson, P.;
Lawrence,

[137537]C. W.: A human homolog of the *Saccharomyces cerevisiae*
REV3 gene,

[137538]which encodes the catalytic subunit of DNA polymerase
zeta. Proc.

[137539]Nat. Acad. Sci. 95: 6876–6880, 1998.

[137540]

[137541]

[137542]12623.

[137543]Morelli, C.; Mungall, A. J.; Negrini, M.; Barbanti-Brodano,
G.;

[137544]Croce, C. M.: Alternative splicing, genomic structure, and
fine chromosome

[137545]localization of REV3L. Cytogenet. Cell Genet. 83: 18–20,
1998.

[137546]

[137547]

[137548]12624.

[137549]Xiao, W.; Lechler, T.; Chow, B. L.; Fontanie, T.; Agustus, M.;

[137550]Carter, K. C.; Wei, Y.-F.: Identification, chromosomal mapping and

[137551]tissue-specific expression of hREV3 encoding a putative human DNA

[137552]polymerase zeta. Carcinogenesis 19: 945-949, 1998.

[137553]

[137554]

[137555]12625.

[137556]Kahn, M. L.; Zheng, Y.-W.; Huang, W.; Bigornia, V.; Zeng, D.; Moff,

[137557]S.; Farese, R. V., Jr.; Tam, C.; Coughlin, S. R.: A dual thrombin

[137558]receptor system for platelet activation. Nature 394: 690-694, 1998.

[137559]

[137560]

[137561]12626.

[137562]Xu, W.-F.; Andersen, H.; Whitmore, T. E.; Presnell, S. R.;
Yee,

[137563]D. P.; Ching, A.; Gilbert, T.; Davie, E. W.; Foster, D. C.:
Cloning

[137564]and characterization of human protease-activated recep-
tor 4. Proc.

[137565]Nat. Acad. Sci. 95: 6642-6646, 1998.

[137566]

[137567]

[137568]12627.

[137569]Santoro, B.; Grant, S. G. N.; Bartsch, D.; Kandel, E. R.: In-
teractive

[137570]cloning with the SH3 domain of N-src identifies a new
brain specific

[137571]ion channel protein, with homology to Eag and cyclic nu-
cleotide-gated

[137572]channels. Proc. Nat. Acad. Sci. 94: 14815-14820, 1997.

[137573]

[137574]

[137575]12628.

[137576]Santoro, B.; Liu, D. T.; Yao, H.; Bartsch, D.; Kandel, E. R.;
Siegelbaum,

[137577]S. A.; Tibbs, G. R.: Identification of a gene encoding a hy-
perpolarization-activated

[137578]pacemaker channel of brain. Cell 93: 717–729, 1998.

[137579]

[137580]

[137581]12629.

[137582]Stevens, D. R.; Seifert, R.; Bufe, B.; Muller, F.; Kremmer, E.;

[137583]Gauss, R.; Meyerhof, W.; Kaupp, U. B.; Lindemann, B.: Hy-
perpolarization-activated

[137584]channels HCN1 and HCN4 mediate responses to sour
stimuli. Nature 413:

[137585]631–635, 2001.

[137586]

[137587]

[137588]12630.

[137589]Wainger, B. J.; DeGennaro, M.; Santoro, B.; Siegelbaum, S.
A.;

[137590]Tibbs, G. R.: Molecular mechanism of cAMP modulation of
HCN pacemaker

[137591]channels. Nature 411: 805–810, 2001.

[137592]

[137593]

[137594]12631.

[137595]Ludwig, A.; Zong, X.; Stieber, J.; Hullin, R.; Hofmann, F.;
Biel,

[137596]M.: Two pacemaker channels from human heart with pro-
foundly different

[137597]activation kinetics. EMBO J. 18: 2323–2329, 1999.

[137598]

[137599]

[137600]12632.

[137601]Casari, G.; De Fusco, M.; Ciarmatori, S.; Zeviani, M.; Mora, M.;

[137602]Fernandez, P.; De Michele, G.; Filla, A.; Coccozza, S.; Marconi, R.;

[137603]Durr, A.; Fontaine, B.; Ballabio, A.: Spastic paraplegia and OXPHOS

[137604]impairment caused by mutations in paraplegin, a nuclear-encoded mitochondrial

[137605]metalloprotease. Cell 93: 973–983, 1998.

[137606]

[137607]

[137608]12633.

[137609]De Michele, G.; De Fusco, M.; Cavalcanti, F.; Filla, A.; Marconi,

[137610]R.; Volpe, G.; Monticelli, A.; Ballabio, A.; Casari, G.; Coccozza,

[137611]S.: A new locus for autosomal recessive hereditary spastic paraplegia

[137612]maps to chromosome 16q24.3. Am. J. Hum. Genet. 63:

135–139, 1998.

[137613]

[137614]

[137615]12634.

[137616]Settasatian, C.; Whitmore, S. A.; Crawford, J.; Bilton, R. L.;

[137617]Cleton-Jansen, A.-M.; Sutherland, G. R.; Callen, D. F.: Ge-
nomic structure

[137618]and expression analysis of the spastic paraplegia gene,
SPG7. Hum.

[137619]Genet. 105: 139–144, 1999.

[137620]

[137621]

[137622]12635.

[137623]de Lecea, L.; Criado, J. R.; Prospero-Garcia, O.; Gautvik, K.
M.;

[137624]Schweitzer, P.; Danielson, P. E.; Dunlop, C. L. M.; Siggins,
G. R.;

[137625]Henriksen, S. J.; Sutcliffe, J. G.: A cortical neuropeptide
with

[137626]neuronal depressant and sleep-modulating properties.

Nature 381:

[137627]242–245, 1996.

[137628]

[137629]

[137630]12636.

[137631]Kerjaschki, D.; Farquhar, M. G.: Immunocytochemical localization

[137632]of the Heymann nephritis antigen (GP330) in glomerular epithelial

[137633]cells of normal Lewis rats. J. Exp. Med. 157: 667–686, 1983.

[137634]

[137635]

[137636]12637.

[137637]Korenberg, J. R.; Argraves, K. M.; Chen, X.–N.; Tran, H.; Strickland,

[137638]D.; Argraves, W. S.: Chromosomal localization of human genes for

[137639]the LDL receptor family member glycoprotein 330 (LRP2) and its associated

[137640]protein RAP (LRPAP1). Genomics 22: 88–93, 1994.

[137641]

[137642]

[137643]12638.

[137644]Kounnas, M. Z.; Loukinova, E. B.; Stefansson, S.; Harmony, J. A.

[137645]K.; Brewer, B. H.; Strickland, D. K.; Argraves, W. S.: Identification

[137646]of glycoprotein 330 as an endocytic receptor for apolipoprotein J/clusterin. J.

[137647]Biol. Chem. 270: 13070–13075, 1995.

[137648]

[137649]

[137650]12639.

[137651]Leheste, J.–R.; Rolinski, B.; Vorum, H.; Hilpert, J.; Nykjaer,

[137652]A.; Jacobsen, C.; Aucouturier, P.; Moskaug, J. O.; Otto, A.; Christensen,

[137653]E. I.; Willnow, T. E.: Megalin knockout mice as an animal model of

[137654]low molecular weight proteinuria. Am. J. Path. 155: 1361–1370, 1999.

[137655]

[137656]

[137657]12640.

[137658]Marino, M.; Chiovato, L.; Friedlander, J. A.; Latrofa, F.;
Pinchera,

[137659]A.; McCluskey, R. T.: Serum antibodies against megalin
(GP330) in

[137660]patients with autoimmune thyroiditis. J. Clin. Endocr.
Metab. 84:

[137661]2468–2474, 1999.

[137662]

[137663]

[137664]12641.

[137665]Moestrup, S. K.; Cui, S.; Vorum, H.; Bregengard, C.; Bjorn,
S.

[137666]E.; Norris, K.; Gliemann, J.; Christensen, E. I.: Evidence that
epithelial

[137667]glycoprotein 330/megalin mediates uptake of polybasic
drugs. J. Clin.

[137668]Invest. 96: 1404–1413, 1995.

[137669]

[137670]

[137671]12642.

[137672]Nykjaer, A.; Dragun, D.; Walther, D.; Vorum, H.; Jacobsen, C.;

[137673]Herz, J.: Melsen, F.; Christensen, E. I.; Willnow, T. E.: An endocytic

[137674]pathway essential for renal uptake and activation of the steroid 25-(OH)

[137675]vitamin D3. Cell 96: 507-515, 1999.

[137676]

[137677]

[137678]12643.

[137679]Raychowdhury, R.; Niles, J. L.; McCluskey, R. T.; Smith, J. A.

[137680]: Autoimmune target in Heymann nephritis is a glycoprotein with homology

[137681]to the LDL receptor. Science 244: 1163-1165, 1989.

[137682]

[137683]

[137684]12644.

[137685]Saito, A.; Pietromonaco, S.; Loo, A. K.-C.; Farquhar, M. G.:

[137686]Complete cloning and sequencing of rat gp330/`megalin,'
a distinctive

[137687]member of the low density lipoprotein receptor gene fam-
ily. Proc.

[137688]Nat. Acad. Sci. 91: 9725-9729, 1994.

[137689]

[137690]

[137691]12645.

[137692]Schmitz, C.; Hilpert, J.; Jacobsen, C.; Boensch, C.; Chris-
tensen,

[137693]E. I.; Luft, F. C.; Willnow, T. E.: Megalin deficiency offers
protection

[137694]from renal aminoglycoside accumulation. J. Biol. Chem.
277: 618-622,

[137695]2002.

[137696]

[137697]

[137698]12646.

[137699]Willnow, T. E.; Hilpert, J.; Armstrong, S. A.; Rohlmann, A.;
Hammer,

[137700]R. E.; Burns, D. K.; Herz, J.: Defective forebrain develop-
ment in

[137701]mice lacking gp330/megalin. Proc. Nat. Acad. Sci. 93:
8460–8464,

[137702]1996.

[137703]

[137704]

[137705]12647.

[137706]Gostout, B.; Liu, Q.; Sommer, S. S.: 'Cryptic' repeating
triplets

[137707]of purines and pyrimidines (cRRY(i)) are frequent and
polymorphic:

[137708]analysis of coding cRRY(i) in the proopiomelanocortin
(POMC) and TATA-binding

[137709]protein (TBP) genes. Am. J. Hum. Genet. 52: 1182–1190,
1993.

[137710]

[137711]

[137712]12648.

[137713]Hobbs, N. K.; Bondareva, A. A.; Barnett, S.; Capecchi, M. R.; Schmidt,

[137714]E. E.: Removing the vertebrate-specific TBP N terminus disrupts placental

[137715]beta-2M-dependent interactions with the maternal immune system. Cell 110:

[137716]43-54, 2002.

[137717]

[137718]

[137719]12649.

[137720]Imbert, G.; Trottier, Y.; Beckmann, J.; Mandel, J. L.: The gene

[137721]for the TATA binding protein (TBP) that contains a highly polymorphic

[137722]protein coding CAG repeat maps to 6q27. Genomics 21: 667-668, 1994.

[137723]

[137724]

[137725]12650.

[137726]Jones, A. L.; Middle, F.; Guy, C.; Spurlock, G.; Cairns, N. J.;

[137727]McGuffin, P.; Craddock, N.; Owen, M.; O'Donovan, M. C.:

No evidence

[137728]for expanded polyglutamine sequences in bipolar disorder
and schizophrenia. Molec.

[137729]Psychiat. 2: 478–482, 1997.

[137730]

[137731]

[137732]12651.

[137733]Kao, C. C.; Lieberman, P. M.; Schmidt, M. C.; Zhou, Q.; Pei,
R.;

[137734]Berk, A. J.: Cloning of a transcriptionally active human
TATA binding

[137735]factor. Science 248: 1646–1650, 1990.

[137736]

[137737]

[137738]12652.

[137739]Koide, R.; Kobayashi, S.; Shimohata, T.; Ikeuchi, T.;
Maruyama,

[137740]M.; Saito, M.; Yamada, M.; Takahashi, H.; Tsuji, S.: A neu-
rological

[137741]disease caused by an expanded CAG trinucleotide repeat
in the TATA-binding

[137742]protein gene: a new polyglutamine disease? Hum. Molec.
Genet. 8:

[137743]2047–2053, 1999.

[137744]

[137745]

[137746]12653.

[137747]Nakamura, K.; Jeong, S.-Y.; Uchihara, T.; Anno, M.; Na-
gashima,

[137748]K.; Nagashima, T.; Ikeda, S.; Tsuji, S.; Kanazawa, I.:
SCA17, a novel

[137749]autosomal dominant cerebellar ataxia caused by an ex-
panded polyglutamine

[137750]in TATA-binding protein. Hum. Molec. Genet. 10:
1441–1448, 2001.

[137751]

[137752]

[137753]12654.

[137754]Nikolov, D. B.; Hu, S.-H.; Lin, J.; Gasch, A.; Hoffmann, A.;
Horikoshi,

[137755]M.; Chua, N.-H.; Roeder, R. G.; Burley, S. K.: Crystal structure

[137756]of TFIID TATA-box binding protein. Nature 360: 40-46, 1992.

[137757]

[137758]

[137759]12655.

[137760]Peterson, M. G.; Tanese, N.; Pugh, B. F.; Tjian, R.: Functional

[137761]domains and upstream activation properties of cloned human TATA binding

[137762]protein. Science 248: 1625-1630, 1990.

[137763]

[137764]

[137765]12656.

[137766]Polymeropoulos, M. H.; Rath, D. S.; Xiao, H.; Merrill, C. R.:

[137767]Trinucleotide repeat polymorphism at the human transcription factor

[137768]IID gene. Nucleic Acids Res. 19: 4307 only, 1991.

[137769]

[137770]

[137771]12657.

[137772]Rosen, D. R.; Trofatter, J. A.; Brown, R. H., Jr.: Mapping of
[137773]the human TATA-binding protein gene (TBP) to chromo-
some 6qter. Cytogenet.

[137774]Cell Genet. 69: 279–280, 1995.

[137775]

[137776]

[137777]12658.

[137778]Rubinsztein, D. C.; Leggo, J.; Crow, T. J.; DeLisi, L. E.;
Walsh,

[137779]C.; Jain, S.; Paykel, E. S.: Analysis of polyglutamine-coding
repeats

[137780]in the TATA-binding protein in different human popula-
tions and in

[137781]patients with schizophrenia and bipolar affective disorder.
Am. J.

[137782]Med. Genet. 67: 495–498, 1996.

[137783]

[137784]

[137785]12659.

[137786]Saito, F.; Yamamoto, T.; Horikoshi, M.; Ikeuchi, T.: Direct mapping

[137787]of the human TATA box-binding protein (TBP) gene to 6q27 by fluorescence

[137788]in situ hybridization. Jpn. J. Hum. Genet. 39: 421-425, 1994.

[137789]

[137790]

[137791]12660.

[137792]Serrano, M.; Hannon, G. J.; Beach, D.: A new regulatory motif

[137793]in cell-cycle control causing specific inhibition of cyclin D/CDK4. Nature 366:

[137794]704-707, 1993.

[137795]

[137796]

[137797]12661.

[137798]Testa, J. R.; Zhou, J.; Bell, D. W.; Yen, T. J.: Chromosomal localization

[137799]of the genes encoding the kinetochore proteins CENPE
and CENPF to

[137800]human chromosomes 4q24–q25 and 1q32–q41, respec-
tively, by fluorescence

[137801]in situ hybridization. Genomics 23: 691–693, 1994.

[137802]

[137803]

[137804]12662.

[137805]Pennica, D.; Swanson, T. A.; Shaw, K. J.; Kuang, W.–J.;
Gray, C.

[137806]L.; Beatty, B. G.; Wood, W. I.: Human cardiotrophin–1: pro-
tein and

[137807]gene structure, biological and binding activities, and
chromosomal

[137808]localization. Cytokine 8: 183–189, 1996.

[137809]

[137810]

[137811]12663.

[137812]Fukudome, K.; Esmon, C. T.: Identification, cloning, and
regulation

[137813]of a novel endothelial cell protein C/activated protein C
receptor. J.

[137814]Biol. Chem. 269: 26486–26491, 1994.

[137815]

[137816]

[137817]12664.

[137818]Hayashi, T.; Nakamura, H.; Okada, A.; Takebayashi, S.;
Wakita,

[137819]T.; Yuasa, H.; Okumura, K.; Suzuki, K.: Organization and
chromosomal

[137820]localization of the human endothelial protein C receptor
gene. Gene 238:

[137821]367–373, 1999.

[137822]

[137823]

[137824]12665.

[137825]Mincheva, A.; Rothbarth, K.; Werner, D.; Lichter, P.: As-
signment

[137826]of the gene encoding centrosome-associated protein
CCD41 to mouse

[137827]chromosome 2H. Mammalian Genome 6: 444 only, 1995.

[137828]

[137829]

[137830]12666.

[137831]Rothbarth, K.; Dabaghian, A. R. H.; Stammer, H.; Werner, D.: One

[137832]single mRNA encodes the centrosomal protein CCD41 and the endothelial

[137833]cell protein C receptor (EPCR). FEBS Lett. 458: 77–80, 1999.

[137834]

[137835]

[137836]12667.

[137837]Rothbarth, K.; Petzelt, C.; Lu, X.; Todorov, I. T.; Joswig, G.;

[137838]Pepperkok, R.; Ansorge, W.; Werner, D.: cDNA–derived molecular characteristics

[137839]and antibodies to a new centrosome–associated and G2/M phase–prevalent

[137840]protein. J. Cell Sci. 104: 19–30, 1993.

[137841]

[137842]

[137843]12668.

[137844]Simmonds, R. E.; Lane, D. A.: Structural and functional im-
plications

[137845]of the intron/exon organization of the human endothelial
cell protein

[137846]C/activated protein C receptor (EPCR) gene: comparison
with the structure

[137847]of CD1/major histocompatibility complex alpha-1 and al-
pha-2 domains. Blood 94:

[137848]632-641, 1999.

[137849]

[137850]

[137851]12669.

[137852]Ye, X.; Fukudome, K.; Tsuneyoshi, N.; Satoh, T.; Toku-
naga, O.;

[137853]Sugawara, K.; Mizokami, H.; Kimoto, M.: The endothelial
cell protein

[137854]C receptor (EPCR) functions as a primary receptor for pro-
tein C activation

[137855]on endothelial cells in arteries, veins, and capillaries.

Biochem.

[137856]Biophys. Res. Commun. 259: 671–677, 1999.

[137857]

[137858]

[137859]12670.

[137860]Purrello, M.; Pietro, C. D.; Mirabile, E.; Rapisarda, A.; Rimini,

[137861]R.; Tine, A.; Pavone, L.; Motta, S.; Grzeschik, K.–H.; Sichel, G.

[137862]: Physical mapping at 6q27 of the locus for the TATA box-binding protein,

[137863]the DNA-binding subunit of TFIID and a component of SL1 and TFIIIB,

[137864]strongly suggests that it is single copy in the human genome. Genomics 22:

[137865]94–100, 1994.

[137866]

[137867]

[137868]12671.

[137869]Candia, A. F.; Hu, J.; Crosby, J.; Lalley, P. A.; Noden, D.;

Nadeau,

[137870]J. H.; Wright, C. V. E.: Mox-1 and Mox-2 define a novel homeobox

[137871]gene subfamily and are differentially expressed during early mesodermal

[137872]patterning in mouse embryos. Development 116: 1123-1136, 1992.

[137873]

[137874]

[137875]12672.

[137876]Gorski, D. H.; LePage, D. F.; Patel, C. V.; Copeland, N. G.; Jenkins,

[137877]N. A.; Walsh, K.: Molecular cloning of a diverged homeobox gene that

[137878]is rapidly down-regulated during the G0/G1 transition in vascular

[137879]smooth muscle cells. Molec. Cell Biol. 13: 3722-3733, 1993.

[137880]

[137881]

[137882]12673.

[137883]Grigoriou, M.; Kastrinaki, M.-C.; Modi, W. S.; Theodorakis, K.;

[137884]Mankoo, B.; Pachnis, V.; Karagogeos, D.: Isolation of the human MOX2

[137885]homeobox gene and localization to chromosome 7p22.1-p21.3. Genomics 26:

[137886]550-555, 1995.

[137887]

[137888]

[137889]12674.

[137890]LePage, D. F.; Altomare, D. A.; Testa, J. R.; Walsh, K.: Molecular

[137891]cloning and localization of the human GAX gene to 7p21. Genomics 24:

[137892]535-540, 1994.

[137893]

[137894]

[137895]12675.

[137896]Mankoo, B. S.; Collins, N. S.; Ashby, P.; Grigorievea, E.; Pevny,

[137897]L. H.; Candia, A.; Wright, C. V. E.; Rigby, P. W. J.; Pachnis, V.

[137898]: Mox2 is a component of the genetic hierarchy controlling limb muscle

[137899]development. Nature 400: 69–73, 1999.

[137900]

[137901]

[137902]12676.

[137903]Lindor, N. M.; Furuichi, Y.; Kitao, S.; Shimamoto, A.; Arndt, C.;

[137904]Jalal, S.: Rothmund–Thomson syndrome due to RECQ4 helicase mutations:

[137905]report and clinical and molecular comparisons with Bloom syndrome

[137906]and Werner syndrome. Am. J. Med. Genet. 90: 223–228, 2000.

[137907]

[137908]

[137909]12677.

[137910]Puranam, K. L.; Blackshear, P. J.: Cloning and characterization

[137911]of RECQL, a potential human homologue of the Escherichia coli DNA

[137912]helicase RecQ. J. Biol. Chem. 269: 29838–29845, 1994.

[137913]

[137914]

[137915]12678.

[137916]Puranam, K. L.; Kennington, E.; J.–Sait, S. N.; Shows, T. B.; Rochelle,

[137917]J. M.; Seldin, M. F.; Blackshear, P. J.: Chromosomal localization

[137918]of the gene encoding the human DNA helicase RECQL and its mouse homologue. Genomics 26:

[137919]595–598, 1995.

[137920]

[137921]

[137922]12679.

[137923]Gollner, H.; Bouwman, P.; Mangold, M.; Karis, A.; Braun, H.; Rohner,

[137924]J.; Del Rey, A.; Besedovsky, H.–O.; Meinhardt, A.; van den Broek,

[137925]M.; Cutforth, T.; Grosveld, F.; Philipsen, S.; Suske, G.:

Complex

[137926]phenotype of mice homozygous for a null mutation in the

Sp4 transcription

[137927]factor gene. Genes Cells 6: 689–697, 2001.

[137928]

[137929]

[137930]12680.

[137931]Hagen, G.; Dennig, J.; Preiss, M.; Beato, M.; Suske, G.:

Functional

[137932]analyses of the transcription factor Sp4 reveal properties

distinct

[137933]from Sp1 and Sp3. J. Biol. Chem. 270: 24989–24994,

1995.

[137934]

[137935]

[137936]12681.

[137937]Hagen, G.; Muller, S.; Beato, M.; Suske, G.: Cloning by

recognition

[137938]site screening of two novel GT box binding proteins: a

family of Sp1

[137939]related genes. Nucleic Acids Res. 20: 5519–5525, 1992.

[137940]

[137941]

[137942]12682.

[137943]Kalff–Suske, M.; Kunz, J.; Grzeschik, K.–H.; Suske, G.: Human
man

[137944]Sp4 transcription factor gene (SP4) maps to chromosome
7p15. Genomics 26:

[137945]631–633, 1995.

[137946]

[137947]

[137948]12683.

[137949]Nguyen–Tran, V. T. B.; Kubalak, S. W.; Minamisawa, S.;
Fiset, C.;

[137950]Wollert, K. C.; Brown, A. B.; Ruiz–Lozano, P.; Barrere–
Lemaire, S.;

[137951]Kondo, R.; Norman, L. W.; Gourdie, R. G.; Rahme, M. M.;
Feld, G. K.;

[137952]Clark, R. B.; Giles, W. R.; Chien, K. R.: A novel genetic
pathway

[137953]for sudden cardiac death via defects in the transition between ventricular

[137954]and conduction system cell lineages. Cell 102: 671–682, 2000.

[137955]

[137956]

[137957]12684.

[137958]Lamour, V.; Lecluse, Y.; Desmaze, C.; Spector, M.; Bodescot, M.;

[137959]Aurias, A.; Osley, M. A.; Lipinski, M.: A human homolog of the S.

[137960]cerevisiae HIR1 and HIR2 transcriptional repressors cloned from the

[137961]DiGeorge syndrome critical region. Hum. Molec. Genet. 4: 791–799,

[137962]1995.

[137963]

[137964]

[137965]12685.

[137966]Llevadot, R.; Scambler, P.; Estivill, X.; Pritchard, M.: Genomic

[137967]organization of TUPLE1/HIRA: a gene implicated in DiGeorge syndrome. Mammalian

[137968]Genome 7: 911–914, 1996.

[137969]

[137970]

[137971]12686.

[137972]Lorain, S.; Quivy, J.-P.; Monier-Gavelle, F.; Scamps, C.;
Lecluse,

[137973]Y.; Almouzni, G.; Lipinski, M.: Core histones and HIRIP3, a
novel

[137974]histone-binding protein, directly interact with WD repeat
protein

[137975]HIRA. Molec. Cell. Biol. 18: 5546–5556, 1998.

[137976]

[137977]

[137978]12687.

[137979]Magnaghi, P.; Roberts, C.; Lorain, S.; Lipinski, M.; Scambler,

[137980]P. J.: HIRA, a mammalian homologue of *Saccharomyces cerevisiae* transcriptional

[137981]co-repressors, interacts with Pax3. Nature Genet. 20:
74–77, 1998.

[137982]

[137983]

[137984]12688.

[137985]Mattei, M.–G.; Halford, S.; Scambler, P. J.: Mapping of the
Tuple1

[137986]gene to mouse chromosome 16A–B1. Genomics 23:
717–718, 1994.

[137987]

[137988]

[137989]12689.

[137990]Roberts, C.; Daw, S. C. M.; Halford, S.; Scambler, P. J.:
Cloning

[137991]and developmental expression analysis of chick Hira
(Chira), a candidate

[137992]gene for DiGeorge syndrome. Hum. Molec. Genet. 6:
237–245, 1997.

[137993]

[137994]

[137995]12690.

[137996]Wilming, L. G.; Snoeren, C. A. S.; van Rijswijk, A.;

Grosveld,

[137997]F.; Meijers, C.: The murine homologue of HIRA, a DiGeorge syndrome

[137998]candidate gene, is expressed in embryonic structures affected in human

[137999]CATCH22 patients. Hum. Molec. Genet. 6: 247-258, 1997.

[138000]

[138001]

[138002]12691.

[138003]Kim, I.-G.; Gorman, J. J.; Park, S.-C.; Chung, S.-I.; Steinert,

[138004]P. M.: The deduced sequence of the novel protransglutaminase E (TGase3)

[138005]of human and mouse. J. Biol. Chem. 268: 12682-12690, 1993.

[138006]

[138007]

[138008]12692.

[138009]Kim, I.-G.; Lee, S.-C.; Lee, J.-H.; Yang, J.-M.; Chung, S.-I.;

[138010]Steinert, P. M.: Structure and organization of the human transglutaminase

[138011]3 gene: evolutionary relationship to the transglutaminase family. J.

[138012]Invest. Derm. 103: 137–142, 1994.

[138013]

[138014]

[138015]12693.

[138016]Marchese, A.; Docherty, J. M.; Nguyen, T.; Heiber, M.; Cheng, R.;

[138017]Heng, H. H. Q.; Tsui, L.-C.; Shi, X.; George, S. R.; O'Dowd, B. F.

[138018]: Cloning of human genes encoding novel G protein-coupled receptors. Genomics 23:

[138019]609–618, 1994.

[138020]

[138021]

[138022]12694.

[138023]Homey, B.; Alenius, H.; Muller, A.; Soto, H.; Bowman, E. P.; Yuan,

[138024]W.; McEvoy, L.; Lauerma, A. I.; Assmann, T.; Bunemann, E.;
Lehto,
[138025]M.; Wolff, H.; Yen, D.; Marxhausen, H.; To, W.; Sedgwick,
J.; Ruzicka,
[138026]T.; Lehmann, P.; Zlotnik, A.: CCL27–CCR10 interactions
regulate T
[138027]cell–mediated skin inflammation. Nature Med. 8:
157–165, 2002.
[138028]
[138029]
[138030]12695.
[138031]Homey, B.; Wang, W.; Soto, H.; Buchanan, M. E.; Wiesen-
born, A.;
[138032]Catron, D.; Muller, A.; McClanahan, T. K.; Dieu–Nosjean,
M.–C.; Orozco,
[138033]R.; Ruzicka, T.; Lehmann, P.; Oldham, E.; Zlotnik, A.: Cut-
ting edge:
[138034]the orphan chemokine receptor G protein–coupled recep-
tor–2 (GPR–2,
[138035]CCR10) binds the skin–associated chemokine CCL27
(CTACK/ALP/ILC). J.
[138036]Immun. 164: 3465–3470, 2000.

[138037]

[138038]

[138039]12696.

[138040]Jarmin, D. I.; Rits, M.; Bota, D.; Gerard, N. P.; Graham, G. J.;

[138041]Clark-Lewis, I.; Gerard, C.: Cutting edge: identification of the

[138042]orphan receptor G-protein-coupled receptor 2 as CCR10, a specific

[138043]receptor for the chemokine ESKine. J. Immun. 164: 3460-3464, 2000.

[138044]

[138045]

[138046]12697.

[138047]Wang, W.; Soto, H.; Oldham, E. R.; Buchanan, M. E.; Homey, B.;

[138048]Catron, D.; Jenkins, N.; Copeland, N. G.; Gilbert, D. J.; Nguyen,

[138049]N.; Abrams, J.; Kershenovich, D.; Smith, K.; McClanahan, T.; Vicari,

[138050]A. P.; Zlotnik, A.: Identification of a novel chemokine

(CCL28),

[138051]which binds CCR10 (GPR2). J. Biol. Chem. 275:
22313–22323, 2000.

[138052]

[138053]

[138054]12698.

[138055]Akao, Y.; Matsuda, Y.: Identification and chromosome
mapping of

[138056]the mouse homologue of the human gene (DDX6) that en-
codes a putative

[138057]RNA helicase of the DEAD box protein family. Cytogenet.
Cell Genet. 75:

[138058]38–44, 1996.

[138059]

[138060]

[138061]12699.

[138062]Akao, Y.; Seto, M.; Takahashi, T.; Kubonishi, I.; Miyoshi, I.;

[138063]Nakazawa, S.; Tsujimoto, Y.; Croce, C. M.; Ueda, R.: Molec-
ular cloning

[138064]of the chromosomal breakpoint of a B-cell lymphoma with

the t(11;14)(q23;q32)

[138065]chromosome translocation. Cancer Res. 51: 1574–1576,
1991.

[138066]

[138067]

[138068]12700.

[138069]Akao, Y.; Seto, M.; Yamamoto, K.; Iida, S.; Nakazawa, S.;
Inazawa,

[138070]J.; Abe, T.; Takahashi, T.; Ueda, R.: The RCK gene associ-
ated with

[138071]t(11;14) translocation is distinct from the MLL/ALL-1 gene
with t(4;11)

[138072]and t(11;19) translocations. Cancer Res. 52: 6083–6087,
1992.

[138073]

[138074]

[138075]12701.

[138076]Akao, Y.; Tsujimoto, Y.; Finan, J.; Nowell, P. C.; Croce, C.
M.

[138077]: Molecular characterization of a t(11;14)(q23;q32) chro-
mosome translocation

[138078]in a B-cell lymphoma. Cancer Res. 50: 4856–4859, 1990.

[138079]

[138080]

[138081]12702.

[138082]Lu, D.; Yunis, J. J.: Cloning, expression and localization of

[138083]an RNA helicase gene from a human lymphoid cell line
with chromosomal

[138084]breakpoint 11q23.3. Nucleic Acids Res. 20: 1967–1972,
1992.

[138085]

[138086]

[138087]12703.

[138088]Seto, M.; Yamamoto, K.; Takahashi, T.; Ueda, R.: Cloning
and expression

[138089]of a murine cDNA homologous to the human RCK/P54, a
lymphoma-linked

[138090]chromosomal translocation junction gene on 11q23. Gene
166: 293–296,

[138091]1995.

[138092]

[138093]

[138094]12704.

[138095]Tunnacliffe, A.; Perry, H.; Radice, P.; Budarf, M. L.;

Emanuel,

[138096]B. S.: A panel of sequence tagged sites for chromosome

band 11q23. Genomics 17:

[138097]744–747, 1993.

[138098]

[138099]

[138100]12705.

[138101]Hallier, E.; Jager, R.; Deutschmann, S.; Bolt, H. M.; Peter, H.

[138102]: Glutathione conjugation and cytochrome P–450

metabolism of methyl

[138103]chloride in vitro. Toxicol. in Vitro 4: 513–517, 1990.

[138104]

[138105]

[138106]12706.

[138107]Pemble, S.; Schroeder, K. R.; Spencer, S. R.; Meyer, D. J.;

Hallier,

[138108]E.; Bolt, H. M.; Ketterer, B.; Taylor, J. B.: Human glutathione S-transferase

[138109]theta (GSTT1): cDNA cloning and the characterization of a genetic

[138110]polymorphism. Biochem. J. 300: 271–276, 1994.

[138111]

[138112]

[138113]12707.

[138114]Peter, H.; Deutschmann, S.; Reichel, C.; Hallier, E.:
Metabolism

[138115]of methyl chloride by human erythrocytes. Arch. Toxicol.
63: 351–355,

[138116]1989.

[138117]

[138118]

[138119]12708.

[138120]Schroder, K. R.; Wiebel, F. A.; Reich, S.; Dannappel, D.;
Bolt,

[138121]H. M.; Hallier, E.: Glutathione S-transferase (GST) theta
polymorphism

[138122]influences background SCE rate. Arch. Toxicol. 69:
505–507, 1995.

[138123]

[138124]

[138125]12709.

[138126]Webb, G.; Vaska, V.; Coggan, M.; Board, P.: Chromosomal
localization

[138127]of the gene for the human theta class glutathione trans-
ferase (GSTT1). Genomics 33:

[138128]121–123, 1996.

[138129]

[138130]

[138131]12710.

[138132]Wiebel, F. A.; Dommermuth, A.; Thier, R.: The hereditary
transmission

[138133]of the glutathione transferase hGSTT1–1 conjugator phe-
notype in a

[138134]large family. Pharmacogenetics 9: 251–256, 1999.

[138135]

[138136]

[138137]12711.

[138138]Tiranti, V.; Rossi, E.; Ruiz-Carrillo, A.; Rossi, G.; Rocchi, M.;

[138139]DiDonato, S.; Zuffardi, O.; Zeviani, M.: Chromosomal localization

[138140]of mitochondrial transcription factor A (TCF6), single-stranded DNA-binding

[138141]protein (SSBP), and endonuclease G (ENDOG), three human housekeeping

[138142]genes involved in mitochondrial biogenesis. Genomics 25: 559–564,

[138143]1995.

[138144]

[138145]

[138146]12712.

[138147]Li, L. Y.; Luo, X.; Wang, X.: Endonuclease G is an apoptotic DNase

[138148]when released from mitochondria. Nature 412: 95–99, 2001.

[138149]

[138150]

[138151]12713.

[138152]Parrish, J.; Li, L.; Klotz, K.; Ledwich, D.; Wang, X.; Xue, D.

[138153]: Mitochondrial endonuclease G is important for apoptosis
in *C. elegans*. *Nature* 412:

[138154]90–94, 2001.

[138155]

[138156]

[138157]12714.

[138158]Dumon, K. R.; Ishii, H.; Fong, L. Y. Y.; Zanesi, N.; Fidanza,
V.;

[138159]Mancini, R.; Vecchione, A.; Baffa, R.; Trapasso, F.; During,
M. J.;

[138160]Huebner, K.; Croce, C. M.: FHIT gene therapy prevents tu-
mor development

[138161]in Fhit-deficient mice. *Proc. Nat. Acad. Sci.* 98:
3346–3351, 2001.

[138162]

[138163]

[138164]12715.

[138165]Ikeda, A.; Zheng, Q. Y.; Zuberi, A. R.; Johnson, K. R.; Nag-

gert,

[138166]J. K.; Nishina, P. M.: Microtubule-associated protein 1A is
a modifier

[138167]of tubby hearing (moth1). Nature Genet. 30: 401–405,
2002.

[138168]

[138169]

[138170]12716.

[138171]Roy, S. K.; Hu, J.; Meng, Q.; Xia, Y.; Shapiro, P. S.; Reddy, S.

[138172]P. M.; Platanias, L. C.; Lindner, D. J.; Johnson, P. F.;

Pritchard,

[138173]C.; Pages, G.; Pouyssegur, J.; Kalvakolanu, D. V.: MEKK1

plays a

[138174]critical role in activating the transcription factor C/

EBP-beta-dependent

[138175]gene expression in response to IFN-gamma. Proc. Nat.

Acad. Sci. 99:

[138176]7945–7950, 2002.

[138177]

[138178]

[138179]12717.

[138180]Chou, D.; Miyashita, T.; Mollenhauer, H. W.; Ueki, K.;
Kastury,

[138181]K.; Druck, T.; von Deimling, A.; Huebner, K.; Reed, J. C.;
Louis,

[138182]D. N.: The BAX gene maps to the glioma candidate region
at 19q13.3,

[138183]but is not altered in human gliomas. Cancer Genet. Cyto-
genet. 88:

[138184]136–140, 1996.

[138185]

[138186]

[138187]12718.

[138188]Deckwerth, T. L.; Elliott, J. L.; Knudson, C. M.; Johnson, E.
M.,

[138189]Jr.; Snider, W. D.; Korsmeyer, S. J.: BAX is required for
neuronal

[138190]death after trophic factor deprivation and during develop-
ment. Neuron 17:

[138191]401–411, 1996.

[138192]

[138193]

[138194]12719.

[138195]Knudson, C. M.; Tung, K. S. K.; Tourtellotte, W. G.; Brown, G.

[138196]A. J.; Korsmeyer, S. J.: Bax-deficient mice with lymphoid hyperplasia

[138197]and male germ cell death. Science 270: 96–99, 1995.

[138198]

[138199]

[138200]12720.

[138201]LeBlanc, H.; Lawrence, D.; Varfolomeev, E.; Totpal, K.; Morlan,

[138202]J.; Schow, P.; Fong, S.; Schwall, R.; Sinicropi, D.; Ashkenazi, A.

[138203]: Tumor-cell resistance to death receptor-induced apoptosis through

[138204]mutational inactivation of the proapoptotic Bcl-2 homolog Bax. Nature

[138205]Med. 8: 274–281, 2002.

[138206]

[138207]

[138208]12721.

[138209]Lindsten, T.; Ross, A. J.; King, A.; Zong, W.-X.; Rathmell, J.

[138210]C.; Shiels, H. A.; Ulrich, E.; Waymire, K. G.; Mahar, P.;

Frauwirth,

[138211]K.; Chen, Y.; Wei, M.; and 9 others: The combined functions of

[138212]proapoptotic Bcl-2 family members Bak and Bax are essential for normal

[138213]development of multiple tissues. *Molec. Cell* 6: 1389-1399, 2000.

[138214]

[138215]

[138216]12722.

[138217]Marzo, I.; Brenner, C.; Zamzami, N.; Jurgensmeier, J. M.; Susin,

[138218]S. A.; Vieira, H. L. A.; Prevost, M.-C.; Xie, Z.; Matsuyama, S.; Reed,

[138219]J. C.; Kroemer, G.: Bax and adenine nucleotide translocator cooperate

[138220]in the mitochondrial control of apoptosis. *Science* 281: 2027-2031,

[138221]1998.

[138222]

[138223]

[138224]12723.

[138225]Matikainen, T.; Perez, G. I.; Jurisicova, A.; Pru, J. K.;

Schlezingner,

[138226]J. J.; Ryu, H.-Y.; Laine, J.; Sakai, T.; Korsmeyer, S. J.;

Casper,

[138227]R. F.; Sherr, D. H.; Tilly, J. L.: Aromatic hydrocarbon receptor-driven

[138228]Bax gene expression is required for premature ovarian failure caused

[138229]by biohazardous environmental chemicals. Nature Genet. 28: 355–360,

[138230]2001.

[138231]

[138232]

[138233]12724.

[138234]Matsuda, Y.; Kusano, H.; Tsujimoto, Y.: Chromosomal assignment

[138235]of the Bcl2-related genes, Bcl2l and Bax, in the mouse and
rat. Cytogenet.

[138236]Cell Genet. 74: 107–110, 1996.

[138237]

[138238]

[138239]12725.

[138240]Meijerink, J. P. P.; Mensink, E. J. B. M.; Wang, K.; Sedlak, T.

[138241]W.; Sloetjes, A. W.; de Witte, T.; Waksman, G.; Korsmeyer,
S. J.:

[138242]Hematopoietic malignancies demonstrate loss-of-function
mutations

[138243]of BAX. Blood 91: 2991–2997, 1998.

[138244]

[138245]

[138246]12726.

[138247]Miyashita, T.; Reed, J. C.: Tumor suppressor p53 is a di-
rect

[138248]transcriptional activator of the human bax gene. Cell 80:
293–299,

[138249]1995.

[138250]

[138251]

[138252]12727.

[138253]Oltvai, Z. N.; Millman, C. L.; Korsmeyer, S. J.: Bcl-2 heterodimers

[138254]in vivo with a conserved homolog, Bax, that accelerates programmed

[138255]cell death. Cell 74: 609–619, 1993.

[138256]

[138257]

[138258]12728.

[138259]Perez, G. I.; Robles, R.; Knudson, C. M.; Flaws, J. A.; Korsmeyer,

[138260]S. J.; Tilly, J. L.: Prolongation of ovarian lifespan into advanced

[138261]chronological age by Bax-deficiency. Nature Genet. 21: 200–203,

[138262]1999.

[138263]

[138264]

[138265]12729.

[138266]Rampino, N.; Yamamoto, H.; Ionov, Y.; Li, Y.; Sawai, H.;

Reed,

[138267]J. C.; Perucho, M.: Somatic frameshift mutations in the BAX
gene

[138268]in colon cancers of the microsatellite mutator phenotype.

Science 275:

[138269]967–969, 1997.

[138270]

[138271]

[138272]12730.

[138273]Suzuki, M.; Youle, R. J.; Tjandra, N.: Structure of Bax:

coregulation

[138274]of dimer formation and intracellular localization. Cell 103:

645–654,

[138275]2000.

[138276]

[138277]

[138278]12731.

[138279]Wei, M. C.; Zong, W.-X.; Cheng, E. H.-Y.; Lindsten, T.;

Panoutsakopoulou,

[138280]V.; Ross, A. J.; Roth, K. A.; MacGregor, G. R.; Thompson, C.

B.; Korsmeyer,

[138281]S. J.: Proapoptotic BAX or BAK: a requisite gateway to mitochondrial

[138282]dysfunction and death. Science 292: 727–730, 2001.

[138283]

[138284]

[138285]12732.

[138286]Zhang, L.; Yu, J.; Park, B. H.; Kinzler, K. W.; Vogelstein, B.

[138287]: Role of BAX in the apoptotic response to anticancer agents. Science 290:

[138288]989–992, 2000.

[138289]

[138290]

[138291]12733.

[138292]Dasari, V. R.; Sandhu, A. K.; Mills, D. C. B.; Athwal, R. S.; Kunapuli,

[138293]S. P.: Mapping of the P2U purinergic receptor gene to human chromosome

[138294]11q13.5–14.1. Somat. Cell Molec. Genet. 22: 75–79,
1996.

[138295]

[138296]

[138297]12734.

[138298]Katzur, A. C.; Koshimizu, T.–A.; Tomic, M.; Schultze–Mos–
gau, A.;

[138299]Ortmann, O.; Stojilkovic, S. S.: Expression and responsive–
ness of

[138300]P2Y2 receptors in human endometrial cancer cell lines. J.
Clin. Endocr.

[138301]Metab. 84: 4085–4091, 1999.

[138302]

[138303]

[138304]12735.

[138305]Parr, C. E.; Sullivan, D. M.; Paradiso, A. M.; Lazarowski, E.
R.;

[138306]Burch, L. H.; Olsen, J. C.; Erb, L.; Weisman, G. A.; Boucher,
R. C.;

[138307]Turner, J. T.: Cloning and expression of a human P(2U)
nucleotide

[138308]receptor, a target for cystic fibrosis pharmacotherapy.

Proc. Nat.

[138309]Acad. Sci. 91: 3275–3279, 1994.

[138310]

[138311]

[138312]12736.

[138313]Somers, G. R.; Hammet, F.; Woollatt, E.; Richards, R. I.;
Southey,

[138314]M. C.; Venter, D. J.: Chromosomal localization of the hu-
man P2Y(6)

[138315]purinoceptor gene and phylogenetic analysis of the P2y
purinoceptor

[138316]family. Genomics 44: 127–130, 1997.

[138317]

[138318]

[138319]12737.

[138320]Tai, C.-J.; Kang, S. K.; Cheng, K. W.; Choi, K.-C.; Nathwani,
P.

[138321]S.; Leung, P. C. K.: Expression and regulation of
P2U–purinergic

[138322]receptor in human granulosa–luteal cells. J. Clin. Endocr.

Metab. 85:

[138323]1591–1597, 2000.

[138324]

[138325]

[138326]12738.

[138327]Santamarina–Fojo, S.; Peterson, K.; Knapper, C.; Qiu, Y.;
Freeman,

[138328]L.; Cheng, J.–F.; Osorio, J.; Remaley, A.; Yang, X.–P.; Hau–
denschild,

[138329]C.; Prades, C.; Chimini, G.; Blackmon, E.; Francois, T.; Du–
verger,

[138330]N.; Rubin, E. M.; Rosier, M.; Denefle, P.; Fredrickson, D. S.;
Brewer,

[138331]H. B., Jr.: Complete genomic sequence of the human
ABCA1 gene: analysis

[138332]of the human and mouse ATP–binding cassette A pro–
moter. Proc. Nat.

[138333]Acad. Sci. 97: 7987–7992, 2000. Note: Erratum: Proc. Nat.
Acad. Sci.

[138334]99: 1098 only, 2002.

[138335]

[138336]

[138337]12739.

[138338]Szakacs, G.; Langmann, T.; Ozvegy, C.; Orso, E.; Schmitz, G.;

[138339]Varadi, A.; Sarkadi, B.: Characterization of the ATPase cycle of

[138340]human ABCA1: implications for its function as a regulator rather than

[138341]an active transporter. Biochem. Biophys. Res. Commun. 288: 1258–1264,

[138342]2001.

[138343]

[138344]

[138345]12740.

[138346]Utech, M.; Hobbel, G.; Rust, S.; Reinecke, H.; Assmann, G.; Walter,

[138347]M.: Accumulation of RhoA, RhoB, RhoG, and Rac1 in fibroblasts from

[138348]Tangier disease subjects suggests a regulatory role of Rho family

[138349]proteins in cholesterol efflux. Biochem. Biophys. Res.

Commun. 280:

[138350]229–236, 2001.

[138351]

[138352]

[138353]12741.

[138354]Zhao, L.-X.; Zhou, C.-J.; Tanaka, A.; Nakata, M.;

Hirabayashi,

[138355]T.; Amachi, T.; Shioda, S.; Ueda, K.; Inagaki, N.: Cloning,

characterization

[138356]and tissue distribution of the rat ATP-binding cassette

(ABC) transporter

[138357]ABC2/ABCA2. Biochem J. 350: 865–872, 2000.

[138358]

[138359]

[138360]12742.

[138361]Zwarts, K. Y.; Clee, S. M.; Zwinderman, A. H.; Engert, J. C.;

[138362]Singaraja, R.; Loubser, O.; James, E.; Roomp, K.; Hudson,

T. J.; Jukema,

[138363]J. W.; Kastelein, J. J. P.; Hayden, M. R.: ABCA1 regulatory

variants

[138364]influence coronary artery disease independent of effects
on plasma

[138365]lipid levels. Clin. Genet. 61: 115–125, 2002.

[138366]

[138367]

[138368]12743.

[138369]Fears, S.; Mathieu, C.; Zeleznik–Le, N.; Huang, S.; Rowley,
J.

[138370]D.; Nucifora, G.: Intergenic splicing of MDS1 and EVI1 oc-
curs in

[138371]normal tissues as well as in myeloid leukemia and pro-
duces a new member

[138372]of the PR domain family. Proc. Nat. Acad. Sci. 93:
1642–1647, 1996.

[138373]

[138374]

[138375]12744.

[138376]Mochizuki, N.; Shimizu, S.; Nagasawa, T.; Tanaka, H.;
Taniwaki,

[138377]M.; Yokota, J.; Morishita, K.: A novel gene, MEL1, mapped

to 1p36.3

[138378]is highly homologous to the MDS1/EVI1 gene and is transcriptionally

[138379]activated in t(1;3)(p36;q21)-positive leukemia cells. Blood 96:

[138380]3209-3214, 2000.

[138381]

[138382]

[138383]12745.

[138384]Ing, Y. L.; Leung, I. W. L.; Heng, H. H. Q.; Tsui, L.-C.; Las-sam,

[138385]N. J.: MLK-3: identification of a widely-expressed protein kinase

[138386]bearing an SH3 domain and a leucine zipper-basic region domain. Oncogene 9:

[138387]1745-1750, 1994.

[138388]

[138389]

[138390]12746.

[138391]Bernard, O. A.; Mauchauffe, M.; Mecucci, C.; Van Den

Berghe, H.;

[138392]Berger, R.: A novel gene, AF-1p, fused to HRX in
t(1;11)(p32;q23),

[138393]is not related to AF-4, AF-9 nor ENL. Oncogene 9:
1039-1045, 1994.

[138394]

[138395]

[138396]12747.

[138397]Fazioli, F.; Minichiello, L.; Matoska, V.; Castagnino, P.;
Miki,

[138398]T.; Wong, W. T.; Di Fiore, P. P.: Eps8, a substrate for the
epidermal

[138399]growth factor receptor kinase, enhances EGF-dependent
mitogenic signals.

[138400]EMBO J. 12: 3799-3808, 1993.

[138401]

[138402]

[138403]12748.

[138404]Fazioli, F.; Minichiello, L.; Matoskova, B.; Wong, W. T.; Di
Fiore,

[138405]P. P.: eps 15, a novel tyrosine kinase substrate, exhibits

transforming

[138406]activity. Molec. Cell. Biol. 13: 5814–5828, 1993.

[138407]

[138408]

[138409]12749.

[138410]Fazioli, F.; Wong, W. T.; Ullrich, S. J.; Sakaguchi, K.; Appella,

[138411]E.; Di Fiore, P. P.: The ezrin-like family of tyrosine kinase substrates:

[138412]receptor-specific pattern of tyrosine phosphorylation and relationship

[138413]to malignant transformation. Oncogene 8: 1335–1345, 1993.

[138414]

[138415]

[138416]12750.

[138417]Wong, W. T.; Kraus, M. H.; Carlomagno, F.; Zelano, A.; Druck, T.;

[138418]Croce, C. M.; Huebner, K.; Di Fiore, P. P.: The human eps15 gene,

[138419]encoding a tyrosine kinase substrate, is conserved in evo-

lution and

[138420]maps to 1p31–p32. Oncogene 9: 1591–1597, 1994.

[138421]

[138422]

[138423]12751.

[138424]Frydman, J.; Nimmesgern, E.; Erdjument–Bromage, H.;
Wall, J. S.;

[138425]Tempst, P.; Hartl, F.–U.: Function in protein folding of
TRiC, a

[138426]cytosolic ring complex containing TCP–1 and structurally
related subunits. EMBO

[138427]J. 11: 4767–4778, 1992.

[138428]

[138429]

[138430]12752.

[138431]Joly, E. C.; Seigny, G.; Todorov, I. T.; Bibor–Hardy, V.:
cDNA

[138432]encoding a novel TCP1–related protein. Biochim. Biophys.
Acta 1217:

[138433]224–226, 1994.

[138434]

[138435]

[138436]12753.

[138437]Sevigny, G.; Joly, E. C.; Bibor-Hardy, V.; Lemieux, N.: Assignment
segment

[138438]of the human homologue of the mTRiC-P5 gene (TRIC5)
to band 1q23 by

[138439]fluorescence in situ hybridization. Genomics 22: 634-636,
1994.

[138440]

[138441]

[138442]12754.

[138443]Sevigny, G.; Lemieux, N.; Steyaert, A.; Bibor-Hardy, V.:
Structure

[138444]of the gene coding for the mouse TRiC-P5 subunit of the
cytosolic

[138445]chaperonin TRiC. Genomics 31: 107-110, 1996.

[138446]

[138447]

[138448]12755.

[138449]Walkley, N. A.; Demaine, A. G.; Malik, A. N.: Cloning,
structure

[138450]and mRNA expression of human Cctg, which encodes the
chaperonin subunit

[138451]CCT-gamma. Biochem. J. 313: 381–389, 1996.

[138452]

[138453]

[138454]12756.

[138455]Kozu, T.; Henrich, B.; Schafer, K. P.: Structure and expres-
sion

[138456]of the gene (HNRPA2B1) encoding the human hnRNP pro-
tein A2/B1. Genomics 25:

[138457]365–371, 1995.

[138458]

[138459]

[138460]12757.

[138461]Kordeli, E.; Lambert, S.; Bennett, V.: Ankyrin-G: a new
ankyrin

[138462]gene with neural-specific isoforms localized at the axonal
initial

[138463]segment and node of Ranvier. J. Biol. Chem. 270:
2352–2359, 1995.

[138464]

[138465]

[138466]12758.

[138467]Hsu, L. C.; Chang, W.–C.; Yoshida, A.: Human aldehyde
dehydrogenase

[138468]genes, ALDH7 and ALDH8: genomic organization and
gene structure comparison. Gene 189:

[138469]89–94, 1997.

[138470]

[138471]

[138472]12759.

[138473]Hsu, L. C.; Chang, W.–C.; Yoshida, A.: Cloning of a cDNA
encoding

[138474]human ALDH7, a new member of the aldehyde dehydro-
genase family. Gene 151:

[138475]285–289, 1994.

[138476]

[138477]

[138478]12760.

[138479]Aruga, J.; Yokota, N.; Hashimoto, M.; Furuichi, T.; Fukuda, M.;

[138480]Mikoshiha, K.: A novel zinc finger protein, Zic, is involved in neurogenesis,

[138481]especially in the cell lineage of cerebellar granule cells. J. Neurochem. 63:

[138482]1880–1890, 1994.

[138483]

[138484]

[138485]12761.

[138486]Salero, E.; Perez–Sen, R.; Aruga, J.; Gimenez, C.; Zafra, F.:

[138487]Transcription factors Zic1 and Zic2 bind and transactivate the apolipoprotein

[138488]E gene promoter. J. Biol. Chem. 276: 1881–1888, 2001.

[138489]

[138490]

[138491]12762.

[138492]Yokota, N.; Aruga, J.; Takai, S.; Yamada, K.; Hamazaki, M.;

Iwase,

[138493]T.; Sugimura, H.; Mikoshiba, K.: Predominant expression
of human

[138494]Zic in cerebellar granule cell lineage and medulloblas-
toma. Cancer

[138495]Res. 56: 377–383, 1996.

[138496]

[138497]

[138498]12763.

[138499]Bevins, C. L.; Jones, D. E.; Dutra, A.; Schaffzin, J.; Muenke,

[138500]M.: Human enteric defensin genes: chromosomal map po-
sition and a

[138501]model for possible evolutionary relationships. Genomics
31: 95–106,

[138502]1996.

[138503]

[138504]

[138505]12764.

[138506]Jones, D. E.; Bevins, C. L.: Defensin–6 mRNA in human
Paneth cells:

[138507]implications for antimicrobial peptides in host defense of
the human

[138508]bowel. FEBS Lett. 315: 187–192, 1993.

[138509]

[138510]

[138511]12765.

[138512]Mallow, E. B.; Harris, A.; Salzman, N.; Russell, J. P.; DeBer-
ardinis,

[138513]R. J.; Ruchelli, E.; Bevins, C. L.: Human enteric defensins:
gene

[138514]structure and developmental expression. J. Biol. Chem.
271: 4038–4045,

[138515]1996.

[138516]

[138517]

[138518]12766.

[138519]Jones, D. E.; Bevins, C. L.: Paneth cells of the human small
intestine

[138520]express an antimicrobial peptide gene. J. Biol. Chem. 267:
23216–23225,

[138521]1992.

[138522]

[138523]

[138524]12767.

[138525]Porter, E. M.; Liu, L.; Oren, A.; Anton, P. A.; Ganz, T.: Localization
calization

[138526]of human intestinal defensin 5 in Paneth cell granules. Infect. Immun. 65:

[138527]2389–2395, 1997.

[138528]

[138529]

[138530]12768.

[138531]Quayle, A. J.; Porter, E. M.; Nussbaum, A. A.; Wang, Y. M.;
Brabec,

[138532]C.; Yip, K.–P.; Mok, S. C.: Gene expression, immunolocalization,
ization,

[138533]and secretion of human defensin–5 in human female reproductive tract. Am.

[138534]J. Path. 152: 1247–1258, 1998.

[138535]

[138536]

[138537]12769.

[138538]Chen, N. N.; Chang, C.-F.; Gallia, G. L.; Kerr, D. A.; Johnson,

[138539]E. M.; Krachmarov, C. P.; Barr, S. M.; Frisque, R. J.; Bollag, B.;

[138540]Khalili, K.: Cooperative action of cellular proteins YB-1 and Pur-alpha

[138541]with the tumor antigen of the human JC polyomavirus determines their

[138542]interaction with the viral lytic control element. Proc. Nat. Acad.

[138543]Sci. 92: 1087-1091, 1995.

[138544]

[138545]

[138546]12770.

[138547]Gallia, G. L.; Johnson, E. M.; Khalili, K.: Pur-alpha: a multifunctional

[138548]single-stranded DNA- and RNA-binding protein. Nucleic Acids Res. 28:

[138549]3197-3205, 2000.

[138550]

[138551]

[138552]12771.

[138553]Kennedy, G. C.; Rutter, W. J.: Pur-1, a zinc-finger protein
that

[138554]binds to purine-rich sequences, transactivates an insulin
promoter

[138555]in heterologous cells. Proc. Nat. Acad. Sci. 89:
11498-11502, 1992.

[138556]

[138557]

[138558]12772.

[138559]Ma, Z.-W.; Pejovic, T.; Najfeld, V.; Ward, D. C.; Johnson, E.
M.

[138560]: Localization of PURA, the gene encoding the sequence-
specific single-stranded-DNA-binding

[138561]protein Pur-alpha, to chromosome band 5q31. Cytogenet.
Cell Genet. 71:

[138562]64-67, 1995.

[138563]

[138564]

[138565]12773.

[138566]Matsuzono, Y.; Kinoshita, N.; Tamura, S.; Shimozawa, N.;
Hamasaki,

[138567]M.; Ghaedi, K.; Wanders, R. J. A.; Suzuki, Y.; Kondo, N.;
Fujiki,

[138568]Y.: Human PEX19: cDNA cloning by functional comple-
mentation, mutation

[138569]analysis in a patient with Zellweger syndrome, and poten-
tial role

[138570]in peroxisomal membrane assembly. Proc. Nat. Acad. Sci.
96: 2116-2121,

[138571]1999.

[138572]

[138573]

[138574]12774.

[138575]Kinoshita, N.; Ghaedi, K.; Shimozawa, N.; Wanders, R. J. A.;
Matsuzono,

[138576]Y.; Imanaka, T.; Okumoto, K.; Suzuki, Y.; Kondo, N.; Fujiki,
Y.:

[138577]Newly identified Chinese hamster ovary cell mutants are

defective

[138578]in biogenesis of peroxisomal membrane vesicles

(peroxisomal ghosts),

[138579]representing a novel complementation group in mammals

J. Biol. Chem. 273:

[138580]24122–24130, 1998.

[138581]

[138582]

[138583]12775.

[138584]Chen, K.–S.; Gunaratne, P. H.; Hoheisel, J. D.; Young, I. G.;

Gabor

[138585]Miklos, G. L.; Greenberg, F.; Shaffer, L. G.; Campbell, H.

D.; Lupski,

[138586]J. R.: The human homologue of the *Drosophila*

melanogaster flightless–I

[138587]gene (*flii*) maps within the Smith–Magenis microdeletion

critical region

[138588]in 17p11.2. Am. J. Hum. Genet. 56: 175–182, 1995.

[138589]

[138590]

[138591]12776.

[138592]Heisterkamp, N.; Kaartinen, V.; van Soest, S.; Bokoch, G.

M.; Groffen,

[138593]J.: Human ABR encodes a protein with GAP-rac activity and
homology

[138594]to the DBL nucleotide exchange factor domain. J. Biol.
Chem. 268:

[138595]16903–16906, 1993.

[138596]

[138597]

[138598]12777.

[138599]Heisterkamp, N.; Morris, C.; Groffen, J.: ABR, an active
BCR-related

[138600]gene. Nucleic Acids Res. 17: 8821–8831, 1989.

[138601]

[138602]

[138603]12778.

[138604]McDonald, J. D.; Daneshvar, L.; Willert, J. R.; Matsumura, K.;

[138605]Waldman, F.; Cogen, P. H.: Physical mapping of chromosome 17p13.3

[138606]in the region of a putative tumor suppressor gene important in medulloblastoma. Genomics 23:

[138607]229-232, 1994.

[138608]

[138609]

[138610]12779.

[138611]Shimomura, H.; Sanke, T.; Hanabusa, T.; Tsunoda, K.; Furuta, H.;

[138612]Nanjo, K.: Nonsense mutation of islet-1 gene (Q310X) found in a type

[138613]2 diabetic patient with a strong family history. Diabetes 49: 1597-1600,

[138614]2000.

[138615]

[138616]

[138617]12780.

[138618]Tanizawa, Y.; Riggs, A. C.; Dagogo-Jack, S.; Vaxillaire, M.;
Froguel,

[138619]P.; Liu, L.; Donis-Keller, H.; Permutt, M. A.: Isolation of the
human

[138620]LIM/homeodomain gene islet-1 and identification of a
simple sequence

[138621]repeat 1. Diabetes 43: 935-941, 1994.

[138622]

[138623]

[138624]12781.

[138625]Kleiman, F. E.; Manley, J. L.: Functional interaction of
BRCA1-associated

[138626]BARD1 with polyadenylation factor CstF-50. Science 285:
1576-1579,

[138627]1999.

[138628]

[138629]

[138630]12782.

[138631]Ishibashi, T.; Bottaro, D. P.; Chan, A.; Miki, T.; Aaronson,

S.

[138632]A.: Expression cloning of a human dual-specificity phosphatase. Proc.

[138633]Nat. Acad. Sci. 89: 12170–12174, 1992.

[138634]

[138635]

[138636]12783.

[138637]Kamb, A.; Futreal, P. A.; Rosenthal, J.; Cochran, C.; Harshman,

[138638]K. D.; Liu, Q.; Phelps, R. S.; Tavtigian, S. V.; Tran, T.; Hussey,

[138639]C.; Bell, R.; Miki, Y.; Swensen, J.; Hobbs, M. R.; Marks, J.; Bennett,

[138640]L. M.; Barrett, J. C.; Wiseman, R. W.; Shattuck-Eidens, D.: Localization

[138641]of the VHR phosphatase gene and its analysis as a candidate for BRCA1. Genomics 23:

[138642]163–167, 1994.

[138643]

[138644]

[138645]12784.

[138646]Corti, O.; Finocchiaro, G.; Rossi, E.; Zuffardi, O.; DiDonato,

[138647]S.: Molecular cloning of cDNAs encoding human carnitine
acetyltransferase

[138648]and mapping of the corresponding gene to chromosome
9q34.1. Genomics 23:

[138649]94–99, 1994.

[138650]

[138651]

[138652]12785.

[138653]Kalaria, R. N.; Harik, S. I.: Carnitine acetyltransferase ac-
tivity

[138654]in the human brain and its microvessels is decreased in
Alzheimer's

[138655]disease. Ann. Neurol. 32: 583–586, 1992.

[138656]

[138657]

[138658]12786.

[138659]van der Leij, F. R.; Huijkman, N. C. A.; Boomsma, C.;
Kuipers,

[138660]J. R. G.; Bartelds, B.: Genomics of the human carnitine

acyltransferase

[138661]genes. Molec. Genet. Metab. 71: 139–153, 2000.

[138662]

[138663]

[138664]12787.

[138665]Couch, F. J.; Rommens, J. M.; Neuhausen, S. L.; Belanger, C.; Dumont,

[138666]M.; Abel, K.; Bell, R.; Berry, S.; Bogden, R.; Cannon-Albright, L.;

[138667]Farid, L.; Frye, C.; and 30 others: Generation of an integrated

[138668]transcription map of the BRCA2 region on chromosome 13q12–q13. Genomics 36:

[138669]86–99, 1996.

[138670]

[138671]

[138672]12788.

[138673]Nelms, K.; Snow, A. J.; Noben-Trauth, K.: Dok1 encoding p62(dok)

[138674]maps to mouse chromosome 6 and human chromosome 2

in a region of translocation

[138675]in chronic lymphocytic leukemia. Genomics 53: 243–245,
1998.

[138676]

[138677]

[138678]12789.

[138679]Wisniewski, D.; Strife, A.; Wojchiewowicz, D.; Lambek, C.;
Clarkson,

[138680]B.: A 62–kilodalton tyrosine phosphoprotein constitutively
present

[138681]in primary chronic phase chronic myelogenous leukemia
enriched lineage

[138682]negative blast populations. Leukemia 8: 688–693, 1994.

[138683]

[138684]

[138685]12790.

[138686]Gengyo–Ando, K.; Kitayama, H.; Mukaida, M.; Ikawa, Y.: A
murine

[138687]neural–specific homolog corrects cholinergic defects in
Caenorhabditis

[138688]elegans unc–18 mutants. J. Neurosci. 16: 6695–6702,

1996.

[138689]

[138690]

[138691]12791.

[138692]Pevsner, J.; Hsu, S.-C.; Scheller, R. H.: n-Sec1: a neural-specific

[138693]syntaxin-binding protein. Proc. Nat. Acad. Sci. 91: 1445-1449, 1994.

[138694]

[138695]

[138696]12792.

[138697]Swanson, D. A.; Steel, J. M.; Valle, D.: Identification and characterization

[138698]of the human ortholog of rat STXBP1, a protein implicated in vesicle

[138699]trafficking and neurotransmitter release. Genomics 48: 373-376,

[138700]1998.

[138701]

[138702]

[138703]12793.

[138704]Verhage, M.; Mala, A. S.; Plomp, J. J.; Brussaard, A. B.;
Heeroma,

[138705]J. H.; Vermeer, H.; Toonen, R. F.; Hammer, R. E.; van den
Berg, T.

[138706]K.; Missler, M.; Geuze, H. J.; Sudhof, T. C.: Synaptic as-
sembly of

[138707]the brain in the absence of neurotransmitter secretion.
Science 287:

[138708]864–869, 2000.

[138709]

[138710]

[138711]12794.

[138712]Cox, P. R.; Zoghbi, H. Y.: Sequencing, expression analysis,
and

[138713]mapping of three unique human tropomodulin genes and
their mouse orthologs. Genomics 63:

[138714]97–107, 2000.

[138715]

[138716]

[138717]12795.

[138718]Watakabe, A.; Kobayashi, R.; Helfman, D. M.: N-tropomodulin: a

[138719]novel isoform of tropomodulin identified as the major binding protein

[138720]to brain tropomyosin. J. Cell Sci. 109: 2299–2310, 1996.

[138721]

[138722]

[138723]12796.

[138724]Galvin, K. M.; Donovan, M. J.; Lynch, C. A.; Meyer, R. I.; Paul,

[138725]R. J.; Lorenz, J. N.; Fairchild–Huntress, V.; Dixon, K. L.; Dunmore,

[138726]J. H.; Gimbrone, M. A., Jr.; Falb, D.; Huszar, D.: A role for Smad6

[138727]in development and homeostasis of the cardiovascular system. Nature

[138728]Genet. 24: 171–174, 2000.

[138729]

[138730]

[138731]12797.

[138732]Topper, J. N.; Cai, J.; Qiu, Y.; Anderson, K. R.; Xu, Y.-Y.;

Deeds,

[138733]J. D.; Feeley, R.; Gimeno, C. J.; Woolf, E. A.; Tayber, O.;

Mays,

[138734]G. G.; Sampson, B. A.; Schoen, F. J.; Gimbrone, M. A., Jr.;

Falb,

[138735]D.: Vascular MADs: two novel MAD-related genes selectively inducible

[138736]by flow in human vascular endothelium. Proc. Nat. Acad.

Sci. 94:

[138737]9314-9319, 1997.

[138738]

[138739]

[138740]12798.

[138741]Festing, M. F. W. (ed.): Inbred strains in biomedical research. New

[138742]York: Oxford University Press , 1979. p. 255.

[138743]

[138744]

[138745]12799.

[138746]Illa, I.; Serrano-Munuera, C.; Gallardo, E.; Lasa, A.; Rojas-Garcia,

[138747]R.; Palmer, J.; Gallano, P.; Baiget, M.; Matsuda, C.; Brown, R. H.

[138748]: Distal anterior compartment myopathy: a dysferlin mutation causing

[138749]a new muscular dystrophy phenotype. Ann. Neurol. 49: 130-134, 2001.

[138750]

[138751]

[138752]12800.

[138753]Illarioshkin, S. N.; Ivanova-Smolenskaya, I. A.; Greenberg, C.

[138754]R.; Nylén, E.; Sukhorukov, V. S.; Poleshchuk, V. V.; Markova, E. D.;

[138755]Wrogemann, K.: Identical dysferlin mutation in limb-girdle muscular

[138756]dystrophy type 2B and distal myopathy. Neurology 55: 1931-1933,

[138757]2000.

[138758]

[138759]

[138760]12801.

[138761]Liu, J.; Wu, C.; Bossie, K.; Bejaoui, K.; Hosler, B. A.; Gingrich,

[138762]J. C.; Ben Hamida, M.; Hentati, F.; Schurr, E.; de Jong, P. J.; Brown,

[138763]R. H., Jr.: Generation of 3-Mb PAC contig spanning the Miyoshi myopathy/limb

[138764]girdle muscular dystrophy (MM/LGMD2B) locus on chromosome 2p13. Genomics 49:

[138765]23–29, 1998.

[138766]

[138767]

[138768]12802.

[138769]Mahjneh, I.; Vannelli, G.; Bushby, K.; Marconi, G. P.: A large

[138770]inbred Palestinian family with two forms of muscular dystrophy. Neuromusc.

[138771]Disord. 2: 277–283, 1992.

[138772]

[138773]

[138774]12803.

[138775]Weiler, T.; Bashir, R.; Anderson, L. V. B.; Davison, K.; Moss,

[138776]J. A.; Britton, S.; Nylen, E.; Keers, S.; Vafiadaki, E.; Green-
berg,

[138777]C. R.; Bushby, K. M. D.; Wrogemann, K.: Identical mutation
in patients

[138778]with limb girdle muscular dystrophy type 2B or Miyoshi
myopathy suggests

[138779]a role for modifier gene(s). Hum. Molec. Genet. 8:
871–877, 1999.

[138780]

[138781]

[138782]12804.

[138783]Pennica, D.; Arce, V.; Swanson, T. A.; Vejsada, R.; Pollock,
R.

[138784]A.; Armanini, M.; Dudley, K.; Phillips, H. S.; Rosenthal, A.;
Kato,

[138785]A. C.; Henderson, C. E.: Cardiotrophin–1, a cytokine
present in embryonic

[138786]muscle, supports long-term survival of spinal motoneurons. Neuron 17:

[138787]63–74, 1996.

[138788]

[138789]

[138790]12805.

[138791]Pennica, D.; King, K. L.; Shaw, K. J.; Luis, E.; Rullamas, J.;

[138792]Luoh, S.-M.; Darbonne, W. C.; Knutzon, D. S.; Yen, R.;
Chien, K. R.;

[138793]Baker, J. B.; Wood, W. I.: Expression cloning of cardiotrophin 1,

[138794]a cytokine that induces cardiac myocyte hypertrophy.
Proc. Nat. Acad.

[138795]Sci. 92: 1142–1146, 1995.

[138796]

[138797]

[138798]12806.

[138799]Ghosh, A.: Learning more about NMDA receptor regulation. Science 295:

[138800]449–451, 2002.

[138801]

[138802]

[138803]12807.

[138804]Grunwald, I. C.; Korte, M.; Wolfer, D.; Wilkinson, G. A.; Unsicker,

[138805]K.; Lipp, H.-P.; Bonhoeffer, T.; Klein, R.: Kinase-independent requirement

[138806]of EphB2 receptors in hippocampal synaptic plasticity.

Neuron 32:

[138807]1027–1040, 2001.

[138808]

[138809]

[138810]12808.

[138811]Henderson, J. T.; Georgiou, J.; Jia, Z.; Robertson, J.; Elowe,

[138812]S.; Roder, J. C.; Pawson, T.: The receptor tyrosine kinase EphB2

[138813]regulates NMDA-dependent synaptic function. Neuron 32: 1041–1056,

[138814]2001.

[138815]

[138816]

[138817]12809.

[138818]Himanen, J.-P.; Rajashankar, K. R.; Lackmann, M.; Cowan, C. A.;

[138819]Henkemeyer, M.; Nikolov, D. B.: Crystal structure of an Eph receptor–ephrin

[138820]complex. Nature 414: 933–938, 2001.

[138821]

[138822]

[138823]12810.

[138824]Takasu, M. A.; Dalva, M. B.; Zigmond, R. E.; Greenberg, M. E.

[138825]: Modulation of NMDA receptor–dependent calcium influx and gene expression

[138826]through EphB receptors. Science 295: 491–495, 2002.

[138827]

[138828]

[138829]12811.

[138830]Doudney, K.; Murdoch, J. N.; Paternotte, C.; Bentley, L.;

Gregory,

[138831]S.; Copp, A. J.; Stanier, P.: Comparative physical and transcript

[138832]maps of approximately 1 Mb around loop-tail, a gene for severe neural

[138833]tube defects on distal mouse chromosome 1 and human chromosome 1q22-q23. Genomics 72:

[138834]180-192, 2001.

[138835]

[138836]

[138837]12812.

[138838]Kibar, Z.; Vogan, K. J.; Groulx, N.; Justice, M. J.; Underhill,

[138839]D. A.; Gros, P.: Ltap, a mammalian homolog of Drosophila Strabismus/Van

[138840]Gogh, is altered in the mouse neural tube mutant loop-tail. Nature

[138841]Genet. 28: 251-255, 2001.

[138842]

[138843]

[138844]12813.

[138845]Mullick, A.; Trasler, D.; Gros, P.: High-resolution linkage map

[138846]in the vicinity of the Lp locus. Genomics 26: 479–488, 1995.

[138847]

[138848]

[138849]12814.

[138850]Murdoch, J. N.; Doudney, K.; Paternotte, C.; Copp, A. J.; Stanier,

[138851]P.: Severe neural tube defects in the loop-tail mouse result from

[138852]mutation of Lpp1, a novel gene involved in floor plate specification. Hum.

[138853]Molec. Genet. 10: 2593–2601, 2001.

[138854]

[138855]

[138856]12815.

[138857]Stanier, P.; Henson, J. N.; Eddleston, J.; Moore, G. E.; Copp,

[138858]A. J.: Genetic basis of neural tube defects: the mouse gene loop-tail

[138859]maps to a region of chromosome 1 syntenic with human
1q21–q23. Genomics 26:
[138860]473–478, 1995.

[138861]

[138862]

[138863]12816.

[138864]Yu, J. X.; Chao, L.; Chao, J.: Prostasin is a novel human
serine

[138865]proteinase from seminal fluid: purification, tissue distri-
bution,

[138866]and localization in prostate gland. J. Biol. Chem. 269:
18843–18848,

[138867]1994.

[138868]

[138869]

[138870]12817.

[138871]Yu, J. X.; Chao, L.; Chao, J.: Molecular cloning, tissue-
specific

[138872]expression, and cellular localization of human prostasin
mRNA. J.

[138873]Biol. Chem. 270: 13483–13489, 1995.

[138874]

[138875]

[138876]12818.

[138877]Yu, J. X.; Chao, L.; Ward, D. C.; Chao, J.: Structure and
chromosomal

[138878]localization of the human prostatic (PRSS8) gene. Ge-
nomics 32: 334–340,

[138879]1996.

[138880]

[138881]

[138882]12819.

[138883]Arber, S.; Halder, G.; Caroni, P.: Muscle LIM protein, a
novel

[138884]essential regulator of myogenesis, promotes myogenic
differentiation. Cell 79:

[138885]221–231, 1994.

[138886]

[138887]

[138888]12820.

[138889]Fung, Y. W.; Wang, R. X.; Heng, H. H. Q.; Liew, C. C.: Map-
ping

[138890]of a human LIM protein (CLP) to human chromosome
11p15.1 by fluorescence

[138891]in situ hybridization. Genomics 28: 602–603, 1995.

[138892]

[138893]

[138894]12821.

[138895]Becker–Andre, M.; Andre, E.; DeLamarter, J. F.: Identifica-
tion

[138896]of nuclear receptor mRNAs by RT–PCR amplification of
conserved zinc–finger

[138897]motif sequences. Biochem. Biophys. Res. Commun. 194:
1371–1379,

[138898]1993.

[138899]

[138900]

[138901]12822.

[138902]Carlberg, C.; Hooft van Huijsduijnen, R.; Staple, J. K.; De–

Lamarter,

[138903]J. F.; Becker–Andre, M.: RZR_s, a new family of retinoid-related orphan

[138904]receptors that function as both monomers and homodimers. *Molec. Endocr.* 8:

[138905]757–770, 1994.

[138906]

[138907]

[138908]12823.

[138909]Giguere, V.; Beatty, B.; Squire, J.; Copeland, N. G.; Jenkins,

[138910]N. A.: The orphan nuclear receptor ROR- α (RORA) maps to a conserved

[138911]region of homology on human chromosome 15q21–q22 and mouse chromosome

[138912]9. *Genomics* 28: 596–598, 1995.

[138913]

[138914]

[138915]12824.

[138916]Giguere, V.; Tini, M.; Flock, G.; Ong, E.; Evans, R. M.; Otu-lakowski,

[138917]G.: Isoform-specific amino-terminal domains dictate
DNA-binding properties

[138918]of ROR-alpha, a novel family of orphan hormone nuclear
receptors. Genes

[138919]Dev. 8: 538-553, 1994.

[138920]

[138921]

[138922]12825.

[138923]Hamilton, B. A.; Frankel, W. N.; Kerrebrock, A. W.;
Hawkins, T.

[138924]L.; FitzHugh, W.; Kusumi, K.; Russell, L. B.; Mueller, K. L.;
van

[138925]Berkel, V.; Birren, B. W.; Kruglyak, L.; Lander, E. S.: Disrup-
tion

[138926]of the nuclear hormone receptor ROR-alpha in staggerer
mice. Nature 379:

[138927]736-739, 1996.

[138928]

[138929]

[138930]12826.

[138931]Matysiak-Scholze, U.; Nehls, M.: The structural integrity of ROR-alpha

[138932]isoforms is mutated in staggerer mice: cerebellar coexpression of

[138933]ROR-alpha-1 and ROR-alpha-4. Genomics 43: 78-84, 1997.

[138934]

[138935]

[138936]12827.

[138937]Meyer, T.; Kneissel, M.; Mariani, J.; Fournier, B.: In vitro and

[138938]in vivo evidence for orphan nuclear receptor ROR-alpha function in

[138939]bone metabolism. Proc. Nat. Acad. Sci. 97: 9197-9202, 2000.

[138940]

[138941]

[138942]12828.

[138943]Ueda, H. R.; Chen, W.; Adachi, A.; Wakamatsu, H.; Hayashi, S.;

[138944]Takasugi, T.; Nagano, M.; Nakahama, K.; Suzuki, Y.; Sug-

ano, S.; Iino,

[138945]M.; Shigeyoshi, Y.; Hashimoto, S.: A transcription factor
response

[138946]element for gene expression during circadian night. Na-
ture 418:

[138947]534–539, 2002.

[138948]

[138949]

[138950]12829.

[138951]Zanjani, H. S.; Herrup, K.; Guastavino, J.–M.; Delhayé–
Bouchaud,

[138952]N.; Mariani, J.: Developmental studies of the inferior oli-
vary nucleus

[138953]in staggerer mutant mice. Develop. Brain Res. 82: 18–28,
1994.

[138954]

[138955]

[138956]12830.

[138957]Masternak, K.; Barras, E.; Zufferey, M.; Conrad, B.;
Corthals,

[138958]G.; Aebersold, R.; Sanchez, J.–C.; Hochstrasser, D. F.;

Mach, B.;

[138959]Reith, W.: A gene encoding a novel RFX-associated trans-activator

[138960]is mutated in the majority of MHC class II deficiency patients. Nature

[138961]Genet. 20: 273–277, 1998.

[138962]

[138963]

[138964]12831.

[138965]Prange, C. K.; Pennacchio, L. A.; Lieuallen, K.; Fan, W.; Lennon,

[138966]G. G.: Characterization of the human neurocan gene, CSPG3. Gene 221:

[138967]199–205, 1998.

[138968]

[138969]

[138970]12832.

[138971]Rauch, U.; Grimpe, B.; Kulbe, G.; Arnold–Ammer, I.; Beier, D. R.;

[138972]Fassler, R.: Structure and chromosomal localization of the

mouse

[138973]neurocan gene. Genomics 28: 405–410, 1995.

[138974]

[138975]

[138976]12833.

[138977]Rauch, U.; Karthikeyan, L.; Maurel, P.; Margolis, R. U.;
Margolis,

[138978]R. K.: Cloning and primary structure of neurocan, a devel-
opmentally

[138979]regulated, aggregating chondroitin sulfate proteoglycan of
brain. J.

[138980]Biol. Chem. 267: 19536–19547, 1992.

[138981]

[138982]

[138983]12834.

[138984]Clement, S.; Krause, U.; Desmedt, F.; Tanti, J.–F.;
Behrends, J.;

[138985]Pesesse, X.; Sasaki, T.; Penninger, J.; Doherty, M.;
Malaisse, W.;

[138986]Dumont, J. E.; Le Marchand–Brustel, Y.; Erneux, C.; Hue,

L.; Schurmans,

[138987]S.: The lipid phosphatase SHIP2 controls insulin sensitivity. Nature 409:

[138988]92–97, 2001.

[138989]

[138990]

[138991]12835.

[138992]Habib, T.; Hejna, J. A.; Moses, R. E.; Decker, S. J.: Growth factors

[138993]and insulin stimulate tyrosine phosphorylation of the 51C/SHIP2 protein. J.

[138994]Biol. Chem. 273: 18605–18609, 1998.

[138995]

[138996]

[138997]12836.

[138998]Hejna, J. A.; Saito, H.; Merkens, L. S.; Tittle, T. V.; Jakobs,

[138999]P. M.; Whitney, M. A.; Grompe, M.; Friedberg, A. S.; Moses, R. E.

[139000]: Cloning and characterization of a human cDNA (INPPL1) sharing homology

[139001]with inositol polyphosphate phosphatases. Genomics 29:
285–287,

[139002]1995.

[139003]

[139004]

[139005]12837.

[139006]Pesesse, X.; Deleu, S.; De Smedt, F.; Drayer, L.; Erneux, C.:

[139007]Identification of a second SH2–domain–containing protein
closely related

[139008]to the phosphatidylinositol polyphosphate 5–phosphatase
SHIP. Biochem.

[139009]Biophys. Res. Commun. 239: 697–700, 1997.

[139010]

[139011]

[139012]12838.

[139013]Schurmans, S.; Carrio, R.; Behrends, J.; Pouillon, V.;
Merino,

[139014]J.; Clement, S.: The mouse SHIP2 (Inpp1) gene: comple-
mentary DNA,

[139015]genomic structure, promoter analysis, and gene expres-

sion in the embryo

[139016]and adult mouse. Genomics 62: 260–271, 1999.

[139017]

[139018]

[139019]12839.

[139020]Drummond, J. T.; Genschel, J.; Wolf, E.; Modrich, P.: DHFR/
MSH3

[139021]amplification in methotrexate-resistant cells alters the
hMutS-alpha/hMutS-beta

[139022]ratio and reduces the efficiency of base-base mismatch
repair. Proc.

[139023]Nat. Acad. Sci. 94: 10144–10149, 1997.

[139024]

[139025]

[139026]12840.

[139027]Fujii, H.; Shimada, T.: Isolation and characterization of
cDNA

[139028]clones derived from the divergently transcribed gene in
the region

[139029]upstream from the human dihydrofolate reductase gene. J.

Biol. Chem. 264:

[139030]10057–10064, 1989.

[139031]

[139032]

[139033]12841.

[139034]Inokuchi, K.; Ikejima, M.; Watanabe, A.; Nakajima, E.;
Orimo, H.;

[139035]Nomura, T.; Shimada, T.: Loss of expression of the human
MSH3 gene

[139036]in hematological malignancies. Biochem. Biophys. Res.
Commun. 214:

[139037]171–179, 1995.

[139038]

[139039]

[139040]12842.

[139041]Linton, J. P.; Yen, J. Y.–J.; Selby, E.; Chen, Z.; Chinsky, J.

[139042]M.; Liu, K.; Kellems, R. E.; Crouse, G. F.: Dual bidirectional
promoters

[139043]at the mouse dhfr locus: cloning and characterization of
two mRNA

[139044]classes of the divergently transcribed Rep-1 gene. Molec.
Cell. Biol. 9:

[139045]3058-3072, 1989.

[139046]

[139047]

[139048]12843.

[139049]Marra, G.; Iaccarino, I.; Lettieri, T.; Roscilli, G.; Delmastro,

[139050]P.; Jiricny, J.: Mismatch repair deficiency associated with
overexpression

[139051]of the MSH3 gene. Proc. Nat. Acad. Sci. 95: 8568-8573,
1998.

[139052]

[139053]

[139054]12844.

[139055]Nakajima, E.; Orimo, H.; Ikejima, M.; Shimada, T.: Nine-bp
repeat

[139056]polymorphism in exon 1 of the hMSH3 gene. Jpn. J. Hum.
Genet. 40:

[139057]343-345, 1995.

[139058]

[139059]

[139060]12845.

[139061]Risinger, J. I.; Umar, A.; Boyd, J.; Berchuck, A.; Kunkel, T.
A.;

[139062]Barrett, J. C.: Mutation of MSH3 in endometrial cancer and
evidence

[139063]for its functional role in heteroduplex repair. Nature
Genet. 14:

[139064]102–109, 1996.

[139065]

[139066]

[139067]12846.

[139068]Smith, M. L.; Mitchell, P. J.; Crouse, G. F.: Analysis of the

[139069]mouse Dhfr/Rep–3 major promoter region by using
linker–scanning and

[139070]internal deletion mutations and DNase I footprinting.
Molec. Cell.

[139071]Biol. 10: 6003–6012, 1990.

[139072]

[139073]

[139074]12847.

[139075]Watanabe, A.; Ikejima, M.; Suzuki, N.; Shimada, T.: Ge-
nomic organization

[139076]and expression of the human MSH3 gene. Genomics 31:
311–318, 1996.

[139077]

[139078]

[139079]12848.

[139080]Yin, J.; Kong, D.; Wang, S.; Zou, T.–T. Souza, R. F.; Smolin-
ski,

[139081]K. N.; Lynch, P. M.; Hamilton, S. R.; Sugimura, H.; Powell,
S. M.;

[139082]Young, J.; Abraham, J. M.; Meltzer, S. J.: Mutation of
hMSH3 and

[139083]hMSH6 mismatch repair genes in genetically unstable hu-
man colorectal

[139084]and gastric carcinomas. Hum. Mutat. 10: 474–478, 1997.

[139085]

[139086]

[139087]12849.

[139088]Chan, A. M-L.; McGovern, E. S.; Catalano, G.; Fleming, T. P.; Miki,

[139089]T.: Expression cDNA cloning of a novel oncogene with sequence similarity

[139090]to regulators of small GTP-binding proteins. Oncogene 9: 1057-1063,

[139091]1994.

[139092]

[139093]

[139094]12850.

[139095]Miki, T.; Fleming, T. P.; Crescenzi, M.; Molloy, C. J.; Blam, S.

[139096]B.; Reynolds, S. H.; Aaronson, S. A.: Development of a highly efficient

[139097]expression cDNA cloning system: application to oncogene isolation. Proc.

[139098]Nat. Acad. Sci. 88: 5167-5171, 1991.

[139099]

[139100]

[139101]12851.

[139102]Takai, S.; Chan, A. M.-L.; Yamada, K.; Miki, T.: Assignment
of

[139103]the human TIM proto-oncogene to 7q33-q35. Cancer
Genet. Cytogenet. 83:

[139104]87-89, 1995.

[139105]

[139106]

[139107]12852.

[139108]Burger, B.; Uhlhaas, S.; Mangold, E.; Propping, P.; Friedl,
W.;

[139109]Jenne, D.; Dockter, G.; Back, W.: Novel de novo mutation
of MADH4/SMAD4

[139110]in a patient with juvenile polyposis. (Letter) Am. J. Med.
Genet. 110:

[139111]289-291, 2002.

[139112]

[139113]

[139114]12853.

[139115]Kurima, K.; Peters. L. M.; Yang, Y.; Riazuddin, S.; Ahmed,
Z. M.;

[139116]Naz, S.; Arnaud, D.; Drury, S.; Mo, J.; Makishima, T.;
Ghosh, M.;

[139117]Menon, P. S. N.; and 13 others: Dominant and recessive
deafness

[139118]caused by mutations of a novel gene, TMC1, required for
cochlear hair-cell

[139119]function. Nature Genet. 30: 277–284, 2002.

[139120]

[139121]

[139122]12854.

[139123]Bryant, P. J.; Huettner, B.; Held, L. I., Jr.; Ryerse, J.; Szi-
donya,

[139124]J.: Mutations at the fat locus interfere with cell prolifera-
tion

[139125]control and epithelial morphogenesis in Drosophila. Dev.
Biol. 129:

[139126]541–554, 1988.

[139127]

[139128]

[139129]12855.

[139130]Dunne, J.; Hanby, A. M.; Poulson, R.; Jones, T. A.; Sheer, D.;

[139131]Chin, W. G.; Da, S. M.; Zhao, Q.; Beverley, P. C. L.; Owen, M. J.

[139132]: Molecular cloning and tissue expression of FAT, the human homologue

[139133]of the Drosophila fat gene that is located on chromosome 4q34–q35

[139134]and encodes a putative adhesion molecule. Genomics 30: 207–223,

[139135]1995.

[139136]

[139137]

[139138]12856.

[139139]Hortsch, M.; Goodman, C. S.: Cell and substrate adhesion molecules

[139140]in Drosophila. Ann. Rev. Cell. Biol. 7: 505–557, 1991.

[139141]

[139142]

[139143]12857.

[139144]Mahoney, P. A.; Weber, U.; Onofrechuk, P.; Biessmann, H.;
Bryant,

[139145]P. J.; Goodman, C. S.: The fat tumor suppressor gene in
Drosophila

[139146]encodes a novel member of the cadherin gene superfam-
ily. Cell 67:

[139147]853–868, 1991.

[139148]

[139149]

[139150]12858.

[139151]Crowe, P. D.; VanArsdale, T. L.; Walter, B. N.; Ware, C. F.;
Hession,

[139152]C.; Ehrenfels, B.; Browning, J. L.; Din, W. S.; Goodwin, R. G;
Smith,

[139153]C. A.: A lymphotoxin–beta–specific receptor. Science 264:
707–710,

[139154]1994.

[139155]

[139156]

[139157]12859.

[139158]Nakamura, T.; Tashiro, K.; Nazarea, M.; Nakano, T.;
Sasayama, S.;

[139159]Honjo, T.: The murine lymphotoxin-beta receptor cDNA:
isolation by

[139160]the signal sequence trap and chromosomal mapping. Ge-
nomics 30: 312-319,

[139161]1995.

[139162]

[139163]

[139164]12860.

[139165]Tashiro, K.; Tada, H.; Heilker, R.; Shirozu, M.; Nakano, T.;
Honjo,

[139166]T.: Signal sequence trap: a cloning strategy for secreted
proteins

[139167]and type I membrane proteins. Science 261: 600-603,
1993.

[139168]

[139169]

[139170]12861.

[139171]George, A.; Sabsay, B.; Simonian, P. A.; Veis, A.: Charac-
terization

[139172]of a novel dentin matrix acidic phosphoprotein: implications for induction

[139173]of biomineralization . J. Biol. Chem. 268: 12624–12630, 1993.

[139174]

[139175]

[139176]12862.

[139177]Hirst, K. L.; Simmons, D.; Feng, J.; Aplin, H.; Dixon, M. J.; MacDougall,

[139178]M.: Elucidation of the sequence and the genomic organization of the

[139179]human dentin matrix acidic phosphoprotein 1 (DMP1) gene: exclusion

[139180]of the locus from a causative role in the pathogenesis of dentinogenesis

[139181]imperfecta type II. Genomics 42: 38–45, 1997.

[139182]

[139183]

[139184]12863.

[139185]MacDougall, M.; DuPont, B. R.; Simmons, D.; Leach, R. J.: Assignment

[139186]of DMP1 to human chromosome 4 band q21 by in situ hybridization. Cytogenet.

[139187]Cell. Genet. 74: 189 only, 1996.

[139188]

[139189]

[139190]12864.

[139191]Lu, Z.; Xu, S.; Joazeiro, C.; Cobb, M. H.; Hunter, T.: The PHD

[139192]domain of MEKK1 acts as an E3 ubiquitin ligase and mediates ubiquitination

[139193]and degradation of ERK1/2. Molec. Cell 9: 945–956, 2002.

[139194]

[139195]

[139196]12865.

[139197]Bohm, H.; Benndorf, R.; Gaestel, M.; Gross, B.; Nurnberg, P.; Kraft,

[139198]R.; Otto, A.; Bielka, H.: The growth-related protein P23 of the Ehrlich

[139199]ascites tumor: translational control, cloning and primary structure. Biochem.

[139200]Int. 19: 277–286, 1989.

[139201]

[139202]

[139203]12866.

[139204]Barnard, R. C.; Pascall, J. C.; Brown, K. D.; McKay, I. A.;
Williams,

[139205]N. S.; Bustin, S. A.: Coding sequence of ERF–1, the human
homologue

[139206]of Tis11b/cMG1, members of the Tis11 family of early re–
sponse genes. Nucleic

[139207]Acids Res. 21: 3580 only, 1993.

[139208]

[139209]

[139210]12867.

[139211]Bustin, S. A.; Xiao–Feng, N.; Barnard, R. C.; Kumar, V.;
Pascall,

[139212]J. C.; Brown, K. D.; Leigh, I. M.; Williams, N. S.; McKay, I.
A.:

[139213]Cloning and characterisation of ERF1, a human member of
the Tis11

[139214]family of early-response genes. DNA Cell Biol. 13:
449–459, 1994.

[139215]

[139216]

[139217]12868.

[139218]Maclean, K. N.; See, C. G.; McKay, I. A.; Bustin, S. A.: The
human

[139219]immediate early gene BRF1 maps to chromosome
14q22–q24. Genomics 30:

[139220]89–90, 1995.

[139221]

[139222]

[139223]12869.

[139224]Ning, Z.–Q.; Norton, J. D.; Li, J.; Murphy, J. J.: Distinct
mechanisms

[139225]for rescue from apoptosis in Ramos human B cells by sig-
naling through

[139226]CD40 and interleukin–4 receptor: a role for inhibition of
an early

[139227]response gene, Berg36. Europ. J. Immun. 26: 2356–2363,
1996.

[139228]

[139229]

[139230]12870.

[139231]Chihade, J. W.; Brown, J. R.; Schimmel, P. R.; Ribas de Pouplana,

[139232]L.: Origin of mitochondria in relation to evolutionary history of

[139233]eukaryotic alanyl-tRNA synthetase. Proc. Nat. Acad. Sci. 97: 12153–12157,

[139234]2000.

[139235]

[139236]

[139237]12871.

[139238]Maas, S.; Kim, Y.-G.; Rich, A.: Genomic clustering of tRNA-specific

[139239]adenosine deaminase ADAT1 and two tRNA synthetases. Mammalian Genome 12:

[139240]387–393, 2001.

[139241]

[139242]

[139243]12872.

[139244]Derynck, R.; Gelbart, W. M.; Harland, R. M.; Heldin, C.-H.;
Kern,

[139245]S. E.; Massague, J.; Melton, D. A.; Mlodzik, M.; Padgett, R.
W.; Roberts,

[139246]A. B.; Smith, J.; Thomsen, G. H.; Vogelstein, B.; Wang,
X.-F.: Nomenclature:

[139247]vertebrate mediators of TGF-beta family signals. (Letter)
Cell 87:

[139248]173 only, 1996.

[139249]

[139250]

[139251]12873.

[139252]Friedl, W.; Kruse, R.; Uhlhaas, S.; Stolte, M.; Schartmann,
B.;

[139253]Keller, K. M.; Jungck, M.; Stern, M.; Loff, S.; Back, W.;
Propping,

[139254]P.; Jenne, D. E.: Frequent 4-bp deletion in exon 9 of the
SMAD4/MADH4

[139255]gene in familial juvenile polyposis patients. Genes Chro-
mosomes Cancer 25:

[139256]403–406, 1999.

[139257]

[139258]

[139259]12874.

[139260]Hahn, S. A.; Schutte, M.; Hoque, T. M. S.; Moskaluk, C. A.;
da

[139261]Costa, L. T.; Rozenblum, E.; Weinstein, C. L.; Fischer, A.;
Yeo, C.

[139262]J.; Hruban, R. H.; Kern, S. E.: DPC4, a candidate tumor
suppressor

[139263]gene at human chromosome 18q21.1. Science 271:
350–354, 1996.

[139264]

[139265]

[139266]12875.

[139267]Howe, J. R.; Shellnut, J.; Wagner, B.; Ringold, J. C.; Sayed,

[139268]M. G.; Ahmed, A. F.; Lynch, P. M.; Amos, C. I.; Sistonen, P.;
Aaltonen,

[139269]L. A.: Common deletion of SMAD4 in juvenile polyposis is
a mutational

[139270]hotspot. Am. J. Hum. Genet. 70: 1357–1362, 2002.

[139271]

[139272]

[139273]12876.

[139274]Inman, G. J.; Nicolas, F. J.; Hill, C. S.: Nucleocytoplasmic

[139275]shuttling of Smads 2, 3, and 4 permits sensing of TGF-
beta receptor

[139276]activity. Molec. Cell 10: 283–294, 2002.

[139277]

[139278]

[139279]12877.

[139280]Kim, S. K.; Fan, Y.; Papadimitrakopoulou, V.; Clayman, G.;
Hittelman,

[139281]W. N.; Hong, W. K.; Lotan, R.; Mao, L.: DPC4, a candidate
tumor suppressor

[139282]gene, is altered infrequently in head and neck squamous
cell carcinoma. Cancer

[139283]Res. 56: 2519–2521, 1996.

[139284]

[139285]

[139286]12878.

[139287]Kinzler, K. W.; Vogelstein, B.: Landscaping the cancer terrain. Science 280:

[139288]1036–1037, 1998.

[139289]

[139290]

[139291]12879.

[139292]MacGrogan, D.; et al.; et al: Comparative mutational analysis

[139293]of DPC4 (Smad4) in prostatic and colorectal carcinomas. Oncogene 15:

[139294]1111–1114, 1997.

[139295]

[139296]

[139297]12880.

[139298]Roth, S.; Johansson, M.; Loukola, A.; Peltomaki, P.; Jarvinen,

[139299]H.; Mecklin, J.-P.; Aaltonen, L. A.: Mutation analysis of

SMAD2,

[139300]SMAD3, and SMAD4 genes in hereditary non-polyposis colorectal cancer. J.

[139301]Med. Genet. 37: 298–300, 2000.

[139302]

[139303]

[139304]12881.

[139305]Schutte, M.; Hruban, R. H.; Hedrick, L.; Cho, K. R.;
Nadasdy,

[139306]G. M.; Weinstein, C. L.; Bova, G. S.; Isaacs, W. B.; Cairns, P.;
Nawroz,

[139307]H.; Sidransky, D.; Casero, R. A., Jr.; Meltzer, P. S.; Hahn, S.
A.;

[139308]Kern, S. E.: DPC4 gene in various tumor types. Cancer Res.
56:

[139309]2527–2530, 1996.

[139310]

[139311]

[139312]12882.

[139313]Shioda, T.; Lechleider, R. J.; Dunwoodie, S. L.; Li, H.; Ya–

hata,

[139314]T.; de Caestecker, M. P.; Fenner, M. H.; Roberts, A. B.; Is-selbacher,

[139315]K. J.: Transcriptional activating activity of Smad4: roles of SMAD

[139316]hetero-oligomerization and enhancement by an associating transactivator. Proc.

[139317]Nat. Acad. Sci. 95: 9785–9790, 1998.

[139318]

[139319]

[139320]12883.

[139321]Sirard, C.; de la Pompa, J. L.; Elia, A.; Itie, A.; Mirtsos, C.;

[139322]Cheung, A.; Hahn, S.; Wakeham, A.; Schwartz, L.; Kern, S. E.; Rossant,

[139323]J.; Mak, T. W.: The tumor suppressor gene Smad4/Dpc4 is required

[139324]for gastrulation and later for anterior development of the mouse embryo. Genes

[139325]Dev. 12: 107–119, 1998.

[139326]

[139327]

[139328]12884.

[139329]Tagaki, Y.; et al.; et al: Somatic alterations of the DPC4
gene

[139330]in human colorectal cancers in vivo. Gastroenterology
111: 1369–1372,

[139331]1996.

[139332]

[139333]

[139334]12885.

[139335]Takaku, K.; Oshima, M.; Miyoshi, H.; Matsui, M.; Seldin, M.
F.;

[139336]Taketo, M. M.: Intestinal tumorigenesis in compound mu-
tant mice of

[139337]both Dpc4 (Smad4) and Apc genes. Cell 92: 645–656,
1998.

[139338]

[139339]

[139340]12886.

[139341]Thiagalingam, S.; Lengauer, C.; Leach, F. S.; Schutte, M.;
Hahn,

[139342]S. A.; Overhauser, J.; Willson, J. K. V.; Markowitz, S.;
Hamilton,

[139343]S. R.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Evaluation
of

[139344]candidate tumour suppressor genes on chromosome 18 in
colorectal cancers. Nature

[139345]Genet. 13: 343–346, 1996.

[139346]

[139347]

[139348]12887.

[139349]Zawel, L.; Dai, J. L.; Buckhaults, P.; Zhou, S.; Kinzler, K. W.;

[139350]Vogelstein, B.; Kern, S. E.: Human Smad3 and Smad4 are
sequence-specific

[139351]transcription activators. Molec. Cell 1: 611–617, 1998.

[139352]

[139353]

[139354]12888.

[139355]Zhou, S.; Buckhaults, P.; Zawel, L.; Bunz, F.; Riggins, G.; Le

[139356]Dai, J.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Targeted

deletion

[139357]of Smad4 shows it is required for transforming growth factor beta

[139358]and activin signaling in colorectal cancer cells. Proc. Nat. Acad.

[139359]Sci. 95: 2412–2416, 1998.

[139360]

[139361]

[139362]12889.

[139363]Abbs, S.; Roberts, R. G.; Mathew, C. G.; Bentley, D. R.; Bobrow,

[139364]M.: Accurate assessment of intragenic recombination frequency within

[139365]the Duchenne muscular dystrophy gene. Genomics 7: 602–606, 1990.

[139366]

[139367]

[139368]12890.

[139369]Ahn, A. H.; Kunkel, L. M.: The structural and functional diversity

[139370]of dystrophin. Nature Genet. 3: 283–291, 1993.

[139371]

[139372]

[139373]12891.

[139374]Alwine, J. C.; Kemp, D. J.; Stark, G. R.: Method for detection
tion

[139375]of specific RNAs in agarose gels by transfer to diazoben-
zyloxymethyl-paper

[139376]and by hybridization with DNA probes. Proc. Nat. Acad.
Sci. 74:

[139377]5350-5354, 1977.

[139378]

[139379]

[139380]12892.

[139381]Angelini, C.; Beggs, A. H.; Hoffman, E. P.; Fanin, M.;
Kunkel,

[139382]L. M.: Enormous dystrophin in a patient with Becker mus-
cular dystrophy. Neurology 40:

[139383]808-812, 1990.

[139384]

[139385]

[139386]12893.

[139387]Badorff, C.; Berkely, N.; Mehrotra, S.; Talhouk, J. W.;

Rhoads,

[139388]R. E.; Knowlton, K. U.: Enteroviral protease 2A directly
cleaves

[139389]dystrophin and is inhibited by a dystrophin-based sub-
strate analogue. J.

[139390]Biol. Chem. 275: 11191–11197, 2000.

[139391]

[139392]

[139393]12894.

[139394]Badorff, C.; Lee, G.–H.; Lamphear, B. J.; Martone, M. E.;

Campbell,

[139395]K. P.; Rhoads, R. E.; Knowlton, K. U.: Enteroviral protease
2A cleaves

[139396]dystrophin: evidence of cytoskeletal disruption in an ac-
quired cardiomyopathy. Nature

[139397]Med. 5: 320–326, 1999.

[139398]

[139399]

[139400]12895.

[139401]Bakker, E.; Pearson, P. L.: Mutation of the Duchenne muscular

[139402]dystrophy gene associated with meiotic recombination.
(Letter) Clin.

[139403]Genet. 30: 347–349, 1986.

[139404]

[139405]

[139406]12896.

[139407]Bakker, E.; Hofker, M. H.; Goor, N.; Mandel, J. L.; Wroger-

[139408]K.; Davies, K. E.; Kunkel, L. M.; Willard, H. F.; Fenton, W.
A.; Sandkuyl,

[139409]L.; Majoor-Krakauer, D.; van Essen, A. J.; Jahoda, M. G. J.;
Sachs,

[139410]E. S.; van Ommen, G. J. B.; Pearson, P. L.: Prenatal diagnosis and

[139411]carrier detection of Duchenne muscular dystrophy with
closely linked

[139412]RFLPs. Lancet I: 655–658, 1985.

[139413]

[139414]

[139415]12897.

[139416]Bakker, E.; Van Broeckhoven, C.; Bonten, E. J.; van de Vooren,

[139417]M. J.; Veenema, H.; Van Hul, W.; Van Ommen, G. J. B.; Vandenberghe,

[139418]A.; Pearson, P. L.: Germline mosaicism and Duchenne muscular dystrophy

[139419]mutations. Nature 329: 554–556, 1987.

[139420]

[139421]

[139422]12898.

[139423]Bar, S.; Barnea, E.; Levy, Z.; Neuman, S.; Yaffe, D.; Nudel, U.

[139424]: A novel product of the Duchenne muscular dystrophy gene which greatly

[139425]differs from the known isoforms in its structure and tissue distribution. Biochem.

[139426]J. 272: 557–560, 1990.

[139427]

[139428]

[139429]12899.

[139430]Barbieri, A. M.; Soriani, N.; Tubiello, G. M.; Ferrari, M.;
Carrera,

[139431]P.: A nonsense mutation (gln-673-term) in exon 17 of the
human dystrophin

[139432]gene detected by heteroduplex analysis. Hum. Genet. 96:
343-344,

[139433]1995.

[139434]

[139435]

[139436]12900.

[139437]Bartlett, R. J.; Pericak-Vance, M. A.; Koh, J.; Yamaoka, L.
H.;

[139438]Chen, J. C.; Hung, W.-Y.; Speer, M. C.; Wapenaar, M. C.;
Van Ommen,

[139439]G. J. B.; Bakker, E.; Pearson, P. L.; Kandt, R. S.; Siddique,
T.;

[139440]Gilbert, J. R.; Lee, J. E.; Sirotkin-Roses, M. J.; Roses, A. D.:

[139441]Duchenne muscular dystrophy: high frequency of dele-
tions. Neurology 38:

[139442]1-4, 1988.

[139443]

[139444]

[139445]12901.

[139446]Barton-Davis, E. R.; Cordier, L.; Shoturma, D. I.; Leland, S.

[139447]E.; Sweeney, H. L.: Aminoglycoside antibiotics restore dystrophin

[139448]function to skeletal muscles of mdx mice. J. Clin. Invest.
104:

[139449]375-381, 1999.

[139450]

[139451]

[139452]12902.

[139453]Bastianutto, C.; Bestard, J. A.; Lahnakoski, K.; Broere, D.;
De

[139454]Visser, M.; Zaccolo, M.; Pozzan, T.; Ferlini, A.; Muntoni, F.;
Patarnello,

[139455]T.; Klamut, H. J.: Dystrophin muscle enhancer 1 is implicated in

[139456]the activation of non-muscle isoforms in the skeletal

muscle of patients

[139457]with X-linked dilated cardiomyopathy. Hum. Molec. Genet.

10: 2627-2635,

[139458]2001.

[139459]

[139460]

[139461]12903.

[139462]Baumbach, L. L.; Chamberlain, J. S.; Ward, P. A.; Farwell, N.

[139463]J.; Caskey, C. T.: Molecular and clinical correlation of deletion
tion

[139464]leading to Duchenne and Becker muscular dystrophies.

Neurology 39:

[139465]465-474, 1989.

[139466]

[139467]

[139468]12904.

[139469]Baumbach, L. L.; Ward, P. A.; Fenwick, R.; Caskey, C. T.:

Analysis

[139470]of mutations at the Duchenne muscular dystrophy locus
provides no

[139471]evidence for illegitimate recombination in deletion formation. (Abstract) Am.

[139472]J. Hum. Genet. 45 (suppl.): A173, 1989.

[139473]

[139474]

[139475]12905.

[139476]Beggs, A. H.; Koenig, M.; Boyce, F. M.; Kunkel, L. M.: Detection

[139477]of 98% of DMD/BMD gene deletions by polymerase chain reaction. Hum.

[139478]Genet. 86: 45-48, 1990.

[139479]

[139480]

[139481]12906.

[139482]Berko, B. A.; Swift, M.: X-linked dilated cardiomyopathy. New

[139483]Eng. J. Med. 316: 1186-1191, 1987.

[139484]

[139485]

[139486]12907.

[139487]Bettecken, T.; Muller, C. R.: Identification of a 220-kb insertion

[139488]into the Duchenne gene in a family with an atypical course of muscular

[139489]dystrophy. Genomics 4: 592-596, 1989.

[139490]

[139491]

[139492]12908.

[139493]Bies, R. D.: X-linked dilated cardiomyopathy. (Letter) New Eng.

[139494]J. Med. 330: 368-369, 1994.

[139495]

[139496]

[139497]12909.

[139498]Bies, R. D.; Caskey, C. T.; Fenwick, R.: An intact cysteine-rich

[139499]domain is required for dystrophin function. J. Clin. Invest. 90:

[139500]666-672, 1992.

[139501]

[139502]

[139503]12910.

[139504]Bittner, R. E.; Streubel, B.; Shorny, S.; Schaden, G.; Voit, T.;

[139505]Hoger, H.: Coisogenic all-plus-one immunization: a
model for identifying

[139506]missing proteins in null-mutant conditions. Antibodies to
dystrophin

[139507]in mdx mouse after transplantation of muscle from nor-
mal coisogenic

[139508]donor. Neuropediatrics 25: 176-182, 1994.

[139509]

[139510]

[139511]12911.

[139512]Ryu, S.; Zhou, S.; Ladurner, A. G.; Tjian, R.: The transcrip-
tional

[139513]cofactor complex CRSP is required for activity of the en-
hancer-binding

[139514]protein Sp1. Nature 397: 446-450, 1999.

[139515]

[139516]

[139517]12912.

[139518]Mohapatra, B.; Verma, S.; Shankar, S.; Suri, A.: Molecular cloning

[139519]of human testis mRNA specifically expressed in haploid germ cells,

[139520]having structural homology with the A-kinase anchoring proteins. Biochem.

[139521]Biophys. Res. Commun. 244: 540–545, 1998.

[139522]

[139523]

[139524]12913.

[139525]Turner, R. M. O.; Johnson, L. R.; Haig-Ladewig, L.; Gerton, G.

[139526]L.; Moss, S. B.: An X-linked gene encodes a major human sperm fibrous

[139527]sheath protein, hAKAP82: genomic organization, protein kinase A-RII

[139528]binding, and distribution of the precursor in the sperm tail. J.

[139529]Biol. Chem. 273: 32135–32141, 1998.

[139530]

[139531]

[139532]12914.

[139533]Dever, T. E.; Wei, C.-L.; Benkowski, L. A.; Browning, K.;
Merrick,

[139534]W. C.; Hershey, J. W. B.: Determination of the amino acid
sequence

[139535]of rabbit, human, and wheat germ protein synthesis factor
eIF-4C by

[139536]cloning and chemical sequencing. J. Biol. Chem. 269:
3212-3218,

[139537]1994.

[139538]

[139539]

[139540]12915.

[139541]Dry, K. L.; Aldred, M. A.; Edgar, A. J.; Brown, J.; Manson, F.

[139542]D. C.; Ho, M.-F.; Prosser, J.; Hardwick, L. J.; Lennon, A. A.;
Thomson,

[139543]K.; Van Keuren, M.; Kurnit, D. M.; Bird, A. C.; Jay, M.;
Monaco, A.

[139544]P.; Wright, A. F.: Identification of a novel gene, ETX1, from
Xp21.1,

[139545]a candidate gene for X-linked retinitis pigmentosa (RP3).
Hum. Molec.

[139546]Genet. 4: 2347–2353, 1995.

[139547]

[139548]

[139549]12916.

[139550]Meindl, A.; Carvalho, M. R. S.; Herrmann, K.; Lorenz, B.;
Achatz,

[139551]H.; Lorenz, B.; Apfelstedt-Sylla, E.; Wittwer, B.; Ross, M.;
Meitinger,

[139552]T.: A gene (SRPX) encoding a sushi-repeat-containing
protein is deleted

[139553]in patients with X-linked retinitis pigmentosa. Hum.
Molec. Genet. 4:

[139554]2339–2346, 1995.

[139555]

[139556]

[139557]12917.

[139558]Lee, S.-H.; Kim, W.-H.; Kim, H.-K.; Woo, K.-M.; Nam, H.-S.; Kim,

[139559]H.-S.; Kim, J.-G.; Cho, M.-H.: Altered expression of the fragile

[139560]histidine triad gene in primary gastric adenocarcinomas. Biochem.

[139561]Biophys. Res. Commun. 284: 850–855, 2001.

[139562]

[139563]

[139564]12918.

[139565]Markkanen, A.; Heinonen, K.; Knuutila, S.; de la Chapelle, A.

[139566]: Methotrexate-induced increase in gap formation in human chromosome

[139567]band 3p14. Hereditas 96: 317–319, 1982.

[139568]

[139569]

[139570]12919.

[139571]Markkanen, A.; Knuutila, S.; de la Chapelle, A.: Inducible fragile

[139572]site on chromosome 3. (Letter) Hum. Genet. 65: 217 only,
1983.

[139573]

[139574]

[139575]12920.

[139576]Anderson, D. M.; Kumaki, S.; Ahdieh, M.; Bertles, J.;
Tometsko,

[139577]M.; Loomis, A.; Giri, J.; Copeland, N. G.; Gilbert, D. J.;
Jenkins,

[139578]N. A.; Valentine, V.; Shapiro, D. N.; Morris, S. W.; Park, L.
S.;

[139579]Cosman, D.: Functional characterization of the human in-
terleukin-15

[139580]receptor alpha chain and close linkage of IL15RA and
IL2RA genes. J.

[139581]Biol. Chem. 270: 29862-29869, 1995.

[139582]

[139583]

[139584]12921.

[139585]Giri, J. G.; Kumaki, S.; Ahdieh, M.; Friend, D. J.; Loomis, A.;

[139586]Shanebeck, K.; DuBose, R.; Cosman, D.; Park, L. S.; Anderson, D. M.

[139587]: Identification and cloning of a novel IL-15 binding protein that

[139588]is structurally related to the alpha of the IL-2 receptor.

EMBO J. 14:

[139589]3654-3663, 1995.

[139590]

[139591]

[139592]12922.

[139593]Yasunaga, S.; Grati, M.; Cohen-Salmon, M.; El-Amraoui, A.; Mustapha,

[139594]M.; Salem, N.; El-Zir, E.; Loiselet, J.; Petit, C.: A mutation in

[139595]OTOF, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic

[139596]form of deafness. Nature Genet. 21: 363-369, 1999.

[139597]

[139598]

[139599]12923.

[139600]Berry, A.; Scott, H. S.; Kudoh, J.; Talior, I.; Korostishevsky,

[139601]M.; Wattenhofer, M.; Guipponi, M.; Barras, C.; Rossier, C.;
Shibuya,

[139602]K.; Wang, J.; Kawasaki, K.; Asakawa, S.; Minoshima, S.;
Shimizu, N.;

[139603]Antonarakis, S.; Bonne-Tamir, B.: Refined localization of
autosomal

[139604]recessive nonsyndromic deafness DFNB10 locus using 34
novel microsatellite

[139605]markers, genomic structure, and exclusion of six known
genes in the

[139606]region. Genomics 68: 22–29, 2000.

[139607]

[139608]

[139609]12924.

[139610]Good, P. J.; Chen, Q.; Warner, S. J.; Herring, D. C.: A family

[139611]of human RNA-binding proteins related to the Drosophila
Bruno transcriptional

[139612]regulator. J. Biol. Chem. 275: 28583–28592, 2000.

[139613]

[139614]

[139615]12925.

[139616]Roberts, R.; Timchenko, N. A.; Miller, J. W.; Reddy, S.;
Caskey,

[139617]C. T.; Swanson, M. S.; Timchenko, L. T.: Altered phospho-
rylation

[139618]and intracellular distribution of a (CUG)_n triplet repeat
RNA-binding

[139619]protein in patients with myotonic dystrophy and in my-
otonin protein

[139620]kinase knockout mice. Proc. Nat. Acad. Sci. 94:
13221–13226, 1997.

[139621]

[139622]

[139623]12926.

[139624]Li, Z.; Jiang, H.; Xie, W.; Zhang, Z.; Smrcka, A. V.; Wu, D.:

[139625]Roles of PLC-beta-2 and -beta-3 and PI3K-gamma in
chemoattractant-mediated

[139626]signal transduction. Science 287: 1046-1049, 2000.

[139627]

[139628]

[139629]12927.

[139630]Greger, V.; Knoll, J. H. M.; Woolf, E.; Glatt, K.; Tyndale, R.

[139631]F.; DeLorey, T. M.; Olsen, R. W.; Tobin, A. J.; Sikela, J. M.;
Nakatsu,

[139632]Y.; Brilliant, M. H.; Whiting, P. J.; Lalande, M.: The gamma-
aminobutyric

[139633]acid receptor gamma-3 subunit gene (GABRG3) is tightly
linked to the

[139634]alpha-5 subunit gene (GABRA5) on human chromosome
15q11-q13 and is

[139635]transcribed in the same orientation. Genomics 26:
258-264, 1995.

[139636]

[139637]

[139638]12928.

[139639]Fowler, K. J.; Saffery, R.; Irvine, D. V.; Trowell, H. E.; Choo,

[139640]K. H. A.: Mouse centromere protein F (Cenpf) gene maps
to the distal

[139641]region of chromosome 1 by interspecific backcross analysis. Cytogenet.

[139642]Cell Genet. 82: 180–181, 1998.

[139643]

[139644]

[139645]12929.

[139646]Liao, H.; Winkfein, R. J.; Mack, G.; Rattner, J. B.; Yen, T. J.

[139647]: CENP–F is a protein of the nuclear matrix that assembles
onto kinetochores

[139648]at late G2 and is rapidly degraded after mitosis. J. Cell
Biol. 130:

[139649]507–518, 1995.

[139650]

[139651]

[139652]12930.

[139653]Rattner, J. B.; Rao, A.; Fritzler, M. J.; Valencia, D. W.; Yen,

[139654]T. J.: CENP-F is a ca 400 kDa kinetochore protein that exhibits a

[139655]cell-cycle dependent localization. Cell Motil. Cytoskeleton 26:

[139656]214-226, 1993.

[139657]

[139658]

[139659]12931.

[139660]Panagopoulos, I.; Mencinger, M.; Dietrich, C. U.; Bjerkehan,

[139661]B.; Saeter, G.; Mertens, F.; Mandahl, N.; Heim, S.: Fusion of the

[139662]RBP56 and CHN genes in extraskeletal myxoid chondrosarcomas with translocation

[139663]t(9;17)(q22;q11). Oncogene 18: 7594-7598, 1999.

[139664]

[139665]

[139666]12932.

[139667]Gassmann, M.; Casagrande, F.; Orioli, D.; Simon, H.; Lai, C.; Klein,

[139668]R.; Lemke, G.: Aberrant neural and cardiac development in mice lacking

[139669]the ErbB4 neuregulin receptor. Nature 378: 390–394, 1995.

[139670]

[139671]

[139672]12933.

[139673]Golding, J. P.; Trainor, P.; Krumlauf, R.; Gassmann, M.: Defects

[139674]in pathfinding by cranial neural crest cells in mice lacking the neuregulin

[139675]receptor ErbB4. Nature Cell Biol. 2: 103–109, 2000.

[139676]

[139677]

[139678]12934.

[139679]Zimonjic, D. B.; Alimandi, M.; Miki, T.; Popescu, N. C.; Kraus,

[139680]M. H.: Localization of the human HER4/erbB-4 gene to
chromosome 2. Oncogene 10:

[139681]1235-1237, 1995.

[139682]

[139683]

[139684]12935.

[139685]Adibi, S. A.: The oligopeptide transporter (Pept-1) in hu-
man intestine:

[139686]biology and function. Gastroenterology 113: 332-340,
1997.

[139687]

[139688]

[139689]12936.

[139690]Fei, Y. -J.; Kanai, Y.; Nussberger, S.; Ganapathy, V.;
Leibach,

[139691]F. H.; Romero, M. F.; Singh, S. K.; Boron, W. F.; Hediger, M.
A.:

[139692]Expression cloning of a mammalian proton-coupled
oligopeptide transporter. Nature 368:

[139693]563-566, 1994.

[139694]

[139695]

[139696]12937.

[139697]Liang, R.; Fei, Y.-J.; Prasad, P. D.; Ramamoorthy, S.; Han, H.;

[139698]Yang-Feng, T. L.; Hediger, M. A.; Ganapathy, V.; Leibach, F. H.:

[139699]Human intestinal H(+)/peptide cotransporter: cloning, functional expression,

[139700]and chromosomal localization. J. Biol. Chem. 270: 6456-6463, 1995.

[139701]

[139702]

[139703]12938.

[139704]Heiber, M.; Docherty, J. M.; Shah, G.; Nguyen, T.; Cheng, R.; Heng,

[139705]H. H. Q.; Marchese, A.; Tsui, L.-C.; Shi, X.; George, S. R.; O'Dowd,

[139706]B. F.: Isolation of three novel human genes encoding G protein-coupled

[139707]receptors. DNA Cell Biol. 14: 25-35, 1995.

[139708]

[139709]

[139710]12939.

[139711]Mahadevan, M. S.; Baird, S.; Bailly, J. E.; Shutler, G. G.;
Sabourin,

[139712]L. A.; Tsilfidis, C.; Neville, C. E.; Narang, M.; Korneluk, R.
G.

[139713]: Isolation of a novel G protein-coupled receptor (GPR4)
localized

[139714]to chromosome 19q13.3. Genomics 30: 84-88, 1995.

[139715]

[139716]

[139717]12940.

[139718]Ye, R. D.; Prossnitz, E. R.; Zou, A.; Cochrane, C. G.: Char-
acterization

[139719]of a human cDNA that encodes a functional receptor for
platelet activating

[139720]factor. Biochem. Biophys. Res. Commun. 180: 105-111,
1991.

[139721]

[139722]

[139723]12941.

[139724]Hossain, A.; Saunders, G. F.: The human sex-determining
gene

[139725]SRY is a direct target of WT1. J. Biol. Chem. 276:

16817-16823,

[139726]2001.

[139727]

[139728]

[139729]12942.

[139730]Iida, T.; Nakahori, Y.; Komaki, R.; Mori, E.; Hayashi, N.;
Tsutsumi,

[139731]O.; Taketani, Y.; Nakagome, Y.: A novel missense muta-
tion in the

[139732]HMG box of the SRY gene in a patient with XY sex rever-
sal. Hum. Molec.

[139733]Genet. 3: 1437-1438, 1994.

[139734]

[139735]

[139736]12943.

[139737]Jager, R. J.; Harley, V. R.; Pfeiffer, R. A.; Goodfellow, P. N.;

[139738]Scherer, G.: A familial mutation in the testis-determining
gene SRY

[139739]shared by both sexes. Hum. Genet. 90: 350–355, 1992.

[139740]

[139741]

[139742]12944.

[139743]Kawakura, K.; Miyake, Y.–I.; Murakami, R.–K.; Kondoh, S.;
Hirata,

[139744]T.–I.; Kaneda, Y.: Deletion of the SRY region on the Y
chromosome

[139745]detected in bovine gonadal hypoplasia (XY female) by PCR.
Cytogenet.

[139746]Cell Genet. 72: 183–184, 1996.

[139747]

[139748]

[139749]12945.

[139750]Lahr, G.; Maxson, S. C.; Mayer, A.; Just, W.; Pilgrim, C.;
Reisert,

[139751]I.: Transcription of the Y chromosomal gene, Sry, in adult

mouse

[139752]brain. Molec. Brain Res. 33: 179–182, 1995.

[139753]

[139754]

[139755]12946.

[139756]Li, B.; Zhang, W.; Chan, G.; Jancso–Radek, A.; Liu, S.;
Weiss,

[139757]M. A.: Human sex reversal due to impaired nuclear localization of

[139758]SRY: a clinical correlation. J. Biol. Chem. 276:
46480–46484, 2001.

[139759]

[139760]

[139761]12947.

[139762]Lovell–Badge, R.; Robertson, E.: XY female mice resulting
from

[139763]a heritable mutation in the primary testis–determining
gene, Tdy. Development 109:

[139764]635–646, 1990.

[139765]

[139766]

[139767]12948.

[139768]Margarit, E.; Coll, M. D.; Oliva, R.; Gomez, D.; Soler, A.;
Ballesta,

[139769]F.: SRY gene transferred to the long arm of the X chromo-
some in a

[139770]Y-positive XX true hermaphrodite. Am. J. Med. Genet. 90:
25-28,

[139771]2000.

[139772]

[139773]

[139774]12949.

[139775]Mittwoch, U.: Sex determination and sex reversal: geno-
type, phenotype,

[139776]dogma and semantics. Hum. Genet. 89: 467-479, 1992.

[139777]

[139778]

[139779]12950.

[139780]Muller, J.; Schwartz, M.; Skakkebaek, N. E.: Analysis of the

[139781]sex-determining region of the Y-chromosome (SRY) in
sex reversed patients:

[139782]point-mutation in SRY causing sex-reversion in a 46,XY
female. J.

[139783]Clin. Endocr. Metab. 75: 331-333, 1992.

[139784]

[139785]

[139786]12951.

[139787]Peters, R.; King, C.-Y.; Ukiyama, E.; Falsafi, S.; Donahoe, P.

[139788]K.; Weiss, M. A.: An SRY mutation causing human sex re-
versal resolves

[139789]a general mechanism of structure-specific DNA recogni-
tion: application

[139790]to the four-way DNA junction. Biochemistry 34:
4569-4576, 1995.

[139791]

[139792]

[139793]12952.

[139794]Schaffler, A.; Barth, N.; Winkler, K.; Zietz, B.; Rummele, P.;

[139795]Knuchel, R.; Scholmerich, J.; Palitzsch, K.-D.: Identification

of

[139796]a new missense mutation (gly95glu) in a highly conserved codon within

[139797]the high-mobility group box of the sex-determining region Y gene:

[139798]report on a 46,XY female with gonadal dysgenesis and yolk-sac tumor. J.

[139799]Clin. Endocr. Metab. 85: 2287-2292, 2000.

[139800]

[139801]

[139802]12953.

[139803]Schmitt-Ney, M.; Thiele, H.; Kaltwasser, P.; Bardoni, B.; Cisternino,

[139804]M.; Scherer, G.: Two novel SRY missense mutations reducing DNA binding

[139805]identified in XY females and their mosaic fathers. Am. J. Hum. Genet. 56:

[139806]862-869, 1995.

[139807]

[139808]

[139809]12954.

[139810]Sinclair, A. H.; Berta, P.; Palmer, M. S.; Hawkins, R.; Griffiths,

[139811]B. L.; Smith, M.; Foster, J. W.; Frischauf, A.-M.; Lovell-Badge, R.;

[139812]Goodfellow, P. N.: A gene from the human sex-determining region encodes

[139813]a protein with homology to a conserved DNA-binding motif.. Nature 346:

[139814]240-245, 1990.

[139815]

[139816]

[139817]12955.

[139818]Su, H.; Lau, Y.-F. C.: Identification of the transcriptional

[139819]unit, structural organization, and promoter sequence of the human

[139820]sex-determining region Y (SRY) gene, using a reverse genetic approach. Am.

[139821]J. Hum. Genet. 52: 24-38, 1993.

[139822]

[139823]

[139824]12956.

[139825]Tucker, P. K.; Lundrigan, B. L.: Rapid evolution of the sex
determining

[139826]locus in Old World mice and rats. Nature 364: 715–717,
1993.

[139827]

[139828]

[139829]12957.

[139830]Uehara, S.; Funato, T.; Yaegashi, N.; Suzuki, H.; Sato, J.;
Sasaki,

[139831]T.; Yajima, A.: SRY mutation and tumor formation on the
gonads of

[139832]XY pure gonadal dysgenesis patients. Cancer Genet. Cyto-
genet. 113:

[139833]78–84, 1999.

[139834]

[139835]

[139836]12958.

[139837]Uehara, S.; Hashiyada, M.; Sato, K.; Nata, M.; Funato, T.;
Okamura,

[139838]K.: Complete XY gonadal dysgenesis and aspects of the
SRY genotype

[139839]and gonadal tumor formation. J. Hum. Genet. 47:
279–284, 2002.

[139840]

[139841]

[139842]12959.

[139843]Veitia, R.; Ion, A.; Barbaux, S.; Jobling, M. A.; Souleyreau,

[139844]N.; Ennis, K.; Ostrer, H.; Tosi, M.; Meo, T.; Chibani, J.; Fel-
lous,

[139845]M.; McElreavey, K.: Mutations and sequence variants in the
testis-determining

[139846]region of the Y chromosome in individuals with a 46,XY
female phenotype. Hum.

[139847]Genet. 99: 648–652, 1997.

[139848]

[139849]

[139850]12960.

[139851]Werner, M. H.; Huth, J. R.; Gronenborn, A. M.; Clore, G. M.:

[139852]Molecular basis of human 46X,Y (sic) sex reversal revealed

from the

[139853]three-dimensional solution structure of the human SRY-DNA complex. Cell 81:

[139854]705-714, 1995.

[139855]

[139856]

[139857]12961.

[139858]Whitfield, L. S.; Lovell-Badge, R.; Goodfellow, P. N.: Rapid
[139859]sequence evolution of the mammalian sex-determining
gene SRY. Nature 364:

[139860]713-715, 1993.

[139861]

[139862]

[139863]12962.

[139864]Zeng, Y.; Ren, Z.; Zhang, M.; Huang, Y.; Zeng, F.; Huang,
S.:

[139865]A new de novo mutation (A113T) in HMG box of the SRY
gene leads to

[139866]XY gonadal dysgenesis. J. Med. Genet. 30: 655-657,
1993.

[139867]

[139868]

[139869]12963.

[139870]Zenteno, J. C.; Lopez, M.; Vera, C.; Mendez, J. P.; Kofman-
Alfaro,

[139871]S.: Two SRY-negative XX male brothers without genital
ambiguity. Hum.

[139872]Genet. 100: 606-610, 1997.

[139873]

[139874]

[139875]12964.

[139876]Henkemeyer, M.; Orioli, D.; Henderson, J. T.; Saxton, T.
M.; Roder,

[139877]J.; Pawson, T.; Klein, R.: Nuk controls pathfinding of com-
missural

[139878]axons in the mammalian central nervous system. Cell 86:
35-46, 1996.

[139879]

[139880]

[139881]12965.

- [139882] Ikegaki, N.; Tang, X. X.; Liu, X.-G.; Biegel, J. A.; Allen, C.;
- [139883] Yoshioka, A.; Sulman, E. P.; Brodeur, G. M.; Pleasure, D. E.:
Molecular
[139884] characterization and chromosomal localization of DRT
(EPHT3): a developmentally
[139885] regulated human protein-tyrosine kinase gene of the EPH
family. Hum.
[139886] Molec. Genet. 4: 2033-2045, 1995.
- [139887]
- [139888]
- [139889] 12966.
- [139890] Saito, T.; Seki, N.; Matsuda, Y.; Kitahara, M.; Murata, M.;
Kanda,
[139891] N.; Nomura, N.; Yamamoto, T.; Hori, T.: Identification of
the human
[139892] ERK gene as a putative receptor tyrosine kinase and its
chromosomal
[139893] localization to 1p36.1: a comparative mapping of human,
mouse, and
[139894] rat chromosomes. Genomics 26: 382-384, 1995.
- [139895]

[139896]

[139897]12967.

[139898]Wybenga-Groot, L. E.; Baskin, B.; Ong, S. H.; Tong, J.;

Pawson,

[139899]T.; Sicheri, F.: Structural basis for autoinhibition of the

EphB2

[139900]receptor tyrosine kinase by the unphosphorylated jux-
tamembrane region. Cell 106:

[139901]745–757, 2001.

[139902]

[139903]

[139904]12968.

[139905]Adams, J. W.; Sakata, Y.; Davis, M. G.; Sah, V. P.; Wang, Y.;

Liggett,

[139906]S. B.; Chien, K. R.; Brown, J. H.; Dorn, G. W., II: Enhanced

G-alpha-q

[139907]signaling: a common pathway mediates cardiac hypertro-
phy and apoptotic

[139908]heart failure. Proc. Nat. Acad. Sci. 95: 10140–10145,

1998.

[139909]

[139910]

[139911]12969.

[139912]Dong, Q.; Shenker, A.; Way, J.; Haddad, B. R.; Lin, K.;
Hughes,

[139913]M. R.; McBride, O. W.; Spiegel, A. M.; Battey, J.: Molecular
cloning

[139914]of human G-alpha(q) cDNA and chromosomal localization
of the G-alpha(q)

[139915]gene (GNAQ) and a processed pseudogene. Genomics 30:
470-475, 1995.

[139916]

[139917]

[139918]12970.

[139919]Gabbeta, J.; Yang, X.; Kowalska, M. A.; Sun, L.;
Dhanasekaran,

[139920]N.; Rao, A. K.: Platelet signal transduction defect with G-
alpha

[139921]subunit dysfunction and diminished G-alpha(q) in a pa-
tient with abnormal

[139922]platelet responses. Proc. Nat. Acad. Sci. 94: 8750-8755,
1997.

[139923]

[139924]

[139925]12971.

[139926]Offermanns, S.; Hashimoto, K.; Watanabe, M.; Sun, W.;
Kurihara,

[139927]H.; Thompson, R. F.; Inoue, Y.; Kano, M.; Simon, M. I.: Im-
paired

[139928]motor coordination and persistent multiple climbing fiber
innervation

[139929]of cerebellar Purkinje cells in mice lacking G- α -q.
Proc. Nat.

[139930]Acad. Sci. 94: 14089–14094, 1997.

[139931]

[139932]

[139933]12972.

[139934]Offermanns, S.; Toombs, C. F.; Hu, Y.-H.; Simon, M. I.:
Defective

[139935]platelet activation in G- α -q-deficient mice. Nature
389: 183–186,

[139936]1997.

[139937]

[139938]

[139939]12973.

[139940]Johnson, D. F.; Hamon, M.; Fischel-Ghodsian, N.: Characterization

[139941]of the human mitochondrial ribosomal S12 gene. Genomics 52: 363–368,

[139942]1998.

[139943]

[139944]

[139945]12974.

[139946]Shah, Z. H.; Migliosi, V.; Miller, S. C. M.; Wang, A.; Friedman,

[139947]T. B.; Jacobs, H. T.: Chromosomal locations of three human nuclear

[139948]genes (RPSM12, TUFM, and AFG3L1) specifying putative components of

[139949]the mitochondrial gene expression apparatus. Genomics 48: 384–388,

[139950]1998.

[139951]

[139952]

[139953]12975.

[139954]Honore, B.; Leffers, H.; Madsen, P.; Celis, J. E.: Interferon-
gamma

[139955]up-regulates a unique set of proteins in human ker-
atinocytes: molecular

[139956]cloning and expression of the cDNA encoding the RGD-
sequence-containing

[139957]protein IGUP I5111. Europ. J. Biochem. 218: 421-430,
1993.

[139958]

[139959]

[139960]12976.

[139961]Preckel, T.; Fung-Leung, W.-P.; Cai, Z.; Vitiello, A.; Salter-
Cid,

[139962]L.; Winqvist, O.; Wolfe, T. G.; Von Herrath, M.; Angulo, A.;
Ghazal,

[139963]P.; Lee, J.-D.; Fourie, A. M.; Wu, Y.; Pang, J.; Ngo, K.; Pe-
terson,

[139964]P. A.; Fruh, K.; Yang, Y.: Impaired immunoproteasome as-

sembly and

[139965]immune responses in PA28 $-/-$ mice. Science 286:
2162–2165, 1999.

[139966]

[139967]

[139968]12977.

[139969]Realini, C.; Dubiel, W.; Pratt, G.; Ferrell, K.; Rechsteiner, M.

[139970]: Molecular cloning and expression of a gamma-inter-
feron-inducible

[139971]activator of the multicatalytic protease. J. Biol. Chem. 269:
20727–20732,

[139972]1994.

[139973]

[139974]

[139975]12978.

[139976]Trachtulec, Z.; Forejt, J.: Synteny of orthologous genes
conserved

[139977]in mammals, snake, fly, nematode, and fission yeast.

Mammalian Genome 12:

[139978]227–231, 2001.

[139979]

[139980]

[139981]12979.

[139982]Reif, K.; Ekland, E. H.; Ohl, L.; Nakano, H.; Lipp, M.;

Forster,

[139983]R.; Cyster, J. G.: Balanced responsiveness to chemoattractants from

[139984]adjacent zones determines B-cell position. Nature 416: 94–99, 2002.

[139985]

[139986]

[139987]12980.

[139988]Blanco, G.; Brown, S. D. M.: Genetic mapping of protein kinase

[139989]C theta (Pkcq) to mouse chromosome 2. Mammalian Genome 8: 70–71,

[139990]1997.

[139991]

[139992]

[139993]12981.

[139994]Zhu, K.; Baudhuin, L. M.; Hong, G.; Williams, F. S.; Cristina,
[139995]K. L.; Kabarowski, J. H. S.; Witte, O. N.; Xu, Y.: Sphingo-
sylphosphorylcholine
[139996]and lysophosphatidylcholine are ligands for the G protein-
coupled
[139997]receptor GPR4. J. Biol. Chem. 276: 41325–41335, 2001.

[139998]

[139999]

[140000]12982.

[140001]Vinik, B. S.; Kay, E. S.; Fiedorek, F. T., Jr.: Mapping of the
[140002]MEK kinase gene (Mekk) to mouse chromosome 13 and
human chromosome

[140003]5. Mammalian Genome 6: 782–783, 1995.

[140004]

[140005]

[140006]12983.

[140007]Yujiri, T.; Sather, S.; Fanger, G. R.; Johnson, G. L.: Role of

[140008]MEKK1 in cell survival and activation of JNK and ERK path-
ways defined

[140009]by targeted gene disruption. Science 282: 1911–1914,
1998.

[140010]

[140011]

[140012]12984.

[140013]Arriza, J. L.; Weinberger, C.; Cerelli, G.; Glaser, T. M.; Han-
delin,

[140014]B. L.; Housman, D. E.; Evans, R. M.: Cloning of human
mineralocorticoid

[140015]receptor complementary DNA: structural and functional
kinship with

[140016]the glucocorticoid receptor. Science 237: 268–275, 1987.

[140017]

[140018]

[140019]12985.

[140020]Berger, S.; Bleich, M.; Schmid, W.; Cole, T. J.; Peters, J.;
Watanabe,

[140021]H.; Kriz, W.; Warth, R.; Greger, R.; Schutz, G.: Mineralocor-
ticoid

[140022]receptor knockout mice: pathophysiology of Na⁺
metabolism. Proc.

[140023]Nat. Acad. Sci. 95: 9424–9429, 1998.

[140024]

[140025]

[140026]12986.

[140027]Fan, Y.–S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[140028]M.; Nowak, N. J.; Shows, T. B.: The human mineralocorticoid receptor

[140029]gene (MLR) is located on chromosome 4 at q31.2. Cytogenet. Cell Genet. 52:

[140030]83–84, 1989.

[140031]

[140032]

[140033]12987.

[140034]Geller, D. S.; Farhi, A.; Pinkerton, N.; Fradley, M.; Moritz, M.;

[140035]Spitzer, A.; Meinke, G.; Tsai, F. T. F.; Sigler, P. B.; Lifton, R.

[140036]P.: Activating mineralocorticoid receptor mutation in hypertension

[140037]exacerbated by pregnancy. Science 289: 119–123, 2000.

[140038]

[140039]

[140040]12988.

[140041]Geller, D. S.; Rodriguez–Soriano, J.; Vallo Boado, A.;
Schifter,

[140042]S.; Bayer, M.; Chang, S. S.; Lifton, R. P.: Mutations in the
mineralocorticoid

[140043]receptor gene cause autosomal dominant pseudohypoal–
dosteronism type

[140044]I. Nature Genet. 19: 279–281, 1998.

[140045]

[140046]

[140047]12989.

[140048]Hellal–Levy, C.; Fagart, J.; Souque, A.; Wurtz, J.–M.; Moras,
D.;

[140049]Rafestin–Oblin, M.–E.: Crucial role of the H11–H12 loop in
stabilizing

[140050]the active conformation of the human mineralocorticoid
receptor. Molec.

[140051]Endocr. 14: 1210–1221, 2000.

[140052]

[140053]

[140054]12990.

[140055]Le Menuet, D.; Isnard, R.; Bichara, M.; Viengchareun, S.;
Muffat-Joly,

[140056]M.; Walker, F.; Zennaro, M.-C.; Lombes, M.: Alteration of
cardiac

[140057]and renal functions in transgenic mice overexpressing hu-
man mineralocorticoid

[140058]receptor. J. Biol. Chem. 276: 38911-38920, 2001.

[140059]

[140060]

[140061]12991.

[140062]Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor, J.

[140063]M.: Regional chromosomal assignment of the human min-
eralocorticoid

[140064]receptor gene to 4q31.1. Hum. Genet. 85: 130-132,
1990.

[140065]

[140066]

[140067]12992.

[140068]Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor,

[140069]J. M.: Regional chromosomal assignment of the human
mineralocorticoid

[140070]receptor gene to 4q31.1. (Abstract) Cytogenet. Cell Genet.
51: 1048,

[140071]1989.

[140072]

[140073]

[140074]12993.

[140075]Veenstra, G. J. C.; Weeks, D. L.; Wolffe, A. P.: Distinct roles

[140076]for TBP and TBP-like factor in early embryonic gene tran-
scription

[140077]in *Xenopus*. Science 290: 2312–2314, 2000.

[140078]

[140079]

[140080]12994.

[140081]Zuhlke, C.; Hellenbroich, Y.; Dalski, A.; Kononowa, N.; Ha-
genah,

[140082]J.; Vieregge, P.; Riess, O.; Klein, C.; Schwinger, E.: Different
ent

[140083]types of repeat expansion in the TATA-binding protein
gene are associated

[140084]with a new form of inherited ataxia. Europ. J. Hum. Genet.
9: 160–164,

[140085]2001.

[140086]

[140087]

[140088]12995.

[140089]Charest, A.; Wagner, J.; Muise, E. S.; Heng, H. H. Q.; Trem-
blay,

[140090]M. L.: Structure of the murine MPTP-PEST gene: genomic
organization

[140091]and chromosomal mapping. Genomics 28: 501–507,
1995.

[140092]

[140093]

[140094]12996.

[140095]Takekawa, M.; Itoh, F.; Hinoda, Y.; Adachi, M.; Ariyama, T.;
Inazawa,

[140096]J.; Imai, K.; Yachi, A.: Chromosomal localization of the protein

[140097]tyrosine phosphatase G1 gene and characterization of the aberrant

[140098]transcripts in human colon cancer cells. FEBS Lett. 339: 222–228,

[140099]1994.

[140100]

[140101]

[140102]12997.

[140103]Takekawa, M.; Itoh, F.; Hinoda, Y.; Arimura, Y.; Toyota, M.; Sekiya,

[140104]M.; Adachi, M.; Imai, K.; Yachi, A.: Cloning and characterization

[140105]of a human cDNA encoding a novel putative cytoplasmic protein–tyrosine–phosphatase. Biochem.

[140106]Biophys. Res. Commun. 189: 1223–1230, 1992.

[140107]

[140108]

[140109]12998.

[140110]Yang, Q.; Co, D.; Sommercorn, J.; Tonks, N. K.: Cloning
and expression
[140111]of PTP-PEST: a novel, human, nontransmembrane protein
tyrosine phosphatase. J.
[140112]Biol. Chem. 268: 6622–6628, 1993.

[140113]

[140114]

[140115]12999.

[140116]Coopman, P. J. P.; Do, M. T. H.; Barth, M.; Bowden, E. T.;
Hayes,
[140117]A. J.; Basyuk, E.; Blancato, J. K.; Vezza, P. R.; McLeskey, S.
W.;

[140118]Mangeat, P. H.; Mueller, S. C.: The Syk tyrosine kinase
suppresses
[140119]malignant growth of human breast cancer cells. Nature
406: 742–747,
[140120]2000.

[140121]

[140122]

[140123]13000.

[140124]Inatome, R.; Yanagi, S.; Takano, T.; Yamamura, H.: A critical

[140125]role for Syk in endothelial cell proliferation and migration. Biochem.

[140126]Biophys. Res. Commun. 286: 195–199, 2001.

[140127]

[140128]

[140129]13001.

[140130]Mocsai, A.; Zhou, M.; Meng, F.; Tybulewicz, V. L.; Lowell, C. A.

[140131]: Syk is required for integrin signaling in neutrophils. Immunity 16:

[140132]547–558, 2002.

[140133]

[140134]

[140135]13002.

[140136]Muller, B.; Cooper, L.; Terhorst, C.: Molecular cloning of the

[140137]human homologue to the pig protein–tyrosine kinase syk. Immunogenetics 39:

[140138]359–362, 1994.

[140139]

[140140]

[140141]13003.

[140142]Taniguchi, T.; Kobayashi, T.; Kondo, J.; Takahashi, K.;
Nakamura,

[140143]H.; Suzuki, J.; Nagai, K.; Yamada, T.; Nakamura, S.; Yama-
mura, H.

[140144]: Molecular cloning of a porcine gene syk that encodes a
72-kDa protein-tyrosine

[140145]kinase showing high susceptibility to proteolysis. J. Biol.
Chem. 266:

[140146]15790–15796, 1991.

[140147]

[140148]

[140149]13004.

[140150]Zioncheck, T. F.; Harrison, M. L.; Isaacson, C. C.; Geahlen,
R.

[140151]L.: Generation of an active protein-tyrosine kinase from
lymphocytes

[140152]by proteolysis. J. Biol. Chem. 263: 19195–19202, 1988.

[140153]

[140154]

[140155]13005.

[140156]Satre, M. A.; Zgombic-Knight, M.; Duester, G.: The complete structure

[140157]of human class IV alcohol dehydrogenase (retinol dehydrogenase) determined

[140158]from the ADH7 gene. J. Biol. Chem. 269: 15606–15612, 1994.

[140159]

[140160]

[140161]13006.

[140162]Yokoyama, H.; Baraona, E.; Lieber, C. S.: Molecular cloning and

[140163]chromosomal localization of the ADH7 gene encoding human class IV

[140164](sigma) ADH. Genomics 31: 243–245, 1996.

[140165]

[140166]

[140167]13007.

[140168]Zgombic-Knight, M.; Foglio, M. H.; Duester, G.: Genomic structure

[140169]and expression of the ADH7 gene encoding human class IV alcohol dehydrogenase,

[140170]the form most efficient for retinol metabolism in vitro. J. Biol.

[140171]Chem. 270: 4305-4311, 1995.

[140172]

[140173]

[140174]13008.

[140175]Bashir, M. M.; Abrams, W. R.; Tucker, T.; Sellinger, B.; Budarf,

[140176]M.; Emanuel, B.; Rosenbloom, J.: Molecular cloning and characterization

[140177]of the bovine and human tuftelin genes. Connect. Tissue Res. 39:

[140178]13-24, 1998.

[140179]

[140180]

[140181]13009.

[140182]Deutsch, D.: Structure and function of enamel gene products. Anat.

[140183]Rec. 224: 189–210, 1989.

[140184]

[140185]

[140186]13010.

[140187]Beguín, P.; Nagashima, K.; Gonoï, T.; Shibasaki, T.; Takahashi,

[140188]K.; Kashima, Y.; Ozaki, N.; Geering, K.; Iwanaga, T.; Seino, S.:

[140189]Regulation of Ca(2+) channel expression at the cell surface by the

[140190]small G-protein kir/Gem. Nature 411: 701–706, 2001.

[140191]

[140192]

[140193]13011.

[140194]Maguire, J.; Santoro, T.; Jensen, P.; Siebenlist, U.; Yewdell,

[140195]J.; Kelly, K.: GEM: an induced, immediate early protein be-

longing

[140196]to the Ras family. Science 265: 241–244, 1994.

[140197]

[140198]

[140199]13012.

[140200]Santoro, T.; Maguire, J.; McBride, O. W.; Avraham, K. B.;
Copeland,

[140201]N. G.; Jenkins, N. A.; Kelly, K.: Chromosomal organization
and transcriptional

[140202]regulation of human GEM and localization of the human
and mouse Gem

[140203]loci encoding an inducible Ras–like protein. Genomics 30:
558–564,

[140204]1995.

[140205]

[140206]

[140207]13013.

[140208]Inoue, I.; Taniuchi, I.; Kitamura, D.; Jenkins, N. A.; Gilbert,

[140209]D. J.; Copeland, N. G.; Watanabe, T.: Characteristics of the
mouse

[140210]genomic histamine H1 receptor gene. Genomics 36:
178–181, 1996.

[140211]

[140212]

[140213]13014.

[140214]Le Coniat, M.; Traiffort, E.; Ruat, M.; Arrang, J.–M.; Berger,

[140215]R.: Chromosomal localization of the human histamine
H1–receptor gene. Hum.

[140216]Genet. 94: 186–188, 1994.

[140217]

[140218]

[140219]13015.

[140220]Ma, R. Z.; Gao, J.; Meeker, N. D.; Fillmore, P. D.; Tung, K.
S.

[140221]K.; Watanabe, T.; Zachary, J. F.; Offner, H.; Blankenhorn, E.
P.;

[140222]Teuscher, C.: Identification of Bphs, an autoimmune dis-
ease locus,

[140223]as histamine receptor H–1. Science 297: 620–623, 2002.

[140224]

[140225]

[140226]13016.

[140227]Yamashita, M.; Fukui, H.; Sugama, K.; Horio, Y.; Ito, S.;
Mizuguchi,

[140228]H.; Wada, H.: Expression cloning of a cDNA encoding the
bovine histamine

[140229]H1 receptor. Proc. Nat. Acad. Sci. 88: 11515–11519,
1991.

[140230]

[140231]

[140232]13017.

[140233]Gaudino, G.; Follenzi, A.; Naldini, L.; Collesi, C.; Santoro,
M.;

[140234]Gallo, K. A.; Godowski, P. J.; Comoglio, P. M.: RON is a
heterodimeric

[140235]tyrosine kinase receptor activated by the HGF homologue
MSP. EMBO

[140236]J. 13: 3524–3532, 1994.

[140237]

[140238]

[140239]13018.

[140240]Muraoka, R. S.; Sun, W. Y.; Colbert, M. C.; Waltz, S. E.;

Witte,

[140241]D. P.; Degen, J. L.; Degen, S. J. F.: The Ron/STK receptor
tyrosine

[140242]kinase is essential for peri-implantation development in
the mouse. J.

[140243]Clin. Invest. 103: 1277–1285, 1999.

[140244]

[140245]

[140246]13019.

[140247]Ronsin, C.; Muscatelli, F.; Mattei, M.–G.; Breathnach, R.: A
novel

[140248]putative receptor protein tyrosine kinase of the met fam-
ily. Oncogene 8:

[140249]1195–1202, 1993.

[140250]

[140251]

[140252]13020.

[140253]Santoro, M. M.; Collesi, C.; Grisendi, S.; Gaudino, G.; Co-

moglio,

[140254]P. M.: Constitutive activation of the RON gene promotes
invasive

[140255]growth but not transformation. Molec. Cell. Biol. 16:
7072–7083,

[140256]1996.

[140257]

[140258]

[140259]13021.

[140260]Wang, M.–H.; Ronsin, C.; Gesnel, M.–C.; Coupey, L.; Skeel,
A.;

[140261]Leonard, E. J.; Breathnach, R.: Identification of the RON
gene product

[140262]as the receptor for the human macrophage stimulating
protein. Science 266:

[140263]117–119, 1994.

[140264]

[140265]

[140266]13022.

[140267]Bauer, S.; Groh, V.; Wu, J.; Steinle, A.; Phillips, J. H.; Lanier,

[140268]L. L.; Spies, T.: Activation of NK cells and T cells by
NKG2D, a

[140269]receptor for stress-inducible MICA. Science 285:
727–729, 1999.

[140270]

[140271]

[140272]13023.

[140273]Diefenbach, A.; Jensen, E. R.; Jamieson, A. M.; Raulet, D.
H.:

[140274]Rae1 and H60 ligands of the NKG2D receptor stimulate
tumour immunity. Nature 413:

[140275]165–171, 2001.

[140276]

[140277]

[140278]13024.

[140279]Girardi, M.; Oppenheim, D. E.; Steele, C. R.; Lewis, J. M.;
Glusac,

[140280]E.; Filler, R.; Hobby, P.; Sutton, B.; Tigelaar, R. E.; Hayday,
A.

[140281]C.: Regulation of cutaneous malignancy by gamma–delta
T cells. Science 294:

[140282]605–609, 2001.

[140283]

[140284]

[140285]13025.

[140286]Groh, V.; Rhinehart, R.; Randolph-Habecker, J.; Topp, M.
S.; Riddell,

[140287]S. R.; Spies, T.: Costimulation of CD8–alpha–beta T cells
by NKG2D

[140288]via engagement by MIC induced on virus–infected cells.
Nature Immun. 2:

[140289]255–260, 2001.

[140290]

[140291]

[140292]13026.

[140293]Groh, V.; Steinle, A.; Bauer, S.; Spies, T.: Recognition of
stress–induced

[140294]MHC molecules by intestinal epithelial gamma–delta T
cells. Science 279:

[140295]1737–1740, 1998.

[140296]

[140297]

[140298]13027.

[140299]Groh, V.; Wu, J.; Yee, C.; Spies, T.: Tumour-derived soluble
MIC

[140300]ligands impair expression of NKG2D and T-cell activation.

Nature 419:

[140301]734–738, 2002.

[140302]

[140303]

[140304]13028.

[140305]Li, P.; Morris, D. L.; Willcox, B. E.; Steinle, A.; Spies, T.;

[140306]Strong, R. K.: Complex structure of the activating im-
munoreceptor

[140307]NKG2D and its MHC class I-like ligand MICA. Nature Im-
mun. 2: 443–451,

[140308]2001.

[140309]

[140310]

[140311]13029.

[140312]Nakashima, T.; Sekiguchi, T.; Kuraoka, A.; Fukushima, K.;
Shibata,

[140313]Y.; Komiyama, S.; Nishimoto, T.: Molecular cloning of a
human cDNA

[140314]encoding a novel protein, DAD1, whose defect causes
apoptotic cell

[140315]death in hamster BHK21 cells. Molec. Cell. Biol. 13:
6367–6374,

[140316]1993.

[140317]

[140318]

[140319]13030.

[140320]Ylug, I. G.; See, C. G.; Fisher, E. M. C.: The DAD1 protein,
whose

[140321]defect causes apoptotic cell death, maps to human chro-
mosome 14. Genomics 26:

[140322]433–435, 1995. Note: Erratum: Genomics: 28: 129 only,
1995.

[140323]

[140324]

[140325]13031.

[140326]Yulug, I. G.; See, C. G.; Fisher, E. M. C.: The DAD1 protein,
[140327]whose defect causes apoptotic cell death, maps to human
chromosome

[140328]14. Genomics 26: 433–435, 1995.

[140329]

[140330]

[140331]13032.

[140332]Jepsen, K. J.; Wu, F.; Peragallo, J. H.; Paul, J.; Roberts, L.;

[140333]Ezura, Y.; Oldberg, A.; Birk, D. E.; Chakravarti, S.: A syn-
drome

[140334]of joint laxity and impaired tendon integrity in lumican-
and fibromodulin-deficient

[140335]mice. J. Biol. Chem. 277: 35532–35540, 2002.

[140336]

[140337]

[140338]13033.

[140339]Sztrolovics, R.; Chen, X.-N.; Grover, J.; Roughley, P. J.; Ko-
renberg,

[140340]J. R.: Localization of the human fibromodulin gene (FMOD)

to chromosome

[140341]1q32 and completion of the cDNA sequence. Genomics
23: 715–717,

[140342]1994.

[140343]

[140344]

[140345]13034.

[140346]Dalton, S.; Treisman, R.: Characterization of SAP–1, a pro-
tein

[140347]recruited by serum response factor to the c–fos serum re-
sponse element. Cell 68:

[140348]597–612, 1992.

[140349]

[140350]

[140351]13035.

[140352]Mo, Y.; Vaessen, B.; Johnston, K.; Marmorstein, R.: Struc-
tures

[140353]of SAP–1 bound to DNA targets from the E74 and c–fos
promoters: insights

[140354]into DNA sequence discrimination by Ets proteins. Molec.

Cell 2:

[140355]201–212, 1998.

[140356]

[140357]

[140358]13036.

[140359]Shipley, J.; Sheer, D.; Dalton, S.; Treisman, R.; Patel, K.:

Mapping

[140360]of the human SAP1 (SRF accessory protein 1) gene and

SAP2, a gene

[140361]encoding a related protein, to chromosomal bands 1q32

and 12q23, respectively. Genomics 23:

[140362]710–711, 1994.

[140363]

[140364]

[140365]13037.

[140366]Giovane, A.; Pintzas, A.; Maira, S.–M.; Sobieszczuk, P.;

Wasylyk,

[140367]B.: Net, a new ets transcription factor that is activated by

Ras. Genes

[140368]Dev. 8: 1502–1513, 1994.

[140369]

[140370]

[140371]13038.

[140372]Lopez, M.; Oettgen, P.; Akbarali, Y.; Dendorfer, U.; Libermann,

[140373]T. A.: ERP, a new member of the ets transcription factor/
oncoprotein

[140374]family: cloning, characterization, and differential expression during

[140375]B-lymphocyte development. Molec. Cell. Biol. 14:
3292-3309, 1994.

[140376]

[140377]

[140378]13039.

[140379]Kelner, G. S.; Kennedy, J.; Bacon, K. B.; Kleyensteuber, S.;
Largaespada,

[140380]D. A.; Jenkins, N. A.; Copeland, N. G.; Bazan, J. F.; Moore,
K. W.;

[140381]Schall, T. J.; Zlotnik, A.: Lymphotactin: a cytokine that represents

[140382]a new class of chemokine. Science 266: 1395-1399,

1994.

[140383]

[140384]

[140385]13040.

[140386]Kennedy, J.; Kelner, G. S.; Kleyensteuber, S.; Schall, T. J.;
Weiss,

[140387]M. C.; Yssel, H.; Schneider, P. V.; Cocks, B. G.; Bacon, K. B.;
Zlotnik,

[140388]A.: Molecular cloning and functional characterization of
human lymphotactin. J.

[140389]Immun. 155: 203–209, 1995.

[140390]

[140391]

[140392]13041.

[140393]Muller, S.; Dorner, B.; Korthauer, U.; Mages, H. W.;
D'Apuzzo,

[140394]M.; Senger, G.; Kroczeck, R. A.: Cloning of ATAC, an activa-
tion-induced,

[140395]chemokine-related molecule exclusively expressed in
CD8+ T lymphocytes. Europ.

[140396]J. Immun. 25: 1744–1748, 1995.

[140397]

[140398]

[140399]13042.

[140400]Yoshida, T.; Imai, T.; Kakizaki, M.; Nishimura, M.; Takagi, S.;

[140401]Yoshie, O.: Identification of single C motif-1/lymphotactin receptor

[140402]XCR1. J. Biol. Chem. 273: 16551-16554, 1998.

[140403]

[140404]

[140405]13043.

[140406]Yoshida, T.; Imai, T.; Kakizaki, M.; Nishimura, M.; Yoshie, O.

[140407]: Molecular cloning of a novel C or gamma type chemokine, SCM-1. FEBS

[140408]Lett. 360: 155-159, 1995.

[140409]

[140410]

[140411]13044.

[140412]Yoshida, T.; Imai, T.; Takagi, S.; Nishimura, M.; Ishikawa, I.;

[140413]Yaoi, T.; Yoshie, O.: Structure and expression of two highly related

[140414]genes encoding SCM-1/human lymphotactin. FEBS Lett. 395: 82-88,

[140415]1996.

[140416]

[140417]

[140418]13045.

[140419]Ikeuchi, T.; Asaka, T.; Saito, M.; Tanaka, H.; Higuchi, S.; Tanaka,

[140420]K.; Saida, K.; Uyama, E.; Mizusawa, H.; Fukuhara, N.; Nonaka, I.;

[140421]Takamori, M.; Tsuji, S.: Gene locus for autosomal recessive distal

[140422]myopathy with rimmed vacuoles maps to chromosome 9. Ann. Neurol. 41:

[140423]432-437, 1997.

[140424]

[140425]

[140426]13046.

[140427]Erickson, J. D.; Varoqui, H.; Schafer, M. K.-H.; Modi, W.;
Diebler,

[140428]M.-F.; Weihe, E.; Rand, J.; Eiden, L. E.; Bonner, T. I.; Usdin,
T.

[140429]B.: Functional identification of a vesicular acetylcholine
transporter

[140430]and its expression from a 'cholinergic' gene locus. J. Biol.
Chem. 269:

[140431]21929-21932, 1994.

[140432]

[140433]

[140434]13047.

[140435]Bachvarov, D. R.; Hess, J. F.; Menke, J. G.; Larrivee, J.-F.;
Marceau,

[140436]F.: Structure and genomic organization of the human B(1)
receptor

[140437]gene for kinins (BDKRB1). Genomics 33: 374-381, 1996.

[140438]

[140439]

[140440]13048.

[140441]Chai, K. X.; Ni, A.; Wang, D.; Ward, D. C.; Chao, J.; Chao, L.

[140442]: Genomic DNA sequence, expression, and chromosomal
localization of

[140443]the human B1 bradykinin receptor gene BDKRB1. Ge-
nomics 31: 51-57,

[140444]1996.

[140445]

[140446]

[140447]13049.

[140448]MacNeil, T.; Bierilo, K. K.; Menke, J. G.; Hess, J. F.: Cloning

[140449]and pharmacological characterization of a rabbit
bradykinin B-1 receptor. Biochim.

[140450]Biophys. Acta 1264: 223-228, 1995.

[140451]

[140452]

[140453]13050.

[140454]Menke, J. G.; Borkowski, J. A.; Bierilo, K. K.; MacNeil, T.;
Derrick,

[140455]A. W.; Schneck, K. A.; Ransom, R. W.; Strader, C. D.; Line-
meyer, D.

[140456]L.; Hess, J. F.: Expression cloning of a human B1
bradykinin receptor. J.

[140457]Biol. Chem. 269: 21583–21586, 1994.

[140458]

[140459]

[140460]13051.

[140461]Yang, X.; Polgar, P.: Genomic structure of the human
bradykinin

[140462]B(1) receptor gene and preliminary characterization of its
regulatory

[140463]regions. Biochem. Biophys. Res. Commun. 222: 718–725,
1996.

[140464]

[140465]

[140466]13052.

[140467]Sanders, J.; Maassen, J. A.; Amons, R.; Moller, W.: Nu-
cleotide

[140468]sequence of human elongation factor-1-beta cDNA. Nu-
cleic Acids Res. 19:

[140469]4551, 1991.

[140470]

[140471]

[140472]13053.

[140473]Tanahashi, H.; Tabira, T.: Alzheimer's disease-associated
presenilin

[140474]2 interacts with DRAL, an LIM-domain protein. Hum.
Molec. Genet. 9:

[140475]2281–2289, 2000.

[140476]

[140477]

[140478]13054.

[140479]Erkinheimo, T.–L.; Saukkonen, K.; Narko, K.; Jalkanen, J.;
Ylikorkala,

[140480]O.; Ristimäki, A.: Expression of cyclooxygenase-2 and
prostanoid

[140481]receptors by human myometrium. J. Clin. Endocr. Metab.
85: 3468–3475,

[140482]2000.

[140483]

[140484]

[140485]13055.

[140486]Hla, T.; Neilson, K.: Human cyclooxygenase-2 cDNA. Proc.
Nat.

[140487]Acad. Sci. 89: 7384-7388, 1992.

[140488]

[140489]

[140490]13056.

[140491]Jones, D. A.; Carlton, D. P.; McIntyre, T. M.; Zimmerman,
G. A.;

[140492]Prescott, S. M.: Molecular cloning of human prostaglandin
endoperoxide

[140493]synthase type II and demonstration of expression in re-
sponse to cytokines. J.

[140494]Biol. Chem. 268: 9049-9054, 1993.

[140495]

[140496]

[140497]13057.

[140498]Kraemer, S. A.; Meade, E. A.; DeWitt, D. L.: Prostaglandin

endoperoxide

[140499] synthase gene structure: identification of the transcriptional start

[140500] site and 5-prime-flanking regulatory sequences. Arch. Biochem. Biophys. 293:

[140501] 391–400, 1992.

[140502]

[140503]

[140504] 13058.

[140505] Lassus, P.; Wolff, H.; Andersson, S.: Cyclooxygenase-2 in human

[140506] perinatal lung. Pediat. Res. 47: 602–605, 2000.

[140507]

[140508]

[140509] 13059.

[140510] Lim, H.; Paria, B. C.; Das, S. K.; Dinchuk, J. E.; Langenbach,

[140511] R.; Trzaskos, J. M.; Dey, S. K.: Multiple female reproductive failures

[140512] in cyclooxygenase 2-deficient mice. Cell 91: 197–208, 1997.

[140513]

[140514]

[140515]13060.

[140516]Morrow, J. A.; Collie, I. T.; Dunbar, D. R.; Walker, G. B.;
Shahid,

[140517]M.; Hill, D. R.: Molecular cloning and functional expression
of the

[140518]human glycine transporter GlyT2 and chromosomal localisation of the

[140519]gene in the human genome. FEBS Lett. 439: 334–340,
1998.

[140520]

[140521]

[140522]13061.

[140523]Jacquemin, P.; Durviaux, S. M.; Jensen, J.; Godfraind, C.;
Gradwohl,

[140524]G.; Guillemot, F.; Madsen, O. D.; Carmeliet, P.; Dewerchin,
M.; Collen,

[140525]D.; Rousseau, G. G.; Lemaigre, F. P.: Transcription factor
hepatocyte

[140526]nuclear factor 6 regulates pancreatic endocrine cell differ–

entiation

[140527]and controls expression of the proendocrine gene ngn3.

Molec. Cell.

[140528]Biol. 20: 4445–4454, 2000.

[140529]

[140530]

[140531]13062.

[140532]Jacquemin, P.; Lannoy, V. J.; Rousseau, G. G.; Lemaigre, F.
P.

[140533]: OC–2, a novel mammalian member of the ONECUT class
of homeodomain

[140534]transcription factors whose function in liver partially over-
laps with

[140535]that of hepatocyte nuclear factor–6. J. Biol. Chem. 274:
2665–2671,

[140536]1999.

[140537]

[140538]

[140539]13063.

[140540]Lemaigre, F. P.; Durviaux, S. M.; Truong, O.; Lannoy, V. J.;

Hsuan,

[140541]J. J.; Rousseau, G. G.: Hepatocyte nuclear factor 6, a transcription

[140542]factor that contains a novel type of homeodomain and a single cut

[140543]domain. Proc. Nat. Acad. Sci. 93: 9460–9464, 1996.

[140544]

[140545]

[140546]13064.

[140547]Pierreux, C. E.; Stafford, J.; Demonte, D.; Scott, D. K.; Vandenhaute,

[140548]J.; O'Brien, R. M.; Granner, D. K.; Rousseau, G. G.; Lemaignre, F.

[140549]P.: Antigluocorticoid activity of hepatocyte nuclear factor-6. Proc.

[140550]Nat. Acad. Sci. 96: 8961–8966, 1999.

[140551]

[140552]

[140553]13065.

[140554]Samadani, U.; Costa, R. H.: The transcriptional activator

hepatocyte

[140555]nuclear factor 6 regulates liver gene expression. Molec.

Cell. Biol. 16:

[140556]6273–6284, 1996.

[140557]

[140558]

[140559]13066.

[140560]Vaisse, C.; Kim, J.; Espinosa, R., III; Le Beau, M. M.; Stoffel,

[140561]M.: Pancreatic islet expression studies and polymorphic

DNA markers

[140562]in the genes encoding hepatocyte nuclear factor–3–alpha,

–3–beta,

[140563]–4–gamma, and –6. Diabetes 46: 1364–1367, 1997.

[140564]

[140565]

[140566]13067.

[140567]Iacobazzi, V.; Palmieri, F.; Runswick, M. J.; Walker, J. E.:

Sequences

[140568]of the human and bovine genes for the mitochondrial

2–oxoglutarate

[140569]carrier. DNA Seq. 3: 79–88, 1992.

[140570]

[140571]

[140572]13068.

[140573]Piccininni, S.; Iacobazzi, V.; Lauria, G.; Rocchi, M.;

Palmieri,

[140574]F.: Assignment of the oxoglutarate carrier gene (SLC20A4)
to human

[140575]chromosome 17p13.3. Cytogenet. Cell Genet. 83:
256–257, 1998.

[140576]

[140577]

[140578]13069.

[140579]Abbott, G. W.; Sesti, F.; Splawski, I.; Buck, M. E.; Lehmann,
M.

[140580]H.; Timothy, K. W.; Keating, M. T.; Goldstein, S. A. N.:
MiRP1 forms

[140581]I(Kr) potassium channels with HERG and is associated with
cardiac

[140582]arrhythmia. Cell 97: 175–187, 1999.

[140583]

[140584]

[140585]13070.

[140586]Schroeder, B. C.; Waldegger, S.; Fehr, S.; Bleich, M.; Warth, R.;

[140587]Greger, R.; Jentsch, T. J.: A constitutively open potassium channel

[140588]formed by KCNQ1 and KCNE3. Nature 403: 196–199, 2000.

[140589]

[140590]

[140591]13071.

[140592]Harvey, T. J.; Hooper, J. D.; Myers, S. A.; Stephenson, S.–A.;

[140593]Ashworth, L. K.; Clements, J. A.: Tissue–specific expression patterns

[140594]and fine mapping of the human kallikrein (KLK) locus on proximal 19q13.4. J.

[140595]Biol. Chem. 275: 37397–37406, 2000.

[140596]

[140597]

[140598]13072.

[140599]Chaudhry, F. A.; Reimer, R. J.; Krizaj, D.; Barber, D.;

Storm–Mathisen,

[140600]J.; Copenhagen, D. R.; Edwards, R. H.: Molecular analysis

of system

[140601]N suggests novel physiological roles in nitrogen

metabolism and synaptic

[140602]transmission. Cell 99: 769–780, 1999.

[140603]

[140604]

[140605]13073.

[140606]Yamada, K.; Nishida, K.; Hibi, M.; Hirano, T.; Matsuda, Y.:

Comparative

[140607]FISH mapping of Gab1 and Gab2 genes in human, mouse

and rat. Cytogenet.

[140608]Cell Genet. 94: 39–42, 2001.

[140609]

[140610]

[140611]13074.

- [140612]Afonina, E.; Stauber, R.; Pavlakis, G. N.: The human poly(A)–binding protein 1 shuttles between the nucleus and the cytoplasm. J. Biol. Chem. 273: 13015–13021, 1998.
- [140615]
- [140616]
- [140617]13075.
- [140618]Deo, R. C.; Bonanno, J. B.; Sonenberg, N.; Burley, S. K.: Recognition of polyadenylate RNA by the poly(A)–binding protein. Cell 98: 835–845, 1999.
- [140620]1999.
- [140621]
- [140622]
- [140623]13076.
- [140624]Gorlach, M.; Burd, C. G.; Dreyfuss, G.: The mRNA poly(A)–binding protein: localization, abundance, and RNA–binding specificity. Exp.

[140626]Cell Res. 211: 400–407, 1994.

[140627]

[140628]

[140629]13077.

[140630]Grange, T.; Martins de Sa, C.; Oddos, J.; Pictet, R.: Human mRNA

[140631]polyadenylate binding protein: evolutionary conservation of a nucleic

[140632]acid binding motif. Nucleic Acids Res. 15: 4771–4787, 1987.

[140633]

[140634]

[140635]13078.

[140636]Danpure, C. J.; Purdue, P. E.; Fryer, P.; Griffiths, S.; Allsop,

[140637]J.; Lumb, M. J.; Guttridge, K. M.; Jennings, P. R.; Scheinman, J.

[140638]I.; Mauer, S. M.; Davidson, N. O.: Enzymological and mutational analysis

[140639]of a complex primary hyperoxaluria type I phenotype involving alanine:glyoxylate

[140640]aminotransferase peroxisome-to-mitochondrion mistar-
geting and intraperoxisomal

[140641]aggregation. Am. J. Hum. Genet. 53: 417–432, 1993.

[140642]

[140643]

[140644]13079.

[140645]Mori, M.; Oda, T.; Nishiyama, K.; Serikawa, T.; Yamada, J.;
Ichiyama,

[140646]A.: A single serine:pyruvate aminotransferase gene on rat
chromosome

[140647]9q34–q36. Genomics 13: 686–689, 1992.

[140648]

[140649]

[140650]13080.

[140651]Purdue, P. E.; Allsop, J.; Isaya, G.; Rosenberg, L. E.; Dan-
pure,

[140652]C. J.: Mistargeting of peroxisomal L-alanine:glyoxylate
aminotransferase

[140653]to mitochondria in primary hyperoxaluria patients de-
pends upon activation

[140654]of a cryptic mitochondrial targeting sequence by a point
mutation. Proc.

[140655]Nat. Acad. Sci. 88: 10900–10904, 1991.

[140656]

[140657]

[140658]13081.

[140659]Purdue, P. E.; Lumb, M. J.; Allsop, J.; Minatogawa, Y.; Danpure,

[140660]C. J.: A glycine-to-glutamate substitution abolishes alanine:glyoxylate

[140661]aminotransferase catalytic activity in a subset of patients with primary

[140662]hyperoxaluria type 1. Genomics 13: 215–218, 1992.

[140663]

[140664]

[140665]13082.

[140666]Purdue, P. E.; Takada, Y.; Danpure, C. J.: Identification of

[140667]mutations associated with peroxisome-to-mitochondrion mistargeting

[140668]of alanine/glyoxylate aminotransferase in primary hyperoxaluria type

[140669]1. J. Cell Biol. 111: 2341–2351, 1990.

[140670]

[140671]

[140672]13083.

[140673]Takada, Y.; Kaneko, N.; Esumi, H.; Purdue, P. E.; Danpure,
C.

[140674]J.: Human peroxisomal L-alanine:glyoxylate aminotrans-
ferase: evolutionary

[140675]loss of a mitochondrial targeting signal by point mutation
of the

[140676]initiation codon. Biochem. J. 268: 517–520, 1990.

[140677]

[140678]

[140679]13084.

[140680]von Schnakenburg, C.; Rumsby, G.: Primary hyperoxaluria
type

[140681]1: a cluster of new mutations in exon 7 of the AGXT gene.
J. Med.

[140682]Genet. 34: 489–492, 1997.

[140683]

[140684]

[140685]13085.

[140686]Kobayashi, H.; Hino, M.; Shimodahira, M.; Iwakura, T.;

Ishihara,

[140687]T.; Ikekubo, K.; Ogawa, Y.; Nakao, K.; Kurahachi, H.: Mis-
sense mutation

[140688]of TRPS1 in a family of tricho-rhino-phalangeal syndrome
type III. Am.

[140689]J. Med. Genet. 107: 26-29, 2002.

[140690]

[140691]

[140692]13086.

[140693]Lipkin, S. M.; Moens, P. B.; Wang, V.; Lenzi, M.; Shanmu-

garajah,

[140694]D.; Gilgeous, A.; Thomas, J.; Cheng, J.; Touchman, J. W.;

Green, E.

[140695]D.; Schwartzberg, P.; Collins, F. S.; Cohen, P. E.: Meiotic
arrest

[140696]and aneuploidy in MLH3-deficient mice. Nature Genet. 31:
385-390,

[140697]2002.

[140698]

[140699]

[140700]13087.

[140701]Lipkin, S. M.; Wang, V.; Jacoby, R.; Banerjee-Basu, S.; Bax-
evanis,

[140702]A. D.; Lynch, H. T.; Elliott, R. M.; Collins, F. S.: MLH3: a
DNA

[140703]mismatch repair gene associated with mammalian mi-
crosatellite instability. Nature

[140704]Genet. 24: 27-35, 2000.

[140705]

[140706]

[140707]13088.

[140708]Lipkin, S. M.; Wang, V.; Stoler, D. L.; Anderson, G. R.;
Kirsch,

[140709]I.; Hadley, D.; Lynch, H. T.; Collins, F. S.: Germline and so-
matic

[140710]mutation analyses in the DNA mismatch repair gene
MLH3: evidence for

[140711]somatic mutation in colorectal cancers. Hum. Mutat. 17:
389-396,

[140712]2001.

[140713]

[140714]

[140715]13089.

[140716]Wu, Y.; Berends, M. J. W.; Sijmons, R. H.; Mensink, R. G. J.;
Verlind,

[140717]E.; Kooi, K. A.; van der Sluis, T.; Kempinga, C.; van der
Zee, A.

[140718]G. J.; Hollema, H.; Buys, C. H. C. M.; Kleibeuker, J. H.; Hof-
stra,

[140719]R. M. W.: A role for MLH3 in hereditary nonpolyposis col-
orectal cancer. Nature

[140720]Genet. 29: 137–138, 2001.

[140721]

[140722]

[140723]13090.

[140724]Sahara, S.; Aoto, M.; Eguchi, Y.; Imamoto, N.; Yoneda, Y.;
Tsujimoto,

[140725]Y.: Acinus is a caspase-3-activated protein required for
apoptotic

[140726]chromatin condensation. Nature 401: 168–173, 1999.

[140727]

[140728]

[140729]13091.

[140730]Hata, M.; Ohtsuka, K.: Characterization of HSE sequences
in human

[140731]Hsp40 gene: structural and promoter analysis. Biochim.
Biophys. Acta 1397:

[140732]43–55, 1998.

[140733]

[140734]

[140735]13092.

[140736]Hata, M.; Okumura, K.; Seto, M.; Ohtsuka, K.: Genomic
cloning

[140737]of a human heat shock protein 40 (Hsp40) gene (HSPF1)
and its chromosomal

[140738]localization to 19p13.2. Genomics 38: 446–449, 1996.

[140739]

[140740]

[140741]13093.

[140742]Ohtsuka, K.: Cloning of a cDNA for heat-shock protein
hsp40, a

[140743]human homologue of bacterial DnaJ. Biochem. Biophys.
Res. Commun. 197:

[140744]235–240, 1993.

[140745]

[140746]

[140747]13094.

[140748]Adams, M. D.; Dubnick, M.; Kerlavage, A. R.; Moreno, R.;
Kelley,

[140749]J. M.; Utterback, T. R.; Nagle, J. W.; Fields, C.; Venter, J. C.:

[140750]Sequence identification of 2,375 human brain genes. Na-
ture 355:

[140751]632–634, 1992. Note: Comment: Nature 357: 367–368,
1992.

[140752]

[140753]

[140754]13095.

[140755]Muravenko, O. V.; Gizatullin, R. Z.; Protopopov, A. I.;
Kashuba,

[140756]V. I.; Zabarovsky, E. R.; Zelenin, A. V.: Assignment of
CDK5R2 coding

[140757]for the cyclin-dependent kinase 5, regulatory subunit 2
(NCK5AI protein)

[140758]to human chromosome band 2q35 by fluorescent in situ
hybridization. Cytogenet.

[140759]Cell Genet. 89: 160–161, 2000.

[140760]

[140761]

[140762]13096.

[140763]Nilden, F.; Backstrom, A.; Bark, C.: Molecular cloning and
characterisation

[140764]of a mouse gene encoding an isoform of the neuronal cy-
clin-dependent

[140765]kinase 5 (CDK5) activator. Biochim. Biophys. Acta 1398:
371–376,

[140766]1998.

[140767]

[140768]

[140769]13097.

[140770]Tang, D.; Yeung, J.; Lee, K-Y.; Matsushita, M.; Matsui, H.;
Tomizawa,

[140771]K.; Hatase, O.; Wang, J. H.: An isoform of the neuronal cy-
clin-dependent

[140772]kinase 5 (Cdk5) activator. J. Biol. Chem. 270:
26897-26903, 1995.

[140773]

[140774]

[140775]13098.

[140776]DuPont, B. R.; Hu, C.-C.; Reveles, X.; Simmer, J. P.: As-
signment

[140777]of serine protease 17 (PRSS17) to human chromosome
bands 19q13.3-q13.4

[140778]by in situ hybridization. Cytogenet. Cell Genet. 86:
212-213, 1999.

[140779]

[140780]

[140781]13099.

[140782]Nelson, P. S.; Gan, L.; Ferguson, C.; Moss, P.; Gelinas, R.;
Hood,

[140783]L.; Wang, K.: Molecular cloning and characterization of

prostase,

[140784]an androgen-regulated serine protease with prostate-restricted expression. Proc.

[140785]Nat. Acad. Sci. 96: 3114–3119, 1999.

[140786]

[140787]

[140788]13100.

[140789]Hallas, C.; Pekarsky, Y.; Itoyama, T.; Varnum, J.; Bichi, R.; Rothstein,

[140790]J. L.; Croce, C. M.: Genomic analysis of human and mouse TCL1 loci

[140791]reveals a complex of tightly clustered genes. Proc. Nat. Acad. Sci. 96:

[140792]14418–14423, 1999.

[140793]

[140794]

[140795]13101.

[140796]Pekarsky, Y.; Hallas, C.; Isobe, M.; Russo, G.; Croce, C. M.:

[140797]Abnormalities at 14q32.1 in T cell malignancies involve two oncogenes. Proc.

[140798]Nat. Acad. Sci. 96: 2949–2951, 1999.

[140799]

[140800]

[140801]13102.

[140802]Koga, Y.; Fabrizi, G. M.; Mita, S.; Arnaudo, E.; Lomax, M. I.;

[140803]Aqua, M. S.; Grossman, L. I.; Schon, E. A.: Sequence of a
cDNA specifying

[140804]subunit VIIc of human cytochrome c oxidase. Nucleic
Acids Res. 18:

[140805]684 only, 1990.

[140806]

[140807]

[140808]13103.

[140809]Chaudhary, P. M.; Ferguson, C.; Nguyen, V.; Nguyen, O.;
Massa,

[140810]H. F.; Eby, M.; Jasmin, A.; Trask, B. J.; Hood, L.; Nelson, P.
S.

[140811]: Cloning and characterization of two Toll/interleukin–1
receptor–like

[140812]genes TIL3 and TIL4: evidence for a multi–gene receptor

family in

[140813]humans. Blood 91: 4020–4027, 1998.

[140814]

[140815]

[140816]13104.

[140817]Harju, K.; Glumoff, V.; Hallman, M.: Ontogeny of Toll-like
receptors

[140818]Tlr2 and Tlr4 in mice. Pediat. Res. 49: 81–83, 2001.

[140819]

[140820]

[140821]13105.

[140822]Qu, L. H.; Nicoloso, M.; Michot, B.; Azum, M. C.; Caizer-
gues-Ferrer,

[140823]M.; Renalier, M. H.; Bachellerie, J. P.: U21, a novel small
nucleolar

[140824]RNA with a 13 nt. complementarity to 28S rRNA, is en-
coded in an intron

[140825]of ribosomal protein L5 gene in chicken and mammals.

Nucleic Acids

[140826]Res. 22: 4073–4081, 1994.

[140827]

[140828]

[140829]13106.

[140830]Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.;
Grandy, D.;

[140831]Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an
11q22.3-9p22

[140832]translocation with affective disorder: proximity of the
dopamine D2

[140833]receptor gene relative to the translocation breakpoint.
(Abstract) Am.

[140834]J. Hum. Genet. 45 (suppl.): A220 only, 1989.

[140835]

[140836]

[140837]13107.

[140838]Abuladze, N.; Lee, I.; Newman, D.; Hwang, J.; Boorer, K.;
Pushkin,

[140839]A.; Kurtz, I.: Molecular cloning, chromosomal localization,
tissue

[140840]distribution, and functional expression of the human pan-
creatic sodium

[140841]bicarbonate cotransporter. J. Biol. Chem. 273:
17689–17695, 1998.

[140842]

[140843]

[140844]13108.

[140845]Burnham, C. E.; Amlal, H.; Wang, Z.; Shull, G. E.; Soleimani,
M.

[140846]: Cloning and functional expression of a human kidney
Na⁺:HCO₃⁻ cotransporter. J.

[140847]Biol. Chem. 272: 19111–19114, 1997.

[140848]

[140849]

[140850]13109.

[140851]Choi, I.; Romero, M. F.; Khandoudi, N.; Bril, A.; Boron, W.
F.

[140852]: Cloning and characterization of a human electrogenic
Na⁽⁺⁾–HCO₃⁽⁻⁾

[140853]cotransporter isoform (hhNBC). Am. J. Physiol. 276:
C576–C584, 1999.

[140854]

[140855]

[140856]13110.

[140857]Igarashi, T.; Inatomi, J.; Sekine, T.; Cha, S. H.; Kanai, Y.;

Kunimi,

[140858]M.; Tsukamoto, K.; Satoh, H.; Shimadzu, M.; Tozawa, F.;

Mori, T.;

[140859]Shiobara, M.; Seki, G.; Endou, H.: Mutations in SLC4A4

cause permanent

[140860]isolated proximal renal tubular acidosis with ocular ab-

normalities.

[140861](Letter) Nature Genet. 23: 264–265, 1999.

[140862]

[140863]

[140864]13111.

[140865]Romero, M. F.; Boron, W. F.: Electrogenic Na(+)/HCO(3-)

cotransporters:

[140866]cloning and physiology. Annu. Rev. Physiol. 61: 699–723,

1999.

[140867]

[140868]

[140869]13112.

[140870]Soleimani, M.; Burnham, C. E.: Physiologic and molecular aspects

[140871]of the Na(+):HCO(3-) cotransporter in health and disease processes. Kidney

[140872]Int. 57: 371–384, 2000.

[140873]

[140874]

[140875]13113.

[140876]Usui, T.; et al.; et al. :Pflugers Arch. 438: 458–462, 1999.

[140877]

[140878]

[140879]13114.

[140880]Zhou, Y.–D.; Barnard, M.; Tian, H.; Li, X.; Ring, H. Z.;
Francke,

[140881]U.; Shelton, J.; Richardson, J.; Russell, D. W.; McKnight, S.
L.:

[140882]Molecular characterization of two mammalian bHLH–PAS
domain proteins

[140883]selectively expressed in the central nervous system. Proc.

Nat. Acad.

[140884]Sci. 94: 713–718, 1997.

[140885]

[140886]

[140887]13115.

[140888]Garcia, J. A.; Zhang, D.; Estill, S. J.; Michnoff, C.; Rutter,

[140889]J.; Reick, M.; Scott, K.; Diaz–Arrastia, R.; McKnight, S. L.:

Impaired

[140890]cued and contextual memory in NPAS2–deficient mice.

Science 288:

[140891]2226–2230, 2000.

[140892]

[140893]

[140894]13116.

[140895]Spencer, E.; Jiang, J.; Chen, Z. J.: Signal–induced ubiquiti–

nation

[140896]of I–kappa–B–alpha by the F–box protein Slimb/

beta–TrCP. Genes Dev. 13:

[140897]284–294, 1999.

[140898]

[140899]

[140900]13117.

[140901]Yaron, A.; Hatzubai, A.; Davis, M.; Lavon, I.; Amit, S.; Manning,

[140902]A. M.; Andersen, J. S.; Mann, M.; Mercurio, F.; Ben-Neriah, Y.: Identification

[140903]of the receptor component of the I-kappa-B-alpha-ubiquitin ligase. Nature 396:

[140904]590-594, 1998.

[140905]

[140906]

[140907]13118.

[140908]Di Fruscio, M.; Gilchrist, C. A.; Baker, R. T.; Gray, D. A.: Genomic

[140909]structure of Unp, a murine gene encoding a ubiquitin-specific protease. Biochim.

[140910]Biophys. Acta 1398: 9-17, 1998.

[140911]

[140912]

[140913]13119.

[140914]Frederick, A.; Rolfe, M.; Chiu, M. I.: The human UNP locus
at

[140915]3p21.31 encodes two tissue-selective, cytoplasmic iso-
forms with deubiquitinating

[140916]activity that have reduced expression in small cell lung
carcinoma

[140917]cell lines. Oncogene 16: 153–165, 1998.

[140918]

[140919]

[140920]13120.

[140921]Gray, D. A.; Inazawa, J.; Gupta, K.; Wong, A.; Ueda, R.;
Takahashi,

[140922]T.: Elevated expression of Unph, a proto-oncogene at
3p21.3, in human

[140923]lung tumors. Oncogene 10: 2179–2183, 1995.

[140924]

[140925]

[140926]13121.

[140927]Gupta, K.; Chevrette, M.; Gray, D. A.: The Unp proto-

oncogene

[140928]encodes a nuclear protein. Oncogene 9: 1729–1731,
1994.

[140929]

[140930]

[140931]13122.

[140932]Gupta, K.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.;
Gray,

[140933]D. A.: Unp, a mouse gene related to the tre oncogene.

Oncogene 8:

[140934]2307–2310, 1993.

[140935]

[140936]

[140937]13123.

[140938]Lucas, C. A.; Rughani, A.; Hoh, J. F. Y.: Expression of ex-
traocular

[140939]myosin heavy chain in rabbit laryngeal muscle. J. Muscle
Res. Cell

[140940]Motil. 16: 368–378, 1995.

[140941]

[140942]

[140943]13124.

[140944]Wieczorek, D. F.; Periasamy, M.; Butler-Browne, G. S.;
Whalen,

[140945]R. G.; Nadal-Ginard, B.: Co-expression of multiple myosin
heavy chain

[140946]genes, in addition to a tissue-specific one, in extraocular
musculature. J.

[140947]Cell Biol. 101: 618–629, 1985.

[140948]

[140949]

[140950]13125.

[140951]Winters, L. M.; Briggs, M. M.; Schachat, F.: The human ex-
traocular

[140952]muscle myosin heavy chain gene (MYH13) maps to the
cluster of fast

[140953]and developmental myosin genes on chromosome 17. Ge-
nomics 54: 188–189,

[140954]1998.

[140955]

[140956]

[140957]13126.

[140958]Briskin, C.; Heineman, A.; Chavarria, T.; Elenbaas, B.; Tan, J.;

[140959]Dey, S. K.; McMahon, J. A.; McMahon, A. P.; Weinberg, R. A.: Essential

[140960]function of Wnt-4 in mammary gland development downstream of progesterone

[140961]signaling. Genes Dev. 14: 650-654, 2000.

[140962]

[140963]

[140964]13127.

[140965]Jordan, B. K.; Mohammed, M.; Ching, S. T.; Delot, E.; Chen, X.-N.;

[140966]Dewing, P.; Swain, A.; Rao, P. N.; Elejalde, B. R.; Vilain, E.: Up-regulation

[140967]of WNT-4 signaling and dosage-sensitive sex reversal in humans. Am.

[140968]J. Hum. Genet. 68: 1102-1109, 2001.

[140969]

[140970]

[140971]13128.

[140972]Stark, K.; Vainio, S.; Vassileva, G.; McMahon, A. P.: Epithelial

[140973]transformation of metanephric mesenchyme in the developing kidney

[140974]regulated by Wnt-4. Nature 372: 679-683, 1994.

[140975]

[140976]

[140977]13129.

[140978]Vainio, S.; Heikkila, M.; Kispert, A.; Chin, N.; McMahon, A. P.

[140979]: Female development in mammals is regulated by Wnt-4 signalling. Nature 397:

[140980]405-409, 1999.

[140981]

[140982]

[140983]13130.

[140984]Vilain, A.; Apiou, F.; Dutrillaux, B.; Malfoy, B.: Assignment

[140985]of candidate DNA methyltransferase gene (DNMT2) to human chromosome

[140986]band 10p15.1 by in situ hybridization. Cytogenet. Cell Genet. 82:

[140987]120 only, 1998.

[140988]

[140989]

[140990]13131.

[140991]Yoder, J. A.; Bestor, T. H.: A candidate mammalian DNA methyltransferase

[140992]related to pmt1p of fission yeast. Hum. Molec. Genet. 7: 279-284,

[140993]1998.

[140994]

[140995]

[140996]13132.

[140997]Avraham, K. B.; Cho, B. C.; Gilbert, D.; Fujii, H.; Okamoto, K.;

[140998]Shimazaki, T.; Ito, T.; Shoji, H.; Wakamatsu, Y.; Kondoh, H.; Takahashi,

[140999]N.; Muramatsu, M.; Hamada, H.; Copeland, N. G.; Jenkins,
N. A.: Murine

[141000]chromosomal location of four class III POU transcription
factors. Genomics 18:

[141001]131–133, 1993.

[141002]

[141003]

[141004]13133.

[141005]Egan, S.; Herbrick, J.–A.; Tsui, L.–C.; Cohen, B.; Flock, G.;
Beatty,

[141006]B.; Scherer, S. W.: Mapping of the human lunatic fringe
(LFNG) gene

[141007]to 7p22 and manic fringe (MFNG) to 22q12. Genomics 54:
576–577,

[141008]1998.

[141009]

[141010]

[141011]13134.

[141012]Johnston, S. H.; Rauskolb, C.; Wilson, R.; Prabhakaran, B.;
Irvine,

[141013]K. D.; Vogt, T. F.: A family of mammalian Fringe genes implicated

[141014]in boundary determination and the Notch pathway. Development 124:

[141015]2245–2254, 1997.

[141016]

[141017]

[141018]13135.

[141019]Moran, J. L.; Johnston, S. H.; Rauskolb, C.; Bhalerao, J.; Bowcock,

[141020]A. M.; Vogt, T. F.: Genomic structure, mapping, and expression analysis

[141021]of the mammalian lunatic, manic, and radical fringe genes. Mammalian

[141022]Genome 10: 535–541, 1999.

[141023]

[141024]

[141025]13136.

[141026]Van Tine, B. A.; Knops, J.; Shaw, G. M.; May, W. A.: Assignment

[141027]of human MFNG, manic fringe Drosophila homolog, to
22q13.1 using tyramide

[141028]fluorescence in situ hybridization (T-FISH). Cytogenet. Cell
Genet. 87:

[141029]132–133, 1999.

[141030]

[141031]

[141032]13137.

[141033]Laufer, E.; Dahn, R.; Orozco, O. E.; Yeo, C.–Y.; Piseni, J.;
Henrique,

[141034]D.; Abbott, U. K.; Fallon, J. F.; Tabin, C.: Expression of
Radical

[141035]fringe in limb–bud ectoderm regulates apical ectodermal
ridge formation. Nature 386:

[141036]366–373, 1997.

[141037]

[141038]

[141039]13138.

[141040]Rodriguez–Esteban, C.; Schwabe, J. W. R.; De La Pena, J.;
Foy,

[141041]B.; Eshelman, B.; Izpisua Belmonte, J. C.: Radical fringe positions

[141042]the apical ectodermal ridge at the dorsoventral boundary of the vertebrate

[141043]limb. Nature 386: 360–366, 1997.

[141044]

[141045]

[141046]13139.

[141047]Hofmann, Y.; Becker, J.; Wright, F.; Avner, E. D.; Mrug, M.; Guay–Woodford,

[141048]L. M.; Somlo, S.; Zerres, K.; Germino, G. G.; Onuchic, L. F.: Genomic

[141049]structure of the gene for the human P1 protein (MCM3) and its exclusion

[141050]as a candidate for autosomal recessive polycystic kidney disease. Europ.

[141051]J. Hum. Genet. 8: 163–166, 2000.

[141052]

[141053]

[141054]13140.

[141055]Starborg, M.; Brundell, E.; Gell, K.; Larsson, C.; White, I.;
Daneholt,

[141056]B.; Hoog, C.: A murine replication protein accumulates
temporarily

[141057]in the heterochromatic regions of nuclei prior to initiation
of DNA

[141058]replication. J. Cell Sci. 108: 927–934, 1995.

[141059]

[141060]

[141061]13141.

[141062]Chicheportiche, Y.; Bourdon, P. R.; Xu, H.; Hsu, Y.–M.;
Scott,

[141063]H.; Hession, C.; Garcia, I.; Browning, J. L.: TWEAK, a new
secreted

[141064]ligand in the tumor necrosis factor family that weakly in-
duces apoptosis. J.

[141065]Biol. Chem. 272: 32401–32410, 1997.

[141066]

[141067]

[141068]13142.

[141069]Kaptein, A.; Jansen, M.; Dilaver, G.; Kitson, J.; Dash, L.;
Wang,

[141070]E.; Owen, M. J.; Bodmer, J.-L.; Tschopp, J.; Farrow, S. N.:
Studies

[141071]on the interaction between TWEAK and the death receptor
WSL-1/TRAMP

[141072](DR3). FEBS Lett. 485: 135-141, 2000.

[141073]

[141074]

[141075]13143.

[141076]Marsters, S. A.; Sheridan, J. P.; Pitti, R. M.; Brush, J.; God-
dard,

[141077]A.; Ashkenazi, A.: Identification of a ligand for the death-
domain-containing

[141078]receptor Apo3. Curr. Biol. 8: 525-528, 1998.

[141079]

[141080]

[141081]13144.

[141082]Schneider, P.; Schwenzer, R.; Haas, E.; Muhlenbeck, F.;
Schubert,

[141083]G.; Scheurich, P.; Tschopp, J.; Wajant, H.: TWEAK can induce cell

[141084]death via endogenous TNF and TNF receptor 1. *Europ. J. Immun.* 29:

[141085]1785–1792, 1999.

[141086]

[141087]

[141088]13145.

[141089]Wiley, S. R.; Cassiano, L.; Lofton, T.; Davis–Smith, T.; Winkles,

[141090]J. A.; Lindner, V.; Liu, H.; Daniel, T. O.; Smith, C. A.; Fanslow,

[141091]W. C.: A novel TNF receptor family member binds TWEAK and is implicated

[141092]in angiogenesis. *Immunity* 15: 837–846, 2001.

[141093]

[141094]

[141095]13146.

[141096]Hoey, T.; Sun, Y.–L.; Williamson, K.; Xu, X.: Isolation of two

[141097]new members of the NF–AT gene family and functional

characterization

[141098]of the NF-AT proteins. Immunity 2: 461–472, 1995.

[141099]

[141100]

[141101]13147.

[141102]Hoey, T.; Sun, Y. L.; Williamson, K.; Xu, X.: Isolation of two

[141103]new members of the NF-AT gene family and functional
characterization

[141104]of the NF-AT proteins. Immunity 2: 461–472, 1995.

[141105]

[141106]

[141107]13148.

[141108]Arany, Z.; Newsome, D.; Oldread, E.; Livingston, D. M.;
Eckner,

[141109]R.: A family of transcriptional adaptor proteins targeted by
the

[141110]E1A oncoprotein. Nature 374: 81–84, 1995.

[141111]

[141112]

[141113]13149.

[141114]de Lecea, L.; Ruiz-Lozano, P.; Danielson, P. E.; Peelle-Kirley,

[141115]J.; Foye, P. E.; Frankel, W. N.; Sutcliffe, J. G.: Cloning, mRNA

[141116]expression, and chromosomal mapping of mouse and human preprocortistatin. Genomics 42:

[141117]499–506, 1997.

[141118]

[141119]

[141120]13150.

[141121]Ejeskar, K.; Abel, F.; Sjoberg, R.-M.; Backstrom, J.; Kogner, P.;

[141122]Martinsson, T.: Fine mapping of the human preprocortistatin gene

[141123](CORT) to neuroblastoma consensus deletion region 1p36.3–p36.2, but

[141124]absence of mutations in primary tumors. Cytogenet. Cell Genet. 89:

[141125]62–66, 2000.

[141126]

[141127]

[141128]13151.

[141129]Fukusumi, S.; Kitada, C.; Takekawa, S.; Kizawa, H.;
Sakamoto, J.;

[141130]Miyamoto, M.; Hinuma, S.; Kitano, K.; Fujino, M.: Identifi-
cation

[141131]and characterization of a novel human cortistatin-like
peptide. Biochem.

[141132]Biophys. Res. Commun. 232: 157–163, 1997.

[141133]

[141134]

[141135]13152.

[141136]Zhong, R.; Roeder, R. G.; Heintz, N.: The primary structure
and

[141137]expression of four cloned human histone genes. Nucleic
Acids Res. 11:

[141138]7409–7425, 1983.

[141139]

[141140]

[141141]13153.

[141142]Dong, C.; Zhu, S.; Wang, T.; Yoon, W.; Li, Z.; Alvarez, R. J.;

[141143]ten Dijke, P.; White, B.; Wigley, F. M.; Goldschmidt-Clermont, P.

[141144]J.: Deficient Smad7 expression: a putative molecular defect in scleroderma. Proc.

[141145]Nat. Acad. Sci. 99: 3908–3913, 2002.

[141146]

[141147]

[141148]13154.

[141149]Kavsak, P.; Rasmussen, R. K.; Causing, C. G.; Bonni, S.; Zhu, H.;

[141150]Thomsen, G. H.; Wrana, J. L.: Smad7 binds to Smurf2 to form an E3

[141151]ubiquitin ligase that targets the TGF-beta receptor for degradation. Molec.

[141152]Cell 6: 1365–1375, 2000.

[141153]

[141154]

[141155]13155.

[141156]Lallemant, F.; Mazars, A.; Prunier, C.; Bertrand, F.; Korn-

prost,

[141157]M.; Gallea, S.; Roman–Roman, S.; Cherqui, G.; Atfi, A.:

Smad7 inhibits

[141158]the survival nuclear factor kappa–B and potentiates apop–

tosis in epithelial

[141159]cells. *Oncogene* 20: 879–884, 2001.

[141160]

[141161]

[141162]13156.

[141163]Roijs, E.; Moren, A.; ten Dijke, P.; Stenman, G.: Assign–

ment

[141164]of the Smad7 gene (MADH7) to human chromosome

18q21.1 by fluorescence

[141165]in situ hybridization. *Cytogenet. Cell Genet.* 81: 189–190,

1998.

[141166]

[141167]

[141168]13157.

[141169]Cravatt, B. F.; Giang, D. K.; Mayfield, S. P.; Boger, D. L.;

Lerner,

[141170]R. A.; Gilula, N. B.: Molecular characterization of an en–

zyme that

[141171]degrades neuromodulatory fatty-acid amides. Nature 384:
83-87, 1996.

[141172]

[141173]

[141174]13158.

[141175]Giang, D. K.; Cravatt, B. F.: Molecular characterization of
human

[141176]and mouse fatty acid amide hydrolases. Proc. Nat. Acad.
Sci. 94:

[141177]2238-2242, 1997.

[141178]

[141179]

[141180]13159.

[141181]Sipe, J. C.; Chiang, K.; Gerber, A. L.; Beutler, E.; Cravatt, B.

[141182]F.: A missense mutation in human fatty acid amide hydro-
lase associated

[141183]with problem drug use. Proc. Nat. Acad. Sci. 99:
8394-8399, 2002.

[141184]

[141185]

[141186]13160.

[141187]Wan, M.; Cravatt, B. F.; Ring, H. Z.; Zhang, X.; Francke, U.:

[141188]Conserved chromosomal location and genomic structure
of human and

[141189]mouse fatty-acid amide hydrolase genes and evaluation of
clasper as

[141190]a candidate neurological mutation. Genomics 54:
408-414, 1998.

[141191]

[141192]

[141193]13161.

[141194]Falany, C. N.; Johnson, M. R.; Barnes, S.; Diasio, R. B.:
Glycine

[141195]and taurine conjugation of bile acids by a single enzyme:
molecular

[141196]cloning and expression of a human liver bile acid
CoA:amino acid N-acyltransferase. J.

[141197]Biol. Chem. 269: 19375-19379, 1994.

[141198]

[141199]

[141200]13162.

[141201]Johnson, M. R.; Barnes, S.; Kwakye, J. B.; Diasio, R. B.: Pu-
rification

[141202]and characterization of bile acid-CoA:amino acid N-
acyltransferase

[141203]from human liver. J. Biol. Chem. 266: 10227-10233,
1991.

[141204]

[141205]

[141206]13163.

[141207]Baier, M.; Bannert, N.; Werner, A.; Lang, K.; Kurth, R.:
Molecular

[141208]cloning, sequence, expression, and processing of the in-
terleukin 16

[141209]precursor. Proc. Nat. Acad. Sci. 94: 5273-5277, 1997.

[141210]

[141211]

[141212]13164.

[141213]Roh, M. H.; Makarova, O.; Liu, C.-J.; Shin, K.; Lee, S.; Lau-

rinec,

[141214]S.; Goyal, M.; Wiggins, R.; Margolis, B.: The Maguk protein, Pals1,

[141215]functions as an adapter, linking mammalian homologues of Crumbs and

[141216]Discs Lost. J. Cell Biol. 157: 161–172, 2002.

[141217]

[141218]

[141219]13165.

[141220]Bannert, N.; Baier, M.; Werner, A.; Kurth, R.: Interleukin–16

[141221]or not? Nature 381: 30 only, 1996.

[141222]

[141223]

[141224]13166.

[141225]Bannert, N.; Kurth, R.; Baier, M.: The gene encoding mouse interleukin–16

[141226]consists of seven exons and maps to Chromosome 7 D2–D3. Immunogenetics 49:

[141227]704–706, 1999.

[141228]

[141229]

[141230]13167.

[141231]Cruikshank, W.; Center, D. M.: Modulation of lymphocyte
migration

[141232]by human lymphokines. II. Purification of a lymphotactic
factor (LCF). J.

[141233]Immun. 128: 2569–2574, 1982.

[141234]

[141235]

[141236]13168.

[141237]Cruikshank, W. W.; Center, D. M.; Nisar, N.; Wu, M.; Natke,
B.;

[141238]Theodore, A. C.; Kornfeld, H.: Molecular and functional
analysis

[141239]of a lymphocyte chemoattractant factor: association of bi-
ologic function

[141240]with CD4 expression. Proc. Nat. Acad. Sci. 91:
5109–5113, 1994.

[141241]

[141242]

[141243]13169.

[141244]Drwinga, H. L.; Toji, L. H.; Kim, C. H.; Greene, A. E.;

Mulivor,

[141245]R. A.: NIGMS human/rodent somatic cell hybrid mapping

panels 1 and

[141246]2. Genomics 16: 311–314, 1993.

[141247]

[141248]

[141249]13170.

[141250]Hudson, T. J.; Stein, L. D.; Gerety, S. S.; Ma, J.; Castle, A.

[141251]B.; Silva, J.; Slonim, D. K.; Baptista, R.; Kruglyak, L.; Xu, S.

H.;

[141252]Hu, X.; Colbert, A. M. E.; and 39 others: An STS-based

map of the

[141253]human genome. Science 270: 1945–1954, 1995.

[141254]

[141255]

[141256]13171.

[141257]Keane, J.; Nicoll, J.; Kim, S.; Wu, D. M. H.; Cruikshank, W.
W.;

[141258]Brazer, W.; Natke, B.; Zhang, Y.; Center, D. M.; Kornfeld,
H.: Conservation

[141259]of structure and function between human and murine IL-
16. J. Immun. 160:

[141260]5945–5954, 1998.

[141261]

[141262]

[141263]13172.

[141264]Tarnasky, H.; Gill, D.; Murthy, S.; Shao, X.; Demetrick, D.
J.;

[141265]van der Hoorn, F. A.: A novel testis-specific gene, SPAG4,
whose

[141266]product interacts specifically with outer dense fiber pro-
tein ODF27,

[141267]maps to human chromosome 20q11.2. Cytogenet. Cell
Genet. 81: 65–67,

[141268]1998.

[141269]

[141270]

[141271]13173.

[141272]Scheuerpflug, C. G.; Lichter, P.; Debatin, K.-M.; Mincheva, A.

[141273]: Assignment of TRADD to human chromosome band 16q22 by in situ hybridization. Cytogenet.

[141274]Cell Genet. 92: 347-348, 2001.

[141275]

[141276]

[141277]13174.

[141278]Lennon-Dumenil, A.-M.; Barbouche, M.-R.; Vedrenne, J.; Prod'Homme,

[141279]T.; Bejaoui, M.; Ghariani, S.; Charron, D.; Fellous, M.; Del-lagi,

[141280]K.; Alcaide-Loridan, C.: Uncoordinated HLA-D gene expression in a

[141281]RFXANK-defective patient with MHC class II deficiency. J. Immun. 166:

[141282]5681-5687, 2001.

[141283]

[141284]

[141285]13175.

[141286]Lisowska-Groszpiere, B.; Fondaneche, M. C.; Rols, M. P.;
Griscelli,

[141287]C.; Fischer, A.: Two complementation groups account for
most cases

[141288]of inherited MHC class II deficiency. Hum. Molec. Genet. 3:
953–958,

[141289]1994.

[141290]

[141291]

[141292]13176.

[141293]Boll, W.; Wagner, P.; Mantei, N.: Structure of the chromo-
somal

[141294]gene and cDNAs coding for lactase-phlorizin hydrolase in
humans with

[141295]adult-type hypolactasia or persistence of lactase. Am. J.
Hum. Genet. 48:

[141296]889–902, 1991.

[141297]

[141298]

[141299]13177.

[141300]Wiszniewski, W.; Fondaneche, M.-C.; Lambert, N.; Master-
nak, K.;

[141301]Picard, C.; Notarangelo, L.; Schwartz, K.; Bal, J.; Reith, W.;
Alcaide,

[141302]C.; de Saint Basile, G.; Fischer, A.; Lisowska-GrosPierre, B.:
Founder

[141303]effect for a 26-bp deletion in the RFXANK gene in North
African major

[141304]histocompatibility complex class II-deficient patients be-
longing to

[141305]complementation group B. Immunogenetics 51: 261-267,
2000.

[141306]

[141307]

[141308]13178.

[141309]Harvey, C. B.; Fox, M. F.; Jeggo, P. A.; Mantei, N.; Povey, S.;

[141310]Swallow, D. M.: Regional localization of the lactase-
phlorizin hydrolase

[141311]gene, LCT, to chromosome 2q21. Ann. Hum. Genet. 57:
179-185, 1993.

[141312]

[141313]

[141314]13179.

[141315]Kruse, T. A.; Bolund, L.; Byskov, A.; Sjostrom, H.; Noren, O.;

[141316]Mantei, N.; Semenza, G.: Mapping of the human lactase-phlorizin hydrolase

[141317]gene to chromosome 2. (Abstract) Cytogenet. Cell Genet. 51: 1026,

[141318]1989.

[141319]

[141320]

[141321]13180.

[141322]Kruse, T. A.; Bolund, L.; Grzeschik, K.-H.; Ropers, H. H.; Sjostrom,

[141323]H.; Noren, O.; Mantei, N.; Semenza, G.: The human lactase-phlorizin

[141324]hydrolase gene is located on chromosome 2. FEBS Lett. 240: 123-126,

[141325]1988.

[141326]

[141327]

[141328]13181.

[141329]Mantei, N.; Villa, M.; Enzler, T.; Wacker, H.; Boll, W.; James,

[141330]O.; Hunziker, W.; Semenza, G.: Complete primary structure of human

[141331]and rabbit lactase-phlorizin hydrolase: implications for biosynthesis,

[141332]membrane anchoring and evolution of the enzyme. EMBO J. 7: 2705-2713,

[141333]1988.

[141334]

[141335]

[141336]13182.

[141337]Stec, I.; Wright, T. J.; van Ommen, G.-J. B.; de Boer, P. A. J.;

[141338]van Haeringen, A.; Moorman, A. F. M.; Altherr, M. R.; den Dunnen,

[141339]J. T.: WHSC1, a 90 kb SET domain-containing gene, expressed in early

[141340]development and homologous to a Drosophila dysmorphy

gene maps in

[141341]the Wolf-Hirschhorn syndrome critical region and is fused to IgH in

[141342]t(4;14) multiple myeloma. Hum. Molec. Genet. 7: 1071-1082, 1998.

[141343]

[141344]

[141345]13183.

[141346]Ame, J.-C.; Apiou, F.; Jacobson, E. L.; Jacobson, M. K.: Assignment

[141347]of the poly(ADP-ribose) glycohydrolase gene (PARG) to human chromosome

[141348]10q11.23 and mouse chromosome 14B by in situ hybridization. Cytogenet.

[141349]Cell Genet. 85: 269-270, 1999.

[141350]

[141351]

[141352]13184.

[141353]Lin, W.; Ame, J.-C.; Aboul-Ela, N.; Jacobson, E. L.; Jacobson,

[141354]M. K.: Isolation and characterization of the cDNA encoding

bovine

[141355]poly(ADP-ribose) glycohydrolase. J. Biol. Chem. 272:

11895-11901,

[141356]1997.

[141357]

[141358]

[141359]13185.

[141360]Brown, S. D.; Twells, R. C. J.; Hey, P. J.; Cox, R. D.; Levy, E.

[141361]R.; Soderman, A. R.; Metzker, M. L.; Caskey, C. T.; Todd, J.

A.; Hess,

[141362]J. F.: Isolation and characterization of LRP6, a novel mem-

ber of

[141363]the low density lipoprotein receptor gene family. Biochem.

Biophys.

[141364]Res. Commun. 248: 879-888, 1998.

[141365]

[141366]

[141367]13186.

[141368]Mao, B.; Wu, W.; Li, Y.; Hoppe, D.; Stannek, P.; Glinka, A.;

Niehrs,

[141369]C.: LDL-receptor-related protein 6 is a receptor for Dickkopf proteins. Nature 411: 321–325, 2001.

[141371]

[141372]

[141373]13187.

[141374]Pinson, K. I.; Brennan, J.; Monkley, S.; Avery, B. J.; Skarnes,

[141375]W. C.: An LDL-receptor-related protein mediates Wnt signalling in

[141376]mice. Nature 407: 535–538, 2000.

[141377]

[141378]

[141379]13188.

[141380]Tamai, K.; Semenov, M.; Kato, Y.; Spokony, R.; Liu, C.; Katsuyama,

[141381]Y.; Hess, F.; Saint-Jeannet, J.-P.; He, X.: LDL-receptor-related

[141382]proteins in Wnt signal transduction. Nature 407: 530–535, 2000.

[141383]

[141384]

[141385]13189.

[141386]Wehrli, M.; Dougan, S. T.; Caldwell, K.; O'Keefe, L.;
Schwartz,

[141387]S.; Vaizel-Ohayon, D.; Schejter, E.; Tomlinson, A.; Di-
Nardo, S.:

[141388]Arrow encodes an LDL-receptor-related protein essential
for Wingless

[141389]signalling. Nature 407: 527-530, 2000. Note: Erratum:
Nature 410:

[141390]847 only, 2001.

[141391]

[141392]

[141393]13190.

[141394]Vinkemeier, U.; Obermann, W.; Weber, K.; Furst, D. O.: The
globular

[141395]head domain of titin extends into the center of the sar-
comeric M band:

[141396]cDNA cloning, epitope mapping and immunoelectron mi-
croscopy of two

[141397]titin-associated proteins. J. Cell Sci. 106: 319-330, 1993.

[141398]

[141399]

[141400]13191.

[141401]van der Ven, P. F. M.; Speel, E. J. M.; Albrechts, J. C. M.;
Ramaekers,

[141402]F. C. S.; Hopman, A. H. N.; Furst, D. O.: Assignment of the
human

[141403]gene for endosarcomeric cytoskeletal M-protein (MYOM2)
to 8p23.3. Genomics 55:

[141404]253–255, 1999.

[141405]

[141406]

[141407]13192.

[141408]Black, R. A.; Rauch, C. T.; Kozlosky, C. J.; Peschon, J. J.;
Slack,

[141409]J. L.; Wolfson, M. F.; Castner, B. J.; Stocking, K. L.; Reddy,
P.;

[141410]Srinivasan, S.; Nelson, N.; Boiani, N.; Schooley, K. A.; Ger-
hart,

[141411]M.; Davis, R.; Fitzner, J. N.; Johnson, R. S.; Paxton, R. J.;
March,

[141412]C. J.; Cerretti, D. P.: A metalloproteinase disintegrin that releases

[141413]tumour–necrosis factor–alpha from cells. Nature 385: 729–733, 1997.

[141414]

[141415]

[141416]13193.

[141417]Hirohata, S.; Seldin, M. F.; Apte, S. S.: Chromosomal assignment

[141418]of two ADAM genes, TACE (ADAM17) and MLTNB (ADAM19), to human chromosomes

[141419]2 and 5, respectively, and of MltNb to mouse chromosome 11. Genomics 54:

[141420]178–179, 1998.

[141421]

[141422]

[141423]13194.

[141424]Moss, M. L.; Jin, S.–L. C.; Milla, M. E.; Burkhart, W.; Carter,

[141425]H. L.; Chen, W.–J.; Clay, W. C.; Didsbury, J. R.; Hassler, D.; Hoffman,

[141426]C. R.; Kost, T. A.; Lambert, M. H.; and 13 others: Cloning
of a

[141427]disintegrin metalloproteinase that processes precursor tu-
mour-necrosis

[141428]factor-alpha. Nature 385: 733-736, 1997.

[141429]

[141430]

[141431]13195.

[141432]Patel, I. R.; Attur, M. G.; Patel, R. N.; Stuchin, S. A.;
Abagyan,

[141433]R. A.; Abramson, S. B.; Amin, A. R.: TNF-alpha convertase
enzyme

[141434]from human arthritis-affected cartilage: isolation of cDNA
by differential

[141435]display, expression of the active enzyme, and regulation
of TNF-alpha. J.

[141436]Immun. 160: 4570-4579, 1998.

[141437]

[141438]

[141439]13196.

[141440]Peschon, J. J.; Slack, J. L.; Reddy, P.; Stocking, K. L.; Sun-
narborg,

[141441]S. W.; Lee, D. C.; Russell, W. E.; Castner, B. J.; Johnson, R.
S.;

[141442]Fitzner, J. N.; Boyce, R. W.; Nelson, N.; Kozlosky, C. J.;
Wolfson,

[141443]M. F.; Rauch, C. T.; Cerretti, D. P.; Paxton, R. J.; March, C.
J.;

[141444]Black, R. A.: An essential role for ectodomain shedding in
mammalian

[141445]development. Science 282: 1281–1284, 1998.

[141446]

[141447]

[141448]13197.

[141449]Yamazaki, K.; Mizui, Y.; Sagane, K.; Tanaka, I.: Genetic
mapping

[141450]of mouse tumor necrosis factor–alpha converting enzyme
(Tace) to chromosome

[141451]12. Genomics 49: 336–337, 1998.

[141452]

[141453]

[141454]13198.

[141455]Inoue, D.; Reid, M.; Lum, L.; Kratzschmar, J.; Weskamp, G.;
Myung,

[141456]Y. M.; Baron, R.; Blobel, C. P.: Cloning and initial charac-
terization

[141457]of mouse meltrin beta and analysis of the expression of
four metalloprotease-disintegrins

[141458]in bone cells. J. Biol. Chem. 273: 4180–4187, 1998.

[141459]

[141460]

[141461]13199.

[141462]Petruzzella, V.; Tiranti, V.; Fernandez, P.; Ianna, P.; Car-
rozzo,

[141463]R.; Zeviani, M.: Identification and characterization of hu-
man cDNAs

[141464]specific to BCS1, PET112, SCO1, COX15, and COX11, five
genes involved

[141465]in the formation and function of the mitochondrial respi-
ratory chain. Genomics 54:

[141466]494–504, 1998.

[141467]

[141468]

[141469]13200.

[141470]Iyer, N. V.; Leung, S. W.; Semenza, G. L.: The human hypoxia-inducible

[141471]factor 1-alpha gene: HIF1A structure and evolutionary conservation. Genomics 52:

[141472]159-165, 1998.

[141473]

[141474]

[141475]13201.

[141476]Kline, D. D.; Peng, Y.-J.; Manalo, D. J.; Semenza, G. L.; Prabhakar,

[141477]N. R.: Defective carotid body function and impaired ventilatory responses

[141478]to chronic hypoxia in mice partially deficient for hypoxia-inducible

[141479]factor 1-alpha. Proc. Nat. Acad. Sci. 99: 821-826, 2002.

[141480]

[141481]

[141482]13202.

[141483]Lando, D.; Peet, D. J.; Whelan, D. A.; Gorman, J. J.;

Whitelaw,

[141484]M. L.: Asparagine hydroxylation of the HIF transactivation domain:

[141485]a hypoxic switch. Science 295: 858–861, 2002.

[141486]

[141487]

[141488]13203.

[141489]Marti, H. H.; Katschinski, D. M.; Wagner, K. F.; Schaffer, L.;

[141490]Stier, B.; Wenger, R. H.: Isoform–specific expression of hypoxia–inducible

[141491]factor–1–alpha during the late stages of mouse spermiogenesis. Molec.

[141492]Endocr. 16: 234–243, 2002.

[141493]

[141494]

[141495]13204.

[141496]Maxwell, P. H.; Wiesener, M. S.; Chang, G.–W.; Clifford, S. C.;

[141497]Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.;

Maher, E.

[141498]**R.; Ratcliffe, P. J.: The tumour suppressor protein VHL tar-
gets hypoxia-inducible**

[141499]**factors for oxygen-dependent proteolysis. Nature 399:
271-275, 1999.**

[141500]

[141501]

[141502]**13205.**

[141503]**Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W. G., Jr.;**

[141504]**Pavletich, N. P.: Structure of an HIF-1-alpha-pVHL com-
plex: hydroxyproline**

[141505]**recognition in signaling. Science 296: 1886-1889, 2002.**

[141506]

[141507]

[141508]**13206.**

[141509]**Semenza, G. L.: HIF-1 and human disease: one highly in-
volved**

[141510]**factor. Genes Dev. 14: 1983-1991, 2000.**

[141511]

[141512]

[141513]13207.

[141514]Semenza, G. L.; Rue, E. A.; Iyer, N. V.; Pang, M. G.; Kearns,

[141515]W. G.: Assignment of the hypoxia-inducible factor 1- α gene to

[141516]a region of conserved synteny on mouse chromosome 12 and human chromosome

[141517]14q. Genomics 34: 437-439, 1996.

[141518]

[141519]

[141520]13208.

[141521]Sutter, C. H.; Laughner, E.; Semenza, G. L.: Hypoxia-inducible

[141522]factor 1- α protein expression is controlled by oxygen-regulated

[141523]ubiquitination that is disrupted by deletions and missense mutations. Proc.

[141524]Nat. Acad. Sci. 97: 4748-4753, 2000.

[141525]

[141526]

[141527]13209.

[141528]Wenger, R. H.; Rolfs, A.; Kvietikova, I.; Spielmann, P.; Zimmermann,

[141529]D. R.; Gassmann, M.: The mouse gene for hypoxia-inducible factor-1-alpha--genomic

[141530]organization, expression and characterization of an alternative first

[141531]exon and 5-prime flanking sequence. Europ. J. Biochem. 246: 155-165,

[141532]1997.

[141533]

[141534]

[141535]13210.

[141536]Wenger, R. H.; Rolfs, A.; Marti, H. H.; Guenet, J.-L.; Gassmann,

[141537]M.: Nucleotide sequence, chromosomal assignment and mRNA expression

[141538]of mouse hypoxia-inducible factor-1-alpha. Biochem. Biophys. Res.

[141539]Commun. 223: 54-59, 1996.

[141540]

[141541]

[141542]13211.

[141543]Wenger, R. H.; Rolfs, A.; Spielmann, P.; Zimmermann, D.
R.; Gassmann,

[141544]M.: Mouse hypoxia-inducible factor-1-alpha is encoded
by two different

[141545]mRNA isoforms: expression from a tissue-specific and a
housekeeping-type

[141546]promoter. Blood 91: 3471-3480, 1998.

[141547]

[141548]

[141549]13212.

[141550]Tian, H.; Hammer, R. E.; Matsumoto, A. M.; Russell, D. W.;
McKnight,

[141551]S. L.: The hypoxia-responsive transcription factor EPAS1 is
essential

[141552]for catecholamine homeostasis and protection against
heart failure

[141553]during embryonic development. Genes Dev. 12:
3320-3324, 1998.

[141554]

[141555]

[141556]13213.

[141557]Tian, H.; McKnight, S. L.; Russell, D. W.: Endothelial PAS domain

[141558]protein 1 (EPAS1), a transcription factor selectively expressed in

[141559]endothelial cells. Genes Dev. 11: 72–82, 1997.

[141560]

[141561]

[141562]13214.

[141563]Hovanessian, A. G.; Laurent, A. G.; Chebath, J.; Galabru, J.; Robert,

[141564]N.; Svab, J.: Identification of 69–kd and 100–kd forms of 2–5A synthetase

[141565]in interferon–treated human cells by specific monoclonal antibodies. EMBO

[141566]J. 6: 1273–1280, 1987.

[141567]

[141568]

[141569]13215.

[141570]Marie, I.; Galabru, J.; Svab, J.; Hovanessian, A. G.: Preparation

[141571]and characterization of polyclonal antibodies specific for the 69

[141572]and 100 k-dalton forms of human 2-5A synthetase.

Biochem. Biophys.

[141573]Res. Commun. 160: 580-587, 1989.

[141574]

[141575]

[141576]13216.

[141577]Marie, I.; Hovanessian, A. G.: The 69-kDa 2-5A synthetase is composed

[141578]of two homologous and adjacent functional domains. J.

Biol. Chem. 267:

[141579]9933-9939, 1992.

[141580]

[141581]

[141582]13217.

[141583]Kools, P.; Van Imschoot, G.; van Roy, F.: Characterization of

[141584]three novel human cadherin genes (CDH7, CDH19, and CDH20) clustered

[141585]on chromosome 18q22–q23 and with high homology to chicken cadherin–7. Genomics 68:

[141586]283–295, 2000.

[141587]

[141588]

[141589]13218.

[141590]Dantzig, A. H.; Hoskins, J.; Tabas, L. B.; Bright, S.; Shepard,

[141591]R. L.; Jenkins, I. L.; Duckworth, D. C.; Sportsman, J. R.; Mackensen,

[141592]D.; Rosteck, P. R., Jr.; Skatrud, P. L.: Association of intestinal

[141593]peptide transport with a protein related to the cadherin superfamily. Science 264:

[141594]430–433, 1994.

[141595]

[141596]

[141597]13219.

[141598]Shibata, T.; Shimoyama, Y.; Gotoh, M.; Hirohashi, S.: Identification

[141599]of human cadherin-14, a novel neurally specific type II cadherin,

[141600]by protein interaction cloning. J. Biol. Chem. 272: 5236-5240, 1997.

[141601]

[141602]

[141603]13220.

[141604]Burn, T. C.; Connors, T. D.; Van Raay, T. J.; Dackowski, W. R.;

[141605]Millholland, J. M.; Klinger, K. W.; Landes, G. M.: Generation of

[141606]a transcriptional map for a 700-kb region surrounding the polycystic

[141607]kidney disease type 1 (PKD1) and tuberous sclerosis type 1 (TSC2)

[141608]disease genes on human chromosome 16p13.3. Genome Res. 6: 525-537,

[141609]1996.

[141610]

[141611]

[141612]13221.

[141613]Rooney, R. J.; Daniels, R. R.; Jenkins, N. A.; Gilbert, D. J.;

[141614]Rothhammer, K.; Morris, S. W.; Higgs, D. R.; Copeland, N.
G.: Chromosomal

[141615]location and tissue expression of the gene encoding the
adenovirus

[141616]E1A-regulated transcription factor E4F in humans and
mice. Mammalian

[141617]Genome 9: 320–323, 1998.

[141618]

[141619]

[141620]13222.

[141621]Saccone, S.; Sandy, P.; Meroni, G.; Gostissa, M.; Della
Valle,

[141622]G.; Del Sal, G.: Assignment of the E1A-regulated tran-
scription factor

[141623]E4F gene (E4F1) to human chromosome band 16p13.3 by
in situ hybridization

[141624]and somatic cell hybrids. Cytogenet. Cell Genet. 82:
99–100, 1998.

[141625]

[141626]

[141627]13223.

[141628]Barabino, S. M. L.; Hubner, W.; Jenny, A.; Minvielle-Sebastia,
tia,

[141629]L.; Keller, W.: The 30 kDa subunit of mammalian cleavage
and polyadenylation

[141630]specificity factor and its yeast homolog are RNA binding
zinc finger

[141631]proteins. Genes Dev. 11: 1703–1716, 1997.

[141632]

[141633]

[141634]13224.

[141635]Nemeroff, M. E.; Barabino, S. M. L.; Li, Y.; Keller, W.; Krug,

[141636]R. M.: Influenza virus NS1 protein interacts with the cellular 30
lar 30

[141637]kDa subunit of CPSF and inhibits 3–prime end formation
of cellular

[141638]pre-mRNAs. Molec. Cell 1: 991–1000, 1998.

[141639]

[141640]

[141641]13225.

[141642]Dutnall, R. N.; Tafrov, S. T.; Sternglanz, R.; Ramakrishnan, V.

[141643]: Structure of the histone acetyltransferase Hat1: a paradigm for

[141644]the GCN5-related N-acetyltransferase superfamily. Cell 94: 427–438,

[141645]1998.

[141646]

[141647]

[141648]13226.

[141649]Verreault, A.; Kaufman, P. D.; Kobayashi, R.; Stillman, B.: Nucleosomal

[141650]DNA regulates the core-histone-binding subunit of the human Hat1 acetyltransferase. Curr.

[141651]Biol. 8: 96–108, 1998.

[141652]

[141653]

[141654]13227.

[141655]Hsu, D. R.; Economides, A. N.; Wang, X.; Eimon, P. M.;
Harland,

[141656]R. M.: The *Xenopus* dorsalizing factor gremlin identifies a
novel

[141657]family of secreted proteins that antagonize BMP activities.
Molec.

[141658]Cell 1: 673–683, 1998.

[141659]

[141660]

[141661]13228.

[141662]Topol, L. Z.; Modi, W. S.; Koochekpour, S.; Blair, D. G.:

DRM-Gremlin

[141663](CKTSF1B1) maps to human chromosome 15 and is highly
expressed in

[141664]adult and fetal brain. Cytogenet. Cell Genet. 89: 79–84,
2000.

[141665]

[141666]

[141667]13229.

[141668]Beaconsfield, P.; Rainsbury, R.; Kalton, G.: Glucose–

6–phosphate

[141669]dehydrogenase deficiency and the incidence of cancer.

Oncologia 19:

[141670]11–19, 1965.

[141671]

[141672]

[141673]13230.

[141674]Ben-Bassat, J.; Ben-Ishay, D.: Hereditary hemolytic anemia

associated

[141675]with glucose-6-phosphate dehydrogenase deficiency
(Mediterranean type). Israel

[141676]J. Med. Sci. 5: 1053-1059, 1969.

[141677]

[141678]

[141679]13231.

[141680]Benabadji, M.; Merad, F.; Benmoussa, M.; Trabuchet, G.;
Junien,

[141681]C.; Dreyfus, J. C.; Kaplan, J. C.: Heterogeneity of glucose-
6-phosphate

[141682]dehydrogenase deficiency in Algeria. Hum. Genet. 40:
177-184, 1978.

[141683]

[141684]

[141685]13232.

[141686]Benohr, H. C.; Klumpp, F.; Waller, H. D.: Glucose-
6-phosphat-Dehydrogenase

[141687]Typ Schwaben. Dtsch. Med. Wschr. 96: 1029-1032, 1971.

[141688]

[141689]

[141690]13233.

[141691]Benohr, H. C.; Waller, H. D.: Eigenschaften der Glucose-
6-p-dehydrogenase,

[141692]Typ Tübingen. Klin. Wschr. 48: 71–74, 1970.

[141693]

[141694]

[141695]13234.

[141696]Benohr, H. C.; Waller, H. D.; Arnold, H.; Blume, K. G.; Lohr,

[141697]G. W.: Glucose-6-P-Dehydrogenase Typ Bodensee (eine
neue Enzymvariante). Klin.

[141698]Wschr. 49: 1058–1062, 1971.

[141699]

[141700]

[141701]13235.

[141702]Beutler, E.: Glucose 6-phosphate dehydrogenase defi-
ciency, a

[141703]new Indian variant, G6PD Jammu. In: Sen, N. N.; Basu, A.
K.: Trends

[141704]in Haematology. Calcutta: Chatterjea Memorial Committee
(pub.)

[141705]1975. Pp. 279–283.

[141706]

[141707]

[141708]13236.

[141709]Beutler, E.: Personal Communication. La Jolla, Calif.
11/12/1990.

[141710]

[141711]

[141712]13237.

[141713]Beutler, E.: G6PD deficiency. Blood 84: 3613–3636, 1994.

[141714]

[141715]

[141716]13238.

[141717]Beutler, E.: Glucose–6–phosphate dehydrogenase defi-
ciency. New

[141718]Eng. J. Med. 324: 169–174, 1991.

[141719]

[141720]

[141721]13239.

[141722]Beutler, E.: Selectivity of proteases as a basis for tissue distribution

[141723]of enzymes in hereditary deficiencies. Proc. Nat. Acad. Sci. 80:

[141724]3767–3768, 1983.

[141725]

[141726]

[141727]13240.

[141728]Beutler, E.: Glucose–6–phosphate dehydrogenase deficiency.In:

[141729]Wintrobe, M. M.: Red Cell Metabolism in Hemolytic Anemia. New York:

[141730]Plenum Press (pub.) 1978.

[141731]

[141732]

[141733]13241.

[141734]Beutler, E.: The hemolytic effect of primaquine and related

compounds:

[141735]a review. Blood 14: 103–139, 1959.

[141736]

[141737]

[141738]13242.

[141739]Beutler, E.; Grooms, A. M.; Morgan, S. K.; Trinidad, F.:

Chronic

[141740]severe hemolytic anemia due to G-6-PD Charleston: a new
deficiency

[141741]variant. J. Pediat. 80: 1005–1009, 1972.

[141742]

[141743]

[141744]13243.

[141745]Beutler, E.; Hartman, K.; Gelbart, T.; Forman, L.: G-6-PD

Walter

[141746]Reed: possible insight into 'structural' NADP in G-6-PD.

Am. J. Hemat. 23:

[141747]25–30, 1986.

[141748]

[141749]

[141750]13244.

[141751]Beutler, E.; Keller, J. W.; Matsumoto, F.: A new glucose
6-phosphate

[141752]dehydrogenase (G6PD) variant associated with nonsphero-
cytic hemolytic

[141753]anemia: G6PD Atlanta. I.R.C.S. 4: 479, 1976.

[141754]

[141755]

[141756]13245.

[141757]Beutler, E.; Kuhl, W.: The NT 1311 polymorphism of G6PD:
G6PD

[141758]Mediterranean mutation may have originated indepen-
dently in Europe

[141759]and Asia. Am. J. Hum. Genet. 47: 1008-1012, 1990.

[141760]

[141761]

[141762]13246.

[141763]Beutler, E.; Kuhl, W.: Linkage between a PvuII restriction

fragment

[141764]length polymorphism and G6PD A- (202A/376G): evidence for a single

[141765]origin of the common G6PD A- mutation. Hum. Genet. 85: 9-11, 1990.

[141766]

[141767]

[141768]13247.

[141769]Beutler, E.; Kuhl, W.; Gelbart, T.; Forman, L.: DNA sequence

[141770]abnormalities of human glucose-6-phosphate dehydrogenase variants. J.

[141771]Biol. Chem. 266: 4145-4150, 1991.

[141772]

[141773]

[141774]13248.

[141775]Beutler, E.; Kuhl, W.; Ramirez, E.; Lisker, R.: Some Mexican

[141776]glucose-6-phosphate dehydrogenase variants revisited. Hum. Genet. 86:

[141777]371-374, 1991.

[141778]

[141779]

[141780]13249.

[141781]Beutler, E.; Kuhl, W.; Saenz, G. F.; Rodriguez, W.: Mutation

[141782]analysis of glucose-6-phosphate dehydrogenase (G6PD)
variants in Costa

[141783]Rica. Hum. Genet. 87: 462-464, 1991.

[141784]

[141785]

[141786]13250.

[141787]Beutler, E.; Mathai, C. K.; Smith, J. E.: Biochemical variants

[141788]of glucose-6-phosphate dehydrogenase giving rise to
congenital nonspherocytic

[141789]hemolytic disease. Blood 31: 131-150, 1968.

[141790]

[141791]

[141792]13251.

[141793]Beutler, E.; Matsumoto, F.: A new glucose 6-phosphate

dehydrogenase

[141794]variant: G6PD (-) Los Angeles. I.R.C.S. 5: 89, 1977.

[141795]

[141796]

[141797]13252.

[141798]Beutler, E.; Matsumoto, F.; Daiber, A.: Nonspherocytic
hemolytic

[141799]anemia due to G-6-PD Panama. I.R.C.S. 2: 1389, 1974.

[141800]

[141801]

[141802]13253.

[141803]Beutler, E.; Rosen, R.: Nonspherocytic congenital
hemolytic anemia

[141804]due to a new G-6-PD variant: G-6-PD Alhambra. Pedi-
atrics 45: 230-235,

[141805]1970.

[141806]

[141807]

[141808]13254.

- [141809]Beutler, E.; Westwood, B.; Prchal, J. T.; Vaca, G.; Bartsocas,
[141810]C. S.; Baronciani, L.: New glucose-6-phosphate dehydro-
genase mutations
[141811]from various ethnic groups. Blood 80: 255-256, 1992.
[141812]
[141813]
[141814]13255.
[141815]Beutler, E.; Yoshida, A.: Human glucose-6-phosphate de-
hydrogenase
[141816]variants: a supplementary tabulation. Ann. Hum. Genet.
37: 151-156,
[141817]1973.
[141818]
[141819]
[141820]13256.
[141821]Boivin, P.; Galand, C.: Nouvelles variantes de la glucose-
6-phosphate
[141822]dehydrogenase erythrocytaire. Rev. Franc. Etud. Clin. Biol.
13:
[141823]30-39, 1968.

[141824]

[141825]

[141826]13257.

[141827]Botha, M. C.; Dern, R. J.; Mitchell, M.; West, C.; Beutler, E.

[141828]: G6PD Capetown, a variant of glucose-6-phosphate dehydrogenase. Am.

[141829]J. Hum. Genet. 21: 547-551, 1969.

[141830]

[141831]

[141832]13258.

[141833]Boyer, S. H.; Graham, J. B.: Linkage between the X chromosome

[141834]loci for glucose-6-phosphate dehydrogenase electrophoretic variation

[141835]and hemophilia A. Am. J. Hum. Genet. 17: 320-324, 1965.

[141836]

[141837]

[141838]13259.

[141839]Boyer, S. H.; Porter, I. H.; Weilbaecher, R. G.: Elec-

trophoretic

[141840]heterogeneity of glucose-6-phosphate dehydrogenase
and its relationship

[141841]to enzyme deficiency in man. Proc. Nat. Acad. Sci. 48:
1868-1876,

[141842]1962.

[141843]

[141844]

[141845]13260.

[141846]Busch, D.; Bote, K.: Glucose-
6-phosphate-dehydrogenase-Defect

[141847]in Deutschland. II. Bigenschabten des Enzyms (Typ
Freiburg). Klin.

[141848]Wschr. 48: 74-78, 1970.

[141849]

[141850]

[141851]13261.

[141852]Calabro, V.; Giacobbe, A.; Vallone, D.; Montanaro, V.; Cas-
cone,

[141853]A.; Filosa, S.; Battistuzzi, G.: Genetic heterogeneity at the

glucose-6-phosphate

[141854]dehydrogenase locus in southern Italy: a study on a population from

[141855]the Matera district. Hum. Genet. 86: 49-53, 1990.

[141856]

[141857]

[141858]13262.

[141859]Cappadoro, M.; Giribaldi, G.; O'Brien, E.; Turrini, F.;
Mannu,

[141860]F.; Ulliers, D.; Simula, G.; Luzzatto, L.; Arese, P.: Early
phagocytosis

[141861]of glucose-6-phosphate dehydrogenase (G6PD)-deficient
erythrocytes

[141862]parasitized by Plasmodium falciparum may explain
malaria protection

[141863]in G6PD deficiency. Blood 92: 2527-2534, 1998.

[141864]

[141865]

[141866]13263.

[141867]Cappellini, M. D.; Sampietro, M.; Toniolo, D.; Carandina,

G.;

[141868]Pittalis, S.; Martinez di Montemuros, F.; Tavazzi, D.;
Fiorelli, G.

[141869]: Biochemical and molecular characterization of a new
sporadic glucose-6-phosphate

[141870]dehydrogenase variant described in Italy: G6PD Modena.

Brit. J. Haemat. 87:

[141871]209-211, 1994.

[141872]

[141873]

[141874]13264.

[141875]Carandina, G.; Moretto, E.; Zecchi, G.; Conighi, C.: Glucose

[141876]6-phosphate dehydrogenase Ferrara. A new variant of
G6PD identified

[141877]in Northern Italy. Acta Haemat. 56: 116-122, 1976.

[141878]

[141879]

[141880]13265.

[141881]Carson, P. E.; Flanagan, C. L.; Ickes, C. E.; Alving, A. S.:

[141882]Enzymatic deficiency in primaquine-sensitive erythrocytes. Science 124:

[141883]484–485, 1956.

[141884]

[141885]

[141886]13266.

[141887]Poy, F.; Yaffe, M. B.; Sayos, J.; Saxena, K.; Morra, M.; Sumegi,

[141888]J.; Cantley, L. C.; Terhorst, C.; Eck, M. J.: Crystal structures

[141889]of the XLP protein SAP reveal a class of SH2 domains with extended,

[141890]phosphotyrosine-independent sequence recognition.

Molec. Cell 4:

[141891]555–561, 1999.

[141892]

[141893]

[141894]13267.

[141895]Provisor, A. J.; Iaccone, J. J.; Chilcote, R. R.; Neiburger, R.

[141896]G.; Crussi, F. G.; Baehner, R. L.: Acquired agammaglobu-

linemia after

[141897]a life-threatening illness with clinical and laboratory features of

[141898]infectious mononucleosis in three related male children.

New Eng.

[141899]J. Med. 293: 62–65, 1975.

[141900]

[141901]

[141902]13268.

[141903]Purtilo, D. T.: Pathogenesis and phenotypes of an X-linked recessive

[141904]lymphoproliferative syndrome. Lancet II: 882–885, 1976.

[141905]

[141906]

[141907]13269.

[141908]Purtilo, D. T.: X-linked lymphoproliferative syndrome: an immunodeficiency

[141909]disorder with acquired agammaglobulinemia, fatal infectious mononucleosis,

[141910]or malignant lymphoma. Arch. Path. Lab. Med. 105:

119-121, 1981.

[141911]

[141912]

[141913]13270.

[141914]Purtilo, D. T.; Bhawan, J.; Hutt, L. M.; De Nicola, L.; Szymanski,

[141915]I.; Yang, J. P. S.; Boto, W.; Naier, R.; Thorley-Lawson, D.:
Epstein-Barr

[141916]virus in the X-linked recessive lymphoproliferative syndrome. Lancet I:

[141917]798-801, 1978.

[141918]

[141919]

[141920]13271.

[141921]Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.: Fatal infectious

[141922]mononucleosis in familial lymphohistiocytosis. (Letter)
New Eng.

[141923]J. Med. 201: 736 only, 1974.

[141924]

[141925]

[141926]13272.

[141927]Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.; Harper, R.;
Stephenson,

[141928]S. R.; Landing, B. H.; Vewter, G. F.: X-linked recessive pro-
gressive

[141929]combined variable immunodeficiency (Duncan's disease).
Lancet I:

[141930]935-941, 1975.

[141931]

[141932]

[141933]13273.

[141934]Purtilo, D. T.; DeFlorio, D., Jr.; Hutt, L. M.; Bhawan, J.;
Yang,

[141935]J. P. S.; Otto, R. L.; Edwards, W.: Variable phenotypic ex-
pression

[141936]of an X-linked recessive lymphoproliferative syndrome.
New Eng. J.

[141937]Med. 297: 1077-1081, 1977.

[141938]

[141939]

[141940]13274.

[141941]Purtilo, D. T.; Grierson, H. L.: Methods of detection of new
[141942]families with X-linked lymphoproliferative disease. Cancer
Genet.

[141943]Cytogenet. 51: 143–153, 1991.

[141944]

[141945]

[141946]13275.

[141947]Purtilo, D. T.; Sakamoto, K.; Barnabei, V.; Seeley, J.; Bech-
told,

[141948]T.; Rogers, G.; Yetz, J.; Harada, S.; the XLP collaborators:
Epstein–Barr

[141949]virus-induced diseases in boys with the X-linked lympho-
proliferative

[141950]syndrome (XLP): update on studies of the registry. Am. J.
Med. 73:

[141951]49–56, 1982.

[141952]

[141953]

[141954]13276.

[141955]Purtilo, D. T.; Yang, J. P. S.; Allegra, S.; DeFlorio, D.; Hutt,

[141956]L. M.; Soltani, M.; Vawter, G. F.: Hematopathology and
pathogenesis

[141957]of the X-linked recessive lymphoproliferative syndrome.

Am. J. Med. 62:

[141958]225-233, 1977.

[141959]

[141960]

[141961]13277.

[141962]Sanger, W. G.; Grierson, H. L.; Skare, J.; Wyandt, H.; Pirruccello,

[141963]S.; Fordyce, R.; Purtilo, D. T.: Partial Xq25 deletion in a
family

[141964]with the X-linked lymphoproliferative disease (XLP). Can-
cer Genet.

[141965]Cytogenet. 47: 163-169, 1990.

[141966]

[141967]

[141968]13278.

[141969]Sayos, J.; Wu, C.; Morra, M.; Wang, N.; Zhang, X.; Allen, D.;

[141970]van Schaik, S.; Notarangelo, L.; Gehat, R.; Roncarolo, M.

G.; Oettgen,

[141971]H.; De Vries, J. E.; Aversall, G.; Terhorst, C.: The X-linked lymphoproliferative-disease

[141972]gene product SAP regulates signals induced through the co-receptor

[141973]SLAM. Nature 395: 462–469, 1998.

[141974]

[141975]

[141976]13279.

[141977]Scher, I.: The CBA/N mouse strain: an experimental model illustrating

[141978]the influence of the X-chromosome on immunity. Adv. Immun. 33: 1–71,

[141979]1982.

[141980]

[141981]

[141982]13280.

[141983]Schuster, V.; Kreth, H. W.: X-linked lymphoproliferative

disease.In:

[141984]Ochs, H. D.; Smith, C. I. E.; Puck, J. M. (eds.): Primary Immunodeficiency

[141985]Diseases: A Molecular and Genetic Approach. New York: Oxford University

[141986]Press 1999. Pp. 222–232.

[141987]

[141988]

[141989]13281.

[141990]Seemayer, T. A.; Gross, T. G.; Egeler, R. M.; Pirruccello, S.

[141991]J.; Davis, D. J.; Kelly, C. M.; Okano, M.; Lanyi, A.; Sumegi, J.:

[141992]X-linked lymphoproliferative disease: twenty-five years after the

[141993]discovery. Pediat. Res. 38: 471–478, 1995.

[141994]

[141995]

[141996]13282.

[141997]Skare, J.; Grierson, H.; Wyandt, H.; Sanger, W.; Milunsky, J.;

[141998]Purtilo, D.; Sullivan, J.; Milunsky, A.: Genetics of the X-linked

[141999]lymphoproliferative syndrome. (Abstract) Am. J. Hum. Genet. 45 (suppl.):

[142000]A161 only, 1989.

[142001]

[142002]

[142003]13283.

[142004]Skare, J.; Madan, S.; Glaser, J.; Purtilo, D.; Nitowsky, H.; Pulijaal,

[142005]V.; Milunsky, A.: First prenatal diagnosis of X-linked lymphoproliferative

[142006]disease. Am. J. Med. Genet. 44: 79–81, 1992.

[142007]

[142008]

[142009]13284.

[142010]Skare, J.; Milunsky, A.; Byron, K.; Sullivan, J.: The mutation

[142011]causing X-linked lymphoproliferative syndrome lies in Xq26. (Abstract) Am.

[142012]J. Hum. Genet. 41: A185 only, 1987.

[142013]

[142014]

[142015]13285.

[142016]Skare, J.; Wu, B.-L.; Madan, S.; Pulijaal, V.; Purtilo, D.;
Haber,

[142017]D.; Nelson, D.; Sylla, B.; Grierson, H.; Nitowsky, H.; Glaser,
J.;

[142018]Wissink, J.; White, B.; Holden, J.; Housman, D.; Lenoir, G.;
Wyandt,

[142019]H.; Milunsky, A.: Characterization of three overlapping
deletions

[142020]causing X-linked lymphoproliferative disease. Genomics
16: 254-255,

[142021]1993.

[142022]

[142023]

[142024]13286.

[142025]Skare, J. C.; Grierson, H. L.; Sullivan, J. L.; Nussbaum, R. L.;

[142026]Purtilo, D. T.; Sylla, B. S.; Lenoir, G. M.; Reilly, D. S.; White,

[142027]B. N.; Milunsky, A.: Linkage analysis of seven kindreds

with the

[142028]X-linked lymphoproliferative syndrome (XLP) confirms
that the XLP

[142029]locus is near DXS42 and DXS37. Hum. Genet. 82:
354–358, 1989.

[142030]

[142031]

[142032]13287.

[142033]Skare, J. C.; Milunsky, A.; Byron, K. S.; Sullivan, J. L.: Map-
ping

[142034]the X-linked lymphoproliferative syndrome. Proc. Nat.
Acad. Sci. 84:

[142035]2015–2018, 1987.

[142036]

[142037]

[142038]13288.

[142039]Skare, J. C.; Sullivan, J. L.; Milunsky, A.: Mapping the mu-
tation

[142040]causing the X-linked lymphoproliferative syndrome in re-
lation to restriction

[142041]fragment length polymorphisms on Xq. Hum. Genet. 82:

349–353, 1989.

[142042]

[142043]

[142044]13289.

[142045]Steinherz, R.; Levy, Y.; Litwin, A.; Nitzan, M.; Friedman, E.;

[142046]Levin, S.: X-linked lymphoproliferative syndrome: a new
kindred with

[142047]variable phenotypic expression. Am. J. Dis. Child. 139:
191–193,

[142048]1985.

[142049]

[142050]

[142051]13290.

[142052]Nehlin, J. O.; Hara, E.; Kuo, W.-L.; Collins, C.; Campisi, J.:

[142053]Genomic organization, sequence, and chromosomal local-
ization of the

[142054]human helix–loop–helix Id1 gene. Biochem. Biophys. Res.
Commun. 231:

[142055]628–634, 1997.

[142056]

[142057]

[142058]13291.

[142059]Loftin, C. D.; Trivedi, D. B.; Tiano, H. F.; Clark, J. A.; Lee,

[142060]C. A.; Epstein, J. A.; Morham, S. G.; Breyer, M. D.; Nguyen,
M.; Hawkins,

[142061]B. M.; Goulet, J. L.; Smithies, O.; Koller, B. H.; Langenbach,
R.

[142062]: Failure of ductus arteriosus closure and remodeling in
neonatal

[142063]mice deficient in cyclooxygenase-1 and cyclooxygenase-2.
Proc. Nat.

[142064]Acad. Sci. 98: 1059-1064, 2001.

[142065]

[142066]

[142067]13292.

[142068]Macchia, L.; Di Paola, R.; Guerrese, M.-C.; Chiechi, L. M.;
Tursi,

[142069]A.; Caiaffa, M. F.; Haeggstrom, J. Z.: Expression of
prostaglandin

[142070]endoperoxide H synthase 1 and 2 in human placenta at term. Biochem.

[142071]Biophys. Res. Commun. 233: 496–501, 1997.

[142072]

[142073]

[142074]13293.

[142075]Morham, S. G.; Langenbach, R.; Loftin, C. D.; Tiano, H. F.; Vouloumanos,

[142076]N.; Jennette, J. C.; Mahler, J. F.; Kluckman, K. D.; Ledford, A.;

[142077]Lee, C. A.; Smithies, O.: Prostaglandin synthase 2 gene disruption

[142078]causes severe renal pathology in the mouse. Cell 83: 473–482, 1995.

[142079]

[142080]

[142081]13294.

[142082]Neufang, G.; Furstenberger, G.; Heidt, M.; Marks, F.; Muller–Decker,

[142083]K.: Abnormal differentiation of epidermis in transgenic mice constitutively

[142084]expressing cyclooxygenase-2 in skin. Proc. Nat. Acad. Sci.
98: 7629-7634,
[142085]2001.

[142086]

[142087]

[142088]13295.

[142089]O'Banion, M. K.; Sadowski, H. B.; Winn, V.; Young, D. A.: A
serum-

[142090]and glucocorticoid-regulated 4-kilobase mRNA encodes a
cyclooxygenase-related

[142091]protein. J. Biol. Chem. 266: 23261-23267, 1991.

[142092]

[142093]

[142094]13296.

[142095]O'Banion, M. K.; Winn, V. D.; Young, D. A.: cDNA cloning
and

[142096]functional activity of a glucocorticoid-regulated inflam-
matory cyclooxygenase. Proc.

[142097]Nat. Acad. Sci. 89: 4888-4892, 1992.

[142098]

[142099]

[142100]13297.

[142101]Oshima, M.; Dinchuk, J. E.; Kargman, S. L.; Oshima, H.;
Hancock,

[142102]B.; Kwong, E.; Trzaskos, J. M.; Evans, J. F.; Taketo, M. M.:
Suppression

[142103]of intestinal polyposis in Apc(delta-716) knockout mice by
inhibition

[142104]of cyclooxygenase 2 (COX-2). Cell 87: 803-809, 1996.

[142105]

[142106]

[142107]13298.

[142108]Salmenkivi, K.; Haglund, C.; Ristimäki, A.; Arola, J.;
Heikkilä,

[142109]P.: Increased expression of cyclooxygenase-2 in malignant pheochromocytomas. J.

[142110]Clin. Endocr. Metab. 86: 5615-5619, 2001.

[142111]

[142112]

[142113]13299.

[142114]Tay, A.; Squire, J. A.; Goldberg, H.; Skorecki, K.: Assign-
ment

[142115]of the human prostaglandin-endoperoxide synthase 2
(PTGS2) gene to

[142116]1q25 by fluorescence in situ hybridization. Genomics 23:
718-719,

[142117]1994.

[142118]

[142119]

[142120]13300.

[142121]Swan, S. K.; Rudy, D. W.; Lasseter, K. C.; Ryan, C. F.;
Buechel,

[142122]K. L.; Lambrecht, L. J.; Pinto, M. B.; Dilzer, S. C.; Obrda, O.;
Sundblad,

[142123]K. J.; Gumbs, C. P.; Ebel, D. L.; Quan, H.; Larson, P. J.;
Schwartz,

[142124]J. I.; Musliner, T. A.; Gertz, B. J.; Brater, D. C.; Yao, S.-L.:

[142125]Effect of cyclooxygenase-2 inhibition on renal function in
elderly

[142126]persons receiving a low-salt diet: a randomized controlled
trial. Ann.

[142127]Intern. Med. 133: 1–9, 2000.

[142128]

[142129]

[142130]13301.

[142131]Tazawa, R.; Xu, X.–M.; Wu, K. K.; Wang, L.–H.: Characteri-
zation

[142132]of the genomic structure, chromosomal location and pro-
moter of human

[142133]prostaglandin H synthase–2 gene. Biochem. Biophys. Res.
Commun. 203:

[142134]190–199, 1994.

[142135]

[142136]

[142137]13302.

[142138]Tsujii, M.; DuBois, R. N.: Alterations in cellular adhesion
and

[142139]apoptosis in epithelial cells overexpressing prostaglandin
endoperoxide

[142140]synthase 2. Cell 83: 493–501, 1995.

[142141]

[142142]

[142143]13303.

[142144]Zhou, X.-L.; Lei, Z. M.; Rao, C. V.: Treatment of human
endometrial

[142145]gland epithelial cells with chorionic gonadotropin/luteiniz-
ing hormone

[142146]increases the expression of the cyclooxygenase-2 gene. J.
Clin. Endocr.

[142147]Metab. 84: 3364-3377, 1999.

[142148]

[142149]

[142150]13304.

[142151]Erdel, M.; Baier-Bitterlich, G.; Duba, C.; Isakov, N.; Altman,

[142152]A.; Utermann, G.; Baier, G.: Mapping of the human protein
kinase

[142153]C-theta (PRKCQ) gene locus to the short arm of chromo-
some 10 (10p15)

[142154]by FISH. Genomics 25: 595-597, 1995.

[142155]

[142156]

[142157]13305.

[142158]Sun, Z.; Arendt, C. W.; Ellmeier, W.; Schaeffer, E. M.; Sunshine,

[142159]M. J.; Gandhi, L.; Annes, J.; Petrzilka, D.; Kupfer, A.; Schwartzberg,

[142160]P. L.; Littman, D. R.: PKC-theta is required for TCR-induced NF-kappa-B

[142161]activation in mature but not immature T lymphocytes. Nature 404:

[142162]402-407, 2000.

[142163]

[142164]

[142165]13306.

[142166]Maho, A.; Bensimon, A.; Vassart, G.; Parmentier, M.: Mapping of

[142167]the CCXCR1, CX3CR1, CCBP2 and CCR9 genes to the CCR cluster within

[142168]the 3p21.3 region of the human genome. Cytogenet. Cell Genet. 87:

[142169]265-268, 1999.

[142170]

[142171]

[142172]13307.

[142173]Anderson, D. M.; Johnson, L.; Glaccum, M. B.; Copeland,
N. G.;

[142174]Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Kirstein, M. N.;
Shapiro,

[142175]D. N.; Morris, S. W.; Grabstein, K.; Cosman, D.: Chromoso-
mal assignment

[142176]and genomic structure of IL15. Genomics 25: 701–706,
1995.

[142177]

[142178]

[142179]13308.

[142180]Grabstein, K. H.; Eisenman, J.; Shanebeck, K.; Rauch, C.;
Srinivasan,

[142181]S.; Fung, V.; Beers, C.; Richardson, J.; Schoenborn, M. A.;
Ahdieh,

[142182]M.; Johnson, L.; Alderson, M. R.; Watson, J. D.; Anderson,
D. M.;

[142183]Giri, J. G.: Cloning of a T cell growth factor that interacts
with

[142184]the beta chain of the interleukin-2 receptor. Science 264:
965-968,
[142185]1994.

[142186]

[142187]

[142188]13309.

[142189]Roberts, A. I.; Lee, L.; Schwarz, E.; Groh, V.; Spies, T.;
Ebert,

[142190]E. C.; Jabri, B.: Cutting edge: NKG2D receptors induced by
IL-15

[142191]costimulate CD28-negative effector CTL in the tissue mi-
croenvironment. J.

[142192]Immun. 167: 5527-5530, 2001.

[142193]

[142194]

[142195]13310.

[142196]Robinson, P.; Okhuysen, P. C.; Chappell, C. L.; Lewis, D. E.;
Shahab,

[142197]I.; Lahoti, S.; White, A. C., Jr.: Expression of IL-15 and IL-4
in

[142198]IFN-gamma-independent control of experimental human
Cryptosporidium

[142199]parvum infection. Cytokine 15: 39-46, 2001.

[142200]

[142201]

[142202]13311.

[142203]Chen, X.; Vinkemeier, U.; Zhao, Y.; Jeruzalmi, D.; Darnell,
J.

[142204]E., Jr.; Kuriyan, J.: Crystal structure of a tyrosine phospho-
rylated

[142205]STAT-1 dimer bound to DNA. Cell 93: 827-839, 1998.

[142206]

[142207]

[142208]13312.

[142209]Copeland, N. G.; Gilbert, D. J.; Schindler, C.; Zhong, Z.;
Wen,

[142210]Z.; Darnell, J. E., Jr.; Mui, A. L.-F.; Miyajima, A.; Quelle, F.
W.;

[142211]Ihle, J. N.; Jenkins, N. A.: Distribution of the mammalian
Stat gene

[142212]family in mouse chromosomes. Genomics 29: 225–228,
1995.

[142213]

[142214]

[142215]13313.

[142216]Darnell, J. E., Jr.; Kerr, I. M.; Stark, G. M.: Jak–STAT path–
ways

[142217]and transcriptional activation in response to IFNs and
other extracellular

[142218]signaling proteins. Science 264: 1415–1421, 1994.

[142219]

[142220]

[142221]13314.

[142222]Durbin, J. E.; Hackenmiller, R.; Simon, M. C.; Levy, D. E.:
Targeted

[142223]disruption of the mouse Stat1 gene results in compro–
mised innate immunity

[142224]to viral disease. Cell 84: 443–450, 1996.

[142225]

[142226]

[142227]13315.

[142228]Haddad, B.; Pabon-Pena, C. R.; Young, H.; Sun, W. H.: Assignment
signment

[142229]of STAT1 to human chromosome 2q32 by FISH and radiation hybrids. Cytogenet.

[142230]Cell Genet. 83: 58-59, 1998.

[142231]

[142232]

[142233]13316.

[142234]Ihle, J. N.: STATs: signal transducers and activators of
transcription. Cell 84:

[142235]331-334, 1996.

[142236]

[142237]

[142238]13317.

[142239]Meraz, M. A.; White, J. M.; Sheehan, K. C. F.; Bach, E. A.;
Rodig,

[142240]S. J.; Dighe, A. S.; Kaplan, D. H.; Riley, J. K.; Greenlund, A.
C.

[142241]Campbell, D.; Carver-Moore, K.; DuBois, R. N.; Clark, R.;

Aguet, M.;

[142242]Schreiber, R. D.: Targeted disruption of the Stat1 gene in mice reveals

[142243]unexpected physiologic specificity in the JAK-STAT signaling pathway. Cell 84:

[142244]431-442, 1996.

[142245]

[142246]

[142247]13318.

[142248]Mowen, K. A.; Tang, J.; Zhu, W.; Schurter, B. T.; Shuai, K.; Herschman,

[142249]H. R.; David, M.: Arginine methylation of STAT1 modulates IFN-alpha/beta-induced

[142250]transcription. Cell 104: 731-741, 2001.

[142251]

[142252]

[142253]13319.

[142254]Ramana, C. V.; Chatterjee-Kishore, M.; Nguyen, H.; Stark, G. R.

[142255]: Complex roles of Stat1 in regulating gene expression.

Oncogene 19:

[142256]2619–2627, 2000.

[142257]

[142258]

[142259]13320.

[142260]Yamamoto, K.; Kobayashi, H.; Arai, A.; Miura, O.; Hiro-
sawa, S.;

[142261]Miyasaka, N.: cDNA cloning, expression and chromosome
mapping of

[142262]the human STAT4 gene: both STAT4 and STAT1 genes are
mapped to 2q32.2–q32.3. Cytogenet.

[142263]Cell Genet. 77: 207–210, 1997.

[142264]

[142265]

[142266]13321.

[142267]Ware, R. E.; Howard, T. A.; Kamitani, T.; Chang, H.–M.;
Yeh, E.

[142268]T. H.; Seldin, M. F.: Chromosomal assignment of genes in-
volved in

[142269]glycosylphosphatidylinositol anchor biosynthesis: implica-

tions for

[142270]the pathogenesis of paroxysmal nocturnal hemoglobin-
uria. Blood 83:

[142271]3753–3757, 1994.

[142272]

[142273]

[142274]13322.

[142275]Watanabe, R.; Inoue, N.; Westfall, B.; Taron, C. H.; Orlean,
P.;

[142276]Takeda, J.; Kinoshita, T.: The first step of glycosylphos-
phatidylinositol

[142277]biosynthesis is mediated by a complex of PIG-A, PIG-H,
PIG-C and GPI1. EMBO

[142278]J. 17: 877–885, 1998.

[142279]

[142280]

[142281]13323.

[142282]Watanabe, R.; Kinoshita, T.; Masaki, R.; Yamamoto, A.;
Takeda,

[142283]J.; Inoue, N.: PIG-A and PIG-H, which participate in glyco-

syphosphatidylinositol

[142284]anchor biosynthesis, form a protein complex in the endoplasmic reticulum. J.

[142285]Biol. Chem. 271: 26868–26875, 1996.

[142286]

[142287]

[142288]13324.

[142289]Chernajovsky, Y.; Brown, A.; Clark, J.: Human kinesin light (beta)

[142290]chain gene: DNA sequence and functional characterization of its promoter

[142291]and first exon. DNA Cell Biol. 15: 965–974, 1996.

[142292]

[142293]

[142294]13325.

[142295]Goedert, M.; Marsh, S.; Carter, N.: Localization of the human

[142296]kinesin light chain gene (KNS2) to chromosome 14q32.3 by fluorescence

[142297]in situ hybridization. Genomics 32: 173–175, 1996.

[142298]

[142299]

[142300]13326.

[142301]Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.; Goldstein, L. S.:

[142302]Axonal transport of amyloid precursor protein is mediated
by direct

[142303]binding to the kinesin light chain subunit of kinesin-I.

Neuron 28:

[142304]449–459, 2000.

[142305]

[142306]

[142307]13327.

[142308]Butzow, R.; Huhtala, M.-L.; Bohn, H.; Virtanen, I.; Seppala,
M.

[142309]: Purification and characterization of placental protein 5.
Biochem.

[142310]Biophys. Res. Commun. 150: 483–490, 1988.

[142311]

[142312]

[142313]13328.

[142314]Kisiel, W.; Sprecher, C. A.; Foster, D. C.: Evidence that a
second

[142315]human tissue factor pathway inhibitor (TFPI-2) and human
placental

[142316]protein 5 are equivalent. (Letter) Blood 84: 4384-4385,
1994.

[142317]

[142318]

[142319]13329.

[142320]Miyagi, Y.; Yasumitsu, H.; Eki, T.; Miyata, S.; Koshikawa,
N.;

[142321]Hirahara, F.; Aoki, I.; Misugi, K.; Miyazaki, K.: Assignment
of the

[142322]human PP5/TFPI-2 gene to 7q22 by FISH and PCR-based
human/rodent cell

[142323]hybrid mapping panel analysis. Genomics 35: 267-268,
1996.

[142324]

[142325]

[142326]13330.

[142327]Miyagi, Y.; Yasumitsu, H.; Mizushima, H.; Koshikawa, N.;
Matsuda,

[142328]Y.; Itoh, H.; Hori, T.-A.; Aoki, I.; Misugi, K.; Miyazaki, K.:
Cloning

[142329]of the cDNA encoding mouse PP5/TFPI-2 and mapping of
the gene to chromosome

[142330]6. DNA and Cell Biology 15: 947-954, 1996.

[142331]

[142332]

[142333]13331.

[142334]Siiteri, J. E.; Koistinen, R.; Salem, H. T.; Bohn, H.; Seppala,

[142335]M.: Placental protein 5 is related to blood coagulation and
fibrinolytic

[142336]systems. Life Sci. 30: 1885-1891, 1982.

[142337]

[142338]

[142339]13332.

[142340]Sprecher, C. A.; Kisiel, W.; Mathewes, S.; Foster, D. C.:
Molecular

[142341]cloning, expression, and partial characterization of a sec-

ond human

[142342]tissue-factor-pathway inhibitor. Proc. Nat. Acad. Sci. 91:
3353–3357,

[142343]1994.

[142344]

[142345]

[142346]13333.

[142347]Maglott, D. R.; Durkin, A. S.; Lane, S. A.; Callen, D. F.;
Feldblyum,

[142348]T. V.; Nierman, W. C.: The gene for membrane protein E16
(D16S469E)

[142349]maps to human chromosome 16q24.3 and is expressed in
human brain,

[142350]thymus, and retina. Genomics 23: 303–304, 1994.

[142351]

[142352]

[142353]13334.

[142354]Kastury, K.; Druck, T.; Huebner, K.; Barletta, C.; Acampora,
D.;

[142355]Simeone, A.; Faiella, A.; Boncinelli, E.: Chromosome loca–

tions of

[142356]human EMX and OTX genes. Genomics 22: 41–45, 1994.

[142357]

[142358]

[142359]13335.

[142360]Simeone, A.; Acampora, D.; Gulisano, M.; Stornaiuolo, A.;
Boncinelli,

[142361]E.: Nested expression domains of four homeobox genes in
developing

[142362]rostral brain. Nature 358: 687–690, 1992.

[142363]

[142364]

[142365]13336.

[142366]Bishop, K. M.; Goudreau, G.; O'Leary, D. D. M.: Regulation
of

[142367]area identity in the mammalian neocortex by Emx2 and
Pax6. Science 288:

[142368]344–349, 2000.

[142369]

[142370]

[142371]13337.

[142372]Boncinelli, E.; Gulisano, M.; Spada, F.; Broccoli, V.: Emx
and

[142373]Otx gene expression in the developing mouse brain. Ciba
Found. Symp. 193:

[142374]100–116, 1995.

[142375]

[142376]

[142377]13338.

[142378]Bosetti, A.; Faiella, A.; Boncinelli, E.; Consalez, G. G.: Link-
age

[142379]mapping of Emx2 to mouse chromosome 19. Mammalian
Genome 8: 71–72,

[142380]1997.

[142381]

[142382]

[142383]13339.

[142384]Noonan, F. C.; Mutch, D. G.; Mallon, M. A.; Goodfellow, P.

J.:

[142385]Characterization of the homeodomain gene EMX2: sequence conservation,
[142386]expression analysis, and a search for mutations in endometrial cancers. Genomics 76:
[142387]37–44, 2001.

[142388]

[142389]

[142390]13340.

[142391]Acampora, D.; Mazan, S.; Avantaggiato, V.; Barone, P.;
Tuorto,

[142392]F.; Lallemand, Y.; Brulet, P.; Simeone, A.: Epilepsy and
brain abnormalities

[142393]in mice lacking the Otx1 gene. Nature Genet. 14:
218–222, 1996.

[142394]

[142395]

[142396]13341.

[142397]Boncinelli, E.; Gulisano, M.; Broccoli, V.: Emx and Otx
homeobox

[142398]genes in the developing mouse brain. J. Neurobiol. 24:

1356–1366,

[142399]1993.

[142400]

[142401]

[142402]13342.

[142403]Frantz, G. D.; Weimann, J. M.; Levin, M. E.; McConnell, S.
K.:

[142404]Otx1 and Otx2 define layers and regions in developing
cerebral cortex

[142405]and cerebellum. J. Neurosci. 14: 5725–5740, 1994.

[142406]

[142407]

[142408]13343.

[142409]Avraham, S.; Jiang, S.; Ota, S.; Fu, Y.; Deng, B.; Dowler, L.
L.;

[142410]White, R. A.; Avraham, H.: Structural and functional stud-
ies of the

[142411]intracellular tyrosine kinase MATK gene and its translated
product. J.

[142412]Biol. Chem. 270: 1833–1842, 1995.

[142413]

[142414]

[142415]13344.

[142416]Bennett, B. D.; Cowley, S.; Jiang, S.; London, R.; Deng, B.;
Grabarek,

[142417]J.; Groopman, J. E.; Goeddel, D. V.; Avraham, H.: Identifi-
cation

[142418]and characterization of a novel tyrosine kinase from
megakaryocytes. J.

[142419]Biol. Chem. 269: 1068–1074, 1994.

[142420]

[142421]

[142422]13345.

[142423]Klages, S.; Adam, D.; Class, K.; Fargnoli, J.; Bolen, J. B.;
Penhallow,

[142424]R. C.: Ctk: a protein–tyrosine kinase related to Csk that
defines

[142425]an enzyme family. Proc. Nat. Acad. Sci. 91: 2597–2601,
1994.

[142426]

[142427]

[142428]13346.

[142429]Sakano, S.; Iwama, A.; Inazawa, J.; Ariyama, T.; Ohno, M.;
Suda,

[142430]T.: Molecular cloning of a novel non-receptor tyrosine ki-
nase, HYL

[142431](hematopoietic consensus tyrosine-lacking kinase). Onco-
gene 9: 1155-1161,

[142432]1994.

[142433]

[142434]

[142435]13347.

[142436]Zrihan-Licht, S.; Lim, J.; Keydar, I.; Sliwkowski, M. X.;
Groopman,

[142437]J. E.; Avraham, H.: Association of Csk-homologous kinase
(CHK) (formerly

[142438]MATK) with HER-2/ErbB-2 in breast cancer cells. J. Biol.
Chem. 272:

[142439]1856-1863, 1997.

[142440]

[142441]

[142442]13348.

[142443]Shimizu, S.; Narita, M.; Tsujimoto, Y.: Bcl-2 family proteins

[142444]regulate the release of apoptogenic cytochrome c by the mitochondrial

[142445]channel VDAC. Nature 399: 483-487, 1999.

[142446]

[142447]

[142448]13349.

[142449]Nichols, A. F.: Personal Communication. Berkeley, Calif.
10/4/1995.

[142450]

[142451]

[142452]13350.

[142453]Campbell, H. D.; Schimansky, T.; Claudianos, C.; Ozsarac, N.; Kasprzak,

[142454]A. B.; Cotsell, J. N.; Young, I. G.; de Couet, H. G.; Gabor Miklos,

[142455]G. L.: The Drosophila melanogaster flightless-I gene in-

volved in

[142456]gastrulation and muscle degeneration encodes gelsolin-like and leucine-rich

[142457]repeat domains and is conserved in *Caenorhabditis elegans* and humans. *Proc.*

[142458]*Nat. Acad. Sci.* 90: 11386–11390, 1993.

[142459]

[142460]

[142461]13351.

[142462]Harmon, D. L.; Gardner-Medwin, D.; Stirling, J. L.: Two new mutations

[142463]in a late infantile Tay–Sachs patient are both in exon 1 of the beta-hexosaminidase

[142464]alpha subunit gene. *J. Med. Genet.* 30: 123–128, 1993.

[142465]

[142466]

[142467]13352.

[142468]Buckle, V.; Mondello, C.; Darling, S.; Craig, I. W.; Goodfellow,
low,

[142469]P. N.: Homologous expressed genes in the human sex

chromosome pairing

[142470]region. Nature 317: 739–741, 1985.

[142471]

[142472]

[142473]13353.

[142474]Slaugenhaupt, S. A.; Roca, A. L.; Liebert, C. B.; Altherr, M.
R.;

[142475]Gusella, J. F.; Reppert, S. M.: Mapping of the gene for the
Mel1a-melatonin

[142476]receptor to human chromosome 4 (MTNR1A) and mouse
chromosome 8 (Mtnr1a). Genomics 27:

[142477]355–357, 1995.

[142478]

[142479]

[142480]13354.

[142481]Hattori, Y.; Yamashiro, Y.; Ohba, Y.; Miyaji, T.; Morishita,
M.;

[142482]Yamamoto, K.; Yamamoto, K.; Narai, S.; Kimura, A.: A new
beta-thalassemia

[142483]mutation (initiation codon ATG-to-GTG) found in the

Japanese population. Hemoglobin 15:

[142484]317–325, 1991.

[142485]

[142486]

[142487]13355.

[142488]Holmes, L. B.: Norrie's disease--an X-linked syndrome of
retinal

[142489]malformation, mental retardation and deafness. New Eng.
J. Med. 284:

[142490]367–368, 1971.

[142491]

[142492]

[142493]13356.

[142494]Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Doi, N.;
Nakagawa,

[142495]M.; Ozawa, M.; Kuroda, N.: Novel mutation at the initiation
codon

[142496]in the Norrie disease gene in two Japanese families. Hum.
Genet. 95:

[142497]105–108, 1995.

[142498]

[142499]

[142500]13357.

[142501]Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Hotta, Y.;
Hayakawa,

[142502]M.; Fujiki, K.; Tanabe, U.: Mutations in the Norrie disease
gene:

[142503]a new mutation in a Japanese family. (Letter) Brit. J. Oph-
thal. 79:

[142504]703–708, 1995.

[142505]

[142506]

[142507]13358.

[142508]Johnson, K.; Mintz–Hittner, H. A.; Conley, Y. P.; Ferrell, R.

[142509]E.: X–linked exudative vitreoretinopathy caused by an
arginine to

[142510]leucine substitution (R121L) in the Norrie disease protein.
Clin.

[142511]Genet. 50: 113–115, 1996.

[142512]

[142513]

[142514]13359.

[142515]Johnston, S. S.; Hanna, J. E.; Nevin, N. C.; Bryars, J. H.:
Norrie's

[142516]disease. Birth Defects Orig. Art. Ser. 18(6): 729–738,
1982.

[142517]

[142518]

[142519]13360.

[142520]Katayama, S.; Wohlferd, M.; Golbus, M. S.: First demon-
stration

[142521]of recombination between the gene for Norrie disease and
probe L1.28. Am.

[142522]J. Med. Genet. 30: 967–970, 1988.

[142523]

[142524]

[142525]13361.

[142526]Kivlin, J. D.; Sanborn, G. E.; Wright, E.; Cannon, L.; Carey,

[142527]J.: Further linkage data on Norrie disease. Am. J. Med.

Genet. 26:

[142528]733–736, 1987.

[142529]

[142530]

[142531]13362.

[142532]Lindsay, S.; Thiselton, D. L.; Bateman, J. B.; Ngo, J. T.;
Sparkes,

[142533]R. S.; Coleman, M.; Davies, K. E.; Bhattacharya, S. S.: Lo-
calisation

[142534]of the gene for Norrie disease to between DXS7 and
DXS426 on Xp. Hum.

[142535]Genet. 88: 349–350, 1992.

[142536]

[142537]

[142538]13363.

[142539]Meindl, A.; Berger, W.; Meitinger, T.; van de Pol, D.;
Achatz,

[142540]H.; Dorner, C.; Haasemann, M.; Hellebrand, H.; Gal, A.;
Cremers, F.;

[142541]Ropers, H.–H.: Norrie disease is caused by mutations in an

extracellular

[142542]protein resembling C-terminal globular domain of
mucins. Nature Genet. 2:

[142543]139–143, 1992.

[142544]

[142545]

[142546]13364.

[142547]Meindl, A.; Lorenz, B.; Achatz, H.; Hellebrand, H.;
Schmitz-Valckenberg,

[142548]P.; Meitinger, T.: Missense mutations in the NDP gene in
patients

[142549]with a less severe course of Norrie disease. Hum. Molec.
Genet. 4:

[142550]489–490, 1995.

[142551]

[142552]

[142553]13365.

[142554]Meitinger, T.; Meindl, A.; Bork, P.; Rost, B.; Sander, C.;
Haasemann,

[142555]M.; Murken, J.: Molecular modeling of the Norrie disease

protein

[142556]predicts a cystine knot growth factor tertiary structure.

Nature

[142557]Genet. 5: 376–380, 1993.

[142558]

[142559]

[142560]13366.

[142561]Moreira-Filho, C. A.; Neustein, I.: A presumptive new variant

[142562]of Norrie's disease. J. Med. Genet. 16: 125–128, 1979.

[142563]

[142564]

[142565]13367.

[142566]Nance, W. E.; Hara, S.; Hansen, A.; Elliott, J.; Lewis, M.; Chown,

[142567]B.: Genetic linkage studies in a Negro kindred with Norrie's disease. Am.

[142568]J. Hum. Genet. 21: 423–429, 1969.

[142569]

[142570]

[142571]13368.

[142572]Ngo, J.; Spence, M. A.; Cortessis, V.; Bateman, J. B.;
Sparkes,

[142573]R. S.: Duplicate report crossing over in Norrie disease
family. (Letter) Am.

[142574]J. Med. Genet. 33: 286, 1989.

[142575]

[142576]

[142577]13369.

[142578]Ngo, J. T.; Bateman, J. B.; Cortessis, V.; Sparkes, R. S.; Mo-
handas,

[142579]T.; Inana, G.; Spence, M. A.: Norrie disease: linkage analy-
sis using

[142580]a 4.2-kb RFLP detected by a human ornithine aminotrans-
ferase cDNA

[142581]probe. Genomics 4: 539-545, 1989.

[142582]

[142583]

[142584]13370.

[142585]Ngo, J. T.; Spence, M. A.; Cortessis, V.; Sparkes, R. S.;
Bateman,

[142586]J. B.: Recombinational event between Norrie disease and
DXS7 loci. Clin.

[142587]Genet. 34: 43–47, 1988.

[142588]

[142589]

[142590]13371.

[142591]Norrie, G.: Nogle Blindhedsaarsager: en oversigt. Hospi-
talstidende 76:

[142592]141–147, 1933.

[142593]

[142594]

[142595]13372.

[142596]Ohba, N.; Yamashita, T.: Primary vitreoretinal dysplasia
resembling

[142597]Norrie's disease in a female: association with X autosome
chromosomal

[142598]translocation. Brit. J. Ophthal. 70: 64–71, 1986.

[142599]

[142600]

[142601]13373.

[142602]Phillips, C. I.; Newton, M.; Duvall, J.; Holloway, S.; Levy, A.

[142603]M.: Probably Norrie's disease due to mutation: two sporadic sibships

[142604]of two males each, a necropsy of one case, and, given Norrie's disease,

[142605]a calculation of the gene mutation frequency. Brit. J. Ophthalmol. 70:

[142606]305-313, 1986.

[142607]

[142608]

[142609]13374.

[142610]Rehm, H. L.; Gutierrez-Espeleta, G. A.; Garcia, R.; Jimenez, G.;

[142611]Khetarpal, U.; Priest, J. M.; Sims, K. B.; Keats, B. J. B.; Morton,

[142612]C. C.: Norrie disease gene mutation in a large Costa Rican kindred

[142613]with a novel phenotype including venous insufficiency. Hum. Mutat.

[142614]9: 402–408, 1997.

[142615]

[142616]

[142617]13375.

[142618]Schuback, D. E.; Chen, Z. Y.; Craig, I. W.; Breakefield, X.
O.;

[142619]Sims, K. B.: Mutations in the Norrie disease gene. Hum.
Mutat. 5:

[142620]285–292, 1995.

[142621]

[142622]

[142623]13376.

[142624]Shastry, B. S.: Identification of a recurrent missense muta-
tion

[142625]in the Norrie disease gene associated with a simplex case
of exudative

[142626]vitreoretinopathy. Biochem. Biophys. Res. Commun. 246:
35–38, 1998.

[142627]

[142628]

[142629]13377.

[142630]Shastry, B. S.; Hejtmancik, J. F.; Plager, D. A.; Hartzler, M.

[142631]K.; Trese, M. T.: Linkage and candidate gene analysis in X-linked

[142632]familial exudative vitreoretinopathy. Genomics 27: 341-344, 1995.

[142633]

[142634]

[142635]13378.

[142636]Shastry, B. S.; Hejtmancik, J. F.; Trese, M. T.: Identification

[142637]of novel missense mutations in the Norrie disease gene associated

[142638]with one X-linked and four sporadic cases of familial exudative vitreoretinopathy. Hum.

[142639]Mutat. 9: 396-401, 1997.

[142640]

[142641]

[142642]13379.

[142643]Sims, K. B.; Lebo, R. V.; Benson, G.; Shalish, C.; Schuback,

D.;

[142644]Chen, Z. Y.; Bruns, G.; Craig, I. W.; Golbus, M. S.; Breakefield,

[142645]X. O.: The Norrie disease gene maps to a 150 kb region on chromosome

[142646]Xp11.3. Hum. Molec. Genet. 1: 83–89, 1992.

[142647]

[142648]

[142649]13380.

[142650]Sims, K. B.; Ozelius, L.; Corey, T.; Rinehart, W. B.; Liberafarb,

[142651]R.; Haines, J.; Chen, W. J.; Norio, R.; Sankila, E.; de la Chapelle,

[142652]A.; Murphy, D. L.; Gusella, J.; Breakefield, X. O.: Norrie disease

[142653]gene is distinct from the monoamine oxidase genes. Am. J. Hum. Genet. 45:

[142654]424–434, 1989.

[142655]

[142656]

[142657]13381.

[142658]Taylor, P. J.; Coates, T.; Newhouse, M. L.: Episkopi blindness:
ness:

[142659]hereditary blindness in a Greek Cypriot family. Brit. J.
Ophthalm. 43:

[142660]340–344, 1959.

[142661]

[142662]

[142663]13382.

[142664]Torrente, I.; Mangino, M.; Gennarelli, M.; Novelli, G.; Gian-
notti,

[142665]A.; Vadala, P.; Dallapiccola, B.: Two new missense muta-
tions (A105T

[142666]and C110G) in the norrin gene in two Italian families with
Norrie

[142667]disease and familial exudative vitreoretinopathy. (Letter)
Am. J.

[142668]Med. Genet. 72: 242–244, 1997.

[142669]

[142670]

[142671]13383.

[142672]Burgoyne, P. S.: Genetic homology and crossing over in
the X and

[142673]Y chromosomes of mammals. Hum. Genet. 61: 85–90,
1982.

[142674]

[142675]

[142676]13384.

[142677]Cooke, H. J.; Brown, W. R. A.; Rappold, G. A.: Hypervari-
able telomeric

[142678]sequences from the human sex chromosomes are pseu-
doautosomal. Nature 317:

[142679]687–692, 1985.

[142680]

[142681]

[142682]13385.

[142683]Darling, S. M.; Banting, G. S.; Pym, B.; Wolfe, J.; Goodfellow,
low,

[142684]P. N.: Cloning an expressed gene shared by the human
sex chromosomes. Proc.

[142685]Nat. Acad. Sci. 83: 135–139, 1986.

[142686]

[142687]

[142688]13386.

[142689]Dracopoli, N. C.; Rettig, W. J.; Albino, A. P.; Esposito, D.;
Archidiacono,

[142690]N.; Rocchi, M.; Siniscalco, M.; Old, L. J.: Genes controlling
gp25/30

[142691]cell–surface molecules map to chromosomes X and Y and
escape X–inactivation. Am.

[142692]J. Hum. Genet. 37: 199–207, 1985.

[142693]

[142694]

[142695]13387.

[142696]Gelin, C.; Aubrit, F.; Phalipon, A.; Raynal, B.; Cole, S.; Kaczorek,

[142697]M.; Bernard, A.: The E2 antigen, a 32 kD glycoprotein involved in

[142698]T-cell adhesion processes, is the MIC2 gene product.

EMBO J. 8:

[142699]3253–3259, 1989.

[142700]

[142701]

[142702]13388.

[142703]Geller, R. L.; Shapiro, L. J.; Mohandas, T. K.: Fine mapping of

[142704]the distal short arm of the human X chromosome using X/Y translocations. Am.

[142705]J. Hum. Genet. 38: 884–890, 1986.

[142706]

[142707]

[142708]13389.

[142709]Goodfellow, P.; Banting, G.; Sheer, D.; Ropers, H. H.;
Caine,

[142710]A.; Ferguson-Smith, M. A.; Povey, S.; Voss, R.: Genetic evidence

[142711]that a Y-linked gene in man is homologous to a gene on
the X chromosome. Nature 302:

[142712]346–349, 1983.

[142713]

[142714]

[142715]13390.

[142716]Goodfellow, P.; Pym, B.; Mohandas, T.; Shapiro, L. J.: The
cell

[142717]surface antigen locus, MIC2X, escapes X-inactivation. Am.
J. Hum.

[142718]Genet. 36: 777–782, 1984.

[142719]

[142720]

[142721]13391.

[142722]Goodfellow, P. J.; Darling, S. M.; Thomas, N. S.; Goodfellow,
low,

[142723]P. N.: A pseudoautosomal gene in man. Science 234:
740–743, 1986.

[142724]

[142725]

[142726]13392.

[142727]Goodfellow, P. N.; Tippet, P.: A human quantitative poly-
morphism

[142728]related to Xg blood groups. Nature 289: 404–405, 1981.

[142729]

[142730]

[142731]13393.

[142732]Levy, R.; Dilley, J.; Fox, R. I.; Warnke, R.: A human thymus-
leukemia

[142733]antigen defined by hybridoma monoclonal antibodies.
Proc. Nat. Acad.

[142734]Sci. 76: 6552–6556, 1979.

[142735]

[142736]

[142737]13394.

[142738]Pettersen, R. D.; Bernard, G.; Olafsen, M. K.; Pourtein, M.;
Lie,

[142739]S. O.: CD99 signals caspase-independent T cell death. J.
Immun. 166:

[142740]4931-4942, 2001.

[142741]

[142742]

[142743]13395.

[142744]Ropers, H. H.; Zimmer, J.; Strobl, G.; Goodfellow, P.: The
MIC2X

[142745](12E7) locus maps distally from STS on Xp. (Abstract) Cy-
togenet.

[142746]Cell Genet. 40: 736 only, 1985.

[142747]

[142748]

[142749]13396.

[142750]Simmler, M.-C.; Rouyer, F.; Vergnaud, G.; Nystrom-Lahti,
M.; Ngo,

[142751]K. Y.; de la Chapelle, A.; Weissenbach, J.: Pseudoautosomal
DNA sequences

[142752]in the pairing region of the human sex chromosomes. Nature 317:

[142753]692–697, 1985.

[142754]

[142755]

[142756]13397.

[142757]Tippett, P.; Shaw, M.–A.; Green, C. A.; Daniels, G. L.: The 12E7

[142758]red cell quantitative polymorphism: control by the Y–borne locus,

[142759]Yg. Ann. Hum. Genet. 50: 339–347, 1986.

[142760]

[142761]

[142762]13398.

[142763]Janz, R.; Sudhof, T. C.; Hammer, R. E.; Unni, V.; Siegelbaum, S.

[142764]A.; Bolshakov, V. Y.: Essential roles in synaptic plasticity for

[142765]synaptogyrin I and synaptophysin I. Neuron 24: 687–700, 1999.

[142766]

[142767]

[142768]13399.

[142769]McMahon, H. T.; Bolshakov, V. Y.; Janz, R.; Hammer, R. E.;
Siegelbaum,

[142770]S. A.; Sudhof, T. C.: Synaptophysin, a major synaptic vesicle
protein,

[142771]is not essential for neurotransmitter release. Proc. Nat.
Acad. Sci. 93:

[142772]4760–4764, 1996.

[142773]

[142774]

[142775]13400.

[142776]Brown, C. J.; Sekiguchi, T.; Nishimoto, T.; Willard, H. F.:
Regional

[142777]localization of CCG1 gene which complements hamster
cell cycle mutation

[142778]BN462 to Xq11–Xq13. Somat. Cell Molec. Genet. 15:
93–96, 1989.

[142779]

[142780]

[142781]13401.

[142782]Derry, J. M. J.; Barnard, P. J.: Localization of the Ccg1 gene

[142783]on the mouse X chromosome. (Abstract) Cytogenet. Cell
Genet. 51:

[142784]988, 1989.

[142785]

[142786]

[142787]13402.

[142788]Dikstein, R.; Ruppert, S.; Tjian, R.: TAFII250 is a bipartite

[142789]protein kinase that phosphorylates the base transcription
factor RAP74. Cell 84:

[142790]781–790, 1996.

[142791]

[142792]

[142793]13403.

[142794]Giles, R. E.; Ruddle, F. H.: X-linkage of a human genetic
locus

[142795]that corrects the DNA synthesis lesion in tsC1AGOH

mouse cells. Genetics 93:

[142796]975–996, 1979.

[142797]

[142798]

[142799]13404.

[142800]Hisatake, K.; Hasegawa, S.; Takada, R.; Nakatani, Y.;
Horikoshi,

[142801]M.; Roeder, R. G.: The p250 subunit of native TATA box–
binding factor

[142802]TFIID is the cell–cycle regulatory protein CCG1. Nature
362: 179–181,

[142803]1993.

[142804]

[142805]

[142806]13405.

[142807]Jacobson, R. H.; Ladurner, A. G.; King, D. S.; Tjian, R.:
Structure

[142808]and function of a human TAFII250 double bromodomain
module. Science 288:

[142809]1422–1425, 2000.

[142810]

[142811]

[142812]13406.

[142813]Jha, K. K.; Ozer, H. L.: Genetic studies with a mutant mouse cell,

[142814]ts-2 Balb-3T3, with a temperature-sensitive defect in DNA synthesis.

[142815](Abstract) Genetics 86: s32-s33, 1977.

[142816]

[142817]

[142818]13407.

[142819]Jha, K. K.; Siniscalco, M.; Ozer, H. L.: Temperature-sensitive

[142820]mutants of Balb-3T3 cells. III. Hybrids between ts2 and other mouse

[142821]mutant cells affected in DNA synthesis and correction of ts2 defect

[142822]by human X chromosome. Somat. Cell Genet. 6: 603-614, 1980.

[142823]

[142824]

[142825]13408.

[142826]Almind, K.; Ahlgren, M. G.; Hansen, T.; Urhammer, S. A.;
Clausen,

[142827]J. O.; Pedersen, O.: Discovery of a Met300Val variant in
Shc and

[142828]studies of its relationship to birth weight and length, im-
paired insulin

[142829]secretion, insulin resistance, and type 2 diabetes mellitus.
J. Clin.

[142830]Endocr. Metab. 84: 2241–2244, 1999.

[142831]

[142832]

[142833]13409.

[142834]Harun, R. B.; Smith, K. K.; Leek, J. P.; Markham, A. F.; Nor-
ris,

[142835]A.; Morrison, J. F. J.: Characterization of human SHC p66
cDNA and

[142836]its processed pseudogene mapping to Xq12–q13.1. Ge-
nomics 42: 349–352,

[142837]1997.

[142838]

[142839]

[142840]13410.

[142841]McGlade, J.; Cheng, A.; Pelicci, G.; Pelicci, P. G.; Pawson, T.

[142842]: Shc proteins are phosphorylated and regulated by the v-src and v-fps

[142843]protein-tyrosine-kinases. Proc. Nat. Acad. Sci. 89:
8869-8873, 1992.

[142844]

[142845]

[142846]13411.

[142847]Migliaccio, E.; Giorgio, M.; Mele, S.; Pelicci, G.; Reboldi, P.;

[142848]Pandolfi, P. P.; Lanfranccone, L.; Pelicci, P. G.: The p66(shc)
adaptor

[142849]protein controls oxidative stress response and life span in
mammals. Nature 402:

[142850]309-313, 1999.

[142851]

[142852]

[142853]13412.

[142854]Nemoto, S.; Finkel, T.: Redox regulation of forkhead proteins

[142855]through a p66shc-dependent signaling pathway. Science
295: 2450–2452,

[142856]2002.

[142857]

[142858]

[142859]13413.

[142860]Pelicci, G.; Lanfrancone, L.; Grignani, F.; McGlade, J.; Cavallo,

[142861]F.; Forni, G.; Nicoletti, I.; Grignani, F.; Pawson, T.; Pelicci, P.

[142862]G.: A novel transforming protein (SHC) with an SH2 domain is implicated

[142863]in mitogenic signal transduction. Cell 70: 93–104, 1992.

[142864]

[142865]

[142866]13414.

[142867]Yulug, I. G.; Egan, S. E.; See, C. G.; Fisher, E. M. C.: A hu-

man

[142868]SHC-related sequence maps to chromosome 17, the SHC
gene maps to chromosome

[142869]1. Hum. Genet. 96: 245–248, 1995.

[142870]

[142871]

[142872]13415.

[142873]Zhang, L.; Camerini, V.; Bender, T. P.; Ravichandran, K. S.:
A

[142874]nonredundant role for the adapter protein Shc in thymic T
cell development. Nature

[142875]Immun. 3: 749–755, 2002.

[142876]

[142877]

[142878]13416.

[142879]Abramovitz, M.; Boie, Y.; Nguyen, T.; Rushmore, T. H.;
Bayne, M.

[142880]A.; Metters, K. M.; Slipetz, D. M.; Grygorczyk, R.: Cloning
and expression

[142881]of a cDNA for the human prostanoid FP receptor. J. Biol.

Chem. 269:

[142882]2632–2636, 1994.

[142883]

[142884]

[142885]13417.

[142886]Betz, R.; Lagercrantz, J.; Kedra, D.; Dumanski, J. P.; Nordenskjold,

[142887]A.: Genomic structure, 5–prime flanking sequences, and precise localization

[142888]in 1p31.1 of the human prostaglandin F receptor gene.

Biochem. Biophys.

[142889]Res. Commun. 254: 413–416, 1999.

[142890]

[142891]

[142892]13418.

[142893]Sugimoto, Y.; Yamasaki, A.; Segi, E.; Tsuboi, K.; Aze, Y.; Nishimura,

[142894]T.; Oida, H.; Yoshida, N.; Tanaka, T.; Katsuyama, M.; Hashimoto, K.;

[142895]Murata, T.; Hirata, M.; Ushikubi, F.; Negishi, M.; Ichikawa,

A.; Narumiya,

[142896]S.: Failure of parturition in mice lacking the prostaglandin
F receptor. Science 277:

[142897]681–683, 1997.

[142898]

[142899]

[142900]13419.

[142901]Geppert, M.; Khvotchev, M.; Krasnoperov, V.; Goda, Y.;
Missler,

[142902]M.; Hammer, R. E.; Ichtchenko, K.; Petrenko, A. G.; Sudhof,
T. C.

[142903]: Neurexin I–alpha is a major alpha–latrotoxin receptor
that cooperates

[142904]in alpha–latrotoxin action. J. Biol. Chem. 273: 1705–1710,
1998.

[142905]

[142906]

[142907]13420.

[142908]Ichtchenko, K.; Hata, Y.; Nguyen, T.; Ullrich, B.; Missler,
M.;

[142909]Moomaw, C.; Sudhof, T. C.: Neuroligin 1: a splice site–

specific ligand

[142910]for beta-neurexins. Cell 81: 435–443, 1995.

[142911]

[142912]

[142913]13421.

[142914]Kleiderlein, J. J.; Nisson, P. E.; Jessee, J.; Li, W.-B.; Becker,

[142915]K. G.; Derby, M. L.; Ross, C. A.; Margolis, R. L.: CCG re-
peats in

[142916]cDNAs from human brain. Hum. Genet. 103: 666–673,
1998. Note: Erratum:

[142917]Hum. Genet. 104: 113 only, 1999.

[142918]

[142919]

[142920]13422.

[142921]Apte, S. S.; Mattei, M.-G.; Olsen, B. R.: Mapping of human
BAX

[142922]gene to chromosome 19q13.3–q13.4 and isolation of a
novel alternatively

[142923]spliced transcript, BAX–delta. Genomics 26: 592–594,
1995.

[142924]

[142925]

[142926]13423.

[142927]Margolis, R.: Personal Communication. Baltimore, Md.
3/29/2000.

[142928]

[142929]

[142930]13424.

[142931]Missler, M.; Sudhof, T. C.: Neurexins: three genes and
1001 products. Trends

[142932]Genet. 14: 20–26, 1998.

[142933]

[142934]

[142935]13425.

[142936]Ikeshima, H.; Imai, S.; Shimoda, K.; Hata, J.; Takano, T.:
Expression

[142937]of a MADS box gene, MEF2D, in neurons of the mouse
central nervous

[142938]system: implication of its binary function in myogenic and

neurogenic

[142939]cell lineages. Neurosci. Lett. 200: 117–120, 1995.

[142940]

[142941]

[142942]13426.

[142943]Delhase, M.; Hayakawa, M.; Chen, Y.; Karin, M.: Positive
and negative

[142944]regulation of I-kappa-B kinase activity through IKK-beta
subunit phosphorylation. Science 284:

[142945]309–312, 1999.

[142946]

[142947]

[142948]13427.

[142949]DiDonato, J. A.; Hayakawa, M.; Rothwarf, D. M.; Zandi, E.;
Karin,

[142950]M.: A cytokine-responsive IkappaB kinase that activates
the transcription

[142951]factor NF-kappaB. Nature 388: 548–554, 1997.

[142952]

[142953]

[142954]13428.

[142955]Mercurio, F.; Zhu, H.; Murray, B. W.; Shevchenko, A.; Bennett,

[142956]B. L.; Li, J.; Young, D. B.; Barbosa, M.; Mann, M.; Manning, A.; Rao,

[142957]A.: IKK-1 and IKK-2: cytokine-activated I-kappa-B kinases essential

[142958]for NF-kappa-B activation. Science 278: 860-866, 1997.

[142959]

[142960]

[142961]13429.

[142962]Matsushima, A.; Kaisho, T.; Rennert, P. D.; Nakano, H.; Kurosawa,

[142963]K.; Uchida, D.; Takeda, K.; Akira, S.; Matsumoto, M.: Essential role

[142964]of nuclear factor (NF)-kappa-B-inducing kinase and inhibitor of kappa-B

[142965](I-kappa-B) kinase alpha in NF-kappa-B activation through lymphotoxin

[142966]beta receptor, but not through tumor necrosis factor re-

ceptor I. J.

[142967]Exp. Med. 193: 631–636, 2001.

[142968]

[142969]

[142970]13430.

[142971]Brzezinski, A.: Melatonin in humans. New Eng. J. Med.
336: 186–195,

[142972]1997.

[142973]

[142974]

[142975]13431.

[142976]Liu, R.–Y.; Zhou, J.–N.; van Heerikhuize, J.; Hofman, M. A.;
Swaab,

[142977]D. F.: Decreased melatonin levels in postmortem cere-
brospinal fluid

[142978]in relation to aging, Alzheimer's disease, and apolipopro-
tein E-epsilon-4/4

[142979]genotype. J. Clin. Endocr. Metab. 84: 323–327, 1999.

[142980]

[142981]

[142982]13432.

[142983]Nelson, C. S.; Ikeda, M.; Gompf, H. S.; Robinson, M. L.;
Fuchs,

[142984]N. K.; Yoshioka, T.; Neve, K. A.; Allen, C. N.: Regulation of
melatonin

[142985]1a receptor signaling and trafficking by asparagine-124.
Molec. Endocr. 15:

[142986]1306–1317, 2001.

[142987]

[142988]

[142989]13433.

[142990]Reppert, S. M.; Weaver, D. R.: Melatonin madness. Cell 83:
1059–1062,

[142991]1995.

[142992]

[142993]

[142994]13434.

[142995]Reppert, S. M.; Weaver, D. R.; Ebisawa, T.: Cloning and

characterization

[142996]of a mammalian melatonin receptor that mediates reproductive and circadian

[142997]responses. Neuron 13: 1177–1185, 1994.

[142998]

[142999]

[143000]13435.

[143001]Muller–Pillasch, F.; Zimmerhackl, F.; Lacher, U.; Schultz, N.;

[143002]Hameister, H.; Varga, G.; Friess, H.; Buchler, M.; Adler, G.; Gress,

[143003]T. M.: Cloning of novel transcripts of the human guanine–nucleotide–exchange

[143004]factor Mss4: in situ chromosomal mapping and expression in pancreatic

[143005]cancer. Genomics 46: 389–396, 1997.

[143006]

[143007]

[143008]13436.

[143009]Yu, H.; Schreiber, S. L.: Cloning, Zn(2+) binding, and

structural

[143010]characterization of the guanine nucleotide exchange factor human Mss4. Biochemistry 34:

[143011]9103–9110, 1995.

[143012]

[143013]

[143014]13437.

[143015]Yabe, D.; Nakamura, T.; Kanazawa, N.; Tashiro, K.; Honjo, T.:

[143016]Calumenin, a $\text{Ca}(2+)$ –binding protein retained in the endoplasmic reticulum

[143017]with a novel carboxyl–terminal sequence, HDEF. J. Biol. Chem. 272:

[143018]18232–18239, 1997.

[143019]

[143020]

[143021]13438.

[143022]Yabe, D.; Taniwaki, M.; Nakamura, T.; Kanazawa, N.; Tashiro, K.;

[143023]Honjo, T.: Human calumenin gene (CALU): cDNA isolation

and chromosomal

[143024]mapping to 7q32. Genomics 49: 331–333, 1998.

[143025]

[143026]

[143027]13439.

[143028]Angelin–Duclos, C.; Cattoretti, G.; Lin, K.–I.; Calame, K.:
Commitment

[143029]of B lymphocytes to a plasma cell fate is associated with
Blimp–1

[143030]expression in vivo. J. Immun. 165: 5462–5471, 2000.

[143031]

[143032]

[143033]13440.

[143034]Chang, D. H.; Angelin–Duclos, C.; Calame, K.: BLIMP–1:
trigger

[143035]for differentiation of myeloid lineage. Nature Immun. 1:
169–176,

[143036]2000.

[143037]

[143038]

[143039]13441.

[143040]Huang, S.: Blimp-1 is the murine homolog of the human transcriptional

[143041]repressor PRDI-BF1. (Letter) Cell 78: 9 only, 1994.

[143042]

[143043]

[143044]13442.

[143045]Keller, A. D.; Maniatis, T.: Identification and characterization

[143046]of a novel repressor of beta-interferon gene expression. Genes Dev. 5:

[143047]868-879, 1991.

[143048]

[143049]

[143050]13443.

[143051]Mock, B. A.; Liu, L.; Le Paslier, D.; Huang, S.: The B-lymphocyte

[143052]maturation promoting transcription factor

BLIMP1/PRDI-BF1 maps to

[143053]D6S447 on human chromosome 6q21-q22.1 and the syntenic region of mouse

[143054]chromosome 10. Genomics 37: 24-28, 1996.

[143055]

[143056]

[143057]13444.

[143058]Ito, T.; Yang, M.; May, W. S.: RAX, a cellular activator for double-stranded

[143059]RNA-dependent protein kinase during stress signaling. J. Biol. Chem. 274:

[143060]15427-15432, 1999.

[143061]

[143062]

[143063]13445.

[143064]Patel, R. C.; Sen, G. C.: PACT, a protein activator of the interferon-induced

[143065]protein kinase, PKR. EMBO J. 17: 4379-4390, 1998.

[143066]

[143067]

[143068]13446.

[143069]Scott, A. F.: Personal Communication. Baltimore, Md.
2001.

[143070]

[143071]

[143072]13447.

[143073]Shearman, L. P.; Zylka, M. J.; Weaver, D. R.; Kolakowski, L.
F.,

[143074]Jr.; Reppert, S. M.: Two period homologs: circadian ex-
pression and

[143075]photic regulation in the suprachiasmatic nuclei. Neuron
19: 1261–1269,

[143076]1997.

[143077]

[143078]

[143079]13448.

[143080]Toh, K. L.; Jones, C. R.; He, Y.; Eide, E. J.; Hinz, W. A.; Vir-
shup,

[143081]D. M.; Ptacek, L. J.; Fu, Y.-H.: An hPer2 phosphorylation

site mutation

[143082]in familial advanced sleep phase syndrome. Science 291:
1040–1043,
[143083]2001.

[143084]

[143085]

[143086]13449.

[143087]Engelender, S.; Wanner, T.; Kleiderlein, J. J.; Wakabayashi,
K.;

[143088]Tsuji, S.; Takahashi, H.; Ashworth, R.; Margolis, R. L.;
Ross, C.

[143089]A.: Organization of the human synphilin–1 gene, a candi-
date for Parkinson's

[143090]disease. Mammalian Genome 11: 763–766, 2000.

[143091]

[143092]

[143093]13450.

[143094]Chien, W.; Pei, L.: A novel binding factor facilitates nuclear

[143095]translocation and transcriptional activation function of the
pituitary

[143096]tumor-transforming gene product. J. Biol. Chem. 275:
19422–19427,
[143097]2000.

[143098]

[143099]

[143100]13451.

[143101]Yaspo, M.–L.; Aaltonen, J.; Horelli–Kuitunen, N.; Peltonen,
L.;

[143102]Lehrach, H.: Cloning of a novel human putative type Ia in-
tegral membrane

[143103]protein mapping to 21q22.3. Genomics 49: 133–136,
1998.

[143104]

[143105]

[143106]13452.

[143107]Yaspo, M.–L.; Gellen, L.; Mott, R.; Korn, B.; Nizetic, D.;
Poustka,

[143108]A. M.; Lehrach, H.: Model for a transcript map of human
chromosome

[143109]21: isolation of new coding sequences from exon and en-
riched cDNA

[143110]libraries. Hum. Molec. Genet. 4: 1291–1304, 1995.

[143111]

[143112]

[143113]13453.

[143114]Ghiso, J. A.; Holton, J.; Miravalle, L.; Calero, M.; Lashley, T.;

[143115]Vidal, R.; Houlden, H.; Wood, N.; Neubert, T. A.; Rostagno, A.; Plant,

[143116]G.; Revesz, T.; Frangione, B.: Systemic amyloid deposits in familial

[143117]British dementia. J. Biol. Chem. 276: 43909–43914, 2001.

[143118]

[143119]

[143120]13454.

[143121]Stromgren, E.; Dalby, A.; Dalby, M.; Ranheim, B. :Acta Neurol.

[143122]Scand. 46 (suppl. 43): 97–98, 1970.

[143123]

[143124]

[143125]13455.

[143126]Gandhi, R.; Elble, R. C.; Gruber, A. D.; Schreur, K. D.; Ji, H.-L.;

[143127]Fuller, C. M.; Pauli, B. U.: Molecular and functional characterization

[143128]of a calcium-sensitive chloride channel from mouse lung. J. Biol.

[143129]Chem. 273: 32096-32101, 1998.

[143130]

[143131]

[143132]13456.

[143133]Gruber, A. D.; Elble, R. C.; Ji, H.-L.; Schreur, K. D.; Fuller,

[143134]C. M.; Pauli, B. U.: Genomic cloning, molecular characterization,

[143135]and functional analysis of human CLCA1, the first human member of

[143136]the family of Ca^{2+} -activated Cl^- channel proteins. Genomics 54:

[143137]200-214, 1998.

[143138]

[143139]

[143140]13457.

[143141]Gruber, A. D.; Schreur, K. D.; Ji, H.-L.; Fuller, C. M.; Pauli,

[143142]B. U.: Molecular cloning and transmembrane structure of
hCLCA2 from

[143143]human lung, trachea, and mammary gland. Am. J. Physiol.
276: C1261–C1270,

[143144]1999.

[143145]

[143146]

[143147]13458.

[143148]Liao, C.; Wang, X. Y.; Wei, H. Q.; Li, S. Q.; Merghoub, T.;
Pandolfi,

[143149]P. P.; Wolgemuth, D. J.: Altered myelopoiesis and the de-
velopment

[143150]of acute myeloid leukemia in transgenic mice overex-
pressing cyclin

[143151]A1. Proc. Nat. Acad. Sci. 98: 6853–6858, 2001.

[143152]

[143153]

[143154]13459.

[143155]Liu, D.; Matzuk, M. M.; Sung, W. K.; Guo, Q.; Wang, P.;

Wolgemuth,

[143156]D. J.: Cyclin A1 is required for meiosis in the male mouse.

Nature

[143157]Genet. 20: 377–380, 1998.

[143158]

[143159]

[143160]13460.

[143161]Muller, C.; Yang, R.; Beck–von–Peccoz, L.; Idos, G.; Ver–

beek, W.;

[143162]Koeffler, H. P.: Cloning of the cyclin A1 genomic structure

and characterization

[143163]of the promoter region: GC boxes are essential for cell cy–

cle–regulated

[143164]transcription of the cyclin A1 gene. J. Biol. Chem. 274:

11220–11228,

[143165]1999.

[143166]

[143167]

[143168]13461.

[143169]Albertsen, H. M.; Smith, S. A.; Mazoyer, S.; Fujimoto, E.;
Stevens,

[143170]J.; Williams, B.; Rodriguez, P.; Cropp, C. S.; Slijepcevic, P.;
Carlson,

[143171]M.; Robertson, M.; Bradley, P.; and 9 others: A physical
map and

[143172]candidate genes in the BRCA1 region on chromosome
17q12–21. Nature

[143173]Genet. 7: 472–479, 1994.

[143174]

[143175]

[143176]13462.

[143177]Han, H.–J.; Sudo, K.; Inazawa, J.; Nakamura, Y.: Isolation
and

[143178]mapping of a human gene (RABL) encoding a small GTP–
binding protein

[143179]homologous to the Ras–related RAB gene. Cytogenet. Cell
Genet. 73:

[143180]137–139, 1996.

[143181]

[143182]

[143183]13463.

[143184]Csoka, A. B.; Scherer, S. W.; Stern, R.: Expression analysis
of

[143185]six paralogous human hyaluronidase genes clustered on
chromosomes

[143186]3p21 and 7q31. Genomics 60: 356–361, 1999.

[143187]

[143188]

[143189]13464.

[143190]Heckel, D.; Comtesse, N.; Brass, N.; Blin, N.; Zang, K. D.;
Meese,

[143191]E.: Novel immunogenic antigen homologous to
hyaluronidase in meningioma. Hum.

[143192]Molec. Genet. 7: 1859–1872, 1998.

[143193]

[143194]

[143195]13465.

[143196]Dolganov, G. M.; Maser, R. S.; Novikov, A.; Tosto, L.;

Chong, S.;

[143197]Bressan, D. A.; Petrini, J. H. J.: Human Rad50 is physically associated

[143198]with human Mre11: identification of a conserved multi-protein complex

[143199]implicated in recombinational DNA repair. Molec. Cell Biol. 16:

[143200]4832–4841, 1996.

[143201]

[143202]

[143203]13466.

[143204]Hopfner, K.–P.; Craig, L.; Moncalian, G.; Zinkel, R. A.; Usui,

[143205]T.; Owen, B. A. L.; Karcher, A.; Henderson, B.; Bodmer, J.–L.; McMurray,

[143206]C. T.; Carney, J. P.; Petrini, J. H. J.; Tainer, J. A.: The Rad50

[143207]zinc–hook is a structure joining Mre11 complexes in DNA recombination

[143208]and repair. Nature 418: 562–566, 2002.

[143209]

[143210]

[143211]13467.

[143212]Luo, G.; Yao, M. S.; Bender, C. F.; Mills, M.; Bladl, A. R.;
Bradley,

[143213]A.; Petrini, J. H. J.: Disruption of mRad50 causes embry-
onic stem

[143214]cell lethality, abnormal embryonic development, and sen-
sitivity to

[143215]ionizing radiation. Proc. Nat. Acad. Sci. 96: 7376–7381,
1999.

[143216]

[143217]

[143218]13468.

[143219]Trujillo, K. M.; Yuan, S.–S. F.; Lee, E. Y.–H. P.; Sung, P.:
Nuclease

[143220]activities in a complex of human recombination and DNA
repair factors

[143221]Rad50, Mre11, and p95. J. Biol. Chem. 273:
21447–21450, 1998.

[143222]

[143223]

[143224]13469.

[143225]Clark, J.; Lu, Y. J.; Sidhar, S. K.; Parker, C.; Gill, S.; Smedley,

[143226]D.; Hamoudi, R.; Linehan, W.; Shipley, J.; Cooper, C.: Fusion of

[143227]splicing factor genes PSF and NonO (p54nrb) to the TFE3 gene in papillary

[143228]renal cell carcinoma. *Oncogene* 15: 2233–2239, 1997.

[143229]

[143230]

[143231]13470.

[143232]Heimann, P.; El Housni, H.; Ogur, G.; Weterman, M. A. J.; Petty,

[143233]E. M.; Vassart, G.: Fusion of a novel gene, RCC17, to the TFE3 gene

[143234]in t(X;17)(p11.2;q25.3)–bearing papillary renal cell carcinomas. *Cancer*

[143235]*Res.* 61: 4130–4135, 2001.

[143236]

[143237]

[143238]13471.

[143239]Henthorn, P. S.; Stewart, C. C.; Kadesch, T.; Puck, J. M.:

The

[143240]gene encoding human TFE3, a transcription factor that
binds the immunoglobulin

[143241]heavy-chain enhancer, maps to Xp11.22. Genomics 11:
374–378, 1991.

[143242]

[143243]

[143244]13472.

[143245]Joyama, S.; Ueda, T.; Shimizu, K.; Kudawara, I.; Mano, M.;
Funai,

[143246]H.; Takemura, K.; Yoshikawa, H.: Chromosome rearrange-
ment at 17q25

[143247]and Xp11.2 in alveolar soft-part sarcoma: a case report
and review

[143248]of the literature. Cancer 86: 1246–1250, 1999.

[143249]

[143250]

[143251]13473.

[143252]Ladanyi, M.; Lui, M. Y.; Antonescu, C. R.; Krause-Boehm,
A.; Meindl,

[143253]A.; Argani, P.; Healey, J. H.; Ueda, T.; Yoshikawa, H.; Meloni-Ehrig,

[143254]A.; Sorensen, P. H. B.; Mertens, F.; Mandahl, N.; van den Berghe,

[143255]H.; Sciot, R.; Dal Cin, P.; Bridge, J.: The der(17)t(X;17)(p11;q25)

[143256]of human alveolar soft part sarcoma fuses the TFE3 transcription factor

[143257]gene to ASPL, a novel gene at 17q25. Oncogene 20: 48–57, 2001.

[143258]

[143259]

[143260]13474.

[143261]Macchi, P.; Notarangelo, L.; Giliani, S.; Strina, D.; Repetto,

[143262]M.; Sacco, M. G.; Vezzoni, P.; Villa, A.: The genomic organization

[143263]of the human transcription factor 3 (TFE3) gene. Genomics 28: 491–494,

[143264]1995.

[143265]

[143266]

[143267]13475.

[143268]Shipley, J. M.; Birdsall, S.; Clark, J.; Crew, J.; Gill, S.; Linehan,

[143269]M.; Gnarr, J.; Gisher, S.; Craig, I. W.; Cooper, C. S.: Mapping

[143270]the X chromosome breakpoint in two papillary renal cell carcinoma

[143271]cell lines with a t(X;1)(p11.2;q21.2) and the first report of a female

[143272]case. Cytogenet. Cell. Genet. 71: 280–284, 1995.

[143273]

[143274]

[143275]13476.

[143276]Sidhar, S. K.; Clark, J.; Gill, S.; Hamoudi, R.; Crew, A. J.;

[143277]Gwilliam, R.; Ross, M.; Linehan, W. M.; Birdsall, S.; Shipley, J.;

[143278]Cooper, C. S.: The t(X;1)(p11.2;q21.2) translocation in papillary

[143279]renal cell carcinoma fuses a novel gene PRCC to the TFE3 transcription

[143280]factor gene. Hum. Molec. Genet. 5: 1333–1338, 1996.

[143281]

[143282]

[143283]13477.

[143284]Larola, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.: Control

[143285]of mRNA decay by heat shock-ubiquitin-proteasome

pathway. Science 284:

[143286]499-502, 1999.

[143287]

[143288]

[143289]13478.

[143290]Deutsch, D.; Palmon, A.; Dafni, L.; Fisher, L.; Termine, J.

D.;

[143291]Young, M.: Cloning, sequencing, and characterization of

tuftelin:

[143292]a novel acidic enamel protein. Connect. Tissue Res. 27:

121 only,

[143293]1992.

[143294]

[143295]

[143296]13479.

[143297]Deiss, L. P.; Feinstein, E.; Berissi, H.; Cohen, O.; Kimchi, A.

[143298]: Identification of a novel serine/threonine kinase and a novel 15-kD

[143299]protein as potential mediators of the gamma interferon-induced cell

[143300]death. Genes Dev. 9: 15-30, 1995.

[143301]

[143302]

[143303]13480.

[143304]Feinstein, E.; Druck, T.; Kastury, K.; Berissi, H.; Goodart, S.

[143305]A.; Overhauser, J.; Kimchi, A.; Huebner, K.: Assignment of DAP1 and

[143306]DAPK: genes that positively mediate programmed cell death triggered

[143307]by IFN-gamma--to chromosome regions 5p12.2 (sic) and 9q34.1, respectively. Genomics 29:

[143308]305-307, 1995.

[143309]

[143310]

[143311]13481.

[143312]Fink, T. M.; Vaesen, M.; Kratzin, H. D.; Lichter, P.; Zimmer, M.

[143313]: Localization of the gene encoding the putative human HLA class II

[143314]associated protein (PHAPI) to chromosome 15q22.3–q23 by fluorescence

[143315]in situ hybridization. Genomics 29: 309–310, 1995.

[143316]

[143317]

[143318]13482.

[143319]Li, M.; Makkinje, A.; Damuni, Z.: Molecular identification of

[143320]I–1(PP2A), a novel potent heat–stable inhibitor protein of protein

[143321]phosphatase 2A. Biochemistry 35: 6998–7002, 1996.

[143322]

[143323]

[143324]13483.

[143325]Matilla, A.; Koshy, B. T.; Cummings, C. J.; Isobe, T.; Orr, H.

[143326]T.; Zoghbi, H. Y.: The cerebellar leucine-rich acidic nuclear protein

[143327]interacts with ataxin-1. Nature 389: 974-978, 1997.

Note: Erratum:

[143328]Nature 391: 818 only, 1998.

[143329]

[143330]

[143331]13484.

[143332]Sandberg, A. A.; Morgan, R.; McCallister, J. A.; Kaiser-McCaw,

[143333]B.; Hecht, F.: Acute myeloblastic leukemia (AML) with t(6;9)(p23;q34):

[143334]a specific subgroup of AML? Cancer Genet. Cytogenet. 10: 139-142,

[143335]1983.

[143336]

[143337]

[143338]13485.

[143339]Vaesen, M.; Barnikol-Watanabe, S.; Gotz, H.; Awni, L. A.; Cole,

[143340]T.; Zimmermann, B; Kratzin, H. D.; Hilschmann, N.: Purification and

[143341]characterization of two putative HLA class II associated proteins:

[143342]PHAPI and PHAPII. Biol. Chem. Hoppe-Seyler 375: 113-126, 1994.

[143343]

[143344]

[143345]13486.

[143346]von Lindern, M.; van Baal, S.; Wiegant, J.; Raap, A.; Hagemijer,

[143347]A.; Grosveld, G.: Can, a putative oncogene associated with myeloid

[143348]leukemogenesis may be activated by fusion of its 3-prime half to different

[143349]genes: characterization of the set gene. Molec. Cell Biol. 12: 3346-3355,

[143350]1992.

[143351]

[143352]

[143353]13487.

[143354]Hughes, K. A.; Hurlstone, A. F. L.; Tobias, E. S.; McFarlane, R.;

[143355]Black, D. M.: Absence of ST7 mutations in tumor-derived cell lines

[143356]and tumors. Nature Genet. 29: 380–381, 2001.

[143357]

[143358]

[143359]13488.

[143360]Thomas, N. A.; Choong, D. Y. H.; Jokubaitis, V. J.; Neville, P.

[143361]J.; Campbell, I. G.: Mutation of the ST7 tumor suppressor gene on

[143362]7q31.1 is rare in breast, ovarian and colorectal cancers.

Nature

[143363]Genet. 29: 379–380, 2001.

[143364]

[143365]

[143366]13489.

[143367]Zenklusen, J. C.; Conti, C. J.; Green, E. D.: Mutational and functional

[143368]analyses reveal that ST7 is a highly conserved tumor-suppressor gene

[143369]on human chromosome 7q31. Nature Genet. 27: 392-398, 2001.

[143370]

[143371]

[143372]13490.

[143373]Zenklusen, J. C.; Rodriguez, L. V.; LaCava, M.; Wang, Z.; Goldstein,

[143374]L. S.; Conti, C. J.: Novel susceptibility locus for mouse hepatomas:

[143375]evidence for a conserved tumor suppressor gene. Genome Res. 6: 1070-1076,

[143376]1996.

[143377]

[143378]

[143379]13491.

[143380]Zenklusen, J. C.; Weitzel, J. N.; Ball, H. G.; Conti, C. J.: Allelic

[143381]loss at 7q31.1 in human primary ovarian carcinomas suggests the existence

[143382]of a tumor suppressor gene. Oncogene 11: 359–363,
1995.

[143383]

[143384]

[143385]13492.

[143386]Lee, P. L.; Gelbart, T.; West, C.; Adams, M.; Blackstone, R.;
Beutler,

[143387]E.: Three genes encoding zinc finger proteins on human
chromosome

[143388]6p21.3: members of a new subclass of the Kruppel gene
family containing

[143389]the conserved SCAN box domain. Genomics 43: 191–201,
1997.

[143390]

[143391]

[143392]13493.

[143393]Tirosvoutis, K. N.; Divane, A.; Jones, M.; Affara, N. A.:
Characterization

[143394]of a novel zinc finger gene (ZNF165) mapping to 6p21
that is expressed

[143395]specifically in testis. Genomics 28: 485–490, 1995.

[143396]

[143397]

[143398]13494.

[143399]Deutsch, D.; Palmon, A.; Fisher, L. W.; Kolodny, N.; Ter-
mine, J.

[143400]D.; Young, M. F.: Sequencing of bovine enamelin (tuftelin),
a novel

[143401]acidic enamel protein. J. Biol. Chem. 266: 16021–16028,
1991.

[143402]

[143403]

[143404]13495.

[143405]Deutsch, D.; Palmon, A.; Young, M. F.; Selig, S.; Kearns, W.
G.;

[143406]Fisher, L. W.: Mapping of the human tuftelin (TUFT1) gene
to chromosome

[143407]1 by fluorescence in situ hybridization. (Abstract) Mam-
malian Genome 5:

[143408]461–462, 1994.

[143409]

[143410]

[143411]13496.

[143412]MacDougall, M.; Simmons, D.; Dodds, A.; Knight, C.; Luan, X.; Zeichner-David,

[143413]M.; Zhang, C.; Ryu, O. H.; Qian, Q.; Simmer, J. P.; Hu, C.-C.: Cloning,

[143414]characterization, and tissue expression pattern of mouse tuftelin

[143415]cDNA. J. Dent. Res. 77: 1970–1978, 1998.

[143416]

[143417]

[143418]13497.

[143419]Dickeson, S. K.; Helmkamp, G. M., Jr.; Yarbrough, L. R.: Sequence

[143420]of a human cDNA encoding phosphatidylinositol transfer protein and

[143421]occurrence of a related sequence in widely divergent eukaryotes. Gene 142:

[143422]301–305, 1994.

[143423]

[143424]

[143425]13498.

[143426]Dickeson, S. K.; Lim, C. N.; Schuyler, G. T.; Dalton, T. P.;
Helmkamp,

[143427]G. M., Jr.; Yarbrough, L. R.: Isolation and sequence of
cDNA clones

[143428]encoding rat phosphatidylinositol transfer protein. J. Biol.
Chem. 264:

[143429]16557–16564, 1989.

[143430]

[143431]

[143432]13499.

[143433]Hay, J. C.; Martin, T. F. J.: Phosphatidylinositol transfer
protein

[143434]required for ATP–dependent priming of Ca^{2+} –activated
secretion. Nature 366:

[143435]572–575, 1993.

[143436]

[143437]

[143438]13500.

[143439]Vihtelic, T. S.; Goebel, M.; Milligan, S.; O'Tousa, J. E.; Hyde,
[143440]D. R.: Localization of Drosophila retinal degeneration B, a
membrane-associated
[143441]phosphatidylinositol transfer protein. J. Cell Biol. 122:
1013-1022,
[143442]1993.
[143443]
[143444]
[143445]13501.
[143446]Wirtz, K. W. A.: Phospholipid transfer proteins. Annu. Rev.
Biochem. 60:
[143447]73-99, 1991.
[143448]
[143449]
[143450]13502.
[143451]Fitzgibbon, J.; Pilz, A.; Gayther, S.; Appukuttan, B.; Dulai,
K.
[143452]S.; Delhanty, J. D. A.; Helmkamp, G. M., Jr.; Yarbrough, L.
R.; Hunt,
[143453]D. M.: Localization of the gene encoding human phos-

phatidylinositol

[143454]transfer protein (PITPN) to 17p13.3: a gene showing homology to the

[143455]*Drosophila* retinal degeneration B gene (rdgB). *Cytogenet. Cell Genet.* 67:

[143456]205–207, 1994.

[143457]

[143458]

[143459]13503.

[143460]Hamilton, B. A.; Smith, D. J.; Mueller, K. L.; Kerrebrock, A. W.;

[143461]Bronson, R. T.; van Berkel, V.; Daly, M. J.; Kruglyak, L.; Reeve,

[143462]M. P.; Nemhauser, J. L.; Hawkins, T. L.; Rubin, E. M.; Lander, E.

[143463]S.: The vibrator mutation causes neurodegeneration via reduced expression

[143464]of PITP- α : positional complementation cloning and extragenic suppression. *Neuron* 18:

[143465]711–722, 1997.

[143466]

[143467]

[143468]13504.

[143469]Hara, T.; Yamauchi, M.; Takahashi, E.; Hoshino, M.; Aoki, K.; Ayusawa,

[143470]D.; Kawakita, M.: The UDP–galactose translocator gene is mapped to

[143471]band Xp11.23–p11.22 containing the Wiskott–Aldrich syndrome locus. Somat.

[143472]Cell Molec. Genet. 19: 571–575, 1993.

[143473]

[143474]

[143475]13505.

[143476]Ishida, N.; Miura, N.; Yoshioka, S.; Kawakita, M.: Molecular cloning

[143477]and characterization of a novel isoform of the human UDP–galactose

[143478]transporter, and of related complementary DNAs belonging to the nucleotide–sugar

[143479]transporter gene family. J. Biochem. 120: 1074–1078, 1996.

[143480]

[143481]

[143482]13506.

[143483]Miura, N.; Ishida, N.; Hoshino, M.; Yamauchi, M.; Hara, T.;

Ayusawa,

[143484]D.; Kawakita, M.: Human UDP-galactose translocator:

molecular cloning

[143485]of a complementary DNA that complements the genetic

defect of a mutant

[143486]cell line deficient in UDP-galactose translocator. J.

Biochem. 120:

[143487]236-241, 1996.

[143488]

[143489]

[143490]13507.

[143491]Lund, A.; Udd, B.; Juvonen, V.; Andersen, P. M.; Ced-

erquist, K.;

[143492]Davis, M.; Gellera, C.; Kolmel, C.; Ronnevi, L.-O.; Sperfeld,

A.-D.;

[143493]Sorensen, S.-A.; Tranebjaerg, L.; Van Maldergem, L.;

Watanabe, M.;

[143494]Weber, M.; Yeung, L.; Savontaus, M.-L.: Multiple founder

effects

[143495]in spinal and bulbar muscular atrophy (SBMA, Kennedy disease) around

[143496]the world. *Europ. J. Hum. Genet.* 9: 431–436, 2001.

[143497]

[143498]

[143499]13508.

[143500]Edelhoff, S.; Ayer, D. E.; Zervos, A. S.; Steingrimsson, E.; Jenkins,

[143501]N. A.; Copeland, N. G.; Eisenman, R. N.; Brent, R.; Disteche, C. M.

[143502]: Mapping of two genes encoding members of a distinct subfamily of

[143503]MAX interacting proteins: MAD to human chromosome 2 and mouse chromosome

[143504]6, and MXI1 to chromosome 10 and mouse chromosome 19. *Oncogene* 9:

[143505]665–668, 1994.

[143506]

[143507]

[143508]13509.

[143509]Ferguson, G. D.; Anagnostaras, S. G.; Silva, A. J.; Herschman,

[143510]H. R.: Deficits in memory and motor performance in synaptotagmin

[143511]IV mutant mice. Proc. Nat. Acad. Sci. 97: 5598–5603, 2000.

[143512]

[143513]

[143514]13510.

[143515]Ferguson, G. D.; Chen, X.–N.; Korenberg, J. R.; Herschman, H. R.

[143516]: The human synaptotagmin IV gene defines an evolutionary break point

[143517]between syntenic mouse and human chromosome regions but retains ligand

[143518]inducibility and tissue specificity. J. Biol. Chem. 275: 36920–36926,

[143519]2000.

[143520]

[143521]

[143522]13511.

[143523]Holterhus, P.-M.; Wiebel, J.; Sinnecker, G. H. G.; Bruggen-
wirth,

[143524]H. T.; Sippell, W. G.; Brinkmann, A. O.; Kruse, K.; Hiort, O.:
Clinical

[143525]and molecular spectrum of somatic mosaicism in andro-
gen insensitivity

[143526]syndrome. *Pediat. Res.* 46: 684–690, 1999.

[143527]

[143528]

[143529]13512.

[143530]Foresta, C.; Bertella, A.; Moro, E.; Roverato, A.; Merico, M.;

[143531]Ferlin, A.: Sertoli cell function in infertile patients with and

[143532]without microdeletions of the azoospermia factors on the
Y chromosome

[143533]long arm. *J. Clin. Endocr. Metab.* 86: 2414–2419, 2001.

[143534]

[143535]

[143536]13513.

[143537]Kobayashi, K.; Mizuno, K.; Hida, A.; Komaki, R.; Tomita,

- K.; Matsushita,
[143538]I.; Namiki, M.; Iwamoto, T.; Tamura, S.; Minowada, S.;
Nakahori, Y.;
- [143539]Nakagome, Y.: PCR analysis of the Y chromosome long
arm in azoospermic
[143540]patients: evidence for a second locus required for sper-
matogenesis. Hum.
[143541]Molec. Genet. 3: 1965–1967, 1994.
- [143542]
- [143543]
- [143544]13514.
- [143545]Agulnik, A. I.; Zharkikh, A.; Boettger–Tong, H.; Bourgeron,
T.;
- [143546]McElreavey, K.; Bishop, C. E.: Evolution of the DAZ gene
family suggests
[143547]that Y–linked DAZ plays little, or a limited, role in sper-
matogenesis
[143548]but underlines a recent African origin for human popula-
tions. Hum.
[143549]Molec. Genet. 7: 1371–1377, 1998.
- [143550]

[143551]

[143552]13515.

[143553]Anonymous: A missing piece on the Y. (Editorial) Nature
Genet. 10:

[143554]367–368, 1995.

[143555]

[143556]

[143557]13516.

[143558]Cooke, H. J.; Lee, M.; Kerr, S.; Ruggiu, M.: A murine homo-
logue

[143559]of the human DAZ gene is autosomal and expressed only
in male and

[143560]female gonads. Hum. Molec. Genet. 5: 513–516, 1996.

[143561]

[143562]

[143563]13517.

[143564]Foresta, C.; Moro, E.; Garolla, A.; Onisto, M.; Ferlin, A.: Y

[143565]chromosome microdeletions in cryptorchidism and idio-
pathic infertility. J.

[143566]Clin. Endocr. Metab. 84: 3660–3665, 1999.

[143567]

[143568]

[143569]13518.

[143570]Makova, K. D.; Li, W.–H.: Strong male–driven evolution of
DNA

[143571]sequences in humans and apes. Nature 416: 624–626,
2002.

[143572]

[143573]

[143574]13519.

[143575]Menke, D. B.; Mutter, G. L.; Page, D. C.: Expression of
DAZ, an

[143576]azoospermia factor candidate, in human spermatogonia.
(Letter) Am.

[143577]J. Hum. Genet. 60: 237–241, 1997.

[143578]

[143579]

[143580]13520.

[143581]Moro, E.; Ferlin, A.; Yen, P. H.; Franchi, P. G.; Palka, G.;
Foresta,

[143582]C.: Male infertility caused by a de novo partial deletion of
the

[143583]DAZ cluster on the Y chromosome. J. Clin. Endocr. Metab.
85: 4069–4073,

[143584]2000.

[143585]

[143586]

[143587]13521.

[143588]Reijo, R.; Lee, T.–Y.; Salo, P.; Alagappan, R.; Brown, L. G.;

[143589]Rosenberg, M.; Rozen, S.; Jaffe, T.; Straus, D.; Hovatta, O.;

de la

[143590]Chapelle, A.; Silber, S.; Page, D. C.: Diverse spermatogenic
defects

[143591]in humans caused by Y chromosome deletions encom-
passing a novel RNA-binding

[143592]protein gene. Nature Genet. 10: 383–393, 1995.

[143593]

[143594]

[143595]13522.

[143596]Reijo, R.; Seligman, J.; Dinulos, M. B.; Jaffe, T.; Brown, L.

[143597]G.; Disteché, C. M.; Page, D. C.: Mouse autosomal homolog of DAZ,

[143598]a candidate male sterility gene in humans, is expressed in male germ

[143599]cells before and after puberty. Genomics 35: 346–352, 1996.

[143600]

[143601]

[143602]13523.

[143603]Saxena, R.; Brown, L. G.; Hawkins, T.; Alagappan, R. K.; Skaletsky,

[143604]H.; Reeve, M. P.; Reijo, R.; Rozen, S.; Dinulos, M. B.; Disteché,

[143605]C. M.; Page, D. C.: The DAZ gene cluster on the human Y chromosome

[143606]arose from an autosomal gene that was transposed, repeatedly amplified

[143607]and pruned. Nature Genet. 14: 292–299, 1996.

[143608]

[143609]

[143610]13524.

[143611]Najmabadi, H.; Huang, V.; Yen, P.; Subbarao, M. N.;
Bhasin, D;

[143612]Banaag, L.; Naseeruddin, S.; de Kretser, D. M.; Baker, H. W.
G.; McLachlan,

[143613]R. I.; Loveland, K. A.; Bhasin, S.: Substantial prevalence of
microdeletions

[143614]of the Y-chromosome in infertile men with idiopathic
azoospermia and

[143615]oligozoospermia detected using a sequence-tagged site-
based mapping

[143616]strategy. J. Clin. Endocr. Metab. 81: 1347-1352, 1996.

[143617]

[143618]

[143619]13525.

[143620]Wang, G.; Sawai, N.; Kotliarova, S.; Kanazawa, I.; Nukina,
N.:

[143621]Ataxin-3, the MJD1 gene product, interacts with the two
human homologs

[143622]of yeast DNA repair protein RAD23, HHR23A and HHR23B.

Hum. Molec.

[143623]Genet. 9: 1795–1803, 2000.

[143624]

[143625]

[143626]13526.

[143627]Ng, J. M. Y.; Vrieling, H.; Sugasawa, K.; Ooms, M. P.;
Grootegoed,

[143628]J. A.; Vreeburg, J. T. M.; Visser, P.; Beems, R. B.; Gorgels,
T. G.

[143629]M. F.; Hanaoka, F.; Hoeijmakers, J. H. J.; van der Horst, G.
T. J.

[143630]: Developmental defects and male sterility in mice lacking
the ubiquitin-like

[143631]DNA repair gene mHR23B. Molec. Cell. Biol. 22:
1233–1245, 2002.

[143632]

[143633]

[143634]13527.

[143635]Volker, M.; Mone, M. J.; Karmakar, P.; van Hoffen, A.;
Schul, W.;

[143636]Vermeulen, W.; Hoeijmakers, J. H. J.; van Driel, R.; van

Zeeland,

[143637]A. A.; Mullenders, L. H. F.: Sequential assembly of the nucleotide

[143638]excision repair factors in vivo. Molec. Cell 8: 213–224, 2001.

[143639]

[143640]

[143641]13528.

[143642]Abramson, J. S.; Mills, E. L.; Sawyer, M. K.; Regelman, W. R.;

[143643]Nelson, J. D.; Quie, P. G.: Recurrent infections and delayed separation

[143644]of the umbilical cord in an infant with abnormal phagocytic cell locomotion

[143645]and oxidative response during particle phagocytosis. J. Pediat. 99:

[143646]887–894, 1981.

[143647]

[143648]

[143649]13529.

[143650]Allende, L. M.; Hernandez, M.; Corell, A.; Garcia-Perez, M.

A.;

[143651]Varela, P.; Moreno, A.; Caragol, I.; Garcia-Martin, F.;

Guillen-Perales,

[143652]J.; Olive, T.; Espanol, T.; Arnaiz-Villena, A.: A novel CD18

genomic

[143653]deletion in a patient with severe leucocyte adhesion defi-

ciency: a

[143654]possible CD2/lymphocyte function-associated antigen-1

functional association

[143655]in humans. Immunology 99: 440-450, 2000.

[143656]

[143657]

[143658]13530.

[143659]Anderson, D. C.; Springer, T. A.: Leukocyte adhesion defi-

ciency:

[143660]an inherited defect in the Mac-1, LFA-1, and p150,95 gly-

coproteins. Annu.

[143661]Rev. Med. 38: 175-194, 1987.

[143662]

[143663]

[143664]13531.

[143665]Arnaout, M. A.; Dana, N.; Gupta, S. K.; Tenen, D. G.;

Fathallah,

[143666]D. M.: Point mutations impairing cell surface expression of
the common

[143667]beta subunit (CD18) in a patient with leukocyte adhesion
molecule

[143668](Leu-CAM) deficiency. J. Clin. Invest. 85: 977-981, 1990.

[143669]

[143670]

[143671]13532.

[143672]Arnaout, M. A.; Pitt, J.; Cohen, H. J.; Melamed, J.; Rosen, F.

[143673]S.; Colten, H. R.: Deficiency of a granulocyte-membrane
glycoprotein

[143674](gp150) in a boy with recurrent bacterial infections. New
Eng. J.

[143675]Med. 306: 693-699, 1982.

[143676]

[143677]

[143678]13533.

[143679]Arnaout, M. A.; Spits, H.; Terhorst, C.; Pitt, J.; Todd, R. F.,

[143680]III: Deficiency of a leukocyte surface glycoprotein (LFA-1)
in two

[143681]patients with Mo1 deficiency: effects of cell activation on
Mo1/LFA-1

[143682]surface expression in normal and deficient leukocytes. J.
Clin. Invest. 74:

[143683]1291-1300, 1984.

[143684]

[143685]

[143686]13534.

[143687]Anderson, D. C.; Schmalstieg, F. C.; Finegold, M. J.;
Hughes, B.

[143688]J.; Rothlein, R.; Miller, L. J.; Kohl, S.; Tosi, M. F.; Jacobs, R.

[143689]L.; Waldrop, T. C.; Goldman, A. S.; Shearer, W. T.; Springer,
T. A.

[143690]: The severe and moderate phenotypes of heritable Mac-
1, LFA-1 deficiency:

[143691]their quantitative definition and relation to leukocyte dys-
function

[143692]and clinical features. J. Infect. Dis. 152: 668-689, 1985.

[143693]

[143694]

[143695]13535.

[143696]Breviario, F.; d'Aniello, E. M.; Golay, J.; Peri, G.; Bottazzi,

[143697]B.; Bairoch, A.; Saccone, S.; Marzella, R.; Predazzi, V.; Rocchi,

[143698]M.; Della Valle, G.; Dejana, E.; Mantovani, A.; Introna, M.:
Interleukin-1-inducible

[143699]genes in endothelial cells: cloning of a new gene related
to C-reactive

[143700]protein and serum amyloid P component. J. Biol. Chem.
267: 22190-22197,

[143701]1992.

[143702]

[143703]

[143704]13536.

[143705]Lee, T. H.; Lee, G. W.; Ziff, E. B.; Vilcek, J.: Isolation and

[143706]characterization of eight tumor necrosis factor-induced
gene sequences

[143707]from human fibroblasts. Molec. Cell. Biol. 10: 1982-1988,
1990.

[143708]

[143709]

[143710]13537.

[143711]Pallini, R.; Guazzi, G. C.; Cannella, C.; Cacace, M. G.:
Cloning

[143712]and sequence analysis of the human liver rhodanese:

comparison with

[143713]the bovine and chicken enzymes. Biochem. Biophys. Res.

Commun. 180:

[143714]887–893, 1991.

[143715]

[143716]

[143717]13538.

[143718]Greco, A.; Mariani, C.; Miranda, C.; Lupas, A.; Pagliardini, S.;

[143719]Pomati, M.; Pierotti, M. A.: The DNA rearrangement that generates

[143720]the TRK–T3 oncogene involves a novel gene on chromosome 3 whose product

[143721]has a potential coiled–coil domain. Molec. Cell Biol. 15: 6118–6127,

[143722]1995.

[143723]

[143724]

[143725]13539.

[143726]Mencinger, M.; Panagopoulos, I.; Andreasson, P.; Lassen, C.; Mitelman,

[143727]F.; Aman, P.: Characterization and chromosomal mapping of the human

[143728]TFG gene involved in thyroid carcinoma. Genomics 41: 327–331, 1997.

[143729]

[143730]

[143731]13540.

[143732]Kramps, T.; Peter, O.; Brunner, E.; Nellen, D.; Froesch, B.; Chatterjee,

[143733]S.; Murone, M.; Zullig, S.; Basler, K.: Wnt/Wingless signaling requires

[143734]BCL9/legless-mediated recruitment of pygopus to the nuclear beta-catenin-TCF

[143735]complex. Cell 109: 47–60, 2002.

[143736]

[143737]

[143738]13541.

[143739]Willis, T. G.; Zalcberg, I. R.; Coignet, L. J. A.; Wlodarska, M.;

[143740]Stul, D. M.; Jadayel, D. M.; Bastard, C.; Treleaven, J. G.;
Catovsky,

[143741]D.; Silva, M. L. M.; Dyer, M. J. S.: Molecular cloning of
translocation

[143742]t(1;14)(q21;q32) defines a novel gene (BCL9) at chromo-
some 1q21. Blood 91:

[143743]1873–1881, 1998.

[143744]

[143745]

[143746]13542.

[143747]Kim, D.-H.; Iijima, H.; Goto, K.; Sakai, J.; Ishii, H.; Kim,
H.-J.;

[143748]Suzuki, H.; Kondo, H.; Saeki, S.; Yamamoto, T.: Human
apolipoprotein

[143749]E receptor 2: a novel lipoprotein receptor of the low den-
sity lipoprotein

[143750]receptor family predominantly expressed in brain. J. Biol.
Chem. 271:

[143751]8373–8380, 1996.

[143752]

[143753]

[143754]13543.

[143755]Kim, D.-H.; Magoori, K.; Inoue, T. R.; Mao, C. C.; Kim, H.-J.;

[143756]Suzuki, H.; Fujita, T.; Endo, Y.; Saeki, S.; Yamamoto, T. T.:
Exon/intron

[143757]organization, chromosome localization, alternative splicing, and transcription

[143758]units of the human apolipoprotein E receptor 2 gene. J. Biol. Chem. 272:

[143759]8498–8504, 1997.

[143760]

[143761]

[143762]13544.

[143763]Davidson, R. G.; Nitowsky, H. M.; Childs, B.: Demonstration of

[143764]two populations of cells in the human female heterozygous for glucose-6-phosphate

[143765]dehydrogenase variants. Proc. Nat. Acad. Sci. 50:
481–485, 1963.

[143766]

[143767]

[143768]13545.

[143769]De Flora, A.; Morelli, A.; Benatti, U.; Giuntini, P.; Ferraris,

[143770]A. M.; Galiano, S.; Ravazzolo, R.; Gaetani, G. F.: G6PD

Napoli and

[143771]Ferrara II: two new glucose-6-phosphate dehydrogenase

variants having

[143772]similar characteristics but different intracellular lability

and specific

[143773]activity. Brit. J. Haemat. 48: 417-423, 1981.

[143774]

[143775]

[143776]13546.

[143777]Dern, R. J.; McCurdy, P. R.; Yoshida, A.: A new structural

variant

[143778]of glucose-6-phosphate dehydrogenase with a high pro-

duction rate (G6PD

[143779]Hektoen). J. Lab. Clin. Med. 73: 283-290, 1969.

[143780]

[143781]

[143782]13547.

[143783]De Vita, G.; Alcalay, M.; Sampietro, M.; Cappelini, M. D.;
Fiorelli,

[143784]G.; Toniolo, D.: Two point mutations are responsible for
G6PD polymorphism

[143785]in Sardinia. Am. J. Hum. Genet. 44: 233–240, 1989.

[143786]

[143787]

[143788]13548.

[143789]Du, C.; Xu, Y.; Hua, X.; Wu, Q.; Liu, L.; Wu, M.: Studies on

[143790]erythrocyte glucose–6–phosphate dehydrogenase variants
in Chinese.

[143791]III: Gd(–) Miaozi–Baisha. Acta Genet. Sinica 11(2):
153–158, 1984.

[143792]

[143793]

[143794]13549.

[143795]Du, C.–S.; Hua, X.–Y.; Wu, Q.–L.; Li, C.–Q.; Zheng, J.–F.; Li,

[143796]H.–L.: Studies on erythrocyte glucose–6–phosphate dehy-
drogenase variants

[143797]in Chinese. IV. Gd(–) Gaohe associated with paroxysmal

nocturnal hemoglobinuria. Chinese

[143798]J. Pathophysiol. 1: 12–15, 1985.

[143799]

[143800]

[143801]13550.

[143802]Du, C. S.; Xu, Y. K.; Hua, X. Y.; Wu, Q. L.; Liu, L. B.: Glucose-6-phosphate

[143803]dehydrogenase variants and their frequency in Guangdong, China. Hum.

[143804]Genet. 80: 385–388, 1988.

[143805]

[143806]

[143807]13551.

[143808]Eber, S. W.; Gahr, M.; Schroter, W.: Glucose-6-phosphate dehydrogenase

[143809](G6PD) Iserlohn and G6PD Regensburg: two new severe enzyme defects

[143810]in German families. Blut 51: 109–115, 1985.

[143811]

[143812]

[143813]13552.

[143814]Echard, G.; Gillois, M.: G6PD--alpha-GAL-PGK--HPRT
syntenry in

[143815]the rabbit, *Oryctolagus cuniculus*. (Abstract) Cytogenet.
Cell Genet. 25:

[143816]148-149, 1979.

[143817]

[143818]

[143819]13553.

[143820]Elizondo, J.; Saenz, G. F.; Paez, C. A.; Ramon, M.; Garcia,
M.;

[143821]Gutierrez, A.; Estrada, M.: G6PD Puerto Limon: a new defi-
cient variant

[143822]of glucose-6-phosphate dehydrogenase associated with
congenital nonspherocytic

[143823]hemolytic anemia. Hum. Genet. 62: 110-112, 1982.

[143824]

[143825]

[143826]13554.

[143827]Engstrom, P. F.; Beutler, E.: G-6-PD Tripler: a unique variant

[143828]associated with chronic hemolytic disease. Blood 36:
10-13, 1970.

[143829]

[143830]

[143831]13555.

[143832]Epstein, C. J.: Mammalian oocytes: X-chromosome activity. Science 163:

[143833]1078-1079, 1969.

[143834]

[143835]

[143836]13556.

[143837]Ermakov, N. V.; Chernyak, N. B.; Tokarev, Y. N.: Properties of

[143838]new variant of glucose-6-phosphate dehydrogenase
(Regar variant):

[143839]glucose metabolism in erythrocytes containing abnormal
enzyme. Biokhimiia 48:

[143840]577-583, 1983.

[143841]

[143842]

[143843]13557.

[143844]Estrada, M.; Garcia, M.; Gutierrez, A.; Quintero, I.; Gonzalez,
lez,

[143845]R.: G6PD Varadero. Vox Sang. 43: 102–104, 1982.

[143846]

[143847]

[143848]13558.

[143849]Fairbanks, V. F.; Nepo, A. G.; Beutler, E.; Dickson, E. R.;
Honig,

[143850]G.: Glucose–6–phosphate dehydrogenase variants: reex–
amination of

[143851]G6PD Chicago and Cornell and a new variant (G6PD Pea
Ridge) resembling

[143852]G6PD Chicago. Blood 55: 216–220, 1980.

[143853]

[143854]

[143855]13559.

[143856]Faust, C. J.; Levinson, B.; Gitschier, J.; Herman, G. E.: Ex-
tension

[143857]of the physical map in the region of the mouse X chromo-
some homologous

[143858]to human Xq28 and identification of an exception to con-
served linkage. Genomics 13:

[143859]1289-1295, 1992.

[143860]

[143861]

[143862]13560.

[143863]Feldman, R.; Gromisch, D. S.; Luhby, A. L.; Beutler, E.:
Congenital

[143864]nonspherocytic hemolytic anemia due to glucose-
6-phosphate dehydrogenase

[143865]East Harlem: a new deficient variant. J. Pediat. 90: 89-91,
1977.

[143866]

[143867]

[143868]13561.

[143869]Fernandez, M.; Fairbanks, V. F.: Glucose-6-phosphate de-
hydrogenase

[143870]deficiency in the Philippines: report of a new variant–
–G6PD Panay. Mayo

[143871]Clin. Proc. 43: 645–660, 1968.

[143872]

[143873]

[143874]13562.

[143875]Ferraris, A. M.; Broccia, G.; Meloni, T.; Forteleoni, G.; Gae–
tani,

[143876]G. F.: Glucose–6–phosphate dehydrogenase deficiency
and incidence

[143877]of hematologic malignancy. Am. J. Hum. Genet. 42:
516–520, 1988.

[143878]

[143879]100. Filosa, E.: Personal Communication. La Jolla, Calif.
1989.

[143880]

[143881]101. Filosa, S.; Calabro, V.; Vallone, D.; Poggi, V.; Mason,
P.; Pagnini,

[143882]D.; Alfinito, F.; Rotoli, B.; Martini, G.; Luzzatto, L.; Battis–
tuzzi,

[143883]G.: Molecular basis of chronic non–spherocytic haemolytic

anaemia:

[143884]a new G6PD variant (393arg-to-his) with abnormal K(m)
G6P and marked

[143885]in vivo instability. Brit. J. Haemat. 80: 111–116, 1992.

[143886]

[143887]102. Filosa, S.; Giacometti, N.; Wangwei, C.; De Mattia, D.;
Pagnini,

[143888]D.; Alfinito, F.; Schettini, F.; Luzzatto, L.; Martini, G.: So-
matic-cell

[143889]selection is a major determinant of the blood-cell pheno-
type in heterozygotes

[143890]for glucose-6-phosphate dehydrogenase mutations caus-
ing severe enzyme

[143891]deficiency. Am. J. Hum. Genet. 59: 887–895, 1996.

[143892]

[143893]103. Fiorelli, G.; Manoussakis, C.; Sampietro, M.; Pittalis,
S.; Guglielmino,

[143894]C. R.; Cappellini, M. D.: Different polymorphic variants of
glucose-6-phosphate

[143895]dehydrogenase (G6PD) in Italy. Ann. Hum. Genet. 53:
229–236, 1989.

[143896]

[143897]104. Fite, E.; Morell, F.; Zuazu, J.; Julia, A.; Morera, J.: Leucocyte

[143898]glucose-6-phosphate dehydrogenase deficiency and necrotizing pneumonia. Europ.

[143899]J. Resp. Dis. 64: 150-154, 1983.

[143900]

[143901]105. Francke, U.; Bakay, B.; Connor, J. D.; Coldwell, J. G.; Nyhan,

[143902]W. L.: Linkage relationships of X-linked enzymes glucose-6-phosphate

[143903]dehydrogenase and hypoxanthine guanine phosphoribosyltransferase. Am.

[143904]J. Hum. Genet. 26: 512-522, 1974.

[143905]

[143906]106. Friedman, M. J.; Trager, W.: The biochemistry of resistance

[143907]to malaria. Sci. Am. 244(3): 154-164, 1981.

[143908]

[143909]107. Frigerio, R.; Sole, G.; Olla, N.; Lovicu, M.; Passiu, G.; Carcassi,

[143910]U.: Cagliari II: a new G-6-PD variant. Haematologica 72:

241–243,

[143911]1987.

[143912]

[143913]108. Fujii, H.; Miwa, S.; Takegawa, S.; Takahashi, K.; Hirono, A.;

[143914]Takizawa, T.; Morisaki, T.; Kanno, H.; Taguchi, T.; Okamura, J.:

[143915]Gd(–) Gifu and Gd(–) Fukuoka: two new variants of glucose-6-phosphate

[143916]dehydrogenase found in Japan. Hum. Genet. 66: 276–278, 1984.

[143917]

[143918]109. Fujii, H.; Miwa, S.; Tani, K.; Takegawa, S.; Fujinami, N.; Takahashi,

[143919]K.; Nakayama, S.; Konno, M.; Sato, T.: Glucose 6-phosphate dehydrogenase

[143920]variants: a unique variant (G6PD Kobe) showed an extremely increased

[143921]affinity for galactose 6-phosphate and a new variant (G6PD Sapporo)

[143922]resembling G6PD Pea Ridge. Hum. Genet. 58: 405–407, 1981.

[143923]

[143924]110. Gaetani, G. F.; Galiano, S.; Melani, C.; Miglino, M.;
Forni,

[143925]G. L.; Napoli, G.; Perrone, L.; Ferraris, A. M.: A new glu-
cose-6-phosphate

[143926]dehydrogenase variant with congenital nonspherocytic
hemolytic anemia

[143927](G6PD Genova): biochemical characterization and mo-
saicism expression

[143928]in the heterozygote. Hum. Genet. 84: 337-340, 1990.

[143929]

[143930]111. Gahr, M.; Bornhalm, D.; Schroeter, W.: Biochemische
Eigenschaften

[143931]einer neuen Variante des Glucose-
6-phosphadehydrogenase (G6PD) Mangels

[143932]mit Favismus: G6PD Bielefeld. Klin. Wschr. 55: 379-384,
1977.

[143933]

[143934]112. Gahr, M.; Bornhalm, D.; Schroeter, W.: Haemolytic
anemia due

[143935]to glucose 6-phosphate dehydrogenase (G6PD) deficiency:
demonstration

[143936]of two new biochemical variants, G6PD Hamm and G6PD
Tarsus. Brit.

[143937]J. Haemat. 33: 363–370, 1976.

[143938]

[143939]113. Gahr, M.; Schroeter, W.: Glucose 6–phosphate dehydrogenase (G6PD)

[143940]Hamburg, a new variant with chronic nonspherocytic
hemolytic anemia. Europ.

[143941]J. Clin. Invest. 4: 187–191, 1974.

[143942]

[143943]114. Gahr, M.; Schroeter, W.; Sturzenegger, M.; Bornhalm,
D.; Marti,

[143944]H. R.: Glucose 6–phosphate dehydrogenase (G6PD) deficiency in Switzerland. Helv.

[143945]Paediat. Acta 31: 159–166, 1976.

[143946]

[143947]115. Geerdink, R. A.; Horst, R.; Staal, G. E.: An Iraqi Jewish
family

[143948]with a new red cell glucose 6–phosphate dehydrogenase
variant (Gd–Bagdad)

[143949]and kernicterus. Israel J. Med. Sci. 9: 1040–1043, 1973.

[143950]

[143951]116. Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors-Cayla, M.

[143952]C.; Gillois, M.: Pig gene mapping: synteny, attempt to assign the

[143953]histocompatibility complex (SLA). (Abstract) Cytogenet. Cell Genet. 25:

[143954]159, 1979.

[143955]

[143956]117. Gomez-Gallego, F.; Garrido-Pertierra, A.; Bautista, J. M.: Structural

[143957]defects underlying protein dysfunction in human glucose-6-phosphate

[143958]dehydrogenase A- deficiency. J. Biol. Chem. 275: 9256-9262, 2000.

[143959]

[143960]118. Gonzalez, R.; Estrada, M.; Garcia, M.; Gutierrez, A.: G6PD Ciudad

[143961]de la Habana: a new slow variant with deficiency found in a Cuban

[143962]family. Hum. Genet. 55: 133-135, 1980.

[143963]

[143964]119. Gonzalez, R.; Wade, M.; Estrada, M.; Svarch, E.;
Colombo, B.

[143965]: G6PD Pinar del Rio: a new variant discovered in a Cuban
family. Biochem.

[143966]Genet. 15: 909–913, 1977.

[143967]

[143968]120. Goss, S. J.; Harris, H.: Gene transfer by means of cell
fusion.

[143969]I. Statistical mapping of the human X–chromosome by
analysis of radiation

[143970]induced gene segregation. J. Cell Sci. 25: 17–37, 1977.

[143971]

[143972]121. Gourdin, D.; Vergnes, H.; Bouloux, C.; Ruffie, J.;
Gherardi,

[143973]M.: Polymorphism of erythrocyte G6PD in the baboon. Am.
J. Phys.

[143974]Anthrop. 37: 281–288, 1972.

[143975]

[143976]122. Gray, G. R.; Stamatoyannopoulos, G.; Naiman, S. C.;
Kliman, M.

[143977]R.; Klebanoff, S. J.; Austin, T.; Yoshida, A.; Robinson, G. C.
G.

[143978]: Neutrophil dysfunction, chronic granulomatous disease,
and non-spherocytic

[143979]haemolytic anaemia caused by complete deficiency of glu-
cose-6-phosphate

[143980]dehydrogenase. Lancet II: 530-534, 1973.

[143981]

[143982]123. Grossman, A.; Ramanathan, K.; Justice, P.; Gordon, J.;
Shahidi,

[143983]N. T.; Hsia, D. Y. Y.: Congenital nonspherocytic hemolytic
anemia

[143984]associated with erythrocyte G-6-PD deficiency in a Negro
family. Pediatrics 37:

[143985]624-629, 1966.

[143986]

[143987]124. Gutierrez, A.; Garcia, M.; Estrada, M.; Quintero, I.;
Gonzalez,

[143988]R.: Glucose-6-phosphate dehydrogenase (G6PD) Guan-
tanamo and G6PD

[143989]Caujeri: two new glucose-6-phosphate dehydrogenase-defi-
cient variants

[143990]found in Cuba. Biochem. Genet. 25: 231–238, 1987.

[143991]

[143992]125. Hall, K.; Schreeder, M. T.; Prchal, J. T.: G6PD

Huntsville:

[143993]a new glucose–6–phosphate dehydrogenase associated
with chronic hemolytic

[143994]anemia. Hum. Genet. 79: 90–91, 1988.

[143995]

[143996]126. Harkonen, M.; Vuopio, P.: Red cell glucose–

6–phosphate dehydrogenase

[143997]deficiency in Finland. Ann. Clin. Res. 6: 187–197, 1974.

[143998]

[143999]127. Harley, J. D.; Agar, N. S.; Yoshida, A.: Glucose

6–phosphate

[144000]dehydrogenase variant Gd(+) Alexandra associated with
neonatal jaundice

[144001]and Gd(–) Camperdown in a young man with lamellar
cataracts. J. Lab.

[144002]Clin. Med. 91: 295–300, 1978.

[144003]

[144004]128. Helge, H.; Borner, K.: Kongenitale nichtsphaerozytare
haemolytische

[144005]Anaemie, Katarakt und Glucose-
6-phosphat-dehydrogenase-mangel. Dtsch.

[144006]Med. Wschr. 91: 1584-1589, 1966.

[144007]

[144008]129. Hirono, A.; Beutler, E.: Alternative splicing of human
glucose-6-phosphate

[144009]dehydrogenase messenger RNA in different tissues. J.
Clin. Invest. 83:

[144010]343-346, 1989.

[144011]

[144012]130. Hirono, A.; Beutler, E.: Molecular cloning and nu-
cleotide sequence

[144013]of cDNA for human glucose-6-phosphate dehydrogenase
variant A(-). Proc.

[144014]Nat. Acad. Sci. 85: 3951-3954, 1988.

[144015]

[144016]131. Hirono, A.; Fujii, H.; Shima, M.; Miwa, S.: G6PD Nara:
a new

[144017]class 1 glucose-6-phosphate dehydrogenase variant with

an eight amino

[144018]acid deletion. Blood 82: 3250–3252, 1998.

[144019]

[144020]132. Hirono, A.; Kawate, K.; Honda, A.; Fujii, H.; Miwa, S.:

A single

[144021]mutation 202G–A in the human glucose–6–phosphate de–
hydrogenase gene

[144022](G6PD) can cause acute hemolysis by itself. (Letter) Blood
99: 1498

[144023]only, 2002.

[144024]

[144025]133. Hirono, A.; Kuhl, W.; Gelbart, T.; Forman, L.; Fair–
banks, V.

[144026]F.; Beutler, E.: Identification of the binding domain for
NADP(+)

[144027]of human glucose–6–phosphate dehydrogenase by se–
quence analysis of

[144028]mutants. Proc. Nat. Acad. Sci. 86: 10015–10017, 1989.

[144029]

[144030]134. Hitzeroth, H. W.; Bender, K.: Age–dependency of so–
matic selection

[144031]in South African Negro G-6-PD heterozygotes. Hum.

Genet. 58: 338-343,

[144032]1981.

[144033]

[144034]135. Honig, G. R.; Habacon, E.; Vida, L. N.; Matsumoto, F.;

Beutler,

[144035]E.: Three new variants of glucose-6-phosphate dehydro-
genase associated

[144036]with chronic nonspherocytic hemolytic anemia: G6PD Lin-
coln Park, G6PD

[144037]Arlington Heights, and G6PD West Town. Am. J. Hemat. 6:
353-360,

[144038]1979.

[144039]

[144040]136. Hook, E. B.; Stamatoyannopoulos, G.; Yoshida, A.;

Motulsky, A.

[144041]G.: Glucose-6-phosphate dehydrogenase Madrona: a slow
electrophoretic

[144042]glucose-6-phosphate dehydrogenase variant with kinetic
characteristics

[144043]similar to those of normal type. J. Lab. Clin. Med. 72:
404-409,

[144044]1968.

[144045]

[144046]137. Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Benne, F.: Cattle

[144047]gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

[144048]Cell Genet. 25: 165–166, 1979.

[144049]

[144050]138. Howell, E. B.; Nelson, A. J.; Jones, O. W.: A new G–
6–PD variant

[144051]associated with chronic non–spherocytic haemolytic
anaemia in a Negro

[144052]family. J. Med. Genet. 9: 160–164, 1972.

[144053]

[144054]139. Hutz, M. H.; Yoshida, A.; Salzano, F. M.: Three rare
G–6–PD

[144055]variants from Porto Alegre, Brazil. Hum. Genet. 39:
191–197, 1977.

[144056]

[144057]140. Ishwad, C. S.; Naik, S. N.: A new glucose–
6–phosphate dehydrogenase

[144058]variant (G-6-PD Kalyan) found in a Koli family. Hum.

Genet. 66:

[144059]171-175, 1984.

[144060]

[144061]141. Johnson, G. J.; Kaplan, M. E.; Beutler, E.: G6PD Long

Prairie:

[144062]a new mutant exhibiting normal sensitivity to inhibition by

NADPH

[144063]and accompanied by nonspherocytic hemolytic anemia.

Blood 49: 247-251,

[144064]1977.

[144065]

[144066]142. Johnston, P. G.; VandeBerg, J. L.; Sharman, G. B.: In-

heritance

[144067]of erythrocyte glucose 6-phosphate dehydrogenase in the

red-necked

[144068]wallaby, *Macropus rufogriseus* (Desmarest) consistent

with paternal

[144069]X inactivation. Biochem. Genet. 13: 235-242, 1975.

[144070]

[144071]143. Junien, C.; Kaplan, J.-C.; Meienhofer, M. C.; Maigret,

P.; Sender,

[144072]A.: G6PD Baudelocque: a new unstable variant characterized in cultured

[144073]fibroblasts. Enzyme 18: 48–59, 1974.

[144074]

[144075]144. Kaeda, J. S.; Chhotray, G. P.; Ranjit, M. R.; Bautista, J. M.;

[144076]Reddy, P. H.; Stevens, D.; Naidu, J. M.; Britt, R. P.; Vulliamy, T.

[144077]J.; Luzzatto, L.; Mason, P. J.: A new glucose–6–phosphate dehydrogenase

[144078]variant, G6PD Orissa (44 ala–gly), is the major polymorphic variant

[144079]in tribal populations in India. Am. J. Hum. Genet. 57: 1335–1341,

[144080]1995.

[144081]

[144082]145. Kageoka, T.; Satoh, C.; Goriki, K.; Fujita, M.; Neriishi, S.;

[144083]Yamamura, K.; Kaneko, J.; Masunari, N.: Electrophoretic variants

[144084]of blood proteins in Japanese. IV. Prevalence and enzymo–

logic characteristics

[144085]of G6PD variants in Hiroshima and Nagasaki. Hum. Genet.
70: 101–108,

[144086]1985.

[144087]

[144088]146. Kahn, A.; Bernard, J.–F.; Cottreau, D.; Mazie, J.;
Boivin, P.

[144089]: Gd(–) Abrami, a deficient G6PD variant with hemizygous
expression

[144090]in blood cells of a woman with primary myelofibrosis. Hu–
mangenetik 30:

[144091]41–46, 1975.

[144092]

[144093]147. Kahn, A.; Boivin, P.; Hakim, J.; Lagneau, J.: Hetero–
geneite

[144094]des glucose–6–phosphate dehydrogenase erythrocytaire
deficitaires

[144095]dans la race noire: etude cinetique et description de deux
nouvelles

[144096]variants Gd(–) Dakar et Gd(–) Mali. Nouv. Rev. Franc.
Hemat. 11:

[144097]741–758, 1971.

[144098]

[144099]148. Kahn, A.; Boivin, P.; Lagneau, J.: Phenotypes de la
glucose-6-phosphate

[144100]dehydrogenase erythrocytaire dans la race noire. Human-
genetik 18:

[144101]261-270, 1973.

[144102]

[144103]149. Kahn, A.; Boulard, M.; Hakim, J.; Schaison, G.; Boivin,
P.; Bernard,

[144104]J.: Anemie hemolytique congenitale non spherocytaire par
deficit

[144105]en glucose 6-phosphate dehydrogenase erythrocytaire:
description de

[144106]deux nouvelles variants: Gd(-) Saint Louis (Paris) et Gd(-)
Hayem. Nouv.

[144107]Rev. Franc. Hemat. 14: 587-600, 1974.

[144108]

[144109]150. Kahn, A.; Dao, C.; Cottreau, D.; Bilski-Pasquier, G.:
'Gd(-)

[144110]Hotel Dieu': a new G-6PD variant with chronic hemolysis
in a Negro

[144111]patient from Senegal. Hum. Genet. 39: 353–358, 1977.

[144112]

[144113]151. Kahn, A.; Exters, A.; Habedank, M.: Gd(–) Aachen, a
new variant

[144114]of deficient glucose–6–phosphate dehydrogenase. Hu-
mangenetik 32:

[144115]171–180, 1976.

[144116]

[144117]152. Kahn, A.; Hakim, J.; Cottreau, D.; Boivin, P.: Gd (–)
Matam.

[144118]An African glucose 6–phosphate dehydrogenase variant
with enzyme deficiency.

[144119]Biochemical and immunological properties in various
hemopoietic tissues. Clin.

[144120]Chim. Acta 59: 183–190, 1975.

[144121]

[144122]153. Kahn, A.; North, M. L.; Cottreau, D.; Giron, G.; Lang,
J. M.

[144123]: G6PD Vientiane: a new glucose–6–phosphate dehydro-
genase variant

[144124]with increased stability. Hum. Genet. 43: 85–89, 1978.

[144125]

[144126]154. Kahn, A.; North, M. L.; Messer, J.; Boivin, P.: G-6PD
'Ankara':

[144127]a new G-6PD variant with deficiency found in a Turkish
family. Humangenetik 27:

[144128]247-250, 1975.

[144129]

[144130]155. Kanno, H.; Huang, I.-Y.; Kan, Y. W.; Yoshida, A.: Two
structural

[144131]genes on different chromosomes are required for encod-
ing the major

[144132]subunit of human red cell glucose-6-phosphate dehydro-
genase. Cell 58:

[144133]595-606, 1989.

[144134]

[144135]156. Kanno, H.; Takano, T.; Fujii, H.; Tani, K.; Morisaki, T.;
Hirono,

[144136]A.; Kumakawa, T.; Ogura, H.; Takahashi, K.; Tsutsumi, H.;
Miwa, S.

[144137]: A new glucose-6-phosphate variant (G6PD Iwate) associ-
ated with congenital

[144138]non-spherocytic hemolytic anemia. Acta Haemat. Jpn. 51:
715-719,
[144139]1987.

[144140]

[144141]157. Kaplan, J. C.; Hanzlickova-Leroux, A.; Nicholas, A.
M.; Rosa,

[144142]R.; Weiler, C.; Lepercq, G.: A new glucose-6-phosphate
dehydrogenase

[144143]variant (G6PD Port-Royal). Enzyme 12: 25-32, 1971.

[144144]

[144145]158. Kaplan, J. C.; Rosa, R.; Seringe, P.; Hoeffel, J. C.: Le
polymorphisme

[144146]genetique de la glucose-6-phosphate deshydrogenase
erythrocytaire

[144147]chez l'homme. II. Etude d'une nouvelle variete a activite
diminuee:

[144148]le type 'Kabyle.'. Enzym. Biol. Clin. 8: 332-340, 1967.

[144149]

[144150]159. Kaplan, M.; Hammerman, C.; Vreman, H. J.; Steven-
son, D. K.; Beutler,

[144151]E.: Acute hemolysis and severe neonatal hyperbilirubine-

mia in glucose-6-phosphate

[144152]dehydrogenase deficient heterozygotes. J. Pediat. 139:
137-140,

[144153]2001.

[144154]

[144155]160. Kaplan, M.; Renbaum, P.; Levy-Lahad, E.; Hammer-
man, C.; Lahad,

[144156]A. Beutler, E.: Gilbert syndrome and glucose-6-phosphate
dehydrogenase

[144157]deficiency: a dose-dependent genetic interaction crucial
to neonatal

[144158]hyperbilirubinemia. Proc. Nat. Acad. Sci. 94:
12128-12132, 1997.

[144159]

[144160]161. Kappas, A.; Drummond, G. S.; Valaes, T.: A single
dose of Sn-mesoporphyrin

[144161]prevents development of severe hyperbilirubinemia in
glucose-6-phosphate

[144162]dehydrogenase-deficient newborns. Pediatrics 108:
25-30, 2001.

[144163]

[144164]162. Karadsheh, N. S.; Awidi, A. S.; Tarawneh, M. S.: Two

new glucose-6

[144165]phosphate dehydrogenase (G6PD) variants associated with hemolytic

[144166]anemia: G6PD Amman-1 and G6PD Amman-2. Am. J. Hemat. 22: 185-192,

[144167]1986.

[144168]

[144169]163. Kay, A. C.; Kuhl, W.; Prchal, J.; Beutler, E.: The origin of

[144170]glucose-6-phosphate-dehydrogenase (G6PD) polymorphisms in African-Americans. Am.

[144171]J. Hum. Genet. 50: 394-398, 1992.

[144172]

[144173]164. Kirkman, H. N.; Kidson, C.; Kennedy, M.: Variants of human glucose-6-phosphate

[144174]dehydrogenase. Studies of samples from New Guinea. In: Beutler, E.

[144175]: Hereditary Disorders of Erythrocyte Metabolism. New York: Grune

[144176]and Stratton (pub.) 1968. Pp. 126-145.

[144177]

[144178]165. Kirkman, H. N.; Lie-Injo, L. E.: Variants of glucose
6-phosphate

[144179]dehydrogenase in Indonesia. Nature 221: 959, 1969.

[144180]

[144181]166. Kirkman, H. N.; McCurdy, P. R.; Naiman, J. L.: Func-
tionally

[144182]abnormal glucose-6-phosphate dehydrogenases. Cold
Spring Harbor Symp.

[144183]Quant. Biol. 29: 391-398, 1964.

[144184]

[144185]167. Kirkman, H. N.; Ramot, B.; Lee, J. T.: Altered aggre-
gational

[144186]properties in a genetic variant of human glucose-
6-phosphate dehydrogenase. Biochem.

[144187]Genet. 3: 137-150, 1969.

[144188]

[144189]168. Kirkman, H. N.; Riley, H. D., Jr.: Congenital nonsphe-
rocytic

[144190]hemolytic anemia. Am. J. Dis. Child. 102: 313-320, 1961.

[144191]

[144192]169. Kirkman, H. N.; Rosenthal, I. M.; Simon, E. B.; Carson, P. E.;

[144193]Brinson, A. G.: 'Chicago I' variant of glucose-6-phosphate dehydrogenase

[144194]in congenital hemolytic disease. J. Lab. Clin. Med. 63: 715-725,

[144195]1964.

[144196]

[144197]170. Kirkman, H. N.; Schettini, E.; Pickard, B. M.: Mediter-ranean

[144198]variant of glucose-6-phosphate dehydrogenase. J. Lab. Clin. Med. 63:

[144199]726-735, 1964.

[144200]

[144201]171. Kirkman, H. N.; Simon, E. R.; Pickard, B. M.: Seattle variant

[144202]of glucose-6-phosphate dehydrogenase. J. Lab. Clin. Med. 66: 834-840,

[144203]1965.

[144204]

[144205]172. Kissin, C.; Cotte, J.: Etude d'un variant de glucose-

6-phosphate

[144206]deshydrogenase: I B type Constantine. Enzyme 11:
277-284, 1970.

[144207]

[144208]173. Kitao, T.; Ito, K.; Hattori, K.; Matsuki, T.; Yoneyama,
Y.:

[144209]G6PD Kanazawa: a new variant of glucose-6-phosphate
dehydrogenase

[144210]associated with congenital nonspherocytic hemolytic ane-
mia. Acta

[144211]Haemat. 68: 131-135, 1982.

[144212]

[144213]174. Kojima, H.: Congenital nonspherocytic hemolytic dis-
ease (CNHD)

[144214]due to a G-6-PD variant: G-6-PD Kyoto. Acta Haemat.
Jpn. 35: 32-38,

[144215]1972.

[144216]

[144217]175. Koliakos, G.; Kalomenopoulou, M.; Grammatikos, P.;
Dimitriadou,

[144218]A.; Kouzi-Koliakos, K.; Zacharaki, R.; Skaragas, G.; Kokka,

A.; Trakatellis,

[144219]A.: A new glucose 6-phosphate dehydrogenase variant
(G6PD Thessaloniki)

[144220]in a patient with idiopathic myelofibrosis. Hum. Hered. 39:
141-149,

[144221]1989.

[144222]

[144223]176. Krasnopol'skaya, K. D.; Bochkov, N. P.: Genetic het-
erogeneity

[144224]of hereditary enzymopathies. Vestn. Akad. Med. Nauk.
SSSR 9: 56-64,

[144225]1982.

[144226]

[144227]177. Krasnopol'skaya, K. D.; Shatskaya, T. L.; Filippov, I.
K.; Annenkov,

[144228]G. A.; Zakharova, T. V.; Mekhtiev, N. K.; Movsum-Zade, K.
M.: Genetic

[144229]heterogeneity of G6PD deficiency: study of mutant alleles
in Shekii

[144230]district of Azerbaijan. Genetika 13: 1455-1461, 1977.

[144231]

[144232]178. Kumakawa, T.; Suzuki, S.; Fujii, H.; Miwa, S.: Frequency of
[144233]glucose 6-phosphate dehydrogenase (G6PD) deficiency in
Tokyo and a
[144234]new variant: G6PD Musashino. *Acta Haemat. Jpn.* 50:
25-28, 1987.

[144235]

[144236]179. Kurdi-Haidar, B.; Mason, P. J.; Berrebi, A.; Ankra-
Badu, G.;
[144237]Al-Ali, A.; Oppenheim, A.; Luzzatto, L.: Origin and spread
of the
[144238]glucose-6-phosphate dehydrogenase variant
(G6PD-Mediterranean) in
[144239]the Middle East. *Am. J. Hum. Genet.* 47: 1013-1019,
1990.

[144240]

[144241]180. Kwiatkowska, J.; Kacprzak-Bergman, I.: New erythro-
cyte glucose
[144242]6-phosphate dehydrogenase variant. *Acta Haemat.* 46:
188-192, 1971.

[144243]

[144244]181. Kwok, C. J.; Martin, A. C. R.; Au, S. W. N.; Lam, V. M.

S.:

[144245]G6PDdb, an integrated database of glucose-6-phosphate dehydrogenase

[144246](G6PD) mutations. Hum. Mutat. 19: 217-224, 2002.

[144247]

[144248]182. Lee, K. T.; Thomas, W. A.; Janakidevi, K.; Kroms, M.; Reiner,

[144249]J. M.; Borg, K. Y.: Mosaicism in female hybrid hares heterozygous

[144250]for glucose-6-phosphate dehydrogenase (G-6-PD). I. General properties

[144251]of a hybrid hare model with special reference to atherogenesis. Exp.

[144252]Molec. Path. 34: 191-201, 1981.

[144253]

[144254]183. Lenzerini, L.; Meera Khan, P.; Filippi, G.; Rattazzi, M. C.;

[144255]Rat, A. K.: Characterization of glucose-6-phosphate dehydrogenase

[144256]variants. I. Occurrence of a G6PD Seattle-like variant in Sardinia

[144257]and its interaction with G6PD Mediterranean variant. Am.

J. Hum.

[144258]Genet. 21: 142–153, 1969.

[144259]

[144260]184. Lisker, R.; Linares, C.; Motulsky, A. G.: Glucose–
6–phosphate

[144261]dehydrogenase Mexico, a new variant with enzyme defi–
ciency, abnormal

[144262]mobility and absence of hemolysis. J. Lab. Clin. Med. 29:
788–793,

[144263]1972.

[144264]

[144265]185. Lisker, R.; Perez–Briceno, R.; Beutler, E.: A new glu–
cose–6–phosphate

[144266]dehydrogenase variant, Gd(–) Tepic, characterized by
moderate enzyme

[144267]deficiency and mild episodes of hemolytic anemia. Hum.
Genet. 69:

[144268]19–21, 1985.

[144269]

[144270]186. Lisker, R.; Perez–Briceno, R.; Rave, V.; Yoshida, A.:
Glucosa–6–fosfato

[144271]deshidrogenasa Gd(-) Distrito Federal: nueva variante asociada a deficiencia

[144272]enzimatica moderada y anemia hemolitica ocasional. Rev. Invest. Clin. 33:

[144273]209-211, 1981.

[144274]

[144275]187. Lisker, R.; Perez Briceno, R.; Agrilar, L.; Yoshida, A.: A variant

[144276]glucose-6-phosphate dehydrogenase Gd(-) Chiapas associated with moderate

[144277]enzyme deficiency and occasional hemolytic anemia. Hum. Genet. 43:

[144278]81-84, 1978.

[144279]

[144280]188. Lisker, R.; Perez Briceno, R.; Zavala, C.; Navarrette, J. I.;

[144281]Wessels, M.; Yoshida, A.: A glucose 6-phosphate dehydrogenase Gd(-)

[144282]Castilla variant characterized by mild deficiency associated with

[144283]drug-induced hemolytic anemia. J. Lab. Clin. Med. 90: 754-759, 1977.

[144284]

[144285]189. Liu, Y.; Phelan, J.; Go, R. C. P.; Prchal, J. F.; Prchal, J.

[144286]T.: Rapid determination of clonality by detection of two
closely-linked

[144287]X chromosome exonic polymorphisms using allele-spe-
cific PCR. J. Clin.

[144288]Invest. 99: 1984-1990, 1997.

[144289]

[144290]190. Long, W. K.; Kirkman, H. N.; Sutton, H. H.: Elec-
trophoretically

[144291]slow variants of glucose-6-phosphate dehydrogenase
from red cells

[144292]of Negroes. J. Lab. Clin. Med. 65: 81-87, 1965.

[144293]

[144294]191. Longo, L.; Vanegas, O. C.; Patel, M.; Rosti, V.; Li, H.;
Waka,

[144295]J.; Merghoub, T.; Pandolfi, P. P.; Notaro, R.; Manova, K.;
Luzzatto,

[144296]L.: Maternally transmitted severe glucose 6-phosphate
dehydrogenase

[144297]deficiency is an embryonic lethal. EMBO J. 21: 4229-4239,

2002.

[144298]

[144299]192. Luzzatto, L.: Genetic heterogeneity and pathophysiology of G6PD

[144300]deficiency. Brit. J. Haemat. 28: 151–156, 1974.

[144301]

[144302]193. Luzzatto, L.: Personal Communication. London, England 4/1990.

[144303]

[144304]194. Luzzatto, L.; Afolayam, A.: Enzyme properties of different types

[144305]of human erythrocyte glucose-6-phosphate dehydrogenase with characterization

[144306]of two new genetic variants. J. Clin. Invest. 47: 1833–1842, 1968.

[144307]

[144308]195. Luzzatto, L.; Martini, G.: X-Linked Wiskott–Aldrich syndrome

[144309]in a girl. (Letter) New Eng. J. Med. 338: 1850–1851, 1998.

[144310]

[144311]196. Luzzatto, L.; Usanga, E. A.; Bienzle, U.; Esan, G. F. J.;

Fusuan,

[144312]F. A.: Imbalance in X-chromosome expression: evidence
for a human

[144313]X-linked gene affecting growth of hemopoietic cells. Sci-
ence 205:

[144314]1418–1420, 1979.

[144315]

[144316]197. Luzzatto, L.; Usanga, E. A.; Reddy, S.: Glucose-
6-phosphate

[144317]dehydrogenase deficient red cells: resistance to infection
by malarial

[144318]parasites. Science 164: 839–842, 1969.

[144319]

[144320]198. MacDonald, D.; Town, M.; Mason, P.; Vulliamy, T.;
Luzzatto, L.;

[144321]Goff, D. K.: Deficiency in red blood cells. (Letter) Nature
350:

[144322]115, 1991.

[144323]

[144324]199. Maeda, M.; Constantoulakis, P.; Chen, C.-S.; Stama-
toyannopoulos,

[144325]G.; Yoshida, A.: Molecular abnormalities of a human glucose-6-phosphate

[144326]dehydrogenase variant associated with undetectable enzyme activity

[144327]and immunologically cross-reacting material. Am. J. Hum. Genet. 51:

[144328]386-395, 1992.

[144329]

[144330]200. Mallouh, A. A.; Abu-Osba, Y. K.: Bacterial infections in children

[144331]with glucose-6-phosphate dehydrogenase deficiency. J. Pediat. 111:

[144332]850-852, 1987.

[144333]

[144334]201. Mamlok, R. J.; Mamlok, V.; Mills, G. C.; Daeschner, C. W., III;

[144335]Schmalstieg, F. C.; Anderson, D. C.: Glucose-6-phosphate dehydrogenase

[144336]deficiency, neutrophil dysfunction and Chromobacterium violaceum sepsis. J.

[144337]Pediat. 111: 852-854, 1987.

[144338]

[144339]202. Mamlok, R. J.; Mills, G. C.; Goldblum, R. M.;

Daeschner, C. W.

[144340]: Glucose-6-phosphate dehydrogenase Beaumont: a new variant with severe

[144341]enzyme deficiency and chronic nonspherocytic hemolytic anemia. Enzyme 34:

[144342]15-21, 1985.

[144343]

[144344]203. Mandelli, F.; Amadori, S.; DeLaurenzi, A.; Kahn, A.;

Isacchi,

[144345]G.; Papa, G.: Glucose-6-phosphate dehydrogenase Vellettri: a new variant

[144346]with reduced activity in a patient with congenital nonspherocytic

[144347]haemolytic anemia. Acta Haemat. 57: 121-126, 1977.

[144348]

[144349]204. Marks, P. A.; Banks, J.; Gross, R.: Genetic heterogeneity of

[144350]glucose-6-phosphate dehydrogenase deficiency. Nature 194: 454-456,

[144351]1962.

[144352]

[144353]205. Martin, S. K.; Miller, L. H.; Alling, D.; Okoye, V. C.;
Esan,

[144354]G. J. F.; Osunkoya, B. O.; Deane, M.: Severe malaria and
glucose-6-phosphate-dehydrogenase

[144355]deficiency: a reappraisal of the malaria-G6PD hypothesis.
Lancet I:

[144356]524-526, 1979.

[144357]

[144358]206. Martini, G.; Toniolo, D.; Vulliamy, T.; Luzzatto, L.;
Dono, R.;

[144359]Viglietto, G.; Paonessa, G.; D'Urso, M.; Persico, M. G.:
Structural

[144360]analysis of the X-linked gene encoding human glucose
6-phosphate dehydrogenase. EMBO

[144361]J. 5: 1849-1855, 1986.

[144362]

[144363]207. Mason, P. J.: New insights into G6PD deficiency.
(Annotation) Brit.

[144364]J. Haemat. 94: 585-591, 1996.

[144365]

[144366]208. Mason, P. J.; Stevens, D. J.; Luzzatto, L.; Brenner, S.; Aparicio,

[144367]S.: Genomic structure and sequence of the Fugu rubripes glucose-6-phosphate

[144368]dehydrogenase gene (G6PD). Genomics 26: 587-591, 1995.

[144369]

[144370]209. McCann, S. R.; Smithwick, A. M.; Temperley, I. J.; Tipton, K.

[144371]: G6PD (Dublin): chronic non-spherocytic haemolytic anaemia resulting

[144372]from glucose-6-phosphate dehydrogenase deficiency in an Irish kindred. J.

[144373]Med. Genet. 17: 191-193, 1980.

[144374]

[144375]210. McCurdy, P. R.: Use of genetic linkage for the detection of

[144376]female carriers of hemophilia. New Eng. J. Med. 285: 218-219, 1971.

[144377]

[144378]211. McCurdy, P. R.; Blackwell, R. Q.; Todd, D.; Tso, S. C.;
Tuchinda,

[144379]S.: Further studies on glucose-6-phosphate dehydroge-
nase deficiency

[144380]in Chinese subjects. J. Lab. Clin. Med. 75: 788-797, 1970.

[144381]

[144382]212. McCurdy, P. R.; Kamel, K.; Selim, O.: Heterogeneity of
red cell

[144383]glucose 6-phosphate dehydrogenase (G6PD) deficiency in
Egypt. J.

[144384]Lab. Clin. Med. 84: 673-680, 1974.

[144385]

[144386]213. McCurdy, P. R.; Kirkman, H. N.; Naiman, J. L.; Jim, R.
T. S.;

[144387]Pickard, B. M.: A Chinese variant of glucose-6-phosphate
dehydrogenase. J.

[144388]Lab. Clin. Med. 67: 374-385, 1966.

[144389]

[144390]214. McCurdy, P. R.; Mahmood, L.: Red cell glucose-
6-phosphate dehydrogenase

[144391]deficiency in Pakistan. J. Lab. Clin. Med. 76: 943-948,

1970.

[144392]

[144393]215. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.: Variants of

[144394]glucose-6-phosphate dehydrogenase (G-6-PD) associated with G-6-PD

[144395]deficiency in Puerto Ricans. J. Lab. Clin. Med. 82: 432-437, 1973.

[144396]

[144397]216. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.; Conrad, M. E.

[144398]: Variants of glucose-6-phosphate dehydrogenase (G-6-PD) associated

[144399]with G-6-PD deficiency in Puerto Ricans. J. Lab. Clin. Med. 82:

[144400]432-437, 1973.

[144401]

[144402]217. Meloni, T.; Carta, F.; Forteleoni, G.; Carta, A.; Ena, F.; Meloni,

[144403]G. F.: Glucose 6-phosphate dehydrogenase deficiency and cataract

[144404]of patients in northern Sardinia. Am. J. Ophthal. 110:

661–664,

[144405]1990.

[144406]

[144407]218. Mentzer, W. C., Jr.; Warner, R.; Addiego, J.; Smith, B.;
Walter,

[144408]T.: G6PD San Francisco: a new variant of glucose–
6–phosphate dehydrogenase

[144409]associated with congenital nonspherocytic hemolytic ane–
mia. Blood 55:

[144410]195–198, 1980.

[144411]

[144412]219. Mesbah–Namin, S. A.; Sanati, M. H.; Mowjoodi, A.;
Mason, P. J.;

[144413]Vulliamy, T. J.; Noori–Daloii, M. R.: Three major glucose–
6–phosphate

[144414]dehydrogenase–deficient polymorphic variants identified
in Mazandaran

[144415]state of Iran. Brit. J. Haemat. 117: 763–764, 2002.

[144416]

[144417]220. Miller, D. R.; Wollman, M. R.: A new variant of glu–
cose 6–phosphate

[144418]dehydrogenase deficiency hereditary hemolytic anemia,
G6PD Cornell:

[144419]erythrocyte, leukocyte and platelet studies. Blood 44:
277-284,

[144420]1974.

[144421]

[144422]221. Mills, G. C.; Alperin, J. B.; Trimmer, K. B.: Studies on
variant

[144423]glucose-6-phosphate dehydrogenase: G6PD Fort Worth.
Biochem. Med. 13:

[144424]264-275, 1975.

[144425]

[144426]222. Milner, G.; Delamore, I. W.; Yoshida, A.: G-6-PD
Manchester:

[144427]a new variant associated with chronic nonspherocytic
hemolytic anemia. Blood 43:

[144428]271-276, 1974.

[144429]

[144430]223. Miwa, S.; Fujii, H.: Molecular basis of erythroenzy-
mopathies

[144431]associated with hereditary hemolytic anemia: tabulation of

mutant

[144432]enzymes. Am. J. Hemat. 51: 122–132, 1996.

[144433]

[144434]224. Miwa, S.; Fujii, H.; Nakashima, K.; Miura, Y.; Yamada, K.; Hagiwara,

[144435]T.; Fukuda, M.: Three new electrophoretically normal glucose–6–phosphate

[144436]dehydrogenase variants associated with congenital nonspherocytic hemolytic

[144437]anemia found in Japan: G6PD Ogikubo, Yokohama, and Akita. Hum. Genet. 45:

[144438]11–17, 1978.

[144439]

[144440]225. Miwa, S.; Fujii, H.; Nakatsuji, T.; Ishida, Y.; Oda, E.; Kaneto,

[144441]A.; Motokawa, M.; Ariga, Y.; Fukuchi, S.; Sasai, S.; Hiraoka, K.;

[144442]Kashii, H.; Kodama, T.: Four new electrophoretically slow-moving

[144443]glucose–6–phosphate dehydrogenase variants associated with congenital

[144444]nonspherocytic hemolytic anemia found in Japan: Gd(–)

Kurume, Gd(-)

[144445]Fukushima, Gd(-) Yamaguchi, and Gd(-) Wakayama. Am. J. Hemat. 5:

[144446]131-138, 1978.

[144447]

[144448]226. Miwa, S.; Nakashima, K.; Ono, J.; Fujii, H.; Suzuki, E.: Three

[144449]glucose 6-phosphate dehydrogenase variants found in Japan. Hum. Genet. 36:

[144450]327-334, 1977.

[144451]

[144452]227. Miwa, S.; Ono, J.; Nakashima, K.; Abe, S.; Kageoka, T.; Shinohara,

[144453]K.; Isobe, J.; Yamaguchi, H.: Two new glucose 6-phosphate dehydrogenase

[144454]variants associated with congenital nonspherocytic hemolytic anemia

[144455]found in Japan: Gd(-) Tokushima and Gd(-) Tokyo. Am. J. Hemat. 1:

[144456]433-442, 1976.

[144457]

[144458]228. Modiano, G.; Battistuzzi, G.; Esan, G. J. F.; Testa, U.;
Luzzatto,

[144459]L.: Genetic heterogeneity of 'normal' human erythrocyte
glucose-6-phosphate

[144460]dehydrogenase: an isoelectrophoretic polymorphism.
Proc. Nat. Acad.

[144461]Sci. 76: 852-856, 1979.

[144462]

[144463]229. Mohrenweiser, H. W.; Neel, J. V.: Frequency of ther-
mostability

[144464]variants: estimation of total 'rare' variant frequency in hu-
man populations. Proc.

[144465]Nat. Acad. Sci. 78: 5729-5733, 1981.

[144466]

[144467]230. Morelli, A.; Benatti, U.; Guida, L.; De Flora, A.: G6PD
Cagliari:

[144468]a new low activity glucose 6-phosphate dehydrogenase
variant characterized

[144469]by enhanced intracellular lability. Hum. Genet. 66: 62-65,
1984.

[144470]

[144471]231. Morisaki, T.; Fujii, H.; Takegawa, S.; Tani, K.; Hirono, A.;

[144472]Takizawa, T.; Takahashi, K.; Shinogi, M.; Teshirogi, T.; Miwa, S.

[144473]: G6PD Sendagi: a new glucose-6-phosphate dehydrogenase variant associated

[144474]with congenital hemolytic anemia. Hum. Genet. 65: 214-215, 1983.

[144475]

[144476]232. Nafa, K.; Reghis, A.; Osmani, N.; Baghli, L.; Benabadji, M.;

[144477]Kaplan, J.-C.; Vulliamy, T. J.; Luzzatto, L.: G6PD Aures: a new mutation

[144478](48 ile-to-thr) causing mild G6PD deficiency is associated with favism. Hum.

[144479]Molec. Genet. 2: 81-82, 1993.

[144480]

[144481]233. Nagel, R. L.; Ranney, H. M.: Genetic epidemiology of structural

[144482]mutations of the beta-globin gene. Semin. Hemat. 27: 342-359, 1990.

[144483]

[144484]234. Nakai, T.; Yoshida, A.: G6PD Heian. A glucose-6-phosphate dehydrogenase

[144485]variant associated with hemolytic anemia found in Japan. Clin. Chim.

[144486]Acta 51: 199-203, 1974.

[144487]

[144488]235. Nakashima, K.; Ono, J.; Abe, S.; Miwa, S.; Yoshida, A.: G6PD

[144489]Ube: a glucose 6-phosphate dehydrogenase variant found in four unrelated

[144490]Japanese families. Am. J. Hum. Genet. 29: 24-30, 1977.

[144491]

[144492]236. Nakatsuji, T.; Miwa, S.: Incidence and characteristics of glucose-6-phosphate

[144493]dehydrogenase variants in Japan. Hum. Genet. 51: 297-305, 1979.

[144494]

[144495]237. Nance, W. E.: Turner's syndrome, twinning, and an unusual variant

[144496]of glucose-6-phosphate dehydrogenase. Am. J. Hum. Genet. 16: 380-392,

[144497]1964.

[144498]

[144499]238. Necheles, T. F.; Snyder, L. M.; Strauss, W.: Glucose-6-phosphate

[144500]dehydrogenase Boston. A new variant associated with congenital nonspherocytic

[144501]hemolytic disease. Humangenetik 13: 218-221, 1971.

[144502]

[144503]239. Niazi, G.; Adeyokunu, A.; Westwood, B.; Beutler, E.: G6PD Aures:

[144504]a rare mutant of G6PD in Saudi Arabia: molecular and clinical presentations. Saudi

[144505]Med. J. 17: 311-314, 1996.

[144506]

[144507]240. Ninfali, P.; Baronciani, L.; Bardoni, A.; Bresolin, N.: Muscle

[144508]expression of glucose-6-phosphate dehydrogenase deficiency in different

[144509]variants. Clin. Genet. 48: 232-237, 1995.

[144510]

[144511]241. Notaro, R.; Afolayan, A.; Luzzatto, L.: Human mutations in glucose

[144512]6-phosphate dehydrogenase reflect evolutionary history. FASEB J. 14:

[144513]485–494, 2000.

[144514]

[144515]242. Nowicki, L.; Strobel, S.; Martin, H.; Koschwitz, U.: Ueber eine

[144516]neue erythrocytaere glucose 6-phosphatdehydrogenase Variante, Typ

[144517]Frankfurt. Klin. Wschr. 52: 478–484, 1974.

[144518]

[144519]243. Nsouly, G. M.; Prchal, J. T.: Characterization of a new G6PD

[144520]variant and its associated oxidative damage. Clin. Res. 29: 829,

[144521]1981.

[144522]

[144523]244. O'Brien, S. J.: The extent and character of biochemical genetic

[144524]variation in the domestic cat. J. Hered. 71: 2–8, 1980.

[144525]

[144526]245. Ogura, H.; Morisaki, T.; Tani, K.; Kanno, H.; Tsutsumi, H.; Takahashi,

[144527]K.; Miyamori, T.; Fujii, H.; Miwa, S.: A new glucose-6-phosphate

[144528]dehydrogenase variant (G6PD Tsukui) associated with congenital hemolytic

[144529]anemia. Hum. Genet. 78: 369-371, 1988.

[144530]

[144531]246. Ohno, S.: Sex Chromosomes and Sex-linked Genes. Berlin and

[144532]New York: Springer (pub.) 1967.

[144533]

[144534]247. Orzalesi, N.; Sorcinelli, R.; Guiso, G.: Increased incidence

[144535]of cataracts in male subjects deficient in glucose-6-phosphate dehydrogenase. Arch.

[144536]Ophthal. 99: 69-70, 1981.

[144537]

[144538]248. Othieno-Obel, A.: East African variant of glucose-6-phosphate

[144539]dehydrogenase. East Afr. Med. J. 49: 230–234, 1972.

[144540]

[144541]249. Pai, G. S.; Sprenkle, J. A.; Do, T. T.; Maren, C. E.; Mi-
geon,

[144542]B. R.: Localization of loci for hypoxanthine phosphoribo-
syltransferase

[144543]and glucose-6-phosphate dehydrogenase and biochemi-
cal evidence of

[144544]nonrandom X chromosome expression from studies of a
human X-autosome

[144545]translocation. Proc. Nat. Acad. Sci. 77: 2810–2813, 1980.

[144546]

[144547]250. Panich, V.: G6PD variants in Laotians. Hum. Hered.
24: 285–290,

[144548]1974.

[144549]

[144550]251. Panich, V.: G6PD Intanon, a new glucose
6-phosphate dehydrogenase

[144551]variant. Humangenetik 21: 203–205, 1974.

[144552]

[144553]252. Panich, V.: G6PD characterization in Thailand. *Genetics* 74

[144554](suppl.): s208 only, 1973.

[144555]

[144556]253. Panich, V.: Glucose-6-phosphate dehydrogenase in Thailand. *Hum.*

[144557]*Genet.* 53: 227-228, 1980.

[144558]

[144559]254. Panich, V.; Bumrungtrakul, P.; Jitjai, C.; Kamolmatayakul, S.;

[144560]Khoprasert, B.; Klaisuvan, C.; Kongmuang, U.; Maneechai, P.; Pornpatkul,

[144561]M.; Ruengrairatanaroje, P.; Surapruk, P.; Viriyayudhakorn, S.: Glucose-6-phosphate

[144562]dehydrogenase deficiency in South Vietnamese. *Hum.*

Hered. 30: 361-364,

[144563]1980.

[144564]

[144565]255. Panich, V.; Na-Nakorn, S.: G6PD variants in Thailand. *J. Med.*

[144566]*Assoc. Thai.* 63: 537-543, 1980.

[144567]

[144568]256. Panich, V.; Sungnate, T.: Characterization of glucose 6-phosphate

[144569]dehydrogenase in Thailand: the occurrence of 6 variants among 50 G6PD

[144570]deficient Thai. Humangenetik 18: 39-46, 1973.

[144571]

[144572]257. Panich, V.; Sungnate, T.; Na-Nakorn, S.: Acute intravascular

[144573]hemolysis and renal failure in a new glucose 6-phosphate dehydrogenase

[144574]variant: G6PD Siriraj. J. Med. Assoc. Thai. 55: 726-731, 1972.

[144575]

[144576]258. Panich, V.; Sungnate, T.; Wasi, P.; Na-Nakorn, S.: G-6-PD Mahidol:

[144577]the most common glucose-6-phosphate dehydrogenase variant in Thailand. J.

[144578]Med. Assoc. Thai. 55: 576-585, 1972.

[144579]

[144580]259. Pawlak, A. L.; Mazurkiewicz, C. A.; Ordynski, J.;

Ruzynkova,

[144581]D.; Horst, A.: G6PD Poznan, variant with severe enzyme deficiency. Humangenetik 28:

[144582]163–165, 1975.

[144583]

[144584]260. Pawlak, A. L.; Zagorski, Z.; Ruzynkova, D.; Horst, A.: Polish

[144585]variant of glucose–6–phosphate dehydrogenase (G–6–PD Lublin). Humangenetik 10:

[144586]340–343, 1970.

[144587]

[144588]261. Pekrun, A.; Eber, S. W.; Schroter, W.: G6PD Avenches and G6PD

[144589]Moosburg: biochemical and erythrocyte membrane characterization. Blut 58:

[144590]11–14, 1989.

[144591]

[144592]262. Perroni, L.; Tassara, P.; Baldi, M.; Reali, R.; Scartezzini,

[144593]P.: G6PD variants detected in Genoa area. In: Weatherall, D. J.; Fiorelli,

[144594]G.; Gorini, S.: Advances in Red Blood Cell Biology. New York: Raven

[144595]Press (pub.) 1982. Pp. 409–416.

[144596]

[144597]263. Persico, M. G.; Toniolo, D.; Nobile, C.; D'Urso, M.; Luzzatto,

[144598]L.: cDNA sequences of human glucose 6–phosphate dehydrogenase cloned

[144599]in pBR322. Nature 294: 778–780, 1981.

[144600]

[144601]264. Persico, M. G.; Viglietto, G.; Martini, G.; Toniolo, D.; Paonessa,

[144602]G.; Moscatelli, C.; Dono, R.; Vulliamy, T.; Luzzatto, L.; D'Urso,

[144603]M.: Isolation of human glucose–6–phosphate dehydrogenase (G6PD) cDNA

[144604]clones: primary structure of the protein and unusual 5–prime non–coding

[144605]region. Nucleic Acids Res. 14: 2511–2522, 1986.

[144606]

[144607]265. Picat, C.; Etiemble, J.; Boivin, P.; Le Prise, P.–Y.: Gd(–)

[144608]Rennes, a new deficient variant of glucose-6-phosphate dehydrogenase

[144609]associated with congenital nonspherocytic hemolytic anemia found in

[144610]France. Hum. Genet. 55: 125-127, 1980.

[144611]

[144612]266. Pinto, P. V. C.; Newton, W. A., Jr.; Richardson, K. E.: Evidence

[144613]for four types of erythrocyte glucose-6-phosphate dehydrogenase from

[144614]G-6-PD deficient human subjects. J. Clin. Invest. 45: 823-831, 1966.

[144615]

[144616]267. Poggi, V.: Personal Communication. London, England 1989.

[144617]

[144618]268. Poon, M.-C.; Hall, K.; Scott, C. W.; Prchal, J. T.: G6PD Viangchan:

[144619]a new glucose 6-phosphate dehydrogenase variant from Laos. Hum. Genet. 78:

[144620]98-99, 1988.

[144621]

[144622]269. Porter, I. H.; Boyer, S. H.; Watson-Williams, E. J.;

Adam, A.;

[144623]Szeinberg, A.; Siniscalco, M.: Variation of glucose-
6-phosphate dehydrogenase

[144624]in different populations. Lancet I: 895-899, 1964.

[144625]

[144626]270. Porter, I. H.; Schulze, J.; McKusick, V. A.: Genetical
linkage

[144627]between the loci for glucose-6-phosphate dehydrogenase
deficiency

[144628]and colour-blindness in American Negroes. Ann. Hum.
Genet. 26: 107-122,

[144629]1962.

[144630]

[144631]271. Prchal, J.; Moreno, H.; Conrad, M.; Vitek, A.: G-6-PD

Dothan:

[144632]a new variant associated with chronic hemolytic anemia.

I.R.C.S. 7:

[144633]348, 1979.

[144634]

[144635]272. Prchal, J. T.: Personal Communication. Birmingham, Ala. 1985.

[144636]

[144637]273. Prchal, J. T.; Crist, W. M.; Malluh, A.; Vitek, A.; Tauxe, W.

[144638]N.; Carroll, A. J.: A new glucose-6-phosphate dehydrogenase deficient

[144639]variant in a patient with Chediak-Higashi syndrome. Blood 56: 476-480,

[144640]1980.

[144641]

[144642]274. Prchal, J. T.; Hall, K.; Csepregy, M.; Lilly, M.; Berkow, R.;

[144643]Scott, C. W.: Two apparent glucose-6-phosphate dehydrogenase variants

[144644]in normal XY males: G6PD Alabama. Am. J. Med. 84: 517-523, 1988.

[144645]

[144646]275. Pretsch, W.; Charles, D. J.; Merkle, S.: X-linked glucose-6-phosphate-dehydrogenase

[144647]deficiency in *Mus musculus*. Biochem. Genet. 26: 89-103, 1988.

[144648]

[144649]276. Puck, J. M.; Willard, H. F.: X inactivation in females with

[144650]X-linked disease. New Eng. J. Med. 338: 325–327, 1998.

[144651]

[144652]277. Ramot, B.; Ben-Bassat, I.; Shchory, M.: New glucose-6-phosphate

[144653]dehydrogenase variants observed in Israel and their association with

[144654]congenital nonspherocytic hemolytic disease. J. Lab. Clin. Med. 74:

[144655]895–901, 1969.

[144656]

[144657]278. Ramot, B.; Brok, F.: A new glucose-6-phosphate dehydrogenase

[144658]mutant (Tel-Hashomer mutant). Ann. Hum. Genet. 28: 167–172, 1964.

[144659]

[144660]279. Rattazzi, M. C.; Corash, L. M.; Van Zzanen, G. E.; Jaffe, E.

[144661]R.; Piomelli, S.: G6PD deficiency and chronic hemolysis:

four new

[144662]mutants--relationships between clinical syndrome and enzyme kinetics. Blood 38:

[144663]205-218, 1971.

[144664]

[144665]280. Rattazzi, M. C.; Lenzerini, L.; Meera Khan, P.; Luz-
zatto, L.

[144666]: Characterization of glucose-6-phosphate dehydroge-
nase variants.

[144667]II. G6PD Kephalaria, G6PD Attica, and G6PD 'Seattle-like'
found in

[144668]Greece. Am. J. Hum. Genet. 21: 154-167, 1969.

[144669]

[144670]281. Ravindranath, Y.; Beutler, E.: Two new variants of
glucose-6-phosphate

[144671]dehydrogenase associated with hereditary non-
spherocytic hemolytic

[144672]anemia: G6PD Wayne and G6PD Huron. Am. J. Hemat. 24:
357-363, 1987.

[144673]

[144674]282. Reys, L.; Manso, C.; Stamatoyannopoulos, G.: Genetic

studies

[144675]on Southeastern Bantu of Mozambique. I. Variants of glucose-6-phosphate

[144676]dehydrogenase. Am. J. Hum. Genet. 22: 203-215, 1970.

[144677]

[144678]283. Rinaldi, A.; Filippi, G.; Siniscalco, M.: Variability of red

[144679]cell phenotypes between and within individuals in an unbiased sample

[144680]of 77 heterozygotes for G6PD deficiency in Sardinia. Am. J. Hum.

[144681]Genet. 28: 496-505, 1976.

[144682]

[144683]284. Rosenstrauss, M.; Chasin, L. A.: Isolation of mammalian cell

[144684]mutants deficient in glucose-6-phosphate dehydrogenase activity: linkage

[144685]to hypoxanthine phosphoribosyl transferase. Proc. Nat. Acad. Sci. 72:

[144686]493-497, 1975.

[144687]

[144688]285. Roth, E. F., Jr.; Raventos-Suarez, C.; Rinaldi, A.;
Nagel, R.

[144689]L.: Glucose-6-phosphate dehydrogenase deficiency in-
hibits in vitro

[144690]growth of *Plasmodium falciparum*. Proc. Nat. Acad. Sci.
80: 298-299,

[144691]1983.

[144692]

[144693]286. Roychoudhury, A. K.; Nei, M.: Human Polymorphic
Genes: World

[144694]Distribution. New York: Oxford Univ. Press (pub.) 1988.

[144695]

[144696]287. Ruwando, C.; Khea, S. C.; Snow, R. W.; Yates, S. N. R.;
Kwiatkoweld,

[144697]D.; Gupta, S.; Warn, P.; Alisopp, G. E. M.; Gilbert, S. C.;
Peschu,

[144698]N.; Newbold, C. I.; Greenwood, S. M.; Marsh, K.; Hill, A. V.
S.:

[144699]Natural selection of hemi- and heterozygotes for G6PD
deficiency in

[144700]Africa by resistance to severe malaria. Nature 376:
246-249, 1995.

[144701]

[144702]288. Saenz, G. F.; Chaves, M.; Berrantes, A.; Elizondo, J.;
Montero,

[144703]A. G.; Yoshida, A.: A glucose-6-phosphate dehydrogenase
variant,

[144704]Gd(-) Santamaria found in Costa Rica. Acta Haemat. 72:
37-40, 1984.

[144705]

[144706]289. Saidi, N.; Hors-Cayla, M. C.; Van Cong, N.; Benne, F.:
Sheep

[144707]gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

[144708]Cell Genet. 25: 200, 1979.

[144709]

[144710]290. Samuel, A. P. W.; Saha, N.; Omer, A.; Hoffbrand, A.
V.: Quantitative

[144711]expression of G6PD activity of different phenotypes of
G6PD and haemoglobin

[144712]in a Sudanese population. Hum. Hered. 31: 110-115,
1981.

[144713]

[144714]291. Sansone, G.; Perroni, L.; Testa, U.; Mareni, C.; Luz-
zatto, L.

[144715]: New genetic variants of glucose 6-phosphate dehydro-
genase (G6PD)

[144716]in Italy. Ann. Hum. Genet. 45: 97-104, 1981.

[144717]

[144718]292. Sansone, G.; Perroni, L.; Yoshida, A.: Glucose-6-phosphate dehydrogenase

[144719]variants from Italian subjects associated with severe neonatal jaundice. Brit.

[144720]J. Haemat. 31: 159-165, 1975.

[144721]

[144722]293. Sansone, G.; Perroni, L.; Yoshida, A.; Dave, V.: A new glucose

[144723]6-phosphate dehydrogenase variant (Gd Trinacria) in two unrelated

[144724]families of Sicilian ancestry. Ital. J. Biochem. 26: 44-50, 1977.

[144725]

[144726]294. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Annenkov, G. A.: A

[144727]description of new mutant forms of erythrocyte glucose-6-phosphate

[144728]dehydrogenase isolated at the territory of the Soviet Union. Genetika 11:

[144729]116-122, 1975.

[144730]

[144731]295. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Idelson, L. J.:

The

[144732]new form of glucose 6-phosphate dehydrogenase (G6PD

'Kaluga') from

[144733]erythrocytes of a patient with chronic non-spherocytic

hemolytic anemia. Vopr.

[144734]Med. Khim. 22: 764-768, 1976.

[144735]

[144736]296. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Idelson, L. J.:

Mutant

[144737]forms of erythrocyte glucose 6-phosphate dehydrogenase

in Ashkenazi:

[144738]description of two new variants, G6PD Kirovograd and

G6PD Zhitomir. Humangenetik 33:

[144739]175-178, 1976.

[144740]

[144741]297. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Tzoneva, M.;

Mavrudieva,

[144742]M.; Toncheva, D.: Variants of erythrocyte glucose-

6-phosphate dehydrogenase

[144743](G6PD) in Bulgarian populations. Hum. Genet. 54:
115–117, 1980.

[144744]

[144745]298. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Zakharova, T.
V.: Regularities

[144746]of distribution of Gd–alleles in Azerbaijan. II. Identification
of

[144747]G6PD mutant forms. Genetika 16: 2217–2225, 1980.

[144748]

[144749]299. Shows, T. B.; Brown, J. A.: Human X–linked genes re-
gionally

[144750]mapped utilizing X–autosome translocations and somatic
cell hybrids. Proc.

[144751]Nat. Acad. Sci. 72: 2125–2129, 1975.

[144752]

[144753]300. Shows, T. B.; Brown, J. A.; Chapman, V. M.: Compara-
tive gene

[144754]mapping of HPRT, G6PD and PGK in man, mouse, and
Muntjac deer. Birth

[144755]Defects Orig. Art. Ser. XII(7): 436–439, 1976.

[144756]

[144757]301. Shows, T. B.; Tashian, R. E.; Brewer, G. J.: Erythrocyte glucose-6-phosphate

[144758]dehydrogenase in Caucasians: new inherited variant. Science 145:

[144759]1056-1057, 1964.

[144760]

[144761]302. Sidi, Y.; Aderka, D.; Brok-Simoni, F.; Benjamin, D.; Ramot, B.;

[144762]Pinkhas, J.: Viral hepatitis with extreme hyperbilirubinaemia, massive

[144763]hemolysis and encephalopathy in a patient with a new G6PD variant. Israel

[144764]J. Med. Sci. 16: 130-133, 1980.

[144765]

[144766]303. Siegel, N. H.; Beutler, E.: Hemolytic anemia caused by G-6-PD

[144767]Carswell, a new variant. Ann. Intern. Med. 75: 437-439, 1971.

[144768]

[144769]304. Smith, J. E.; Ryer, K.; Wallace, L.: Glucose-6-phosphate dehydrogenase

[144770]deficiency in a dog. Enzyme 21: 379–382, 1976.

[144771]

[144772]305. Snyder, L. M.; Necheles, T. F.; Reddy, W. J.: G-6-PD

Worcester:

[144773]a new variant, associated with X-linked optic atrophy. Am.

J. Med. 49:

[144774]125–132, 1970.

[144775]

[144776]306. Stamatoyannopoulos, G.; Kotsakis, P.; Voightlander,

V.; Akrivakis,

[144777]A.; Motulsky, A. G.: Electrophoretic diversity of glucose–

6–phosphate

[144778]dehydrogenase among Greeks. Am. J. Hum. Genet. 22:

587–596, 1970.

[144779]

[144780]307. Stamatoyannopoulos, G.; Voightlander, V.; Akrivakis,

A.: Thessaly

[144781]variant of glucose–6–phosphate dehydrogenase. Human–

genetik 9: 23–25,

[144782]1970.

[144783]

[144784]308. Stamatoyannopoulos, G.; Voigtlander, V.; Kotsakis, P.; Akrivakis,

[144785]A.: Genetic diversity of the 'Mediterranean' glucose-6-phosphate

[144786]dehydrogenase deficiency phenotype. J. Clin. Invest. 50: 1253-1261,

[144787]1971.

[144788]

[144789]309. Stamatoyannopoulos, G.; Yoshida, A.; Bacopoulos, C.; Motulsky,

[144790]A. G.: Athens variant of glucose-6-phosphate dehydrogenase. Science 157:

[144791]831-833, 1967.

[144792]

[144793]310. Stevens, D. J.; Wanachiwanawin, W.; Mason, P. J.; Vulliamy, T.

[144794]J.; Luzzatto, L.: G6PD Canton a common deficient variant in South

[144795]East Asia caused by a 459 arg-to-leu mutation. Nucleic Acids Res. 18:

[144796]7190, 1990.

[144797]

[144798]3 11. Stocco dos Santos, R. C.; Barretto, O. C. O.;

Nonoyama, K.; Castro,

[144799]N. H. C.; Ferraz, O. P.; Walter-Moura, J.; Vescio, C. C. S.;

Becak,

[144800]W.: X-linked syndrome: mental retardation, hip luxation,
and G6PD

[144801]variant (Gd(+)) Butantan). Am. J. Med. Genet. 39: 133-136,
1991.

[144802]

[144803]3 12. Stockham, S. L.; Harvey, J. W.; Kinden, D. A.: Equine
glucose-6-phosphate

[144804]dehydrogenase deficiency. Vet. Path. 31: 518-527, 1994.

[144805]

[144806]3 13. Streiff, F.; Vigneron, C.: Anemie hemolytique
chronique par

[144807]deficit en glucose 6-phosphate deshydrogenase dans une
famille d'origine

[144808]Lorraine. Nouv. Rev. Franc. Hemat. 11: 279-290, 1971.

[144809]

[144810]3 14. Takahashi, K.; Fujii, H.; Takegawa, S.; Tani, K.; Hi-

rono, A.;

[144811]Takizawa, T.; Kawakatsu, T.; Miwa, S.: A new glucose-6-phosphate

[144812]dehydrogenase variant (G6PD Nagano) associated with congenital hemolytic

[144813]anemia. Hum. Genet. 62: 368–370, 1982.

[144814]

[144815]315. Takizawa, T.; Fujii, H.; Takegawa, S.; Takahashi, K.; Hirono,

[144816]A.; Morisaki, T.; Kanno, H.; Oka, R.; Yoshioka, H.; Miwa, S.: A unique

[144817]electrophoretic slow-moving glucose 6-phosphate dehydrogenase variant

[144818](G6PD Asahikawa) with a markedly acidic pH optimum. Hum. Genet. 68:

[144819]70–72, 1984.

[144820]

[144821]316. Takizawa, T.; Huang, I.-Y.; Ikuta, T.; Yoshida, A.: Human glucose-6-phosphate

[144822]dehydrogenase: primary structure and cDNA cloning. Proc. Nat. Acad.

[144823]Sci. 83: 4157–4161, 1986.

[144824]

[144825]317. Takizawa, T.; Yoneyama, Y.; Miwa, S.; Yoshida, A.: A
single

[144826]nucleotide base transition is the basis of the common hu-
man glucose-6-phosphate

[144827]dehydrogenase variant A(+). Genomics 1: 228-231, 1987.

[144828]

[144829]318. Takizawa, T.; Yoshida, A.: Molecular abnormality of
the common

[144830]glucose-6-phosphate dehydrogenase variant, G6PD A(+),
and restriction-fragment-length

[144831]polymorphism. (Abstract) Am. J. Hum. Genet. 41: A241,
1987.

[144832]

[144833]319. Talalak, P.; Beutler, E.: G-6-PD Bangkok: a new vari-
ant found

[144834]in congenital nonspherocytic hemolytic disease (CNHD).
Blood 33:

[144835]772-776, 1969.

[144836]

[144837]320. Tanaka, K. R.; Beutler, E.: Hereditary hemolytic ane-

mia due

[144838]to glucose-6-phosphate dehydrogenase Torrance: a new variant. J.

[144839]Lab. Clin. Med. 73: 657-667, 1969.

[144840]

[144841]321. Tang, T. K.; Huang, C.-S.; Huang, M. J.; Tam, K.-B.; Yeh, C.-H.;

[144842]Tang, C.-J. C.: Diverse point mutations result in glucose-6-phosphate

[144843]dehydrogenase (G6PD) polymorphism in Taiwan. Blood 79: 2135-2140,

[144844]1992.

[144845]

[144846]322. Testa, U.; Meloni, T.; Lania, A.; Battistuzzi, G.; Cuttillo, S.;

[144847]Luzzatto, L.: Genetic heterogeneity of glucose-6-phosphate dehydrogenase

[144848]deficiency in Sardinia. Hum. Genet. 56: 99-105, 1980.

[144849]

[144850]323. Thigpen, J. T.; Steinberg, M. H.; Beutler, E.; Gillespie, G.

[144851]T., Jr.; Dreiling, B. J.; Morrison, F. S.: Glucose-6-phosphate dehydrogenase

[144852]Jackson, a new variant associated with hemolytic anemia. Acta Haemat. 51:

[144853]310-314, 1974.

[144854]

[144855]324. Tishkoff, S. A.; Varkonyi, R.; Cahinhinan, N.; Abbes, S.; Argyropoulos,

[144856]G.; Destro-Bisol, G.; Drousiotou, A.; Dangerfield, B.; Lefranc, G.;

[144857]Loiselet, J.; Piro, A.; Stoneking, M.; Tagarelli, A.; Tagarelli, G.;

[144858]Touma, E. H.; Williams, S. M.; Clark, A. G.: Haplotype diversity

[144859]and linkage disequilibrium at human G6PD: recent origin of alleles

[144860]that confer malarial resistance. Science 293: 455-462, 2001.

[144861]

[144862]325. Tokarev, Y. N.; Chernyak, N. B.; Batischev, A. I.; Lanzina, N.

[144863]V.; Alexeyev, G. A.: Etude des proprietes electrophore-

tiques et cinetiques

[144864]de la glucose-6-phosphate deshydrogenase (Gd)

d'erythrocytes dans

[144865]les deficits hereditaires de l'enzyme: description d'une
nouvelle

[144866]variante de glucose-6-phosphate deshydrogenase: la Gd
Kremenchug. Nouv.

[144867]Rev. Franc. Hemat. 20: 557-564, 1978.

[144868]

[144869]326. Toncheva, D.: Variants of glucose-6-phosphate de-
hydrogenase

[144870]in a Vietnamese population. Hum. Hered. 36: 348-351,
1986.

[144871]

[144872]327. Toncheva, D.; Tzoneva, M.: Genetic polymorphism of
G6PD in a

[144873]Bulgarian population. Hum. Genet. 67: 340-342, 1984.

[144874]

[144875]328. Toniolo, D.; Martini, G.; Migeon, B. R.; Dono, R.: Ex-
pression

[144876]of the G6PD locus on the human X chromosome is associ-

ated with demethylation

[144877]of three CpG islands within 100 kb of DNA. EMBO J. 7:
401–406, 1988.

[144878]

[144879]329. Town, M.; Athanasiou–Metaxa, M.; Luzzatto, L.: In–
tragenic interspecific

[144880]complementation of glucose 6–phosphate dehydrogenase
in human–hamster

[144881]cell hybrids. Somat. Cell Molec. Genet. 16: 97–108, 1990.

[144882]

[144883]330. Town, M.; Bautista, J. M.; Mason, P. J.; Luzzatto, L.:
Both

[144884]mutations in G6PD A– are necessary to produce the G6PD
deficient phenotype. Hum.

[144885]Molec. Genet. 1: 171–174, 1992.

[144886]

[144887]331. Usanga, E. A.; Bienzle, U.; Cancedda, K.; Fasuan, F.
A.; Ajayi,

[144888]O.; Luzzatto, L.: Genetic variants of human erythrocyte
glucose 6–phosphate

[144889]dehydrogenase: new variants in West Africa characterized

by column

[144890]chromatography. Ann. Hum. Genet. 40: 279–286, 1977.

[144891]

[144892]332. Vaca, G.; Ibarra, B.; Garcia Cruz, D.; Medina, C.;
Romero, F.;

[144893]Cantu, J. M.; Beutler, E.: G–6–PD Jalisco and G–6–PD
Morelia: two

[144894]new Mexican variants. Hum. Genet. 71: 82–85, 1985.

[144895]

[144896]333. Vaca, G.; Ibarra, B.; Romero, F.; Olivares, N.; Cantu, J.
M.;

[144897]Beutler, E.: G–6–PD Guadalajara: a new mutant associated
with chronic

[144898]nonspherocytic hemolytic anemia. Hum. Genet. 61:
175–176, 1982.

[144899]

[144900]334. Ventura, A.; Panizon, F.; Soranzo, M. R.; Veneziano,
G.; Sansone,

[144901]G.; Testa, U.; Luzzatto, L.: Congenital dyserythropoietic
anaemia

[144902]type II associated with a new type of G6PD deficiency

(G6PD Gabrovizza). Acta

[144903]Haemat. 71: 227–234, 1984.

[144904]

[144905]335. Vergnes, H.; Gherardi, M.; Bouloux, C.: Erythrocyte
glucose–6–phosphate

[144906]dehydrogenase in the Niokolonko (Malinke of the Niokolo)
of the Eastern

[144907]Senegal: identification of a slow variant with normal activ–
ity (Tacoma–like). Hum.

[144908]Hered. 25: 80–87, 1975.

[144909]

[144910]336. Vergnes, H.; Gherardi, M.; Quilici, J. C.; Yoshida, A.;
Giacardy,

[144911]R.: G6PD Luz–Saint–Sauveur: a new variant with abnormal
electrophoretic

[144912]mobility, mild enzyme deficiency and absence of
hemolytic disorders. I.R.C.S. 7:

[144913]14, 1973.

[144914]

[144915]337. Vergnes, H.; Gherardi, M.; Yoshida, A.: G6PD Lozere
and Trinacria–like:

[144916]segregation of two non-hemolytic variants in a French family. Hum.

[144917]Genet. 34: 293-298, 1976.

[144918]

[144919]338. Vergnes, H.; Ribet, A.; Bommelaer, G.; Amadieu, J.; Brun, H.

[144920]: GD(-) Muret and GD(-) Colomiers, two new variants of glucose-6-phosphate

[144921]dehydrogenase associated with favism. Hum. Genet. 57: 332-334, 1981.

[144922]

[144923]339. Vergnes, H.; Yoshida, A.; Gourdin, D.; Gherardi, M.; Bierme,

[144924]R.; Ruffie, J.: Glucose 6-phosphate dehydrogenase Toulouse: a new

[144925]variant with marked instability and severe deficiency discovered in

[144926]a family of Mediterranean ancestry. Acta Haemat. 51: 240-249, 1974.

[144927]

[144928]340. Viglietto, G.; Montanaro, V.; Calabro, V.; Vallone, D.; D'Urso,

[144929]M.; Persico, M. G.; Battistuzzi, G.: Common glucose-6-phosphate dehydrogenase

[144930](G6PD) variants from the Italian population: biochemical and molecular

[144931]characterization. Ann. Hum. Genet. 54: 1-15, 1990.

[144932]

[144933]341. Vives-Corrons, J.-L.; Kuhl, W.; Pujades, M. A.; Beutler, E.:

[144934]Molecular genetics of the glucose-6-phosphate dehydrogenase (G6PD)

[144935]Mediterranean variant and description of a new G6PD mutant, G6PD Andalus(1361A). Am.

[144936]J. Hum. Genet. 47: 575-579, 1990.

[144937]

[144938]342. Vives-Corrons, J. L.; Feliu, E.; Pujades, M. A.; Cardellach,

[144939]F.; Rozman, C.; Carreras, A.; Jou, J. M.; Vallespi, M. T.; Zuazu,

[144940]F. J.: Severe glucose-6-phosphate dehydrogenase (G6PD) deficiency

[144941]associated with chronic hemolytic anemia, granulocyte dysfunction,

[144942]and increased susceptibility to infections: description of a new molecular

[144943]variant (G6PD Barcelona). Blood 59: 428–434, 1982.

[144944]

[144945]343. Vives–Corrons, J. L.; Pujades, A.: Heterogeneity of 'Mediterranean

[144946]type' glucose–6–phosphate dehydrogenase (G6PD) deficiency in Spain

[144947]and description of two new variants associated with favism. Hum.

[144948]Genet. 60: 216–221, 1982.

[144949]

[144950]344. Vives–Corrons, J. L.; Pujades, A.; Curia, M. D.: Caracterizacion

[144951]molecular de la glucosa–6–fosfato deshidrogenasa (G6PD) en 24 casos

[144952]de deficit enzimatico y descripcion de una nueva variante (G6PD–Betica). Sangre 25:

[144953]1049–1064, 1980.

[144954]

[144955]345. Vives–Corrons, J. L.; Pujades, M. A.; Petit, J.; Colomer,

D.;

[144956]Corbella, M.; Aguilar i Bascompte, J. L.; Merino, A.:

Chronic nonspherocytic

[144957]hemolytic anemia (CNSHA) and glucose 6 phosphate dehydrogenase (G6PD)

[144958]deficiency in a patient with familial amyloidotic polyneuropathy (FAP):

[144959]molecular study of a new variant (G6PD Clinic) with markedly acidic

[144960]pH optimum. Hum. Genet. 81: 161–164, 1989.

[144961]

[144962]346. Vulliamy, T.; Beutler, E.; Luzzatto, L.: Variants of glucose–6–phosphate

[144963]dehydrogenase are due to missense mutations spread throughout the

[144964]coding region of the gene. Hum. Mutat. 2: 159–167, 1993.

[144965]

[144966]347. Vulliamy, T.; Mason, P.; Luzzatto, L.: The molecular basis of

[144967]glucose–6–phosphate dehydrogenase deficiency. Trends Genet. 8: 138–143,

[144968]1992.

[144969]

[144970]348. Vulliamy, T.; Rovira, A.; Yusoff, N.; Colomer, D.; Luzzatto,

[144971]L.; Vives-Corrons, J.-L.: Independent origin of single and double

[144972]mutations in the human glucose 6-phosphate dehydrogenase gene. Hum.

[144973]Mutat. 8: 311-318, 1996.

[144974]

[144975]349. Vulliamy, T. J.: Personal Communication. London, England

[144976]1989.

[144977]

[144978]350. Vulliamy, T. J.; D'Urso, M.; Battistuzzi, G.; Estrada, M.; Foulkes,

[144979]N. S.; Martini, G.; Calabro, V.; Poggi, V.; Giordano, R.; Town, M.;

[144980]Luzzatto, L.; Persico, M. G.: Diverse point mutations in the human

[144981]glucose-6-phosphate dehydrogenase gene cause enzyme

deficiency and

[144982]mild or severe hemolytic anemia. Proc. Nat. Acad. Sci. 85:
5171–5175,

[144983]1988.

[144984]

[144985]351. Vulliamy, T. J.; Kaeda, J. S.; Ait-Chafa, D.; Mangerini,
R.;

[144986]Roper, D.; Barbot, J.; Mehta, A. B.; Athanassiou–Metaxa,
M.; Luzzatto,

[144987]L.; Mason, P. J.: Clinical and haematological consequences
of recurrent

[144988]G6PD mutations and a single new mutation causing
chronic nonspherocytic

[144989]haemolytic anaemia. Brit. J. Haemat. 101: 670–675, 1998.

[144990]

[144991]352. Vulliamy, T. J.; Othman, A.; Town, M.; Nathwani, A.;
Falusi,

[144992]A. G.; Mason, P. J.; Luzzatto, L.: Polymorphic sites in the
African

[144993]population detected by sequence analysis of the glucose–
6–phosphate

[144994]dehydrogenase gene outline the evolution of the variants

A and A-. Proc.

[144995]Nat. Acad. Sci. 88: 8568–8571, 1991.

[144996]

[144997]353. Vulliamy, T. J.; Wanachiwanawin, W.; Mason, P. J.;
Luzzatto,

[144998]L.: G6PD Mahidol, a common deficient variant in South
East Asia is

[144999]caused by a (163)glycine-to-serine mutation. Nucleic
Acids Res. 17:

[145000]5868, 1989.

[145001]

[145002]354. Vuopio, P.; Harkonen, M.; Helske, T.; Naeveri, H.: Red
cell

[145003]glucose-6-phosphate dehydrogenase deficiency in Fin-
land: characterization

[145004]of a new variant with severe enzyme deficiency. Scand. J.
Haemat. 15:

[145005]145–152, 1975.

[145006]

[145007]355. Vuopio, P.; Harkonen, M.; Johnsson, P.; Nuutinen, M.:
Red cell

[145008]glucose-phosphate dehydrogenase deficiency in Finland.

Ann. Clin.

[145009]Res. 5: 168-178, 1973.

[145010]

[145011]356. Waitz, R.; Boivin, P.; Oberling, F.; Casenave, J. P.;

North,

[145012]M. L.; Mayer, S.: Variante Gd(-) Strasbourg de la glucose-

6-phosphate

[145013]dehydrogenase. Nouv. Rev. Franc. Hemat. 10: 312-314,

1970.

[145014]

[145015]357. Wang, Y. M.; Patterson, J. H.; Van Eys, J.: The poten-

tial use

[145016]of xylitol in glucose-6-phosphate dehydrogenase defi-

ciency anemia. J.

[145017]Clin. Invest. 50: 1421-1428, 1971.

[145018]

[145019]358. Weimer, T. A.; Salzano, F. M.; Hutz, M. H.: Erythro-

cyte isozymes

[145020]and hemoglobin types in a southern Brazilian population.

J. Hum.

[145021]Evol. 10: 319–328, 1981.

[145022]

[145023]359. Weimer, T. A.; Schuler, L.; Beutler, E.; Salzano, F. M.:
Gd(+)

[145024]Laguna, a new rare glucose–6–phosphate dehydrogenase
variant from

[145025]Brazil. Hum. Genet. 65: 402–404, 1984.

[145026]

[145027]360. Weinreich, J.; Busch, D.; Gottstein, U.; Schaefer, J.;
Rohr,

[145028]J.: Ueber zwei neue Faelle von hereditaerer nichtsphaero-
cytaerer

[145029]haemolytischer Anaemie bei glucose–
6–phosphat–dehydrogenase–Defekt

[145030]in einer Nord Deutschen Familie. Klin. Wschr. 46:
146–149, 1968.

[145031]

[145032]361. Welch, S. G.; McGregor, I. A.; Williams, K.: A new
variant of

[145033]human erythrocyte G6PD occurring at a high frequency
amongst the population

[145034]of two villages in The Gambia, West Africa. Hum. Genet.
40: 305–309,
[145035]1978.

[145036]

[145037]362. Westring, D. W.; Pisciotto, A. V.: Anemia, cataracts,
and seizures

[145038]in patients with glucose–6–phosphate dehydrogenase de-
ficiency. Arch.

[145039]Intern. Med. 118: 385–390, 1966.

[145040]

[145041]363. WHO: Nomenclature of glucose–6–phosphate dehy-
drogenase in

[145042]man. Bull. WHO 36: 319–322, 1967. Note: See Also:
Canad. Med. Assoc.

[145043]J. 97: 422–424, 1967.

[145044]

[145045]364. WHO: Scientific group on the standardization of pro-
cedures

[145046]for the study of glucose–6–phosphate dehydrogenase.
WHO Techn. Rep.

[145047](pub.) Ser. No. 366: 1967.

[145048]

[145049]365. Wilson, W. W.: Congenital hemolytic anemia due to a deficiency

[145050]of glucose 6-phosphate dehydrogenase. Rocky Mt. Med. J. 73: 160-162,

[145051]1976.

[145052]

[145053]366. Witt, I.; Yoshioka, S.: Biochemical characterization of a glucose-6-phosphate

[145054]dehydrogenase variant with favism: G-6-PD Zaehringen. Klin. Wschr. 50:

[145055]205-209, 1972.

[145056]

[145057]367. Wong, P. W. K.; Shih, L.-Y.; Hsia, D. Y. Y.: Characterization

[145058]of glucose-6-phosphate dehydrogenase among Chinese. Nature 208:

[145059]1323-1324, 1965.

[145060]

[145061]368. Yermakov, N.; Tokarev, J.; Chernjak, N.; Schoenian, G.; Grieger,

[145062]M.; Guckler, G.; Jacobasch, G.; Mahmudova, M.; Bahramov,
S.: New

[145063]stable mutant Gd(–) variants: G6PD Tashkent and G6PD
Nucus: molecular

[145064]basis of hereditary enzyme deficiency. Acta Biol. Med. Ger.
40:

[145065]559–562, 1981.

[145066]

[145067]369. Yoshida, A.: Personal Communication. Duarte, Calif.
2/26/1996.

[145068]

[145069]370. Yoshida, A.: Amino acid substitution (histidine to ty-
rosine)

[145070]in a glucose–6–phosphatate dehydrogenase variant (G6PD
Hektoen) associated

[145071]with over–production. J. Mol. Biol. 52: 483–490, 1970.

[145072]

[145073]371. Yoshida, A.: A single amino acid substitution
(asparagine to

[145074]aspartic acid) between normal (B plus) and the common
Negro variant

[145075](A plus) of human glucose-6-phosphate dehydrogenase.

Proc. Nat. Acad.

[145076]Sci. 57: 835-840, 1967.

[145077]

[145078]372. Yoshida, A.: Human glucose-6-phosphate dehydrogenase: purification

[145079]and characterization of Negro type variant (A+) and comparison with

[145080]normal enzyme (B+). Biochem. Genet. 1: 81-99, 1967.

[145081]

[145082]373. Yoshida, A.; Baur, E. W.; Motulsky, A. G.: A Philippino glucose-6-phosphate

[145083]dehydrogenase variant (G6PD Union) with enzyme deficiency and altered

[145084]substrate specificity. Blood 35: 506-513, 1970.

[145085]

[145086]374. Yoshida, A.; Beutler, E.: Human glucose-6-phosphate dehydrogenase

[145087]variants: a supplementary tabulation. Ann. Hum. Genet. 41: 347-355,

[145088]1978.

[145089]

[145090]375. Yoshida, A.; Beutler, E.; Motulsky, A. G.: Table of human glucose-6-phosphate

[145091]dehydrogenase variants. Bull. WHO 45: 243-253, 1971.

[145092]

[145093]376. Yoshida, A.; Stamatoyannopoulos, G.; Motulsky, A. G.: Negro

[145094]variant of glucose-6-phosphate dehydrogenase deficiency (A-) in man. Science 155:

[145095]97-99, 1967.

[145096]

[145097]377. Yoshida, A.; Takizawa, T.: The same extra FokI cleavage site

[145098]exists in glucose-6-phosphate dehydrogenase variants A(+) and A(-). Am.

[145099]J. Hum. Genet. 43: 131-133, 1988.

[145100]

[145101]378. Yoshida, A.; Takizawa, T.; Prchal, J. T.: RFLP of the X chromosome-linked

[145102]glucose-6-phosphate dehydrogenase locus in blacks. Am. J. Hum. Genet. 42:

[145103]872–876, 1988.

[145104]

[145105]379. Zinkham, W. H.: A deficiency of glucose–
6–phosphate dehydrogenase

[145106]activity in lens from individuals with primaquine–sensitive
erythrocytes. Bull.

[145107]Johns Hopkins Hosp. 109: 206–216, 1961.

[145108]

[145109]380. Zuo, L.; Chen, E.; Du, C. S.; Chang, C. N.; Chiu, D. T.
Y.:

[145110]Genetic study of Chinese G6PD variants by direct PCR se–
quencing. (Abstract) Blood 76

[145111](suppl. 1): 51A, 1990.

[145112]

[145113]

[145114]13563.

[145115]Campuzano, V.; Montermini, L.; Molto, M. D.; Pianese, L.;
Cossee,

[145116]M.; Cavalcanti, F.; Monros, E.; Rodius, F.; Duclos, F.; Mon–
ticelli,

[145117]A.; Zara, F.; Canizares, J.; Koutnikova, H.; Bidichandani, S.
I.;

[145118]Gellera, C.; Brice, A.; Trouillas, P.; De Michele, G.; Filla, A.;

[145119]De Frutos, R.; Palau, F.; Patel, P. I.; Di Donato, S.; Mandel,
J.

[145120]–L.; Coccozza, S.; Koenig, M.; Pandolfo, M.: Friedreich's
ataxia:

[145121]autosomal recessive disease caused by an intronic GAA
triplet repeat

[145122]expansion. Science 271: 1423–1427, 1996.

[145123]

[145124]

[145125]13564.

[145126]Ilyin, G. P.; Rialland, M.; Pigeon, C.; Guguen–Guillouzo, C.:

[145127]cDNA cloning and expression analysis of new members of
the mammalian

[145128]F–box protein family. Genomics 67: 40–47, 2000.

[145129]

[145130]

[145131]13565.

[145132]Inase, N.; Fushimi, K.; Ishibashi, K.; Uchida, S.; Ichioka, M.;

[145133]Sasaki, S.; Marumo, F.: Isolation of human aquaporin 3
gene. J.

[145134]Biol. Chem. 270: 17913–17916, 1995.

[145135]

[145136]

[145137]13566.

[145138]Ishibashi, K.; Sasaki, S.; Saito, F.; Ikeuchi, T.; Marumo, F.:

[145139]Structure and chromosomal localization of a human water
channel (AQP3)

[145140]gene. Genomics 27: 352–354, 1995. Note: Erratum: Ge-
nomics 30: 633

[145141]only, 1995.

[145142]

[145143]

[145144]13567.

[145145]Ma, T.; Hara, M.; Sougrat, R.; Verbavatz, J.–M.; Verkman,
A. S.

[145146]: Impaired stratum corneum hydration in mice lacking epi–

dermal water

[145147]channel aquaporin-3. J. Biol. Chem. 277: 17147-17153, 2002.

[145148]

[145149]

[145150]13568.

[145151]Ishibashi, K.; Sasaki, S.; Fushimi, K.; Uchida, S.; Kuwahara, M.;

[145152]Saito, H.; Furukawa, T.; Nakajima, K.; Yamaguchi, Y.; Gojobori, T.;

[145153]Marumo, F.: Molecular cloning and expression of a member of the aquaporin

[145154]family with permeability to glycerol and urea in addition to water

[145155]expressed at the basolateral membrane of kidney collecting duct cells. Proc.

[145156]Nat. Acad. Sci. 91: 6269-6273, 1994.

[145157]

[145158]

[145159]13569.

[145160]Ma, T.; Song, Y.; Yang, B.; Gillespie, A.; Carlson, E. J.; Ep-

stein,

[145161]C. J.; Verkman, A. S.: Nephrogenic diabetes insipidus in mice lacking

[145162]aquaporin-3 water channels. Proc. Nat. Acad. Sci. 97: 4386-4391,

[145163]2000.

[145164]

[145165]

[145166]13570.

[145167]Mulders, S. M.; Olde Weghuis, D.; van Boxtel, J. A. F.; Geurts

[145168]van Kessel, A.; Echevarria, M.; van Os, C. H.; Deen, P. M. T.: Localization

[145169]of the human gene for aquaporin 3 (AQP3) to chromosome 9, region p21-p12,

[145170]using fluorescent in situ hybridization. Cytogenet. Cell Genet. 72:

[145171]303-305, 1996.

[145172]

[145173]

[145174]13571.

[145175]Sougrat, R.; Morand, M.; Gondran, C.; Barre, P.; Gobin, R.;
Bonte,

[145176]F.; Dumas, M.; Verbavatz, J.–M.: Functional expression of
AQP3 in

[145177]human skin epidermis and reconstructed epidermis. J. In-
vest. Derm. 118:

[145178]678–685, 2002.

[145179]

[145180]

[145181]13572.

[145182]Andersson, P.; McGuire, J.; Rubio, C.; Gradin, K.; Whitelaw,
M.

[145183]L.; Pettersson, S.; Hanberg, A.; Poellinger, L.: A constitu-
tively

[145184]active dioxin/aryl hydrocarbon receptor induces stomach
tumors. Proc.

[145185]Nat. Acad. Sci. 99: 9990–9995, 2002.

[145186]

[145187]

[145188]13573.

[145189]Ema, M.; Matsushita, N.; Sogawa, K.; Ariyama, T.; Inazawa, J.;

[145190]Nemoto, T.; Ota, M.; Oshimura, M.; Fujii-Kuriyama, Y.:
Human arylhydrocarbon

[145191]receptor: functional expression and chromosomal assignment to 7p21. J.

[145192]Biochem. 116: 845–851, 1994.

[145193]

[145194]

[145195]13574.

[145196]Le Beau, M. M.; Carver, L. A.; Espinosa, R., III; Schmidt, J. V.;

[145197]Bradfield, C. A.: Chromosomal localization of the human
AHR locus

[145198]encoding the structural gene for the Ah receptor to
7p21–p15. Cytogenet.

[145199]Cell Genet. 66: 172–176, 1994.

[145200]

[145201]

[145202]13575.

[145203]Micka, J.; Milatovich, A.; Menon, A.; Grabowski, G. A.;
Puga, A.;

[145204]Nebert, D. W.: Human Ah receptor (AHR) gene: localization
to 7p15

[145205]and suggestive correlation of polymorphism with CYP1A1
inducibility. Pharmacogenetics 7:

[145206]95–101, 1997.

[145207]

[145208]

[145209]13576.

[145210]Shimizu, Y.; Nakatsuru, Y.; Ichinose, M.; Takahashi, Y.;
Kume,

[145211]H.; Mimura, J.; Fujii–Kuriyama, Y.; Ishikawa, T.:
Benzo[a]pyrene

[145212]carcinogenicity is lost in mice lacking the aryl hydrocar-
bon receptor. Proc.

[145213]Nat. Acad. Sci. 97: 779–782, 2000.

[145214]

[145215]

[145216]13577.

[145217]Adams, M. D.; Kelley, J. M.; Gocayne, J. D.; Dubnick, M.;
Polymeropoulos,

[145218]M. H.; Xiao, H.; Merril, C. R.; Wu, A.; Olde, B.; Moreno, R.
F.; Kerlavage,

[145219]A. R.; McCombie, W. R.; Venter, J. C.: Complementary DNA
sequencing:

[145220]expressed sequence tags and human genome project. Sci-
ence 252: 1651–1656,

[145221]1991.

[145222]

[145223]

[145224]13578.

[145225]Polvi, A.; Armstrong, E.; Lai, C.; Lemke, G.; Huebner, K.;
Spritz,

[145226]R. A.; Guida, L. C.; Nicholls, R. D.; Alitalo, K.: The human
TYRO3

[145227]gene and pseudogene are located in chromosome
15q14–q25. Gene 134:

[145228]289–293, 1993.

[145229]

[145230]

[145231]13579.

[145232]Chen, P.; Hao, W.; Rife, L.; Wang, X. P.; Shen, D.; Chen, J.;
Ogden,

[145233]T.; Van Boemel, G. B.; Wu, L.; Yang, M.; Fong, H. K. W.: A
photic

[145234]visual cycle of rhodopsin regeneration is dependent on
Rgr. Nature

[145235]Genet. 28: 256–260, 2001.

[145236]

[145237]

[145238]13580.

[145239]Chen, X.–N.; Korenberg, J. R.; Jiang, M.; Shen, D.; Fong, H.
K.

[145240]W.: Localization of the human RGR opsin gene to chromo-
some 10q23. Hum.

[145241]Genet. 97: 720–722, 1996.

[145242]

[145243]

[145244]13581.

[145245]Jiang, M.; Pandey, S.; Fong, H. K. W.: An opsin homologue

in the

[145246]retina and pigment epithelium. Invest. Ophthalm. Visual Sci.

34:

[145247]3669–3678, 1993.

[145248]

[145249]

[145250]13582.

[145251]Morimura, H.; Saindelle–Ribeaudéau, F.; Berson, E. L.;

Dryja, T.

[145252]P.: Mutations in RGR, encoding a light–sensitive opsin homologue,

[145253]in patients with retinitis pigmentosa. (Letter) Nature

Genet. 23:

[145254]393–394, 1999.

[145255]

[145256]

[145257]13583.

[145258]Shen, D.; Jiang, M.; Hao, W.; Tao, L.; Salazar, M.; Fong, H.

K.

[145259]W.: A human opsin–related gene that encodes a retinalde–

hyde-binding

[145260]protein. Biochemistry 33: 13117-13125, 1994.

[145261]

[145262]

[145263]13584.

[145264]Asano, H.; Ishida, A.; Hasegawa, M.; Ono, T.; Yoshida, M.
C.; Taniguchi,

[145265]M.; Kanno, M.: The mouse Mel-18 'RING-finger' gene: genomic organization,

[145266]promoter analysis and chromosomal assignment. DNA
Seq. 3: 369-377,

[145267]1993.

[145268]

[145269]

[145270]13585.

[145271]Ishida, A.; Asano, H.; Hasegawa, M.; Koseki, H.; Ono, T.;
Yoshida,

[145272]M. C.; Taniguchi, M.; Kanno, M.: Cloning and chromosome
mapping of

[145273]the human Mel-18 gene which encodes a DNA-binding

protein with a new

[145274]'RING-finger' motif. Gene 129: 249-255, 1993.

[145275]

[145276]

[145277]13586.

[145278]Tagawa, M.; Sakamoto, T.; Shigemoto, K.; Matsubara, H.;
Tamura,

[145279]Y.; Ito, T.; Nakamura, I.; Okitsu, A.; Imai, K.; Taniguchi, M.:
Expression

[145280]of novel DNA-binding protein with zinc finger structure in
various

[145281]tumor cells. J. Biol. Chem. 265: 20021-20026, 1990.

[145282]

[145283]

[145284]13587.

[145285]Hara, E.; Yamaguchi, T.; Nojima, H.; Ide, T.; Campisi, J.;
Okayama,

[145286]H.; Oda, K.: Id-related genes encoding helix-loop-helix
proteins

[145287]are required for G1 progression and are repressed in

senescent human

[145288]fibroblasts. J. Biol. Chem. 269: 2139–2145, 1994.

[145289]

[145290]

[145291]13588.

[145292]Mathew, S.; Chen, W.; Murty, V. V. V. S.; Benezra, R.; Chaganti,

[145293]R. S. K.: Chromosomal assignment of human ID1 and ID2 genes. Genomics 30:

[145294]385–387, 1995.

[145295]

[145296]

[145297]13589.

[145298]Chen, M. X.; McPartlin, A. E.; Brown, L.; Chen, Y. H.; Barker,

[145299]H. M.; Cohen, P. T. W.: A novel human protein serine/threonine phosphatase,

[145300]which possesses four tetratricopeptide repeat motifs and localizes

[145301]to the nucleus. EMBO J. 13: 4278–4290, 1994.

[145302]

[145303]

[145304]13590.

[145305]Xu, X.; Lagercrantz, J.; Zickert, P.; Bajalica-Lagercrantz, S.;

[145306]Zetterberg, A.: Chromosomal localization and 5-prime sequence of

[145307]the human protein serine/threonine phosphatase 5-prime gene. Biochem.

[145308]Biophys. Res. Commun. 218: 514-517, 1996.

[145309]

[145310]

[145311]13591.

[145312]Yong, W. H.; Ueki, K.; Chou, D.; Reeves, S. A.; von Deimling, A.;

[145313]Gusella, J. F.; Mohrenweiser, H. W.; Buckler, A. J.; Louis, D. N.

[145314]: Cloning of a highly conserved human protein serine-threonine phosphatase

[145315]gene from the glioma candidate region on chromosome 19q13.3. Genomics 29:

[145316]533–536, 1995.

[145317]

[145318]

[145319]13592.

[145320]Gaubatz, S.; Lindeman, G. J.; Ishida, S.; Jakoi, L.; Nevins, J.

[145321]R.; Livingston, D. M.; Rempel, R. E.: E2F4 and E2F5 play an
essential

[145322]role in pocket protein-mediated G1 control. Molec. Cell 6:
729–735,

[145323]2000.

[145324]

[145325]

[145326]13593.

[145327]Ginsberg, D.; Vairo, G.; Chittenden, T.; Xiao, Z.-X.; Xu, G.;
Wydner,

[145328]K. L.; DeCaprio, J. A.; Lawrence, J. B.; Livingston, D. M.:
E2F-4,

[145329]a new member of the E2F transcription factor family, in-
teracts with

[145330]p107. Genes Dev. 8: 2665–2679, 1994.

[145331]

[145332]

[145333]13594.

[145334]Humbert, P. O.; Rogers, C.; Ganiatsas, S.; Landsberg, R. L.;
Trimarchi,

[145335]J. M.; Dandapani, S.; Brugnara, C.; Erdman, S.; Schrenzel,
M.; Bronson,

[145336]R. T.; Lees, J. A.: E2F4 is essential for normal erythrocyte
maturation

[145337]and neonatal viability. Molec. Cell 6: 281–291, 2000.

[145338]

[145339]

[145340]13595.

[145341]Rempel, R. E.; Saenz–Robles, M. T.; Storms, R.; Morham,
S.; Ishida,

[145342]S.; Engel, A.; Jakoi, L.; Melhem, M. F.; Pipas, J. M.; Smith,
C.;

[145343]Nevins, J. R.: Loss of E2F4 activity leads to abnormal de-
velopment

[145344]of multiple cellular lineages. Molec. Cell 6: 293–306,
2000.

[145345]

[145346]

[145347]13596.

[145348]Sardet, C.; Vidal, M.; Cobrinik, D.; Geng, Y.; Onufryk, C.;
Chen,

[145349]A.; Weinberg, R. A.: E2F-4 and E2F-5, two members of the
E2F family,

[145350]are expressed in the early phases of the cell cycle. Proc.
Nat. Acad.

[145351]Sci. 92: 2403-2407, 1995.

[145352]

[145353]

[145354]13597.

[145355]Breitbart, R. E.; Liang, C.; Smoot, L. B.; Laheru, D. A.; Mah-
davi,

[145356]V.; Nadal-Ginard, B.: A fourth human MEF2 transcription
factor, hMEF2D,

[145357]is an early marker of the myogenic lineage. Development
118: 1095-1106,

[145358]1993.

[145359]

[145360]

[145361]13598.

[145362]Hobson, G. M.; Krahe, R.; Garcia, E.; Siciliano, M. J.; Funanage,

[145363]V. L.: Regional chromosomal assignments for four members of the MADS

[145364]domain transcription enhancer factor 2 (MEF2) gene family to human

[145365]chromosomes 15q26, 19p12, 5q14, and 1q12–q23. Genomics 29: 704–711,

[145366]1995.

[145367]

[145368]

[145369]13599.

[145370]Mao, Z.; Bonni, A.; Xia, F.; Nadal-Vicans, M.; Greenberg, M. E.

[145371]: Neuronal activity–dependent cell survival mediated by transcription

[145372]factor MEF2. Science 286: 785–790, 1999.

[145373]

[145374]

[145375]13600.

[145376]Martin, J. F.; Miano, J. M.; Hustad, C. M.; Copeland, N. G.;
Jenkins,

[145377]N. A.; Olson, E. N.: A Mef2 gene that generates a muscle-
specific

[145378]isoform via alternative mRNA splicing. Molec. Cell. Biol.
14: 1647–1656,

[145379]1994.

[145380]

[145381]

[145382]13601.

[145383]Molkentin, J. D.; Black, B. L.; Martin, J. F.; Olson, E. N.: Co-
operative

[145384]activation of muscle gene expression by MEF2 and myo-
genic bHLH proteins. Cell 83:

[145385]1125–1136, 1995.

[145386]

[145387]

[145388]13602.

[145389]Naya, F. J.; Black, B. L.; Wu, H.; Bassel-Duby, R.; Richardson,
son,

[145390]J. A.; Hill, J. A.; Olson, E. N.: Mitochondrial deficiency and
cardiac

[145391]sudden death in mice lacking the MEF2A transcription fac-
tor. Nature

[145392]Med. 15Oct, 2002. Note: Advance Electronic Publication.

[145393]

[145394]

[145395]13603.

[145396]Pollock, R.; Treisman, R.: Human SRF-related proteins:
DNA-binding

[145397]properties and potential regulatory targets. Genes Dev. 5:
2327-2341,

[145398]1991.

[145399]

[145400]

[145401]13604.

[145402]Suzuki, E.; Lowry, J.; Sonoda, G.; Testa, J. R.; Walsh, K.:

Structures

[145403]and chromosome locations of the human MEF2A gene and
a pseudogene

[145404]MEF2AP. Cytogenet. Cell Genet. 73: 244–249, 1996.

[145405]

[145406]

[145407]13605.

[145408]Yu, Y.–T.; Breitbart, R. E.; Smoot, L. B.; Lee, Y.; Mahdavi,
V.;

[145409]Nadal–Ginard, B.: Human myocyte–specific enhancer fac–
tor 2 comprises

[145410]a group of tissue–restricted MADS box transcription fac–
tors. Genes

[145411]Dev. 6: 1783–1798, 1992.

[145412]

[145413]

[145414]13606.

[145415]Youn, H.–D.; Sun, L.; Prywes, R.; Liu, J. O.: Apoptosis of T
cells

[145416]mediated by Ca(2+)–induced release of the transcription

factor MEF2. Science 286:

[145417]790–793, 1999.

[145418]

[145419]

[145420]13607.

[145421]Krainc, D.; Haas, M.; Ward, D. C.; Lipton, S. A.; Bruns, G.;
Leifer,

[145422]D.: Assignment of human myocyte-specific enhancer
binding factor

[145423]2C (hMEF2C) to human chromosome 5q14 and evidence
that MEF2C is evolutionarily

[145424]conserved. Genomics 29: 809–811, 1995.

[145425]

[145426]

[145427]13608.

[145428]Leifer, D.; Krainc, D.; Yu, Y.–T.; McDermott, J.; Breitbart, R.

[145429]E.; Heng, J.; Neve, R. L.; Kosofsky, B.; Nadal-Ginard, B.;
Lipton,

[145430]S. A.: MEF2C, a MADS/MEF2-family transcription factor
expressed in

[145431]a laminar distribution in cerebral cortex. Proc. Nat. Acad.

Sci. 90:

[145432]1546–1550, 1993.

[145433]

[145434]

[145435]13609.

[145436]Greengard, P.; Valtorta, F.; Czernik, A. J.; Benfenati, F.:

Synaptic

[145437]vesicle phosphoproteins and regulation of synaptic function. Science 259:

[145438]780–785, 1993.

[145439]

[145440]

[145441]13610.

[145442]Li, L.; Chin, L.-S.; Greengard, P.; Copeland, N. G.; Gilbert, D.

[145443]J.; Jenkins, N. A.: Localization of the synapsin II (SYN2) gene to

[145444]human chromosome 3 and mouse chromosome 6. Genomics 28: 365–366,

[145445]1995.

[145446]

[145447]

[145448]13611.

[145449]McCright, B.; Rivers, A. M.; Audlin, S.; Virshup, D. M.: The
B56

[145450]family of protein phosphatase 2A (PP2A) regulatory sub-
units encodes

[145451]differentiation-induced phosphoproteins that target PP2A
to both nucleus

[145452]and cytoplasm. J. Biol. Chem. 271: 22081-22089, 1996.

[145453]

[145454]

[145455]13612.

[145456]Van Hoof, C.; Aly, M. S.; Garcia, A.; Cayla, X.; Cassiman, J.
J.;

[145457]Merlevede, W.; Goris, J.: Structure and chromosomal local-
ization

[145458]of the human gene of the phosphotyrosyl phosphatase
activator (PTPA)

[145459]of protein phosphatase 2A. Genomics 28: 261–272, 1995.

[145460]

[145461]

[145462]13613.

[145463]Andre, E.; Becker–Andre, M.: Expression of an N–
terminally truncated

[145464]form of human focal adhesion kinase in brain. Biochem.
Biophys. Res.

[145465]Commun. 190: 140–147, 1993.

[145466]

[145467]

[145468]13614.

[145469]Fiedorek, F. T., Jr.; Kay, E. S.: Mapping of the focal adhe–
sion

[145470]kinase (Fadk) gene to mouse chromosome 15 and human
chromosome 8. Mammalian

[145471]Genome 6: 123–126, 1995.

[145472]

[145473]

[145474]13615.

[145475]Polte, T. R.; Hanks, S. K.: Interaction between focal adhesion

[145476]kinase and Crk-associated tyrosine kinase substrate
p130-Cas. Proc.

[145477]Nat. Acad. Sci. 92: 10678–10682, 1995.

[145478]

[145479]

[145480]13616.

[145481]Schaller, M. D.; Borgman, C.; Cobb, B. S.; Vines, R. R.;
Reynolds,

[145482]A. B.; Parsons, J. T.: pp125(FAK), a structurally distinctive
protein-tyrosine

[145483]kinase associated with focal adhesions. Proc. Nat. Acad.
Sci. 89:

[145484]5192–5196, 1992.

[145485]

[145486]

[145487]13617.

[145488]Arnemann, J.; Epplen, J. T.; Cooke, H. J.; Sauermann, U.;

Engel,

[145489]W.; Schmidtke, J.: A human Y-chromosomal DNA sequence expressed in

[145490]testicular tissue. Nucleic Acids Res. 15: 8713–8724, 1987.

[145491]

[145492]

[145493]13618.

[145494]Arnemann, J.; Jakubiczka, S.; Thuring, S.; Schmidtke, J.:
Cloning

[145495]and sequence analysis of a human Y-chromosome-derived, testicular

[145496]cDNA, TSPY. Genomics 11: 108–114, 1991.

[145497]

[145498]

[145499]13619.

[145500]Guttenbach, M.; Muller, U.; Schmid, M.: A human moderately repeated

[145501]Y-specific DNA sequence is evolutionarily conserved in
the Y chromosome

[145502]of the great apes. Genomics 13: 363–367, 1992.

[145503]

[145504]

[145505]13620.

[145506]Jakubiczka, S.; Schnieders, F.; Schmidtke, J.: A bovine homologue

[145507]of the human TSPY gene. Genomics 17: 732–735, 1993.

[145508]

[145509]

[145510]13621.

[145511]Lau, Y.–F. C.: Sex chromosome genetics '99: gonadoblastoma, testicular

[145512]and prostate cancers, and the TSPY gene. Am. J. Hum. Genet. 64:

[145513]921–927, 1999.

[145514]

[145515]

[145516]13622.

[145517]Manz, E.; Schnieders, F.; Brechlin, A. M.; Schmidtke, J.: TSPY-related

[145518]sequences represent a microheterogeneous gene family organized as

[145519]constitutive elements in DYZ5 tandem repeat units on the human Y chromosome. Genomics 17:

[145520]726–731, 1993.

[145521]

[145522]

[145523]13623.

[145524]Mazeyrat, S.; Mitchell, M. J.: Rodent Y chromosome TSPY gene is

[145525]functional in rat and non-functional in mouse. Hum. Molec. Genet. 7:

[145526]557–562, 1998.

[145527]

[145528]

[145529]13624.

[145530]Schnieders, F.; Dork, T.; Arnemann, J.; Vogel, T.; Werner, M.;

[145531]Schmidtke, J.: Testis-specific protein, Y-encoded (TSPY) expression

[145532]in testicular tissues. Hum. Molec. Genet. 5: 1801–1807,
1996.

[145533]

[145534]

[145535]13625.

[145536]Vogel, T.; Boettger–Tong, H.; Nanda, I.; Dechend, F.; Ag–
ulnik,

[145537]A. I.; Bishop, C. E.; Schmid, M.; Schmidtke, J.: A murine
TSPY. Chromosome

[145538]Res. 6: 35–40, 1998.

[145539]

[145540]

[145541]13626.

[145542]Zhang, J. S.; Yang–Feng, T. L.; Muller, U.; Mohandas, T. K.;
de

[145543]Jong, P. J.; Lau, Y.–F. C.: Molecular isolation and charac–
terization

[145544]of an expressed gene from the human Y–chromosome.
Hum. Molec. Genet. 1:

[145545]717–726, 1992.

[145546]

[145547]

[145548]13627.

[145549]Donnelly, P.; Tavaré, S.; Balding, D. J.; Griffiths, R. C.: Estimating
mating

[145550]the age of the common ancestor of men from the ZFY intron.(Letter) Science 272:

[145551]1357–1359, 1996.

[145552]

[145553]

[145554]13628.

[145555]Dorit, R. L.; Akashi, H.; Gilbert, W.: Absence of polymorphism
phism

[145556]at the ZFY locus on the human Y chromosome. Science
268: 1183–1185,

[145557]1995.

[145558]

[145559]

[145560]13629.

[145561]Dorit, R. L.; Akashi, H.; Gilbert, W.: Estimating the age of the

[145562]common ancestor of men from the ZFY intron.(Letter) Science 272:

[145563]1361–1362, 1996.

[145564]

[145565]

[145566]13630.

[145567]Fu, Y.–X.; Li, W.–H.: Estimating the age of the common ancestor

[145568]of men from the ZFY intron.(Letter) Science 272: 1356–1357, 1996.

[145569]

[145570]

[145571]13631.

[145572]Mahaffey, C. L.; Bayleran, J. K.; Yeh, G. Y.; Lee, T. C.; Page,

[145573]D. C.; Simpson, E. M.: Intron/exon structure confirms that mouse

[145574]Zfy1 and Zfy2 are members of the ZFY gene family. Genomics 41: 123–127,

[145575]1997.

[145576]

[145577]

[145578]13632.

[145579]Palmer, M. S.; Berta, P.; Sinclair, A. H.; Pym, B.; Goodfel-
low,

[145580]P. N.: Comparison of human ZFY and ZFX transcripts. Proc.
Nat. Acad.

[145581]Sci. 87: 1681–1685, 1990.

[145582]

[145583]

[145584]13633.

[145585]Rogers, J.; Samollow, P. B.; Comuzzie, A. G.: Estimating
the

[145586]age of the common ancestor of men from the ZFY in-
tron.(Letter) Science 272:

[145587]1360–1361, 1996.

[145588]

[145589]

[145590]13634.

[145591]Sinclair, A. H.; Foster, J. W.; Spencer, J. A.; Page, D. C.;
Palmer,

[145592]M.; Goodfellow, P. N.; Graves, J. A. M.: Sequences homolo-
gous to

[145593]ZFY, a candidate human sex-determining gene, are auto-
somal in marsupials. Nature 336:

[145594]780-783, 1988.

[145595]

[145596]

[145597]13635.

[145598]Slattery, J. P.; Sanner-Wachter, L.; O'Brien, S. J.: Novel gene

[145599]conversion between X-Y homologues located in the non-
recombining region

[145600]of the Y chromosome in Felidae (mammalia). Proc. Nat.
Acad. Sci. 97:

[145601]5307-5312, 2000.

[145602]

[145603]

[145604]13636.

[145605]Weiss, G.; von Haeseler: Estimating the age of the common ancestor

[145606]of men from the ZFY intron.(Letter) Science 272: 1359–1360, 1996.

[145607]

[145608]

[145609]13637.

[145610]Zambrowicz, B. P.; Findley, S. D.; Simpson, E. M.; Page, D. C.;

[145611]Palmiter, R. D.: Characterization of the murine Zfy1 and Zfy2 promoters. Genomics 24:

[145612]406–408, 1994.

[145613]

[145614]

[145615]13638.

[145616]Mach, B.; Steimle, V.; Martinez–Soria, E.; Reith, W.: Regulation

[145617]of MHC class II genes: lessons from a disease. Annu. Rev. Immun. 14:

[145618]301–331, 1996.

[145619]

[145620]

[145621]13639.

[145622]Reith, W.: Personal Communication. Geneva, Switzerland
5/30/1997.

[145623]

[145624]

[145625]13640.

[145626]Scholl, T.; Mahanta, S. K.; Strominger, J. L.: Specific complex
plex

[145627]formation between the type II bare lymphocyte syndrome-
associated

[145628]transactivators CIITA and RFX5. Proc. Nat. Acad. Sci. 94:
6330-6334,

[145629]1997.

[145630]

[145631]

[145632]13641.

[145633]Emery, P.; Durand, B.; Mach, B.; Reith, W.: RFX proteins, a

novel

[145634]family of DNA binding proteins conserved in the eukary-
otic kingdom. Nucleic

[145635]Acids Res. 24: 803-807, 1996.

[145636]

[145637]

[145638]13642.

[145639]Pugliati, L.; Reith, W.; Fey, S.; Mach, B.: Mapping the RF-X
gene,

[145640]encoding a DNA-binding protein controlling HLA class II
gene expression,

[145641]to 19p13.(Abstract) Cytogenet. Cell Genet. 51: 1061 only,
1989.

[145642]

[145643]

[145644]13643.

[145645]Mizuta, M.; Inagaki, N.; Nemoto, Y.; Matsukura, S.; Taka-
hashi,

[145646]M.; Seino, S.: Synaptotagmin III is a novel isoform of rat
synaptotagmin

[145647]expressed in endocrine and neuronal cells. J. Biol. Chem.

269: 11675–11678,

[145648]1994.

[145649]

[145650]

[145651]13644.

[145652]Li, X.-J.; Wang, D.-Y.; Zhu, Y.; Guo, R.-J.; Wang, X.-D.;

Lubomir,

[145653]K.; Mukai, K.; Sasaki, H.; Yoshida, H.; Oka, T.; Machinami,

R.; Shinmura,

[145654]K.; Tanaka, M.; Sugimura, H.: Mxi1 mutations in human
neurofibrosarcomas. Jpn.

[145655]J. Cancer Res. 90: 740–746, 1999.

[145656]

[145657]

[145658]13645.

[145659]Prochownik, E. V.; Grove, L. E.; Deubler, D.; Zhu, X. L.;

Stephenson,

[145660]R. A.; Rohr, L. R.; Yin, X.; Brothman, A. R.: Commonly oc-
curring

[145661]loss and mutation of the MXI1 gene in prostate cancer.

Genes Chromosomes

[145662]Cancer 22: 295–304, 1998.

[145663]

[145664]

[145665]13646.

[145666]Schreiber–Agus, N.; Meng, Y.; Hoang, T.; Hou, H., Jr.;
Chen, K.;

[145667]Greenberg, R.; Cordon–Cardo, C.; Lee, H.–W.; DePinho, R.
A.: Role

[145668]of Mxi1 in ageing organ systems and the regulation of
normal and neoplastic

[145669]growth. Nature 393: 483–487, 1998.

[145670]

[145671]

[145672]13647.

[145673]Shapiro, D. N.; Valentine, V.; Eagle, L.; Yin, X.; Morris, S.
W.;

[145674]Prochownik, E. V.: Assignment of the human MAD and
MXI1 genes to

[145675]chromosomes 2p12–p13 and 10q24–q25. Genomics 23:

282–285, 1994.

[145676]

[145677]

[145678]13648.

[145679]Wechsler, D. S.; Hawkins, A. L.; Li, X.; Jabs, E. W.; Griffin,

[145680]C. A.; Dang, C. V.: Localization of the human Mxi1 tran-
scription

[145681]factor gene (MXI1) to chromosome 10q24–q25. Genomics
21: 669–672,

[145682]1994.

[145683]

[145684]

[145685]13649.

[145686]Wechsler, D. S.; Shelly, C. A.; Dang, C. V.: Genomic orga-
nization

[145687]of human MXI1, a putative tumor suppressor gene. Ge-
nomics 32: 466–470,

[145688]1996.

[145689]

[145690]

[145691]13650.

[145692]Zervos, A. S.; Gyuris, J.; Brent, R.: Mxi1, a protein that
specifically

[145693]interacts with Max to bind Myc–Max recognition sites. Cell
72: 223–232,

[145694]1993. Note: Erratum: Cell 79: 389 only, 1994.

[145695]

[145696]

[145697]13651.

[145698]Boie, Y.; Rushmore, T. H.; Darmon–Goodwin, A.; Grygor–
czyk, R.;

[145699]Slipetz, D. M.; Metters, K. M.; Abramovitz, M.: Cloning and
expression

[145700]of a cDNA for the human prostanoid IP receptor. J. Biol.
Chem. 269:

[145701]12173–12178, 1994.

[145702]

[145703]

[145704]13652.

[145705]Ishikawa, T.; Tamai, Y.; Rochelle, J. M.; Hirata, M.; Namba, T.;

[145706]Sugimoto, Y.; Ichikawa, A.; Narumiya, S.; Taketo, M. M.; Seldin, M.

[145707]F.: Mapping of the genes encoding mouse prostaglandin D, E, and F

[145708]and prostacyclin receptors. Genomics 32: 285–288, 1996.

[145709]

[145710]

[145711]13653.

[145712]Murata, T.; Ushikubi, F.; Matsuoka, T.; Hirata, M.; Yamasaki, A.;

[145713]Sugimoto, Y.; Ichikawa, A.; Aze, Y.; Tanaka, T.; Yoshida, N.; Ueno,

[145714]A.; Oh-ishi, S.; Narumiya, S.: Altered pain perception and inflammatory

[145715]response in mice lacking prostacyclin receptor. Nature 388: 678–682,

[145716]1997.

[145717]

[145718]

[145719]13654.

[145720]Ogawa, Y.; Tanaka, I.; Inoue, M.; Yoshitake, Y.; Isse, N.;
Nakagawa,

[145721]O.; Usui, T.; Itoh, H.; Yoshimasa, T.; Narumiya, S.; Nakao,
K.: Structural

[145722]organization and chromosomal assignment of the human
prostacyclin

[145723]receptor gene. Genomics 27: 142–148, 1995.

[145724]

[145725]

[145726]13655.

[145727]Hoffman, I.; Balling, R.: Chromosomal localization of the
murine

[145728]cadherin–11. Mammalian Genome 6: 304 only, 1995.

[145729]

[145730]

[145731]13656.

[145732]Okazaki, M.; Takeshita, S.; Kawai, S.; Kikuno, R.; Tsu–
jimura, A.;

[145733]Kudo, A.; Amann, E.: Molecular cloning and characteriza–

tion of OB-cadherin,

[145734]a new member of cadherin family expressed in osteoblasts. J. Biol.

[145735]Chem. 269: 12092–12098, 1994.

[145736]

[145737]

[145738]13657.

[145739]Tanihara, H.; Sano, K.; Heimark, R. L.; St. John, T.; Suzuki, S.

[145740]: Cloning of five human cadherins clarifies characteristic features

[145741]of cadherin extracellular domain and provides further evidence for

[145742]two structurally different types of cadherin. Cell Adhes. Commun. 2:

[145743]15–26, 1994.

[145744]

[145745]

[145746]13658.

[145747]Christie, P. T.; Curley, A.; Nesbit, M. A.; Chapman, C.; Genet,

[145748]S.; Harper, P. S.; Keeling, S. L.; Wilkie, A. O. M.; Winter, R. M.;

[145749]Thakker, R. V.: Mutational analysis in X-linked spondy-

loepiphyseal

[145750]dysplasia tarda. J. Clin. Endocr. Metab. 86: 3233–3236,
2001.

[145751]

[145752]

[145753]13659.

[145754]Fiedler, J.; Bittner, M.; Puhl, W.; Brenner, R. E.: Mutations

[145755]in the X-linked spondyloepiphyseal dysplasia tarda (SEDL)
coding sequence

[145756]are not a common cause of early primary osteoarthritis in
men. (Letter) Clin.

[145757]Genet. 62: 94–95, 2002.

[145758]

[145759]

[145760]13660.

[145761]Gecz, J.; Hillman, M. A.; Gedeon, A. K.; Cox, T. C.; Baker,
E.;

[145762]Mulley, J. C.: Gene structure and expression study of the
SEDL gene

[145763]for spondyloepiphyseal dysplasia tarda. Genomics 69:

242–251, 2000.

[145764]

[145765]

[145766]13661.

[145767]Gedeon, A. K.; Colley, A.; Jamieson, R.; Thompson, E. M.;
Rogers,

[145768]J.; Sillence, D.; Tiller, G. E.; Mulley, J. C.; Gecz, J.: Identifi-
cation

[145769]of the gene (SEDL) causing X-linked spondyloepiphyseal
dysplasia tarda. Nature

[145770]Genet. 22: 400–404, 1999.

[145771]

[145772]

[145773]13662.

[145774]Gedeon, A. K.; Tiller, G. E.; Le Merrer, M.; Heuertz, S.;
Tranebjaerg,

[145775]L.; Chitayat, D.; Robertson, S.; Glass, I. A.; Savarirayan, R.;
Cole,

[145776]W. G.; Rimoin, D. L.; Kousseff, B. G.; Ohashi, H.; Zabel, B.;
Munnich,

[145777]A.; Gecz, J.; Mulley, J. C.: The molecular basis of X-linked

spondyloepiphyseal

[145778]dysplasia tarda. Am. J. Hum. Genet. 68: 1386–1397,
2001.

[145779]

[145780]

[145781]13663.

[145782]Grunebaum, E.; Arpaia, E.; MacKenzie, J. J.; Fitzpatrick, J.;
Ray,

[145783]P. N.; Roifman, C. M.: A missense mutation in the SEDL
gene results

[145784]in delayed onset of X linked spondyloepiphyseal dysplasia
in a large

[145785]pedigree. (Letter) J. Med. Genet. 38: 409–411, 2001.

[145786]

[145787]

[145788]13664.

[145789]Mumm, S.; Christie, P. T.; Finnegan, P.; Jones, J.; Dixon, P.
H.;

[145790]Pannett, A. A. J.; Harding, B.; Gottesman, G. S.; Thakker, R.
V.;

[145791]Whyte, M. P.: A five–base pair deletion in the sedlin gene

causes

[145792]spondyloepiphyseal dysplasia tarda in a six-generation
Arkansas kindred. J.

[145793]Clin. Endocr. Metab. 85: 3343–3347, 2000.

[145794]

[145795]

[145796]13665.

[145797]Mumm, S.; Zhang, X.; Vacca, M.; D'Esposito, M.; Whyte, M.
P.:

[145798]The sedlin gene for spondyloepiphyseal dysplasia tarda
escapes X-inactivation

[145799]and contains a non-canonical splice site. Gene 273:
285–293, 2001.

[145800]

[145801]

[145802]13666.

[145803]Takahashi, T.; Takahashi, I.; Tsuchida, S.; Oyama, K.; Ko-
matsu,

[145804]M.; Saito, H.; Takada, G.: An SEDL gene mutation in a
Japanese kindred

[145805]of X-linked spondyloepiphyseal dysplasia tarda. (Letter)

Clin. Genet. 61:

[145806]319–320, 2002.

[145807]

[145808]

[145809]13667.

[145810]Tiller, G. E.; Hannig, V. L.; Dozier, D.; Carrel, L.; Trevarthen,

[145811]K. C.; Wilcox, W. R.; Mundlos, S.; Haines, J. L.; Gedeon, A. K.; Gecz,

[145812]J.: A recurrent RNA-splicing mutation in the SEDL gene causes X-linked

[145813]spondyloepiphyseal dysplasia tarda. Am. J. Hum. Genet. 68: 1398–1407,

[145814]2001.

[145815]

[145816]

[145817]13668.

[145818]Whyte, M. P.; Gottesman, G. S.; Eddy, M. C.; McAlister, W. H.

[145819]: X-linked recessive spondyloepiphyseal dysplasia tarda:

clinical

[145820]and radiographic evolution in a 6-generation kindred and
review of

[145821]the literature. *Medicine* 78: 9–25, 1999.

[145822]

[145823]

[145824]13669.

[145825]Becker, M. A.; Heidler, S. A.; Bell, G. I.; Seino, S.; Le Beau,

[145826]M. M.; Westbrook, C. A.; Neuman, W.; Shapiro, L. J.; Mo-
handas, T.

[145827]K.; Roessler, B. J.; Palella, T. D.: Cloning of cDNAs for hu-
man phosphoribosylpyrophosphate

[145828]synthetases 1 and 2 and X chromosome localization of
PRPS1 and PRPS2

[145829]genes. *Genomics* 8: 555–561, 1990.

[145830]

[145831]

[145832]13670.

[145833]Buchner, G.; Montini, E.; Andolfi, G.; Quaderi, N.; Cainarca,
S.;

[145834]Messali, S.; Bassi, M. T.; Ballabio, A.; Meroni, G.; Franco, B.:

[145835]MID2, a homologue of the Opitz syndrome gene MID1: similarities in

[145836]subcellular localization and differences in expression during development. Hum.

[145837]Molec. Genet. 8: 1397–1407, 1999.

[145838]

[145839]

[145840]13671.

[145841]Braverman, N.; Lin, P.; Moebius, F. F.; Obie, C.; Moser, A.; Glossmann,

[145842]H.; Wilcox, W. R.; Rimoïn, D. L.; Smith, M.; Kratz, L.; Kelley, R.

[145843]I.; Valle, D.: Mutations in the gene encoding 3–beta–hydroxysteroid–delta(8),delta(7)–isomerase

[145844]cause X–linked dominant Conradi–Hunermann syndrome. Nature Genet. 22:

[145845]291–294, 1999.

[145846]

[145847]

[145848]13672.

[145849]Cho, S. Y.; Kim, J. H.; Paik, Y. K.: Cholesterol biosynthesis

[145850]from lanosterol: differential inhibition of sterol delta

8-isomerase

[145851]and other lanosterol-converting enzymes by tamoxifen.

Molec. Cells 8:

[145852]233-239, 1998.

[145853]

[145854]

[145855]13673.

[145856]Clayton, P. T.; Kalter, D. C.; Atherton, D. J.; Besley, G. T.;

[145857]Broadhead, D. M.: Peroxisomal enzyme deficiency in X-

linked dominant

[145858]Conradi-Hunermann syndrome. J. Inherit. Metab. Dis. 12:

358-360,

[145859]1989.

[145860]

[145861]

[145862]13674.

- [145863]Derry, J. M. J.; Gormally, E.; Means, G. D.; Zhao, W.;
Meindl,
[145864]A.; Kelley, R. I.; Boyd, Y.; Herman, G. E.: Mutations in a
delta(8)-delta(7)
[145865]sterol isomerase in the tattered mouse and X-linked dom-
inant chondrodysplasia
[145866]punctata. Nature Genet. 22: 286-290, 1999.
- [145867]
- [145868]
- [145869]13675.
- [145870]Grange, D. K.; Kratz, L. E.; Braverman, N. E.; Kelley, R. I.:
[145871]CHILD syndrome caused by deficiency of
3-beta-hydroxysteroid-delta-8,delta-7-isomerase. Am.
[145872]J. Med. Genet. 90: 328-335, 2000.
- [145873]
- [145874]
- [145875]13676.
- [145876]Hanner, M.; Moebius, F. F.; Weber, F.; Grabner, M.; Striess-
nig,
[145877]J.; Glossmann, H.: Phenylalkylamine Ca(2+) antagonist

binding protein:

[145878]molecular cloning, tissue distribution, and heterologous expression. J.

[145879]Biol. Chem. 270: 7551–7557, 1995.

[145880]

[145881]

[145882]13677.

[145883]Has, C.; Bruckner–Tuderman, L.; Muller, D.; Floeth, M.; Folkers,

[145884]E.; Donnai, D.; Traupe, H.: The Conradi–Hunermann–Happle syndrome

[145885](CDPX2) and emopamil binding protein: novel mutations, and somatic

[145886]and gonadal mosaicism. Hum. Molec. Genet. 9: 1951–1955, 2000.

[145887]

[145888]

[145889]13678.

[145890]Holmes, R. D.; Wilson, G. N.; Hajra, A. K.: Peroxisomal enzyme

[145891]deficiency in the Conradi–Hunerman (sic) form of chon–

drodysplasia

[145892]punctata. New Eng. J. Med. (Letter) 316: 1608 only, 1987.

[145893]

[145894]

[145895]13679.

[145896]Ikegawa, S.; Ohashi, H.; Ogata, T.; Honda, A.; Tsukahara,
M.; Kubo,

[145897]T.; Kimizuka, M.; Shimode, M.; Hasegawa, T.; Nishimura,
G.; Nakamura,

[145898]Y.: Novel and recurrent EBP mutations in X-linked domi-
nant chondrodysplasia

[145899]punctata. Am. J. Med. Genet. 94: 300–305, 2000.

[145900]

[145901]

[145902]13680.

[145903]Kelley, R. I.; Wilcox, W. G.; Smith, M.; Kratz, L. E.; Moser,

[145904]A.; Rimoin, D. S.: Abnormal sterol metabolism in patients
with Conradi–Hunermann–Happle

[145905]syndrome and sporadic lethal chondrodysplasia punctata.
Am. J. Med.

[145906]Genet. 83: 213–219, 1999.

[145907]

[145908]

[145909]13681.

[145910]Aksoy, S.; Brandriff, B. F.; Ward, A.; Little, P. F. R.; Weinshilboum,

[145911]R. M.: Human nicotinamide N-methyltransferase gene: molecular cloning,

[145912]structural characterization and chromosomal localization. Genomics 29:

[145913]555–561, 1995.

[145914]

[145915]

[145916]13682.

[145917]Aksoy, S.; Szumlanski, C. L.; Weinshilboum, R. M.: Human liver

[145918]nicotinamide N-methyltransferase: cDNA cloning, expression, and biochemical

[145919]characterization. J. Biol. Chem. 269: 14835–14840, 1994.

[145920]

[145921]

[145922]13683.

[145923]Bennett, B. D.; Wang, Z.; Kuang, W.-J.; Wang, A.; Groopman, J.

[145924]E.; Goeddel, D. V.; Scadden, D. T.: Cloning and characterization

[145925]of HTK, a novel transmembrane tyrosine kinase of the EPH subfamily. J.

[145926]Biol. Chem. 269: 14211-14218, 1994.

[145927]

[145928]

[145929]13684.

[145930]Berclaz, G.; Andres, A.-C.; Albrecht, D.; Dreher, E.; Ziemiecki,

[145931]A.; Gusterson, B. A.; Crompton, M. R.: Expression of the receptor

[145932]protein tyrosine kinase myk-1/htk in normal and malignant mammary

[145933]epithelium. Biochem. Biophys. Res. Commun. 226: 869-875, 1996.

[145934]

[145935]

[145936]13685.

[145937]Gerety, S. S.; Wang, H. U.; Chen, Z.-F.; Anderson, D. J.:

Symmetrical

[145938]mutant phenotypes of the receptor EphB4 and its specific
transmembrane

[145939]ligand ephrin-B2 in cardiovascular development. Molec.
Cell 4: 403-414,

[145940]1999.

[145941]

[145942]

[145943]13686.

[145944]Moynihan, T. P.; Ardley, H. C.; Leek, J. P.; Thompson, J.;
Brindle,

[145945]N. S.; Markham, A. F.; Robinson, P. A.: Characterization of
a human

[145946]ubiquitin-conjugating enzyme gene UBE2L3. Mammalian
Genome 7: 520-525,

[145947]1996.

[145948]

[145949]

[145950]13687.

[145951]Flanagan, J. R.; Becker, K. G.; Ennist, D. L.; Gleason, S. L.;

[145952]Driggers, P. H.; Levi, B.-Z.; Appella, E.; Ozato, K.: Cloning
of

[145953]a negative transcription factor that binds to the upstream
conserved

[145954]region of Moloney murine leukemia virus. Molec. Cell.
Biol. 12:

[145955]38–44, 1992.

[145956]

[145957]

[145958]13688.

[145959]Hariharan, N.; Kelley, D. E.; Perry, R. P.: Delta, a transcrip-
tion

[145960]factor that binds to downstream elements in several poly-
merase II

[145961]promoters, is a functionally versatile zinc finger protein.
Proc.

[145962]Nat. Acad. Sci. 88: 9799–9803, 1991.

[145963]

[145964]

[145965]13689.

[145966]Oei, S. L.; Shi, Y.: Transcription factor Yin Yang 1 stimulates

[145967]poly(ADP-ribosyl)ation and DNA repair. Biochem. Biophys. Res. Commun. 284:

[145968]450-454, 2001.

[145969]

[145970]

[145971]13690.

[145972]Park, K.; Atchison, M. L.: Isolation of a candidate repressor/activator,

[145973]NF-E1 (YY-1, delta), that binds to the immunoglobulin kappa 3-prime

[145974]enhancer and the immunoglobulin heavy-chain micro-E1 site.. Proc.

[145975]Nat. Acad. Sci. 88: 9804-9808, 1991.

[145976]

[145977]

[145978]13691.

[145979]Shi, Y.; Seto, E.; Chang, L.-S.; Shenk, T.: Transcriptional repression

[145980]by YY1, a human GLI-Kruppel-related protein, and relief of repression

[145981]by adenovirus E1A protein. Cell 67: 377-388, 1991.

[145982]

[145983]

[145984]13692.

[145985]Yao, Y.-L.; Dupont, B. R.; Ghosh, S.; Fang, Y.; Leach, R. J.; Seto,

[145986]E.: Cloning, chromosomal localization and promoter analysis of the

[145987]human transcription factor YY1. Nucleic Acids Res. 26: 3776-3783,

[145988]1998.

[145989]

[145990]

[145991]13693.

[145992]Zhu, W.; Lossie, A. C.; Camper, S. A.; Gumucio, D. L.:

Chromosomal

[145993] localization of the transcription factor YY1 in the mouse and human. Mammalian

[145994] Genome 5: 234–236, 1994.

[145995]

[145996]

[145997] 13694.

[145998] Hakimi, M.–A.; Bochar, D. A.; Schmiesing, J. A.; Dong, Y.; Barak,

[145999] O. G.; Speicher, D. W.; Yokomori, K.; Shiekhattar, R.: A chromatin

[146000] remodelling complex that loads cohesin onto human chromosomes. Nature 418:

[146001] 994–998, 2002.

[146002]

[146003]

[146004] 13695.

[146005] Muchardt, C.; Reyes, J. C.; Bourachot, B.; Leguoy, E.; Yaniv, M.

[146006]: The hbrm and BRG–1 proteins, components of the hu–

man SNF/SWI complex,
[146007]are phosphorylated and excluded from the condensed
chromosomes during
[146008]mitosis. EMBO J. 15: 3394–3402, 1996.

[146009]

[146010]

[146011]13696.

[146012]Muchardt, C.; Yaniv, M.: A human homologue of *Saccharomyces cerevisiae*
SNF2/SWI2 and *Drosophila* brm genes potentiates tran-
scriptional activation
[146013]by the glucocorticoid receptor. EMBO J. 12: 4279–4290,
1993.

[146015]

[146016]

[146017]13697.

[146018]Muchardt, C.; Yaniv, M.; Mattei, M.–G.: Assignment of
HBRM, the
[146019]human homolog of *S. cerevisiae* SNF2/SWI2 and
Drosophila brm genes,
[146020]to chromosome region 9p23–p24, by in situ hybridization.

Mammalian

[146021]Genome 5: 241–243, 1994.

[146022]

[146023]

[146024]13698.

[146025]Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.;

Kawarabayasi,

[146026]Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.:

Prediction

[146027]of the coding sequences of unidentified human genes. I.

The coding

[146028]sequences of 40 new genes (KIAA0001–KIAA0040) de-

duced by analysis

[146029]of randomly samples cDNA clones from human immature

myeloid cell line

[146030]KG-1. DNA Res. 1: 27–35, 1994.

[146031]

[146032]

[146033]13699.

[146034]Kikuno, R.; Nagase, T.; Ishikawa, K.; Hirose, M.; Miya-

jima, N.;

[146035]Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction
of the

[146036]coding sequences of unidentified human genes. XIV. The
complete sequences

[146037]of 100 new cDNA clones from brain which code for large
proteins in

[146038]vitro. DNA Res. 6: 197–205, 1999.

[146039]

[146040]

[146041]13700.

[146042]Polsky, D.; Young, A. Z.; Busam, K. J.; Alani, R. M.: The
transcriptional

[146043]repressor of p16/Ink4a, Id1, is up-regulated in early
melanomas. Cancer

[146044]Res. 61: 6008–6011, 2001.

[146045]

[146046]

[146047]13701.

[146048]Walker, J. L.; Dixon, J.; Fenton, C. R.; Hungerford, J.;

Lynch,

[146049]S. A.; Stenhouses, S. A. R.; Christian, A.; Craig, I. W.: Two
new

[146050]mutations in exon 3 of the NDP gene: S73X and S101F as-
sociated with

[146051]severe and less severe ocular phenotype, respectively.

Hum. Mutat. 9:

[146052]53–56, 1997.

[146053]

[146054]

[146055]13702.

[146056]Lee, D. K.; Horikoshi, M.; Roeder, R. G.: Interaction of
TFIID

[146057]in the minor groove of the TATA element. Cell 67:
1241–1250, 1991.

[146058]

[146059]

[146060]13703.

[146061]Wilson, R. B.; Kiledjian, M.; Shen, C.–P.; Benezra, R.;
Zwollo,

[146062]P.; Dymecki, S. M.; Desiderio, S. V.; Kadesch, T.: Repres–

sion of

[146063]immunoglobulin enhancers by the helix-loop-helix protein Id: implications

[146064]for B-lymphoid-cell development. Molec. Cell. Biol. 11: 6185-6191,

[146065]1991.

[146066]

[146067]

[146068]13704.

[146069]Battaglia, G.; Princivale, A.; Forti, F.; Lizier, C.; Zeviani,

[146070]M.: Expression of the SMN gene, the spinal muscular atrophy determining

[146071]gene, in the mammalian central nervous system. Hum. Molec. Genet. 6:

[146072]1961-1971, 1997.

[146073]

[146074]

[146075]13705.

[146076]Burglen, L.; Lefebvre, S.; Clermont, O.; Burlet, P.; Viollet, L.;

[146077]Cruaud, C.; Munnich, A.; Melki, J.: Structure and organization of

[146078]the human survival motor neurone (SMN) gene. Genomics 32: 479–482,

[146079]1996.

[146080]

[146081]

[146082]13706.

[146083]Burglen, L.; Spiegel, R.; Ignatius, J.; Cobben, J. M.; Landrieu,

[146084]P.; Lefebvre, S.; Munnich, A.; Melki, J.: SMN gene deletion in variant

[146085]of infantile spinal muscular atrophy. (Letter) Lancet 346: 316–317,

[146086]1995.

[146087]

[146088]

[146089]13707.

[146090]Bussaglia, E.; Clermont, O.; Tizzano, E.; Lefebvre, S.; Burglen,

[146091]L.; Cruaud, C.; Urtizberea, J. A.; Colomer, J.; Munnich, A.;
Baiget,

[146092]M.; Melki, J.: A frame-shift deletion in the survival motor
neuron

[146093]gene in Spanish spinal muscular atrophy patients. Nature
Genet. 11:

[146094]335–337, 1995.

[146095]

[146096]

[146097]13708.

[146098]Callebaut, I.; Mornon, J. P.: The human EBNA-2 coactivator
p100:

[146099]multidomain organization and relationship to the staphy-
lococcal nuclease

[146100]fold and to the tudor protein involved in Drosophila
melanogaster

[146101]development. Biochem. J. 321: 125–132, 1997.

[146102]

[146103]

[146104]13709.

[146105]Campbell, L.; Daniels, R. J.; Dubowitz, V.; Davies, K. E.:

Maternal

[146106]mosaicism for a second mutational event in a type I spinal
muscular

[146107]atrophy family. Am. J. Hum. Genet. 63: 37–44, 1998.

[146108]

[146109]

[146110]13710.

[146111]Campbell, L.; Hunter, K. M. D.; Mohaghegh, P.; Tinsley, J.
M.;

[146112]Brasch, M. A.; Davies, K. E.: Direct interaction of Smn with
dp103,

[146113]a putative RNA helicase: a role for Smn in transcription
regulation? Hum.

[146114]Molec. Genet. 9: 1093–1100, 2000.

[146115]

[146116]

[146117]13711.

[146118]Chen, K.–L.; Wang, Y. L.; Rennert, H.; Joshi, I.; Mills, J. K.;

[146119]Leonard, D. G. B.; Wilson, R. B.: Duplications and de novo

deletions

[146120]of the SMNt gene demonstrated by fluorescence-based
carrier testing

[146121]for spinal muscular atrophy. Am. J. Med. Genet. 85:
463–469, 1999.

[146122]

[146123]

[146124]13712.

[146125]Warburg, M.: Norrie's disease: a new hereditary bilateral
pseudotumour

[146126]of the retina. Acta Ophthal. (Copenh) 39: 757–772, 1961.

[146127]

[146128]

[146129]13713.

[146130]Warburg, M.: Norrie's disease (atrofia bulborum heredi-
taria). Acta

[146131]Ophthal. 41: 134–146, 1963.

[146132]

[146133]

[146134]13714.

[146135]Warburg, M.: Norrie's disease, a congenital progressive
oculo-acoustico-cerebral

[146136]degeneration. Acta Ophthal. 89 (suppl.): 1-147, 1966.

[146137]

[146138]

[146139]13715.

[146140]Warburg, M.; Hauge, M.; Sanger, R.: Norrie's disease and
the

[146141]Xg blood group system: linkage data. Acta Genet. Statist.
Med. 15:

[146142]103-115, 1965.

[146143]

[146144]

[146145]13716.

[146146]Whitnall, S. E.; Norman, R. M.: Microphthalmia and the vi-
sual

[146147]pathways: a case associated with blindness and imbecility,
and sex-linked. Brit.

[146148]J. Ophthal. 24: 229-244, 1940.

[146149]

[146150]

[146151]13717.

[146152]Wilson, W. M. G.: Congenital blindness (pseudoglioma) occurring

[146153]as a sex-linked developmental anomaly. Canad. Med. Assoc. J. 60:

[146154]580-584, 1949.

[146155]

[146156]

[146157]13718.

[146158]Wolff, G.; Mayerova, A.; Wienker, T. F.; Atalianis, P.; Ioannou,

[146159]P.; Warburg, M.: Clinical reinvestigation and linkage analysis in

[146160]the family with Episkopi blindness (Norrie disease). J. Med. Genet. 29:

[146161]816-819, 1992.

[146162]

[146163]

[146164]13719.

[146165]Wong, F.; Goldberg, M. F.; Hao, Y.: Identification of a non-sense

[146166]mutation at codon 128 of the Norrie's disease gene in a male infant. Arch.

[146167]Ophthal. 111: 1553-1557, 1993.

[146168]

[146169]

[146170]13720.

[146171]Woodruff, G.; Newbury-Ecob, R.; Plaha, D. S.; Young, I. D.: Manifesting

[146172]heterozygosity in Norrie's disease?. Brit. J. Ophthal. 77: 813-814,

[146173]1993.

[146174]

[146175]

[146176]13721.

[146177]Zhu, D.; Antonarakis, S. E.; Schmeckpeper, B. J.; Diergaarde,

[146178]P. J.; Greb, A. E.; Maumenee, I. H.: Microdeletion in the X-

chromosome

[146179]and prenatal diagnosis in a family with Norrie disease.

Am. J. Med.

[146180]Genet. 33: 485–488, 1989.

[146181]

[146182]

[146183]13722.

[146184]Avner, P.; Bucan, M.; Arnaud, D.; Lehrach, H.; Rapp, U.: A-
raf

[146185]oncogene localizes on mouse X chromosome to region
some 10–17 centimorgans

[146186]proximal to hypoxanthine phosphoribosyltransferase
gene. Somat. Cell

[146187]Molec. Genet. 13: 267–272, 1987.

[146188]

[146189]

[146190]13723.

[146191]Beck, T. W.; Huleihel, M.; Gunnell, M.; Bonner, T. I.; Rapp,
U.

[146192]R.: The complete coding sequence of the human A-raf-1

oncogene and

[146193]transforming activity of a human A-raf carrying retrovirus.

Nucleic

[146194]Acids Res. 15: 595–609, 1987.

[146195]

[146196]

[146197]13724.

[146198]Huebner, K.; ar-Rushdi, A.; Griffin, C. A.; Isobe, M.; Kozak, C.;

[146199]Emanuel, B. S.; Nagarajan, L.; Cleveland, J. L.; Bonner, T. I.; Goldsborough,

[146200]M. D.; Croce, C. M.; Rapp, U.: Actively transcribed genes in the

[146201]raf oncogene group, located on the X chromosome in mouse and human. Proc.

[146202]Nat. Acad. Sci. 83: 3934–3938, 1986.

[146203]

[146204]

[146205]13725.

[146206]Lee, J.-E.; Beck, T. W.; Brennscheidt, U.; DeGennaro, L. J.;

Rapp,

[146207]U. R.: The complete sequence and promoter activity of the
human A-raf-1

[146208]gene (ARAF1). Genomics 20: 43-55, 1994.

[146209]

[146210]

[146211]13726.

[146212]Mark, G. E.; Seeley, T. W.; Shows, T. B.; Mountz, J. D.: Pks,

[146213]a raf-related sequence in humans. Proc. Nat. Acad. Sci.

83: 6312-6316,

[146214]1986.

[146215]

[146216]

[146217]13727.

[146218]Popescu, N. C.; Mark, G. E.: Localization of the pKs gene, a
raf

[146219]related sequence on human chromosomes X and 7. Onco-
gene 4: 517-519,

[146220]1989.

[146221]

[146222]

[146223]13728.

[146224]Dowdy, S. F.; Fasching, C. L.; Araujo, D.; Lai, K.-M.; Li-
vanos,

[146225]E.; Weissman, B. E.; Stanbridge, E. J.: Suppression of tu-
morigenicity

[146226]in Wilms' tumor by the p15.5-p14 region of chromosome
11. Science 254:

[146227]293-295, 1991.

[146228]

[146229]

[146230]13729.

[146231]Dowdy, S. F.; Lai, K.-M.; Weissman, B. E.; Matsui, Y.;
Hogan, B.

[146232]L. M.; Stanbridge, E. J.: The isolation and characterization
of a

[146233]novel cDNA demonstrating an altered mRNA level in non-
tumorigenic Wilms'

[146234]microcell hybrid cells. Nucleic Acids Res. 19: 5763-5769,
1991.

[146235]

[146236]

[146237]13730.

[146238]Farmer, A. A.; Loftus, T. M.; Mills, A. A.; Sato, K. Y.; Neill,

[146239]J. D.; Tron, T.; Yang, M.; Trumpower, B. L.; Stanbridge, E.

J.: Extreme

[146240]evolutionary conservation of QM, a novel c-Jun associated
transcription

[146241]factor. Hum. Molec. Genet. 3: 723–728, 1994.

[146242]

[146243]

[146244]13731.

[146245]Kaneko, K.; Kobayashi, H.; Onodera, O.; Miyatake, T.;

Tsuji, S.

[146246]: Genomic organization of a cDNA (QM) demonstrating an
altered mRNA

[146247]level in nontumorigenic Wilms' microcell hybrid cells and
its localization

[146248]to Xq28. Hum. Molec. Genet. 1: 529–533, 1992.

[146249]

[146250]

[146251]13732.

[146252]Korn, B.; Sedlacek, Z.; Manca, A.; Kioschis, P.; Konecki, D.;
Lehrach,

[146253]H.; Poustka, A.: A strategy for the selection of transcribed
sequences

[146254]in the Xq28 region. Hum. Molec. Genet. 1: 235–242,
1992.

[146255]

[146256]

[146257]13733.

[146258]van den Ouweland, A. M. W.; Verdijk, M.; Mannens, M. M.
A. M.;

[146259]van Oost, B. A.: The QM gene is X-linked and therefore
not involved

[146260]in suppression of tumorigenesis in Wilms' tumor. Hum.
Molec. Genet. 90:

[146261]144–146, 1992.

[146262]

[146263]

[146264]13734.

[146265]Weissman, B. E.; Saxon, P. J.; Pasquale, S. R.; Jones, G. R.; Geiser,

[146266]A. G.; Stanbridge, E. J.: Introduction of a normal human chromosome

[146267]11 into a Wilms' tumor cell line controls its tumorigenic expression. Science 236:

[146268]175–180, 1987.

[146269]

[146270]

[146271]13735.

[146272]Nishimoto, T.; Sekiguchi, T.; Kai, R.; Yamashita, K.; Takahashi,

[146273]T.; Sekiguchi, M.: Large-scale selection and analysis of temperature-sensitive

[146274]mutants for cell reproduction from BHK cells. Somat. Cell Genet. 8:

[146275]811–812, 1982.

[146276]

[146277]

[146278]13736.

[146279]O'Brien, T.; Tjian, R.: Functional analysis of the human
TAFII250

[146280]N-terminal kinase domain. Molec. Cell 1: 905–911, 1998.

[146281]

[146282]

[146283]13737.

[146284]Ruppert, S.; Wang, E. H.; Tjian, R.: Cloning and expression
of

[146285]human TAF(II)250: a TBP-associated factor implicated in
cell-cycle

[146286]regulation. Nature 362: 175–179, 1993.

[146287]

[146288]

[146289]13738.

[146290]Schwartz, H. E.; Holmes, S.; Meiss, H. K.: Assignment of
temperature-sensitive

[146291]mutations of BHK cells to the X-chromosome. (Abstract) J.
Cell Biol. 75:

[146292]393A, 1977.

[146293]

[146294]

[146295]13739.

[146296]Schwartz, H. E.; Moser, G. C.; Holmes, S.; Meiss, H. K.: Assignment
of temperature-sensitive mutations of BHK cells to the X-

[146297]chromosome. Somat.
Cell Genet. 5: 217-224, 1979.

[146298]

[146299]

[146300]

[146301]13740.

[146302]Sekiguchi, T.; Miyata, T.; Nishimoto, T.: Molecular cloning
of

[146303]the cDNA of human X chromosomal gene (CCG1) which
complements the

[146304]temperature-sensitive G(1) mutants, tsBN462 and ts13, of
the BHK cell

[146305]line. EMBO J. 7: 1683-1687, 1988.

[146306]

[146307]

[146308]13741.

[146309]Sekiguchi, T.; Nohiro, Y.; Nakamura, Y.; Hisamoto, N.;
Nishimoto,

[146310]T.: The human CCG1 gene, essential for progression of
the G1 phase,

[146311]encodes a 210-kilodalton nuclear DNA-binding protein.
Molec. Cell.

[146312]Biol. 11: 3317-3325, 1991.

[146313]

[146314]

[146315]13742.

[146316]Sekiguchi, T.; Yoshida, M. C.; Sekiguchi, M.; Nishimoto, T.:

[146317]Isolation of a human X chromosome-linked gene essential
for progression

[146318]from G1 to S phase of the cell cycle. Exp. Cell Res. 169:
395-407,

[146319]1987.

[146320]

[146321]

[146322]13743.

[146323]Simchen, G.: Cell cycle mutants. Annu. Rev. Genet. 12:
161–191,
[146324]1978.

[146325]

[146326]

[146327]13744.

[146328]Slater, M. L.; Ozer, H. L.: Temperature-sensitive mutants
of

[146329]Balb–3T3 cells: description of a mutant affected in cellular
and polyoma

[146330]virus DNA synthesis. Cell 7: 289–295, 1976.

[146331]

[146332]

[146333]13745.

[146334]Starr, D. B.; Hawley, D. K.: TFIID binds in the minor groove

[146335]of the TATA box. Cell 67: 1231–1240, 1991.

[146336]

[146337]

[146338]13746.

[146339]Abel, A.; Walcott, J.; Woods, J.; Duda, J.; Merry, D. E.: Ex-
pression

[146340]of expanded repeat androgen receptor produces neuro-
logic disease in

[146341]transgenic mice. Hum. Molec. Genet. 10: 107–116, 2001.

[146342]

[146343]

[146344]13747.

[146345]Bailey, C. K.; Andriola, I. F. M.; Kampinga, H. H.; Merry, D.
E.

[146346]: Molecular chaperones enhance the degradation of ex-
panded polyglutamine

[146347]repeat androgen receptor in a cellular model of spinal and
bulbar

[146348]muscular atrophy. Hum. Molec. Genet. 11: 515–523,
2002.

[146349]

[146350]

[146351]13748.

[146352]Batch, J. A.; Williams, D. M.; Davies, H. R.; Brown, B. D.;
Evans,

[146353]B. A. J.; Hughes, I. A.; Patterson, M. N.: Androgen receptor gene

[146354]mutations identified by SSCP in fourteen subjects with androgen insensitivity

[146355]syndrome. Hum. Molec. Genet. 1: 497–503, 1992.

[146356]

[146357]

[146358]13749.

[146359]Beitel, L. K.; Kazemi–Esfarjani, P.; Kaufman, M.; Lumbroso, R.;

[146360]DiGeorge, A. M.; Killinger, D. W.; Trifiro, M. A.; Pinsky, L.: Substitution

[146361]of arginine–839 by cysteine or histidine in the androgen receptor

[146362]causes different receptor phenotypes in cultured cells and coordinate

[146363]degrees of clinical androgen resistance. J. Clin. Invest. 94: 546–554,

[146364]1994.

[146365]

[146366]

[146367]13750.

[146368]Belsham, D. D.; Pereira, F.; Greenberg, C. R.; Liao, S.; Wro-
gemann,

[146369]K.: Leu676-to-pro mutation of the androgen receptor
causes complete

[146370]androgen insensitivity syndrome in a large Hutterite kin-
dred. Hum.

[146371]Mutat. 5: 28-33, 1995.

[146372]

[146373]

[146374]13751.

[146375]Bevan, C. L.; Brown, B. B.; Davies, H. R.; Evans, B. A. J.;
Hughes,

[146376]I. A.; Patterson, M. N.: Functional analysis of six androgen
receptor

[146377]mutations identified in patients with partial androgen in-
sensitivity

[146378]syndrome. Hum. Molec. Genet. 5: 265-273, 1996.

[146379]

[146380]

[146381]13752.

[146382]Biancalana, V.; Serville, F.; Pommier, J.; Julien, J.; Hanauer,

[146383]A.; Mandel, J. L.: Moderate instability of the trinucleotide
repeat

[146384]in spinobulbar muscular atrophy. Hum. Molec. Genet. 1:
255–258,

[146385]1992.

[146386]

[146387]

[146388]13753.

[146389]Boehmer, A. L. M.; Brinkmann, A. O.; Niermeijer, M. F.;
Bakker,

[146390]L.; Halley, D. J. J.; Drop, S. L. S.: Germ–line and somatic
mosaicism

[146391]in the androgen insensitivity syndrome: implications for
genetic counseling.

[146392](Letter) Am. J. Hum. Genet. 60: 1003–1006, 1997.

[146393]

[146394]

[146395]13754.

[146396]Boehmer, A. L. M.; Brinkmann, A. O.; Nijman, R. M.; Ver-
leun-Mooijman,

[146397]M. C. T.; de Ruiter, P.; Niermeijer, M. F.; Drop, S. L. S.:
Phenotypic

[146398]variation in a family with partial androgen insensitivity
syndrome

[146399]explained by differences in 5-alpha dihydrotestosterone
availability. J.

[146400]Clin. Endocr. Metab. 86: 1240-1246, 2001.

[146401]

[146402]

[146403]13755.

[146404]Bouvattier, C.; Carel, J.-C.; Lecoindre, C.; David, A.; Sultan,

[146405]C.; Bertrand, A.-M.; Morel, Y.; Chaussain, J.-L.: Postnatal
changes

[146406]of T, LH, and FSH in 46,XY infants with mutations in the
AR gene. J.

[146407]Clin. Endocr. Metab. 87: 29-32, 2002.

[146408]

[146409]

[146410]13756.

[146411]Brown, C. J.; Goss, S. J.; Lubahn, D. B.; Joseph, D. R.; Wilson,
son,

[146412]E. M.; French, F. S.; Willard, H. F.: Androgen receptor locus
on

[146413]the human X chromosome: regional localization to
Xq11-12 and description

[146414]of a DNA polymorphism. Am. J. Hum. Genet. 44: 264-269,
1989.

[146415]

[146416]

[146417]13757.

[146418]Brown, T. R.; Lubahn, D. B.; Wilson, E. M.; Joseph, D. R.;
French,

[146419]F. S.; Migeon, C. J.: Deletion of the steroid-binding do-
main of the

[146420]human androgen receptor gene in one family with com-
plete androgen

[146421]insensitivity syndrome: evidence for further genetic het-
erogeneity

[146422]in this syndrome. Proc. Nat. Acad. Sci. 85: 8151-8155,
1988.

[146423]

[146424]

[146425]13758.

[146426]Bruggenwirth, H. T.; Boehmer, A. L. M.; Ramnarain, S.;
Verleun-Mooijman,

[146427]M. C. T.; Satijn, D. P. E.; Trapman, J.; Grootegoed, J. A.;
Brinkmann,

[146428]A. O.: Molecular analysis of the androgen-receptor gene
in a family

[146429]with receptor-positive partial androgen insensitivity: an
unusual

[146430]type of intronic mutation. Am. J. Hum. Genet. 61:
1067-1077, 1997.

[146431]

[146432]

[146433]13759.

[146434]Buchanan, G.; Yang, M.; Harris, J. M.; Nahm, H. S.; Han, G.;
Moore,

[146435]N.; Bentel, J. M.; Matusik, R. J.; Horsfall, D. J.; Marshall, V.
R.;

[146436]Greenberg, N. M.; Tilley, W. D.: Mutations at the boundary

of the

[146437]hinge and ligand binding domain of the androgen receptor confer increased

[146438]transactivation function. *Molec. Endocr.* 15: 46–56, 2001.

[146439]

[146440]

[146441]13760.

[146442]Butler, R.; Leigh, P. N.; McPhaul, M. J.; Gallo, J.-M.: Truncated

[146443]forms of the androgen receptor are associated with polyglutamine expansion

[146444]in X-linked spinal and bulbar muscular atrophy. *Hum.*

Molec. Genet. 7:

[146445]121–127, 1998.

[146446]

[146447]

[146448]13761.

[146449]Muller, G.; Schempp, W.: Mapping the human ZFX locus to Xp21.3

[146450]by in situ hybridization. *Hum. Genet.* 82: 82–84, 1989.

[146451]

[146452]

[146453]13762.

[146454]Palmer, M. S.; Sinclair, A. H.; Berta, P.; Ellis, N. A.; Good-fellow,

[146455]P. N.; Abbas, N. E.; Fellous, M.: Genetic evidence that ZFY is not

[146456]the testis-determining factor. Nature 342: 937-939, 1989.

[146457]

[146458]

[146459]13763.

[146460]Masutani, C.; Sugasawa, K.; Yanagisawa, J.; Sonoyama, T.; Ui, M.;

[146461]Enomoto, T.; Takio, K.; Tanaka, K.; van der Spek, P. J.; Bootsma,

[146462]D.; Hoeijmakers, J. H. J.; Hanaoka, F.: Purification and cloning

[146463]of a nucleotide excision repair complex involving the xeroderma pigmentosum

[146464]group C protein and a human homologue of yeast RAD23.

EMBO J. 13:

[146465]1831–1843, 1994.

[146466]

[146467]

[146468]13764.

[146469]Coover, D. D.; Le, T. T.; McAndrew, P. E.; Strasswimmer,
J.;

[146470]Crawford, T. O.; Mendell, J. R.; Coulson, S. E.; Androphy, E.
J.;

[146471]Prior, T. W.; Burghes, A. H. M.: The survival motor neuron
protein

[146472]in spinal muscular atrophy. Hum. Molec. Genet. 6:
1205–1214, 1997.

[146473]

[146474]

[146475]13765.

[146476]Campbell, H. D.; Fountain, S.; McLennan, I. S.; Berven, L.
A.;

[146477]Crouch, M. F.; Davy, D. A.; Hooper, J. A.; Waterford, K.;
Chen, K.–S.;

[146478]Lupski, J. R.; Ledermann, B.; Young, I. G.; Matthaei, K. I.:

Fliih,

[146479]a gelsolin-related cytoskeletal regulator essential for early mammalian

[146480]embryonic development. Molec. Cell. Biol. 22: 3518–3526, 2002.

[146481]

[146482]

[146483]13766.

[146484]Campbell, H. D.; Fountain, S.; Young, I. G.; Claudianos, C.; Hoheisel,

[146485]J. D.; Chen, K.-S.; Lupski, J. R.: Genomic structure, evolution,

[146486]and expression of human FLII, a gelsolin and leucine-rich-repeat family

[146487]member: overlap with LLGL. Genomics 42: 46–54, 1997.

[146488]

[146489]

[146490]13767.

[146491]Connelly, M. A.; Zhang, H.; Kieleczawa, J.; Anderson, C. W.: The

[146492]promoters for human DNA-PK(cs) (PRKDC) and MCM4: di-

vergently transcribed

[146493]genes located at chromosome 8 band q11. Genomics 47:
71–83, 1998.

[146494]

[146495]

[146496]13768.

[146497]Aksoy, I. A.; Wood, T. C.; Weinshilboum, R.: Human liver
estrogen

[146498]sulfotransferase: identification by cDNA cloning and ex-
pression. Biochem.

[146499]Biophys. Res. Commun. 200: 1621–1629, 1994.

[146500]

[146501]

[146502]13769.

[146503]Bernier, F.; Leblanc, G.; Labrie, F.; Luu–The, V.: Structure
of

[146504]human estrogen and aryl sulfotransferase gene: two
mRNA species issued

[146505]from a single gene. J. Biol. Chem. 269: 28200–28205,
1994.

[146506]

[146507]

[146508]13770.

[146509]Her, C.; Aksoy, I. A.; Kimura, S.; Brandriff, B. F.; Wasmuth, J.

[146510]J.; Weinshilboum, R. M.: Human estrogen sulfotransferase gene (STE):

[146511]cloning, structure, and chromosomal localization. Genomics 29: 16–23,

[146512]1995.

[146513]

[146514]

[146515]13771.

[146516]Weinshilboum, R.: Personal Communication. Rochester, Minn.

[146517]10/9/1995.

[146518]

[146519]

[146520]13772.

[146521]Weinshilboum, R. M.; Otterness, D. M.; Aksoy, I. A.; Wood,

T. C.;

[146522]Her, C.; Raftogianis, R. B.: Sulfotransferase molecular biology:

[146523]cDNAs and genes. FASEB J. 11: 3–14, 1997.

[146524]

[146525]

[146526]13773.

[146527]Bartley, T. D.; Bogenberger, J.; Hunt, P.; Li, Y.–S.; Lu, H. S.;

[146528]Martin, F.; Chang, M.–S.; Samal, B.; Nichol, J. L.; Swift, S.;
Johnson,

[146529]M. J.; Hsu, R.–Y.; and 41 others: Identification and cloning
of

[146530]a megakaryocyte growth and development factor that is a
ligand for

[146531]the cytokine receptor Mpl. Cell 77: 1117–1124, 1994.

[146532]

[146533]

[146534]13774.

[146535]Chang, M.; McNinch, J.; Basu, R.; Shutter, J.; Hsu, R.;
Perkins,

[146536]C.; Mar, V.; Suggs, S.; Welcher, A.; Li, L.; Lu, H.; Bartley, T.;

[146537]Hunt, P.; Martin, F.; Samal, B.; Bogenberger, J.: Cloning
and characterization

[146538]of the human megakaryocyte growth and development
factor (MGDF) gene. J.

[146539]Biol. Chem. 270: 511–514, 1995.

[146540]

[146541]

[146542]13775.

[146543]Chang, M.–S.; Hsu, R.–Y.; McNinch, J.; Copeland, N. G.;
Jenkins,

[146544]N. A.: The gene for murine megakaryocyte growth and de-
velopment factor

[146545](thrombopoietin, Thpo) is located on mouse chromosome
16. Genomics 26:

[146546]636–637, 1995.

[146547]

[146548]

[146549]13776.

[146550]de Sauvage, F. J.; Hass, P. E.; Spencer, S. D.; Malloy, B. E.;

[146551]Gurney, A. L.; Spencer, S. A.; Darbonne, W. C.; Henzel, W. J.; Wong,

[146552]S. C.; Kuang, W.-J.; Oles, K. J.; Hultgren, B.; Solberg, L. A., Jr.;

[146553]Goeddel, D. V.; Eaton, D. L.: Stimulation of megakaryocytopoiesis

[146554]and thrombopoiesis by the c-Mpl ligand. Nature 369: 533–538, 1994.

[146555]

[146556]

[146557]13777.

[146558]Farese, A. M.; Hunt, P.; Grab, L. B.; MacVittie, T. J.: Combined

[146559]administration of recombinant human megakaryocyte growth and development

[146560]factor and granulocyte colony-stimulating factor enhances multilineage

[146561]hematopoietic reconstitution in nonhuman primates after radiation-induced

[146562]marrow aplasia. J. Clin. Invest. 97: 2145–2151, 1996.

[146563]

[146564]

[146565]13778.

[146566]Foster, D. C.; Sprecher, C. A.; Grant, F. J.; Kramer, J. M.;
Kuijper,

[146567]J. L.; Holly, R. D.; Whitmore, T. E.; Heipel, M. D.; Bell, L. A.;

[146568]Ching, A. F. T.; McGrane, V.; Hart, C.; O'Hara, P. J.; Lok, S.:
Human

[146569]thrombopoietin: gene structure, cDNA sequence, expres-
sion, and chromosomal

[146570]localization. Proc. Nat. Acad. Sci. 91: 13023–13027, 1994.

[146571]

[146572]

[146573]13779.

[146574]Ghilardi, N.; Wiestner, A.; Kikuchi, M.; Ohsaka, A.; Skoda,
R.

[146575]C.: Hereditary thrombocythaemia in a Japanese family is
caused by

[146576]a novel point mutation in the thrombopoietin gene. Brit. J.
Haemat. 107:

[146577]310–316, 1999.

[146578]

[146579]

[146580]13780.

[146581]Kaushansky, K.; Lok, S.; Holly, R. D.; Broudy, V. C.; Lin, N.;

[146582]Bailey, M. C.; Forstrom, J. W.; Buddle, M. M.; Oort, P. J.;

Hagen,

[146583]F. S.; Roth, G. J.; Papayannopoulou, T.; Foster, D. C.: Pro-
motion

[146584]of megakaryocyte progenitor expansion and differentia-
tion by the c-Mpl

[146585]ligand thrombopoietin. Nature 369: 568–571, 1994.

[146586]

[146587]

[146588]13781.

[146589]Kondo, T.; Okabe, M.; Sanada, M.; Kurosawa, M.; Suzuki,
S.; Kobayashi,

[146590]M.; Hosokawa, M.; Asaka, M.: Familial essential thrombo-
cythemia associated

[146591]with one-base deletion in the 5-prime-untranslated re-
gion of the thrombopoietin

[146592]gene. Blood 92: 1091–1096, 1998.

[146593]

[146594]

[146595]13782.

[146596]Li, J.; Yang, C.; Xia, Y.; Bertino, A.; Glaspy, J.; Roberts, M.;

[146597]Kuter, D. J.: Thrombocytopenia caused by the develop-
ment of antibodies

[146598]to thrombopoietin. Blood 98: 3241–3248, 2001.

[146599]

[146600]

[146601]13783.

[146602]Lok, S.; Kaushansky, K.; Holly, R. D.; Kuijper, J. L.; Lofton-
Day,

[146603]C. E.; Oort, P. J.; Grant, F. J.; Heipel, M. D.; Burkhead, S. K.;

[146604]Kramer, J. M.; Bell, L. A.; Sprecher, C. A.; Blumberg, H.;
Johnson,

[146605]R.; Prunkard, D.; Ching, A. F. T.; Mathewes, S. L.; Bailey, M.
C.;

[146606]Forstrom, J. W.; Buddle, M. M.; Osborn, S. G.; Evans, S. J.;

Sheppard,

[146607]P. O.; Presnell, S. R.; O'Hara, P. J.; Hagen, F. S.; Roth, G. J.;

[146608]Foster, D. C.: Cloning and expression of murine throm-
bopoietin cDNA

[146609]and stimulation of platelet production in vivo. Nature 369:
565-568,

[146610]1994.

[146611]

[146612]

[146613]13784.

[146614]Metcalf, D.: Thrombopoietin: at last. Nature 369:
519-520,

[146615]1994.

[146616]

[146617]

[146618]13785.

[146619]Methia, N.; Louache, F.; Vainchenker, W.; Wendling, F.:
Oligodeoxynucleotides

[146620]antisense to the proto-oncogene c-mpl specifically inhibit
in vitro

[146621]megakaryocytopoiesis. Blood 82: 1395–1401, 1993.

[146622]

[146623]

[146624]13786.

[146625]Ratajczak, M. Z.; Ratajczak, J.; Marlicz, W.; Pletcher, C. H.,

[146626]Jr.; Machalinski, B.; Moore, J.; Hung, H.; Gewirtz, A. M.:

Recombinant

[146627]human thrombopoietin (TPO) stimulates erythropoiesis by
inhibiting

[146628]erythroid progenitor cell apoptosis. Brit. J. Haemat. 98:
8–17,

[146629]1997.

[146630]

[146631]

[146632]13787.

[146633]Sohma, Y.; Akahori, H.; Seki, N.; Hori, T.; Ogami, K.; Kato,
T.;

[146634]Shimada, Y.; Kawamura, K.; Miyazaki, H.: Molecular
cloning and chromosomal

[146635]localization of the human thrombopoietin gene. FEBS Lett.

353: 57–61,

[146636]1994.

[146637]

[146638]

[146639]13788.

[146640]Wendling, F.; Maraskovsky, E.; Debili, N.; Florindo, C.;
Teepe,

[146641]M.; Titeux, M.; Methia, N.; Breton–Gorius, J.; Cosman, D.;
Vainchenker,

[146642]W.: c–Mpl ligand is a humoral regulator of megakaryocy-
topoiesis. Nature 369:

[146643]571–574, 1994.

[146644]

[146645]

[146646]13789.

[146647]Dualan, R.; Brody, T.; Keeney, S.; Nichols, A. F.; Admon,
A.; Linn,

[146648]S.: Chromosomal localization and cDNA cloning of the
genes (DDB1

[146649]and DDB2) for the p127 and p48 subunits of a human

damage-specific

[146650]DNA binding protein. Genomics 29: 62–69, 1995.

[146651]

[146652]

[146653]13790.

[146654]Koken, M. H. M.; Reynolds, P.; Jaspers–Dekker, I.; Prakash,
L.;

[146655]Prakash, S.; Bootsma, D.; Hoeijmakers, J. H. J.: Structural
and functional

[146656]conservation of two human homologs of the yeast DNA
repair gene RAD6. Proc.

[146657]Nat. Acad. Sci. 88: 8865–8869, 1991.

[146658]

[146659]

[146660]13791.

[146661]Knoll, A.; Schunkert, H.; Reichwald, K.; Danser, A. H. J.;
Bauer,

[146662]D.; Platzter, M.; Stein, G.; Rosenthal, A.: Human renin bind-
ing protein:

[146663]complete genomic sequence and association of an intronic

T/C polymorphism

[146664]with the prorenin level in males. Hum. Molec. Genet. 6:
1527–1534,

[146665]1997.

[146666]

[146667]

[146668]13792.

[146669]Laan, M.; Paabo, S.: Mapping genes by drift-generated
linkage

[146670]disequilibrium. (Letter) Am. J. Hum. Genet. 63: 654–656,
1998.

[146671]

[146672]

[146673]13793.

[146674]Takahashi, S.; Inoue, H.; Miyake, Y.: The human gene for
renin-binding

[146675]protein. J. Biol. Chem. 267: 13007–13013, 1992.

[146676]

[146677]

[146678]13794.

[146679]Terwilliger, J. D.; Zollner, S.; Laan, M.; Paabo, S.: Mapping

[146680]genes through the use of linkage disequilibrium generated by genetic

[146681]drift: 'drift mapping' in small populations with no demographic expansion. Hum.

[146682]Hered. 48: 138–154, 1998.

[146683]

[146684]

[146685]13795.

[146686]van den Ouweland, A. M. W.; Verdijk, M.; Kiochis, P.; Poustka,

[146687]A.; van Oost, B. A.: The human renin-binding protein gene (RENBP)

[146688]maps in Xq28. Genomics 21: 279–281, 1994.

[146689]

[146690]

[146691]13796.

[146692]Warren, S. T.; Knight, S. J. L.; Peters, J. F.; Stayton, C. L.;

[146693]Consalez, G. G.; Zhang, F.: Isolation of the human chromosomal band

[146694]Xq28 within somatic cell hybrids by fragile X site breakage. Proc.

[146695]Nat. Acad. Sci. 87: 3856–3860, 1990.

[146696]

[146697]

[146698]13797.

[146699]Calvo, R. M.; Asuncion, M.; Sancho, J.; San Millan, J. L.; Escobar-Morreale,

[146700]H. F.: The role of the CAG repeat polymorphism in the androgen receptor

[146701]gene and of skewed X-chromosome inactivation, in the pathogenesis

[146702]of hirsutism. J. Clin. Endocr. Metab. 85: 1735–1740, 2000.

[146703]

[146704]

[146705]13798.

[146706]Gao, X.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Gridley,

[146707]T.: Assignment of the murine Notch2 and Notch3 genes to
chromosomes

[146708]3 and 17. Genomics 49: 160–161, 1998.

[146709]

[146710]

[146711]13799.

[146712]Larsson, C.; Lardelli, M.; White, I.; Lendahl, U.: The human
NOTCH1,

[146713]2, and 3 genes are located at chromosome positions
9q34, 1p13–p11,

[146714]and 19p13.2–p13.1 in regions of neoplasia–associated
translocation. Genomics 24:

[146715]253–258, 1994.

[146716]

[146717]

[146718]13800.

[146719]Dichgans, M.; Herzog, J.; Gasser, T.: NOTCH3 mutation in–
volving

[146720]three cysteine residues in a family with typical CADASIL.
Neurology 57:

[146721]1714–1717, 2001.

[146722]

[146723]

[146724]13801.

[146725]Joutel, A.; Corpechot, C.; Ducros, A.; Vahedi, K.; Chabriat, H.;

[146726]Mouton, P.; Alamowitch, S.; Domenga, V.; Cecillion, M.; Marechal,

[146727]E.; Maciazek, J.; Vayssiere, C.; Cruaud, C.; Cabanis, E.–A.; Ruchoux,

[146728]M. M.; Weissenbach, J.; Bach, J. F.; Bousser, M. G.; Tournier–Lasserre,

[146729]E.: Notch3 mutations in CADASIL, a hereditary adult–onset condition

[146730]causing stroke and dementia. Nature 383: 707–710, 1996.

[146731]

[146732]

[146733]13802.

[146734]Joutel, A.; Dodick, D. D.; Parisi, J. E.; Cecillion, M.; Tournier–Lasserre,

[146735]E.; Bousser, M. G.: De novo mutation in the Notch3 gene causing CADASIL. Ann.

[146736]Neurol. 47: 388–391, 2000.

[146737]

[146738]

[146739]13803.

[146740]Joutel, A.; Vahedi, K.; Corpechot, C.; Troesch, A.; Chabriat, H.;

[146741]Vayssiere, C.; Cruaud, C.; Maciazek, J.; Weissenbach, J.; Bousser,

[146742]M.–G.; Bach, J.–F.; Tournier–Lasserre, E.: Strong clustering and

[146743]stereotyped nature of Notch3 mutations in CADASIL patients. Lancet 350:

[146744]1511–1515, 1997.

[146745]

[146746]

[146747]13804.

[146748]Rebay, I.; Fehon, R. G.; Artavanis–Tsakonas, S.: Specific truncations

[146749]of *Drosophila* Notch define dominant activated and dominant negative

[146750]forms of the receptor. *Cell* 74: 319–329, 1993.

[146751]

[146752]

[146753]13805.

[146754]Deed, R. W.; Hirose, T.; Mitchell, E. L. D.; Santibanez-Koref,

[146755]M. F.; Norton, J. D.: Structural organisation and chromosomal mapping

[146756]of the human Id-3 gene. Gene 151: 309-314, 1994.

[146757]

[146758]

[146759]13806.

[146760]Ellmeier, W.; Aguzzi, A.; Kleiner, E.; Kurzbauer, R.; Weith, A.

[146761]: Mutually exclusive expression of a helix-loop-helix gene and N-myc

[146762]in human neuroblastomas and in normal development. EMBO J. 11: 2563-2571,

[146763]1992.

[146764]

[146765]

[146766]13807.

[146767]Kee, B. L.; Rivera, R. R.; Murre, C.: Id3 inhibits B lymphocyte

[146768]progenitor growth and survival in response to TGF-beta. Nature Immun. 2:

[146769]242–247, 2001.

[146770]

[146771]

[146772]13808.

[146773]Lyden, D.; Young, A. Z.; Zagzag, D.; Yan, W.; Gerald, W.;
O'Reilly,

[146774]R.; Bader, B. L.; Hynes, R. O.; Zhuang, Y.; Manova, K.;
Benezra, R.

[146775]: Id1 and Id3 are required for neurogenesis, angiogenesis
and vascularization

[146776]of tumour xenografts. Nature 401: 670–677, 1999.

[146777]

[146778]

[146779]13809.

[146780]Pan, L.; Sato, S.; Frederick, J. P.; Sun, X.–H.; Zhuang, Y.:
Impaired

[146781]immune responses and B–cell proliferation in mice lacking
the Id3

[146782]gene. Molec. Cell. Biol. 19: 5969–5980, 1999.

[146783]

[146784]

[146785]13810.

[146786]Yeh, K.; Lim, R. W.: Genomic organization and promoter analysis

[146787]of the murine Id3 gene. Gene 254: 163–171, 2000.

[146788]

[146789]

[146790]13811.

[146791]Braun, A.; Kammerer, S.; Weissenhorn, W.; Weiss, E. H.; Cleve,

[146792]H.: Sequence of a putative human housekeeping gene (HK33) localized

[146793]on chromosome 1. Gene 146: 291–295, 1994.

[146794]

[146795]

[146796]13812.

[146797]Gotte, K.; Girzalsky, W.; Linkert, M.; Baumgart, E.; Kammerer,

[146798]S.; Kunau, W.–H.; Erdmann, R.: Pex19p, a farnesylated

protein essential

[146799]for peroxisome biogenesis. Molec. Cell. Biol. 18: 616–628, 1998.

[146800]

[146801]

[146802]13813.

[146803]James, G. L.; Goldstein, J. L.; Pathak, R. K.; Anderson, R. G.

[146804]W.; Brown, M. S.: PxF, a prenylated protein of peroxisomes
J. Biol.

[146805]Chem. 269: 14182–14190, 1994.

[146806]

[146807]

[146808]13814.

[146809]Kammerer, S.; Arnold, N.; Gutensohn, W.; Mewes, H.–W.;
Kunau, W.–H.;

[146810]Hofler, G.; Roscher, A. A.; Braun, A.: Genomic organization
and molecular

[146811]characterization of a gene encoding HsPXF, a human per-
oxisomal farnesylated

[146812]protein. Genomics 45: 200–210, 1997.

[146813]

[146814]

[146815]13815.

[146816]Caskey, C. T.; Pizzuti, A.; Fu, Y.-H.; Fenwick, R. G., Jr.;
Nelson,

[146817]D. L.: Triplet repeat mutations in human disease. Science
256:

[146818]784–789, 1992.

[146819]

[146820]

[146821]13816.

[146822]Chang, B.; Zheng, S. L.; Hawkins, G. A.; Isaacs, S. D.; Wiley,

[146823]K. E.; Turner, A.; Carpten, J. D.; Bleecker, E. R.; Walsh, P.
C.;

[146824]Trent, J. M.; Meyers, D. A.; Isaacs, W. B.; Xu, J.: Polymor-
phic GGC

[146825]repeats in the androgen receptor gene are associated with
hereditary

[146826]and sporadic prostate cancer risk. Hum. Genet. 110:
122–129, 2002.

[146827]

[146828]

[146829]13817.

[146830]Chang, C.; Kokontis, J.; Liao, S.: Molecular cloning of human
man

[146831]and rat complementary DNA encoding androgen receptors. Science 240:

[146832]324–326, 1988.

[146833]

[146834]

[146835]13818.

[146836]Choong, C. S.; Kemppainen, J. A.; Zhou, Z.–X.; Wilson, E.
M.:

[146837]Reduced androgen receptor gene expression with first
exon CAG repeat

[146838]expansion. Molec. Endocr. 10: 1527–1535, 1996.

[146839]

[146840]

[146841]13819.

[146842]Choong, C. S.; Quigley, C. A.; French, F. S.; Wilson, E. M.:

[146843]A novel missense mutation in the amino-terminal domain
of the human

[146844]androgen receptor gene in a family with partial androgen
insensitivity

[146845]syndrome causes reduced efficiency of protein translation.
J. Clin.

[146846]Invest. 98: 1423–1431, 1996.

[146847]

[146848]

[146849]13820.

[146850]Chu, J.; Zhang, R.; Zhao, Z.; Zou, W.; Han, Y.; Qi, Q.;
Zhang,

[146851]H.; Wang, J.–C.; Tao, S.; Liu, X.; Luo, Z.: Male fertility is
compatible

[146852]with an Arg840Cys substitution in the AR in a large Chi-
nese family

[146853]with divergent phenotypes of AR insensitivity syndrome. J.
Clin.

[146854]Endocr. Metab. 87: 347–351, 2002.

[146855]

[146856]

[146857]13821.

[146858]Coetzee, G. A.; Ross, R. K.: Re: Prostate cancer and the
androgen

[146859]receptor. (Letter) J. Nat. Cancer Inst. 86: 872–873, 1994.

[146860]

[146861]

[146862]13822.

[146863]Correa–Cerro, L.; Wöhr, G.; Haussler, J.; Berthon, P.;
Drelon,

[146864]E.; Mangin, P.; Fournier, G.; Cussenot, O.; Kraus, P.; Just,
W.; Paiss,

[146865]T.; Cantu, J. M.; Vogel, W.: (CAG)_nCAA and GGN repeats in
the human

[146866]androgen receptor gene are not associated with prostate
cancer in

[146867]a French–German population. Europ. J. Hum. Genet. 7:
347–362, 1999.

[146868]

[146869]

[146870]13823.

[146871]Dowsing, A. T.; Yong, E. L.; Clark, M.; McLachlan, R. I.; de Kretser,

[146872]D. M.; Trounson, A. O.: Linkage between male infertility and trinucleotide

[146873]repeat expansion in the androgen-receptor gene. Lancet 354: 640-643,

[146874]1999.

[146875]

[146876]

[146877]13824.

[146878]Elo, J. P.; Kvist, L.; Leinonen, K.; Isomaa, V.; Hentuu, P.; Lukkarinen,

[146879]O.; Vihko, P.: Mutated human androgen receptor gene detected in a

[146880]prostatic cancer patient is also activated by estradiol. J. Clin.

[146881]Endocr. Metab. 80: 3494-3500, 1995.

[146882]

[146883]

[146884]13825.

[146885]Gaddipati, J. P.; McLeod, D. G.; Heidenberg, H. B.; Sesterhenn,

[146886]I. A.; Finger, M. J.; Moul, J. W.; Srivastava, S.: Frequent detection

[146887]of codon 877 mutation in the androgen receptor gene in advanced prostate

[146888]cancers. Cancer Res. 54: 2861–2864, 1994.

[146889]

[146890]

[146891]13826.

[146892]Gehring, U.; Tomkins, G. M.: Characterization of a hormone receptor

[146893]defect in the androgen–insensitivity mutant. Cell 3: 59–64, 1974.

[146894]

[146895]

[146896]13827.

[146897]Gingrich, J. R.; Barrios, R. J.; Kattan, M. W.; Nahm, H. S.; Finegold,

[146898]M. J.; Greenberg, N. M.: Androgen-independent prostate cancer progression

[146899]in the TRAMP model. Cancer Res. 57: 4687–4691, 1997.

[146900]

[146901]

[146902]13828.

[146903]Gingrich, J. R.; Greenberg, N. M.: A transgenic mouse prostate

[146904]cancer model. Toxicol. Path. 24: 502–504, 1996.

[146905]

[146906]

[146907]13829.

[146908]Giovannucci, E.; Stampfer, M. J.; Krithivas, K.; Brown, M.; Brufsky,

[146909]A.; Talcott, J.; Hennekens, C. H.; Kantoff, P. W.: The CAG repeat

[146910]within the androgen receptor gene and its relationship to prostate

[146911]cancer. Proc. Nat. Acad. Sci. 94: 3320–3323, 1997.

[146912]

[146913]

[146914]13830.

[146915]Gottlieb, B.; Beitel, L. K.; Trifiro, M. A.: Variable expressivity

[146916]and mutation databases: the androgen receptor gene mutations database. Hum.

[146917]Mutat. 17: 382–388, 2001.

[146918]

[146919]

[146920]13831.

[146921]Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Pinsky, L.: The androgen

[146922]receptor gene mutations database. Nucleic Acids Res. 25: 158–162,

[146923]1997.

[146924]

[146925]

[146926]13832.

[146927]Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Vasiliou, D. M.; Pin-

sky,

[146928]L.: The androgen receptor gene mutations database. Nu-
cleic Acids

[146929]Res. 24: 151–154, 1996.

[146930]

[146931]

[146932]13833.

[146933]Griffin, J. E.: Androgen resistance--the clinical and molec-
ular

[146934]spectrum. New Eng. J. Med. 326: 611–618, 1992.

[146935]

[146936]

[146937]13834.

[146938]Grino, P. B.; Isidro-Gutierrez, R. F.; Griffin, J. E.; Wilson,

[146939]J. D.: Androgen resistance associated with a qualitative
abnormality

[146940]of the androgen receptor and responsive to high dose an-
drogen therapy. J.

[146941]Clin. Endocr. Metab. 68: 578–584, 1989.

[146942]

[146943]

[146944]13835.

[146945]Hardy, D. O.; Scher, H. I.; Bogenreider, T.; Sabbatini, P.;
Zhang,

[146946]Z.-F.; Nanus, D. M.; Catterall, J. F.: Androgen receptor
CAG repeat

[146947]lengths in prostate cancer: correlation with age of onset. J.
Clin.

[146948]Endocr. Metab. 81: 4400–4405, 1996.

[146949]

[146950]

[146951]13836.

[146952]Hellwinkel, O. J.-C.; Holterhus, P.-M.; Struve, D.;
Marschke,

[146953]C.; Homburg, N.; Hiort, O.: A unique exonic splicing mu-
tation in

[146954]the human androgen receptor gene indicates a physio-
logic relevance

[146955]of regular androgen receptor transcript variants. J. Clin.
Endocr.

[146956]Metab. 86: 2569–2575, 2001.

[146957]

[146958]

[146959]13837.

[146960]Hickey, T.; Chandy, A.; Norman, R. J.: The androgen receptor

[146961]CAG repeat polymorphism and X-chromosome inactivation in Australian

[146962]Caucasian women with infertility related to polycystic ovary syndrome. J.

[146963]Clin. Endocr. Metab. 87: 161–165, 2002.

[146964]

[146965]

[146966]13838.

[146967]Hiort, O.; Sinnecker, G. H. G.; Holterhus, P.–M.; Nitsche, E.

[146968]M.; Kruse, K.: Inherited and de novo androgen receptor gene mutations:

[146969]investigation of single-case families. J. Pediat. 132: 939–943,

[146970]1998.

[146971]

[146972]

[146973]13839.

[146974]Holterhus, P.-M.; Bruggenwirth, H. T.; Hiort, O.;

Kleinkauf-Houcken,

[146975]A.; Kruse, K.; Sinnecker, G. H. G.; Brinkmann, A. O.: Mo-
saicism due

[146976]to a somatic mutation of the androgen receptor gene de-
termines phenotype

[146977]in androgen insensitivity syndrome. J. Clin. Endocr. Metab.
82:

[146978]3584-3589, 1997.

[146979]

[146980]

[146981]13840.

[146982]Shibuya, H.; Yamaguchi, K.; Shirakabe, K.; Tonegawa, A.;

Gotoh,

[146983]Y.; Ueno, N.; Irie, K.; Nishida, E.; Matsumoto, K.: TAB1: an
activator

[146984]of the TAK1 MAPKKK in TGF-beta signal transduction. Sci-
ence 272:

[146985]1179–1182, 1996.

[146986]

[146987]

[146988]13841.

[146989]Boyd, J. M.; Subramanian, T.; Schaeper, U.; La Regina, M.;
Bayley,

[146990]S.; Chinnadurai, G.: A region in the C-terminus of aden-
ovirus 2/5

[146991]E1a protein is required for association with a cellular
phosphoprotein

[146992]and important for the negative modulation of T24-ras
mediated transformation,

[146993]tumorigenesis and metastasis. EMBO J. 469–478, 1993.

[146994]

[146995]

[146996]13842.

[146997]Furusawa, T.; Moribe, H.; Kondoh, H.; Higashi, Y.: Identifi-
cation

[146998]of CtBP1 and CtBP2 as corepressors of zinc finger-
homeodomain factor

[146999]delta-EF1. Molec. Cell. Biol. 19: 8581–8590, 1999.

[147000]

[147001]

[147002]13843.

[147003]Katsanis, N.; Fisher, E. M. C.: A novel C-terminal binding protein

[147004](CTBP2) is closely related to CTBP1, an adenovirus E1A-binding protein,

[147005]and maps to human chromosome 21q21.3. Genomics 47: 294–299, 1998.

[147006]

[147007]

[147008]13844.

[147009]Schaeper, U.; Boyd, J. M.; Verma, S.; Uhlmann, E.; Subramanian,

[147010]T.; Chinnadurai, G.: Molecular cloning and characterization of a

[147011]cellular phosphoprotein that interacts with a conserved C-terminal

[147012]domain of adenovirus E1A involved in negative modulation of oncogenic

[147013]transformation. Proc. Nat. Acad. Sci. 92: 10467–10471,
1995.

[147014]

[147015]

[147016]13845.

[147017]Sewalt, R. G. A. B.; Gunster, M. J.; van der Vlag, J.; Satijn,

[147018]D. P. E.; Otte, A. P.: C-terminal binding protein is a tran-
scriptional

[147019]repressor that interacts with a specific class of vertebrate
polycomb

[147020]proteins. Molec. Cell. Biol. 19: 777–787, 1999.

[147021]

[147022]

[147023]13846.

[147024]Zhang, Q.; Piston, D. W.; Goodman, R. H.: Regulation of
corepressor

[147025]function by nuclear NADH. Science 295: 1895–1897,
2002.

[147026]

[147027]

[147028]13847.

[147029]Hess, S. D.; Daggett, L. P.; Deal, C.; Lu, C.-C.; Johnson, E. C.;

[147030]Velicelebi, G.: Functional characterization of human N-methyl-D-aspartate

[147031]subtype 1A/2D receptors. J. Neurochem. 70: 1269–1279, 1998.

[147032]

[147033]

[147034]13848.

[147035]Ikeda, K.; Araki, K.; Takayama, C.; Inoue, Y.; Yagi, T.; Aizawa,

[147036]S.; Mishina, M.: Reduced spontaneous activity of mice defective in

[147037]the epsilon 4 subunit of the NMDA receptor channel. Brain Res. Molec.

[147038]Brain Res. 33: 61–71, 1995.

[147039]

[147040]

[147041]13849.

[147042]Watanabe, T.; Inoue, S.; Hiroi, H.; Orimo, A.; Kawashima, H.; Muramatsu,

[147043]M.: Isolation of estrogen-responsive genes with a CpG island library. Molec.

[147044]Cell. Biol. 18: 442–449, 1998.

[147045]

[147046]

[147047]13850.

[147048]Adra, C. N.; Kobayashi, H.; Rowley, J. D.; Lim, B.: Assignment

[147049]of the human GDID4 gene, a GDP/GTP-exchange regulator, to chromosome

[147050]12p12.3. Genomics 24: 188–190, 1994.

[147051]

[147052]

[147053]13851.

[147054]Leffers, H.; Nielsen, M. S.; Andersen, A. H.; Honore, B.; Madsen,

[147055]P.; Vandekerckhove, J.; Celis, J. E.: Identification of two human

[147056]Rho GDP dissociation inhibitor proteins whose overexpression leads

[147057]to disruption of the actin cytoskeleton. Exp. Cell Res. 209: 165–174,

[147058]1993.

[147059]

[147060]

[147061]13852.

[147062]Lelias, J.–M.; Adra, C. N.; Wulf, G. M.; Guillemot, J.–C.; Khagad,

[147063]M.; Caput, D.; Lim, B.: cDNA cloning of a human mRNA preferentially

[147064]expressed in hematopoietic cells and with homology to a GDP–dissociation

[147065]inhibitor for the rho GTP–binding proteins. Proc. Nat. Acad. Sci. 90:

[147066]1479–1483, 1993.

[147067]

[147068]

[147069]13853.

[147070]Scherle, P.; Behrens, T.; Staudt, L. M.: Ly-GDI, a GDP-dissociation

[147071]inhibitor of the RhoA GTP-binding protein, is expressed preferentially

[147072]in lymphocytes. Proc. Nat. Acad. Sci. 90: 7568–7572, 1993.

[147073]

[147074]

[147075]13854.

[147076]Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.; Tanaka, A.;

[147077]Kotani, H.; Nomura, N.; Ohara, O.: Prediction of coding sequences

[147078]of unidentified human genes. X. The complete sequences of 100 new

[147079]cDNA clones from brain which can code for large proteins in vitro. DNA

[147080]Res. 5: 169–176, 1998.

[147081]

[147082]

[147083]13855.

[147084]McMillan, D. R.; Kayes–Wandover, K. M.; Richardson, J. A.;
White,

[147085]P. C.: Very large G protein–coupled receptor–1, the largest
known

[147086]cell surface protein, is highly expressed in the developing
central

[147087]nervous system. J. Biol. Chem. 277: 785–792, 2002.

[147088]

[147089]

[147090]13856.

[147091]Nikkila, H.; McMillan, D. R.; Nunez, B. S.; Pascoe, L.;
Curnow,

[147092]K. M.; White, P. C.: Sequence similarities between a novel
putative

[147093]G protein–coupled receptor and Na(+)/Ca(2+) exchangers
define a cation

[147094]binding domain. Molec. Endocr. 14: 1351–1364, 2000.

[147095]

[147096]

[147097]13857.

[147098]Skradski, S. L.; Clark, A. M.; Jiang, H.; White, H. S.; Fu, Y.-H.;

[147099]Ptacek, L. J.: A novel gene causing a mendelian audiogenic mouse

[147100]epilepsy. Neuron 31: 537–544, 2001.

[147101]

[147102]

[147103]13858.

[147104]Skradski, S. L.; White, H. S.; Ptacek, L. J.: Genetic mapping

[147105]of a locus (mass1) causing audiogenic seizures in mice.

Genomics 49:

[147106]188–192, 1998.

[147107]

[147108]

[147109]13859.

[147110]McLennan, A. G.; Flannery, A. V.; Morten, J. E. N.; Ridanpaa, M.

[147111]: Chromosomal localization of the human diadenosine 5–prime,5–triple

[147112]prime–P(1),P(4)–tetrphosphate pyrophosphohydrolase

(Ap4A hydrolase)

[147113]gene (APAH1) to 9p13. Genomics 47: 307–309, 1998.

[147114]

[147115]

[147116]13860.

[147117]Thorne, M. H.; Hankin, S.; Wilkinson, M. C.; Nunez, C.;
Barraclough,

[147118]R.; McLennan, A. G.: Human diadenosine 5–prime,5–triple
prime–P(1),P(4)–tetrphosphate

[147119]pyrophosphohydrolase is a member of the MutT family of
nucleotide

[147120]pyrophosphatases. Biochem. J. 311: 717–721, 1995.

[147121]

[147122]

[147123]13861.

[147124]Waldegger, S.; Erdel, M.; Nagl, U. O.; Barth, P.; Raber, G.;
Steuer,

[147125]S.; Utermann, G.; Paulmichl, M.; Lang, F.: Genomic organi-
zation and

[147126]chromosomal localization of the human SGK protein ki-

nase gene. Genomics 51:

[147127]299–302, 1998.

[147128]

[147129]

[147130]13862.

[147131]Webster, M. K.; Goya, L.; Ge, Y.; Maiyar, A. C.; Firestone,
G.

[147132]L.: Characterization of sgk, a novel member of the serine/
threonine

[147133]protein kinase gene family which is transcriptionally in-
duced by glucocorticoids

[147134]and serum. Molec. Cell. Biol. 13: 2031–2040, 1993.

[147135]

[147136]

[147137]13863.

[147138]Anand, N.; Murthy, S.; Amann, G.; Wernick, M.; Porter, L.
A.; Cukier,

[147139]J. H.; Collins, C.; Gray, J. W.; Diebold, J.; Demetrick, D. J.;
Lee,

[147140]J. M.: Protein elongation factor EEF1A2 is a putative onco-

gene in

[147141]ovarian cancer. Nature Genet. 31: 301–305, 2002.

[147142]

[147143]

[147144]13864.

[147145]Ann, D. K.; Moutsatsos, I. K.; Nakamura, T.; Lin, H. H.;
Mao, P.-L.;

[147146]Lee, M.-J.; Chin, S.; Liem, R. K. H.; Wang, E.: Isolation and
characterization

[147147]of the rat chromosomal gene for a polypeptide (p21) anti-
genically

[147148]related to statin. J. Biol. Chem. 266: 10429–10437, 1991.

[147149]

[147150]

[147151]13865.

[147152]Bischoff, C.; Kahns, S.; Lund, A.; Jorgensen, H. F.; Praeste-
gaard,

[147153]M.; Clark, B. F. C.; Leffers, H.: The human elongation fac-
tor 1 A-2

[147154]gene (EEF1A2): complete sequence and characterization of

gene structure

[147155]and promoter activity. Genomics 68: 63–70, 2000.

[147156]

[147157]

[147158]13866.

[147159]Chambers, D. M.; Peters, J.; Abbott, C. M.: The lethal mutation

[147160]of the mouse wasted (wst) is a deletion that abolishes expression

[147161]of a tissue-specific isoform of translation elongation factor 1-alpha,

[147162]encoded by the Eef1a2 gene. Proc. Nat. Acad. Sci. 95: 4463–4468,

[147163]1998.

[147164]

[147165]

[147166]13867.

[147167]Shultz, L. D.; Sweet, H. O.; Davisson, M. T.; Coman, D. R.: 'Wasted,'

[147168]a new mutant of the mouse with abnormalities character-

istic of ataxia

[147169]telangiectasia. Nature 297: 402–404, 1982.

[147170]

[147171]

[147172]13868.

[147173]Aoki, K.; Ishida, R.; Kasai, M.: Isolation and characteriza-
tion

[147174]of a cDNA encoding a translin-like protein, TRAX. FEBS
Lett. 401:

[147175]109–112, 1997.

[147176]

[147177]

[147178]13869.

[147179]Meng, G.; Aoki, K.; Tokura, K.; Nakahara, K.; Inazawa, J.;
Kasai,

[147180]M.: Genomic structure and chromosomal localization of
the gene encoding

[147181]TRAX, a translin-associated factor X. J. Hum. Genet. 45:
305–308,

[147182]2000.

[147183]

[147184]

[147185]13870.

[147186]Eudy, J. D.; Yao, S.; Weston, M. D.; Ma-Edmonds, M.; Tal-
madge,

[147187]C. B.; Cheng, J. J.; Kimberling, W. J.; Sumegi, J.: Isolation of

[147188]a gene encoding a novel member of the nuclear receptor
superfamily

[147189]from the critical region of Usher syndrome type IIa at
1q41. Genomics 50:

[147190]382-384, 1998.

[147191]

[147192]

[147193]13871.

[147194]Greschik, H.; Wurtz, J.-M.; Sanglier, S.; Bourguet, W.; van
Dorsselaer,

[147195]A.; Moras, D.; Renaud, J.-P.: Structural and functional evi-
dence

[147196]for ligand-independent transcriptional activation by the
estrogen-related

[147197]receptor 3. Molec. Cell 9: 303–313, 2002.

[147198]

[147199]

[147200]13872.

[147201]Quintana, D. G.; Hou, Z.; Thome, K. C.; Hendricks, M.;
Saha, P.;

[147202]Dutta, A.: Identification of HsORC4, a member of the hu-
man origin

[147203]of replication recognition complex. J. Biol. Chem. 272:
28247–28251,

[147204]1997.

[147205]

[147206]

[147207]13873.

[147208]Eki, T.; Dean, F. B.; Kohda, A.; Okumura, K.; Abe, M.; Mu-
rakami,

[147209]Y.; Ishiai, M.; Satomoto, K.; Hurwitz, J.; O'Donnell, M.;
Hanaoka,

[147210]F.: Assignment1 of the homologue of the yeast origin
recognition

[147211]complex subunit ORC4 (ORC4L) to human chromosome
band 2q22–q23 by

[147212]in situ hybridization and somatic cell hybrid analysis. Cy-
togenet.

[147213]Cell Genet. 81: 89–90, 1998.

[147214]

[147215]

[147216]13874.

[147217]Matsuyoshi, N.; Imamura, S.: Multiple cadherins are ex-
pressed

[147218]in human fibroblasts. Biochem. Biophys. Res. Commun.
235: 355–358,

[147219]1997.

[147220]

[147221]

[147222]13875.

[147223]Heron, L.; Virsolvy, A.; Peyrolier, K.; Gribble, F. M.; Le
Cam,

[147224]A.; Ashcroft, F. M.; Bataille, D.: Human alpha–endosulfine,
a possible

[147225]regulator of sulfonylurea-sensitive K(ATP) channel: molecular cloning,

[147226]expression and biological properties. Proc. Nat. Acad. Sci. 95:

[147227]8387–8391, 1998.

[147228]

[147229]

[147230]13876.

[147231]Scott, A. F.: Personal Communication. Baltimore, Md. 1/11/2001.

[147232]

[147233]

[147234]13877.

[147235]Bertilsson, G.; Heidrich, J.; Svensson, K.; Asman, M.; Jen-deberg,

[147236]L.; Sydow-Backman, M.; Ohlsson, R.; Postlind, H.;

Blomquist, P.; Berkenstam,

[147237]A.: Identification of a human nuclear receptor defines a new signaling

[147238]pathway for CYP3A induction. Proc. Nat. Acad. Sci. 95: 12208–12213,

[147239]1998.

[147240]

[147241]

[147242]13878.

[147243]Blumberg, B.; Sabbagh, W., Jr.; Juguilon, H.; Bolado, J., Jr.;

[147244]van Meter, C. M.; Ong, E. S.; Evans, R. M.: SXR, a novel
steroid

[147245]and xenobiotic-sensing nuclear receptor. Genes Dev. 12:
3195–3205,

[147246]1998.

[147247]

[147248]

[147249]13879.

[147250]Watkins, R. E.; Wisely, G. B.; Moore, L. B.; Collins, J. L.;
Lambert,

[147251]M. H.; Williams, S. P.; Willson, T. M.; Kliewer, S. A.; Red-
inbo, M.

[147252]R.: The human nuclear xenobiotic receptor PXR: structural
determinants

[147253]of directed promiscuity. Science 292: 2329–2333, 2001.

[147254]

[147255]

[147256]13880.

[147257]Grenet, J.; Valentine, V.; Kitson, J.; Li, H.; Farrow, S. N.;
Kidd,

[147258]V. J.: Duplication of the DR3 gene on human chromosome
1p36 and its

[147259]deletion in human neuroblastoma. Genomics 49:
385–393, 1998.

[147260]

[147261]

[147262]13881.

[147263]Adato, A.; Raskin, L.; Petit, C.; Bonne–Tamir, B.: Deafness
heterogeneity

[147264]in a Druze isolate from the Middle East: novel OTOF and
PDS mutations,

[147265]low prevalence of GJB2 35delG mutation and indication for
a new DFNB

[147266]locus. Europ. J. Hum. Genet. 8: 437–442, 2000.

[147267]

[147268]

[147269]13882.

[147270]Yasunaga, S.; Grati, M.; Chardenoux, S.; Smith, T. N.;
Friedman,

[147271]T. B.; Lalwani, A. K.; Wilcox, E. R.; Petit, C.: OTOF encodes
multiple

[147272]long and short isoforms: genetic evidence that the long
ones underlie

[147273]recessive deafness DFNB9. Am. J. Hum. Genet. 67:
591–600, 2000.

[147274]

[147275]

[147276]13883.

[147277]Yasunaga, S.; Petit, C.: Physical map of the region sur-
rounding

[147278]the OTOFERLIN locus on chromosome 2p22–p23. Ge-
nomics 66: 110–112,

[147279]2000.

[147280]

[147281]

[147282]13884.

[147283]Niederreither, K.; Subbarayan, V.; Dolle, P.; Chambon, P.:

Embryonic

[147284]retinoic acid synthesis is essential for early mouse post-
implantation

[147285]development. Nature Genet. 21: 444–448, 1999.

[147286]

[147287]

[147288]13885.

[147289]Ono, Y.; Fukuhara, N.; Yoshie, O.: TAL1 and LIM-only pro-
teins

[147290]synergistically induce retinaldehyde dehydrogenase 2 ex-
pression in

[147291]T-cell acute lymphoblastic leukemia by acting as cofactors
for GATA3. Molec.

[147292]Cell. Biol. 18: 6939–6950, 1998.

[147293]

[147294]

[147295]13886.

[147296]Perlmann, T.: Retinoid metabolism: a balancing act. Na-

ture Genet. 31:

[147297]7–9, 2002. Note: Erratum: Nature Genet. 31: 221 only,
2002.

[147298]

[147299]

[147300]13887.

[147301]Wang, X.; Penzes, P.; Napoli, J. L.: Cloning of a cDNA en-
coding

[147302]an aldehyde dehydrogenase and its expression in Es-
cherichia coli:

[147303]recognition of retinal as substrate. J. Biol. Chem. 271:
16288–16293,

[147304]1996.

[147305]

[147306]

[147307]13888.

[147308]Zhao, D.; McCaffery, P.; Ivins, K. J.; Neve, R. L.; Hogan, P.;

[147309]Chin, W. W.; Drager, U. C.: Molecular identification of a
major retinoic-acid-synthesizing

[147310]enzyme, a retinaldehyde-specific dehydrogenase. Europ.

J. Biochem. 240:

[147311]15–22, 1996.

[147312]

[147313]

[147314]13889.

[147315]McGwire, G. B.; Tan, F.; Michel, B.; Rehli, M.; Skidgel, R. A.

[147316]: Identification of a membrane-bound carboxypeptidase
as the mammalian

[147317]homolog of duck gp180, a hepatitis B virus-binding pro-
tein. Life

[147318]Sci. 60: 715–724, 1997.

[147319]

[147320]

[147321]13890.

[147322]Riley, D. A.; Tan, F.; Miletich, D. J.; Skidgel, R. A.: Chromo-
somal

[147323]localization of the genes for human carboxypeptidase D
(CPD) and the

[147324]active 50-kilodalton subunit of human carboxypeptidase
N (CPN1). Genomics 50:

[147325]105–108, 1998.

[147326]

[147327]

[147328]13891.

[147329]Tan, F.; Rehli, M.; Krause, S. W.; Skidgel, R. A.: Sequence
of

[147330]human carboxypeptidase D reveals it to be a member of
the regulatory

[147331]carboxypeptidase family with three tandem active site do-
mains. Biochem.

[147332]J. 327: 81–87, 1997.

[147333]

[147334]

[147335]13892.

[147336]Gebhard, W.; Schube, M.; Eulitz, M.: cDNA cloning and
complete

[147337]primary structure of the small, active subunit of human
carboxypeptidase

[147338]N (kininase 1). Eur. J. Biochem. 178: 603–607, 1989.

[147339]

[147340]

[147341]13893.

[147342]Skidgel, R. A.; Bennett, C. D.; Schilling, J. W.; Tan, F.;
Weerasinghe,

[147343]D. K.; Erdos, E. G.: Amino acid sequence of the N-
terminus and selected

[147344]tryptic peptides of the active subunit of human plasma
carboxypeptidase

[147345]N: comparison with other carboxypeptidases. Biochem.
Biophys. Res.

[147346]Commun. 154: 1323–1329, 1988.

[147347]

[147348]

[147349]13894.

[147350]Rajadhyaksha, A. Riviere, M.; Van Vooren, P.; Szpirer, J.;
Szpirer,

[147351]C.; Babin, J.; Bina, M.: Assignment of AR1, transcription
factor

[147352]20 (TCF20), to human chromosome 22q13.3 with somatic
cell hybrids

[147353]and in situ hybridization. Cytogenet. Cell Genet. 81:

176–177, 1998.

[147354]

[147355]

[147356]13895.

[147357]Sanz, L.; Moscat, J.; Diaz–Meco, M. T.: Molecular characterization

[147358]of a novel transcription factor that controls stromelysin expression. Molec.

[147359]Cell. Biol. 15: 3164–3170, 1995.

[147360]

[147361]

[147362]13896.

[147363]Ding, H.; Descheemaeker, K.; Marynen, P.; Nelles, L.; Carvalho,

[147364]T.; Carmo–Fonseca, M.; Collen, D.; Belayew, A.: Characterization

[147365]of a helicase–like transcription factor involved in the expression

[147366]of the human plasminogen activator inhibitor–1 gene.

DNA Cell Biol. 15:

[147367]429–442, 1996.

[147368]

[147369]

[147370]13897.

[147371]Lin, Y.; Sheridan, P. L.; Jones, K. A.; Evans, G. A.: The
HIP116

[147372]SNF2/SWI2-related transcription factor gene (SNF2L3) is
located on

[147373]human chromosome 3q25.1-q26.1 Genomics 27:
381-382, 1995.

[147374]

[147375]

[147376]13898.

[147377]Moinova, H. R.; Chen, W.-D.; Shen, L.; Smiraglia, D.;
Olechnowicz,

[147378]J.; Ravi, L.; Kasturi, L.; Myeroff, L.; Plass, C.; Parsons, R.;
Minna,

[147379]J.; Willson, J. K. V.; Green, S. B.; Issa, J.-P.; Markowitz, S. D.

[147380]: HLTF gene silencing in human colon cancer. Proc. Nat.
Acad. Sci. 99:

[147381]4562-4567, 2002.

[147382]

[147383]

[147384]13899.

[147385]Sheridan, P. L.; Schorpp, M.; Voz, M. L.; Jones, K. A.:

Cloning

[147386]of an SNF2/SWI2-related protein that binds specifically to
the SPH

[147387]motifs of the SV40 enhancer and to the HIV-1 promoter. J.
Biol. Chem. 270:

[147388]4575-4587, 1995.

[147389]

[147390]

[147391]13900.

[147392]Ambros, P. F.; Schmid, J.; Rumpler, S.; Binder, B. R.; de
Martin,

[147393]R.: Localization of the human I-kappa-B kinase-beta
(IKBKB) to chromosome

[147394]8p11.2 by fluorescence in situ hybridization and radiation
hybrid

[147395]mapping. Genomics 54: 575-576, 1998.

[147396]

[147397]

[147398]13901.

[147399]Pasparakis, M.; Courtois, G.; Hafner, M.; Schmidt-Sup-
prian, M.;

[147400]Nenci, A.; Toksoy, A.; Krampert, M.; Goebeler, M.;
Gillitzer, R.;

[147401]Israel, A.; Krieg, T.; Rajewsky, K.; Haase, I.: TNF-mediated
inflammatory

[147402]skin disease in mice with epidermis-specific deletion of
IKK2. Nature 417:

[147403]861–866, 2002.

[147404]

[147405]

[147406]13902.

[147407]Rossi, A.; Kapahi, P.; Natoli, G.; Takahashi, T.; Chen, Y.;
Karin,

[147408]M.; Santoro, M. G.: Anti-inflammatory cyclopentenone
prostaglandins

[147409]are direct inhibitors of I-kappa-B kinase. Nature 403:
103–108,

[147410]2000.

[147411]

[147412]

[147413]13903.

[147414]Shindo, M.; Nakano, H.; Sakon, S.; Yagita, H.; Mihara, M.;
Okumura,

[147415]K.: Assignment of I-kappa-B kinase beta (IKBKB) to human
chromosome

[147416]band 8p12-p11 by in situ hybridization. Cytogenet. Cell
Genet. 82:

[147417]32-33, 1998.

[147418]

[147419]

[147420]13904.

[147421]Tang, G.; Yang, J.; Minemoto, Y.; Lin, A.: Blocking cas-
pase-3-mediated

[147422]proteolysis of IKK-beta suppresses TNF-alpha-induced
apoptosis. Molec.

[147423]Cell 8: 1005-1016, 2001.

[147424]

[147425]

[147426]13905.

[147427]Woronicz, J. D.; Gao, X.; Cao, Z.; Rothe, M.; Goeddel, D. V.:

[147428]IkappaB kinase-beta: NF-kappa-B activation and complex formation with

[147429]IkappaB kinase-alpha and NIK. Science 278: 866-869, 1997.

[147430]

[147431]

[147432]13906.

[147433]Yin, M.-J.; Yamamoto, Y.; Gaynor, R. B.: The anti-inflammatory

[147434]agents aspirin and salicylate inhibit the activity of I-kappa-B kinase-beta. Nature 396:

[147435]77-80, 1998.

[147436]

[147437]

[147438]13907.

[147439]Zandi, E.; Rothwarf, D. M.; Delhase, M.; Hayakawa, M.;

Karin,

[147440]M.: The IkappaB kinase complex (IKK) contains two kinase subunits,

[147441]IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappa-B

[147442]activation. Cell 91: 243-252, 1997.

[147443]

[147444]

[147445]13908.

[147446]Aoki, M.; Hamada, F.; Sugimoto, T.; Sumida, S.; Akiyama, T.; Toyoshima,

[147447]K.: The human cot proto-oncogene encodes two protein serine/threonine

[147448]kinases with different transforming activities by alternative initiation

[147449]of translation. J. Biol. Chem. 268: 22723-22732, 1993.

[147450]

[147451]

[147452]13909.

[147453]Kitson, J.; Raven, T.; Jiang, Y.-P.; Goeddel, D. V.; Giles, K.

[147454]M.; Pun, K.-T.; Grinham, C. J.; Brown, R.; Farrow, S. N.: A
death-domain-containing

[147455]receptor that mediates apoptosis. Nature 384: 372–375,
1996.

[147456]

[147457]

[147458]13910.

[147459]Marsters, S. A.; Sheridan, J. P.; Donahue, C. J.; Pitti, R. M.;

[147460]Gray, C. L.; Goddard, A. D.; Bauer, K. D.; Ashkenazi, A.:
Apo-3,

[147461]a new member of the tumor necrosis factor receptor fam-
ily, contains

[147462]a death domain and activates apoptosis and NF-kappa-B.
Curr. Biol. 6:

[147463]1669–1676, 1996.

[147464]

[147465]

[147466]13911.

[147467]Screaton, G. R.; Xu, X.-N.; Olsen, A. L.; Cowper, A. E.; Tan,
R.;

[147468]McMichael, A. J.; Bell, J. I.: LARD: a new lymphoid-specific death

[147469]domain containing receptor regulated by alternative pre-mRNA splicing. Proc.

[147470]Nat. Acad. Sci. 94: 4615–4619, 1997.

[147471]

[147472]

[147473]13912.

[147474]Wang, E. C. Y.; Thern, A.; Denzel, A.; Kitson, J.; Farrow, S. N.;

[147475]Owen, M. J.: DR3 regulates negative selection during thymocyte development. Molec.

[147476]Cell. Biol. 21: 3451–3461, 2001.

[147477]

[147478]

[147479]13913.

[147480]Bataille, V.; Hiles, R.; Bishop, J. A. N.: Retinoblastoma, melanoma

[147481]and the atypical mole syndrome. Brit. J. Derm. 132: 134–138, 1995.

[147482]

[147483]

[147484]13914.

[147485]Chilosi, M.; Doglioni, C.; Yan, Z.; Lestani, M.; Menestrina, F.;

[147486]Sorio, C.; Benedetti, A.; Vinante, F.; Pizzolo, G.; Inghirami, G.

[147487]: Differential expression of cyclin-dependent kinase 6 in cortical

[147488]thymocytes and T-cell lymphoblastic lymphoma/leukemia. Am. J. Pathol. 152:

[147489]209–217, 1998.

[147490]

[147491]

[147492]13915.

[147493]Costello, J. F.; Plass, C.; Arap, W.; Chapman, V. M.; Held, W.

[147494]A.; Berger, M. S.; Su Huang, H. J.; Cavenee, W. K.: Cyclin-dependent

[147495]kinase 6 (cdk6) amplification in human gliomas identified using two-dimensional

[147496]separation of genomic DNA. Cancer Res. 57: 1250–1254,
1997.

[147497]

[147498]

[147499]13916.

[147500]Guan, K.–L.; Jenkins, C. W.; Li, Y.; Nichols, M. A.; Wu, X.;
O'Keefe,

[147501]C. L.; Matera, A. G.; Xiong, Y.: Growth suppression by p18,
a p16(INK4/MTS1)–

[147502]and p14(INK4B/MTS2)–related CDK6 inhibitor, correlates
with wild–type

[147503]pRb function. Genes Dev. 8: 2939–2952, 1994.

[147504]

[147505]

[147506]13917.

[147507]Lien, H.–C.; Lin, C.–W.; Huang, P.–H.; Chang, M.–L.; Hsu,
S.–M.

[147508]: Expression of cyclin–dependent kinase 6 (cdk6) and fre–
quent loss

[147509]of CD44 in nasal–nasopharyngeal NK/T–cell lymphomas:
comparison with

[147510]CD56-negative peripheral T-cell lymphomas. Lab. Invest.
80: 893-900,
[147511]2000.

[147512]

[147513]

[147514]13918.

[147515]Meyerson, M.; Harlow, E.: Identification of G1 kinase activ-
ity

[147516]for cdk6, a novel cyclin D partner. Molec. Cell. Biol. 14:
2077-2086,

[147517]1994.

[147518]

[147519]

[147520]13919.

[147521]Shennan, M. G.; Badin, A.-C.; Walsh, S.; Summers, A.;
From, L.;

[147522]McKenzie, M.; Goldstein, A. M.; Tucker, M. A.; Hogg, D.;
Lassam, N.

[147523]: Lack of germline CDK6 mutations in familial melanoma.
Oncogene 19:

[147524]1849–1852, 2000.

[147525]

[147526]

[147527]13920.

[147528]Traboulsi, E. I.; Zimmerman, L. E.; Manz, H. J.: Cutaneous
malignant

[147529]melanoma in survivors of heritable retinoblastoma. Arch.
Ophthalmol. 106:

[147530]1059–1061, 1988.

[147531]

[147532]

[147533]13921.

[147534]Blais, A.; Labrie, Y.; Pouliot, F.; Lachance, Y.; Labrie, C.:

[147535]Structure of the gene encoding the human cyclin-de-
pendent kinase inhibitor

[147536]p18 and mutational analysis in breast cancer. Biochem.
Biophys. Res.

[147537]Commun. 247: 146–153, 1998.

[147538]

[147539]

[147540]13922.

[147541]Lapointe, J.; Lachance, Y.; Labrie, Y.; Labrie, C.: A p18 mutant

[147542]defective in CDK6 binding in human breast cancer cells.

Cancer Res. 56:

[147543]4586–4589, 1996.

[147544]

[147545]

[147546]13923.

[147547]Bontemps, Y.; Maquart, F.-X.; , Wegrowski, Y.: Human UDP-glucose

[147548]dehydrogenase gene: complete cloning and transcription start mapping. Biochem.

[147549]Biophys. Res. Commun. 275: 981–985, 2000.

[147550]

[147551]

[147552]13924.

[147553]Hempel, J.; Perozich, J.; Romovacek, H.; Hinich, A.; Kuo, I.;

Feingold,

[147554]D. S.: UDP-glucose dehydrogenase from bovine liver: primary structure

[147555]and relationship to other dehydrogenases. Protein Sci. 3: 1074-1080,

[147556]1994.

[147557]

[147558]

[147559]13925.

[147560]Marcu, O.; Stathakis, D. G.; Marsh, J. L.: Assignment of the UGDH

[147561]locus encoding UDP-glucose dehydrogenase to human chromosome band

[147562]4p15.1 by radiation hybrid mapping. Cytogenet. Cell Genet. 86: 244-245,

[147563]1999.

[147564]

[147565]

[147566]13926.

[147567]Spicer, A. P.; Kaback, L. A.; Smith, T. J.; Seldin, M. F.:

Molecular

[147568]cloning and characterization of the human and mouse
UDP-glucose dehydrogenase

[147569]genes. J. Biol. Chem. 273: 25117-25124, 1998.

[147570]

[147571]

[147572]13927.

[147573]Walsh, E. C.; Stainier, D. Y. R.: UDP-glucose dehydroge-
nase required

[147574]for cardiac valve formation in zebrafish. Science 293:
1670-1674,

[147575]2001.

[147576]

[147577]

[147578]13928.

[147579]Bernard, M.; Sanseau, P.; Henry, C.; Couturier, A.; Prigent,
C.

[147580]: Cloning of STK13, a third human protein kinase related
to Drosophila

[147581]Aurora and budding yeast Ipl1 that maps on chromosome

19q13.3-ter. Genomics 53:

[147582]406-409, 1998.

[147583]

[147584]

[147585]13929.

[147586]Tseng, T.-C.; Chen, S.-H.; Hsu, Y.-P. P.; Tang, T. K.: Protein
tein

[147587]kinase profile of sperm and eggs: cloning and characteri-
zation of

[147588]two novel testis-specific protein kinases (AIE1, AIE2) re-
lated to

[147589]yeast and fly chromosome segregation regulators. DNA
Cell Biol. 17:

[147590]823-833, 1998.

[147591]

[147592]

[147593]13930.

[147594]Becker, W.; Weber, Y.; Wetzel, K.; Eirmbter, K.; Tejedor, F.
J.;

[147595]Joost, H.-G.: Sequence characteristics, subcellular local-

ization,

[147596]and substrate specificity of DYRK-related kinases, a novel family

[147597]of dual specificity protein kinases. J. Biol. Chem. 273: 25893–25902,

[147598]1998.

[147599]

[147600]

[147601]13931.

[147602]Hsu, H.; Xiong, J.; Goeddel, D. V.: The TNF receptor 1-associated

[147603]protein TRADD signals cell death and NF-kappa-B activation. Cell 81:

[147604]495–504, 1995.

[147605]

[147606]

[147607]13932.

[147608]Pan, M.-G.; Xiong, J.; Copeland, N. G.; Gilbert, D. J.; Jenkins, ins,

[147609]N. A.; Goeddel, D. V.: Sequence, genomic organization,

and chromosome

[147610] localization of the mouse TRADD gene. J. Inflamm. 46:
168–175, 1996.

[147611]

[147612]

[147613] 13933.

[147614] Bao, J.; Zervos, A. S.: Isolation and characterization of
Nmi,

[147615] a novel partner of Myc proteins. Oncogene 12:
2171–2176, 1996.

[147616]

[147617]

[147618] 13934.

[147619] Zhu, M.; John, S.; Berg, M.; Leonard, W. J.: Functional as-
sociation

[147620] of Nmi with Stat5 and Stat1 in IL-2- and IFN-
gamma-mediated signaling. Cell 96:

[147621] 121–130, 1999.

[147622]

[147623]

[147624]13935.

[147625]Phillips, N. J.; Zeigler, M. R.; Deaven, L. L.: A cDNA from
the

[147626]ovarian cancer critical region of deletion on chromosome
17p13.3. Cancer

[147627]Lett. 102: 85–90, 1996.

[147628]

[147629]

[147630]13936.

[147631]Schultz, D. C.; Vanderveer, L.; Berman, D. B.; Hamilton, T.
C.;

[147632]Wong, A. J.; Godwin, A. K.: Identification of two candidate
tumor

[147633]suppressor genes on chromosome 17p13.3. Cancer Res.
56: 1997–2002,

[147634]1996.

[147635]

[147636]

[147637]13937.

[147638]Kirchhausen, T.; Davis, A. C.; Frucht, S.; O'Brine Greco, B.;

Payne,

[147639]G. S.; Tubb, B.: AP17 and AP19, the mammalian small chains of the

[147640]clathrin-associated protein complexes show homology to Yap17p, their

[147641]putative homolog in yeast. J. Biol. Chem. 266: 11153–11157, 1991.

[147642]

[147643]

[147644]13938.

[147645]Peyrard, M.; Parveneh, S.; Lagercrantz, S.; Ekman, M.; Fransson,

[147646]I.; Sahlen, S.; Dumanski, J. P.: Cloning, expression pattern, and

[147647]chromosomal assignment to 16q23 of the human gamma-adaptin gene (ADTG). Genomics 50:

[147648]275–280, 1998.

[147649]

[147650]

[147651]13939.

[147652]Takatsu, H.; Sakurai, M.; Shin, H.-W.; Murakami, K.;

Nakayama,

[147653]K.: Identification and characterization of novel clathrin adaptor-related

[147654]proteins. J. Biol. Chem. 273: 24693–24700, 1998.

[147655]

[147656]

[147657]13940.

[147658]Albarosa, R.; Colombo, B. M.; Roz, L.; Magnani, I.; Pollo, B.;

[147659]Cirenei, N.; Giani, C.; Conti, A. M.; DiDonato, S.; Finocchiario, G.

[147660]: Deletion mapping of gliomas suggests the presence of two small regions

[147661]for candidate tumor-suppressor genes in a 17-cM interval on chromosome

[147662]10q. Am. J. Hum. Genet. 58: 1260–1267, 1996.

[147663]

[147664]

[147665]13941.

[147666]Maier, D.; Zhang, Z.; Taylor, E.; Hamou, M.–F.; Gratzl, O.;

Van

[147667]Meir, E. G.; Scott, R. J.; Merlo, A.: Somatic deletion mapping on

[147668]chromosome 10 and sequence analysis of PTEN/MMAC1 point to the 10q25–26

[147669]region as the primary target in low–grade and high–grade gliomas. Oncogene 16:

[147670]3331–3335, 1998.

[147671]

[147672]

[147673]13942.

[147674]Rasheed, B. K.; McLendon, R. E.; Friedman, H. S.; Friedman, A.

[147675]H.; Fuchs, H. E.; Bigner, D. D.; Bigner, S. H.: Chromosome 10 deletion

[147676]mapping in human gliomas: a common deletion region in 10q25. Oncogene 10:

[147677]2243–2246, 1995.

[147678]

[147679]

[147680]13943.

[147681]Andre, E.; Conquet, F.; Steinmayr, M.; Stratton, S. C.; Porciatti,

[147682]V.; Becker–Andre, M.: Disruption of retinoid–related orphan receptor

[147683]beta changes circadian behavior, causes retinal degeneration and leads

[147684]to vacillans phenotype in mice. EMBO J. 17: 3867–3877, 1998.

[147685]

[147686]

[147687]13944.

[147688]Paravicini, G.; Steinmayr, M.; Andre, E.; Becker–Andre, M.: The

[147689]metastasis suppressor candidate nucleotide diphosphate kinase NM23

[147690]specifically interacts with members of the ROR/RZR nuclear orphan

[147691]receptor subfamily. Biochem. Biophys. Res. Commun. 227: 82–87, 1996.

[147692]

[147693]

[147694]13945.

[147695]Sirlin, J. L.: Vacillans, a neurological mutant in the house
mouse

[147696]linked with brown. J. Genet. 54: 42–48, 1956.

[147697]

[147698]

[147699]13946.

[147700]Vidal–Puig, A.; Solanes, G.; Grujic, D.; Flier, J. S.; Lowell,

[147701]B. B.: UCP3: an uncoupling protein homologue expressed
preferentially

[147702]and abundantly in skeletal muscle and brown adipose tis-
sue. Biochem.

[147703]Biophys. Res. Commun. 235: 79–82, 1997.

[147704]

[147705]

[147706]13947.

[147707]Vidal–Puig, A. J.; Grujic, D.; Zhang, C.–Y.; Hagen, T.; Boss,

[147708]O.; Ido, Y.; Szczepanik, A.; Wade, J.; Mootha, V.; Cortright,
R.;

[147709]Muoio, D. M.; Lowell, B. B.: Energy metabolism in uncou-
pling protein

[147710]3 gene knockout mice. J. Biol. Chem. 275: 16258–16266,
2000.

[147711]

[147712]

[147713]13948.

[147714]Lovering, R.; Hanson, I. M.; Borden, K. L. B.; Martin, S.;
O'Reilly,

[147715]N. J.; Evan, G. I.; Rahman, D.; Pappin, D. J. C.; Trowsdale,
J.; Freemont,

[147716]P. S.: Identification and preliminary characterization of a
protein

[147717]motif related to the zinc finger. Proc. Nat. Acad. Sci. 90:
2112–2116,

[147718]1993.

[147719]

[147720]

[147721]13949.

[147722]Bourdi, M.; Demady, D.; Martin, J. L.; Jabbour, S. K.; Mar-
tin,

[147723]B. M.; George, J. W.; Pohl, L. R.: cDNA cloning and baculovirus expression

[147724]of the human liver endoplasmic reticulum P58: characterization as

[147725]a protein disulfide isomerase isoform, but not as a protease or a

[147726]carnitine acyltransferase. Arch. Biochem. Biophys. 323: 397–403,

[147727]1995.

[147728]

[147729]

[147730]13950.

[147731]Briquet–Laugier, V.; Xia, Y.–R.; Rooke, K.; Mehrabian, M.; Lysis,

[147732]A. J.; Doolittle, M. H.: Mapping of three members of the mouse protein

[147733]disulfide isomerase family. Mammalian Genome 9: 176–177, 1998.

[147734]

[147735]

[147736]13951.

[147737]Hirano, N.; Shibasaki, F.; Sakai, R.; Tanaka, T.; Nishida, J.;

[147738]Yazaki, Y.; Takenawa, T.; Hirai, H.: Molecular cloning of
the human

[147739]glucose-regulated protein ERp57/GRP58, a thiol-
dependent reductase:

[147740]identification of its secretory form and inducible expres-
sion by the

[147741]oncogenic transformation. Europ. J. Biochem. 234:
336-342, 1995.

[147742]

[147743]

[147744]13952.

[147745]Koivunen, P.; Helaakoski, T.; Annunen, P.; Veijola, J.;
Raisanen,

[147746]S.; Pihlajaniemi, T.; Kivirikko, K. I.: ERp60 does not substi-
tute

[147747]for protein disulphide isomerase as the beta-subunit of
prolyl 4-hydroxylase. Biochem.

[147748]J. 316: 599-605, 1996.

[147749]

[147750]

[147751]13953.

[147752]Koivunen, P.; Horelli-Kuitunen, N.; Helaakoski, T.; Karvonen, P.;

[147753]Jaakkola, M.; Palotie, A.; Kivirikko, K. I.: Structures of the human

[147754]gene for the protein disulfide isomerase-related polypeptide ERp60

[147755]and a processed gene and assignment of these genes to 15q15 and 1q21. Genomics 42:

[147756]397–404, 1997.

[147757]

[147758]

[147759]13954.

[147760]Oliver, J. D.; van der Wal, F. J.; Bulleid, N. J.; High, S.: Interaction

[147761]of the thiol-dependent reductase ERp57 with nascent glycoproteins. Science 275:

[147762]86–88, 1997.

[147763]

[147764]

[147765]13955.

[147766]Didsbury, J.; Weber, R. F.; Bokoch, G. M.; Evans, T.; Snyderman,

[147767]R.: rac, a novel ras-related family of proteins that are botulinum

[147768]toxin substrates. J. Biol. Chem. 264: 16378–16382, 1989.

[147769]

[147770]

[147771]13956.

[147772]Palmiter, R. D.; Cole, T. B.; Quaife, C. J.; Findley, S. D.: ZnT3,

[147773]a putative transporter of zinc into synaptic vesicles. Proc. Nat.

[147774]Acad. Sci. 93: 14934–14939, 1996.

[147775]

[147776]

[147777]13957.

[147778]Palmiter, R. D.; Findley, S. D.: Cloning and functional characterization

[147779]of a mammalian zinc transporter that confers resistance to zinc. EMBO

[147780]J. 14: 639–649, 1995.

[147781]

[147782]

[147783]13958.

[147784]Vallee, B. L.; Falchuk, K. H.: The biochemical basis of zinc

[147785]physiology. Physiol. Rev. 73: 79–118, 1993.

[147786]

[147787]

[147788]13959.

[147789]Berthelsen, J.; Viggiano, L.; Schulz, H.; Ferretti, E.; Con–

salez,

[147790]G. G.; Rocchi, M.; Blasi, F.: PKNOX1, a gene encoding
PREP1, a new

[147791]regulator of Pbx activity, maps on human chromosome
21q22.3 and murine

[147792]chromosome 17B/C. Genomics 47: 323–324, 1998.

[147793]

[147794]

[147795]13960.

[147796]Chen, H.; Rossier, C.; Nakamura, Y.; Lynn, A.; Chakravarti,
A.;

[147797]Antonarakis, S. E.: Cloning of a novel homeobox–contain–
ing gene,

[147798]PKNOX1, and mapping to human chromosome 21q22.3.
Genomics 41: 193–200,

[147799]1997.

[147800]

[147801]

[147802]13961.

[147803]Morita, K.; Furuse, M.; Fujimoto, K.; Tsukita, S.: Claudin

multigene

[147804]family encoding four-transmembrane domain protein
components of tight

[147805]junction strands. Proc. Nat. Acad. Sci. 96: 511-516, 1999.

[147806]

[147807]

[147808]13962.

[147809]Chiang, P.-W.; Fogel, E.; Jackson, C. L.; Lieuallen, K.;
Lennon,

[147810]G.; Qu, X.; Wang, S.-Q.; Kurnit, D. M.: Isolation, sequenc-
ing, and

[147811]mapping of the human homologue of the yeast transcrip-
tion factor,

[147812]SPT5. Genomics 38: 421-424, 1996.

[147813]

[147814]

[147815]13963.

[147816]Chiang, P.-W.; Stubbs, L.; Zhang, L.; Kurnit, D. M.: Isola-
tion

[147817]of murine SPT5 homologue: completion of the isolation

and characterization

[147818]of human and murine homologues of yeast chromatin
structural protein

[147819]complex SPT4, SPT5, and SPT6. Genomics 47: 426–428,
1998.

[147820]

[147821]

[147822]13964.

[147823]Grover, J.; Chen, X.–N.; Korenberg, J. R.; Roughley, P. J.:
The

[147824]structure and chromosome location of the human chon-
droadherin gene

[147825](CHAD). Genomics 45: 379–385, 1997.

[147826]

[147827]

[147828]13965.

[147829]Larsson, T.; Sommarin, Y.; Paulsson, M.; Antonsson, P.;
Hedbom,

[147830]E.; Wendel, M.; Heinegard, D.: Cartilage matrix proteins: a
basic

[147831]36–kDa protein with a restricted distribution to cartilage

and bone. J.

[147832]Biol. Chem. 266: 20428–20433, 1991.

[147833]

[147834]

[147835]13966.

[147836]Lienard, P.; Riviere, M.; Van Vooren, P.; Szpirer, C.;
Szpirer,

[147837]J.: Assignment of SND1, the gene encoding coactivator
p100, to human

[147838]chromosome 7q31.3 and rat chromosome 4q23 by in situ
hybridization. Cytogenet.

[147839]Cell Genet. 90: 253–254, 2000.

[147840]

[147841]

[147842]13967.

[147843]Tong, X.; Drapkin, R.; Yalamanchili, R.; Mosialos, G.; Kieff,
E.

[147844]: The Epstein–Barr virus nuclear protein 2 acidic domain
forms a complex

[147845]with a novel cellular coactivator that can interact with

TFIIE. Molec.

[147846]Cell. Biol. 15: 4735–4744, 1995.

[147847]

[147848]

[147849]13968.

[147850]Tribioli, C.; Frasch, M.; Lufkin, T.: Bapx1: an evolutionary conserved

[147851]homologue of the Drosophila bagpipe homeobox gene is expressed in

[147852]splanchnic mesoderm and the embryonic skeleton. Mech. Dev. 65: 145–162,

[147853]1997.

[147854]

[147855]

[147856]13969.

[147857]Tribioli, C.; Lufkin, T.: Molecular cloning, chromosomal mapping

[147858]and developmental expression of BAPX1, a novel human homeobox-containing

[147859]gene homologous to Drosophila bagpipe. Gene 203:

225–233, 1997.

[147860]

[147861]

[147862]13970.

[147863]Yoshiura, K.-I.; Murray, J. C.: Sequence and chromosomal assignment

[147864]of human BAPX1, a bagpipe-related gene, to 4p16.1: a candidate gene

[147865]for skeletal dysplasia. Genomics 45: 425–428, 1997.

[147866]

[147867]

[147868]13971.

[147869]de Coo, R. F. M.; Buddiger, P.; Smeets, H. J. M.; van Oost, B.

[147870]A.: Molecular cloning and characterization of the human mitochondrial

[147871]NADH:oxidoreductase 10-kDa gene (NDUFV3). Genomics 45: 434–437,

[147872]1997.

[147873]

[147874]

[147875]13972.

[147876]Gautvik, K. M.; de Lecea, L.; Gautvik, V. T.; Danielson, P. E.;

[147877]Tranque, P.; Dopazo, A.; Bloom, F. E.; Sutcliffe, J. G.:
Overview

[147878]of the most prevalent hypothalamus-specific mRNAs as
identified by

[147879]directional tag PCR subtraction. Proc. Nat. Acad. Sci. 93:
8733–8738,

[147880]1996.

[147881]

[147882]

[147883]13973.

[147884]Fritzler, M. J.; Lung, C.-C.; Hamel, J. C.; Griffith, K. J.;
Chan,

[147885]E. K. L.: Molecular characterization of golgin-245, a novel
Golgi

[147886]complex protein containing a granin signature. J. Biol.
Chem. 270:

[147887]31262–31268, 1995.

[147888]

[147889]

[147890]13974.

[147891]Gencik, M.; Dahmen, N.; Wieczorek, S.; Kasten, M.; Bierbrauer,

[147892]J.; Anghelescu, I.; Szegedi, A.; Menezes Saecker, A. M.; Epplen, J.

[147893]T.: A prepro-orexin gene polymorphism is associated with narcolepsy. *Neurology* 56:

[147894]115–117, 2001.

[147895]

[147896]

[147897]13975.

[147898]Hagan, J. J.; Leslie, R. A.; Patel, S.; Evans, M. L.; Wattam, T.

[147899]A.; Holmes, S.; Benham, C. D.; Taylor, S. G.; Routledge, C.; Hemmati,

[147900]P.; Munton, R. P.; Ashmeade, T. E.; Shah, A. S.; Hatcher, J. P.; Hatcher,

[147901]P. D.; Jones, D. N. C.; Smith, M. I.; Piper, D. C.; Hunter, A. J.;

[147902]Porter, R. A.; Upton, N.: Orexin A activates locus coeruleus cell

[147903]firing and increases arousal in the rat. Proc. Nat. Acad. Sci. 96:

[147904]10911–10916, 1999.

[147905]

[147906]

[147907]13976.

[147908]Hara, J.; Beuckmann, C. T.; Nambu, T.; Willie, J. T.; Chemelli,

[147909]R. M.; Sinton, C. M.; Sugiyama, F.; Yagami, K.; Goto, K.; Yanagisawa,

[147910]M.; Sakurai, T.: Genetic ablation of orexin neurons in mice results

[147911]in narcolepsy, hypophagia, and obesity. Neuron 30: 345–354, 2001.

[147912]

[147913]

[147914]13977.

[147915]Hungs, M.; Lin, L.; Okun, M.; Mignot, E.: Polymorphisms in the

[147916]vicinity of the hypocretin/orexin are not associated with human narcolepsy. Neurology 57:

[147917]1893–1895, 2001.

[147918]

[147919]

[147920]13978.

[147921]Nishino, S.; Ripley, B.; Overeem, S.; Lammers, G. J.; Mignot, E.

[147922]: Hypocretin (orexin) deficiency in human narcolepsy. Lancet 355:

[147923]39–40, 2000.

[147924]

[147925]

[147926]13979.

[147927]Chen, L.–S. K.; Lo, C. F.; Numann, R.; Cuddy, M.: Characterization

[147928]of the human and rat phospholemman (PLM) cDNAs and localization of

[147929]the human PLM gene to chromosome 19q13.1. Genomics 41: 435–443,

[147930]1997.

[147931]

[147932]

[147933]13980.

[147934]Bianchi, M. C.; Tosetti, M.; Fornai, F.; Alessandri, M. G.;
Cipriani,

[147935]P.; De Vito, G.; Canapicchi, R.: Reversible brain creatine
deficiency

[147936]in two sisters with normal blood creatine level. Ann. Neu-
rol. 47:

[147937]511–513, 2000.

[147938]

[147939]

[147940]13981.

[147941]Humm, A.; Fritsche, E.; Mann, K.; Gohl, M.; Huber, R.: Re-
combinant

[147942]expression and isolation of human L-arginine:glycine
amidinotransferase

[147943]and identification of its active-site cysteine residue.
Biochem.

[147944]J. 322: 771–776, 1997.

[147945]

[147946]

[147947]13982.

[147948]Humm, A.; Huber, R.; Mann, K.: The amino acid sequences
of human

[147949]and pig L-arginine:glycine amidinotransferase. FEBS Lett.
339: 101–107,

[147950]1994.

[147951]

[147952]

[147953]13983.

[147954]Sakurai, T.; Amemiya, A.; Ishii, M.; Matsuzaki, I.; Chemelli,

[147955]R. M.; Tanaka, H.; Williams, S. C.; Richardson, J. A.; Ko-
złowski,

[147956]G. P.; Wilson, S.; Arch, J. R. S.; Buckingham, R. E.; Haynes,
A. C.;

[147957]Carr, S. A.; Annan, R. S.; McNulty, D. E.; Liu, W.–S.; Terrett,
J.

[147958]A.; Elshourbagy, N. A.; Bergsma, D. J.; Yanagisawa, M.:

Orexins and

[147959] orexin receptors: a family of hypothalamic neuropeptides
and G protein-coupled

[147960] receptors that regulate feeding behavior. Cell 92:
573–585, 1998.

[147961]

[147962]

[147963] 13984.

[147964] Item, C. B.; Stockler-Ipsiroglu, S.; Stromberger, C.; Muhl,
A.;

[147965] Alessandri, M. G.; Bianchi, M. C.; Tosetti, M.; Fornai, F.;

[147966] Cioni,
G.: Arginine:glycine amidinotransferase deficiency: the
third inborn

[147967] error of creatine metabolism in humans. Am. J. Hum.
Genet. 69: 1127–1133,

[147968] 2001.

[147969]

[147970]

[147971] 13985.

[147972] Yasuda, H.; Shima, N.; Nakagawa, N.; Yamaguchi, K.; Ki–

nosaki,

[147973]M.; Mochizuki, S.; Tomoyasu, A.; Yano, K.; Goto, M.; Murakami, A.;

[147974]Tsuda, E.; Morinaga, T.; Higashio, K.; Udagawa, N.; Takahashi, N.;

[147975]Suda, T.: Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory

[147976]factor and is identical to TRANCE/RANKL. Proc. Nat. Acad. Sci. 95:

[147977]3597-3602, 1998.

[147978]

[147979]

[147980]13986.

[147981]Bey, F.; Silva Pereira, I.; Coux, O.; Viegas-Pequignot, E.; Recillas

[147982]Targa, F.; Nothwang, H. G.; Dutrillaux, B.; Scherrer, K.: The prosomal

[147983]RNA-binding protein p27K is a member of the alpha-type human prosomal

[147984]gene family. Molec. Gen. Genet. 237: 193-205, 1993.

[147985]

[147986]

[147987]13987.

[147988]Heard, D. J.; Norby, P. L.; Holloway, J.; Vissing, H.: Human

ERR-gamma,

[147989]a third member of the estrogen receptor-related receptor

(ERR) subfamily

[147990]of orphan nuclear receptors: tissue-specific isoforms are

expressed

[147991]during development in the adult. Molec. Endocr. 14:

382-392, 2000.

[147992]

[147993]

[147994]13988.

[147995]Hong, H.; Yang, L.; Stallcup, M. R.: Hormone-independent

transcriptional

[147996]activation and coactivator binding by novel orphan nuclear

receptor

[147997]ERR3. J. Biol. Chem. 274: 22618-22626, 1999.

[147998]

[147999]

[148000]13989.

[148001]Fisher, D. A.; Smith, J. F.; Pillar, J. S.; St. Denis, S. H.;
Cheng,

[148002]J. B.: Isolation and characterization of PDE9A, a novel hu-
man cGMP-specific

[148003]phosphodiesterase. J. Biol. Chem. 273: 15559-15564,
1998.

[148004]

[148005]

[148006]13990.

[148007]Guipponi, M.; Scott, H. S.; Kudoh, J.; Kawasaki, K.;
Shibuya, K.;

[148008]Shintani, A.; Asakawa, S.; Chen, H.; Lalioti, M. D.; Rossier,
C.;

[148009]Minoshima, S.; Shimizu, N.; Antonarakis, S. E.: Identifica-
tion and

[148010]characterization of a novel cyclic nucleotide phosphodi-
esterase gene

[148011](PDE9A) that maps to 21q22.3: alternative splicing of
mRNA transcripts,

[148012]genomic structure and sequence. Hum. Genet. 103:
386-392, 1998.

[148013]

[148014]

[148015]13991.

[148016]Soderling, S. H.; Bayuga, S. J.; Beavo, J. A.: Identification

[148017]and characterization of a novel family of cyclic nucleotide
phosphodiesterases. J.

[148018]Biol. Chem. 273: 15553–15558, 1998.

[148019]

[148020]

[148021]13992.

[148022]Ishibashi, K.; Kuwahara, M.; Gu, Y.; Kageyama, Y.;

Tohsaka, A.;

[148023]Suzuki, F.; Marumo, F.; Sasaki, S.: Cloning and functional
expression

[148024]of a new water channel abundantly expressed in the testis
permeable

[148025]to water, glycerol, and urea. J. Biol. Chem. 272:
20782–20786, 1997.

[148026]

[148027]

[148028]13993.

[148029]Ishibashi, K.; Yamauchi, K.; Kageyama, Y.; Saito–Ohara, F.;
Ikeuchi,

[148030]T.; Marumo, F.; Sasaki, S.: Molecular characterization of
human aquaporin–7

[148031]gene and its chromosomal mapping. Biochim. Biophys.
Acta 1399: 62–66,

[148032]1998.

[148033]

[148034]

[148035]13994.

[148036]Kuriyama, H.; Kawamoto, S.; Ishida, N.; Ohno, I.; Mita, S.;
Matsuzawa,

[148037]Y.; Matsubara, K.; Okubo, K.: Molecular cloning and ex-
pression of

[148038]a novel human aquaporin from adipose tissue with glyc-
erol permeability. Biochem.

[148039]Biophys. Res. Commun. 241: 53–58, 1997.

[148040]

[148041]

[148042]13995.

[148043]Billin, A. N.; Eilers, A. L.; Queva, C.; Ayer, D. E.: Mlx, a novel

[148044]Max-like BHLHZip protein that interacts with the Max network of transcription

[148045]factors. J. Biol. Chem. 274: 36344–36350, 1999.

[148046]

[148047]

[148048]13996.

[148049]Bjerknes, M.; Cheng, H.: TCFL4: a gene at 17q21.1 encoding a putative

[148050]basic helix-loop-helix leucine-zipper transcription factor.

Gene 181:

[148051]7–11, 1996.

[148052]

[148053]

[148054]13997.

[148055]Cairo, S.; Merla, G.; Urbinati, F.; Ballabio, A.; Reymond, A.:

[148056]WBSCR14, a gene mapping to the Williams–Beuren syn–

drome deleted region,

[148057]is a new member of the Mlx transcription factor network.

Hum. Molec.

[148058]Genet. 10: 617–627, 2001.

[148059]

[148060]

[148061]13998.

[148062]Fukuoka, S.–I.; Freedman, S. D.; Scheele, G. A.: A single
gene

[148063]encodes membrane–bound and free forms of GP–2, the
major glycoprotein

[148064]in pancreatic secretory (zymogen) granule membranes.

Proc. Nat. Acad.

[148065]Sci. 88: 2898–2902, 1991.

[148066]

[148067]

[148068]13999.

[148069]Fukuoka, S.–I.; Freedman, S. D.; Yu, H.; Sukhatme, V. P.;
Scheele,

[148070]G. A.: GP–2/THP gene family encodes self–binding glyco–

syphosphatidylinositol-anchored

[148071]proteins in apical secretory compartments of pancreas
and kidney. Proc.

[148072]Nat. Acad. Sci. 89: 1189-1193, 1992.

[148073]

[148074]

[148075]14000.

[148076]Fukuoka, S.-I.; Suzuki, M.; Okabayashi, K.; Takahashi, E.:
Assignment

[148077]of pancreatic zymogen granule membrane protein GP2
(GP2) to human

[148078]chromosome band 9q21.11 to q21.2 by in situ hybridiza-
tion. Cytogenet.

[148079]Cell Genet. 79: 231-232, 1997.

[148080]

[148081]

[148082]14001.

[148083]Kooy, J.; Toh, B.-H.; Pettitt, J. M.; Erlich, R.; Gleeson, P. A.

[148084]: Human autoantibodies as reagents to conserved Golgi
components:

[148085]characterization of a peripheral, 230-kDa compartment-specific Golgi

[148086]protein. J. Biol. Chem. 267: 20255–20263, 1992.

[148087]

[148088]

[148089]14002.

[148090]Snow, B. E.; Antonio, L.; Suggs, S.; Gutstein, H. B.;
Siderovski,

[148091]D. P.: Molecular cloning and expression analysis of rat
Rgs12 and

[148092]Rgs14. Biochem. Biophys. Res. Commun. 233: 770–777,
1997.

[148093]

[148094]

[148095]14003.

[148096]Kimple, R. J.; Kimple, M. E.; Betts, L.; Sondek, J.;
Siderovski,

[148097]D. P.: Structural determinants for GoLoco-induced inhibition of nucleotide

[148098]release by G-alpha subunits. Nature 416: 878–881, 2002.

[148099]

[148100]

[148101]14004.

[148102]Takesono, A.; et al; et al: Receptor-independent activators
of

[148103]heterotrimeric G-protein signalling pathways. J. Biol.
Chem. 274:

[148104]33202–33205, 1999.

[148105]

[148106]

[148107]14005.

[148108]Traver, S.; Bidot, C.; Spassky, N.; Baltauss, T.; de Tand,
M.–F.;

[148109]Thomas, J.–L.; Zalc, B.; Janoueix–Lerosey, I.; de Gunzburg,
J.: RGS14

[148110]is a novel Rap effector that preferentially regulates the
GTPase activity

[148111]of G-alpha-0. Biochem. J. 350: 19–29, 2000.

[148112]

[148113]

[148114]14006.

[148115]Buckbinder, L.; Velasco-Miguel, S.; Chen, Y.; Xu, N.; Tal-
bott,

[148116]R.; Gelbert, L.; Gao, J.; Seizinger, B. R.; Gutkind, J. S.; Kley,

[148117]N.: The p53 tumor suppressor targets a novel regulator of
G protein

[148118]signaling. Proc. Nat. Acad. Sci. 94: 7868–7872, 1997.

[148119]

[148120]

[148121]14007.

[148122]Chen, C.-K.; Wieland, T.; Simon, M. I.: RGS-r, a retinal
specific

[148123]RGS protein, binds an intermediate conformation of trans-
ducin and

[148124]enhances recycling. Proc. Nat. Acad. Sci. 93:
12885–12889, 1996.

[148125]

[148126]

[148127]14008.

[148128]Snow, B. E.; Antonio, L.; Suggs, S.; Siderovski, D. P.:

Cloning

[148129]of a retinally abundant regulator of G-protein signaling
(RGS-r/RGS16):

[148130]genomic structure and chromosomal localization of the
human gene. Gene 206:

[148131]247–253, 1998.

[148132]

[148133]

[148134]14009.

[148135]Zheng, B.; Chen, D.; Farquhar, M. G.: MIR16, a putative
membrane

[148136]glycerophosphodiester phosphodiesterase, interacts with
RGS16. Proc.

[148137]Nat. Acad. Sci. 97: 3999–4004, 2000.

[148138]

[148139]

[148140]14010.

[148141]LeBoeuf, R. C.; Caldwell, M.; Guo, Y.; Metz, C.; Davitz, M.
A.;

[148142]Olson, L. K.; Deeg, M. A.: Mouse glycosylphosphatidyli-

inositol-specific

[148143]phospholipase D (Gpld1) characterization. Mammalian
Genome 9: 710–714,

[148144]1998.

[148145]

[148146]

[148147]14011.

[148148]Scallan, B. J.; Fung, W.-J. C.; Tsang, T. C.; Li, S.; Kado-
Fong,

[148149]H.; Huang, K.-S.; Kochan, J. P.: Primary structure and
functional

[148150]activity of a phosphatidylinositol-glycan-specific phos-
pholipase D. Science 252:

[148151]446–448, 1991.

[148152]

[148153]

[148154]14012.

[148155]Schofield, J. N.; Rademacher, T. W.: Structure and expres-
sion

[148156]of the human glycosylphosphatidylinositol phospholipase

D1 (GPLD1)

[148157]gene. Biochim. Biophys. Acta 1494: 189–194, 2000.

[148158]

[148159]

[148160]14013.

[148161]Tsang, T. C.; Fung, W.-J.; Levine, J.; Metz, C. N.; Davitz, M.

[148162]A.; Burns, D. K.; Huang, K.-S.; Kochan, J. P.: Isolation and
expression

[148163]of two human glycosylphosphatidylinositol phospholipase
D (GPI-PLD)

[148164]cDNAs. (Abstract) FASEB J. (supp.) 6: A1922 only, 1992.

[148165]

[148166]

[148167]14014.

[148168]Schmitz, F.; Konigstorfer, A.; Sudhof, T. C.: RIBEYE, a com-
ponent

[148169]of synaptic ribbons: a protein's journey through evolution
provides

[148170]insight into synaptic ribbon function. Neuron 28:
857–872, 2000.

[148171]

[148172]

[148173]14015.

[148174]Bergelson, J. M.; Cunningham, J. A.; Droguett, G.; Kurt-Jones,

[148175]E. A.; Krithivas, A.; Hong, J. S.; Horwitz, M. S.; Crowell, R. L.;

[148176]Finberg, R. W.: Isolation of a common receptor for coxsackie B viruses

[148177]and adenoviruses 2 and 5. Science 275: 1320–1323, 1997.

[148178]

[148179]

[148180]14016.

[148181]Bowles, K. R.; Gibson, J.; Wu, J.; Shaffer, L. G.; Towbin, J. A.;

[148182]Bowles, N. E.: Genomic organization and chromosomal localization

[148183]of the human Coxsackievirus B–adenovirus receptor gene. Hum. Genet. 105:

[148184]354–359, 1999.

[148185]

[148186]

[148187]14017.

[148188]Carson, S. D.; Chapman, N. N.; Tracy, S. M.: Purification of
the

[148189]putative coxsackievirus B receptor from HeLa cells.

Biochem. Biophys.

[148190]Res. Commun. 233: 325–328, 1997.

[148191]

[148192]

[148193]14018.

[148194]Griffin, L. D.; Kearney, D.; Ni, J.; Jaffe, R.; Fricker, F. J.;

[148195]Webber, S.; Demmler, G.; Gelb, B. D.; Towbin, J. A.: Analy-
sis of

[148196]formalin–fixed and frozen myocardial autopsy samples for
viral genome

[148197]in childhood myocarditis and dilated cardiomyopathy with
endocardial

[148198]fibroelastosis using polymerase chain reaction (PCR). Car-
diovasc.

[148199]Path. 4: 3–11, 1995.

[148200]

[148201]

[148202]14019.

[148203]Martin, A. B.; Webber, S.; Fricker, F. J.; Jaffe, R.; Demmler,

[148204]G.; Kearney, D.; Zhang, Y.–H.; Bodurtha, J.; Gelb, B.; Ni, J.;
Bricker,

[148205]J. T.; Towbin, J. A.: Acute myocarditis: rapid diagnosis by
PCR in

[148206]children. Circulation 90: 330–339, 1994.

[148207]

[148208]

[148209]14020.

[148210]Pauschinger, M.; Bowles, N. E.; Fuentes–Garcia, F. J.;

Pham, V.;

[148211]Kuhl, U.; Schwimmbeck, P. L.; Schultheiss, H.–P.; Towbin,
J. A.:

[148212]Detection of adenoviral genome in the myocardium of
adult patients

[148213]with idiopathic left ventricular dysfunction. Circulation 99:

1348–1354,

[148214]1999.

[148215]

[148216]

[148217]14021.

[148218]Tomko, R. P.; Xu, R.; Philipson, L.: HCAR and MCAR: the
human

[148219]and mouse receptors for subgroup C adenoviruses and
group B coxsackieviruses. Proc.

[148220]Nat. Acad. Sci. 94: 3352–3356, 1997.

[148221]

[148222]

[148223]14022.

[148224]Margolis, R. L.; Abraham, M. R.; Gatchell, S. B.; Li, S.–H.;
Kidwai,

[148225]A. S.; Breschel, T. S.; Stine, O. C.; Callahan, C.; McInnis, M.
G.;

[148226]Ross, C. A.: cDNAs with long CAG trinucleotide repeats
from human

[148227]brain. Hum. Genet. 100: 114–122, 1997.

[148228]

[148229]

[148230]14023.

[148231]Dale, M.; Hammond, D. W.; Cox, A.; Nicklin, M. J. H.: The
human

[148232]gene encoding the interleukin-1 receptor accessory pro-
tein (IL1RAP)

[148233]maps to chromosome 3q28 by fluorescence in situ hy-
bridization and

[148234]radiation hybrid mapping. *Genomics* 47: 325-326, 1998.

[148235]

[148236]

[148237]14024.

[148238]Huang, J.; Gao, X.; Li, S.; Cao, Z.: Recruitment of IRAK to
the

[148239]interleukin 1 receptor complex requires interleukin 1 re-
ceptor accessory

[148240]protein. *Proc. Nat. Acad. Sci.* 94: 12829-12832, 1997.

[148241]

[148242]

[148243]14025.

[148244]Wesche, H.; Korherr, C.; Kracht, M.; Falk, W.; Resch, K.;
Martin,

[148245]M. U.: The interleukin-1 receptor accessory protein
(IL-1RAcP) is

[148246]essential for IL-1-induced activation of interleukin-1 re-
ceptor-associated

[148247]kinase (IRAK) and stress-activated protein kinases (SAP ki-
nases). J.

[148248]Biol. Chem. 272: 7727-7731, 1997.

[148249]

[148250]

[148251]14026.

[148252]Saha, P.; Chen, J.; Thome, K. C.; Lawlis, S. J.; Hou, Z.-H.;
Hendricks,

[148253]M.; Parvin, J. D.; Dutta, A.: Human CDC6/Cdc18 associates
with Orc1

[148254]and cyclin-cdk and is selectively eliminated from the nu-
cleus at the

[148255]onset of S phase. Molec. Cell. Biol. 18: 2758-2767, 1998.

[148256]

[148257]

[148258]14027.

[148259]Williams, R. S.; Shohet, R. V.; Stillman, B.: A human protein
[148260]related to yeast Cdc6p. Proc. Nat. Acad. Sci. 94: 142–147,
1997.

[148261]

[148262]

[148263]14028.

[148264]Yan, Z.; DeGregori, J.; Shohet, R.; Leone, G.; Stillman, B.;
Nevins,

[148265]J. R.; Williams, R. S.: Cdc6 is regulated by E2F and is es-
sential

[148266]for DNA replication in mammalian cells. Proc. Nat. Acad.
Sci. 95:

[148267]3603–3608, 1998.

[148268]

[148269]

[148270]14029.

[148271]Brais, B.; Bouchard, J.–P.; Xie, Y.–G.; Rochefort, D. L.;

Chretien,

[148272]N.; Tome, F. M. S.; Lafreniere, R. G.; Rommens, J. M.;

Uyama, E.;

[148273]Nohira, O.; Blumen, S.; Korczyn, A. D.; Heutink, P.; Mathieu, J.;

[148274]Duranceau, A.; Codere, F.; Fardeau, M.; Rouleau, G. A.:

Short GCG

[148275]expansions in the PABP2 gene cause oculopharyngeal muscular dystrophy. Nature

[148276]Genet. 18: 164–167, 1998. Note: Erratum: Nature Genet. 19: 404 only,

[148277]1998.

[148278]

[148279]

[148280]14030.

[148281]Calado, A.; Tome, F. M. S.; Brais, B.; Rouleau, G. A.; Kuhn, U.;

[148282]Wahle, E.; Carmo-Fonseca, M.: Nuclear inclusions in oculopharyngeal

[148283]muscular dystrophy consist of poly(A) binding protein 2 aggregates

[148284]which sequester poly(A) RNA. Hum. Molec. Genet. 9:

2321–2328, 2000.

[148285]

[148286]

[148287]14031.

[148288]Fan, X.; Dion, P.; Laganier, J.; Brais, B.; Rouleau, G. A.:

Oligomerization

[148289]of polyalanine expanded PABPN1 facilitates nuclear protein aggregation

[148290]that is associated with cell death. Hum. Molec. Genet. 10:

2341–2351,

[148291]2001.

[148292]

[148293]

[148294]14032.

[148295]Ikeda, S.; Shiva, N.; Ikeda, A.; Smith, R. S.; Nusinowitz, S.;

[148296]Yan, G.; Lin, T. R.; Chu, S.; Heckenlively, J. R.; North, M. A.;

Naggert,

[148297]J. K.; Nishina, P. M.; Duyao, M. P.: Retinal degeneration but not

[148298]obesity is observed in null mutants of the tubby-like pro-

tein 1 gene. Hum.

[148299]Molec. Genet. 9: 155–163, 2000.

[148300]

[148301]

[148302]14033.

[148303]North, M. A.; Naggert, J. K.; Yan, Y.; Noben–Trauth, K.;
Nishina,

[148304]P. M.: Molecular characterization of TUB, TULP1, and
TULP2, members

[148305]of the novel tubby gene family and their possible relation
to ocular

[148306]diseases. Proc. Nat. Acad. Sci. 94: 3128–3133, 1997.

[148307]

[148308]

[148309]14034.

[148310]Paloma, E.; Hjelmqvist, L.; Bayes, M.; Garcia–Sandoval, B.;
Ayuso,

[148311]C.; Balcells, S.; Gonzalez–Duarte, R.: Novel mutations in
the TULP1

[148312]gene causing autosomal recessive retinitis pigmentosa.

Invest. Ophthal.

[148313]Vis. Sci. 41: 656–659, 2000.

[148314]

[148315]

[148316]14035.

[148317]Aoki, N.; Ishii, T.; Ohira, S.; Yamaguchi, Y.; Negi, M.;
Adachi,

[148318]T.; Nakamura, R.; Matsuda, T.: Stage specific expression
of milk

[148319]fat globule membrane glycoproteins in mouse mammary
gland: comparison

[148320]of MFG–E8, butyrophilin, and CD36 with a major milk pro-
tein, beta–casein. Biochim.

[148321]Biophys. Acta 1334: 182–190, 1997.

[148322]

[148323]

[148324]14036.

[148325]Collins, C.; Nehlin, J. O.; Stubbs, J. D.; Kowbel, D.; Kuo,
W.–L.;

[148326]Parry, G.: Mapping of a newly discovered human gene ho–

mologous to

[148327]the apoptosis associated–murine mammary protein, MFG–
E8, to chromosome

[148328]15q25. Genomics 39: 117–118, 1997.

[148329]

[148330]

[148331]14037.

[148332]Haggqvist, B.; Naslund, J.; Sletten, K.; Westermarck, G. T.;
Mucchiano,

[148333]G.; Tjernberg, L. O.; Nordstedt, C.; Engstrom, U.; Wester–
mark, P.

[148334]: Medin: an integral fragment of aortic smooth muscle
cell–produced

[148335]lactadherin forms the most common human amyloid.
Proc. Nat. Acad.

[148336]Sci. 96: 8669–8674, 1999.

[148337]

[148338]

[148339]14038.

[148340]Hanayama, R.; Tanaka, M.; Miwa, K.; Shinohara, A.; Iwa–

matsu, A.;

[148341]Nagata, S.: Identification of a factor that links apoptotic cells

[148342]to phagocytes. Nature 417: 182–187, 2002.

[148343]

[148344]

[148345]14039.

[148346]Larocca, D.; Peterson, J. A.; Urrea, R.; Kuniyoshi, J.; Bis-train,

[148347]A. M.; Ceriani, R. L.: A M(r) 46,000 human milk fat globule protein

[148348]that is highly expressed in human breast tumors contains factor VIII-like

[148349]domains. Cancer Res. 51: 4994–4998, 1991.

[148350]

[148351]

[148352]14040.

[148353]Stubbs, J. D.; Lekutis, C.; Singer, K. L.; Bui, A.; Yuzuki, D.;

[148354]Srinivasan, U.; Parry, G.: cDNA cloning of a mouse mammary epithelial

[148355]cell surface protein reveals the existence of epidermal growth factor-like

[148356]domains linked to factor VIII-like sequences. Proc. Nat. Acad. Sci. 87:

[148357]8417–8421, 1990.

[148358]

[148359]

[148360]14041.

[148361]Zhang, J.; Kuehl, P.; Green, E. D.; Touchman, J. W.; Watkins, P.

[148362]B.; Daly, A.; Hall, S. D.; Maurel, P.; Relling, M.; Brimer, C.; Yasuda,

[148363]K.; Wrighton, S. A.; Hancock, M.; Kim, R. B.; Strom, S.; Thummel,

[148364]K.; Russell, C. G.; Hudson, J. R., Jr.; Schuetz, E. G.; Boguski, M.

[148365]S.: The human pregnane X receptor: genomic structure and identification

[148366]and functional characterization of natural allelic variants. Pharmacogenetics 11:

[148367]555–572, 2001.

[148368]

[148369]

[148370]14042.

[148371]Su, L.-K.; Qi, Y.: Characterization of human MAPRE genes
and their

[148372]proteins. Genomics 71: 143–149, 2001.

[148373]

[148374]

[148375]14043.

[148376]Arai, T.; Akiyama, Y.; Okabe, S.; Ando, M.; Endo, M.;
Yuasa, Y.

[148377]: Genomic structure of the human Smad3 gene and its in-
frequent alterations

[148378]in colorectal cancers. Cancer Lett. 122: 157–163, 1998.

[148379]

[148380]

[148381]14044.

[148382]Zhang, Y.; Feng, X.-H.; Wu, R.-Y.; Derynck, R.: Receptor-
associated

[148383]Mad homologues synergize as effectors of the TGF-beta response. Nature 383:

[148384]168-172, 1996.

[148385]

[148386]

[148387]14045.

[148388]Zhu, Y.; Richardson, J. A.; Parada, L. F.; Graff, J. M.: Smad3

[148389]mutant mice develop metastatic colorectal cancer. Cell 94:
703-714,

[148390]1998.

[148391]

[148392]

[148393]14046.

[148394]Bruno, E.; Horrigan, S. K.; Van Den Berg, D.; Rozler, E.; Fitting,

[148395]P. R.; Moss, S. T.; Westbrook, C.; Hoffman, R.: The Smad5 gene is

[148396]involved in the intracellular signaling pathways that mediate the

[148397]inhibitory effects of transforming growth factor-beta on

human hematopoiesis. Blood 91:

[148398]1917–1923, 1998.

[148399]

[148400]

[148401]14047.

[148402]Gemma, A.; Hagiwara, K.; Vincent, F.; Ke, Y.; Hancock, A.
R.; Nagashima,

[148403]M.; Bennett, W. P.; Harris, C. C.: hSmad5 gene, a human
hSmad family

[148404]member: its full length cDNA, genomic structure, pro-
moter region and

[148405]mutation analysis in human tumors. Oncogene 16:
951–956, 1998.

[148406]

[148407]

[148408]14048.

[148409]Cole, A. M.; Kim, Y.-H.; Tahk, S.; Hong, T.; Weis, P.; War-
ing,

[148410]A. J.; Ganz, T.: Calcitermin, a novel antimicrobial peptide
isolated

[148411]from human airway secretions. FEBS Lett. 504: 5–10,

2001.

[148412]

[148413]

[148414]14049.

[148415]Guignard, F.; Mauel, J.; Markert, M.: Identification and
characterization

[148416]of a novel human neutrophil protein related to the S100
family. Biochem.

[148417]J. 309: 395–401, 1995.

[148418]

[148419]

[148420]14050.

[148421]Ilg, E. C.; Troxler, H.; Burgisser, D. M.; Kuster, T.; Markert,

[148422]M.; Guignard, F.; Hunziker, P.; Birchler, N.; Heizmann, C.
W.: Amino

[148423]acid sequence determination of human S100A12 (P6, Cal-
granulin C, CGRP,

[148424]CAAF1) by tandem mass spectrometry. Biochem. Biophys.
Res. Commun. 225:

[148425]146–150, 1996.

[148426]

[148427]

[148428]14051.

[148429]Marti, T.; Erttmann, K. D.; Gallin, M. Y.: Host-parasite interaction

[148430]in human onchocerciasis: identification and sequence analysis of a

[148431]novel human calgranulin. Biochem. Biophys. Res. Commun. 221: 454-458,

[148432]1996.

[148433]

[148434]

[148435]14052.

[148436]Wicki, R.; Marenholz, I.; Mischke, D.; Schafer, B. W.; Heizmann,

[148437]C. W.: Characterization of the human S100A12 (calgranulin C, p6,

[148438]CAAF1, CGRP) gene, a new member of the S100 gene cluster on chromosome

[148439]1q21. Cell Calcium 20: 459-464, 1996.

[148440]

[148441]

[148442]14053.

[148443]Tan, K. B.; Harrop, J.; Reddy, M.; Young, P.; Terrett, J.;
Emery,

[148444]J.; Moore, G.; Truneh, A.: Characterization of a novel TNF-
like ligand

[148445]and recently described TNF ligand and TNF receptor su-
perfamily genes

[148446]and their constitutive and inducible expression in
hematopoietic and

[148447]non-hematopoietic cells. Gene 204: 35-46, 1997.

[148448]

[148449]

[148450]14054.

[148451]Gruter, P.; Tabernero, C.; von Kobbe, C.; Schmitt, C.;
Saavedra,

[148452]C.; Bachi, A.; Wilm, M.; Felber, B. K.; Izaurralde, E.: TAP,
the

[148453]human homolog of Mex67p, mediates CTE-dependent
RNA export from the

[148454]nucleus. Molec. Cell 1: 649–659, 1998.

[148455]

[148456]

[148457]14055.

[148458]Segref, A.; Sharma, K.; Doye, V.; Hellwig, A.; Huber, J.;
Luhrmann,

[148459]R.; Hurt, E.: Mex67p, a novel factor for nuclear mRNA ex-
port, binds

[148460]to both poly(A)+ RNA and nuclear pores. EMBO J. 16:
3256–3271, 1997.

[148461]

[148462]

[148463]14056.

[148464]Yoon, D.-W.; Lee, H.; Seol, W.; DeMaria, M.; Rosenzweig,
M.; Jung,

[148465]J. U.: Tap: a novel cellular protein that interacts with Tip of
herpesvirus

[148466]saimiri and induces lymphocyte aggregation. Immunity 6:
571–582,

[148467]1997.

[148468]

[148469]

[148470]14057.

[148471]Bonini, J. A.; Martin, S. K.; Dralyuk, F.; Roe, M. W.; Philipson,

[148472]L. H.; Steiner, D. F.: Cloning, expression, and chromosomal mapping

[148473]of a novel human CC-chemokine receptor (CCR10) that displays high-affinity

[148474]binding for MCP-1 and MCP-3. DNA Cell Biol. 16: 1249-1256, 1997.

[148475]

[148476]

[148477]14058.

[148478]Nibbs, R. J. B.; Wylie, S. M.; Yang, J.; Landau, N. R.; Graham,

[148479]G. J.: Cloning and characterization of a novel promiscuous human

[148480]beta-chemokine receptor D6. J. Biol. Chem. 272: 32078-32083, 1997.

[148481]

[148482]

[148483]14059.

[148484]Zieger, B.; Tran, H.; Hainmann, I.; Wunderle, D.; Zgaga-

Griesz,

[148485]A.; Blaser, S.; Ware, J.: Characterization and expression
analysis

[148486]of two human septin genes, PNUTL1 and PNUTL2. Gene
261: 197–203,

[148487]2000.

[148488]

[148489]

[148490]14060.

[148491]Brandt, S.; Jentsch, T. J.: ClC–6 and ClC–7 are two novel
broadly

[148492]expressed members of the CLC chloride channel family.
FEBS Lett. 377:

[148493]15–20, 1995.

[148494]

[148495]

[148496]14061.

[148497] Eggermont, J.; Buyse, G.; Voets, T.; Tytgat, J.; De Smedt, H.;

[148498] Droogmans, G.: Alternative splicing of ClC-6 (a member of the ClC

[148499] chloride-channel family) transcripts generates three truncated isoforms

[148500] one of which, ClC-6c, is kidney-specific. *Biochem. J.* 325: 269-276,

[148501] 1997.

[148502]

[148503]

[148504] 14062.

[148505] Cleiren, E.; Benichou, O.; Van Hul, E.; Gram, J.; Bollerslav, J.;

[148506] Singer, F. R.; Beaverson, K.; Aledo, A.; Whyte, M. P.; Yoneyama, T.;

[148507] deVernejou, M.-C.; Van Hul, W.: Albers-Schonberg disease (autosomal

[148508] dominant osteopetrosis, type II) results from mutations in the CLCN7

[148509] chloride channel gene. *Hum. Molec. Genet.* 10: 2861-2867, 2001.

[148510]

[148511]

[148512]14063.

[148513]Hedblom, E.; Kirkness, E. F.: A novel class of GABA-A receptor

[148514]subunit in tissues of the reproductive system. J. Biol. Chem. 272:

[148515]15346–15350, 1997.

[148516]

[148517]

[148518]14064.

[148519]Caloca, M. J.; Garcia-Bermejo, M. L.; Blumberg, P. M.; Lewin, N.

[148520]E.; Kremmer, E.; Mischak, H.; Wang, S.; Nacro, K.; Bienfait, B.; Marquez,

[148521]V. E.; Kazanietz, M. G.: Beta-2-chimaerin is a novel target for diacylglycerol:

[148522]binding properties and changes in subcellular localization mediated

[148523]by ligand binding to its C1 domain. Proc. Nat. Acad. Sci. 96: 11854–11859,

[148524]1999.

[148525]

[148526]

[148527]14065.

[148528]Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Cerebellar beta-
2-chimaerin,

[148529]a GTPase-activating protein for p21 Ras-related Rac is
specifically

[148530]expressed in granule cells and has a unique N-terminal
SH2 domain J.

[148531]Biol. Chem. 269: 12888-12892, 1994.

[148532]

[148533]

[148534]14066.

[148535]Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Germ cell beta-
chimaerin,

[148536]a new GTPase-activating protein for p21rac, is specifically
expressed

[148537]during the acrosomal assembly stage in rat testis. J. Biol.
Chem. 268:

[148538]3813–3816, 1993.

[148539]

[148540]

[148541]14067.

[148542]Yuan, S.; Miller, D. W.; Barnett, G. H.; Hahn, J. F.; Williams,

[148543]B. R.: Identification and characterization of human beta
2-chimaerin:

[148544]association with malignant transformation in astrocytoma.
Cancer

[148545]Res. 55: 3456–3461, 1995.

[148546]

[148547]

[148548]14068.

[148549]Chung, E.; Hanukoglu, A.; Rees, M.; Thompson, R.; Dillon,
M.; Hanukoglu,

[148550]I.; Bistrizter, T.; Kuhnle, U.; Seckl, J.; Gardiner, R. M.: Ex-
clusion

[148551]of the locus for autosomal recessive pseudohypoaldos-
teronism type

[148552]1 from the mineralocorticoid receptor gene region on hu-

man chromosome

[148553]4q by linkage analysis. J. Clin. Endocr. Metab. 80:
3341–3345, 1995.

[148554]

[148555]

[148556]14069.

[148557]Schwienbacher, C.; Sabbioni, S.; Campi, M.; Veronese, A.;
Bernardi,

[148558]G.; Menegatti, A.; Hatada, I.; Mukai, T.; Ohashi, H.; Bar-
banti–Brodano,

[148559]G.; Croce, C. M.; Negrini, M.: Transcriptional map of
170–kb region

[148560]at chromosome 11p15.5: identification and mutational
analysis of the

[148561]BWR1A gene reveals the presence of mutations in tumor
samples. Proc.

[148562]Nat. Acad. Sci. 95: 3873–3878, 1998.

[148563]

[148564]

[148565]14070.

[148566]Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hiro–

sawa, M.;

[148567]Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara,
O.: Prediction

[148568]of the coding sequences of unidentified human genes.

XIII. The complete

[148569]sequences of 100 new cDNA clones from brain which code
for large proteins

[148570]in vitro. DNA Res. 6: 63–70, 1999.

[148571]

[148572]

[148573]14071.

[148574]Philibert, R. A.; Winfield, S. L.; Sandhu, H. K.; Martin, B. M.;

[148575]Ginns, E. I.: The structure and expression of the human
neuroligin–3

[148576]gene. Gene 246: 303–310, 2000.

[148577]

[148578]

[148579]14072.

[148580]Arico, M.; Imashuku, S.; Clementi, R.; Hibi, S.; Teramura,
T.;

[148581]Danesino, C.; Haber, D. A.; Nichols, K. E.: Hemophagocytic lymphohistiocytosis

[148582]due to germline mutations in SH2D1A, the X-linked lymphoproliferative

[148583]disease gene. Blood 97: 1131–1133, 2001.

[148584]

[148585]

[148586]14073.

[148587]Arkwright, P. D.; Makin, G.; Will, A. M.; Ayres, M.; Gokhale, D.

[148588]A.; Fergusson, W. D.; Taylor, G. M.: X linked lymphoproliferative

[148589]disease in a United Kingdom family. Arch. Dis. Child. 79: 52–55,

[148590]1998.

[148591]

[148592]

[148593]14074.

[148594]Bar, R. S.; DeLor, C. J.; Clausen, K. P.; Hurtubise, P.; Henle,

[148595]W.; Hewetson, J. F.: Fatal infectious mononucleosis in a

family. New

[148596]Eng. J. Med. 290: 363–367, 1974.

[148597]

[148598]

[148599]14075.

[148600]Benoit, L.; Wang, X.; Pabst, H. F.; Dutz, J.; Tan, R.: Cutting

[148601]edge: defective NK cell activation in X-linked lymphoproliferative

[148602]disease. J. Immun. 165: 3549–3553, 2000.

[148603]

[148604]

[148605]14076.

[148606]Brandau, O.; Schuster, V.; Weiss, M.; Hellebrand, H.; Fink, F.

[148607]M.; Kreczy, A.; Friedrich, W.; Strahm, B.; Niemeyer, C.; Belohradsky,

[148608]B. H.; Meindl, A.: Epstein–Barr virus–negative boys with non-Hodgkin

[148609]lymphoma are mutated in the SH2D1A gene, as are patients with X-linked

[148610]lymphoproliferative disease (XLP). Hum. Molec. Genet. 8:
2407–2413,
[148611]1999.

[148612]

[148613]

[148614]14077.

[148615]Coffey, A. J.; Brooksbank, R. A.; Brandau, O.; Oohashi, T.;
Howell,

[148616]G. R.; Bye, J. M.; Cahn, A. P.; Durham, J.; Heath, P.; Wray,
P.; Pavitt,

[148617]R.; Wilkinson, J.; and 31 others: Host response to EBV in-
fection

[148618]in X-linked lymphoproliferative disease results from mu-
tations in

[148619]an SH2-domain encoding gene. Nature Genet. 20:
129–135, 1998.

[148620]

[148621]

[148622]14078.

[148623]Czar, M. J.; Kersh, E. N.; Mijares, L. A.; Lanier, G.; Lewis, J.;

[148624]Yap, G.; Chen, A.; Sher, A.; Duckett, C. S.; Ahmed, R.;
Schwartzberg,

[148625]P. L.: Altered lymphocyte responses and cytokine produc-
tion in mice

[148626]deficient in the X-linked lymphoproliferative disease gene
SH2D1A/DSHP/SAP. Proc.

[148627]Nat. Acad. Sci. 98: 7449–7454, 2001.

[148628]

[148629]

[148630]14079.

[148631]Dutz, J. P.; Benoit, L.; Wang, X.; Demetrick, D. J.; Junker,
A.;

[148632]de Sa, D.; Tan, R.: Lymphocytic vasculitis in X-linked lym-
phoproliferative

[148633]disease. Blood 97: 95–100, 2001.

[148634]

[148635]

[148636]14080.

[148637]Grierson, H. L.; Skare, J.; Church, J.; Silberman, T.; Davis, J.

[148638]R.; Kobrinsky, N.; McGregor, R.; Israels, S.; McCarty, J.;

Andrews,

[148639]L. G.; Blecha, T.; Erdman, S.; Obringer, A.; Scharnhorst, D.;

Purtilo,

[148640]D. T.: Evaluation of families wherein a single male mani-

festes a phenotype

[148641]of X-linked lymphoproliferative disease (XLP). Am. J. Med.

Genet. 47:

[148642]458-463, 1993.

[148643]

[148644]

[148645]14081.

[148646]Hambleton, G.; Cottom, D. G.: Familial lymphoma. Proc.

Roy.

[148647]Soc. Med. 62: 1095 only, 1969.

[148648]

[148649]

[148650]14082.

[148651]Hamilton, J. K.; Paquin, L. A.; Sullivan, J. L.; Maurer, H. S.;

[148652]Cruzi, F. G.; Provisor, A. J.; Steuber, C. P.; Hawkins, E.;

Yawn,

[148653]D.; Cornet, J.; Clausen, K.; Finkelstein, G. Z.; Landing, B.;
Grunnet,

[148654]M.; Purtilo, D. T.: X-linked lymphoproliferative syndrome
registry

[148655]report. J. Pediat. 96: 669–673, 1980.

[148656]

[148657]

[148658]14083.

[148659]Harris, A.; Docherty, Z.: X-linked lymphoproliferative dis-
ease:

[148660]a karyotype analysis. Cytogenet. Cell Genet. 47: 92–94,
1988.

[148661]

[148662]

[148663]14084.

[148664]Harris, A.; Lenoir, G. M.; Lankester, S. A.: X-linked lym-
phoproliferative

[148665]disease: linkage studies using DNA probes. Clin. Genet.
33: 162–168,

[148666]1988.

[148667]

[148668]

[148669]14085.

[148670]Hayoz, D.; Lenoir, G. M.; Nicole, A.; Pugin, P.; Regamey, C.:

[148671]X-linked lymphoproliferative syndrome: identification of a large family

[148672]in Switzerland. Am. J. Med. 84: 529–534, 1988.

[148673]

[148674]

[148675]14086.

[148676]Klein, G.; Klein, E.: Sinking surveillance's flagship. Nature 395:

[148677]441–445, 1998.

[148678]

[148679]

[148680]14087.

[148681]Levine, P. H.; Kamaraju, L. S.; Connelly, R. R.; Berard, C. W.;

[148682]Dorfman, R. F.; Magrath, I.; Easton, J. M.: The American
Burkitt's

[148683]Lymphoma Registry: eight years' experience. Cancer 49:
1016–1022,

[148684]1982.

[148685]

[148686]

[148687]14088.

[148688]Loeffel, S.; Chang, C.–H.; Heyn, R.; Harada, S.; Lipscomb,
H.;

[148689]Sinangil, F.; Volsky, D. J.; McClain, K.; Ochs, H.; Purtilo, D.
T.

[148690]: Necrotizing lymphoid vasculitis in X–linked lymphopro-
liferative

[148691]syndrome. Arch. Path. Lab. Med. 109: 546–550, 1985.

[148692]

[148693]

[148694]14089.

[148695]Lyon, M. F.; Loutit, J. F.: X–linked factor in acquired im-
munodeficiency

[148696]syndrome?. (Letter) Lancet I: 768 only, 1983.

[148697]

[148698]

[148699]14090.

[148700]Mulley, J. C.; Turner, A. M.; Gedeon, A. K.; Berdoukas, V.
A.;

[148701]Huang, T. H. M.; Ledbetter, D. H.; Grierson, H.; Purtilo, D.
T.:

[148702]X-linked lymphoproliferative disease: prenatal detection
of an unaffected

[148703]histocompatible male. Clin. Genet. 42: 76–79, 1992.

[148704]

[148705]

[148706]14091.

[148707]Eva, A.; Aaronson, S. A.: Isolation of a new human onco-
gene from

[148708]a diffuse B-cell lymphoma. Nature 316: 273–275, 1985.

[148709]

[148710]

[148711]14092.

[148712]Anderson, S. R.; Warburg, M.: Norrie's disease. Arch. Ophthal. 66:

[148713]614–618, 1961.

[148714]

[148715]

[148716]14093.

[148717]Bergen, A. A. B.; Wapenaar, M. C.; Schuurman, E. J. M.;
Diergaarde,

[148718]P. J.; Lerach, H.; Monaco, A. P.; Bakker, E.; Bleeker–
Wagemakers,

[148719]E. M.; van Ommen, G. J. B.: Detection of a new submicro–
scopic Norrie

[148720]disease deletion interval with a novel DNA probe isolated
by differential

[148721]Alu PCR fingerprint cloning. Cytogenet. Cell Genet. 62:
231–235,

[148722]1993.

[148723]

[148724]

[148725]14094.

[148726]Berger, W.; Meindl, A.; van de Pol, T. J. R.; Cremers, F. P. M.;

[148727]Ropers, H. H.; Doerner, C.; Monaco, A.; Bergen, A. A. B.; Lebo, R.;

[148728]Warburg, M.; Zergollern, L.; Lorenz, B.; Gal, A.; Bleeker-Wagemakers,

[148729]E. M.; Meitinger, T.: Isolation of a candidate gene for Norrie disease

[148730]by positional cloning. Nature Genet. 1: 199–203, 1992.

[148731]

[148732]

[148733]14095.

[148734]Berger, W.; van de Pol, D.; Bachner, D.; Oerlemans, F.; Winkens,

[148735]H.; Hameister, H.; Wieringa, B.; Hendriks, W.; Ropers, H.–H.: An

[148736]animal model for Norrie disease (ND): gene targeting of the mouse

[148737]ND gene. Hum. Molec. Genet. 5: 51–59, 1996.

[148738]

[148739]

[148740]14096.

[148741]Eva, A.; Vecchio, G.; Rao, C. D.; Tronick, S. R.; Aaronson, S.

[148742]A.: The predicted DBL oncogene product defines a distinct class of

[148743]transforming proteins. Proc. Nat. Acad. Sci. 85: 2061–2065, 1988.

[148744]

[148745]

[148746]14097.

[148747]Galland, F.; Stefanova, M.; Lafage, M.; Birnbaum, D.: Localization

[148748]of the 5–prime end of the MCF2 oncogene to human chromosome 15q15–q23.

[148749]Cytogenet. Cell Genet. 60: 114–116, 1992.

[148750]

[148751]

[148752]14098.

[148753]Grant, S. G.; Mattei, M.–G.; Galland, F.; Stephenson, D. A.;

Keitz,

[148754]B. T.; Birnbaum, D.; Chapman, V. M.: Localization of the mouse Mcf-2

[148755](Dbl) protooncogene within a conserved linkage group on the mouse

[148756]X chromosome. Cytogenet. Cell Genet. 54: 175-181, 1990.

[148757]

[148758]

[148759]14099.

[148760]Nguyen, C.; Pontarotti, P.; Birnbaum, D.; Chimini, G.; Rey, J.

[148761]A.; Mattei, J.-F.; Jordan, B. R.: Large scale physical mapping in

[148762]the q27 region of the human X chromosome: the coagulation factor IX

[148763]gene and the mcf.2 transforming sequence are separated by at most

[148764]270 kilobase pairs and are surrounded by several 'HTF islands.' EMBO

[148765]J. 6: 3285-3289, 1987.

[148766]

[148767]

[148768]14100.

[148769]Nguyen, C.; Poustka, A.-M.; Djabali, M.; Roux, D.; Mattei, J.-F.;

[148770]Lehrach, H.; Jordan, B. R.: Large-scale mapping and chromosome jumping

[148771]in the q27 region of the human X chromosome. Genomics 5: 298-303,

[148772]1989.

[148773]

[148774]

[148775]14101.

[148776]Noguchi, T.; Mattei, M.-G.; Oberle, I.; Planche, J.; Imbert, J.;

[148777]Pelassy, C.; Birg, F.; Birnbaum, D.: Localization of the mcf.2 transforming

[148778]sequence to the X chromosome. EMBO J. 6: 1301-1307, 1987.

[148779]

[148780]

[148781]14102.

[148782]Ron, D.; Tronick, S. R.; Aaronson, S. A.; Eva, A.: Molecular cloning

[148783]and characterization of the human DBL proto-oncogene: evidence that

[148784]its overexpression is sufficient to transform NIH/3T3 cells.

EMBO

[148785]J. 7: 2465–2473, 1988.

[148786]

[148787]

[148788]14103.

[148789]Srivastava, S. K.; Wheelock, R. H. P.; Aaronson, S. A.; Eva, A.

[148790]: Identification of the protein encoded by the human diffuse B-cell

[148791]lymphoma (dbl) oncogene. Proc. Nat. Acad. Sci. 83: 8868–8872, 1986.

[148792]

[148793]

[148794]14104.

[148795]Tronick, S. R.; McBride, O. W.; Popescu, N. C.; Eva, A.:
Chromosomal

[148796]localization of DBL oncogene sequences. Genomics 5:
546–553, 1989.

[148797]

[148798]

[148799]14105.

[148800]Giovane, A.; Sobieszczuk, P.; Mignon, C.; Mattei, M.–G.;
Wasylyk,

[148801]B.: Locations of the ets subfamily members net, elk1, and
sap1 (ELK3,

[148802]ELK1, and ELK4) on three homologous regions of the
mouse and human

[148803]genomes. Genomics 29: 769–772, 1995.

[148804]

[148805]

[148806]14106.

[148807]Janz, M.; Lehmann, U.; Olde Weghuis, D.; de Leeuw, B.;

Geurts van

[148808]Kessel, A.; Gilgenkrantz, S.; Hipskind, R. A.; Nordheim, A.:

Refined

[148809]mapping of the human Ets-related gene Elk-1 to

Xp11.2-p11.4, distal

[148810]to the OATL1 region. Hum. Genet. 94: 442-444, 1994.

[148811]

[148812]

[148813]14107.

[148814]Rao, V. N.; Huebner, K.; Isobe, M.; ar-Rushdi, A.; Croce, C.

M.;

[148815]Reddy, E. S. P.: Elk, tissue-specific ets-related genes on

chromosomes

[148816]X and 14 near translocation breakpoints. Science 244:

66-70, 1989.

[148817]

[148818]

[148819]14108.

[148820]Tamai, Y.; Taketo, M.; Nozaki, M.; Seldin, M. F.: Mouse Elk

oncogene

[148821]maps to chromosome X and a novel Elk oncogene (Elk3)

maps to chromosome

[148822]10. Genomics 26: 414–416, 1995.

[148823]

[148824]

[148825]14109.

[148826]Yamauchi, T.; Toko, M.; Suga, M.; Hatakeyama, T.; Isobe, M.: Structural

[148827]organization of the human Elk1 gene and its processed pseudogene Elk2. DNA

[148828]Res. 6: 21–27, 1999.

[148829]

[148830]

[148831]14110.

[148832]Bermingham, J. R., Jr.; Scherer, S. S.; O'Connell, S.; Arroyo,

[148833]E.; Kalla, K. A.; Powell, F. L.; Rosenfeld, M. G.: Tst–
1/Oct–6/SCIP

[148834]regulates a unique step in peripheral myelination and is required

[148835]for normal respiration. Genes Dev. 10: 1751–1762, 1996.

[148836]

[148837]

[148838]14111.

[148839]Jaegle, M.; Mandemakers, W.; Broos, L.; Zwart, R.; Karis, A.; Visser,

[148840]P.; Grosveld, F.; Meijer, D.: The POU factor Oct-6 and Schwann cell

[148841]differentiation. Science 273: 507-510, 1996.

[148842]

[148843]

[148844]14112.

[148845]Sumiyama, K.; Washio-Watanabe, K.; Ono, T.; Yoshida, M. C.; Hayakawa,

[148846]T.; Ueda, S.: Human class III POU genes, POU3F1 and POU3F3, map to

[148847]chromosomes 1p34.1 and 3p14.1. Mammalian Genome 9: 180-181, 1998.

[148848]

[148849]

[148850]14113.

- [148851]Costache, M.; Apoil, P.-A.; Cailleau, A.; Elmgren, A.; Lar-
son,
[148852]G.; Henry, S.; Blancher, A.; Iordachescu, D.; Oriol, R.; Mol-
licone,
[148853]R.: Evolution of fucosyltransferase genes in vertebrates. J.
Biol.
[148854]Chem. 272: 29721-29728, 1997.
- [148855]
- [148856]
- [148857]14114.
- [148858]Yanagidani, S.; Uozumi, N.; Ihara, Y.; Miyoshi, E.; Yam-
aguchi,
[148859]N.; Taniguchi, N.: Purification and cDNA cloning of GDP-
L-Fuc:N-acetyl-beta-D-glucosaminide:alpha-1-6
[148860]fucosyltransferase (alpha-1-6 FucT) from human gastric
cancer MKN45
[148861]cells. J. Biochem. 121: 626-632, 1997.
- [148862]
- [148863]
- [148864]14115.

[148865]Bekri, S.; Adelaide, J.; Merscher, S.; Grosgeorge, J.; Caroli-Bosc,

[148866]F.; Perucca-Lostanlen, D.; Kelley, P. M.; Pebusque, M.-J.; Theillet,

[148867]C.; Birnbaum, D.; Gaudray, P.: Detailed map of a region commonly

[148868]amplified at 11q13-q14 in human breast carcinoma. Cytogenet. Cell

[148869]Genet. 79: 125-131, 1997.

[148870]

[148871]

[148872]14116.

[148873]Goodwin, R. G.; Alderson, M. R.; Smith, C. A.; Armitage, R. J.;

[148874]VandenBos, T.; Jerzy, R.; Tough, T. W.; Schoenborn, M. A.; Davis-Smith,

[148875]T.; Hennen, K.; Falk, B.; Cosman, D.; Baker, E.; Sutherland, G. R.;

[148876]Grabstein, K. H.; Farrah, T.; Giri, J. G.; Beckmann, M. P.: Molecular

[148877]and biological characterization of a ligand for CD27 defines a new

[148878]family of cytokines with homology to tumor necrosis factor. Cell 73:

[148879]447–456, 1993.

[148880]

[148881]

[148882]14117.

[148883]Wright, R. M.; Vaitaitis, G. M.; Wilson, C. M.; Repine, T. B.;

[148884]Terada, L. S.; Repine, J. E.: cDNA cloning, characterization, and

[148885]tissue-specific expression of human xanthine dehydrogenase/xanthine

[148886]oxidase. Proc. Nat. Acad. Sci. 90: 10690–10694, 1993.

[148887]

[148888]

[148889]14118.

[148890]Kawabata, M.; Saeki, K.: Multiple alternative transcripts of the

[148891]human homologue of the mouse TRAD/R51H3/RAD51D gene, a member of the

[148892]rec A/RAD51 gene family. Biochem. Biophys. Res. Com–

mun. 257: 156–162,

[148893]1999.

[148894]

[148895]

[148896]14119.

[148897]Pittman, D. L.; Weinberg, L. R.; Schimenti, J. C.: Identifica-
tion,

[148898]characterization, and genetic mapping of Rad51d, a new
mouse and human

[148899]RAD51/RecA-related gene. Genomics 49: 103–111, 1998.

[148900]

[148901]

[148902]14120.

[148903]Bell, B.; Scheer, E.; Tora, L.: Identification of
hTAFII80-delta

[148904]links apoptotic signaling pathways to transcription factor
TFIID function. Molec.

[148905]Cell 8: 591–600, 2001.

[148906]

[148907]

[148908]14121.

[148909]Hisatake, K.; Ohta, T.; Takada, R.; Guermah, M.; Horikoshi, M.;

[148910]Nakatani, Y.; Roeder, R. G.: Evolutionary conservation of human TATA-binding-polypeptide-associated

[148911]factors TAFII31 and TAFII80 and interactions of TAFII80 with other

[148912]TAFs and with general transcription factors. Proc. Nat. Acad. Sci. 92:

[148913]8195–8199, 1995.

[148914]

[148915]

[148916]14122.

[148917]Weinzierl, R. O. J.; Ruppert, S.; Dynlacht, B. D.; Tanese, N.;

[148918]Tjian, R.: Cloning and expression of Drosophila TAFII60 and human

[148919]TAFII70 reveal conserved interactions with other subunits of TFIID. EMBO

[148920]J. 12: 5303–5309, 1993.

[148921]

[148922]

[148923]14123.

[148924]Tsai, K. J.; Chen, S. K.; Ma, Y. L.; Hsu, W. L.; Lee, E. H. Y.

[148925]: *sgk*, a primary glucocorticoid-induced gene, facilitates
memory consolidation

[148926]of spatial learning in rats. *Proc. Nat. Acad. Sci.* 99:
3990–3995,

[148927]2002.

[148928]

[148929]

[148930]14124.

[148931]Waldegger, S.; Barth, P.; Raber, G.; Lang, F.: Cloning and
characterization

[148932]of a putative human serine/threonine protein kinase tran-
scriptionally

[148933]modified during anisotonic and isotonic alterations of cell
volume. *Proc.*

[148934]*Nat. Acad. Sci.* 94: 4440–4445, 1997.

[148935]

[148936]

[148937]14125.

[148938]Mackay, D. J. G.; Coupe, A.-M.; Shield, J. P. H.; Storr, J. N.

[148939]P.; Temple, I. K.; Robinson, D. O.: Relaxation of imprinted
expression

[148940]of ZAC and HYMAI in a patient with transient neonatal di-
abetes mellitus. Hum.

[148941]Genet. 110: 139–144, 2002.

[148942]

[148943]

[148944]14126.

[148945]Spengler, D.; Villalba, M.; Hoffmann, A.; Pantaloni, C.;
Houssami,

[148946]S.; Bockaert, J.; Journot, L.: Regulation of apoptosis and
cell cycle

[148947]arrest by Zac1, a novel zinc finger protein expressed in
the pituitary

[148948]gland and the brain. EMBO J. 16: 2814–2825, 1997.

[148949]

[148950]

[148951]14127.

[148952]Temple, I. K.; James, R. S.; Crolla, J. A.; Sitch, F. L.; Jacobs,

[148953]P. A.; Howell, W. M.; Betts, P.; Baum, J. D.; Shield, J. P. H.:

An

[148954]imprinted gene(s) for diabetes? (Letter) Nature Genet. 9:

110–112,

[148955]1995.

[148956]

[148957]

[148958]14128.

[148959]Varrault, A.; Ciani, E.; Apiou, F.; Bilanges, B.; Hoffmann,

A.;

[148960]Pantaloni, C.; Bockaert, J.; Spengler, D.; Journot, L.: hZAC

encodes

[148961]a zinc finger protein with antiproliferative properties and

maps to

[148962]a chromosomal region frequently lost in cancer. Proc. Nat.

Acad.

[148963]Sci. 95: 8835–8840, 1998.

[148964]

[148965]

[148966]14129.

[148967]Koch–Nolte, F.; Haag, F.; Braren, R.; Kuhl, M.; Hoovers, J.;

Balasubramanian,

[148968]S.; Bazan, F.; Thiele, H.–G.: Two novel human members of
an emerging

[148969]mammalian gene family related to mono–

ADP–ribosylating bacterial toxins. Genomics 39:

[148970]370–376, 1997.

[148971]

[148972]

[148973]14130.

[148974]Jiang, Y.–W.; Veschambre, P.; Erdjument–Bromage, H.;

Tempst, P.;

[148975]Conaway, J. W.; Conaway, R. C.; Kornberg, R. D.: Mam–
malian mediator

[148976]of transcriptional regulation and its possible role as an
end–point

[148977]of signal transduction pathways. Proc. Nat. Acad. Sci. 95:

8538–8543,

[148978]1998.

[148979]

[148980]

[148981]14131.

[148982]Adams, M. D.; Soares, M. B.; Kerlavage, A. R.; Fields, C.;
Venter,

[148983]J. C.: Rapid cDNA sequencing (expressed sequence tags)
from a directionally

[148984]cloned human infant brain cDNA library. Nature Genet. 4:
373–380,

[148985]1993.

[148986]

[148987]

[148988]14132.

[148989]Dai, K.–S.; Liew, C.–C.: A novel human striated muscle
RING zinc

[148990]finger protein, SMRZ, interacts with SMT3b via its RING
domain. J.

[148991]Biol. Chem. 276: 23992–23999, 2001.

[148992]

[148993]

[148994]14133.

[148995]Abdollahi, A.; Godwin, A. K.; Miller, P. D.; Getts, L. A.;
Schultz,

[148996]D. C.; Taguchi, T.; Testa, J. R.; Hamilton, T. C.: Identifica-
tion

[148997]of a gene containing zinc-finger motifs based on lost ex-
pression in

[148998]malignantly transformed rat ovarian surface epithelial
cells. Cancer

[148999]Res. 57: 2029-2034, 1997.

[149000]

[149001]

[149002]14134.

[149003]Abdollahi, A.; Roberts, D.; Godwin, A. K.; Schultz, D. C.;
Sonoda,

[149004]G.; Testa, J. R.; Hamilton, T. C.: Identification of a zinc-
finger

[149005]gene at 6q25: a chromosomal region implicated in devel-
opment of many

[149006]solid tumors. Oncogene 14: 1973-1979, 1997.

[149007]

[149008]

[149009]14135.

[149010]Kamiya, M.; Judson, H.; Okazaki, Y.; Kusakabe, M.; Muramatsu, M.;

[149011]Takada, S.; Takagi, N.; Arima, T.; Wake, N.; Kamimura, K.; Satomura,

[149012]K.; Hermann, R.; Bonthron, D. T.; Hayashizaki, Y.: The cell cycle

[149013]control gene ZAC/PLAGL1 is imprinted: a strong candidate gene for

[149014]transient neonatal diabetes. Hum. Molec. Genet. 9: 453–460, 2000.

[149015]

[149016]

[149017]14136.

[149018]Ide, H.; Katoh, M.; Sasaki, H.; Yoshida, T.; Aoki, K.; Nawa, Y.;

[149019]Osada, Y.; Sugimura, T.; Terada, M.: Cloning of human bone morphogenetic

[149020]protein type 1B receptor (BMPR–1B) and its expression in prostate

[149021]cancer in comparison with other BMPRs. Oncogene 14:
1377–1382, 1997.

[149022]

[149023]

[149024]14137.

[149025]ten Dijke, P.; Yamashita, H.; Ichijo, H.; Franzen, P.; Laiho,
M.;

[149026]Miyazono, K.; Heldin, C.–H.: Characterization of type I re-
ceptors

[149027]for transforming growth factor–beta and activin. Science
264: 101–104,

[149028]1994.

[149029]

[149030]

[149031]14138.

[149032]Blixt, A.; Mahlapuu, M.; Bjursell, C.; Darnfors, C.; Johan-
nesson,

[149033]T.; Enerback, S.; Carlsson, P.: The two–exon gene of the
human forkhead

[149034]transcription factor FREAC–2 (FKHL6) is located at 6p25.3.
Genomics 53:

[149035]387–390, 1998.

[149036]

[149037]

[149038]14139.

[149039]Best, J. L.; Presky, D. H.; Swerlick, R. A.; Burns, D. K.; Chu,

[149040]W.: Cloning of a full-length cDNA sequence encoding a
cdc2-related

[149041]protein kinase from human endothelial cells. Biochem.
Biophys. Res.

[149042]Commun. 208: 562–568, 1995.

[149043]

[149044]

[149045]14140.

[149046]Garriga, J.; Mayol, X.; Grana, X.: The CDC2-related kinase
PITALRE

[149047]is the catalytic subunit of active multimeric protein com-
plexes. Biochem.

[149048]J. 319: 293–298, 1996.

[149049]

[149050]

[149051]14141.

[149052]Grana, X.; De Luca, A.; Sang, N.; Fu, Y.; Claudio, P. P.;

Rosenblatt,

[149053]J.; Morgan, D. O.; Giordano, A.: PITALRE, a nuclear

CDC2-related

[149054]protein kinase that phosphorylates the retinoblastoma
protein in vitro. Proc.

[149055]Nat. Acad. Sci. 91: 3834–3838, 1994.

[149056]

[149057]

[149058]14142.

[149059]Liu, H.; Rice, A. P.: Genomic organization and characteri-
zation

[149060]of promoter function of the human CDK9 gene. Gene 252:
51–59, 2000.

[149061]

[149062]

[149063]14143.

[149064]Triepeles, R.; van den Heuvel, L.; Loeffen, J.; Smeets, R.;

Trijbels,

[149065]F.; Smeitink, J.: The nuclear-encoded human

NADH:ubiquinone oxidoreductase

[149066]NDUFA8 subunit: cDNA cloning, chromosomal localization, tissue distribution,

[149067]and mutation detection in complex-I-deficient patients.

Hum. Genet. 103:

[149068]557–563, 1998.

[149069]

[149070]

[149071]14144.

[149072]Bai, C.; Connolly, B.; Metzker, M. L.; Hilliard, C. A.; Liu, X.;

[149073]Sandig, V.; Soderman, A.; Galloway, S. M.; Liu, Q.; Austin, C. P.;

[149074]Caskey, C. T.: Overexpression of M68/DCR3 in human gastrointestinal

[149075]tract tumors independent of gene amplification and its location in

[149076]a four-gene cluster. Proc. Nat. Acad. Sci. 97: 1230–1235, 2000.

[149077]

[149078]

[149079]14145.

[149080]Pitti, R. M.; Marsters, S. A.; Lawrence, D. A.; Roy, M.;
Kischkel,

[149081]F. C.; Dowd, P.; Huang, A.; Donahue, C. J.; Sherwood, S.
W.; Baldwin,

[149082]D. T.; Godowski, P. J.; Wood, W. I.; Gurney, A. L.; Hillan, K.
J.;

[149083]Cohen, R. L.; Goddard, A. D.; Botstein, D.; Ashkenazi, A.:
Genomic

[149084]amplification of a decoy receptor for Fas ligand in lung
and colon

[149085]cancer. Nature 396: 699–703, 1998.

[149086]

[149087]

[149088]14146.

[149089]Ringstad, N.; Nemoto, Y.; De Camilli, P.: The
SH3p4/Sh3p8/SH3p13

[149090]protein family: binding partners for synaptojanin and dy-
namin via

[149091]a Grb2-like Src homology 3 domain. Proc. Nat. Acad. Sci.

94: 8569–8574,

[149092]1997.

[149093]

[149094]

[149095]14147.

[149096]Sittler, A.; Walter, S.; Wedemeyer, N.; Hasenbank, R.;

Scherzinger,

[149097]E.; Eickhoff, H.; Bates, G. P.; Lehrach, H.; Wanker, E. E.:

SH3GL3

[149098]associates with the huntington exon 1 protein and promotes the formation

[149099]of polyglu–containing protein aggregates. Molec. Cell 2:

427–436,

[149100]1998.

[149101]

[149102]

[149103]14148.

[149104]Sparks, A. B.; Hoffman, N. G.; McConnell, S. J.; Fowlkes, D.

M.;

[149105]Kay, B. K.: Cloning of ligand targets: systematic isolation

of SH3

[149106]domain-containing proteins. Nature Biotech. 14:
741–744, 1996.

[149107]

[149108]

[149109]14149.

[149110]Chinnaiyan, A. M.; O'Rourke, K.; Yu, G.–L.; Lyons, R. H.;
Garg,

[149111]M.; Duan, D. R.; Xing, L.; Gentz, R.; Ni, J.; Dixit, V. M.: Sig-
nal

[149112]transduction by DR3, a death domain-containing receptor
related to

[149113]TNFR–1 and CD95. Science 274: 990–992, 1996.

[149114]

[149115]

[149116]14150.

[149117]Fracchiolla, N. S.; Colombo, G.; Finelli, P.; Maiolo, A. T.;
Neri,

[149118]A.: EHT, a new member of the MTG8/ETO gene family,
maps on 20q11

[149119]region and is deleted in acute myeloid leukemias. (Letter)

Blood 92:

[149120]3481–3484, 1998.

[149121]

[149122]

[149123]14151.

[149124]Kitabayashi, I.; Ida, K.; Morohoshi, F.; Yokoyama, A.; Mit-
suhashi,

[149125]N.; Shimizu, K.; Nomura, N.; Hayashi, Y.; Ohki, M.: The
AML1–MTG8

[149126]leukemic fusion protein forms a complex with a novel
member of the

[149127]MTG8(ETO/CDR) family, MTGR1. Molec. Cell. Biol. 18:
846–858, 1998.

[149128]

[149129]

[149130]14152.

[149131]Levy, I.; Wu, Y.–Q.; Roeckel, N.; Bulle, F.; Pawlak, A.;
Siegrist,

[149132]S.; Mattei, M. G.; Guellaen, G.: Human testis specifically
expresses

[149133]a homologue of the rodent T lymphocytes RT6 mRNA.

FEBS Lett. 382:

[149134]276–280, 1996.

[149135]

[149136]

[149137]14153.

[149138]Nelson, H.; Mandiyan, S.; Noumi, T.; Moriyama, Y.; Miedel, M. C.;

[149139]Nelson, N.: Molecular cloning of cDNA encoding the C subunit of H(+)-ATPase

[149140]from bovine chromaffin granules. J. Biol. Chem. 265: 20390–20393,

[149141]1990.

[149142]

[149143]

[149144]14154.

[149145]van Hille, B.; Vanek, M.; Richener, H.; Green, J. R.; Bilbe, G.

[149146]: Cloning and tissue distribution of subunits C, D, and E of the human

[149147]vacuolar H(+)-ATPase. Biochem. Biophys. Res. Commun. 197: 15–21,

[149148]1993.

[149149]

[149150]

[149151]14155.

[149152]Fujiwara, T.; Suzuki, M.; Tanigami, A.; Ikenoue, T.; Omata, M.;

[149153]Chiba, T.; Tanaka, K.: The BTRC gene, encoding a human F-box/WD40-repeat

[149154]protein, maps to chromosome 10q24-q25. Genomics 58: 104-105, 1999.

[149155]

[149156]

[149157]14156.

[149158]Margottin, F.; Bour, S. P.; Durand, H.; Selig, L.; Benichou, S.;

[149159]Richard, V.; Thomas, D.; Strebel, K.; Benarous, R.: A novel human

[149160]WD protein, h-beta TrCp, that interacts with HIV-1 Vpu connects CD4

[149161]to the ER degradation pathway through an F-box motif. Molec. Cell 1:

[149162]565–574, 1998.

[149163]

[149164]

[149165]14157.

[149166]Fakruddin, J. M.; Chaganti, R. S. K.; Murty, V. V. V. S.: Lack

[149167]of BCL10 mutations in germ cell tumors and B cell lymphomas. Cell 97:

[149168]683–688, 1999.

[149169]

[149170]

[149171]14158.

[149172]Isaacson, P. G.; Spencer, J.: The biology of low grade MALT lymphoma. J.

[149173]Clin. Path. 48: 395–397, 1995.

[149174]

[149175]

[149176]14159.

[149177]Ruland, J.; Duncan, G. S.; Elia, A.; del Barco Barrantes, I.;

Nguyen,

[149178]L.; Plyte, S.; Millar, D. G.; Bouchard, D.; Wakeham, A.;

Ohashi, P.

[149179]S.; Mak, T. W.: Bcl10 is a positive regulator of antigen re-
ceptor-induced

[149180]activation of NF-kappa-B and neural tube closure. Cell
104: 33-42,

[149181]2001.

[149182]

[149183]

[149184]14160.

[149185]Thome, M.; Martinon, F.; Hofmann, K.; Rubio, V.; Steiner,
V.; Schneider,

[149186]P.; Mattmann, C.; Tschopp, J.: Equine herpesvirus-2 E10
gene product,

[149187]but not its cellular homologue, activates NF-kappa-B
transcription

[149188]factor and c-Jun N-terminal kinase. J. Biol. Chem. 274:
9962-9968,

[149189]1999.

[149190]

[149191]

[149192]14161.

[149193]Willis, T. G.; Jadayel, D. M.; Du, M.-Q.; Peng, H.; Perry, A.
R.;

[149194]Abdul-Rauf, M.; Price, H.; Karran, L.; Majekodunmi, O.;
Wlodarska,

[149195]I.; Pan, L.; Crook, T.; Hamoudi, R.; Isaacson, P. G.; Dyer, M.
J.

[149196]S.: Bcl10 is involved in t(1;14)(p22;q32) of MALT B cell
lymphoma

[149197]and mutated in multiple tumor types Cell 96: 35-45,
1999.

[149198]

[149199]

[149200]14162.

[149201]Wotherspoon, A. C.; Pan, L. X.; Diss, T. C.; Isaacson, P. G.:

[149202]Cytogenetic study of B-cell lymphoma of mucosa-as-
sociated lymphoid

[149203]tissue. Cancer Genet. Cytogenet. 58: 35-38, 1992.

[149204]

[149205]

[149206]14163.

[149207]Anderson, P.; Nagler-Anderson, C.; O'Brien, C.; Levine, H.;
Watkins,

[149208]S.; Slayter, H. S.; Blue, M. L.; Schlossman, S. F.: A mono-
clonal

[149209]antibody reactive with a 15-kDa cytoplasmic granule-
associated protein

[149210]defines a subpopulation of CD8+ T lymphocytes. J. Im-
mun. 144: 574-582,

[149211]1990.

[149212]

[149213]

[149214]14164.

[149215]Forch, P.; Puig, O.; Kedersha, N.; Martinez, C.; Granneman,
S.;

[149216]Seraphin, B.; Anderson, P.; Valcarcel, J.: The apoptosis-
promoting

[149217]factor TIA-1 is a regulator of alternative pre-mRNA splic-
ing. Molec.

[149218]Cell 6: 1089-1098, 2000.

[149219]

[149220]

[149221]14165.

[149222]Kawakami, A.; Tian, Q.; Streuli, M.; Poe, M.; Edelhoff, S.;
Disteche,

[149223]C. M.; Anderson, P.: Intron-exon organization and chro-
mosomal localization

[149224]of the human TIA-1 gene. J. Immun. 152: 4937-4945,
1994.

[149225]

[149226]

[149227]14166.

[149228]Tian, Q.; Streuli, M.; Saito, H.; Schlossman, S. F.; Ander-
son,

[149229]P.: A polyadenylate binding protein localized to the gran-
ules of

[149230]cytolytic lymphocytes induces DNA fragmentation in tar-
get cells. Cell 67:

[149231]629-639, 1991.

[149232]

[149233]

[149234]14167.

[149235]Zheng, B.; Albrecht, U.; Kaasik, K.; Sage, M.; Lu, W.; Vaishnav,

[149236]S.; Li, Q.; Sun, Z. S.; Eichele, G.; Bradley, A.; Lee, C. C.:
Nonredundant

[149237]roles of the mPer1 and mPer2 genes in the mammalian
circadian clock. Cell 105:

[149238]683–694, 2001.

[149239]

[149240]

[149241]14168.

[149242]Cheng, J.-F.; Boyartchuk, V.; Zhu, Y.: Isolation and mapping of

[149243]human chromosome 21 cDNA: progress in constructing a
chromosome 21

[149244]expression map. Genomics 23: 75–84, 1994.

[149245]

[149246]

[149247]14169.

[149248]Imamura, Y.; Kubota, R.; Wang, Y.; Asakawa, S.; Kudoh, J.;
Mashima,

[149249]Y.; Oguchi, Y.; Shimizu, N.: Human retina-specific amine
oxidase

[149250](RAO): cDNA cloning, tissue expression, and chromosomal
mapping. Genomics 40:

[149251]277–283, 1997.

[149252]

[149253]

[149254]14170.

[149255]Imamura, Y.; Noda, S.; Mashima, Y.; Kudoh, J.; Oguchi, Y.;
Shimizu,

[149256]N.: Human retina-specific amine oxidase: genomic struc-
ture of the

[149257]gene (AOC2), alternatively spliced variant, and mRNA ex-
pression in

[149258]retina. Genomics 51: 293–298, 1998.

[149259]

[149260]

[149261]14171.

[149262]Sirotkin, H.; O'Donnell, H.; DasGupta, R.; Halford, S.; St.
Jore,

[149263]B.; Puech, A.; Parimoo, S.; Morrow, B.; Skoultchi, A.;
Weissman, S.

[149264]M.; Scambler, P.; Kucherlapati, R.: Identification of a new
human

[149265]catenin gene family member (ARVCF) from the region
deleted in velo-cardio-facial

[149266]syndrome. Genomics 41: 75-83, 1997.

[149267]

[149268]

[149269]14172.

[149270]Andrews, N. C.: Mining copper transport genes. Proc. Nat.
Acad.

[149271]Sci. 98: 6543-6545, 2001.

[149272]

[149273]

[149274]14173.

[149275]Boulton, J.; Strickson, A. J.; Jabs, E. W.; Cheng, J.-F.; Fi-
dler,

[149276]C.; Wainscoat, J. S.: Physical mapping of the human ATX1
homologue

[149277](HAH1) to the critical region of the 5q- syndrome within
5q32, and

[149278]immediately adjacent to the SPARC gene. Hum. Genet.
106: 127-129,

[149279]2000.

[149280]

[149281]

[149282]14174.

[149283]Hamza, I.; Faisst, A.; Prohaska, J.; Chen, J.; Gruss, P.;
Gitlin,

[149284]J. D.: The metallochaperone Atox1 plays a critical role in
perinatal

[149285]copper homeostasis. Proc. Nat. Acad. Sci. 98: 6848-6852,
2001.

[149286]

[149287]

[149288]14175.

[149289]Klomp, L. W. J.; Lin, S.-J.; Yuan, D. S.; Klausner, R. D.; Cu-
lotta,

[149290]V. C.; Gitlin, J. D.: Identification and functional expression
of

[149291]HAH1, a novel human gene involved in copper homeosta-
sis. J. Biol.

[149292]Chem. 272: 9221–9226, 1997.

[149293]

[149294]

[149295]14176.

[149296]Castro, A. M.; Snyder, L. M.: G6PD San Jose: a new variant
characterized

[149297]by NADPH inhibition studies. Humangenetik 21: 361–363,
1974.

[149298]

[149299]

[149300]14177.

[149301]Cayanis, E.; Gomperts, E. D.; Balinsky, D.; Disler, P.; Myers,

[149302]A.: G6PD Hillbrow: a new variant of glucose–6–phosphate
dehydrogenase

[149303]associated with drug–induced haemolytic anaemia. Brit. J.
Haemat. 30:

[149304]343–350, 1975.

[149305]

[149306]

[149307]14178.

[149308]Cayanis, E.; Lane, A. B.; Jenkins, T.; Nurse, G. T.; Balinsky,

[149309]D.: Glucose–6–phosphate dehydrogenase Porbandar: a
new slow variant

[149310]with slightly reduced activity in a South African family of
Indian

[149311]descent. Biochem. Genet. 15: 765–773, 1977.

[149312]

[149313]

[149314]14179.

[149315]Cederbaum, A. I.; Beutler, E.: Nonspherocytic hemolytic
anemia

[149316]due to G6PD Grand Prairie. I.R.C.S. 3: 579, 1975.

[149317]

[149318]

[149319]14180.

[149320]Chan, T. K.; Todd, D.: Characteristics and distribution of
glucose-6-phosphate

[149321]dehydrogenase-deficient variants in South China. Am. J.
Hum. Genet. 24:

[149322]475-484, 1972.

[149323]

[149324]

[149325]14181.

[149326]Chan, T. K.; Todd, D.; Lai, M. C. S.: Glucose 6-phosphate
dehydrogenase:

[149327]identity of erythrocyte and leukocyte enzyme with report
of a new

[149328]variant in Chinese. Biochem. Genet. 6: 119-124, 1972.

[149329]

[149330]

[149331]14182.

[149332]Chang, J.-G.; Chiou, S.-S.; Perng, L.-I.; Chen, T.-C.; Liu,
T.-C.;

[149333]Lee, L.-S.; Chen, P.-H.; Tang, T. K.: Molecular characteri-
zation

[149334]of glucose-6-phosphate dehydrogenase (G6PD) deficiency
by natural

[149335]and amplification created restriction sites: five mutations
account

[149336]for most G6PD deficiency cases in Taiwan. Blood 80:
1079-1082, 1992.

[149337]

[149338]

[149339]14183.

[149340]Chen, E. Y.; Cheng, A.; Lee, A.; Kuang, W.-J.; Hillier, L.;
Green,

[149341]P.; Schlessinger, D.; Ciccodicola, A.; D'Urso, M.: Sequence
of human

[149342]glucose-6-phosphate dehydrogenase cloned in plasmids
and a yeast artificial

[149343]chromosome. Genomics 10: 792-800, 1991.

[149344]

[149345]

[149346]14184.

[149347]Chen, H.-L.; Huang, M.-J.; Huang, C.-S.; Tang, T. K.: G6PD
NanKang

[149348](517 T-to-C; 173 phe-to-leu): a new Chinese G6PD variant associated

[149349]with neonatal jaundice. Hum. Hered. 46: 201–204, 1996.

[149350]

[149351]

[149352]14185.

[149353]Chernyak, N. B.; Batischev, A. I.; Lanzina, N. V.; Tokarev, Y.

[149354]N.; Alexeyev, G. A.: Electrophoretic and kinetic properties of glucose-6-phosphate

[149355]dehydrogenase from erythrocytes or patients with hemolytic anemia,

[149356]related to deficiency of the enzyme activity. Vopr. Med. Khim. 23:

[149357]166–171, 1977.

[149358]

[149359]

[149360]14186.

[149361]Childs, B.; Zinkham, W.; Browne, E. A.; Kimbro, E. L.; Torbert,

[149362]J. V.: A genetic study of a defect in glutathione
metabolism of the

[149363]erythrocytes. Bull. Johns Hopkins Hosp. 102: 21–37,
1958.

[149364]

[149365]

[149366]14187.

[149367]Chiu, D. T. Y.; Zuo, L.; Chao, L.; Chen, E.; Louie, E.; Lubin,

[149368]B.; Liu, T. Z.; Du, C. S.: Molecular characterization of glu-
cose-6-phosphate

[149369]dehydrogenase (G6PD) deficiency in patients of Chinese
descent and

[149370]identification of new base substitutions in the human
G6PD gene. Blood 81:

[149371]2150–2154, 1993.

[149372]

[149373]

[149374]14188.

[149375]Chockkalingam, K.; Board, P. G.: Further evidence for het-
erogeneity

[149376]of glucose-6-phosphate dehydrogenase deficiency in
Papua New Guinea. Hum.

[149377]Genet. 56: 209-212, 1980.

[149378]

[149379]

[149380]14189.

[149381]Chockkalingam, K.; Board, P. G.; Breguet, G.: Glucose-
6-phosphate

[149382]dehydrogenase variants of Bali Island (Indonesia). Hum.
Genet. 60:

[149383]60-62, 1982.

[149384]

[149385]

[149386]14190.

[149387]Chockkalingam, K.; Board, P. G.; Nurse, G. T.: Glucose-
6-phosphate

[149388]dehydrogenase deficiency in Papua New Guinea: the de-
scription of 13

[149389]new variants. Hum. Genet. 60: 189-192, 1982.

[149390]

[149391]

[149392]14191.

[149393]Chuanshu, D.; Yankang, X.; Lin, W. Q. R.; Xiaoyun, H.:
Studies

[149394]on erythrocyte glucose-6-phosphate dehydrogenase variants in Chinese.

[149395]I. Gd(-) Lizu-Baisha. Acta Acad. Med. Zhong Shan 2:
649-658, 1981.

[149396]

[149397]

[149398]14192.

[149399]Cocco, P.; Todde, P.; Fornera, S.; Manca, M. B.; Manca, P.
Sias,

[149400]A. R.: Mortality in a cohort of men expressing the glucose-6-phosphate

[149401]dehydrogenase deficiency. Blood 91: 706-709, 1998.

[149402]

[149403]

[149404]14193.

[149405]Cohn, J.; Carter, N.; Warburg, M.: Glucose-6-phosphate

dehydrogenase

[149406]deficiency in a native Danish family: a new variant. Scand.
J. Haemat. 23:

[149407]403–406, 1979.

[149408]

[149409]

[149410]14194.

[149411]Colonna–Romano, S.; Iolascon, A.; Lippo, S.; Pinto, L.; Cu-
tillo,

[149412]S.; Battistuzzi, G.: Genetic heterogeneity at the glucose-
6-phosphate

[149413]dehydrogenase locus in southern Italy: a study on the
population of

[149414]Naples. Hum. Genet. 69: 228–232, 1985.

[149415]

[149416]

[149417]14195.

[149418]Cooper, D. W.; Johnston, P. G.; Murtagh, C. E.; Sharman, G.
B.;

[149419]Vandeberg, J. L.; Poole, W. E.: Sex-linked isozymes and

sex-chromosome

[149420]evolution and inactivation in kangaroos.In: Markeit, C. L.:
Isozymes.

[149421]Developmental Biology. New York: Academic Press (pub.)
III: 1975.

[149422]pp. 559–573.

[149423]

[149424]

[149425]14196.

[149426]Cooper, M. R.; Dechatelet, L. R.; McCall, C. E.; Lavia, M. F.;

[149427]Spurr, C. L.; Baehner, R. L.: Complete deficiency of leuko-
cyte glucose-6-phosphate

[149428]dehydrogenase with defective bactericidal activity. J. Clin.
Invest. 51:

[149429]769–778, 1972.

[149430]

[149431]

[149432]14197.

[149433]Corash, L.; Spielberg, S.; Bartsocas, C.; Boxer, L.; Stein-
herz,

[149434]R.; Sheetz, M.; Egan, M.; Schlesselman, J.; Schulman, J. D.:

Reduced

[149435]chronic hemolysis during high-dose vitamin E administration in Mediterranean-type

[149436]glucose-6-phosphate dehydrogenase deficiency. New Eng. J. Med. 303:

[149437]416-420, 1980.

[149438]

[149439]

[149440]14198.

[149441]Corcoran, C. M.; Calabro, V.; Tamagnini, G.; Town, M.; Haidar,

[149442]B.; Vulliamy, T. J.; Mason, P. J.; Luzzatto, L.: Molecular heterogeneity

[149443]underlying the G6PD Mediterranean phenotype. Hum. Genet. 88: 688-690,

[149444]1992.

[149445]

[149446]

[149447]14199.

- [149448]Costa, E.; Cabeda, J. M.; Vieira, E.; Pinto, R.; Pereira, S. A.;
- [149449]Ferraz, L.; Santos, R.; Barbot, J.: Glucose-6-phosphate dehydrogenase
- [149450]Aveiro: a de novo mutation associated with chronic non-spherocytic
- [149451]hemolytic anemia. Blood 95: 1499-1501, 2000.
- [149452]
- [149453]
- [149454]14200.
- [149455]Crookston, J. H.; Yoshida, A.; Lin, M.; Booser, D. J.: G6PD
- Toronto. Biochem.
- [149456]Genet. 8: 259-265, 1973.
- [149457]
- [149458]
- [149459]14201.
- [149460]Csepregy, M.; Hall, M. K.; Berkow, R. L.; Jackson, S.; Prchal,
- [149461]J. T.: Characterization of a new G6PD variant: G6PD Titusville. Am.
- [149462]J. Med. Sci. 297: 114-117, 1989.

[149463]

[149464]

[149465]14202.

[149466]Csepreghy, M.; Yeilding, A.; Lilly, M.; Hall, K.; Scott, C. W.;

[149467]Prchal, J. T.: Characterization of a new glucose-
6-phosphate dehydrogenase

[149468]variant: G6PD Central City. Am. J. Hemat. 28: 61-62,
1988.

[149469]

[149470]

[149471]14203.

[149472]D'Urso, M.; Luzzatto, L.; Perroni, L.; Ciccodicola, A.; Gen-
tile,

[149473]G.; Peluso, I.; Persico, M. G.; Pizzella, T.; Toniolo, D.; Vul-
liamy,

[149474]T. J.: An extensive search for RFLP in the human glucose-
6-phosphate

[149475]dehydrogenase locus has revealed a silent mutation in the
coding sequence. Am.

[149476]J. Hum. Genet. 42: 735-741, 1988.

[149477]

[149478]

[149479]14204.

[149480]Basse, F.; Stout, J. G.; Sims, P. J.; Wiedmer, T.: Isolation of

[149481]an erythrocyte membrane protein that mediates

Ca(2+)-dependent transbilayer

[149482]movement of phospholipid. J. Biol. Chem. 271:

17205-17210, 1996.

[149483]

[149484]

[149485]14205.

[149486]Garrett, R. M.; Johnson, J. L.; Graf, T. N.; Feigenbaum, A.;

Rajagopalan,

[149487]K. V.: Human sulfite oxidase R160Q: identification of the

mutation

[149488]in a sulfite oxidase-deficient patient and expression and

characterization

[149489]of the mutant enzyme. Proc. Nat. Acad. Sci. 95:

6394-6398, 1998.

[149490]

[149491]

[149492]14206.

[149493]Kisker, C.; Schindelin, H.; Pacheco, A.; Wehbi, W. A.; Garrett,

[149494]R. M.; Rajagopalan, K. V.; Enemark, J. H.; Rees, D. C.:
Molecular

[149495]basis of sulfite oxidase deficiency from the structure of
sulfite

[149496]oxidase. Cell 91: 973–983, 1997.

[149497]

[149498]

[149499]14207.

[149500]Corral, J.; Forster, A.; Thompson, S.; Lampert, F.; Kaneko,
Y.;

[149501]Slater, R.; Kroes, W. G.; van der Schoot, C. E.; Ludwig,
W.–D.; Karpas,

[149502]A.; Pocock, C.; Cotter, F.; Rabbitts, T. H.: Acute leukemias
of different

[149503]lineages have similar MLL gene fusions encoding related
chimeric proteins

[149504]resulting from chromosomal translocation. Proc. Nat.

Acad. Sci. 90:

[149505]8538–8542, 1993.

[149506]

[149507]

[149508]14208.

[149509]Parry, P.; Wei, Y.; Evans, G.: Cloning and characterization
of

[149510]the t(X;11) breakpoint from a leukemic cell line identify a
new member

[149511]of the forkhead gene family. Genes Chromosomes Cancer
11: 79–84,

[149512]1994.

[149513]

[149514]

[149515]14209.

[149516]Peters, U.; Haberhausen, G.; Kostrzewa, M.; Nolte, D.;
Muller,

[149517]U.: AFX1 and p54(nrb): fine mapping, genomic structure,
and exclusion

[149518]as candidate genes of X-linked dystonia parkinsonism.

Hum. Genet. 100:

[149519]569–572, 1997.

[149520]

[149521]

[149522]14210.

[149523]Bohme, B.; VandenBos, T.; Cerretti, D. P.; Park, L. S.;

Holtrich,

[149524]U.; Rubsamen-Waigmann, H.; Strebhardt, K.: Cell-cell ad-
hesion mediated

[149525]by binding of membrane-anchored ligand LERK-2 to the
EPH-related receptor

[149526]human embryonal kinase 2 promotes tyrosine kinase ac-
tivity. J. Biol.

[149527]Chem. 271: 24747–24752, 1996.

[149528]

[149529]

[149530]14211.

[149531]Fletcher, F. A.; Huebner, K.; Shaffer, L. G.; Fairweather, N.
D.;

[149532]Monaco, A. P.; Muller, U.; Druck, T.; Simoneaux, D. K.;

Chelly, J.;

[149533]Belmont, J. W.; Beckmann, M. P.; Lyman, S. D.: Assignment
of the

[149534]gene (EPLG2) encoding a high-affinity binding protein for
the receptor

[149535]tyrosine kinase Elk to a 200-kilobasepair region in human
chromosome

[149536]Xq12. Genomics 25: 334-335, 1995.

[149537]

[149538]

[149539]14212.

[149540]Fletcher, F. A.; Renshaw, B.; Hollingsworth, T.; Baum, P.;
Lyman,

[149541]S. D.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Davi-
son, B.

[149542]L.: Genomic organization and chromosomal localization of
mouse Eplg2,

[149543]a gene encoding a binding protein for the receptor tyro-
sine kinase

[149544]Elk. Genomics 24: 127-132, 1994.

[149545]

[149546]

[149547]14213.

[149548]Palmer, A.; Zimmer, M.; Erdmann, K. S.; Eulenburg, V.;

Porthin,

[149549]A.; Heumann, R.; Deutsch, U.; Klein, R.: EphrinB phospho-
rylation

[149550]and reverse signaling: regulation by Src kinases and PTP-
BL phosphatase. Molec.

[149551]Cell 9: 725–737, 2002.

[149552]

[149553]

[149554]14214.

[149555]Bourke, E.; Frindt, G.; Flynn, P.; Schreiner, G. E.: Primary
hyperoxaluria

[149556]with normal alpha-ketoglutarate:glyoxylate carboligase
activity: treatment

[149557]with isocarboxazid. Ann. Intern. Med. 76: 279–284, 1972.

[149558]

[149559]

[149560]14215.

[149561]Danpure, C. J.: Personal Communication. Middlesex, England

[149562]6/16/1988.

[149563]

[149564]

[149565]14216.

[149566]Danpure, C. J.: Primary hyperoxaluria type 1 and peroxisome-to-mitochondrion

[149567]mistargeting of alanine:glyoxylate aminotransferase.

Biochimie 75:

[149568]309-315, 1993.

[149569]

[149570]

[149571]14217.

[149572]Danpure, C. J.; Jennings, P. R.: Peroxisomal alanine:glyoxylate

[149573]aminotransferase deficiency in primary hyperoxaluria type

I. FEBS

[149574]Lett. 201: 20-24, 1986.

[149575]

[149576]

[149577]14218.

[149578]Danpure, C. J.; Jennings, P. R.; Watts, R. W. E.: Enzymological
ical

[149579]diagnosis of primary hyperoxaluria type 1 by measurement of hepatic

[149580]alanine:glyoxylate aminotransferase activity. Lancet I:
289-291,

[149581]1987.

[149582]

[149583]

[149584]14219.

[149585]Brunelli, S.; Faiella, A.; Capra, V.; Nigro, V.; Simeone, A.;
Cama,

[149586]A.; Boncinelli, E.: Germline mutation in the homeobox
gene EMX2 in

[149587]patients with severe schizencephaly. Nature Genet. 12:
94-96, 1996.

[149588]

[149589]

[149590]14220.

[149591]Rulten, S.; Thorpe, J.; Kay, J.: Identification of eukaryotic

[149592]parvulin homologues: a new subfamily of peptidylprolyl
cis-trans isomerases. Biochem.

[149593]Biophys. Res. Commun. 259: 557-562, 1999.

[149594]

[149595]

[149596]14221.

[149597]Barton, D. E.; Yang-Feng, T. L.; Francke, U.: The human
tyrosine

[149598]aminotransferase gene mapped to the long arm of chro-
mosome 16 (region

[149599]16q22-q24) by somatic cell hybrid analysis and in situ
hybridization. Hum.

[149600]Genet. 72: 221-224, 1986.

[149601]

[149602]

[149603]14222.

[149604]Beinfang, D. C.; Kuwabara, T.; Pueschel, S. M.: The Rich-

ner-Hanhart

[149605]syndrome: report of a case with associated tyrosinemia.

Arch. Ophthal. 94:

[149606]1133-1137, 1976.

[149607]

[149608]

[149609]14223.

[149610]Bohnert, A.; Anton-Lamprecht, I.: Richner-Hanhart's syndrome:

[149611]ultrastructural abnormalities of epidermal keratinization
indicating

[149612]a causal relationship to high intracellular tyrosine levels. J.
Invest.

[149613]Derm. 79: 68-74, 1982.

[149614]

[149615]

[149616]14224.

[149617]Buist, N.: Phenylketonuria and related problems.In: Nyhan,
W.

[149618]L.: Amino Acid Metabolism and Genetic Variation. New

York: McGraw-Hill

[149619](pub.) 1967. Pp. 117 only.

[149620]

[149621]

[149622]14225.

[149623]Chitayat, D.; Balbul, A.; Hani, V.; Mamer, O. A.; Clow, C.;
Scriver,

[149624]C. R.: Hereditary tyrosinaemia type II in a consanguineous
Ashkenazi

[149625]Jewish family: intrafamilial variation in phenotype; absence
of parental

[149626]phenotype effects on the fetus. J. Inherit. Metab. Dis. 15:
198-203,

[149627]1992.

[149628]

[149629]

[149630]14226.

[149631]Crovato, F.; Desirello, G.; Gatti, R.; Babbini, N.; Rebora, A.

[149632]: Richner-Hanhart syndrome spares a plantar autograft.
Arch. Derm. 121:

[149633]539–540, 1985.

[149634]

[149635]

[149636]14227.

[149637]Fellman, J. H.; Vanbellinghen, P. J.; Jones, R. T.; Koler, R.
D.

[149638]: Soluble and mitochondrial forms of tyrosine aminotrans-
ferase: relationship

[149639]to human tyrosinemia. *Biochemistry* 8: 615–622, 1969.

[149640]

[149641]

[149642]14228.

[149643]Garibaldi, L. R.; Siliato, F.; De Martini, I.; Scarsi, M. R.; Ro-
mano,

[149644]C.: Oculocutaneous tyrosinosis: report of two cases in the
same family. *Helv.*

[149645]*Paediat. Acta* 32: 173–180, 1977.

[149646]

[149647]

[149648]14229.

[149649]Goldsmith, L. A.; Kang, E. S.; Bienfang, D. C.; Jimbow, K.;
Gerald,

[149650]P. S.; Baden, H. P.: Tyrosinemia with plantar and palmar
keratosis

[149651]and keratitis. J. Pediat. 83: 798–805, 1973.

[149652]

[149653]

[149654]14230.

[149655]Hanhart, E.: Neue Sonderformen von Keratosis palmo-
plantaris,

[149656]u.a. eine regelmaessig-dominante mit systematisierten
Lipomen, ferner

[149657]2 einfach-rezessive mit Schwachsinn und z.T. mit Horn-
hautveraenderungen

[149658]des Auges (Ektodermatosyndrom). Dermatologica 94:
286–308, 1947.

[149659]

[149660]

[149661]14231.

[149662]Hunziker, N.: Richner–Hanhart syndrome and tyrosinemia
type II. *Dermatologica* 160:

[149663]180–189, 1980.

[149664]

[149665]

[149666]14232.

[149667]Kennaway, N. G.; Buist, N. R. M.: Metabolic studies in a
patient

[149668]with hepatic cytosol tyrosine aminotransferase deficiency.
Pediat.

[149669]*Res.* 5: 287–297, 1971.

[149670]

[149671]

[149672]14233.

[149673]Muller, G.; Scherer, G.; Zentgraf, H.; Ruppert, S.; Her-
rmann,

[149674]B.; Lehrach, H.; Schutz, G.: Isolation, characterization and
chromosomal

[149675]mapping of the mouse tyrosine aminotransferase gene. *J.*
Molec. Biol. 184:

[149676]367–373, 1985.

[149677]

[149678]

[149679]14234.

[149680]Natt, E.; Kao, F.–T.; Rettenmeier, R.; Scherer, G.: Assign-
ment

[149681]of the human tyrosine aminotransferase gene to chromo-
some 16. Hum.

[149682]Genet. 72: 225–228, 1986.

[149683]

[149684]

[149685]14235.

[149686]Huang, L.; Gitschier, J.: A novel gene involved in zinc
transport

[149687]is deficient in the lethal milk mouse. Nature Genet. 17:
292–297,

[149688]1997.

[149689]

[149690]

[149691]14236.

[149692]McKusick, V. A.: On the X Chromosome of Man. Washington: Am.

[149693]Inst. Biol. Sci. (pub.) 1964.

[149694]

[149695]

[149696]14237.

[149697]Ichinose, H.; Ohye, T.; Suzuki, T.; Sumi-Ichinose, C.; Nomura,

[149698]T.; Hagino, Y.; Nagatsu, T.: Molecular cloning of the human Nurr1

[149699]gene: characterization of the human gene and cDNAs.

Gene 230: 233-239,

[149700]1999.

[149701]

[149702]

[149703]14238.

[149704]Law, S. W.; Conneely, O. M.; DeMayo, F. J.; O'Malley, B. W.: Identification

[149705]of a new brain-specific transcription factor, NURR1.

Molec. Endocr. 6:

[149706]2129–2135, 1992.

[149707]

[149708]

[149709]14239.

[149710]Mages, H. W.; Rilke, O.; Bravo, R.; Senger, G.; Kroczeck, R.
A.

[149711]: NOT, a human immediate–early response gene closely
related to the

[149712]steroid/thyroid hormone receptor NAK1/TR3. Molec. En-
docr. 8: 1583–1591,

[149713]1994.

[149714]

[149715]

[149716]14240.

[149717]McEvoy, A. N.; Murphy, E. A.; Ponnio, T.; Conneely, O. M.;
Bresnihan,

[149718]B.; FitzGerald, O.; Murphy, E. P.: Activation of nuclear or-
phan receptor

[149719]NURR1 transcription by NF–kappaB and cyclic adenosine

5-prime-monophosphate

[149720]response element-binding protein in rheumatoid arthritis
synovial

[149721]tissue. J. Immun. 168: 2979-2987, 2002.

[149722]

[149723]

[149724]14241.

[149725]Okabe, T.; Takayanagi, R.; Imasaki, K.; Haji, M.; Nawata,
H.; Watanabe,

[149726]T.: cDNA cloning of a NGFI-B/nur77-related transcription
factor from

[149727]an apoptotic human T cell line. J. Immun. 154:
3871-3879, 1995.

[149728]

[149729]

[149730]14242.

[149731]Torii, T.; Kawarai, T.; Nakamura, S.; Kawakami, H.: Orga-
nization

[149732]of the human orphan nuclear receptor Nurr1 gene. Gene
230: 225-232,

[149733]1999.

[149734]

[149735]

[149736]14243.

[149737]Xu, P.-Y.; Liang, R.; Jankovic, J.; Hunter, C.; Zeng, Y.-X.;
Ashizawa,

[149738]T.; Lai, D.; Le, W.-D.: Association of homozygous
7048G7049 variant

[149739]in the intron six of Nurr1 gene with Parkinson's disease.

Neurology 58:

[149740]881–884, 2002.

[149741]

[149742]

[149743]14244.

[149744]Hanahan, D.: Signaling vascular morphogenesis and main-
tenance. Science 277:

[149745]48–50, 1997.

[149746]

[149747]

[149748]14245.

[149749]Wong, S. M. E.; Lowe, A. W.: Sequence of the cDNA encoding human

[149750]GP-2, the major membrane protein in the secretory granule of the exocrine

[149751]pancreas. Gene 171: 311-312, 1996.

[149752]

[149753]

[149754]14246.

[149755]Yamamura, T.; Hitomi, J.; Nagasaki, K.; Suzuki, M.; Takahashi,

[149756]E.; Saito, S.; Tsukada, T.; Yamaguchi, K.: Human CAAF1 gene--molecular

[149757]cloning, gene structure, and chromosome mapping. Biochem. Biophys.

[149758]Res. Commun. 221: 356-360, 1996.

[149759]

[149760]

[149761]14247.

[149762]Miyoshi, J.; Higashi, T.; Mukai, H.; Ohuchi, T.; Kakunaga, T.:

[149763]Structure and transforming potential of the human cot
oncogene encoding

[149764]a putative protein kinase. Molec. Cell. Biol. 11:
4088–4096, 1991.

[149765]

[149766]

[149767]14248.

[149768]Ohara, R.; Miyoshi, J.; Aoki, M.; Toyoshima, K.: The murine
cot

[149769]proto-oncogene: genome structure and tissue-specific
expression. Jpn.

[149770]J. Cancer Res. 84: 518–525, 1993.

[149771]

[149772]

[149773]14249.

[149774]Gould, D. B.; Walter, M. A.: Cloning, characterization, lo-
calization,

[149775]and mutational screening of the human BARX1 gene. Ge-
nomics 68: 336–342,

[149776]2000.

[149777]

[149778]

[149779]14250.

[149780]Tissier-Seta, J. P.; Mucchielli, M. L.; Mark, M.; Mattei, M. G.;

[149781]Goridis, C.; Brunet, J. F.: Barx1, a new mouse home-
odomain transcription

[149782]factor expressed in cranio-facial ectomesenchyme and the
stomach. Mech.

[149783]Dev. 51: 3-15, 1995.

[149784]

[149785]

[149786]14251.

[149787]Ishihara, H.; Shibasaki, Y.; Kizuki, N.; Wada, T.; Yazaki, Y.;

[149788]Asano, T.; Oka, Y.: Type I phosphatidylinositol-
4-phosphate 5-kinases:

[149789]cloning of the third isoform and deletion/substitution
analysis of

[149790]members of this novel lipid kinase family. J. Biol. Chem.
273: 8741-8748,

[149791]1998.

[149792]

[149793]

[149794]14252.

[149795]Besset, S.; Vincourt, J.-B.; Amalric, F.; Girard, J.-P.: Nuclear

[149796]localization of PAPS synthetase 1: a sulfate activation
pathway in

[149797]the nucleus of eukaryotic cells. FASEB J. 14: 345-354,
2000.

[149798]

[149799]

[149800]14253.

[149801]Faiyaz ul Haque, M.; King, L. M.; Krakow, D.; Cantor, R. M.;
Rusiniak,

[149802]M. E.; Swank, R. T.; Superti-Furga, A.; Haque, S.; Abbas,
H.; Ahmad,

[149803]W.; Ahmad, M.; Cohn, D. H.: Mutations in orthologous
genes in human

[149804]spondyloepimetaphyseal dysplasia and the brachymorphic
mouse. Nature

[149805]Genet. 20: 157–162, 1998.

[149806]

[149807]

[149808]14254.

[149809]Geller, D. H.; Henry, J. G.; Belch, J.; Schwartz, N. B.: Co-
purification

[149810]and characterization of ATP-sulfurylase and adenosine-
5-prime-phosphosulfate

[149811]kinase from rat chondrosarcoma. J. Biol. Chem. 262:
7374–7382, 1987.

[149812]

[149813]

[149814]14255.

[149815]Girard, J.-P.; Baekkevold, E. S.; Amalric, F.: Sulfation in
high

[149816]endothelial venules: cloning and expression of the human
PAPS synthetase. FASEB

[149817]J. 12: 603–612, 1998.

[149818]

[149819]

[149820]14256.

[149821]Lyle, S.; Stanczak, J.; Ng, K.; Schwartz, N. B.: Rat chon-
drosarcoma

[149822]ATP sulfurylase and adenosine 5–prime–phosphosulfate
kinase reside

[149823]on a single bifunctional protein. Biochemistry 33:
5920–5925, 1994.

[149824]

[149825]

[149826]14257.

[149827]Venkatachalam, K. V.; Akita, H.; Strott, C. A.: Molecular cloning,

[149828]expression, and characterization of human bifunctional 3-prime-phosphoadenosine

[149829]5-prime-phosphosulfate synthase and its functional domains. J. Biol.

[149830]Chem. 273: 19311–19320, 1998.

[149831]

[149832]

[149833]14258.

[149834]Park, Y. C.; Ye, H.; Hsia, C.; Segal, D.; Rich, R. L.; Liou, H.-C.;

[149835]Myszka, D. G.; Wu, H.: A novel mechanism of TRAF signaling revealed

[149836]by structural and functional analyses of the TRADD-TRAF2 interaction. Cell 101:

[149837]777–787, 2000.

[149838]

[149839]

[149840]14259.

[149841]Asao, H.; Sasaki, Y.; Arita, T.; Tanaka, N.; Endo, K.; Kasai, H.;

[149842]Takeshita, T; Endo, Y.; Fujita, T.; Sugamura, K.: Hrs is associated

[149843]with STAM, a signal-transducing adaptor molecule: its suppressive

[149844]effect on cytokine-induced cell growth. J. Biol. Chem. 272: 32785-32791,

[149845]1997.

[149846]

[149847]

[149848]14260.

[149849]Takeshita, T.; Arita, T.; Asao, H.; Tanaka, N.; Higuchi, M.; Kuroda,

[149850]H.; Kaneko, K.; Munakata, H.; Endo, Y.; Fujita, T.; Sugamura, K.:

[149851]Cloning of a novel signal-transducing adaptor molecule containing

[149852]an SH3 domain and ITAM. Biochem. Biophys. Res. Commun. 225: 1035-1039,

[149853]1996.

[149854]

[149855]

[149856]14261.

[149857]Everett, R. D.; Meredith, M.; Orr, A.; Cross, A.; Kathoria, M.;

[149858]Parkinson, J.: A novel ubiquitin-specific protease is dynamically

[149859]associated with the PML nuclear domain and binds to a herpesvirus

[149860]regulatory protein. EMBO J. 16: 1519–1530, 1997. Note: Corrected

[149861]republication of EMBO J.:16: 566–577, 1997.

[149862]

[149863]

[149864]14262.

[149865]Li, M.; Chen, D.; Shiloh, A.; Luo, J.; Nikolaev, A. Y.; Qin, J.;

[149866]Gu, W.: Deubiquitination of p53 by HAUSP is an important pathway

[149867]for p53 stabilization. Nature 416: 648–653, 2002.

[149868]

[149869]

[149870]14263.

[149871]Robinson, P. A.; Lomonte, P.; Leek, J. P.; Markham, A. F.;
Everett,

[149872]R. D.: Assignment of herpesvirus-associated ubiquitin-specific protease

[149873]gene HAUSP to human chromosome band 16p13.3 by in situ hybridization. Cytogenet.

[149874]Cell Genet. 83: 100 only, 1998.

[149875]

[149876]

[149877]14264.

[149878]Zhou, G.; Bao, Z. Q.; Dixon, J. E.: Components of a new
human

[149879]protein kinase signal transduction pathway. J. Biol. Chem.
270:

[149880]12665–12669, 1995.

[149881]

[149882]

[149883]14265.

[149884]Lee, J.-D.; Ulevitch, R. J.; Han, J.: Primary structure of
BMK1:

[149885]a new mammalian MAP kinase. Biochem. Biophys. Res.
Commun. 213:

[149886]715–724, 1995.

[149887]

[149888]

[149889]14266.

[149890]Purandare, S. M.; Lee, J.-D.; Patel, P. I.: Assignment of big

[149891]MAP kinase (PRKM7) to human chromosome 17 band
p11.2 with somatic

[149892]cell hybrids. Cytogenet. Cell Genet. 83: 258–259, 1998.

[149893]

[149894]

[149895]14267.

[149896]Regan, C. P.; Li, W.; Boucher, D. M.; Spatz, S.; Su, M. S.;
Kuida,

[149897]K.: Erk5 null mice display multiple extraembryonic vascu-
lar and embryonic

[149898]cardiovascular defects. Proc. Nat. Acad. Sci. 99:

9248–9253, 2002.

[149899]

[149900]

[149901]14268.

[149902]Barlow, G. M.; Chen, X.-N.; Shi, Z. Y.; Lyons, G. E.; Kurnit,
D.

[149903]M.; Celle, L.; Spinner, N. B.; Zackai, E.; Pettenati, M. J.; Van
Riper,

[149904]A. J.; Vekemans, M. J.; Mjaatvedt, C. H.; Korenberg, J. R.:
Down

[149905]syndrome congenital heart disease: a narrowed region and
a candidate

[149906]gene. Genet. Med. 3: 91–101, 2001.

[149907]

[149908]

[149909]14269.

[149910]Schmucker, D.; Clemens, J. C.; Shu, H.; Worby, C. A.; Xiao,
J.;

[149911]Muda, M.; Dixon, J. E.; Zipursky, S. L.: Drosophila Dscam is
an axon

[149912]guidance receptor exhibiting extraordinary molecular di-

versity. Cell 101:

[149913]671–684, 2000.

[149914]

[149915]

[149916]14270.

[149917]Yamakawa, K.; Huo, Y.–K.; Haendel, M. A.; Hubert, R.;
Chen, X.–N.;

[149918]Lyons, G. E.; Korenberg, J. R.: DSCAM: a novel member of
the immunoglobulin

[149919]superfamily maps in a Down syndrome region and is in-
volved in the

[149920]development of the nervous system. Hum. Molec. Genet.
7: 227–237,

[149921]1998.

[149922]

[149923]

[149924]14271.

[149925]Cowan, N. J.; Dobner, P. R.; Fuchs, E. V.; Cleveland, D. W.:
Expression

[149926]of human alpha–tubulin genes: interspecies conservation

of 3-prime

[149927]untranslated regions. Molec. Cell. Biol. 3: 1738-1745,
1983.

[149928]

[149929]

[149930]14272.

[149931]Gerhard, D. S.; Dobner, P. R.; Bruns, G.: Testis specific al-
pha-tubulin

[149932]is on chromosome 2q. (Abstract) Cytogenet. Cell Genet.
40: 639-640,

[149933]1985.

[149934]

[149935]

[149936]14273.

[149937]Hall, J. L.; Cowan, N. J.: Structural features and restricted

[149938]expression of a human alpha-tubulin gene. Nucleic Acids
Res. 13:

[149939]207-223, 1985.

[149940]

[149941]

[149942]14274.

[149943]Miller, F. D.; Naus, C. C. G.; Durand, M.; Bloom, F. E.; Milner,

[149944]R. J.: Isotypes of alpha-tubulin are differentially regulated during

[149945]neuronal maturation. J. Cell Biol. 105: 3065–3073, 1987.

[149946]

[149947]

[149948]14275.

[149949]Zhu, J.; Petersen, S.; Tessarollo, L.; Nussenzweig, A.: Targeted

[149950]disruption of the Nijmegen breakage syndrome gene NBS1 leads to early

[149951]embryonic lethality in mice. Curr. Biol. 11: 105–109, 2001.

[149952]

[149953]

[149954]14276.

[149955]Cooper, P. R.; Smilinich, N. J.; Day, C. D.; Nowak, N. J.;

Reid,

[149956]L. H.; Pearsall, R. S.; Reece, M.; Prawitt, D.; Landers, J.;

Housman,

[149957]D. E.; Winterpacht, A.; Zabel, B. U.; Pelletier, J.; Weissman,

B.

[149958]E.; Shows, T. B.; Higgins, M. J.: Divergently transcribed
overlapping

[149959]genes expressed in liver and kidney and located in the
11p15.5 imprinted

[149960]domain. Genomics 49: 38–51, 1998.

[149961]

[149962]

[149963]14277.

[149964]Kershaw, D. B.; Beck, S. G.; Wharram, B. L.; Wiggins, J. E.;
Goyal,

[149965]M.; Thomas, P. E.; Wiggins, R. C.: Molecular cloning and
characterization

[149966]of human podocalyxin-like protein: orthologous relation-
ship to rabbit

[149967]PCLP1 and rat podocalyxin. J. Biol. Chem. 272:
15708–15714, 1997.

[149968]

[149969]

[149970]14278.

[149971]Kershaw, D. B.; Wiggins, J. E.; Wharram, B. L.; Wiggins, R.
C.

[149972]: Assignment of the human podocalyxin-like protein
(PODXL) gene to

[149973]7q32-q33. Genomics 45: 239-240, 1997.

[149974]

[149975]

[149976]14279.

[149977]Chan, K. K.; Tsui, S. K. W.; Lee, S. M. Y.; Luk, S. C. W.;
Liew,

[149978]C. C.; Fung, K. P.; Waye, M. M. Y.; Lee, C. Y.: Molecular
cloning

[149979]and characterization of FHL2, a novel LIM domain protein
preferentially

[149980]expressed in human heart. Gene 210: 345-350, 1998.

[149981]

[149982]

[149983]14280.

[149984]Genini, M.; Schwalbe, P.; Scholl, F. A.; Remppis, A.; Mattei, M.-G.;

[149985]Schafer, B. W.: Subtractive cloning and characterization of DRAL,

[149986]a novel LIM-domain protein down-regulated in rhabdomyosarcoma. DNA

[149987]Cell Biol. 16: 433-442, 1997.

[149988]

[149989]

[149990]14281.

[149991]Claverie-Martin, F.; Wang, M.; Cohen, S. N.: ARD-1 cDNA from human

[149992]cells encodes a site-specific single-strand endoribonuclease that

[149993]functionally resembles Escherichia coli RNase E. J. Biol. Chem. 272:

[149994]13823-13828, 1997.

[149995]

[149996]

[149997]14282.

- [149998]Van Eynde, A.; Wera, S.; Beullens, M.; Torrekens, S.; Van Leuven, F.; Stalmans, W.; Bollen, M.: Molecular cloning of NIPP-1, a nuclear inhibitor of protein phosphatase-1, reveals homology with polypeptides involved in RNA processing. J. Biol. Chem. 270: 28068-28074, 1995.
- [150002]
- [150003]
- [150004]14283.
- [150005]Tekin, M.; Dogu, F.; Tacyildiz, N.; Akar, E.; Ikiniciogullari, A.; Ogur, G.; Yavuz, G.; Babacan, E.; Akar, N.: 657del5 mutation in the NBS1 gene is associated with Nijmegen breakage syndrome in a Turkish family. Clin. Genet. 62: 84-88, 2002.
- [150009]
- [150010]
- [150011]14284.

[150012]Varon, R.; Reis, A.; Henze, G.; Einsiedel, H. G.; Sperling, K.;

[150013]Seeger, K.: Mutations in the Nijmegen breakage syndrome
gene (NBS1)

[150014]in childhood acute lymphoblastic leukemia (ALL). Cancer
Res. 61:

[150015]3570–3572, 2001.

[150016]

[150017]

[150018]14285.

[150019]Wilda, M.; Demuth, I.; Concannon, P.; Sperling, K.;
Hameister,

[150020]H.: Expression pattern of the Nijmegen breakage syn-
drome gene, Nbs1,

[150021]during murine development. Hum. Molec. Genet. 9:
1739–1744, 2000.

[150022]

[150023]

[150024]14286.

[150025]Wu, X.; Ranganathan, V.; Weisman, D. S.; Heine, W. F.; Cic-
cone,

[150026]D. N.; O'Neill, T. B.; Crick, K. E.; Pierce, K. A.; Lane, W. S.;
Rathbun,

[150027]G.; Livingston, D. M.; Weaver, D. T.: ATM phosphorylation
of Nijmegen

[150028]breakage syndrome protein is required in a DNA damage
response. Nature 405:

[150029]477–482, 2000.

[150030]

[150031]

[150032]14287.

[150033]Zhao, S.; Weng, Y.–C.; Yuan, S.–S. F.; Lin, Y.–T.; Hsu,
H.–C.;

[150034]Lin, S.–C. J.; Gerbino, E.; Song, M.; Zdzienicka, M. Z.; Gatti,
R.

[150035]A.; Shay, J. W.; Ziv, Y.; Shiloh, Y.; Lee, E. Y.–H. P.: Func-
tional

[150036]link between ataxia–telangiectasia and Nijmegen breakage
syndrome

[150037]gene products. Nature 405: 473–477, 2000.

[150038]

[150039]

[150040]14288.

[150041]Mankodi, A.; Urbinati, C. R.; Yuan, Q.-P.; Moxley, R. T.;

Sansone,

[150042]V.; Krym, M.; Henderson, D.; Schalling, M.; Swanson, M. S.;

Thornton,

[150043]C. A.: Muscleblind localizes to nuclear foci of aberrant

RNA in myotonic

[150044]dystrophy types 1 and 2. Hum. Molec. Genet. 10:

2165-2170, 2001.

[150045]

[150046]

[150047]14289.

[150048]Hittner, H. M.; Kretzer, F. L.; Antoszyk, J. H.; Ferrell, R. E.;

[150049]Mehta, R. S.: Variable expressivity of autosomal dominant

anterior

[150050]segment mesenchymal dysgenesis in six generations. Am.

J. Ophthal. 93:

[150051]57-70, 1982.

[150052]

[150053]

[150054]14290.

[150055]Semina, E. V.; Ferrell, R. E.; Mintz-Hittner, H. A.; Bitoun, P.;

[150056]Alward, W. L. M.; Reiter, R. S.; Funkhauser, C.; Daack-Hirsch, S.;

[150057]Murray, J. C.: A novel homeobox gene PITX3 is mutated in families

[150058]with autosomal-dominant cataracts and ASMD. Nature Genet. 19: 167-170,

[150059]1998.

[150060]

[150061]

[150062]14291.

[150063]Semina, E. V.; Murray, J. C.; Reiter, R.; Hrstka, R. F.; Graw,

[150064]J.: Deletion in the promoter region and altered expression of Pitx3

[150065]homeobox gene in aphakia mice. Hum. Molec. Genet. 9: 1575-1585,

[150066]2000.

[150067]

[150068]

[150069]14292.

[150070]Semina, E. V.; Reiter, R. S.; Murray, J. C.: Isolation of a new

[150071]homeobox gene belonging to the Pitx/Rieg family: expression during

[150072]lens development and mapping to the aphakia region on mouse chromosome

[150073]19. Hum. Molec. Genet. 6: 2109–2116, 1997.

[150074]

[150075]

[150076]14293.

[150077]Kent, J.; Lee, M.; Schedl, A.; Boyle, S.; Fantes, J.; Powell, M.;

[150078]Rushmere, N.; Abbott, C.; van Heyningen, V.; Bickmore, W. A.: The

[150079]reticulocalbin gene maps to the WAGR region in human and to the Small

[150080]eye Harwell deletion in mouse. Genomics 42: 260–267, 1997.

[150081]

[150082]

[150083]14294.

[150084]Ozawa, M.: Cloning of a human homologue of mouse
reticulocalbin

[150085]reveals conservation of structural domains in the novel
endoplasmic

[150086]reticulum resident Ca^{2+} -binding protein with multiple
EF-hand motifs. J.

[150087]Biochem. 117: 1113-1119, 1995.

[150088]

[150089]

[150090]14295.

[150091]Ozawa, M.; Muramatsu, T.: Reticulocalbin, a novel endo-
plasmic

[150092]reticulum resident Ca^{2+} -binding protein with multiple
EF-hand motifs

[150093]and a carboxyl-terminal HDEL sequence. J. Biol. Chem.
268: 699-705,

[150094]1993.

[150095]

[150096]

[150097]14296.

[150098]Felbor, U.; Gehrig, A.; Sauer, C. G.; Marquardt, A.; Kohler, M.;

[150099]Schmid, M.; Weber, B. H. F.: Genomic organization and chromosomal

[150100]localization of the interphotoreceptor matrix proteoglycan-1 (IMPG1)

[150101]gene: a candidate for 6q-linked retinopathies. Cytogenet. Cell Genet. 81:

[150102]12-17, 1998.

[150103]

[150104]

[150105]14297.

[150106]Kuehn, M. H.; Hageman, G. S.: Characterization of a cDNA encoding

[150107]IPM 150, a novel human interphotoreceptor matrix chondroitin 6-sulfate

[150108]proteoglycan. (Abstract) Invest. Ophthalm. Vis. Sci. 36: 510 only,

[150109]1995.

[150110]

[150111]

[150112]14298.

[150113]Luo, G.; Leroy, E.; Kozak, C. A.; Polymeropoulos, M. H.;
Horowitz,

[150114]R.: Mapping of the gene (NRAP) encoding N-RAP in the
mouse and human

[150115]genomes. Genomics 45: 229–232, 1997.

[150116]

[150117]

[150118]14299.

[150119]Luo, G.; Zhang, J. Q.; Nguyen, T.–P.; Herrera, A. H.; Pater–
son,

[150120]B.; Horowitz, R.: Complete cDNA sequence and tissue lo–
calization

[150121]of N-RAP, a novel nebulin-related protein of striated
muscle. Cell

[150122]Motil. Cytoskeleton 38: 75–90, 1997.

[150123]

[150124]

[150125]14300.

[150126]Ichikawa, S.; Ozawa, K.; Hirabayashi, Y.: Assignment of a

UDP-glucose:ceramide

[150127]glucosyltransferase gene (Ugcg) to mouse chromosome
band 4B3 by in

[150128]situ hybridization. Cytogenet. Cell Genet. 83: 14–15,
1998.

[150129]

[150130]

[150131]14301.

[150132]Ichikawa, S.; Ozawa, K.; Hirabayashi, Y.: Assignment of a
UDP-glucose:ceramide

[150133]glucosyltransferase gene (UGCG) to human chromosome
band 9q31 by in

[150134]situ hybridization. Cytogenet. Cell Genet. 79: 233–234,
1997.

[150135]

[150136]

[150137]14302.

[150138]Ichikawa, S.; Sakiyama, H.; Suzuki, G.; Hidari, K. I.–P. J.;
Hirabayashi,

[150139]Y.: Expression cloning of a cDNA for human ceramide glu-
cosyltransferase

[150140]that catalyzes the first glycosylation step of glycosphingolipid synthesis. Proc.

[150141]Nat. Acad. Sci. 93: 4638–4643, 1996.

[150142]

[150143]

[150144]14303.

[150145]Watanabe, R.; Wu, K.; Paul, P.; Marks, D. L.; Kobayashi, T.; Pittelkow,

[150146]M. R.; Pagano, R. E.: Up-regulation of glucosylceramide synthase

[150147]expression and activity during human keratinocyte differentiation. J.

[150148]Biol. Chem. 273: 9651–9655, 1998.

[150149]

[150150]

[150151]14304.

[150152]Strahm, B.; Rittweiler, K.; Duffner, U.; Brandau, O.; Orłowska-Volk,

[150153]M.; Karajannis, M. A.; zur Stadt, U.; Tiemann, M.; Reiter, A.; Brandis,

[150154]M.; Meindl, A.; Niemeyer, C. M.: Recurrent B-cell non-Hodgkin's lymphoma

[150155]in two brothers with X-linked lymphoproliferative disease without

[150156]evidence for Epstein-Barr virus infection. Brit. J. Haemat. 108:

[150157]377-382, 2000.

[150158]

[150159]

[150160]14305.

[150161]Sylla, B. S.; Murphy, K.; Cahir-McFarland, E.; Lane, W. S.; Mosialos,

[150162]G.; Kieff, E.: The X-linked lymphoproliferative syndrome gene product

[150163]SH2D1A associates with p62(dok) (Dok1) and activates NF-kappa-beta. Proc.

[150164]Nat. Acad. Sci. 97: 7470-7475, 2000.

[150165]

[150166]

[150167]14306.

[150168]Sylla, B. S.; Wang, Q.; Hayoz, D.; Lathrop, G. M.; Lenoir, G.

[150169]M.: Multipoint linkage mapping of the Xq25–q26 region in
a family

[150170]affected by the X–linked lymphoproliferative syndrome.
Clin. Genet. 36:

[150171]359–462, 1989.

[150172]

[150173]

[150174]14307.

[150175]Tangye, S. G.; Lazetic, S.; Woollatt, E.; Sutherland, G. R.;
Lanier,

[150176]L. L.; Phillips, J. H.: Cutting edge: human 2B4, an activat–
ing NK

[150177]cell receptor, recruits the protein tyrosine phosphatase
SHP–2 and

[150178]the adaptor signaling protein SAP. J. Immun. 162:
6981–6985, 1999.

[150179]

[150180]

[150181]14308.

[150182]Tangye, S. G.; Phillips, J. H.; Lanier, L. L.; Nichols, K. E.

[150183]: Cutting edge: functional requirement for SAP in

2B4-mediated activation

[150184]of human natural killer cells as revealed by the X-linked

lymphoproliferative

[150185]syndrome. J. Immun. 165: 2932–2936, 2000.

[150186]

[150187]

[150188]14309.

[150189]Thorley-Lawson, D. A.; Schooley, R. T.; Bhan, A. K.;

Nadler, L.

[150190]M.: Epstein-Barr virus superinduces a new human B cell

differentiation

[150191]antigen (B-LAST 1) expressed on transformed lym-

phoblasts. Cell 30:

[150192]415–425, 1982.

[150193]

[150194]

[150195]14310.

[150196]Vowels, M. R.; Lam-Po-Tang, R.; Berdoukas, V.; Ford, D.;

Thierry,

[150197]D.; Purtilo, D.; Gluckman, E.: Correction of X-linked lymphoproliferative

[150198]disease by transplantation of cord-blood stem cells. New Eng. J.

[150199]Med. 329: 1623–1625, 1993.

[150200]

[150201]

[150202]14311.

[150203]Williams, L. L.; Rooney, C. M.; Conley, M. E.; Brenner, M. K.;

[150204]Krance, R. A.; Heslop, H. E.: Correction of Duncan's syndrome by

[150205]allogeneic bone marrow transplantation. Lancet 342: 587–588, 1993.

[150206]

[150207]

[150208]14312.

[150209]Wu, C.; Nguyen, K. B.; Pien, G. C.; Wang, N.; Gullo, C.; Duncan,

[150210]H.; Sosa, M. R.; Edwards, M. J.; Borrow, P.; Satoskar, A. R.;

Sharpe,

[150211]A. H.; Biron, C. A.; Terhorst, C.: SAP controls T cell responses

[150212]to virus and terminal differentiation of T(H)2 cells. Nature Immun. 2:

[150213]410–414, 2001.

[150214]

[150215]

[150216]14313.

[150217]Wyandt, H. E.; Skare, J. C.; Grierson, H. L.; Purtilo, D. T.;

[150218]Milunsky, A.: Detection of a chromosomal deletion of Xq25 in an affected

[150219]male with X-linked lymphoproliferative disease. (Abstract) Am. J.

[150220]Hum. Genet. 45 (suppl.): A108 only, 1989.

[150221]

[150222]

[150223]14314.

[150224]Yin, L.; Ferrand, V.; Lavoue, M.–F.; Hayoz, D.; Philippe, N.;

[150225]Souillet, G.; Seri, M.; Giacchino, R.; Castagnola, E.; Hodgson, S.;

[150226]Sylla, B. S.; Romeo, G.: SH2D1A mutation analysis for diagnosis of

[150227]XLP in typical and atypical patients. Hum. Genet. 105: 501–505,

[150228]1999.

[150229]

[150230]

[150231]14315.

[150232]Yin, L.; Tocco, T.; Pauly, S.; Lenoir, G. M.; Romeo, G.: Absence

[150233]of SH2D1A point mutation in 62 Burkitts lymphoma cell lines. Am.

[150234]J. Hum. Genet. 65 (suppl. 1868): A331 only, 1999.

[150235]

[150236]

[150237]14316.

[150238]Laxminarayan, K. M.; Chan, B. K.; Tetaz, T.; Bird, P. I.; Mitchell,

[150239]C. A.: Characterization of a cDNA encoding the 43-kDa
membrane-associated

[150240]inositol-polyphosphate 5-phosphatase. J. Biol. Chem.
269: 17305-17310,

[150241]1994.

[150242]

[150243]

[150244]14317.

[150245]Fink, J. K.; Jones, S. M.; Esposito, C.; Wilkowski, J.: Human

[150246]microtubule-associated protein 1a (MAP1A) gene: ge-
nomic organization,

[150247]cDNA sequence, and developmental-and tissue-specific
expression. Genomics 35:

[150248]577-585, 1996.

[150249]

[150250]

[150251]14318.

[150252]Asker, C.; Steinitz, M.; Andersson, K.; Sumegi, J.; Klein, G.;

[150253]Ingvarsson, S.: Nucleotide sequence of the rat Bmyc gene..

Oncogene 4:

[150254]1523–1527, 1989.

[150255]

[150256]

[150257]14319.

[150258]Ingvarsson, S.; Sundaresan, S.; Jin, P.; Francke, U.; Asker, C.;

[150259]Sumegi, J.; Klein, G.; Sejersen, T.: Chromosome localization and

[150260]expression pattern of Lmyc and Bmyc in murine embryonal carcinoma

[150261]cells. Oncogene 3: 679–685, 1988.

[150262]

[150263]

[150264]14320.

[150265]Kojima, T.; Inazawa, J.; Takamatsu, J.; Rosenberg, R. D.; Saito,

[150266]H.: Human ryudocan core protein: molecular cloning and characterization

[150267]of the cDNA, and chromosomal localization of the gene.

Biochem. Biophys.

[150268]Res. Commun. 190: 814–822, 1993.

[150269]

[150270]

[150271]14321.

[150272]Kojima, T.; Shworak, N. W.; Rosenberg, R. D.: Molecular
cloning

[150273]and expression of two distinct cDNA–encoding heparan
sulfate proteoglycan

[150274]core proteins from a rat endothelial cell line. J. Biol. Chem.
267:

[150275]4870–4877, 1992.

[150276]

[150277]

[150278]14322.

[150279]Yu, H.; Humphries, D. E.; Watkins, M.; Karlinsky, J. B.:
Molecular

[150280]cloning of the human ryudocan promoter. Biochem. Bio–
phys. Res. Commun. 212:

[150281]1139–1144, 1995.

[150282]

[150283]

[150284]14323.

[150285]McBride, O. W.; Yi, H. F.; Srivastava, M.: The human cytochrome

[150286]b561 gene (CYB561) is located at 17q11-qter. Genomics
21: 662-663,

[150287]1994.

[150288]

[150289]

[150290]14324.

[150291]Srivastava, M.: Genomic structure and expression of the
human

[150292]gene encoding cytochrome b(561), an integral protein of
the cromaffin

[150293]granule membrane. J. Biol. Chem. 270: 22714-22720,
1995.

[150294]

[150295]

[150296]14325.

[150297]Albarosa, R.; DiDonato, S.; Finocchiaro, G.: Redefinition of the

[150298]coding sequence of the MXI1 gene and identification of a polymorphic

[150299]repeat in the 3-prime non-coding region that allows the detection

[150300]of loss of heterozygosity of chromosome 10q25 in glioblastomas. Hum.

[150301]Genet. 95: 709–711, 1995.

[150302]

[150303]

[150304]14326.

[150305]Mitchell, C. A.; Speed, C. J.; Nicholl, J.; Sutherland, G. R.:

[150306]Chromosomal mapping of the gene (INPP5A) encoding the 43-kDa membrane-associated

[150307]inositol polyphosphate 5-phosphatase to 10q26.3 by fluorescence in

[150308]situ hybridization. Genomics 31: 139–140, 1996.

[150309]

[150310]

[150311]14327.

[150312]Sugasawa, K.; Ng, J. M. Y.; Masutani, C.; Iwai, S.; van der Spek,

[150313]P. J.; Eker, A. P. M.; Hanaoka, F.; Bootsma, D.; Hoeijmakers, J. H.

[150314]J.: Xeroderma pigmentosum group C protein complex is the initiator

[150315]of global genome nucleotide excision repair. *Molec. Cell* 2: 223–232,

[150316]1998.

[150317]

[150318]

[150319]14328.

[150320]van der Spek, P. J.; Smit, E. M. E.; Beverloo, H. B.; Sugawasa,

[150321]K.; Masutani, C.; Hanaoka, F.; Hoeijmakers, J. H. J.; Hage-meijer,

[150322]A.: Chromosomal localization of three repair genes: the xeroderma

[150323]pigmentosum group C gene and two human homologs of yeast RAD23. *Genomics* 23:

[150324]651–658, 1994.

[150325]

[150326]

[150327]14329.

[150328]van der Spek, P. J.; Visser, C. E.; Hanaoka, F.; Smit, B.;
Hagemeijer,

[150329]A.; Bootsma, D.; Hoeijmakers, J. H. J.: Cloning, compara-
tive mapping,

[150330]and RNA expression of the mouse homologues of the Sac-
charomyces cerevisiae

[150331]nucleotide excision repair gene RAD23. Genomics 31:
20–27, 1996.

[150332]

[150333]

[150334]14330.

[150335]Netik, A.; Forss-Petter, S.; Holzinger, A.; Molzer, B.; Un-
terrainer,

[150336]G.; Berger, J.: Adrenoleukodystrophy-related protein can
compensate

[150337]functionally for adrenoleukodystrophy protein deficiency
(X-ALD):

[150338]implications for therapy. Hum. Molec. Genet. 8: 907–913,
1999.

[150339]

[150340]

[150341]14331.

[150342]Rajantie, I.; Ekman, N.; Iljin, K.; Arighi, E.; Gunji, Y.;
Kaukonen,

[150343]J.; Palotie, A.; Dewerchin, M.; Carmeliet, P.; Alitalo, K.: Bmx
tyrosine

[150344]kinase has a redundant function downstream of angiopoi-
etin and vascular

[150345]endothelial growth factor receptors in arterial endothe-
lium. Molec.

[150346]Cell. Biol. 21: 4647–4655, 2001.

[150347]

[150348]

[150349]14332.

[150350]Tamagnone, L.; Lahtinen, I.; Mustonen, T.; Virtaneva, K.;
Francis,

[150351]F.; Muscatelli, F.; Alitalo, R.; Edvard Smith, C. I.; Larsson,
C.;

[150352]Alitalo, K.: BMX, a novel nonreceptor tyrosine kinase gene
of the

[150353]BTK/ITK/TEC/TK family located in chromosome Xp22.2.

Oncogene 9:

[150354]3683–3688, 1994.

[150355]

[150356]

[150357]14333.

[150358]Dinulos, M. B.; Bassi, M. T.; Rugarli, E. I.; Chapman, V.;
Ballabio,

[150359]A.; Disteche, C. M.: A new region of conservation is de-
fined between

[150360]human and mouse X chromosomes. Genomics 35:
244–247, 1996.

[150361]

[150362]

[150363]14334.

[150364]Schiaffino, M. V.; Bassi, M. T.; Rugarli, E. I.; Renieri, A.;
Galli,

[150365]L.; Ballabio, A.: Cloning of a human homologue of the
Xenopus laevis

[150366]APX gene from the ocular albinism type 1 critical region.

Hum. Molec.

[150367]Genet. 4: 373–382, 1995.

[150368]

[150369]

[150370]14335.

[150371]Bachner, D.; Sedlacek, Z.; Korn, B.; Hameister, H.; Poustka, A.

[150372]: Expression patterns of two human genes coding for different rab

[150373]GDP–dissociation inhibitors (GDIs), extremely conserved proteins involved

[150374]in cellular transport. Hum. Molec. Genet. 4: 701–708, 1995.

[150375]

[150376]

[150377]14336.

[150378]Black, G. C. M.; Perveen, R.; Bonshek, R.; Cahill, M.; Clayton-Smith,

[150379]J.; Lloyd, I. C.; McLeod, D.: Coats' disease of the retina (unilateral

[150380]retinal telangiectasis) caused by somatic mutation in the
NDP gene:

[150381]a role for norrin in retinal angiogenesis. Hum. Molec.
Genet. 8:

[150382]2031–2035, 1999.

[150383]

[150384]

[150385]14337.

[150386]Mao, M.; Fu, G.; Wu, J.–S.; Zhang, Q.–H.; Zhou, J.; Kan,
L.–X.;

[150387]Huang, Q.–H.; He, K.–L.; Gu, B.–W.; Han, Z.–G.; Shen, Y.;
Gu, J.;

[150388]Yu, Y.–P.; Xu, S.–H.; Wang, Y.–X.; Chen, S.–J.; Chen, Z.:
Identification

[150389]of genes expressed in human CD34+ hematopoietic
stem/progenitor cells

[150390]by expressed sequence tags and efficient full-length
cDNA cloning. Proc.

[150391]Nat. Acad. Sci. 95: 8175–8180, 1998.

[150392]

[150393]

[150394]14338.

[150395]Chen, Y.-T.; Gure, A. O.; Tsang, S.; Stockert, E.; Jager, E.;
Knuth,

[150396]A.; Old, L. J.: Identification of multiple cancer/testis anti-
gens

[150397]by allogeneic antibody screening of a melanoma cell line
library. Proc.

[150398]Nat. Acad. Sci. 95: 6919–6923, 1998.

[150399]

[150400]

[150401]14339.

[150402]Lucas, S.; De Smet, C.; Arden, K. C.; Viars, C. S.; Lethe, B.;

[150403]Lurquin, C.; Boon, T.: Identification of a new MAGE gene
with tumor-specific

[150404]expression by representational difference analysis. Cancer
Res. 58:

[150405]743–752, 1998.

[150406]

[150407]

[150408]14340.

[150409]Banfi, B.; Maturana, A.; Jaconi, S.; Arnaudeau, S.; Laforge, T.;

[150410]Sinha, S.; Ligeti, E.; Demaurex, N.; Krause, K.-H.: A mammalian H⁺

[150411]channel generated through alternative splicing of the NADPH oxidase

[150412]homolog NOH-1. Science 287: 138-141, 2000.

[150413]

[150414]

[150415]14341.

[150416]Kikuchi, H.; Hikage, M.; Miyashita, H.; Fukumoto, M.: NADPH oxidase

[150417]subunit, gp91-phox homologue, preferentially expressed in human colon

[150418]epithelial cells. Gene 254: 237-243, 2000.

[150419]

[150420]

[150421]14342.

[150422]Pearce, W. G.; Sanger, R.; Race, R. R.: Ocular albinism and Xg. Lancet I:

[150423]1282–1283, 1968.

[150424]

[150425]

[150426]14343.

[150427]Rosenberg, T.; Schwartz, M.: X-linked ocular albinism:
prevalence

[150428]and mutations—a national study. Europ. J. Hum. Genet. 6:
570–577,

[150429]1998.

[150430]

[150431]

[150432]14344.

[150433]Schiaffino, M. V.; Bassi, M. T.; Galli, L.; Renieri, A.; Bruttini,

[150434]M.; De Nigris, F.; Bergen, A. A. B.; Charles, S. J.; Yates, J. R.

[150435]W., Meindl, A.; Lewis, R. A.; King, R. A.; Ballabio, A.: Anal-
ysis

[150436]of the OA1 gene reveals mutations in only one-third of
patients with

[150437]X-linked ocular albinism. Hum. Molec. Genet. 4:

2319–2325, 1995.

[150438]

[150439]

[150440]14345.

[150441]Schiaffino, M. V.; d'Addio, M.; Alloni, A.; Baschiroto, C.;
Valetti,

[150442]C.; Cortese, K.; Puri, C.; Bassi, M. T.; Colla, C.; De Luca,
M.; Tacchetti,

[150443]C.; Ballabio, A.: Ocular albinism: evidence for a defect in
an intracellular

[150444]signal transduction system. Nature Genet. 23: 108–112,
1999.

[150445]

[150446]

[150447]14346.

[150448]Schnur, R. E.; Gao, M.; Wick, P. A.; Keller, M.; Benke, P. J.;

[150449]Edwards, M. J.; Grix, A. W.; Hockey, A.; Jung, J. H.; Kidd, K.
K.;

[150450]Kistenmacher, M.; Levin, A. V.; and 11 others: OA1 muta-
tions and

[150451]deletions in X-linked ocular albinism. Am. J. Hum. Genet.

62: 800–809,

[150452]1998.

[150453]

[150454]

[150455]14347.

[150456]Schnur, R. E.; Trask, B. J.; van den Engh, G.; Punnett, H. H.;

[150457]Kistenmacher, M.; Tomeo, M. A.; Naidu, R. E.; Nussbaum,

R. L.: An

[150458]Xp22 microdeletion associated with ocular albinism and

ichthyosis:

[150459]approximation of breakpoints and estimation of deletion

size by using

[150460]cloned DNA probes and flow cytometry. Am. J. Hum.

Genet. 45: 706–720,

[150461]1989.

[150462]

[150463]

[150464]14348.

[150465]Schnur, R. E.; Wick, P. A.; Bailey, C.; Rebbeck, T.; Weleber,

[150466]R. G.; Wagstaff, J.; Grix, A. W.; Pagon, R. A.; Hockey, A.;
Edwards,

[150467]M. J.: Phenotypic variability in X-linked ocular albinism:
relationship

[150468]to linkage genotypes. Am. J. Hum. Genet. 55: 484–496,
1994.

[150469]

[150470]

[150471]14349.

[150472]Vogt, A.: Die Iris: Albinismus solum bulbi. Atlas Spalt-
Lampen-Mikroskopie

[150473](pub.) (3rd ed.): 1942.

[150474]

[150475]

[150476]14350.

[150477]Waardenburg, P. J.; Van den Bosch, J.: X-chromosomal oc-
ular albinism

[150478]in Dutch family. Ann. Hum. Genet. 21: 101–122, 1956.

[150479]

[150480]

[150481]14351.

[150482]Zhang, Y.; McMahon, R.; Charles, S. J.; Green, J. S.; Moore, A.

[150483]T.; Barton, D. E.; Yates, J. R. W.: Genetic mapping of the Kallmann

[150484]syndrome and X linked ocular albinism gene loci. J. Med. Genet. 30:

[150485]923–925, 1993.

[150486]

[150487]

[150488]14352.

[150489]Shapira, E.; Ben-Yoseph, Y.; Eyal, G.; Russell, A.: Enzymatically

[150490]inactive red cell carbonic anhydrase B in a family with renal tubular

[150491]acidosis. J. Clin. Invest. 53: 59–63, 1974.

[150492]

[150493]

[150494]14353.

[150495]Cianfriglia, M.; Miggiano, V. C.; Meo, T.; Muller, H. J.;

Muller,

[150496]E.; Battistuzzi, G.: Evidence for synteny between the rabbit gene

[150497]loci coding for HPRT, PGK and G6PD in mouse–rabbit somatic cell hybrids.

[150498](Abstract) Cytogenet. Cell Genet. 25: 142, 1979.

[150499]

[150500]

[150501]14354.

[150502]Holterhus, P. M.; Sinnecker, G. H. G.; Wollmann, H. A.; Struve,

[150503]D.; Homburg, N.; Kruse, K.; Hiort, O.: Expression of two functionally

[150504]different androgen receptors in a patient with androgen insensitivity. Europ.

[150505]J. Pediat. 158: 702–706, 1999.

[150506]

[150507]

[150508]14355.

[150509]Shimmin, L. C.; Chang, B.–H.; Li, W.–H.: Male–driven evo–

lution

[150510]of DNA sequences. Nature 362: 745–747, 1993.

[150511]

[150512]

[150513]14356.

[150514]Warburg, M.: Personal Communication. Copenhagen, Denmark 1966.

[150515]

[150516]

[150517]14357.

[150518]Imasaki, K.; Okabe, T.; Murakami, H.; Tanaka, Y.; Haji, M.; Takayanagi,

[150519]R.; Nawata, H.: Androgen insensitivity syndrome due to new mutations

[150520]in the DNA-binding domain of the androgen receptor. Molec. Cell.

[150521]Endocr. 120: 15–24, 1996.

[150522]

[150523]

[150524]14358.

[150525]Jakubiczka, S.; Nedel, S.; Werder, E. A.; Schleiermacher, E.;

[150526]Theile, U.; Wolff, G.; Wieacker, P.: Mutations of the androgen receptor

[150527]gene in patients with complete androgen insensitivity.

Hum. Mutat. 9:

[150528]57–61, 1997.

[150529]

[150530]

[150531]14359.

[150532]Kang, H.–Y.; Yeh, S.; Fujimoto, N.; Chang, C.: Cloning and characterization

[150533]of human prostate coactivator ARA54, a novel protein that associates

[150534]with the androgen receptor. J. Biol. Chem. 274: 8570–8576, 1999.

[150535]

[150536]

[150537]14360.

[150538]Kazemi–Esfarjani, P.; Beitel, L. K.; Trifiro, M.; Kaufman, M.;

[150539]Rennie, P.; Sheppard, P.; Matusik, R.; Pinsky, L.: Substitution of
[150540]valine-865 by methionine or leucine in the human androgen receptor
[150541]causes complete or partial androgen insensitivity, respectively with
[150542]distinct androgen receptor phenotypes. *Molec. Endocr.* 7:
37-46,
[150543]1993.

[150544]

[150545]

[150546]14361.

[150547]Kittles, R. A.; Young, D.; Weinrich, S.; Hudson, J.; Argypoulos,
[150548]G.; Ukoli, F.; Adams-Campbell, L.; Dunston, G. M.: Extent of linkage
[150549]disequilibrium between the androgen receptor gene CAG and GGC repeats
[150550]in human populations: implications for prostate cancer risk. *Hum.*
[150551]*Genet.* 109: 253-261, 2001.

[150552]

[150553]

[150554]14362.

[150555]Klocker, H.; Kaspar, F.; Eberle, J.; Uberreiter, S.; Radmayr,

[150556]C.; Bartsch, G.: Point mutation in the DNA binding domain
of the

[150557]androgen receptor in two families with Reifenstein syn-
drome. Am.

[150558]J. Med. Genet. 50: 1318–1327, 1992.

[150559]

[150560]

[150561]14363.

[150562]Knoke, I.; Allera, A.; Wieacker, P.: Significance of the CAG

[150563]repeat length in the androgen receptor gene (AR) for the
transactivation

[150564]function of an M780I mutant AR. Hum. Genet. 104:
257–261, 1999.

[150565]

[150566]

[150567]14364.

[150568]Kobayashi, Y.; Miwa, S.; Merry, D. E.; Kume, A.; Mei, L.;

Doyu,

[150569]M.; Sobue, G.: Caspase-3 cleaves the expanded androgen
receptor protein

[150570]of spinal and bulbar muscular atrophy in a polyglutamine
repeat length-dependent

[150571]manner. Biochem. Biophys. Res. Commun. 252: 145-150,
1998.

[150572]

[150573]

[150574]14365.

[150575]Koivisto, P. A.; Schleutker, J.; Helin, H.; Ehren-van Eekelen,

[150576]C.; Kallioniemi, O.-P.; Trapman, J.: Androgen receptor
gene alterations

[150577]and chromosomal gains and losses in prostate carcinomas
appearing

[150578]during finasteride treatment for benign prostatic hyper-
plasia. Clin.

[150579]Cancer Res. 5: 3578-3582, 1999.

[150580]

[150581]

[150582]14366.

[150583]Kooy, R. F.; Reyniers, E.; Storm, K.; Vits, L.; van Velzen, D.;

[150584]de Ruiter, P. E.; Brinkmann, A. O.; de Paepe, A.; Willems, P.
J.:

[150585]CAG repeat contraction in the androgen receptor gene in
three brothers

[150586]with mental retardation. Am. J. Med. Genet. 85: 209–213,
1999.

[150587]

[150588]

[150589]14367.

[150590]La Spada, A.; Fischbeck, K. H.: Androgen receptor gene
defect

[150591]in X-linked spinal and bulbar muscular atrophy. (Abstract)
Am. J.

[150592]Hum. Genet. 49 (suppl.): 20, 1991.

[150593]

[150594]

[150595]14368.

[150596]Lewis, M.; Kaita, H.; Giblett, E. R.; Anderson, J.; Philipps,
[150597]S.; Steinberg, A. G.; McAlpine, P. J.: Multiplicity of genetic
polymorphisms
[150598]of blood in the Schmiedeleut Hutterites. Am. J. Med.
Genet. 22:
[150599]477–485, 1985.

[150600]

[150601]

[150602]14369.

[150603]Lim, H. N.; Chen, H.; McBride, S.; Dunning, A. M.; Nixon,
R. M.;

[150604]Hughes, I. A.; Hawkins, J. R.: Longer polyglutamine tracts
in the

[150605]androgen receptor are associated with moderate to severe
undermasculinized

[150606]genitalia in XY males. Hum. Molec. Genet. 9: 829–834,
2000.

[150607]

[150608]

[150609]14370.

[150610]Lobaccaro, J.-M.; Lumbroso, S.; Belon, C.; Galtier-Dereure, F.;

[150611]Bringer, J.; Lesimple, T.; Namer, M.; Cutuli, B. F.; Pujol, H.; Sultan,

[150612]C.: Androgen receptor gene mutation in male breast cancer. Hum.

[150613]Molec. Genet. 2: 1799–1802, 1993.

[150614]

[150615]

[150616]14371.

[150617]Lubahn, D. B.; Brown, T. R.; Simental, J. A.; Higgs, H. N.; Migeon,

[150618]C. J.; Wilson, E. M.; French, F. S.: Sequence of the intron/exon

[150619]junctions of the coding region of the human androgen receptor gene

[150620]and identification of a point mutation in a family with complete androgen

[150621]insensitivity. Proc. Nat. Acad. Sci. 86: 9534–9538, 1989.

[150622]

[150623]

[150624]14372.

[150625]Lubahn, D. B.; Joseph, D. R.; Sar, M.; Tan, J.; Higgs, H. N.;

[150626]Larson, R. E.; French, F. S.; Wilson, E. M.: The human androgen receptor:

[150627]complementary deoxyribonucleic acid cloning, sequence analysis and

[150628]gene expression in prostate. Molec. Endocr. 2: 1265–1275, 1988.

[150629]

[150630]

[150631]14373.

[150632]Lubahn, D. B.; Joseph, D. R.; Sullivan, P. M.; Willard, H. F.;

[150633]French, F. S.; Wilson, E. M.: Cloning of human androgen receptor

[150634]complementary DNA and localization to the X chromosome. Science 240:

[150635]327–330, 1988.

[150636]

[150637]

[150638]14374.

[150639]Lumbroso, S.; Lobaccaro, J. M.; Georget, V.; Leger, J.; Poujol,

[150640]N.; Terouanne, B.; Evain-Brion, D.; Czernichow, P.; Sultan, C.: A

[150641]novel substitution (leu707-to-arg) in exon 4 of the androgen receptor

[150642]gene causes complete androgen resistance. J. Clin. Endocr. Metab. 81:

[150643]1984-1988, 1996.

[150644]

[150645]

[150646]14375.

[150647]Stuppia, L.; Calabrese, G.; Franchi, P. G.; Mingarelli, R.; Gatta,

[150648]V.; Palka, G.; Dallapiccola, B.: Widening of a Y-chromosome interval-6

[150649]deletion transmitted from a father to his infertile son accounts for

[150650]an oligozoospermia critical region distal to the RBM1 and DAZ genes.

[150651](Letter) Am. J. Hum. Genet. 59: 1393-1395, 1996.

[150652]

[150653]

[150654]14376.

[150655]Brown, G. M.; Furlong, R. A.; Sargent, C. A.; Erickson, R. P.;

[150656]Longepied, G.; Mitchell, M.; Jones, M. H.; Hargreave, T. B.;

Cooke,

[150657]H. J.; Affara, N. A.: Characterisation of the coding sequence and

[150658]fine mapping of the human DFFRY gene and comparative expression analysis

[150659]and mapping to the Sxr-b interval of the mouse Y chromosome of the

[150660]Dffry gene. Hum. Molec. Genet. 7: 97-107, 1998.

[150661]

[150662]

[150663]14377.

[150664]Ferlin, A.; Moro, E.; Garolla, A.; Foresta, C.: Human male infertility

[150665]and Y chromosome deletions: role of the AZF-candidate genes DAZ, RBM

[150666]and DFFRY. Hum. Reprod. 14: 1710–1716, 1999.

[150667]

[150668]

[150669]14378.

[150670]Foresta, C.; Ferlin, A.; Moro, E.: Deletion and expression analysis

[150671]of AZFa genes on the human Y chromosome revealed a major role for

[150672]DBY in male infertility. Hum. Molec. Genet. 9: 1161–1169, 2000.

[150673]

[150674]

[150675]14379.

[150676]Sargent, C. A.; Boucher, C. A.; Kirsch, S.; Brown, G.; Weiss,

[150677]B.; Trundley, A.; Burgoyne, P.; Saut, N.; Durand, C.; Levy, N.; Terriou,

[150678]P.; Hargreave, T.; Cooke, H.; Mitchell, M.; Rappold, G. A.; Affara,

[150679]N. A.: The critical region of overlap defining the AZFa male infertility

[150680]interval of proximal Yq contains three transcribed sequences. J.

[150681]Med. Genet. 36: 670–677, 1999.

[150682]

[150683]

[150684]14380.

[150685]Shen, P.; Wang, F.; Underhill, P. A.; Franco, C.; Yang, W.–H.;

[150686]Roxas, A.; Sung, R.; Lin, A. A.; Hyman, R. W.; Vollrath, D.; Davis,

[150687]R. W.; Cavalli–Sforza, L. L.; Oefner, P. J.: Population genetic implications

[150688]from sequence variation in four Y chromosome genes. Proc. Nat. Acad.

[150689]Sci. 97: 7354–7359, 2000.

[150690]

[150691]

[150692]14381.

[150693]Sun, C.; Skaletsky, H.; Birren, B.; Devon, K.; Tang, Z.; Silber,

[150694]S.; Oates, R.; Page, D. C.: An azoospermic man with a de novo point

[150695]mutation in the Y-chromosomal gene USP9Y. Nature Genet. 23: 429–432,

[150696]1999.

[150697]

[150698]

[150699]14382.

[150700]Thomson, R.; Pritchard, J. K.; Shen, P.; Oefner, P. J.; Feldman,

[150701]M. W.: Recent common ancestry of human Y chromosomes: evidence from

[150702]DNA sequence data. Proc. Nat. Acad. Sci. 97: 7360–7365, 2000.

[150703]

[150704]

[150705]14383.

[150706]Schiebel, K.; Winkelmann, M.; Mertz, A.; Xu, X.; Page, D. C.; Weil,

[150707]D.; Petit, C.; Rappold, G. A.: Abnormal XY interchange between a

[150708]novel isolated protein kinase gene, PRKY, and its homologue, PRKX,

[150709]accounts for one third of all (Y+)XX males and (Y-)XY females. Hum.

[150710]Molec. Genet. 6: 1985-1989, 1997.

[150711]

[150712]

[150713]14384.

[150714]Greenfield, A.; Scott, D.; Pennisi, D.; Ehrmann, I.; Ellis, P.;

[150715]Cooper, L.; Simpson, E.; Koopman, P.: An H-YDb epitope is encoded

[150716]by a novel mouse Y chromosome gene. Nature Genet. 14: 474-478, 1996.

[150717]

[150718]

[150719]14385.

[150720]Scott, D.; Addey, C.; Ellis, P; James, E.; Mitchell, M. J.; Saut,

[150721]N.; Jurcevic, S.; Simpson, E.: Dendritic cells permit identification

[150722]of genes encoding MHC class II-restricted epitopes of

transplantation

[150723]antigens. Immunity 12: 711–720, 2000.

[150724]

[150725]

[150726]14386.

[150727]Wu, G.; Chai, J.; Suber, T. L.; Wu, J.–W.; Du, C.; Wang, X.;
Shi,

[150728]Y.: Structural basis of IAP recognition by Smac/DIABLO.
Nature 408:

[150729]1008–1012, 2000.

[150730]

[150731]

[150732]14387.

[150733]Macke, J. P.; Hu, N.; Hu, S.; Bailey, M.; King, V. L.; Brown,

[150734]T.; Hamer, D.; Nathans, J.: Sequence variation in the an-
drogen receptor

[150735]gene is not a common determinant of male sexual orien-
tation. Am.

[150736]J. Hum. Genet. 53: 844–852, 1993.

[150737]

[150738]

[150739]14388.

[150740]Yang, R.-B.; Fulle, H.-J.; Garbers, D. L.: Chromosomal localization

[150741]and genomic organization of genes encoding guanylyl cyclase receptors

[150742]expressed in olfactory sensory neurons and retina. Genomics 31:

[150743]367-372, 1996.

[150744]

[150745]

[150746]14389.

[150747]Nakamura, H.; Izumoto, Y.; Kambe, H.; Kuroda, T.; Mori, T.; Kawamura,

[150748]K.; Yamamoto, H.; Kishimoto, T.: Molecular cloning of complementary

[150749]DNA for a novel human hepatoma-derived growth factor: its homology

[150750]with high mobility group-1 protein. J. Biol. Chem. 269: 25143-25149,

[150751]1994.

[150752]

[150753]

[150754]14390.

[150755]Wanschura, S.; Schoenmakers, E. F. P. M.; Huysmans, C.;
Bartnitzke,

[150756]S.; Van de Ven, W. J. M.; Bullerdiek, J.: Mapping of the
gene encoding

[150757]the human hepatoma-derived growth factor (HDGF) with
homology to the

[150758]high-mobility group (HMG)-1 protein to Xq25. Genomics
32: 298-300,

[150759]1996.

[150760]

[150761]

[150762]14391.

[150763]Baens, M.; Marynen, P.: A human homologue (BICD1) of
the Drosophila

[150764]bicaudal-D gene. Genomics 45: 601-606, 1997.

[150765]

[150766]

[150767]14392.

[150768]Bullock, S. L.; Ish-Horowicz, D.: Conserved signals and
machinery

[150769]for RNA transport in Drosophila oogenesis and embryoge-
nesis. Nature 414:

[150770]611–616, 2001.

[150771]

[150772]

[150773]14393.

[150774]McMurtrie, E. B.; Barbosa, M. D. F. S.; Zerial, M.;
Kingsmore,

[150775]S. F.: Rab17 and Rab18, small GTPases with specificity for
polarized

[150776]epithelial cells: genetic mapping in the mouse. Genomics
45: 623–625,

[150777]1997.

[150778]

[150779]

[150780]14394.

[150781]Schafer, U.; Seibold, S.; Schneider, A.; Neugebauer, E.: Iso-
lation

[150782]and characterisation of the human rab18 gene after stim-
ulation of

[150783]endothelial cells with histamine. FEBS Lett. 466: 148–154,
2000.

[150784]

[150785]

[150786]14395.

[150787]Doupnik, C. A.; Davidson, N.; Lester, H. A.: The inward
rectifier

[150788]potassium channel family. Curr. Opin. Neurobiol. 5:
268–277, 1995.

[150789]

[150790]

[150791]14396.

[150792]Tada, Y.; Horio, Y.; Takumi, T.; Terayama, M.; Tsuji, L.;
Copeland,

[150793]N. G.; Jenkins, N. A.; Kurachi, Y.: Assignment of the glial
inwardly

[150794]rectifying potassium channel K(AB)–2/Kir4.1 (Kcnj10)

gene to the distal

[150795]region of mouse chromosome 1. Genomics 45: 629–630,
1997.

[150796]

[150797]

[150798]14397.

[150799]Takumi, T.; Ishii, T.; Horio, Y.; Morishige, K.-I.; Takahashi,

[150800]N.; Yamada, M.; Yamashita, T.; Kiyama, H.; Sohmiya, K.;
Nakanishi,

[150801]S.; Kurachi, Y.: A novel ATP-dependent inward rectifier
potassium

[150802]channel expressed predominantly in glial cells. J. Biol.
Chem. 270:

[150803]16339–16436, 1995.

[150804]

[150805]

[150806]14398.

[150807]Schenker, T.; Trueb, B.: Assignment of the gene for a de-
velopmentally

[150808]regulated GTP-binding protein (DRG2) to human chromo-

some bands 17p13–p12

[150809]by in situ hybridization. Cytogenet. Cell Genet. 79:
274–275, 1997.

[150810]

[150811]

[150812]14399.

[150813]Heikkinen, J.; Risteli, M.; Wang, C.; Latvala, J.; Rossi, M.;
Valtavaara,

[150814]M.; Myllyla, R.: Lysyl hydroxylase 3 is a multifunctional
protein

[150815]possessing collagen glucosyltransferase activity. J. Biol.
Chem. 275:

[150816]36158–36163, 2000.

[150817]

[150818]

[150819]14400.

[150820]Passoja, K.; Rautavuoma, K.; Ala-Kokko, L.; Kosonen, T.;
Kivirikko,

[150821]K. I.: Cloning and characterization of a third human lysyl
hydroxylase

[150822]isoform. Proc. Nat. Acad. Sci. 95: 10482–10486, 1998.

[150823]

[150824]

[150825]14401.

[150826]Valtavaara, M.; Szpirer, C.; Szpirer, J.; Myllyla, R.: Primary

[150827]structure, tissue distribution, and chromosomal localization of a

[150828]novel isoform of lysyl hydroxylase (lysyl hydroxylase 3). J. Biol.

[150829]Chem. 273: 12881–12886, 1998.

[150830]

[150831]

[150832]14402.

[150833]Rohan, P. J.; Davis, P.; Moskaluk, C. A.; Kearns, M.;
Krutzsch,

[150834]H.; Siebenlist, U.; Kelly, K.: PAC-1: a mitogen-induced nuclear protein

[150835]tyrosine phosphatase. Science 259: 1763-1766, 1993.

[150836]

[150837]

[150838]14403.

[150839]Ward, Y.; Gupta, S.; Jensen, P.; Wartmann, M.; Davis, R. J.;
Kelly,

[150840]K.: Control of MAP kinase activation by the mitogen-induced threonine/tyrosine

[150841]phosphatase PAC1. Nature 367: 651-654, 1994.

[150842]

[150843]

[150844]14404.

[150845]Yi, H.; Morton, C. C.; Weremowicz, S.; McBride, O. W.;

Kelly, K.

[150846] Genomic organization and chromosomal localization of
the DUSP2 gene,

[150847] encoding a MAP kinase phosphatase, to human
2p11.2-q11. Genomics 28:

[150848] 92-96, 1995.

[150849]

[150850]

[150851] 14405.

[150852] Ishibashi, T.; Bottaro, D. P.; Michieli, P.; Kelley, C. A.;
Aaronson,

[150853] S. A.: A novel dual specificity phosphatase induced by
serum stimulation

[150854] and heat shock. J. Biol. Chem. 269: 29897-29902, 1994.

[150855]

[150856]

[150857] 14406.

[150858] Kwak, S. P.; Dixon, J. E.: Multiple dual specificity protein
tyrosine

[150859] phosphatases are expressed and regulated differentially in

liver cell

[150860]lines. J. Biol. Chem. 270: 1156–1160, 1995.

[150861]

[150862]

[150863]14407.

[150864]Kovalenko, O. V.; Golub, E. I.; Bray–Ward, P.; Ward, D. C.;
Radding,

[150865]C. M.: A novel nucleic acid–binding protein that interacts
with human

[150866]Rad51 recombinase. Nucleic Acids Res. 25: 4946–4953,
1997.

[150867]

[150868]

[150869]14408.

[150870]Mizuta, R.; LaSalle, J. M.; Cheng, H.–L.; Shinohara, A.;
Ogawa,

[150871]H.; Copeland, N.; Jenkins, N. A.; Lalande, M.; Alt, F. W.:
RAB22

[150872]and RAB163/mouse BRCA2: proteins that specifically in–
teract with the

[150873]RAD51 protein. Proc. Nat. Acad. Sci. 94: 6927–6932,

1997.

[150874]

[150875]

[150876]14409.

[150877]Blasius, R.; Weber, R. G.; Lichter, P.; Ogilvie, A.: A novel orphan

[150878]G protein-coupled receptor primarily expressed in the brain is localized

[150879]on human chromosomal band 2q21. J. Neurochem. 70: 1357-1365, 1998.

[150880]

[150881]

[150882]14410.

[150883]Raport, C. J.; Schweickart, V. L.; Chantry, D.; Eddy, R. L., Jr.;

[150884]Shows, T. B.; Godiska, R.; Gray, P. W.: New members of the chemokine

[150885]receptor gene family. J. Leukoc. Biol. 59: 18-23, 1996.

[150886]

[150887]

[150888]14411.

[150889]Shindo, M.; Nakano, H.; Kuroyanagi, H.; Shirasawa, T.; Mi-
hara,

[150890]M.; Gilbert, D. J.; Jenkins, N. A.; Copeland, N. G.; Yagita,
H.; Okumura,

[150891]K.: cDNA cloning, expression, subcellular localization, and
chromosomal

[150892]assignment of mammalian aurora homologues, aurora-
related kinase (ARK)

[150893]1 and 2. Biochem. Biophys. Res. Commun. 244: 285–292,
1998.

[150894]

[150895]

[150896]14412.

[150897]Adachi, H.; Tsujimoto, M.; Hattori, M.; Arai, H.; Inoue, K.:
cDNA

[150898]cloning of human cytosolic platelet-activating factor
acetylhydrolase

[150899]gamma-subunit and its mRNA expression in human tis-
sues. Biochem.

[150900]Biophys. Res. Commun. 214: 180–187, 1995.

[150901]

[150902]

[150903]14413.

[150904]Presky, D. H.; Yang, H.; Minetti, L. J.; Chua, A. O.; Nabavi, N.;

[150905]Wu, C.-Y.; Gately, M. K.; Gubler, U.: A functional interleukin 12

[150906]receptor complex is composed of two beta-type cytokine receptor subunits. Proc.

[150907]Nat. Acad. Sci. 93: 14002-14007, 1996.

[150908]

[150909]

[150910]14414.

[150911]Yamamoto, K.; Kobayashi, H.; Miura, O.; Hirosawa, S.; Miyasaka,

[150912]N.: Assignment of IL12RB1 and IL12RB2, interleukin-12 receptor beta-1

[150913]and beta-2 chains, to human chromosome 19 band p13.1 and chromosome

[150914]1 band p31.2, respectively, by in situ hybridization. Cytogenet.

[150915]Cell Genet. 77: 257–258, 1997.

[150916]

[150917]

[150918]14415.

[150919]Lafreniere, R. G.; Rochefort, D. L.; Kibar, Z.; Fon, E. A.;
Han,

[150920]F.-Y.; Cochius, J.; Kang, X.; Baird, S.; Korneluk, R. G.; An-
dermann,

[150921]E.; Rommens, J. M.; Rouleau, G. A.: Isolation and charac-
terization

[150922]of GT335, a novel human gene conserved in Escherichia
coli and mapping

[150923]to 21q22.3. Genomics 38: 264–272, 1996.

[150924]

[150925]

[150926]14416.

[150927]Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.;
Asakawa,

[150928]S.; Ito, F.; Shimizu, N.: Isolation of cDNA for a novel hu-
man protein

[150929]KNP-I that is homologous to the E. coli SCRP-27A protein
from the

[150930]autoimmune polyglandular disease type I (APECED) region
of chromosome

[150931]21q22.3. Biochem. Biophys. Res. Commun. 225: 608-616,
1996.

[150932]

[150933]

[150934]14417.

[150935]Scott, H. S.; Chen, H.; Rossier, C.; Lalioti, M. D.; An-
tonarakis,

[150936]S. E.: Isolation of a human gene (HES1) with homology to
an Escherichia

[150937]coli and a zebrafish protein that maps to chromosome
21q22.3. Hum.

[150938]Genet. 99: 616-623, 1997.

[150939]

[150940]

[150941]14418.

[150942]Bernier-Villamor, V.; Sampson, D. A.; Matunis, M. J.; Lima,
C.

[150943]D.: Structural basis for E2-mediated SUMO conjugation revealed by

[150944]a complex between ubiquitin-conjugating enzyme Ubc9 and RanGAP1. Cell 108:

[150945]345–356, 2002.

[150946]

[150947]

[150948]14419.

[150949]Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association of

[150950]FHIT (fragile histidine triad), a candidate tumour suppressor gene,

[150951]with the ubiquitin-conjugating enzyme hUBC9. Biochem. J. 352: 443–448,

[150952]2000.

[150953]

[150954]

[150955]14420.

[150956]Tachibana, M.; Iwata, N.; Watanabe, A.; Nobukuni, Y.; Ploplis,

[150957]B.; Kajigaya, S.: Assignment of the gene for a ubiquitin-conjugating

[150958]enzyme (UBE2I) to human chromosome band 16p13.3 by in situ hybridization. Cytogenet.

[150959]Cell Genet. 75: 222-223, 1996.

[150960]

[150961]

[150962]14421.

[150963]Wang, Z.-Y.; Qiu, Q.-Q.; Seufert, W.; Taguchi, T.; Testa, J. R.;

[150964]Whitmore, S. A.; Callen, D. F.; Welsh, D.; Shenk, T.; Deuel, T. F.

[150965]: Molecular cloning of the cDNA and chromosome localization of the

[150966]gene for human ubiquitin-conjugating enzyme 9. J. Biol. Chem. 271:

[150967]24811-24816, 1996.

[150968]

[150969]

[150970]14422.

[150971]Yasugi, T.; Howley, P. M.: Identification of the structural
and

[150972]functional human homolog of the yeast ubiquitin conjugating enzyme

[150973]UBC9. Nucleic Acids Res. 24: 2005–2010, 1996.

[150974]

[150975]

[150976]14423.

[150977]Bowen, M. A.; Patel, D. D.; Li, X.; Modrell, B.; Malacko, A.
R.;

[150978]Wang, W.–C.; Marquardt, H.; Neubauer, M.; Pesando, J. M.;
Francke,

[150979]U.; Haynes, B. F.; Aruffo, A.: Cloning, mapping, and characterization

[150980]of activated leukocyte–cell adhesion molecule (ALCAM), a
CD6 ligand. J.

[150981]Exp. Med. 181: 2213–2220, 1995.

[150982]

[150983]

[150984]14424.

[150985]Skonier, J. E.; Bowen, M. A.; Emswiler, J.; Aruffo, A.; Bajorath,

[150986]J.: Recognition of diverse proteins by members of the immunoglobulin

[150987]superfamily: delineation of the receptor binding site in the human

[150988]CD6 ligand ALCAM. Biochemistry 35: 12287–12291, 1996.

[150989]

[150990]

[150991]14425.

[150992]Bord, S.; Horner, A.; Beavan, S.; Compston, J.: Estrogen receptors

[150993]alpha and beta are differentially expressed in developing human bone. J.

[150994]Clin. Endocr. Metab. 86: 2309–2314, 2001.

[150995]

[150996]

[150997]14426.

[150998]Watanabe, T. K.; Fujiwara, T.; Kawai, A.; Shimizu, F.; Takami,

[150999]S.; Hirano, H.; Okuno, S.; Ozaki, K.; Takeda, S.; Shimada, Y.; Nagata,

[151000]M.; Takaichi, A.; Takahashi, E.; Nakamura, Y.; Shin, S.: Cloning,

[151001]expression, and mapping of UBE2I, a novel gene encoding a human homologue

[151002]of yeast ubiquitin-conjugating enzymes which are critical for regulating

[151003]the cell cycle. Cytogenet. Cell Genet. 72: 86–89, 1996.

[151004]

[151005]

[151006]14427.

[151007]Chaidarun, S. S.; Swearingen, B.; Alexander, J. M.: Differential ential

[151008]expression of estrogen receptor-beta (ER-beta) in human pituitary

[151009]tumors: functional interactions with ER-alpha and a tumor-specific

[151010]splice variant. J. Clin. Endocr. Metab. 83: 3308–3315, 1998.

[151011]

[151012]

[151013]14428.

[151014]Chu, S.; Mamers, P.; Burger, H. G.; Fuller, P. J.: Estrogen
receptor

[151015]isoform gene expression in ovarian stromal and epithelial
tumors. J.

[151016]Clin. Endocr. Metab. 85: 1200–1205, 2000.

[151017]

[151018]

[151019]14429.

[151020]Enmark, E.; Peltö-Huikko, M.; Grandien, K.; Lagercrantz,
S.; Lagercrantz,

[151021]J.; Fried, G.; Nordenskjöld, M.; Gustafsson, J.-A.: Human
estrogen

[151022]receptor beta-gene structure, chromosomal localization,
and expression

[151023]pattern. J. Clin. Endocr. Metab. 82: 4258–4265, 1997.

[151024]

[151025]

[151026]14430.

[151027]Esmaeli, B.; Harvey, J. T.; Hewlett, B.: Immunohistochemical evidence

[151028]for estrogen receptors in meibomian glands. Ophthalmology 107: 180–184,

[151029]2000.

[151030]

[151031]

[151032]14431.

[151033]Forsell, C.; Enmark, E.; Axelman, K.; Blomberg, M.; Wahlund, L.-O.;

[151034]Gustafsson, J.-A.; Lannfelt, L.: Investigations of a CA repeat in

[151035]the oestrogen receptor beta gene in patients with Alzheimer's disease. Europ.

[151036]J. Hum. Genet. 9: 802–804, 2001.

[151037]

[151038]

[151039]14432.

[151040]Korach, K. S.; Couse, J. F.; Curtis, S. W.; Washburn, T. F.; Lindzey,

[151041]J.; Kimbro, K. S.; Eddy, E. M.; Migliaccio, S.; Snedeker, S. M.; Lubahn,

[151042]D. B.; Schomberg, D. W.; Smith, E. P.: Estrogen receptor gene disruption:

[151043]molecular characterization and experimental and clinical phenotypes. Recent

[151044]Prog. Horm. Res. 51: 159–188, 1996.

[151045]

[151046]

[151047]14433.

[151048]Krege, J. H.; Hodgin, J. B.; Couse, J. F.; Enmark, E.; Warner,

[151049]M.; Mahler, J. F.; Sar, M.; Korach, K. S.; Gustafsson, J.-A.; Smithies,

[151050]O.: Generation and reproductive phenotypes of mice lacking estrogen

[151051]receptor beta. Proc. Nat. Acad. Sci. 95: 15677–15682, 1998.

[151052]

[151053]

[151054]14434.

[151055]Krezel, W.; Dupont, S.; Krust, A.; Chambon, P.; Chapman,
P. F.

[151056]: Increased anxiety and synaptic plasticity in estrogen re-
ceptor beta-deficient

[151057]mice. Proc. Nat. Acad. Sci. 98: 12278–12282, 2001.

[151058]

[151059]

[151060]14435.

[151061]Kuiper, G. G. J. M.; Enmark, E.; Peltö-Huikko, M.; Nilsson,
S.;

[151062]Gustafsson, J.-A.: Cloning of a novel estrogen receptor
expressed

[151063]in rat prostate and ovary. Proc. Nat. Acad. Sci. 93:
5925–5930,

[151064]1996.

[151065]

[151066]

[151067]14436.

[151068]Mosialos, G.; Birkenbach, M.; Yalamanchili, R.; VanArsdale,
T.;

[151069]Ware, C.; Kieff, E.: The Epstein–Barr virus transforming protein

[151070]LMP1 engages signaling proteins for the tumor necrosis factor receptor

[151071]family. Cell 80: 389–399, 1995.

[151072]

[151073]

[151074]14437.

[151075]Rothe, M.; Wong, S. C.; Henzel, W. J.; Goeddel, D. V.: A novel

[151076]family of putative signal transducers associated with the cytoplasmic

[151077]domain of the 75 kDa tumor necrosis factor receptor. Cell 78: 681–692,

[151078]1994.

[151079]

[151080]

[151081]14438.

[151082]Siemienski, K.; Peters, N.; Scheurich, P.; Wajant, H.: Organization

[151083]of the human tumour necrosis factor receptor-associated
factor 1 (TRAF1)

[151084]gene and mapping to chromosome 9q33–34. Gene 195:
35–39, 1997.

[151085]

[151086]

[151087]14439.

[151088]Tsitsikov, E. N.; Laouini, D.; Dunn, I. F.; Sannikova, T. Y.;
Davidson,

[151089]L.; Alt, F. W.; Geha, R. S.: TRAF1 is a negative regulator of
TNF

[151090]signaling: enhanced TNF signaling in TRAF1-deficient
mice. Immunity 15:

[151091]647–657, 2001.

[151092]

[151093]

[151094]14440.

[151095]Jacquemin, P.; Hwang, J.-J.; Martial, J. A.; Dolle, P.; David-
son,

[151096]I.: A novel family of developmentally regulated mam-
malian transcription

[151097]factors containing the TEA/ATTS DNA binding domain. J.

Biol. Chem. 271:

[151098]21775–21785, 1996.

[151099]

[151100]

[151101]14441.

[151102]Bejjani, B. A.; Lewis, R. A.; Tomey, K. F.; Anderson, K. L.;
Dueker,

[151103]D. K.; Jabak, M.; Astle, W. F.; Otterud, B.; Leppert, M.; Lup-
ski,

[151104]J. R.: Mutations in CYP1B1, the gene for cytochrome
P4501B1, are

[151105]the predominant cause of primary congenital glaucoma in
Saudi Arabia. Am.

[151106]J. Hum. Genet. 62: 325–333, 1998.

[151107]

[151108]

[151109]14442.

[151110]Bejjani, B. A.; Stockton, D. W.; Lewis, R. A.; Tomey, K. F.;
Dueker,

[151111]D. K.; Jabak, M.; Astle, W. F.; Lupski, J. R.: Multiple CYP1B1 mutations

[151112]and incomplete penetrance in an inbred population segregating primary

[151113]congenital glaucoma suggest frequent de novo events and a dominant

[151114]modifier locus. Hum. Molec. Genet. 9: 367–374, 2000.

[151115]

[151116]

[151117]14443.

[151118]Lin, J.; Arnold, H. B.; Della-Fera, M. A.; Azain, M. J.; Hartzell,

[151119]D. L.; Baile, C. A.: Myostatin knockout in mice increases myogenesis

[151120]and decreases adipogenesis. Biochem. Biophys. Res. Commun. 291:

[151121]701–706, 2002.

[151122]

[151123]

[151124]14444.

[151125]McPherron, A. C.; Lawler, A. M.; Lee, S.-J.: Regulation of skeletal

[151126]muscle mass in mice by a new TGF-beta superfamily member. Nature 387:

[151127]83-90, 1997.

[151128]

[151129]

[151130]14445.

[151131]Robins, P.; Lindahl, T.: DNA ligase IV from HeLa cell nuclei. J.

[151132]Biol. Chem. 271: 24257-24261, 1996.

[151133]

[151134]

[151135]14446.

[151136]Roldan-Arjona, T.; Wei, Y.-F.; Carter, K. C.; Klungland, A.; Anselmino,

[151137]C.; Wang, R.-P.; Augustus, M.; Lindahl, T.: Molecular cloning and

[151138]functional expression of a human cDNA encoding the antitumor enzyme

[151139]8-hydroxyguanine-DNA glycosylase. Proc. Nat. Acad. Sci.
94: 8016-8020,
[151140]1997.

[151141]

[151142]

[151143]14447.

[151144]Asano, K.; Merrick, W. C.; Hershey, J. W. B.: The translation
[151145]initiation factor eIF3-p48 subunit is encoded by int-6, a
site of

[151146]frequent integration by the mouse mammary tumor virus
genome. J.

[151147]Biol. Chem. 272: 23477-23480, 1997.

[151148]

[151149]

[151150]14448.

[151151]Rasooly, R. S.: Personal Communication. Baltimore, Md.
2/1998.

[151152]

[151153]

[151154]14449.

[151155]Scott, A. F.: Personal Communication. Baltimore, Md.
10/3/2001.

[151156]

[151157]

[151158]14450.

[151159]Todd, S.; Naylor, S. L.: Dinucleotide repeat polymorphism
in the

[151160]human tubulin alpha 1 (testis specific) gene (TUBA1). Nu-
cleic Acids

[151161]Res. 19: 3755 only, 1991.

[151162]

[151163]

[151164]14451.

[151165]Guo, Z.; Turner, C.; Castle, D.: Relocation of the t-SNARE
SNAP-23

[151166]from lamellipodia-like cell surface projections regulates
compound

[151167]exocytosis in mast cells. Cell 94: 537-548, 1998.

[151168]

[151169]

[151170]14452.

[151171]Lazo, P. A.; Nadal, M.; Ferrer, M.; Area, E.; Hernandez-Torres,

[151172]J.; Nabokina, S. M.; Mollinedo, F.; Estivill, X.: Genomic organization,

[151173]chromosomal localization, alternative splicing, and isoforms of the

[151174]human synaptosome-associated protein-23 gene implicated in vesicle-membrane

[151175]fusion processes. Hum. Genet. 108: 211-215, 2001.

[151176]

[151177]

[151178]14453.

[151179]Mollinedo, F.; Lazo, P. A.: Identification of two isoforms of

[151180]the vesicle-membrane fusion protein SNAP-23 in human neutrophils and

[151181]HL-60 cells. Biochem. Biophys. Res. Commun. 231: 808-812, 1997.

[151182]

[151183]

[151184]14454.

[151185]Ravichandran, V.; Chawla, A.; Roche, P. A.: Identification
of

[151186]a novel syntaxin- and synaptobrevin/VAMP-binding pro-
tein, SNAP-23,

[151187]expressed in non-neuronal tissues. J. Biol. Chem. 271:
13300-13303,

[151188]1996.

[151189]

[151190]

[151191]14455.

[151192]Shukla, A.; Corydon, T. J.; Nielsen, S.; Hoffmann, H. J.;
Dahl,

[151193]R.: Identification of three new splice variants of the SNARE
protein

[151194]SNAP-23. Biochem. Biophys. Res. Commun. 285:
320-327, 2001.

[151195]

[151196]

[151197]14456.

[151198]Mugita, N.; Kumura, Y.; Ogawa, M.; Saya, H.; Nakao, M.:
Identification

[151199]of a novel, tissue-specific calpain htra-3; a human homo-
logue of the

[151200]Caenorhabditis elegans sex determination gene. Biochem.
Biophys.

[151201]Res. Commun. 239: 845–850, 1997.

[151202]

[151203]

[151204]14457.

[151205]Sharpless, N. E.; Ferguson, D. O.; O'Hagan, R. C.; Castril-
lon,

[151206]D. H.; Lee, C.; Farazi, P. A.; Alson, S.; Fleming, J.; Morton,
C.

[151207]C.; Frank, K.; Chin, L.; Alt, F. W.; DePinho, R. A.: Impaired
nonhomologous

[151208]end-joining provokes soft tissue sarcomas harboring
chromosomal translocations,

[151209]amplifications, and deletions. Molec. Cell 8: 1187–1196,
2001.

[151210]

[151211]

[151212]14458.

[151213]Bohme, B.; Holtrich, U.; Wolf, G.; Luzius, H.; Grzeschik, K.-H.;

[151214]Strebhardt, K.; Rubsamen-Waigmann, H.: PCR mediated detection of

[151215]a new human receptor-tyrosine-kinase, HEK 2. Oncogene 8: 2857-2862,

[151216]1993.

[151217]

[151218]

[151219]14459.

[151220]Ruiz, J. C.; Conlon, F. L.; Robertson, E. J.: Identification of

[151221]novel protein kinases expressed in the myocardium of the developing

[151222]mouse heart. Mech. Dev. 48: 153-164, 1994.

[151223]

[151224]

[151225]14460.

[151226]Ashbourne, K. J.; Byth, B. C.; Meijers, J. C. M.; Cox, D. W.:

[151227]Polymorphism of the protein C inhibitor (PCI) gene on
chromosome 14. Hum.

[151228]Molec. Genet. 2: 92 only, 1993.

[151229]

[151230]

[151231]14461.

[151232]Sadler, J. E.: Combined factors V and VIII deficiency climbs
onto

[151233]the map. (Editorial) J. Clin. Invest. 99: 555–556, 1997.

[151234]

[151235]

[151236]14462.

[151237]Yasuda, T.; Nadano, D.; Iida, R.; Tanaka, Y.; Nakanaga, M.;
Kishi,

[151238]K.: Discovery of a genetic polymorphism of human plasma
protein C

[151239]inhibitor (PCI): genetic survey utilizing isoelectric focusing

followed

[151240]by immunoblotting, immunological and biochemical characterization. Hum.

[151241]Genet. 89: 265–269, 1992.

[151242]

[151243]

[151244]14463.

[151245]Bui, T. D.; Rankin, J.; Smith, K.; Huguet, E. L.; Ruben, S.; Strachan,

[151246]T.; Harris, A. L.; Lindsay, S.: A novel human Wnt gene, WNT10B, maps

[151247]to 12q13 and is expressed in human breast carcinomas.

Oncogene 14:

[151248]1249–1253, 1997.

[151249]

[151250]

[151251]14464.

[151252]Hardiman, G.; Kastelein, R. A.; Bazan, J. F.: Isolation, characterization

[151253]and chromosomal localization of human WNT10B. Cyto–

genet. Cell Genet. 77:

[151254]278–282, 1997.

[151255]

[151256]

[151257]14465.

[151258]Ross, S. E.; Hemati, N.; Longo, K. A.; Bennett, C. N.; Lucas,
P.

[151259]C.; Erickson, R. L.; MacDougald, O. A.: Inhibition of adipo-
genesis

[151260]by Wnt signaling. Science 289: 950–953, 2000.

[151261]

[151262]

[151263]14466.

[151264]Meyerhardt, J. A.; Look, A. T.; Bigner, S. H.; Fearon, E. R.:

[151265]Identification and characterization of neogenin, a DCC-
related gene. Oncogene 14:

[151266]1129–1136, 1997.

[151267]

[151268]

[151269]14467.

[151270]Vielmetter, J.; Chen, X.-N.; Miskevich, F.; Lane, R. P.; Yamakawa,

[151271]K.; Korenberg, J. R.; Dreyer, W. J.: Molecular characterization of

[151272]human neogenin, a DCC-related protein, and the mapping of its gene

[151273](NEO1) to chromosomal position 15q22.3-q23. Genomics 41: 414-421,

[151274]1997.

[151275]

[151276]

[151277]14468.

[151278]Vielmetter, J.; Kayyem, J. F.; Roman, J. M.; Dreyer, W. J.: Neogenin,

[151279]an avian cell surface protein expressed during terminal neuronal differentiation,

[151280]is closely related to the human tumor suppressor molecule deleted

[151281]in colorectal cancer. J. Cell Biol. 127: 2009-2020, 1994.

[151282]

[151283]

[151284]14469.

[151285]Rosenquist, T. A.; Zharkov, D. O.; Grollman, A. P.: Cloning
and

[151286]characterization of a mammalian 8-oxoguanine DNA gly-
cosylase. Proc.

[151287]Nat. Acad. Sci. 94: 7429-7434, 1997.

[151288]

[151289]

[151290]14470.

[151291]Tani, M.; Shinmura, K.; Kohno, T.; Shiroishi, T.; Wakama,
S.;

[151292]Kim, S.-R.; Nohmi, T.; Kasai, H.; Takenoshita, S.; Naga-
machi, Y.;

[151293]Yokota, J.: Genomic structure and chromosomal localiza-
tion of the

[151294]mouse Ogg1 gene that is involved in the repair of
8-hydroxyguanine

[151295]in DNA damage. Mammalian Genome 9: 32-37, 1998.

[151296]

[151297]

[151298]14471.

[151299]Lim, H. N.; Hawkins, J. R.; Hughes, I. A.: Genetic evidence
to

[151300]exclude the androgen receptor co-factor, ARA70 (NCOA4)
as a candidate

[151301]gene for the causation of undermasculinised genitalia.
(Letter) Clin.

[151302]Genet. 59: 284–286, 2001.

[151303]

[151304]

[151305]14472.

[151306]Santoro, M.; Dathan, N. A.; Berlingieri, M. T.; Bongarzone,
I.;

[151307]Paulin, C.; Grieco, M.; Pierotti, M. A.; Vecchio, G.; Fusco,
A.:

[151308]Molecular characterization of RET/PTC3: a novel rear-
ranged version

[151309]of the RET proto-oncogene in a human thyroid papillary
carcinoma. Oncogene 9:

[151310]509–516, 1994.

[151311]

[151312]

[151313]14473.

[151314]Yeh, S.; Chang, C.: Cloning and characterization of a specific
cific

[151315]coactivator, ARA-70, for the androgen receptor in human
prostate cells. Proc.

[151316]Nat. Acad. Sci. 93: 5517-5521, 1996.

[151317]

[151318]

[151319]14474.

[151320]Donghi, R.; Sozzi, G.; Pierotti, M. A.; Biunno, I.; Miozzo,
M.;

[151321]Fusco, A.; Grieco, M.; Santoro, M.; Vecchio, G.; Spurr, N.
K.; Della

[151322]Porta, G.: The oncogene associated with human papillary
thyroid carcinoma

[151323](PTC) is assigned to chromosome 10 q11-q12 in the same
region as multiple

[151324]endocrine neoplasia type 2A (MEN2A). Oncogene 4:
521-523, 1989.

[151325]

[151326]

[151327]14475.

[151328]Nikiforova, M. N.; Stringer, J. R.; Blough, R.; Medvedovic,
M.;

[151329]Fagin, J. A.; Nikiforov, Y. E.: Proximity of chromosomal loci
that

[151330]participate in radiation-induced rearrangements in human
cells. Science 290:

[151331]138–141, 2000.

[151332]

[151333]

[151334]14476.

[151335]Sozzi, G.; Pierotti, M. A.; Miozzo, M.; Donghi, R.; Radice,
P.;

[151336]De Benedetti, V.; Grieco, M.; Santoro, M.; Fusco, A.; Vec-
chio, G.;

[151337]Mathew, C. G. P.; Ponder, B. A. J.; Spurr, N. K.: Refined lo-
calization

[151338]to contiguous regions on chromosome 10q of the two
genes (H4 and RET)

[151339]that form the oncogenic sequence PTC. Oncogene 6:
339–342, 1991.

[151340]

[151341]

[151342]14477.

[151343]Tong, Q.; Li, Y.; Smanik, P. A.; Fithian, L. J.; Xing, S.; Maz-
zaferri,

[151344]E. L.; Jhiang, S. M.: Characterization of the promoter re-
gion and

[151345]oligomerization domain of H4 (D10S170), a gene fre-
quently rearranged

[151346]with the ret proto-oncogene. Oncogene 10: 1781–1787,
1995.

[151347]

[151348]

[151349]14478.

[151350]Araki, T.; Milbrandt, J.: Ninjurin, a novel adhesion
molecule,

[151351]is induced by nerve injury and promotes axonal growth.

Neuron 17:

[151352]353–361, 1996.

[151353]

[151354]

[151355]14479.

[151356]Araki, T.; Zimonjic, D. B.; Popescu, N. C.; Milbrandt, J.:

Mechanism

[151357]of homophilic binding mediated by ninjurin, a novel
widely expressed

[151358]adhesion molecule. J. Biol. Chem. 272: 21373–21380,
1997.

[151359]

[151360]

[151361]14480.

[151362]Chadwick, B. P.; Heath, S. K.; Williamson, J.; Obermayr, F.;
Patel,

[151363]L.; Sheer, D.; Frischauf, A.–M.: The human homologue of
the ninjurin

[151364]gene maps to the candidate region of hereditary sensory
neuropathy

[151365]type I (HSN1). Genomics 47: 58–63, 1998.

[151366]

[151367]

[151368]14481.

[151369]Mandich, P.; Bellone, E.; Di Maria, E.; Pigullo, S.; Pizzuti, A.;

[151370]Schenone, A.; Soriani, S.; Varese, A.; Windebank, A. J.; Ajmar, F.

[151371]: Exclusion of the ninjurin gene as a candidate for hereditary sensory

[151372]neuropathies type I and type II. Am. J. Med. Genet. 83: 409-410,

[151373]1999.

[151374]

[151375]

[151376]14482.

[151377]McAllister, G.; Whiting, P.; Hammond, E. A.; Knowles, M. R.; Atack,

[151378]J. R.; Bailey, F. J.; Maigetter, R.; Ragan, C. I.: cDNA cloning of

[151379]human and rat brain myo-inositol monophosphatase: expression and characterization

[151380]of the human recombinant enzyme. Biochem. J. 284:

749–754, 1992.

[151381]

[151382]

[151383]14483.

[151384]Sjoholt, G.; Molven, A.; Lovlie, R.; Wilcox, A.; Sikela, J. M.;

[151385]Steen, V. M.: Genomic structure and chromosomal localization of a

[151386]human myo-inositol monophosphatase gene (IMPA). Genomics 45: 113–122,

[151387]1997.

[151388]

[151389]

[151390]14484.

[151391]Steen, V. M.; Gulbrandsen, A. K.; Eiken, H. G.; Berle, J. O.:

[151392]Lack of genetic variation in the coding region of the myo-inositol

[151393]monophosphatase gene in lithium-treated patients with manic depressive

[151394]illness. Pharmacogenetics 6: 113–116, 1996.

[151395]

[151396]

[151397]14485.

[151398]McDonald, M. T.; Flejter, W.; Sheldon, S.; Putzi, M. J.;
Gorski,

[151399]J. L.: XY sex reversal and gonadal dysgenesis due to 9p24
monosomy. Am.

[151400]J. Med. Genet. 73: 321–326, 1997.

[151401]

[151402]

[151403]14486.

[151404]Muroya, K.; Okuyama, T.; Goishi, K.; Ogiso, Y.; Fukuda, S.;
Kameyama,

[151405]J.; Sato, H.; Suzuki, Y.; Terasaki, H.; Gomyo, H.; Wakui, K.;
Fukushima,

[151406]Y.; Ogata, T.: Sex-determining gene(s) on distal 9p: clinical and

[151407]molecular studies in six cases. J. Clin. Endocr. Metab. 85:
3094–3100,

[151408]2000.

[151409]

[151410]

[151411]14487.

[151412]Raymond, C. S.; Shamu, C. E.; Shen, M. M.; Seifert, K. J.;
Hirsch,

[151413]B.; Hodgkin, J.; Zarkower, D.: Evidence for evolutionary
conservation

[151414]of sex-determining genes. Nature 391: 691–695, 1998.

[151415]

[151416]

[151417]14488.

[151418]Shan, Z.; Zabel, B.; Trautmann, U.; Hillig, U.; Ottolenghi,
C.;

[151419]Wang, Y.; Haaf, T.: FISH mapping of the sex-reversal re-
gion on human

[151420]chromosome 9p in two XY females and in primates. Europ.
J. Hum. Genet. 8:

[151421]167–173, 2000.

[151422]

[151423]

[151424]14489.

[151425]Shen, M. M.; Hodgkin, J.: mab-3, a gene required for sex-specific

[151426]yolk protein expression and a male-specific lineage in *C. elegans*. *Cell* 54:

[151427]1019-1031, 1988.

[151428]

[151429]

[151430]14490.

[151431]Smith, C. A.; McClive, P. J.; Western, P. S.; Reed, K. J.; Sinclair,

[151432]A. H.: Conservation of a sex-determining gene. (Letter) *Nature* 402:

[151433]601-602, 1999.

[151434]

[151435]

[151436]14491.

[151437]Veitia, R.; Nunes, M.; Brauner, R.; Doco-Fenzy, M.; Joanny-Flinois,

[151438]O.; Jaubert, F.; Lortat-Jacob, S.; Fellous, M.; McElreavey,

K.: Deletions

[151439]of distal 9p associated with 46,XY male to female sex reversal: definition

[151440]of the breakpoints at 9p23.3–p24.1. Genomics 41: 271–274, 1997.

[151441]

[151442]

[151443]14492.

[151444]Takekawa, M.; Posas, F.; Saito, H.: A human homolog of the yeast

[151445]Ssk2/Ssk22 MAP kinase kinase kinases, MTK1, mediates stress-induced

[151446]activation of the p38 and JNK pathways. EMBO J. 16: 4973–4982, 1997.

[151447]

[151448]

[151449]14493.

[151450]Chapman, D. L.; Agulnik, I.; Hancock, S.; Silver, L. M.; Papaioannou,

[151451]V. E.: Tbx6, a mouse T-box gene implicated in paraxial mesoderm formation

[151452]at gastrulation. Dev. Biol. 180: 534–542, 1996.

[151453]

[151454]

[151455]14494.

[151456]Chapman, D. L.; Papaioannou, V. E.: Three neural tubes in
mouse

[151457]embryos with mutations in the T-box gene Tbx6. Nature
391: 695–697,

[151458]1998.

[151459]

[151460]

[151461]14495.

[151462]Papapetrou, C.; Putt, W.; Fox, M.; Edwards, Y. H.: The hu-
man TBX6

[151463]gene: cloning and assignment to chromosome 16p11.2.
Genomics 55:

[151464]238–241, 1999.

[151465]

[151466]

[151467]14496.

[151468]Spicer, A. P.; Olson, J. S.; McDonald, J. A.: Molecular
cloning

[151469]and characterization of a cDNA encoding the third puta-
tive mammalian

[151470]hyaluronan synthase. J. Biol. Chem. 272: 8957–8961,
1997.

[151471]

[151472]

[151473]14497.

[151474]Bagri, A.; Marin, O.; Plump, A. S.; Mak, J.; Pleasure, S. J.;
Rubenstein,

[151475]J. L. R.; Tessier–Lavigne, M.: Slit proteins prevent midline
crossing

[151476]and determine the dorsoventral position of major axonal
pathways in

[151477]the mammalian forebrain. Neuron 33: 233–248, 2002.

[151478]

[151479]

[151480]14498.

[151481]Dallol, A.; Forgacs, E.; Martinez, A.; Sekido, Y.; Walker, R.;

[151482]Kishida, T.; Rabbitts, P.; Maher, E. R.; Minna, J. D.; Latif, F.:

[151483]Tumour specific promoter region methylation of the human homologue

[151484]of the Drosophila Roundabout gene DUTT1 (ROBO1) in human cancers. *Oncogene* 21:

[151485]3020–3028, 2002.

[151486]

[151487]

[151488]14499.

[151489]Kidd, T.; Brose, K.; Mitchell, K. J.; Fetter, R. D.; Tessier-Lavigne,

[151490]M.; Goodman, C. S.; Tear, G.: Roundabout controls axon crossing of

[151491]the CNS midline and defines a novel subfamily of evolutionarily conserved

[151492]guidance receptors. *Cell* 92: 205–215, 1998.

[151493]

[151494]

[151495]14500.

[151496]Sundaresan, V.; Chung, G.; Heppell–Parton, A.; Xiong, J.;
Grundy,

[151497]C.; Roberts, I.; James, L.; Cahn, A.; Bench, A.; Douglas, J.;
Minna,

[151498]J.; Sekido, Y.; Lerman, M.; Latif, F.; Bergh, J.; Li, H.; Lowe,
N.;

[151499]Ogilvie, D.; Rabbitts, P.: Homozygous deletions at 3p12 in
breast

[151500]and lung cancer. *Oncogene* 17: 1723–1729, 1998.

[151501]

[151502]

[151503]14501.

[151504]Zallen, J. A.; Yi, B. A.; Bargmann, C. I.: The conserved im-
munoglobulin

[151505]superfamily member SAX–3/Robo directs multiple aspects
of axon guidance

[151506]in *C. elegans*. *Cell* 92: 217–227, 1998.

[151507]

[151508]

[151509]14502.

[151510]Fischer, U.; Heckel, D.; Michel, A.; Janka, M.; Hulsebos, T.;
Meese,
[151511]E.: Cloning of a novel transcription factor-like gene am-
plified in
[151512]human glioma including astrocytoma grade I. Hum. Molec.
Genet. 6:
[151513]1817–1822, 1997.

[151514]

[151515]

[151516]14503.

[151517]Gracia, E.; Fischer, U.; Elkahloun, A.; Trent, J. M.; Meese,
E.;

[151518]Meltzer, P. S.: Isolation of genes amplified in human can-
cers by
[151519]microdissection mediated cDNA capture. Hum. Molec.
Genet. 5: 595–600,
[151520]1996.

[151521]

[151522]

[151523]14504.

[151524]Gerard, M.; Hernandez, L.; Wevrick, R.; Stewart, C. L.: Dis-
ruption

[151525]of the mouse necdin gene results in early post-natal
lethality. Nature

[151526]Genet. 23: 199–202, 1999.

[151527]

[151528]

[151529]14505.

[151530]Hurst, L. D.; McVean, G.; Moore, T.: Imprinted genes have
few

[151531]and small introns. (Letter) Nature Genet. 12: 234–237,
1996.

[151532]

[151533]

[151534]14506.

[151535]Jay, P.; Rougeulle, C.; Massacrier, A.; Moncla, A.; Mattei,
M.–G.;

[151536]Malzac, P.; Roeckel, N.; Taviaux, S.; Lefranc, J.–L. B.; Cau,
P.;

[151537]Berta, P.; Lalande, M.; Muscatelli, F.: The human necdin
gene, NDN,

[151538]is maternally imprinted and located in the Prader–Willi
syndrome chromosomal

[151539]region. Nature Genet. 17: 357–360, 1997.

[151540]

[151541]

[151542]14507.

[151543]MacDonald, H. R.; Wevrick, R.: The necdin gene is deleted
in Prader–Willi

[151544]syndrome and is imprinted in human and mouse. Hum.
Molec. Genet. 6:

[151545]1873–1878, 1997.

[151546]

[151547]

[151548]14508.

[151549]Maruyama, K.; Usami, M.; Aizawa, T.; Yoshikawa, K.: A
novel brain–specific

[151550]mRNA encoding nuclear protein (necdin) expressed in
neurally differentiated

[151551]embryonal carcinoma cells. Biochem. Biophys. Res. Comm.
178: 291–296,

[151552]1991.

[151553]

[151554]

[151555]14509.

[151556]Muscatelli, F.; Arous, D. N.; Massacrier, A.; Boccaccio, I.;

Le

[151557]Moal, M.; Cau, P.; Cremer, H.: Disruption of the mouse

Necdin gene

[151558]results in hypothalamic and behavioral alterations remi-

niscent of

[151559]the human Prader-Willi syndrome. Hum. Molec. Genet. 9:

3101-3110,

[151560]2000.

[151561]

[151562]

[151563]14510.

[151564]Nakada, Y.; Taniura, H.; Uetsuki, T.; Inazawa, J.;

Yoshikawa, K.

[151565]: The human chromosomal gene for necdin, a neuronal

growth suppressor,

[151566]in the Prader–Willi syndrome deletion region. Gene 213:
65–72, 1998.

[151567]

[151568]

[151569]14511.

[151570]Nicholls, R. D.: Incriminating gene suspects, Prader–Willi
style. Nature

[151571]Genet. 23: 132–134, 1999.

[151572]

[151573]

[151574]14512.

[151575]Tsai, T.–F.; Armstrong, D.; Beaudet, A. L.: Necdin–defi-
cient mice

[151576]do not show lethality or the obesity and infertility of
Prader–Willi

[151577]syndrome. (Letter) Nature Genet. 22: 15–16, 1999.

[151578]

[151579]

[151580]14513.

[151581]Watrin, F.; Roeckel, N.; Lacroix, L.; Mignon, C.; Mattei, M.-G.;

[151582]Disteche, C.; Muscatelli, F.: The mouse necdin gene is expressed

[151583]from the paternal allele only and lies in the 7C region of the mouse

[151584]chromosome 7, a region of conserved synteny to the human Prader-Willi

[151585]syndrome region. *Europ. J. Hum. Genet.* 5: 324-332, 1997.

[151586]

[151587]

[151588]14514.

[151589]Delmas, V.; Stokes, D. G.; Perry, R. P.: A mammalian DNA-binding

[151590]protein that contains a chromodomain and an SNF2/SWI2-like helicase

[151591]domain. *Proc. Nat. Acad. Sci.* 90: 2414-2418, 1993.

[151592]

[151593]

[151594]14515.

[151595]Woodage, T.; Basrai, M. A.; Baxevanis, A. D.; Hieter, P.;
Collins,

[151596]F. S.: Characterization of the CHD family of proteins. Proc.
Nat.

[151597]Acad. Sci. 94: 11472–11477, 1997.

[151598]

[151599]

[151600]14516.

[151601]Ge, Q.; Nilasena, D. S.; O'Brien, C. A.; Frank, M. B.; Targoff,

[151602]I. N.: Molecular analysis of a major antigenic region of the
240-kD

[151603]protein of Mi-2 autoantigen. J. Clin. Invest. 96:
1730–1737, 1995.

[151604]

[151605]

[151606]14517.

[151607]Seelig, H. P.; Renz, M.; Targoff, I. N.; Ge, Q.; Frank, M. B.:

[151608]Two forms of the major antigenic protein of the dermato-
myositis-specific

[151609]Mi-2 autoantigen. (Letter) Arthritis Rheum. 39:

1769–1771, 1996.

[151610]

[151611]

[151612]14518.

[151613]Miyazaki, S.; Imatani, A.; Ballard, L.; Marchetti, A.; Buttitta,

[151614]F.; Albertsen, H.; Nevanlinna, H. A.; Gallahan, D.; Callahan,
R.:

[151615]The chromosome location of the human homolog of the
mouse mammary

[151616]tumor-associated gene INT6 and its status in human
breast carcinomas. Genomics 46:

[151617]155–158, 1997.

[151618]

[151619]

[151620]14519.

[151621]Hu, G.; Chung, Y.–L.; Glover, T.; Valentine, V.; Look, A T.;
Fearon,

[151622]E. R.: Characterization of human homologs of the
Drosophila seven

[151623]in absentia (sina) gene. Genomics 46: 103–111, 1997.

[151624]

[151625]

[151626]14520.

[151627]Liu, J.; Stevens, J.; Rote, C. A.; Yost, H. J.; Hu, Y.; Neufeld,

[151628]K. L.; White, R. L.; Matsunami, N.: Siah-1 mediates a novel
beta-catenin

[151629]degradation pathway linking p53 to the adenomatous
polyposis coli

[151630]protein. Molec. Cell 7: 927-936, 2001.

[151631]

[151632]

[151633]14521.

[151634]Nemani, M.; Linares-Cruz, G.; Bruzzoni-Giovanelli, H.;
Roperch,

[151635]J.-P.; Tuynder, M.; Bougueleret, L.; Cherif, D.; Medhioub,
M.; Pasturaud,

[151636]P.; Alvaro, V.; der Sarkissan, H.; Cazes, L.; Le Paslier, D.;
Le Gall,

[151637]I.; Israeli, D.; Dausset, J.; Sigaux, F.; Chumakov, I.; Oren,
M.;

[151638]Calvo, F.; Amson, R. B.; Cohen, D.; Telerman, A.: Activation of the

[151639]human homologue of the Drosophila SINA gene in apoptosis and tumor

[151640]suppression. Proc. Nat. Acad. Sci. 93: 9039–9042, 1996.

[151641]

[151642]

[151643]14522.

[151644]Ozkaynak, E.; Schnegelsberg, P. N. J.; Jin, D. F.; Clifford, G.

[151645]M.; Warren, F. D.; Drier, E. A.; Oppermann, H.: Osteogenic protein-2:

[151646]a new member of the transforming growth factor-beta superfamily expressed

[151647]in early embryogenesis. J. Biol. Chem. 267: 25220–25227, 1992.

[151648]

[151649]

[151650]14523.

[151651]Jang, W.; Weber, J. S.; Harkins, E. B.; Meisler, M. H.: Localization

[151652]of the rhotekin gene RTKN on the physical maps of mouse chromosome

[151653]6 and human chromosome 2p13 and exclusion as a candidate for mnd2

[151654]and LGMD2B. Genomics 40: 506–507, 1997.

[151655]

[151656]

[151657]14524.

[151658]Reid, T.; Furuyashiki, T.; Ishizaki, T.; Watanabe, G.; Watanabe,

[151659]N.; Fujisawa, K.; Morii, N.; Madaule, P.; Narumiya, S.: Rhotekin,

[151660]a new putative target for Rho bearing homology to a serine/threonine

[151661]kinase, PKN, and rhophilin in the Rho-binding domain. J. Biol. Chem. 271:

[151662]13556–13560, 1996.

[151663]

[151664]

[151665]14525.

[151666]Janssen, U.; Davis, E. M.; Le Beau, M. M.; Stoffel, W.: Human
man

[151667]mitochondrial enoyl-CoA hydratase gene (ECHS1): structural organization

[151668]and assignment to chromosome 10q26.2-q26.3. Genomics 40: 470-475,

[151669]1997.

[151670]

[151671]

[151672]14526.

[151673]Kanazawa, M.; Ohtake, A.; Abe, H.; Yamamoto, S.; Satoh, Y.; Takayanagi,

[151674]M.; Niimi, H.; Mori, M.; Hashimoto, T.: Molecular cloning and sequence

[151675]analysis of the cDNA for human mitochondrial short-chain enoyl-CoA

[151676]hydratase. Enzyme Protein 47: 9-13, 1993.

[151677]

[151678]

[151679]14527.

[151680]Bingle, C. D.; Gowan, S.: Molecular cloning of the forkhead transcription

[151681]factor HNF-3-alpha from a human pulmonary adenocarcinoma cell line. Biochim.

[151682]Biophys. Acta 1307: 17-20, 1996.

[151683]

[151684]

[151685]14528.

[151686]Hannigan, G. E.; Bayani, J.; Weksberg, R.; Beatty, B.; Pandita,

[151687]A.; Dedhar, S.; Squire, J.: Mapping of the gene encoding the integrin-linked

[151688]kinase, ILK, to human chromosome 11p15.5-p15.4. Genomics 42: 177-179,

[151689]1997.

[151690]

[151691]

[151692]14529.

[151693]Hannigan, G. E.; Leung-Hagesteijn, C.; Fitz-Gibbon, L.; Coppolino,

[151694]M. G.; Radeva, G.; Filmus, J.; Bell, J. C.; Dedhar, S.: Regula-
tion

[151695]of cell adhesion and anchorage-dependent growth by a
new beta-1-integrin-linked

[151696]protein kinase. Nature 379: 91-96, 1996.

[151697]

[151698]

[151699]14530.

[151700]Omeis, I. A.; Hsu, Y.-C.; Perin, M. S.: Mouse and human
neuronal

[151701]pentraxin 1 (NPXT1): conservation, genomic structure, and
chromosomal

[151702]localization. Genomics 36: 543-545, 1996.

[151703]

[151704]

[151705]14531.

[151706]Babic, A. M.; Kireeva, M. L.; Kolesnikova, T. V.; Lau, L. F.:

[151707]CYR61, a product of a growth factor-inducible immediate
early gene,

[151708]promotes angiogenesis and tumor growth. Proc. Nat.

Acad. Sci. 95:

[151709]6355–6360, 1998.

[151710]

[151711]

[151712]14532.

[151713]Jay, P.; Berge–Lefranc, J. L.; Marsollier, C.; Mejean, C.;
Taviaux,

[151714]S.; Berta, P.: The human growth factor–inducible immediate early

[151715]gene, CYR61, maps to chromosome 1p. Oncogene 14:
1753–1757, 1997.

[151716]

[151717]

[151718]14533.

[151719]Kireeva, M. L.; Lam, S. C.–T.; Lau, L. F.: Adhesion of human umbilical

[151720]vein endothelial cells to the immediate–early gene product
Cyr61 is

[151721]mediated through integrin alpha(V)beta(3). J. Biol. Chem.
273: 3090–3096,

[151722]1998.

[151723]

[151724]

[151725]14534.

[151726]Martinerie, C.; Viegas-Pequignot, E.; Nguyen, V. C.; Perbal, B.

[151727]: Chromosomal mapping and expression of the human
cyr61 gene in tumour

[151728]cells from the nervous system. J. Clin. Path. 50: 310–316,
1997.

[151729]

[151730]

[151731]14535.

[151732]Sampath, D.; Zhu, Y.; Winneker, R. C.; Zhang, Z.: Aberrant
expression

[151733]of Cyr61, a member of the CCN

(CTGF/Cyr61/Cef10/NOVH) family, and

[151734]dysregulation by 17-beta-estradiol and basic fibroblast
growth factor

[151735]in human uterine leiomyomas. J. Clin. Endocr. Metab. 86:
1707–1715,

[151736]2001.

[151737]

[151738]

[151739]14536.

[151740]Furuhata, T.; Tokino, T.; Urano, T.; Nakamura, Y.: Isolation
of

[151741]a novel GPI-anchored gene specifically regulated by p53;
correlation

[151742]between its expression and anti-cancer drug sensitivity.

Oncogene 13:

[151743]1965-1970, 1996.

[151744]

[151745]

[151746]14537.

[151747]Kimura, Y.; Furuhata, T.; Urano, T.; Hirata, K.; Nakamura,
Y.;

[151748]Tokino, T.: Genomic structure and chromosomal localization
of GML

[151749](GPI-anchored molecule-like protein), a gene induced by
p53. Genomics 41:

[151750]477-480, 1997.

[151751]

[151752]

[151753]14538.

[151754]Fujino, T.; Kang, M.-J.; Suzuki, H.; Iijima, H.; Yamamoto, T.:

[151755]Molecular characterization and expression of rat acyl-CoA synthetase

[151756]3. J. Biol. Chem. 271: 16748–16752, 1996.

[151757]

[151758]

[151759]14539.

[151760]Minekura, H.; Fujino, T.; Kang, M.-J.; Fujita, T.; Endo, Y.; Yamamoto,

[151761]T. T.: Human acyl-coenzyme A synthetase 3 cDNA and localization of

[151762]its gene (ACS3) to chromosome band 2q34–q35. Genomics 42: 180–181,

[151763]1997.

[151764]

[151765]

[151766]14540.

[151767]Kubota, N.; Terauchi, Y.; Miki, H.; Tamemoto, H.; Yamauchi, T.;

[151768]Komeda, K.; Satoh, S.; Nakano, R.; Ishii, C.; Sugiyama, T.; Eto, K.;

[151769]Tsubamoto, Y.; and 17 others: PPAR-gamma mediates high-fat diet-induced

[151770]adipocyte hypertrophy and insulin resistance. Molec. Cell 4: 597-609,

[151771]1999.

[151772]

[151773]

[151774]14541.

[151775]Lapsys, N. M.; Kriketos, A. D.; Lim-Fraser, M.; Poynten, A. M.;

[151776]Lowy, A.; Furler, S. M.; Chisholm, D. J.; Cooney, G. J.: Expression

[151777]of genes involved in lipid metabolism correlate with peroxisome proliferator-activated

[151778]receptor gamma expression in human skeletal muscle. J. Clin. Endocr.

[151779]Metab. 85: 4293–4297, 2000.

[151780]

[151781]

[151782]14542.

[151783]Lehmann, J. M.; Moore, L. B.; Smith–Oliver, T. A.; Wilkison,
W.

[151784]O.; Willson, T. M.; Kliewer, S. A.: An antidiabetic thiazo-
lidinedione

[151785]is a high affinity ligand for peroxisome proliferator–acti-
vated receptor

[151786]gamma (PPAR gamma). J. Biol. Chem. 270: 12953–12956,
1995.

[151787]

[151788]

[151789]14543.

[151790]Lowell, B. B.: PPAR–gamma: an essential regulator of adi-
pogenesis

[151791]and modulator of fat cell function. Cell 99: 239–242,
1999.

[151792]

[151793]

[151794]14544.

[151795]Martin, G.; Schoonjans, K.; Staels, B.; Auwerx, J.: PPAR-
gamma

[151796]activators improve glucose homeostasis by stimulating
fatty acid uptake

[151797]in the adipocytes. *Atherosclerosis* 137: S75-S80, 1998.

[151798]

[151799]

[151800]14545.

[151801]Meirhaeghe, A.; Fajas, L.; Helbecque, N.; Cottel, D.; Lebel,
P.;

[151802]Dallongeville, J.; Deeb, S.; Auwerx, J.; Amouyel, P.: A ge-
netic polymorphism

[151803]of the peroxisome proliferator-activated receptor gamma
gene influences

[151804]plasma leptin levels in obese tumors. *Hum. Molec. Genet.*
7: 435-440,

[151805]1998.

[151806]

[151807]

[151808]14546.

[151809]Miles, P. D. G.; Barak, Y.; He, W.; Evans, R. M.; Olefsky, J.

[151810]M.: Improved insulin-sensitivity in mice heterozygous for
PPAR-gamma

[151811]deficiency. J. Clin. Invest. 105: 287-292, 2000.

[151812]

[151813]

[151814]14547.

[151815]Mueller, E.; Sarraf, P.; Tontonoz, P.; Evans, R. M.; Martin,
K.

[151816]J.; Zhang, M.; Fletcher, C.; Singer, S.; Spiegelman, B. M.:
Terminal

[151817]differentiation of human breast cancer through PPAR-
gamma. Molec.

[151818]Cell. 1: 465-470, 1998.

[151819]

[151820]

[151821]14548.

[151822]Mueller, E.; Smith, M.; Sarraf, P.; Kroll, T.; Aiyer, A.; Kaufman,

[151823]D. S.; Oh, W.; Demetri, G.; Figg, W. D.; Zhou, X.-P.; Eng, C.; Spiegelman,

[151824]B. M.; Kantoff, P. W.: Effects of ligand activation of peroxisome

[151825]proliferator-activated receptor gamma in human prostate cancer. Proc.

[151826]Nat. Acad. Sci. 97: 10990-10995, 2000.

[151827]

[151828]

[151829]14549.

[151830]Nagy, L.; Tontonoz, P.; Alvarez, J. G. A.; Chen, H.; Evans, R.

[151831]M.: Oxidized LDL regulates macrophage gene expression through ligand

[151832]activation of PPAR-gamma. Cell 93: 229-240, 1998.

[151833]

[151834]

[151835]14550.

[151836]Oh, E. Y.; Min, K. M.; Chung, J. H.; Min, Y.-K.; Lee, M.-S.;
Kim,

[151837]K.-W.; Lee, M.-K.: Significance of pro12ala mutation in
peroxisome

[151838]proliferator-activated receptor-gamma2 in Korean dia-
betic and obese

[151839]subjects. J. Clin. Endocr. Metab. 85: 1801-1804, 2000.

[151840]

[151841]

[151842]14551.

[151843]Pritchard, J. K.; Rosenberg, N. A.: Use of unlinked genetic
markers

[151844]to detect population stratification in association studies.
Am. J.

[151845]Hum. Genet. 65: 220-228, 1999.

[151846]

[151847]

[151848]14552.

[151849]Ristow, M.; Muller-Wieland, D.; Pfeiffer, A.; Krone, W.;
Kahn,

[151850]C. R.: Obesity associated with a mutation in a genetic regulator

[151851]of adipocyte differentiation. New Eng. J. Med. 339: 953–959, 1998.

[151852]

[151853]

[151854]14553.

[151855]Rocchi, S.; Picard, F.; Vamecq, J.; Gelman, L.; Potier, N.; Zeyer,

[151856]D.; Dubuquoy, L.; Bac, P.; Champy, M.–F.; Plunket, K. D.; Leesnitzer,

[151857]L. M.; Blanchard, S. G.; Desreumaux, P.; Moras, D.; Renaud, J.–P.;

[151858]Auwerx, J.: A unique PPAR–gamma ligand with potent insulin–sensitizing

[151859]yet weak adipogenic activity. Molec. Cell 8: 737–747, 2001.

[151860]

[151861]

[151862]14554.

[151863]Rosen, E. D.; Sarraf, P.; Troy, A. E.; Bradwin, G.; Moore, K.;

[151864]Milstone, D. S.; Spiegelman, B. M.; Mortensen, R. M.:

PPAR-gamma

[151865]is required for the differentiation of adipose tissue in vivo
and

[151866]in vitro. Molec. Cell 4: 611-617, 1999.

[151867]

[151868]

[151869]14555.

[151870]Ricote, M.; Huang, J.; Fajas, L.; Li, A.; Welch, J.; Najib, J.;

[151871]Witztum, J. L.; Auwerx, J.; Palinski, W.; Glass, C. K.: Ex-
pression

[151872]of the peroxisome proliferator-activated receptor gamma
(PPAR-gamma)

[151873]in human atherosclerosis and regulation in macrophages
by colony stimulating

[151874]factors and oxidized low density lipoprotein. Proc. Nat.
Acad. Sci. 95:

[151875]7614-7619, 1998.

[151876]

[151877]

[151878]14556.

[151879]Savage, D. B.; Agostini, M.; Barroso, I.; Gurnell, M.; Luan, J.;

[151880]Meirhaeghe, A.; Harding, A.-H.; Ihrke, G.; Rajanayagam, O.; Soos,

[151881]M. A.; George, S.; Berger, D.; and 9 others: Digenic inheritance

[151882]of severe insulin resistance in a human pedigree. *Nature Genet.* 31:

[151883]379–384, 2002. Note: Erratum: *Nature Genet.* 32: 211 only, 2002.

[151884]

[151885]

[151886]14557.

[151887]Smith, W. M.; Zhou, X.-P.; Kurose, K.; Gao, X.; Latif, F.; Kroll,

[151888]T.; Sugano, K.; Cannistra, S. A.; Clinton, S. K.; Maher, E. R.; Prior,

[151889]T. W.; Eng, C.: Opposite association of two PPARG variants with cancer:

[151890]overrepresentation of H449H in endometrial carcinoma cases and underrepresentation

[151891]of P12A in renal cell carcinoma cases. Hum. Genet. 109:
146–151,
[151892]2001.

[151893]

[151894]

[151895]14558.

[151896]Tarrade, A.; Schoonjans, K.; Pavan, L.; Auwerx, J.; Ro-
chette-Egly,

[151897]C.; Evain-Brion, D.; Fournier, T.: PPAR-gamma/RXR-alpha
heterodimers

[151898]control human trophoblast invasion. J. Clin. Endocr.
Metab. 86:

[151899]5017–5024, 2001.

[151900]

[151901]

[151902]14559.

[151903]Tontonoz, P.; Hu, E.; Devine, J.; Beale, E. G.; Spiegelman,
B.

[151904]M.: PPAR gamma 2 regulates adipose expression of the
phosphoenolpyruvate

[151905]carboxykinase gene. Molec. Cell. Biol. 15: 351–357, 1995.

[151906]

[151907]

[151908]14560.

[151909]Tontonoz, P.; Hu, E.; Graves, R. A.; Budavari, A. I.; Spiegelman,

[151910]B. M.: mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. Genes

[151911]Dev. 8: 1224–1234, 1994.

[151912]

[151913]

[151914]14561.

[151915]Tontonoz, P.; Nagy, L.; Alvarez, J. G. A.; Thomazy, V. A.; Evans,

[151916]R. M.: PPAR-gamma promotes monocyte/macrophage differentiation and

[151917]uptake of oxidized LDL. Cell 93: 241–252, 1998.

[151918]

[151919]

[151920]14562.

[151921]Wang, M.: Isolation and characterization of a human gene encoding

[151922]a single-strand-specific endoribonuclease. Ph.D. Thesis: Stanford

[151923]Univ. , 1995.

[151924]

[151925]

[151926]14563.

[151927]Wang, M.; Cohen, S. N.: ard-1: a human gene that reverses the

[151928]effects of temperature-sensitive and deletion mutations in the Escherichia

[151929]coli rne gene and encodes an activity producing RNase E-like cleavages. Proc.

[151930]Nat. Acad. Sci. 91: 10591-10595, 1994.

[151931]

[151932]

[151933]14564.

[151934]Beck, S.; Hanson, I.; Kelly, A.; Pappin, D. J.; Trowsdale, J.:

[151935]A homologue of the Drosophila female sterile homeotic
(fsh) gene in

[151936]the class II region of the human MHC. DNA Sequence 2:
203–210, 1992.

[151937]

[151938]

[151939]14565.

[151940]Denis, G. V.; Green, M. R.: A novel, mitogen-activated nu-
clear

[151941]kinase is related to a Drosophila developmental regulator.
Genes

[151942]Dev. 10: 261–271, 1996.

[151943]

[151944]

[151945]14566.

[151946]Okamoto, N.; Ando, A.; Kawai, J.; Yoshiwara, T.; Tsuji, K.;
Inoko,

[151947]H.: Orientation of HLA–DNA gene and identification of a
CpG island–associated

[151948]gene adjacent to DNA in human major histocompatibility
complex class

[151949]II region. Hum. Immun. 32: 221–228, 1991.

[151950]

[151951]

[151952]14567.

[151953]Thorpe, K. L.; Abdulla, S.; Kaufman, J.; Trowsdale, J.; Beck, S.

[151954]: Phylogeny and structure of the RING3 gene. Immunogenetics 44:

[151955]391–396, 1996.

[151956]

[151957]

[151958]14568.

[151959]Cushman, L. J.; Camper, S. A.: Molecular basis of pituitary dysfunction

[151960]in mouse and human. Mammalian Genome 12: 485–494, 2001.

[151961]

[151962]

[151963]14569.

[151964]Kozak, M.: Interpreting cDNA sequences: some insights
from studies

[151965]on translation. Mammalian Genome 7: 563–574, 1996.

[151966]

[151967]

[151968]14570.

[151969]Pellegrini–Bouiller, I.; Manrique, C.; Gunz, G.; Grino, M.;
Zamora,

[151970]A. J.; Figarella–Branger, D.; Grisoli, F.; Jaquet, P.; Enjalbert,

[151971]A.: Expression of the members of the Ptx family of tran-
scription

[151972]factors in human pituitary adenomas. J. Clin. Endocr.
Metab. 84:

[151973]2212–2220, 1999.

[151974]

[151975]

[151976]14571.

[151977]Brinke, A.; Green, P. M.; Giannelli, F.: Characterization of
the

[151978]gene (VBP1) and transcript for the von Hippel–Lindau

binding protein

[151979]and isolation of the highly conserved murine homologue.

Genomics 45:

[151980]105–112, 1997.

[151981]

[151982]

[151983]14572.

[151984]Brinke, A.; Tagliavacca, L.; Naylor, J.; Green, P.; Gian-
grande,

[151985]P.; Giannelli, F.: Two chimaeric transcription units result
from

[151986]an inversion breaking intron 1 of the factor VIII gene and a
region

[151987]reportedly affected by reciprocal translocations in T-cell
leukaemia. Hum.

[151988]Molec. Genet. 5: 1945–1951, 1996.

[151989]

[151990]

[151991]14573.

[151992]Clifford, S. C.; Walsh, S.; Hewson, K.; Green, E. K.; Brinke,

A.;

[151993]Green, P. M.; Gianelli, F.; Eng, C.; Maher, E. R.: Genomic organization

[151994]and chromosomal localization of the human CUL2 gene and the role of

[151995]von Hippel–Lindau tumor suppressor–binding protein (CUL2 and VBP1)

[151996]mutation and loss in renal–cell carcinoma development. Genes Chromosomes

[151997]Cancer 26: 20–28, 1999.

[151998]

[151999]

[152000]14574.

[152001]Hemberger, M.; Himmelbauer, H.; Neumann, H. P. H.; Plate, K. H.;

[152002]Schwarzkopf, G.; Fundele, R.: Expression of the von Hippel–Lindau–binding

[152003]protein–1 (Vbp1) in fetal and adult mouse tissues. Hum. Molec. Genet. 8:

[152004]229–236, 1999.

[152005]

[152006]

[152007]14575.

[152008]Tsuchiya, H.; Iseda, T.; Hino, O.: Identification of a novel
protein

[152009](VBP-1) binding to the von Hippel-Lindau (VHL) tumor
suppressor gene

[152010]product. Cancer Res. 56: 2881-2885, 1996.

[152011]

[152012]

[152013]14576.

[152014]Corbaz, A.; ten Hove, T.; Herren, S.; Graber, P.; Schwartz-
burd,

[152015]B.; Belzer, I.; Harrison, J.; Plitz, T.; Kosco-Vilbois, M. H.;
Kim,

[152016]S.-H.; Dinarello, C. A.; Novick, D.; van Deventer, S.;
Chvatchko,

[152017]Y.: IL-18-binding protein expression by endothelial cells
and macrophages

[152018]is up-regulated during active Crohn's disease. J. Immun.
168: 3608-3616,

[152019]2002.

[152020]

[152021]

[152022]14577.

[152023]Turner, N.; Mason, P. J.; Brown, R.; Fox, M.; Povey, S.;
Rees,

[152024]A.; Pusey, C. D.: Molecular cloning of the human Good-
pasture antigen

[152025]demonstrates it to be the alpha-3 chain of type IV colla-
gen. J. Clin.

[152026]Invest. 89: 592-601, 1992.

[152027]

[152028]

[152029]14578.

[152030]Hoffmann, M. M.; Jacob, S.; Luft, D.; Schmulling, R.-M.;
Rett,

[152031]K.; Marz, W.; Haring, H.-U.; Matthaei, S.: Type I hyper-
lipoproteinemia

[152032]due to a novel loss of function mutation of lipoprotein li-
pase, cys239-trp,

[152033]associated with recurrent severe pancreatitis. J. Clin. En-
docr. Metab. 85:

[152034]4795–4798, 2000.

[152035]

[152036]

[152037]14579.

[152038]Holt, L. E., Jr.; Aylward, F. X.; Timbers, H. G.: Idiopathic

[152039]familial lipemia. Bull. Johns Hopkins Hosp. 64: 279–314,
1939.

[152040]

[152041]

[152042]14580.

[152043]Ishimura–Oka, K.; Faustinella, F.; Kihara, S.; Smith, L. C.;
Oka,

[152044]K.; Chan, L.: A missense mutation (trp86–to–arg) in exon
3 of the

[152045]lipoprotein lipase gene: a cause of familial chylomicrone–
mia. Am.

[152046]J. Hum. Genet. 50: 1275–1280, 1992.

[152047]

[152048]

[152049]14581.

[152050]Kastelein, J. J. P.; Groenemeyer, B. E.; Hallman, D. M.; Henderson,

[152051]H.; Reymer, P. W. A.; Gagne, S. E.; Jansen, H.; Seidell, J. C.; Kromhout,

[152052]D.; Jukema, J. W.; Bruschke, A. V. G.; Boerwinkle, E.; Hayden, M.

[152053]R.; The Regress Study Group: The asn9 variant of lipoprotein lipase

[152054]is associated with the -93G promoter mutation and an increased risk

[152055]of coronary artery disease. Clin. Genet. 53: 27-33, 1998.

[152056]

[152057]

[152058]14582.

[152059]Kirchgessner, T. G.; Chuat, J.-C.; Heinzmann, C.; Etienne, J.;

[152060]Guilhot, S.; Svenson, K.; Ameis, D.; Pilon, C.; d'Auriol, L.; Andalibi,

[152061]A.; Schotz, M. C.; Galibert, F.; Lusi, A. J.: Organization of the

[152062]human lipoprotein lipase gene and evolution of the lipase
gene family. Proc.

[152063]Nat. Acad. Sci. 86: 9647–9651, 1989.

[152064]

[152065]

[152066]14583.

[152067]Kirchgessner, T. G.; Svenson, K. L.; Lusi, A. J.; Schotz, M.

[152068]C.: The sequence of cDNA encoding lipoprotein lipase: a
member of

[152069]a lipase gene family. J. Biol. Chem. 262: 8463–8466,
1987.

[152070]

[152071]

[152072]14584.

[152073]Kobayashi, J.; Nishida, T.; Ameis, D.; Stahnke, G.; Schotz,
M.

[152074]C.; Hashimoto, H.; Fukamachi, I.; Shirai, K.; Saito, Y.;
Yoshida,

[152075]S.: A heterozygous mutation (the codon for ser477 to a
stop codon)

[152076]in lipoprotein lipase contributes to a defect in lipid inter-
face recognition

[152077]in a case with type I hyperlipidemia. Biochem. Biophys.
Res. Commun. 182:

[152078]70–77, 1992.

[152079]

[152080]

[152081]14585.

[152082]Kobayashi, J.; Sasaki, N.; Tashiro, J.; Inadera, H.; Saito, Y.;

[152083]Yoshida, S.: A missense mutation (ala334–to–thr) in exon
7 of the

[152084]lipoprotein lipase gene in a case with type I hyperlipi-
demia. Biochem.

[152085]Biophys. Res. Commun. 191: 1046–1054, 1993.

[152086]

[152087]

[152088]14586.

[152089]Langlois, S.; Deeb, S.; Brunzell, J.; Kastelein, J. J.; Hayden,

[152090]M. R.: A unique insertion accounts for a significant pro–

portion of

[152091]the mutations in the lipoprotein lipase (LPL)
gene.(Abstract) Am.

[152092]J. Hum. Genet. 43: A191, 1988.

[152093]

[152094]

[152095]14587.

[152096]Langlois, S.; Deeb, S.; Brunzell, J. D.; Kastelein, J. J.; Hay-
den,

[152097]M. R.: A major insertion accounts for a significant propor-
tion of

[152098]mutations underlying human lipoprotein lipase deficiency.
Proc. Nat.

[152099]Acad. Sci. 86: 948–952, 1989.

[152100]

[152101]

[152102]14588.

[152103]Levak–Frank, S.; Radner, H.; Walsh, A.; Stollberger, R.;
Knipping,

[152104]G.; Hoefler, G.; Sattler, W.; Weinstock, P. H.; Breslow, J. L.;

Zechner,

[152105]R.: Muscle-specific overexpression of lipoprotein lipase causes a

[152106]severe myopathy characterized by proliferation of mitochondria and

[152107]peroxisomes in transgenic mice. J. Clin. Invest. 96: 976–986, 1995.

[152108]

[152109]

[152110]14589.

[152111]Li, S.; Oka, K.; Galton, D.; Stocks, J.: Pvu-II RFLP at the human

[152112]lipoprotein lipase (LPL) gene locus. Nucleic Acids Res. 16: 2358,

[152113]1988.

[152114]

[152115]

[152116]14590.

[152117]Lo, J. Y.; Smith, L. C.; Chan, L.: Lipoprotein lipase: role of

[152118]intramolecular disulfide bonds in enzyme catalysis.

Biochem. Biophys.

[152119]Res. Commun. 206: 266–271, 1995.

[152120]

[152121]

[152122]14591.

[152123]Ma, Y.; Henderson, H. E.; Julien, P.; Roederer, G.; Brunzell,

[152124]J.; Hayden, M. R.: A missense mutation (pro-to-leu207) in

[152125]the human lipoprotein lipase gene is the major cause of
type I hyperlipoproteinemia

[152126]in French Canadians. (Series) Miami Short Reports. Ad-
vances

[152127]in Gene Technology: The Molecular Biology of Human Ge-
netic Disease.

[152128]New York: IRL Press (pub.) 1: 1991. Pp. 34 only.

[152129]

[152130]

[152131]14592.

[152132]Ma, Y.; Henderson, H. E.; Ven Murthy, M. R.; Roederer, G.;
Monsalve,

[152133]M. V.; Clarke, L. A.; Normand, T.; Julien, P.; Gagne, C.;
Lambert,

[152134]M.; Davignon, J.; Lupien, P. J.; Brunzell, J.; Hayden, M. R.: A
mutation

[152135]in the human lipoprotein lipase gene as the most common
cause of familial

[152136]chylomicronemia in French Canadians. New Eng. J. Med.
324: 1761–1766,

[152137]1991.

[152138]

[152139]

[152140]14593.

[152141]Ma, Y.; Liu, M.–S.; Chitayat, D.; Bruin, T.; Beisiegel, U.;
Benlian,

[152142]P.; Foubert, L.; De Gennes, J. L.; Funke, H.; Forsythe, I.;
Blaichman,

[152143]S.; Papanikolaou, M.; Erkelens, D. W.; Kastelein, J.; Brun–
zell, J.

[152144]D.; Hayden, M. R.: Recurrent missense mutations at the
first and

[152145]second base of codon arg243 in human lipoprotein lipase
in patients

[152146]of different ancestries. Hum. Mutat. 3: 52–58, 1994.

[152147]

[152148]

[152149]14594.

[152150]Ma, Y.; Liu, M.–S.; Ginzinger, D.; Frohlich, J.; Brunzell, J.

[152151]D.; Hayden, M. R.: Gene–environment interaction in the
conversion

[152152]of a mild–to–severe phenotype in a patient homozygous
for a ser172–to–cys

[152153]mutation in the lipoprotein lipase gene. J. Clin. Invest. 91:
1953–1958,

[152154]1993.

[152155]

[152156]

[152157]14595.

[152158]Ma, Y.; Wilson, B. I.; Bijvoet, S.; Henderson, H. E.; Cramb,
E.;

[152159]Roederer, G.; Ven Murthy, M. R.; Julien, P.; Bakker, H. D.;
Kastelein,

[152160]J. J. P.; Brunzell, J. D.; Hayden, M. R.: A missense mutation

(asp250-to-asn)

[152161]in exon 6 of the human lipoprotein lipase gene causes
chylomicronemia

[152162]in patients of different ancestries. Genomics 13: 649–653,
1992.

[152163]

[152164]

[152165]14596.

[152166]Mattei, M. G.; Etienne, J.; Chuat, J. C.; Nguyen, V. C.;
Brault,

[152167]D.; Bernheim, A.; Galibert, F.: Assignment of the human
lipoprotein

[152168]lipase (LPL) gene to chromosome band 8p22. Cytogenet.
Cell Genet. 63:

[152169]45–46, 1993.

[152170]

[152171]

[152172]14597.

[152173]Miesenbock, G.; Holzl, B.; Foger, B.; Brandstatter, E.;
Paulweber,

[152174]B.; Sandhofer, F.; Patsch, J. R.: Heterozygous lipoprotein

lipase

[152175]deficiency due to a missense mutation as the cause of impaired triglyceride

[152176]tolerance with multiple lipoprotein abnormalities. J. Clin. Invest. 91:

[152177]448–455, 1993.

[152178]

[152179]

[152180]14598.

[152181]Nevin, D. N.; Brunzell, J. D.; Deeb, S. S.: The LPL gene in individuals

[152182]with familial combined hyperlipidemia and decreased LPL activity. Arteriosclerosis

[152183]Thromb. 14: 869–873, 1994.

[152184]

[152185]

[152186]14599.

[152187]Nevin, N. C.; Slack, J.: Hyperlipidaemic xanthomatosis II: mode

[152188]of inheritance in 55 families with essential hyperlipi-

daemia and xanthomatosis. J.

[152189]Med. Genet. 5: 9–28, 1968.

[152190]

[152191]

[152192]14600.

[152193]Nickerson, D. A.; Taylor, S. L.; Weiss, K. M.; Clark, A. G.;
Hutchinson,

[152194]R. G.; Stengard, J.; Salomaa, V.; Vartiainen, E.; Boerwinkle,
E.;

[152195]Sing, C. F.: DNA sequence diversity in a 9.7–kb region of
the human

[152196]lipoprotein lipase gene. Nature Genet. 19: 233–240,
1998.

[152197]

[152198]

[152199]14601.

[152200]Lahn, B. T.; Page, D. C.: Retroposition of autosomal mRNA
yielded

[152201]testis–specific gene family on human Y chromosome. Na-
ture Genet. 21:

[152202]429–433, 1999.

[152203]

[152204]

[152205]14602.

[152206]Yen, P. H.: A long-range restriction map of deletion interval

[152207]6 of the human Y chromosome: a region frequently deleted in azoospermic

[152208]males. Genomics 54: 5-12, 1998.

[152209]

[152210]

[152211]14603.

[152212]MacLean, H. E.; Chu, S.; Warne, G. L.; Zajac, J. D.: Related

[152213]individuals with different androgen receptor gene deletions. J. Clin.

[152214]Invest. 91: 1123-1128, 1993.

[152215]

[152216]

[152217]14604.

[152218]Madgar, I.; Green, L.; Kent-First, M.; Weissenberg, R.; Ger-

shoni-Baruch,

[152219]R.; Goldman, B.; Friedman, E.: Genotyping of Israeli infertile men

[152220]with idiopathic oligozoospermia. Clin. Genet. 62: 203-207, 2002.

[152221]

[152222]

[152223]14605.

[152224]Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Griffin, J. E.; Wilson,

[152225]J. D.; McPhaul, M. J.: Definition of the human androgen receptor

[152226]gene structure permits the identification of mutations that cause

[152227]androgen resistance: premature termination of the receptor protein

[152228]at amino acid residue 588 causes complete androgen resistance. Molec.

[152229]Endocr. 4: 1105-1116, 1990.

[152230]

[152231]

[152232]14606.

[152233]Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Wilson, J. D.; Griffin,

[152234]J. E.; McPhaul, M. J.: A single nucleotide substitution introduces

[152235]a premature termination codon into the androgen receptor gene of a

[152236]patient with receptor-negative androgen resistance. J. Clin. Invest. 85:

[152237]1522-1528, 1990.

[152238]

[152239]

[152240]14607.

[152241]Marcelli, M.; Tilley, W. D.; Zoppi, S.; Griffin, J. E.; Wilson,

[152242]J. D.; McPhaul, M. J.: Androgen resistance associated with a mutation

[152243]of the androgen receptor at amino acid 772 (arg-to-cys) results from

[152244]a combination of decreased messenger ribonucleic acid levels and impairment

[152245]of receptor function. J. Clin. Endocr. 73: 318-325, 1991.

[152246]

[152247]

[152248]14608.

[152249]McCampbell, A.; Taylor, J. P.; Taye, A. A.; Robitschek, J.; Li,

[152250]M.; Walcott, J.; Merry, D.; Chai, Y.; Paulson, H.; Sobue, G.;

Fischbeck,

[152251]K. H.: CREB-binding protein sequestration by expanded
polyglutamine. Hum.

[152252]Molec. Genet. 9: 2197-2202, 2000.

[152253]

[152254]

[152255]14609.

[152256]McPhaul, M. J.; Griffin, J. E.: Male pseudohermaphroditism
caused

[152257]by mutations of the human androgen receptor. J. Clin. En-
docr. Metab. 84:

[152258]3435-3441, 1999.

[152259]

[152260]

[152261]14610.

[152262]McPhaul, M. J.; Marcelli, M.; Tilley, W. D.; Griffin, J. E.;

Isidro-Gutierrez,

[152263]R. F.; Wilson, J. D.: Molecular basis of androgen resistance
in a

[152264]family with a qualitative abnormality of the androgen re-
ceptor and

[152265]responsive to high-dose androgen therapy. J. Clin. Invest.
87: 1413-1421,

[152266]1991.

[152267]

[152268]

[152269]14611.

[152270]McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Griffin, J. E.; Wilson,

[152271]J. D.: Genetic basis of endocrine disease 4: the spectrum
of mutations

[152272]in the androgen receptor gene that causes androgen re-
sistance. J.

[152273]Clin. Endocr. Metab. 76: 17-23, 1993.

[152274]

[152275]

[152276]14612.

[152277]McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Wilson, C. M.; Griffin,

[152278]J. E.; Wilson, J. D.: Mutations in the ligand-binding domain of the

[152279]androgen receptor gene cluster in two regions of the gene. J. Clin.

[152280]Invest. 90: 2097-2101, 1992.

[152281]

[152282]

[152283]14613.

[152284]McPhaul, M. J.; Schweikert, H.-U.; Allman, D. R.: Assessment

[152285]of androgen receptor function in genital skin fibroblasts using a

[152286]recombinant adenovirus to deliver an androgen-responsive reporter

[152287]gene. J. Clin. Endocr. Metab. 82: 1944-1948, 1997.

[152288]

[152289]

[152290]14614.

[152291]Mifsud, A.; Ramirez, S.; Yong, E. L.: Androgen receptor
gene

[152292]CAG trinucleotide repeats in anovulatory infertility and
polycystic

[152293]ovaries. J. Clin. Endocr. Metab. 85: 3484–3488, 2000.

[152294]

[152295]

[152296]14615.

[152297]Migeon, B. R.; Brown, T. R.; Axelman, J.; Migeon, C. J.:
Studies

[152298]of the locus for androgen receptor: localization on the hu-
man X and

[152299]evidence for homology with the Tfm locus in the mouse.
Proc. Nat.

[152300]Acad. Sci. 78: 6339–6343, 1981.

[152301]

[152302]

[152303]14616.

[152304]Mongan, N. P.; Jaaskelainen, J.; Green, K.; Schwabe, J. W.; Shimura,

[152305]N.; Dattani, M.; Hughes, I. A.: Two de novo mutations in the AR gene

[152306]cause the complete androgen insensitivity syndrome in a pair of monozygotic

[152307]twins. J. Clin. Endocr. Metab. 87: 1057–1061, 2002.

[152308]

[152309]

[152310]14617.

[152311]Mononen, N.; Syrjakoski, K.; Matikainen, M.; Tammela, T. L. J.;

[152312]Schleutker, J.; Kallioniemi, O.–P.; Trapman, J.; Koivisto, P. A.:

[152313]Two percent of Finnish prostate cancer patients have a germ–line mutation

[152314]in the hormone–binding domain of the androgen receptor gene. Cancer

[152315]Res. 60: 6479–6481, 2000.

[152316]

[152317]

[152318]14618.

[152319]Muro, K.; Mendonca, B. B.; Arnhold, I. J. P.; Rigon, A. C. M.

[152320]M.; Migeon, C. J.; Brown, T. R.: Human androgen insensitivity due

[152321]to point mutations encoding amino acid substitutions in the androgen

[152322]receptor steroid-binding domain. Hum. Mutat. 6: 152-162, 1995.

[152323]

[152324]

[152325]14619.

[152326]Nakao, R.; Haji, M.; Yanase, T.; Ogo, A.; Takayanagi, R.; Katsube,

[152327]T.; Fukumaki, Y.; Nawata, H.: A single amino acid substitution (met786-to-val)

[152328]in the steroid-binding domain of human androgen receptor leads to

[152329]complete androgen insensitivity syndrome. J. Clin. Endocr. Metab. 74:

[152330]1152-1157, 1992.

[152331]

[152332]

[152333]14620.

[152334]Newmark, J. R.; Hardy, D. O.; Tonb, D. C.; Carter, B. S.; Epstein,
stein,

[152335]J. I.; Isaacs, W. B.; Brown, T. R.; Barrack, E. R.: Androgen
receptor

[152336]gene mutations in human prostate cancer. Proc. Nat.
Acad. Sci. 89:

[152337]6319–6323, 1992.

[152338]

[152339]

[152340]14621.

[152341]Nguyen, D.; Steinberg, S. V.; Rouault, E.; Chagnon, S.;
Gottlieb,

[152342]B.; Pinsky, L.; Trifiro, M.; Mader, S.: A G577R mutation in
the human

[152343]AR P box results in selective decreases in DNA binding
and in partial

[152344]androgen insensitivity syndrome. Molec. Endocr. 15:
1790–1802, 2001.

[152345]

[152346]

[152347]14622.

[152348]Ohno, S.: The Y-linked antigen locus and the X-linked
Tfm locus

[152349]as major regulatory genes of the mammalian sex determining mechanism. J.

[152350]Steroid Biochem. 8: 585–592, 1977.

[152351]

[152352]100. Ohno, S.: Simplicity of mammalian regulatory systems inferred

[152353]by single gene determination of sex phenotypes. Nature
234: 134–137,

[152354]1971.

[152355]

[152356]101. Ong, Y. C.; Wong, H. B.; Adaikan, G.; Yong, E. L.: Directed

[152357]pharmacological therapy of ambiguous genitalia due to an
androgen

[152358]receptor gene mutation. (Letter) Lancet 354: 1444–1445,
1999.

[152359]

[152360]102. Patterson, M. N.; Hughes, I. A.; Gottlieb, B.; Pinsky, L.: The

[152361]androgen receptor gene mutations database. Nucleic Acids Res. 22:

[152362]3560–3562, 1994.

[152363]

[152364]103. Pinsky, L.; Kaufman, M.; Killinger, D. W.; Burko, B.; Shatz,

[152365]D.; Volpe, R.: Human minimal androgen insensitivity with normal dihydrotestosterone-binding

[152366]capacity in cultured genital skin fibroblasts: evidence for an androgen-selective

[152367]qualitative abnormality of the receptor. Am. J. Hum. Genet. 36:

[152368]965–978, 1984.

[152369]

[152370]104. Pinsky, L.; Kaufman, M.; Levitsky, L. L.: Partial androgen resistance

[152371]due to a distinctive qualitative defect of the androgen receptor. Am.

[152372]J. Med. Genet. 27: 459–466, 1987.

[152373]

[152374]105. Pinsky, L.; Kaufman, M.; Summitt, R. L.: Congenital androgen

[152375]insensitivity due to a qualitatively abnormal androgen receptor. Am.

[152376]J. Med. Genet. 10: 91–99, 1981.

[152377]

[152378]106. Pinsky, L.; Trifiro, M.; Sebbaghian, N.; Kaufman, M.; Chang,

[152379]C.; Trapman, J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris, C. J.;

[152380]Brown, C. J.; Willard, H. F.; Sergovich, F.: A deletional alteration

[152381]of the androgen receptor (AR) gene in a sporadic patient with complete

[152382]androgen insensitivity (CAI) who is mentally retarded. (Abstract) Am.

[152383]J. Hum. Genet. 45 (suppl.): A212, 1989.

[152384]

[152385]107. Prior, L.; Bordet, S.; Trifiro, M. A.; Mhatre, A.; Kauf-

man, M.;

[152386]Pinsky, L.; Wrogeman, K.; Belsham, D. D.; Pereira, F.;

Greenberg,

[152387]C.; Trapman, J.; Brinkman, A. O.; Chang, C.; Liao, S.: Re-

placement

[152388]of arginine 773 by cysteine or histidine in the human an-

drogen receptor

[152389]causes complete androgen insensitivity with different re-

ceptor phenotypes. Am.

[152390]J. Hum. Genet. 51: 143-155, 1992.

[152391]

[152392]108. Quigley, C. A.; Friedman, K. J.; Johnson, A.; Lafre-

niere, R.

[152393]G.; Silverman, L. M.; Lubahn, D. B.; Brown, T. R.; Wilson, E.

M.;

[152394]Willard, H. F.; French, F. S.: Complete deletion of the an-

drogen

[152395]receptor gene: definition of the null phenotype of the an-

drogen insensitivity

[152396]syndrome and determination of carrier status. J. Clin. En-

docr. Metab. 74:

[152397]927-933, 1992.

[152398]

[152399]109. Ris-Stalpers, C.; Kuiper, G. G. J. M.; Faber, P. W.;
Schweikert,

[152400]H. U.; van Rooij, H. C. J.; Zegers, N. D.; Hodgins, M. B.;
Degenhart,

[152401]H. J.; Trapman, J.; Brinkmann, A. O.: Aberrant splicing of
androgen

[152402]receptor mRNA results in synthesis of a nonfunctional re-
ceptor protein

[152403]in a patient with androgen insensitivity. Proc. Nat. Acad.
Sci. 87:

[152404]7866–7870, 1990.

[152405]

[152406]110. Rodien, P.; Mebarki, F.; Mowszowicz, I.; Chaussain,
J.-L.; Young,

[152407]J.; Morel, Y.; Schaison, G.: Different phenotypes in a family
with

[152408]androgen insensitivity caused by the same M780I point
mutation in

[152409]the androgen receptor gene. J. Clin. Endocr. Metab. 81:
2994–2998,

[152410]1996.

[152411]

[152412]111. Sai, T.; Seino, S.; Chang, C.; Trifiro, M.; Pinsky, L.;
Mhatre,

[152413]A.; Kaufman, M.; Lambert, B.; Trapman, J.; Brinkmann, A.
O.; Rosenfield,

[152414]R. L.; Liao, S.: An exonic point mutation of the androgen
receptor

[152415]gene in a family with complete androgen insensitivity. Am.
J. Hum.

[152416]Genet. 46: 1095–1100, 1990.

[152417]

[152418]112. Sammarco, I.; Grimaldi, P.; Rossi, P.; Cappa, M.;
Moretti, C.;

[152419]Frajese, G.; Geremia, R.: Novel point mutation in the splice
donor

[152420]site of exon–intron junction 6 of the androgen receptor
gene in a

[152421]patient with partial androgen insensitivity syndrome. J.
Clin. Endocr.

[152422]Metab. 85: 3256–3261, 2000.

[152423]

[152424]113. Schoenberg, M. P.; Hakimi, J. M.; Wang, S.; Bova, G.

S.; Epstein,

[152425]J. I.; Fischbeck, K. H.; Isaacs, W. B.; Walsh, P. C.; Barrack, E.

[152426]R.: Microsatellite mutation (CAG (24-to-18)) in the androgen receptor

[152427]gene in human prostate cancer Biochem. Biophys. Res. Commun. 198:

[152428]74-80, 1994.

[152429]

[152430]114. Shang, Y.; Myers, M.; Brown, M.: Formation of the androgen receptor

[152431]transcription complex. Molec. Cell 9: 601-610, 2002.

[152432]

[152433]115. Simeoni, S.; Mancini, M. A.; Stenoien, D. L.; Marcelli, M.; Weigel,

[152434]N. L.; Zanisi, M.; Martini, L.; Poletti, A.: Motoneuronal cell death

[152435]is not correlated with aggregate formation of androgen receptors containing

[152436]an elongated polyglutamine tract. Hum. Molec. Genet. 9: 133-144,

[152437]2000.

[152438]

[152439]116. Sullivan, D. A.; Sullivan, B. D.; Ullman, M. D.; Rocha,
E. M.;

[152440]Krenzer, K. L.; Cermak, J. M.; Toda, I.; Doane, M. G.; Evans,
J. E.;

[152441]Wickham, L. A.: Androgen influence on the meibomian
gland. Invest.

[152442]Ophthal. Vis. Sci. 41: 3732–3742, 2000.

[152443]

[152444]117. Sultan, C.; Lumbroso, S.; Poujol, N.; Belon, C.;
Boudon, C.;

[152445]Lobaccaro, J.–M.: Mutations of androgen receptor gene in
androgen

[152446]insensitivity syndromes. J. Steroid Biochem. Molec. Biol.
46: 519–530,

[152447]1993.

[152448]

[152449]118. Sutherland, R. W.; Wiener, J. S.; Hicks, J. P.; Marcelli,
M.;

[152450]Gonzales, E. T.; Roth, D. R.; Lamb, D. J.: Androgen recep–

tor gene

[152451]mutations are rarely associated with isolated penile hypospadias. J.

[152452]Urol. 156: 828–831, 1996.

[152453]

[152454]119. Taplin, M.–E.; Bubley, G. J.; Shuster, T. D.; Frantz, M. E.;

[152455]Spooner, A. E.; Ogata, G. K.; Keer, H. N.; Balk, S. P.: Mutation

[152456]of the androgen–receptor gene in metastatic androgen–independent prostate

[152457]cancer. New Eng. J. Med. 332: 1393–1398, 1995.

[152458]

[152459]120. Tilley, W. D.; Marcelli, M.; Wilson, J. D.; McPhaul, M. J.:

[152460]Characterization and expression of a cDNA encoding the human androgen

[152461]receptor. Proc. Nat. Acad. Sci. 86: 327–331, 1989.

[152462]

[152463]121. Trifiro, M.; Prior, L.; Pinsky, L.; Kaufman, M.; Chang, C.; Trapman,

[152464]J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris, C.: A single transition

[152465]at an exonic CpG site apparently abolishes androgen receptor (AR)–binding

[152466]activity in a family with complete androgen insensitivity (CAI). (Abstract) Am.

[152467]J. Hum. Genet. 45 (suppl.): A225, 1989.

[152468]

[152469]122. Trifiro, M.; Prior, R. L.; Sabbaghian, N.; Pinsky, L.; Kaufman,

[152470]M.; Nylén, E. G.; Belsham, D. D.; Greenberg, C. R.; Wrogemann, K.

[152471]: Amber mutation creates a diagnostic Mael site in the androgen receptor

[152472]gene of a family with complete androgen insensitivity. Am. J. Med.

[152473]Genet. 40: 493–499, 1991.

[152474]

[152475]123. Tut, T. G.; Ghadessy, F. J.; Trifiro, M. A.; Pinsky, L.; Yong,

[152476]E. L.: Long polyglutamine tracts in the androgen receptor are associated

[152477]with reduced trans-activation, impaired sperm production, and male

[152478]infertility. J. Clin. Endocr. Metab. 82: 3777–3782, 1997.

[152479]

[152480]124. Visakorpi, T.; Hyytinen, E.; Koivisto, P.; Tanner, M.; Keinanen,

[152481]R.; Palmberg, C.; Palotie, A.; Tammela, T.; Isola, J.; Kallioniemi,

[152482]O.–P.: In vivo amplification of the androgen receptor gene and progression

[152483]of human prostate cancer. Nature Genet. 9: 401–406, 1995.

[152484]

[152485]125. Von Eckardstein, S.; Syska, A.; Gromoll, J.; Kamischke, A.; Simoni,

[152486]M.; Nieschlag, E.: Inverse correlation between sperm concentration

[152487]and number of androgen receptor CAG repeats in normal men. J. Clin.

[152488]Endocr. Metab. 86: 2585–2590, 2001.

[152489]

[152490]126. Wang, Q.; Ghadessy, F. J.; Trounson, A.; de Kretser, D.; McLachlan,

[152491]R.; Ng, S. C.; Yong, E. L.: Azoospermia associated with a mutation

[152492]in the ligand-binding domain of an androgen receptor displaying normal

[152493]ligand binding, but defective trans-activation. J. Clin. Endocr.

[152494]Metab. 83: 4303-4309, 1998.

[152495]

[152496]127. Weidemann, W.; Peters, B.; Romalo, G.; Spindler, K.-D.; Schweikert,

[152497]H.-U.: Response to androgen treatment in a patient with partial androgen

[152498]insensitivity and a mutation in the deoxyribonucleic acid-binding

[152499]domain of the androgen receptor. J. Clin. Endocr. Metab. 83: 1173-1176,

[152500]1998.

[152501]

[152502]128. Welch, W. J.; Diamond, M. I.: Glucocorticoid modulation of androgen

[152503]receptor nuclear aggregation and cellular toxicity is associated with

[152504]distinct forms of soluble expanded polyglutamine protein. Hum. Molec.

[152505]Genet. 10: 3063–3074, 2001.

[152506]

[152507]129. Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.; Landen,

[152508]M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Polymorphisms

[152509]of the androgen receptor gene and the estrogen receptor beta gene

[152510]are associated with androgen levels in women. J. Clin. Endocr. Metab. 86:

[152511]2562–2568, 2001.

[152512]

[152513]130. Wieacker, P.; Breckwoldt, M.; Gal, A.: Testicular feminization:

[152514]diagnosis and search for closely linked restriction fragment length

[152515]polymorphism. Dis. Markers 3: 213–218, 1985.

[152516]

[152517]131. Wieacker, P.; Griffin, J. E.; Wienker, T.; Lopez, J. M.;
Wilson,

[152518]J. D.; Breckwoldt, M.: Linkage analysis with RFLPs in fami-
lies with

[152519]androgen resistance syndromes: evidence for close link-
age between

[152520]the androgen receptor locus and the DXS1 segment. Hum.
Genet. 76:

[152521]248–252, 1987.

[152522]

[152523]132. Wilson, C. M.; McPhaul, M. J.: A and B forms of the
androgen

[152524]receptor are present in human genital skin fibroblasts.
Proc. Nat.

[152525]Acad. Sci. 91: 1234–1238, 1994.

[152526]

[152527]133. Wilson, J. D.: The promiscuous receptor: prostate
cancer comes

[152528]of age. (Editorial) New Eng. J. Med. 332: 1440–1441,
1995.

[152529]

[152530]134. Wilson, J. D.; Carlson, B. R.; Weaver, D. D.; Kovacs, W. J.;

[152531]Griffin, J. E.: Endocrine and genetic characterization of
cousins

[152532]with male pseudohermaphroditism: evidence that the Lubs
phenotype

[152533]can result from a mutation that alters the structure of the
androgen

[152534]receptor. Clin. Genet. 26: 363–370, 1984.

[152535]

[152536]135. Wooster, R.; Mangion, J.; Eeles, R.; Smith, S.;
Dowsett, M.;

[152537]Averill, D.; Barrett–Lee, P.; Easton, D. F.; Ponder, B. A. J.;
Stratton,

[152538]M. R.: A germline mutation in the androgen receptor gene
in two brothers

[152539]with breast cancer and Reifenstein syndrome. Nature
Genet. 2: 132–134,

[152540]1992.

[152541]

[152542]136. Zhang, L.; Leeflang, E. P.; Yu, J.; Arnheim, N.: Study-
ing human

[152543]mutations by sperm typing: instability of CAG trinucleotide repeats

[152544]in the human androgen receptor gene. Nature Genet. 7:
531–535, 1994.

[152545]

[152546]137. Zhu, Y.-S.; Cai, L.-Q.; Cordero, J. J.; Canovatchel, W.
J.; Katz,

[152547]M. D.; Imperato-McGinley, J.: A novel mutation in the CAG
triplet

[152548]region of exon 1 of androgen receptor gene causes complete androgen

[152549]insensitivity syndrome in a large kindred. J. Clin. Endocr.
Metab. 84:

[152550]1590–1594, 1999.

[152551]

[152552]138. Zoppi, S.; Wilson, C. M.; Harbison, M. D.; Griffin, J. E.;
Wilson,

[152553]J. D.; McPhaul, M. J.; Marcelli, M.: Complete testicular feminization

[152554]caused by an amino-terminal truncation of the androgen

receptor with

[152555]downstream initiation. J. Clin. Invest. 91: 1105–1112,
1993.

[152556]

[152557]

[152558]14623.

[152559]Konig, A.; Happle, R.; Fink–Puches, R.; Soyer, H. P.; Born–
holdt,

[152560]D.; Engel, H.; Grzeschik, K.–H.: A novel missense mutation
of NSDHL

[152561]in an unusual case of CHILD syndrome showing bilateral,
almost symmetric

[152562]involvement. J. Am. Acad. Derm. 46: 594–596, 2002.

[152563]

[152564]

[152565]14624.

[152566]Labit–Le Bouteiller, C.; Jamme, M. F.; David, M.; Silve, S.;
Lanau,

[152567]C.; Dhers, C.; Picard, C.; Rahier, A.; Taton, M.; Loison, G.;
Caput,

[152568]D.; Ferrara, P.; Lupker, J.: Antiproliferative effects of

SR31747A

[152569]in animal cell lines are mediated by inhibition of cholesterol biosynthesis

[152570]at the sterol isomerase step. *Europ. J. Biochem.* 256: 342–349, 1998.

[152571]

[152572]

[152573]14625.

[152574]Liu, X. Y.; Dangel, A. W.; Kelley, R. I.; Zhao, W.; Denny, P.;

[152575]Botcherby, M.; Cattanach, B.; Peters, J.; Hunsicker, P. R.; Mallon,

[152576]A.–M.; Strivens, M. A.; Bate, R.; Miller, W.; Rhodes, M.; Brown, S.

[152577]D. M.; Herman, G. E.: The gene mutated in bare patches and striated

[152578]mice encodes a novel 3–beta–hydroxysteroid dehydrogenase. *Nature*

[152579]*Genet.* 22: 182–187, 1999.

[152580]

[152581]

[152582]14626.

[152583]Schindelhauer, D.; Hellebrand, H.; Grimm, L.; Bader, I.;
Meitinger,

[152584]T.; Wehnert, M.; Ross, M.; Meindl, A.: Long-range map of
a 3.5-Mb

[152585]region in Xp11.23-22 with a sequence-ready map from a
1.1-Mb gene-rich

[152586]interval. *Genome Res.* 6: 1056-1069, 1996.

[152587]

[152588]

[152589]14627.

[152590]Silve, S.; Dupuy, P. H.; Labit-Lebouteiller, C.; Kaghad, M.;
Chalon,

[152591]P.; Rahier, A.; Taton, M.; Lupker, J.; Shire, D.; Loison, G.:
Emopamil-binding

[152592]protein, a mammalian protein that binds a series of struc-
turally diverse

[152593]neuroprotective agents, exhibits delta(8)-delta(7) sterol
isomerase

[152594]activity in yeast. *J. Biol. Chem.* 271: 22434-22440, 1996.

[152595]

[152596]

[152597]14628.

[152598]Traupe, H.; Muller, D.; Atherton, D.; Kalter, D. C.; Cremers,

[152599]F. P. M.; van Oost, B. A.; Ropers, H.-H.: Exclusion mapping
of the

[152600]X-linked dominant chondrodysplasia punctata/
ichthyosis/cataract/short

[152601]stature (Happle) syndrome: possible involvement of an
unstable pre-mutation. Hum.

[152602]Genet. 89: 659-665, 1992.

[152603]

[152604]

[152605]14629.

[152606]Jin, H.; Gardner, R. J.; Viswesvaraiah, R.; Muntoni, F.;
Roberts,

[152607]R. G.: Two novel members of the interleukin-1 receptor
gene family,

[152608]one deleted in Xp22.1-Xp21.3 mental retardation. Europ.
J. Hum. Genet. 8:

[152609]87-94, 2000.

[152610]

[152611]

[152612]14630.

[152613]Biamonti, G.; Ruggiu, M.; Saccone, S.; Della Valle, G.; Riva, S.

[152614]: Two homologous genes, originated by duplication, encode the human

[152615]hnRNP proteins A2 and A1. Nucleic Acids Res. 22: 1996–2002, 1994.

[152616]

[152617]

[152618]14631.

[152619]Stoddart, K. L.; Jermak, C.; Nagaraja, R.; Schlessinger, D.; Bech–Hansen,

[152620]N. T.: Physical map covering a 2 Mb region in human Xp11.3 distal

[152621]to DX6849. Gene 227: 111–116, 1999.

[152622]

[152623]

[152624]14632.

[152625]Swanson, D. A.; Freund, C. L.; Ploder, L.; McInnes, R. R.;

Valle,

[152626]D.: A ubiquitin C-terminal hydrolase gene on the proximal short arm

[152627]of the X chromosome: implications for X-linked retinal disorders. Hum.

[152628]Molec. Genet. 5: 533–538, 1996.

[152629]

[152630]

[152631]14633.

[152632]Narayanan, V.; Olinsky, S.; Dahle, E.; Naidu, S.; Zoghbi, H. Y.

[152633]: Mutation analysis of the M6b gene in patients with Rett syndrome. Am.

[152634]J. Med. Genet. 78: 165–168, 1998.

[152635]

[152636]

[152637]14634.

[152638]Olinsky, S.; Loop, B. T.; DeKosky, A.; Ripepi, B.; Weng, W.; Cummins,

[152639]J.; Wenger, S. L.; Yan, Y.; Lagenaur, C.; Narayanan, V.:

Chromosomal

[152640]mapping of the human M6 genes. Genomics 33: 532–536, 1996.

[152641]

[152642]

[152643]14635.

[152644]Yan, Y.; Lagenaur, C.; Narayanan, V.: Molecular cloning of M6:

[152645]identification of a PLP/DM20 gene family. Neuron 11: 423–431, 1993.

[152646]

[152647]

[152648]14636.

[152649]Ciccodicola, A.; D'Esposito, M.; Esposito, T.; Gianfrancesco, F.;

[152650]Migliaccio, C.; Miano, M. G.; Matarazzo, M. R.; Vacca, M.; Franze,

[152651]A.; Cuccurese, M.; Cocchia, M.; Curci, A.: Differentially regulated

[152652]and evolved genes in the fully sequenced Xq/Yq pseudoautosomal region. Hum.

[152653]Molec. Genet. 9: 395–401, 2000.

[152654]

[152655]

[152656]14637.

[152657]D'Esposito, M.; Ciccodicola, A.; Gianfrancesco, F.; Esposito, T.;

[152658]Flagiello, L.; Mazzarella, R.; Schlessinger, D.; D'Urso, M.: A synaptobrevin-like

[152659]gene in the Xq28 pseudoautosomal region undergoes X inactivation. Nature

[152660]Genet. 13: 227–229, 1996.

[152661]

[152662]

[152663]14638.

[152664]Muller, D. J.; Schulze, T. G.; Jahnes, E.; Cichon, S.; Krauss,

[152665]H.; Kesper, K.; Held, T.; Maier, W.; Propping, P.; Nothen, M. M.;

[152666]Rietschel, M.: Association between a polymorphism in the pseudoautosomal

[152667]X-linked gene SYBL1 and bipolar affective disorder. Am. J.

Med. Genet.

[152668](Neuropsychiat. Genet.) 114: 74–78, 2002.

[152669]

[152670]

[152671]14639.

[152672]Saito, T.; Parsia, S.; Papolos, D. F.; Lachman, H. M.: Analysis
sis

[152673]of the pseudoautosomal X-linked gene SYBL1 in bipolar
affective disorder:

[152674]description of a new candidate allele for psychiatric disorders. Am.

[152675]J. Med. Genet. (Neuropsychiat. Genet.) 96: 317–323, 2000.

[152676]

[152677]

[152678]14640.

[152679]Schaefer, L.; Ballabio, A.; Zoghbi, H. Y.: Cloning and characterization
acterization

[152680]of a putative human holocytochrome c-type synthetase
gene (HCCS) isolated

[152681]from the critical region for microphthalmia with linear skin

defects

[152682](MLS). Genomics 34: 166–172, 1996.

[152683]

[152684]

[152685]14641.

[152686]Van den Veyver, I. B.; Subramanian, S.; Zoghbi, H. Y.: Ge-
nomic

[152687]structure of a human holocytochrome c-type synthetase
gene in Xp22.3

[152688]and mutation analysis in patients with Rett syndrome. Am.
J. Med.

[152689]Genet. 78: 179–181, 1998.

[152690]

[152691]

[152692]14642.

[152693]Starace, D. M; Stefani, E.; Bezanilla, F.: Voltage-dependent
proton

[152694]transport by the voltage sensor of the Shaker K⁺ channel.
Neuron 19:

[152695]1319–1327, 1997.

[152696]

[152697]

[152698]14643.

[152699]Andersson, M.; Page, D. C.; Pettay, D.; Subrt, I.; Turleau, C.;

[152700]de Grouchy, J.; de la Chapelle, A.: Y;autosome translocations and

[152701]mosaicism in the aetiology of 45,X maleness: assignment of fertility

[152702]factor to distal Yq11. Hum. Genet. 79: 2–7, 1988.

[152703]

[152704]

[152705]14644.

[152706]Chandley, A. C.; Gosden, J. R.; Hargreave, T. B.; Spowart, G.;

[152707]Speed, R. M.; McBeath, S.: Deleted Yq in the sterile son of a man

[152708]with a satellited Y chromosome (Yqs). J. Med. Genet. 26: 145–153,

[152709]1989.

[152710]

[152711]

[152712]14645.

[152713]Franco, B.; Meroni, G.; Parenti, G.; Levilliers, J.; Bernard, L.;

[152714]Gebbia, M.; Cox, L.; Maroteaux, P.; Sheffield, L.; Rappold,
G. A.;

[152715]Andria, G.; Petit, C.; Ballabio, A.: A cluster of sulfatase
genes

[152716]on Xp22.3: mutations in chondrodysplasia punctata
(CDPX) and implications

[152717]for warfarin embryopathy. Cell 81: 1-20, 1995.

[152718]

[152719]

[152720]14646.

[152721]Meroni, G.; Franco, B.; Archidiacono, N.; Messali, S.; An-
dolfi,

[152722]G.; Rocchi, M.; Ballabio, A.: Characterization of a cluster of
sulfatase

[152723]genes on Xp22.3 suggests gene duplications in an ances-
tral pseudoautosomal

[152724]region. Hum. Molec. Genet. 5: 423–431, 1996.

[152725]

[152726]

[152727]14647.

[152728]Suh, Y.–A.; Arnold, R. S.; Lassegue, B.; Shi, J.; Xu, X.;
Sorescu,

[152729]D.; Chung, A. B.; Griendling, K. K.; Lambeth, J. D.: Cell
transformation

[152730]by the superoxide–generating oxidase Mox1. Nature 401:
79–82, 1999.

[152731]

[152732]

[152733]14648.

[152734]Lee, C. C.; Yazdani, A.; Wehnert, M.; Zhao, Z. Y.; Lindsay,
E.

[152735]A.; Bailey, J.; Coolbaugh, M. I.; Couch, L.; Xiong, M.; Chin–
ault,

[152736]A. C.; Baldini, A.; Caskey, C. T.: Isolation of chromosome–spe–
cific

[152737]genes by reciprocal probing of arrayed cDNA and cosmid
libraries. Hum.

[152738]Molec. Genet. 4: 1373–1380, 1995.

[152739]

[152740]

[152741]14649.

[152742]Patzak, D.; Zhuchenko, O.; Lee, C.–C.; Wehnert, M.: Identification,

[152743]mapping, and genomic structure of a novel X–
chromosomal human gene

[152744](SMPX) encoding a small muscular protein. Hum. Genet.
105: 506–512,

[152745]1999.

[152746]

[152747]

[152748]14650.

[152749]Ballabio, A.; Bardoni, B.; Carrozzo, R.; Andria, G.; Bick, D.;

[152750]Campbell, L.; Hamel, B.; Ferguson–Smith, M. A.; Gimelli,
G.; Fraccaro,

[152751]M.; Maraschio, P.; Zuffardi, O.; Guioli, S.; Camerino, G.:
Contiguous

[152752]gene syndromes due to deletions in the distal short arm

of the human

[152753]X chromosome. Proc. Nat. Acad. Sci. 86: 10001–10005,
1989.

[152754]

[152755]

[152756]14651.

[152757]Fukami, M.; Kirsch, S.; Schiller, S.; Richter, A.; Benes, V.;
Franco,

[152758]B.; Muroya, K.; Rao, E.; Merker, S.; Niesler, B.; Ballabio, A.;
Ansorge,

[152759]W.; Ogata, T.; Rappold, A.: A member of a gene family on
Xp22.3,

[152760]VCX–A, is deleted in patients with X–linked nonspecific
mental retardation. Am

[152761]J. Hum. Genet. 67: 563–573, 2000.

[152762]

[152763]

[152764]14652.

[152765]Lahn, B. T.; Page, D. C.: A human sex–chromosomal gene
family

[152766]expressed in male germ cells and encoding variably

charged proteins. Hum.

[152767]Molec. Genet. 9: 311–319, 2000.

[152768]

[152769]

[152770]14653.

[152771]Fujikawa–Adachi, K.; Nishimori, I.; Taguchi, T.; Onishi, S.:
Human

[152772]mitochondrial carbonic anhydrase VB: cDNA cloning,
mRNA expression,

[152773]subcellular localization, and mapping to chromosome X. J.
Biol. Chem. 274:

[152774]21228–21233, 1999.

[152775]

[152776]

[152777]14654.

[152778]Shah, G. N.; Hewett–Emmett, D.; Grubb, J. H.; Migas, M.
C.; Fleming,

[152779]R. E.; Waheed, A.; Sly, W. S.: Mitochondrial carbonic anhy-
drase CA

[152780]VB: differences in tissue distribution and pattern of evolu-

tion from

[152781]those of CA VA suggest distinct physiological roles. Proc.
Nat. Acad.

[152782]Sci. 97: 1677–1682, 2000.

[152783]

[152784]

[152785]14655.

[152786]Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayasi,
Y.;

[152787]Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura,
N.: Prediction

[152788]of the coding sequences of unidentified human genes. VI.
The coding

[152789]sequences of 80 new genes (KIAA0201–KIAA0280) de-
duced by analysis

[152790]of cDNA clones from cell line KG–1 and brain. DNA Res. 3:
321–329,

[152791]1996. Note: Supplement: DNA Res. 3: 341–354, 1996.

[152792]

[152793]

[152794]14656.

[152795]Numata, M.; Petrecca, K.; Lake, N.; Orlowski, J.: Identifica-
tion

[152796]of a mitochondrial Na⁺/H⁺ exchanger. J. Biol. Chem. 273:
6951-6959,

[152797]1998.

[152798]

[152799]

[152800]14657.

[152801]Buchner, G.; Orfanelli, U.; Quaderi, N.; Bassi, M. T.; An-
dolfi,

[152802]G.; Ballabio, A.; Franco, B.: Identification of a new EGF-
repeat-containing

[152803]gene from human Xp22: a candidate for developmental
disorders. Genomics 65:

[152804]16-23, 2000.

[152805]

[152806]

[152807]14658.

[152808]Yeung, G.; Mulero, J. J.; Berntsen, R. P.; Loeb, D. B.; Dr-
manac,

[152809]R.; Ford, J. E.: Cloning of a novel epidermal growth factor repeat

[152810]containing gene EGFL6: expressed in tumor and fetal tissues. Genomics 62:

[152811]304–307, 1999.

[152812]

[152813]

[152814]14659.

[152815]Asao, H.; Okuyama, C.; Kumaki, S.; Ishii, N.; Tsuchiya, S.; Foster,

[152816]D.; Sugamura, K.: Cutting edge: the common gamma-chain is an indispensable

[152817]subunit of the IL-21 receptor complex. J. Immun. 167: 1–5, 2001.

[152818]

[152819]

[152820]14660.

[152821]Lankes, W.; Griesmacher, A.; Grunwald, J.; Schwartz-Albiez, R.;

[152822]Keller, R.: A heparin-binding protein involved in inhibition of smooth-muscle

[152823]cell proliferation. Biochem. J. 251: 831–842, 1988.

[152824]

[152825]

[152826]14661.

[152827]Lankes, W. T.; Furthmayr, H.: Moesin: a member of the
protein

[152828]4.1–talin–ezrin family of proteins. Proc. Nat. Acad. Sci. 88:
8297–8301,

[152829]1991.

[152830]

[152831]

[152832]14662.

[152833]Shcherbina, A.; Bretscher, A.; Rosen, F. S.; Kenney, D. M.;
Remold–O'Donnell,

[152834]E.: The cytoskeletal linker protein moesin: decreased lev-
els in Wiskott–Aldrich

[152835]syndrome platelets and identification of a cleavage path-
way in normal

[152836]platelets. Brit. J. Haemat. 106: 216–223, 1999.

[152837]

[152838]

[152839]14663.

[152840]Wilgenbus, K. K.; Hsieh, C.-L.; Lankes, W. T.; Milatovich, A.;

[152841]Francke, U.; Furthmayr, H.: Structure and localization on the X chromosome

[152842]of the gene coding for the human filopodial protein moesin (MSN). Genomics 19:

[152843]326–333, 1994.

[152844]

[152845]

[152846]14664.

[152847]Deng, L.; Wang, C.; Spencer, E.; Yang, L.; Braun, A.; You, J.;

[152848]Slaughter, C.; Pickart, C.; Chen, Z. J.: Activation of the I- κ B

[152849]complex by TRAF6 requires a dimeric ubiquitin-conjugating enzyme complex

[152850]and a unique polyubiquitin chain. Cell 103: 351–361, 2000.

[152851]

[152852]

[152853]14665.

[152854]Wang, C.; Deng, L.; Hong, M.; Akkaraju, G. R.; Inoue, J.;
Chen,

[152855]Z. J.: TAK1 is a ubiquitin-dependent kinase of MKK and
IKK. Nature 412:

[152856]346–351, 2001.

[152857]

[152858]

[152859]14666.

[152860]Wong, B. R.; Besser, D.; Kim, N.; Arron, J. R.; Vologodskaia,
M.;

[152861]Hanafusa, H.; Choi, Y.: TRANCE, a TNF family member, ac-
tivates Akt/PKB

[152862]through a signaling complex involving TRAF6 and c-Src.
Molec. Cell 4:

[152863]1041–1049, 1999.

[152864]

[152865]

[152866]14667.

[152867]Nakano, H.; Oshima, H.; Chung, W.; Williams–Abbott, L.;
Ware, C.

[152868]F.; Yagita, H.; Okumura, K.: TRAF5, an activator of NF–
kappaB and

[152869]putative signal transducer for the lymphotoxin-beta receptor. J.

[152870]Biol. Chem. 271: 14661-14664, 1996.

[152871]

[152872]

[152873]14668.

[152874]Nakano, H.; Sakon, S.; Koseki, H.; Takemori, T.; Tada, K.;
Matsumoto,

[152875]M.; Munechika, E.; Sakai, T.; Shirasawa, T.; Akiba, H; Ko-
bata, T.;

[152876]Santee, S. M.; Ware, C. F.; Renner, P. D.; Taniguchi, M.;
Yagita,

[152877]H.; Okumura, K.: Targeted disruption of Traf5 gene causes
defects

[152878]in CD40- and CD27-mediated lymphocyte activation.
Proc. Nat. Acad.

[152879]Sci. 96: 9803-9808, 1999.

[152880]

[152881]

[152882]14669.

[152883]Nakano, H.; Shindo, M.; Yamada, K.; Yoshida, M. C.; Santee, S.

[152884]M.; Ware, C. F.; Jenkins, N. A.; Gilbert, D. J.; Yagita, H.; Copeland,

[152885]N. G.; Okumura, K.: Human TNF receptor-associated factor 5 (TRAF5):

[152886]cDNA cloning, expression and assignment of the TRAF5 gene to chromosome

[152887]1q32. Genomics 42: 26–32, 1997.

[152888]

[152889]

[152890]14670.

[152891]Yost, C.; Farr, G. H., III; Pierce, S. B.; Ferkey, D. M.; Chen,

[152892]M. M.; Kimelman, D.: GBP, an inhibitor of GSK–3, is implicated in

[152893]Xenopus development and oncogenesis. Cell 93: 1031–1041, 1998.

[152894]

[152895]

[152896]14671.

[152897]De Baere, E.; Speleman, F.; Van Roy, N.; De Paepe, A.;
Messiaen,

[152898]L.: Assignment of SHOX2 (alias OG12X and SHOT) to hu-
man chromosome

[152899]bands 3q25–q26.1 by in situ hybridization. Cytogenet.
Cell Genet. 82:

[152900]228–229, 1998.

[152901]

[152902]

[152903]14672.

[152904]Fernandez–Valle, C.; Tang, Y.; Ricard, J.; Rodenas–Ruano,
A.; Taylor,

[152905]A.; Hackler, E.; Biggerstaff, J.; Iacovelli, J.: Paxillin binds
schwannomin

[152906]and regulates its density–dependent localization and ef-
fect on cell

[152907]morphology. Nature Genet. 31: 354–362, 2002.

[152908]

[152909]

[152910]14673.

[152911]Glenney, J. R., Jr.; Zokas, L.: Novel tyrosine kinase substrates

[152912]from Rous sarcoma virus-transformed cells are present in the membrane

[152913]skeleton. J. Cell Biol. 108: 2401–2408, 1989.

[152914]

[152915]

[152916]14674.

[152917]Mazaki, Y.; Hashimoto, S.; Sabe, H.: Monocyte cells and cancer

[152918]cells express novel paxillin isoforms with different binding properties

[152919]to focal adhesion proteins. J. Biol. Chem. 272: 7437–7444, 1997.

[152920]

[152921]

[152922]14675.

[152923]Salgia, R.; Li, J.–L.; Lo, S. H.; Brunkhorst, B.; Kansas, G. S.;

[152924]Sobhany, E. S.; Sun, Y.; Pisick, E.; Hallek, M.; Ernst, T.; Tantravahi,

[152925]R.; Chen, L. B.; Griffin, J. D.: Molecular cloning of human paxillin,

[152926]a focal adhesion protein phosphorylated by P210(BCR/ABL). J. Biol.

[152927]Chem. 270: 5039–5047, 1995.

[152928]

[152929]

[152930]14676.

[152931]Turner, C. E.; Glenney, J. R., Jr.; Burridge, K.: Paxillin: a

[152932]new vinculin-binding protein present in focal adhesions. J. Cell

[152933]Biol. 111: 1059–1068, 1990.

[152934]

[152935]

[152936]14677.

[152937]Nguyen, V. T.; Kiss, T.; Michels, A. A.; Bensaude, O.: 7SK small

[152938]nuclear RNA binds to and inhibits the activity of CDK9/cyclin T complexes. Nature 414:

[152939]322–325, 2001.

[152940]

[152941]

[152942]14678.

[152943]Peng, J.; Zhu, Y.; Milton, J. T.; Price, D. H.: Identification

[152944]of multiple cyclin subunits of human P-TEFb. Genes Dev.

12: 755–762,

[152945]1998.

[152946]

[152947]

[152948]14679.

[152949]Wei, P.; Garber, M. E.; Fang, S.–M.; Fischer, W. H.; Jones, K.

[152950]A.: A novel CDK9-associated C-type cyclin interacts directly with

[152951]HIV–1 Tat and mediates its high–affinity, loop–specific binding to

[152952]TAR RNA. Cell 92: 451–462, 1998.

[152953]

[152954]

[152955]14680.

[152956]Yang, Z.; Zhu, Q.; Luo, K.; Zhou, Q.: The 7SK small nuclear
RNA

[152957]inhibits the CDK9/cyclin T1 kinase to control transcrip-
tion. Nature 414:

[152958]317–322, 2001.

[152959]

[152960]

[152961]14681.

[152962]Adachi, H.; Tsujimoto, M.; Hattori, M.; Arai, H.; Inoue, K.:
Differential

[152963]tissue distribution of the beta- and gamma-subunits of
human cytosolic

[152964]platelet-activating factor acetylhydrolase (isoform 1).
Biochem.

[152965]Biophys. Res. Commun. 233: 10–13, 1997.

[152966]

[152967]

[152968]14682.

[152969]Moro, F.; Arrigo, G.; Fogli, A.; Bernard, L.; Carrozzo, R.:
The

[152970]beta and gamma subunits of the human platelet-activating factor acetyl

[152971]hydrolase isoform Ib (PAFAH1B2 and PAFAH1B3) map to chromosome 11q23

[152972]and 19q13.1, respectively. Genomics 51: 157–159, 1998.

[152973]

[152974]

[152975]14683.

[152976]Daigo, Y.; Isomura, M.; Nishiwaki, T.; Tamari, M.;
Ishikawa, S.;

[152977]Kai, M.; Murata, Y.; Takeuchi, K.; Yamane, Y.; Hayashi, R.;
Minami,

[152978]M.; Fujino, M. A.; Hojo, Y.; Uchiyama, I.; Takagi, T.; Nakamura, Y.

[152979]: Characterization of a 1200-kb genomic segment of
chromosome 3p22–p21.3. DNA

[152980]Res. 6: 37–44, 1999.

[152981]

[152982]

[152983]14684.

[152984]Erlich, R.; Gleeson, P. A.; Campbell, P.; Dietzsch, E.; Toh, B.-H.

[152985]: Molecular characterization of trans-Golgi p230: a human peripheral

[152986]membrane protein encoded by a gene on chromosome 6p12-22 contains

[152987]extensive coiled-coil alpha-helical domains and a granin motif. J.

[152988]Biol. Chem. 271: 8328-8337, 1996.

[152989]

[152990]

[152991]14685.

[152992]Barr, F. A.; Nakamura, N.; Warren, G.: Mapping the interaction

[152993]between GRASP65 and GM130, components of a protein complex involved

[152994]in the stacking of Golgi cisternae. EMBO J. 17: 3258-3268, 1998.

[152995]

[152996]

[152997]14686.

[152998]Boguski, M. S.; Lowe, T. M. J.; Tolstochev, C. M.:

dbEST:database

[152999]for 'expressed sequence tags.' (Letter) Nature Genet. 4:

332–333,

[153000]1993.

[153001]

[153002]

[153003]14687.

[153004]Izon, D. J.; Aster, J. C.; He, Y.; Weng, A.; Karnell, F. G.; Pa-

triub,

[153005]V.; Xu, L.; Bakkour, S.; Rodriguez, C.; Allman, D.; Pear, W.

S.:

[153006]Deltex1 redirects lymphoid progenitors to the B cell lin-

eage by antagonizing

[153007]Notch1. Immunity 16: 231–243, 2002.

[153008]

[153009]

[153010]14688.

[153011]Lee, L.; Dowhanick–Morrissette, J.; Katz, A.; Jukofsky, L.;

Krantz,

[153012]I. D.: Chromosomal localization, genomic characterization,
and mapping

[153013]to the Noonan syndrome critical region of the human Del-
tex (DTX1)

[153014]gene. Hum. Genet. 107: 577–581, 2000.

[153015]

[153016]

[153017]14689.

[153018]Matsuno, K.; Eastman, D.; Mitsiades, T.; Quinn, A. M.; Car-
canciu,

[153019]M. L.; Ordentlich, P.; Kadesch, T.; Artavanis-Tsakonas, S.:
Human

[153020]deltex is a conserved regulator of Notch signalling. Nature
Genet. 19:

[153021]74–78, 1998.

[153022]

[153023]

[153024]14690.

[153025]Chen, J. J.; Reid, C. E.; Band, V.; Androphy, E. J.: Interaction

[153026]of papillomavirus E6 oncoproteins with a putative cal-

cium-binding

[153027]protein. Science 269: 529-531, 1995.

[153028]

[153029]

[153030]14691.

[153031]Imai, T.; Matsuda, K.; Shimojima, T.; Hashimoto, T.; Masuhiro,

[153032]Y.; Kitamoto, T.; Sugita, A.; Suzuki, K.; Matsumoto, H.; Masushige,

[153033]S.; Nogi, Y.; Muramatsu, M.; Handa, H.; Kato, S.: ERC-55, a binding

[153034]protein for the papilloma virus E6 oncoprotein, specifically interacts

[153035]with vitamin D receptor among nuclear receptors. Biochem. Biophys.

[153036]Res. Commun. 233: 765-769, 1997.

[153037]

[153038]

[153039]14692.

[153040]Wang, J. Y.; Zhen, D. K.; Bianchi, D. W.; Androphy, E. J.;

Chen,

[153041]J. J.: Assignment of the gene for ERC-55 (RCN2) to human chromosome

[153042]band 15q22.33-q24.1 by in situ hybridization. Cytogenet. Cell Genet. 79:

[153043]60-61, 1997.

[153044]

[153045]

[153046]14693.

[153047]Weis, K.; Griffiths, G.; Lamond, A. I.: The endoplasmic reticulum

[153048]calcium-binding protein of 55 kDa is a novel EF-hand protein retained

[153049]in the endoplasmic reticulum by a carboxyl-terminal His-Asp-Glu-Leu

[153050]motif. J. Biol. Chem. 269: 19142-19150, 1994.

[153051]

[153052]

[153053]14694.

[153054]Dempsey, C. E.; Sakurai, H.; Sugita, T.; Guesdon, F.: Alter-

native

[153055]splicing and gene structure of the transforming growth
factor beta-activated

[153056]kinase 1. Biochim. Biophys. Acta 1517: 46-52, 2000.

[153057]

[153058]

[153059]14695.

[153060]Kondo, M.; Osada, H.; Uchida, K.; Yanagisawa, K.; Masuda,
A.; Takagi,

[153061]K.; Takahashi, T.; Takahashi, T.: Molecular cloning of hu-
man TAK1

[153062]and its mutational analysis in human lung cancer. Int. J.
Cancer 75:

[153063]559-563, 1998.

[153064]

[153065]

[153066]14696.

[153067]Sakurai, H.; Shigemori, N.; Hasegawa, K.; Sugita, T.: TGF-
beta-activated

[153068]kinase 1 stimulates NF-kappa-B activation by an NF-

kappa-B-inducing

[153069]kinase-independent mechanism. Biochem. Biophys. Res.

Commun. 243:

[153070]545-549, 1998.

[153071]

[153072]

[153073]14697.

[153074]Yamaguchi, K.; Shirakabe, K.; Shibuya, H.; Irie, K.; Oishi, I.;

[153075]Ueno, N.; Taniguchi, T.; Nishida, E.; Matsumoto, K.: Iden-
tification

[153076]of a member of the MAPKKK family as a potential mediator
of TGF-beta

[153077]signal transduction. Science 270: 2008-2011, 1995.

[153078]

[153079]

[153080]14698.

[153081]Chan, A. M.-L.; Chedid, M.; McGovern, E. S.; Popescu, N.
C.; Miki,

[153082]T.; Aaronson, S. A.: Expression cDNA cloning of a serine
kinase transforming

[153083]gene. Oncogene 8: 1329–1333, 1993.

[153084]

[153085]

[153086]14699.

[153087]Garcia-Barcelo, M.; Law, P. T. W.; Tsui, S. K. W.; Fung, K.
P.;

[153088]Lee, C. Y.; Waye, M. M. Y.: Mapping of the human riboso-
mal large

[153089]subunit protein gene RPL29 to human chromosome
3q29–qter. Genomics 46:

[153090]148–151, 1997.

[153091]

[153092]

[153093]14700.

[153094]Kirn-Safran, C. B.; Dayal, S.; Martin-DeLeon, P. A.; Carson,
D.

[153095]D.: Cloning, expression, and chromosome mapping of the
murine Hip/Rpl29

[153096]gene. Genomics 68: 210–219, 2000.

[153097]

[153098]

[153099]14701.

[153100]Law, P. T. W.; Tsui, S. K. W.; Lam, W. Y.; Luk, S. C. W.;

Hwaang,

[153101]D. M.; Liew, C. C.; Lee, C. Y.; Fung, K. P.; Waye, M. M. Y.: A

novel

[153102]cDNA encoding a human homologue of ribosomal protein

L29. Biochim.

[153103]Biophys. Acta 1305: 105–108, 1996.

[153104]

[153105]

[153106]14702.

[153107]Liu, S.; Smith, S. E.; Julian, J.; Rohde, L. H.; Karin, N. J.;

[153108]Carson, D. D.: cDNA cloning and expression of HIP, a

novel cell surface

[153109]heparan sulfate/heparin-binding protein of human uterine

epithelial

[153110]cells and cell lines. J. Biol. Chem. 271: 11817–11823,

1996.

[153111]

[153112]

[153113]14703.

[153114]Rohde, L. H.; Julian, J.; Babaknia, A.; Carson, D. D.: Cell
surface

[153115]expression of HIP, a novel heparin/heparan sulfate-bind-
ing protein,

[153116]of human uterine epithelial cells and cell lines. J. Biol.
Chem. 271:

[153117]11824–11830, 1996.

[153118]

[153119]

[153120]14704.

[153121]Frank, K. M.; Sekiguchi, J. M.; Seidl, K. J.; Swat, W.; Rath-
bun,

[153122]G. A.; Cheng, H.-L.; Davidson, L.; Kangaloo, L.; Alt, F. W.:
Late

[153123]embryonic lethality and impaired V(D)J recombination in
mice lacking

[153124]DNA ligase IV. Nature 396: 173–177, 1998.

[153125]

[153126]

[153127]14705.

[153128]Frank, K. M.; Sharpless, N. E.; Gao, Y.; Sekiguchi, J. M.;

Ferguson,

[153129]D. O.; Zhu, C.; Manis, J. P.; Horner, J.; DePinho, R. A.; Alt,

F.

[153130]W.: DNA ligase IV deficiency in mice leads to defective
neurogenesis

[153131]and embryonic lethality via the p53 pathway. Molec. Cell
5: 993–1002,

[153132]2000.

[153133]

[153134]

[153135]14706.

[153136]Grawunder, U.; Zimmer, D.; Fugmann, S.; Schwarz, K.;

Lieber, M.

[153137]R.: DNA ligase IV is essential for V(D)J recombination and
DNA double-strand

[153138]break repair in human precursor lymphocytes. Molec. Cell
2: 477–484,

[153139]1998.

[153140]

[153141]

[153142]14707.

[153143]O'Driscoll, M.; Cerosaletti, K. M.; Girard, P.-M.; Dai, Y.;
Stumm,

[153144]M.; Kysela, B.; Hirsch, B.; Gennery, A.; Palmer, S. E.; Seidel,
J.;

[153145]Gatti, R. A.; Varon, R.; Oettinger, M. A.; Neitzel, H.; Jeggo,
P.

[153146]A.; Concannon, P.: DNA ligase IV mutations identified in
patients

[153147]exhibiting developmental delay and immunodeficiency.
Molec. Cell 8:

[153148]1175–1185, 2001.

[153149]

[153150]

[153151]14708.

[153152]Riballo, E.; Critchlow, S. E.; Teo, S.-H.; Doherty, A. J.;
Priestley,

[153153]A.; Broughton, B.; Kysela, B.; Beamish, H.; Plowman, N.;
Arlett, C.

[153154]F.; Lehmann, A. R.; Jackson, S. P.; Jeggo, P. A.: Identifica-
tion

[153155]of a defect in DNA ligase IV in a radiosensitive leukaemia
patient. Curr.

[153156]Biol. 9: 699–702, 1999.

[153157]

[153158]

[153159]14709.

[153160]Riballo, E.; Doherty, A. J.; Dai, Y.; Stiff, T.; Oettinger, M.

[153161]A.; Jeggo, P. A.; Kysela, B.: Cellular and biochemical im-
pact of

[153162]a mutation in DNA ligase IV conferring clinical radiosensi-
tivity. J.

[153163]Biol. Chem. 276: 31124–31132, 2001.

[153164]

[153165]

[153166]14710.

[153167]Giachino, C.; Lantelme, E.; Lanzetti, L.; Saccone, S.; Della
Valle,

[153168]G.; Migone, N.: A novel SH3-containing human gene fam-

ily preferentially

[153169]expressed in the central nervous system. Genomics 41:
427–434, 1997.

[153170]

[153171]

[153172]14711.

[153173]Furlan, M.; Robles, R.; Galbusera, M.; Remuzzi, G.; Kyrle,
P. A.;

[153174]Brenner, B.; Krause, M.; Scharrer, I.; Aumann, V.; Mittler,
U.; Solenthaler,

[153175]M.; Lammle, B.: Von Willebrand factor–cleaving protease in
thrombotic

[153176]thrombocytopenic purpura and the hemolytic–uremic syn-
drome. New Eng.

[153177]J. Med. 339: 1578–1584, 1998.

[153178]

[153179]

[153180]14712.

[153181]Furlan, M.; Robles, R.; Lammle, B.: Partial purification and
characterization

[153182]of a protease from human plasma cleaving von Willebrand

factor to

[153183]fragments produced by in vivo proteolysis. Blood 87:
4223–4234,

[153184]1996.

[153185]

[153186]

[153187]14713.

[153188]Nichols, A. F.; Ong, P.; Linn, S.: Mutations specific to the
xeroderma

[153189]pigmentosum group E Ddb– phenotype. J. Biol. Chem.
271: 24317–24320,

[153190]1996.

[153191]

[153192]

[153193]14714.

[153194]Shisheva, A.; Sudhof, T. C.; Czech, M. P.: Cloning, charac-
terization,

[153195]and expression of a novel GDP dissociation inhibitor iso-
form from

[153196]skeletal muscle. Molec. Cell. Biol. 14: 3459–3468, 1994.

[153197]

[153198]

[153199]14715.

[153200]Grieff, M.; Whyte, M. P.; Thakker, R. V.; Mazzaella, R.: Sequence
analysis of 139 kb in Xp22.1 containing spermine syn-

[153201]thase and the
5-prime region of PEX. Genomics 44: 227-231, 1997.

[153203]

[153204]

[153205]14716.

[153206]Korhonen, V. P.; Halmekyto, M.; Kauppinen, L.; Myohanen,
S.; Wahlfors,

[153207]J.; Keinanen, T.; Hyvonen, T.; Alhonen, L.; Eloranta, T.;
Janne, J.

[153208]: Molecular cloning of a cDNA encoding human spermine
synthase. DNA

[153209]Cell Biol. 14: 841-847, 1995.

[153210]

[153211]

[153212]14717.

[153213]Fathi, Z.; Corjay, M. H.; Shapira, H.; Wada, E.; Benya, R.;
Jensen,

[153214]R.; Viallet, J.; Sausville, E. A.; Battey, J. F.: BRS-3: a novel

[153215]bombesin receptor subtype selectively expressed in testis
and lung

[153216]carcinoma cells. J. Biol. Chem. 268: 5979-5984, 1993.

[153217]

[153218]

[153219]14718.

[153220]Gorbulev, V.; Akhundova, A.; Grzeschik, K. H.; Fahrenholz,
F.:

[153221]Organization and chromosomal localization of the gene
for the human

[153222]bombesin receptor subtype expressed in pregnant uterus.
FEBS Lett. 340:

[153223]260-264, 1994.

[153224]

[153225]

[153226]14719.

[153227]Ohki-Hamazaki, H.; Watase, K.; Yamamoto, K.; Ogura, H.; Yamano,

[153228]M.; Yamada, K.; Maeno, H.; Imaki, J.; Kikuyama, S.; Wada, E.; Wada,

[153229]K.: Mice lacking bombesin receptor subtype-3 develop metabolic defects

[153230]and obesity. Nature 390: 166-169, 1997.

[153231]

[153232]

[153233]14720.

[153234]Banfi, S.; Borsani, G.; Bulfone, A.; Ballabio, A.: Drosophila-related

[153235]expressed sequences. Hum. Molec. Genet. 6: 1745-1753, 1997.

[153236]

[153237]

[153238]14721.

[153239]Castrillon, D. H.; Wasserman, S. A.: Diaphanous is required for

[153240]cytokinesis in Drosophila and shares domains of similarity with the

[153241]products of the limb deformity gene. Development 120:
3367–3377,

[153242]1994.

[153243]

[153244]

[153245]14722.

[153246]Philippe, C.; Cremers, F. P. M.; Chery, M.; Bach, I.; Abbadi,
N.;

[153247]Ropers, H. H.; Gilgenkrantz, S.: Physical mapping of DNA
markers

[153248]in the q13–q22 region of the human X chromosome. Ge-
nomics 17: 147–152,

[153249]1993.

[153250]

[153251]

[153252]14723.

[153253]Bione, S.; Sala, C.; Manzini, C.; Arrigo, G.; Zuffardi, O.;
Banfi,

[153254]S.; Borsani, G.; Jonveaux, P.; Philippe, C.; Zuccotti, M.; Bal-
labio,

[153255]A.; Toniolo, D.: A human homologue of the *Drosophila melanogaster*

[153256]diaphanous gene is disrupted in a patient with premature ovarian failure:

[153257]evidence for conserved function in oogenesis and implications for

[153258]human sterility. *Am. J. Hum. Genet.* 62: 533–541, 1998.

[153259]

[153260]

[153261]14724.

[153262]Sala, C.; Arrigo, G.; Torri, G.; Martinazzi, F.; Riva, P.; Larizza,

[153263]L.; Philippe, C.; Jonveaux, P.; Sloan, F.; Labella, T.; Toniolo, D.

[153264]: Eleven X chromosome breakpoints associated with premature ovarian

[153265]failure (POF) map to a 15-Mb YAC contig spanning Xq21. *Genomics* 40:

[153266]123–131, 1997.

[153267]

[153268]

[153269]14725.

[153270]Hanaoka, F.; Tandai, M.; Miyazawa, H.; Murakami, Y.; Hori, T.;

[153271]Yamada, M.: Assignment of human DNA polymerase alpha gene (POLA)

[153272]to the X chromosome. (Abstract) Cytogenet. Cell Genet. 40: 647 only,

[153273]1985.

[153274]

[153275]

[153276]14726.

[153277]Miyazawa, H.; Tandai, M.; Hanaoka, F.; Yamada, M.; Hori, T.; Shimizu,

[153278]K.; Sekiguchi, M.: Identification of a DNA segment containing the

[153279]human DNA polymerase alpha gene. Biochem. Biophys. Res. Commun. 139:

[153280]637-643, 1986.

[153281]

[153282]

[153283]14727.

[153284]Wang, T. S.-F.; Pearson, B. E.; Suomalainen, H. A.; Mohan-
das, T.;

[153285]Shapiro, L. J.; Schroder, J.; Korn, D.: Assignment of the
gene for

[153286]human DNA polymerase alpha to the X chromosome. Proc.
Nat. Acad.

[153287]Sci. 82: 5270-5274, 1985.

[153288]

[153289]

[153290]14728.

[153291]Watson, J. M.; Spencer, J. A.; Riggs, A. D.; Graves, J. A. M.:

[153292]Sex chromosome evolution: platypus gene mapping sug-
gests that part

[153293]of the human X chromosome was originally autosomal.
Proc. Nat. Acad.

[153294]Sci. 88: 11256-11260, 1991.

[153295]

[153296]

[153297]14729.

[153298]Lachner, M.; O'Carroll, D.; Rea, S.; Mechtler, K.; Jenuwein, T.

[153299]: Methylation of histone H3 lysine 9 creates a binding site for HP1

[153300]proteins. Nature 410: 116–120, 2001.

[153301]

[153302]

[153303]14730.

[153304]Melcher, M.; Schmid, M.; Aagaard, L.; Selenko, P.; Laible, G.;

[153305]Jenuwein, T.: Structure–function analysis of SUV39H1 reveals a dominant

[153306]role in heterochromatin organization, chromosome segregation, and

[153307]mitotic progression. Molec. Cell Biol. 20: 3728–3841, 2000.

[153308]

[153309]

[153310]14731.

[153311]Nielsen, S. J.; Schneider, R.; Bauer, U.–M.; Bannister, A. J.;

- [153312]Morrison, A.; O'Carroll, D.; Firestein, R.; Cleary, M.; Jenuwein, T.; Herrera, R. E.; Kouzarides, T.: Rb targets histone H3 methylation and HP1 to promoters. Nature 412: 561–565, 2001.
- [153315]
- [153316]
- [153317]14732.
- [153318]Peters, A. H. F. M.; O'Carroll, D.; Scherthan, H.; Mechtler, K.; Sauer, S.; Schofer, C.; Weipoltshammer, K.; Pagani, M.; Lachner, M.; Kohlmaier, A.; Opravil, S.; Doyle, M.; Sibilia, M.; Jenuwein, T.: Loss of the Suv39h histone methyltransferases impairs mammalian heterochromatin and genome stability. Cell 107: 323–337, 2001.
- [153323]
- [153324]
- [153325]14733.

[153326]Rea, S.; Eisenhaber, F.; O'Carroll, D.; Strahl, B. D.; Sun, Z.-W.;

[153327]Schmid, M.; Opravil, S.; Mechtler, K.; Ponting, C. P.; Allis, C. D.;

[153328]Jenuwein, T.: Regulation of chromatin structure by site-specific

[153329]histone H3 methyltransferases. Nature 406: 593–599, 2000.

[153330]

[153331]

[153332]14734.

[153333]Scott, A. F.: Personal Communication. Baltimore, Md. 8/7/2000.

[153334]

[153335]

[153336]14735.

[153337]Haltiwanger, R. S.; Blomberg, M. A.; Hart, G. W.: Glycosylation

[153338]of nuclear and cytoplasmic proteins: purification and characterization

[153339]of a uridine diphospho-N-acetylglucosamine:polypeptide

beta-N-acetylglucosaminyltransferase. J.

[153340]Biol. Chem. 267: 9005–9013, 1992.

[153341]

[153342]

[153343]14736.

[153344]Kreppel, L. K.; Blomberg, M. A.; Hart, G. W.: Dynamic glycosylation

[153345]of nuclear and cytosolic proteins: cloning and characterization of

[153346]a unique O-GlcNAc transferase with multiple tetratricopeptide repeats. J.

[153347]Biol. Chem. 272: 9308–9315, 1997.

[153348]

[153349]

[153350]14737.

[153351]Lubas, W. A.; Frank, D. W.; Krause, M.; Hanover, J. A.: O-linked

[153352]GlcNAc transferase is a conserved nucleocytoplasmic protein containing

[153353]tetratricopeptide repeats. J. Biol. Chem. 272: 9316–9324,

1997.

[153354]

[153355]

[153356]14738.

[153357]Shafi, R.; Iyer, S. P. N.; Ellies, L. G.; O'Donnell, N.; Marek,

[153358]K. W.; Chui, D.; Hart, G. W.; Marth, J. D.: The O-GlcNAc
transferase

[153359]gene resides on the X chromosome and is essential for
embryonic stem

[153360]cell viability and mouse ontogeny. Proc. Nat. Acad. Sci. 97:
5735–5739,

[153361]2000.

[153362]

[153363]

[153364]14739.

[153365]Yang, X.; Zhang, F.; Kudlow, J. E.: Recruitment of O-
GlcNAc transferase

[153366]to promoters by corepressor mSin3A: coupling protein O-
GlcNAcylation

[153367]to transcriptional repression. Cell 110: 69–80, 2002.

[153368]

[153369]

[153370]14740.

[153371]Furuta, S.; Kobayashi, A.; Miyazawa, S.; Hashimoto, T.:

Cloning

[153372]and expression of cDNA for a newly identified isozyme of
bovine liver

[153373]3-hydroxyacyl-CoA dehydrogenase and its import into
mitochondria. Biochim.

[153374]Biophys. Acta 1350: 317-324, 1997.

[153375]

[153376]

[153377]14741.

[153378]Hansis, C.; Jahner, D.; Spiess, A.-N.; Boettcher, K.; Ivell, R.

[153379]: The gene for the Alzheimer-associated beta-amy-
loid-binding protein

[153380](ERAB) is differentially expressed in the testicular Leydig
cells

[153381]of the azoospermic by w/w(v) mouse. Europ. J. Biochem.
258: 53-60,

[153382]1998.

[153383]

[153384]

[153385]14742.

[153386]He, X.-Y.; Schulz, H.; Yang, S.-Y.: A human brain L-3-hydroxyacyl-coenzyme

[153387]A dehydrogenase is identical to an amyloid beta-peptide-binding protein

[153388]involved in Alzheimer's disease. J. Biol. Chem. 273: 10741-10746,

[153389]1998.

[153390]

[153391]

[153392]14743.

[153393]Yan, S. D.; Fu, J.; Soto, C.; Chen, X.; Zhu, H.; Al-Mohanna, F.;

[153394]Collison, K.; Zhu, A.; Stern, E.; Saido, T.; Tohyama, M.; Ogawa, S.;

[153395]Roher, A.; Stern, D.: An intracellular protein that binds amyloid-beta

[153396]peptide and mediates neurotoxicity in Alzheimer's disease. Nature 389:

[153397]689–695, 1997.

[153398]

[153399]

[153400]14744.

[153401]Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Wilson, W. G.; Leonard,

[153402]J. V.; Chou, J. Y.: Demyelination of the brain is associated with

[153403]methionine adenosyltransferase I/III deficiency. J. Clin. Invest. 98:

[153404]1021–1027, 1996.

[153405]

[153406]

[153407]14745.

[153408]Nishiyama, K.; Funai, T.; Katafuchi, R.; Hattori, F.; Onoyama,

[153409]K.; Ichiyama, A.: Primary hyperoxaluria type I due to a point mutation

[153410]of T to C in the coding region of the serine:pyruvate
aminotransferase

[153411]gene. Biochem. Biophys. Res. Commun. 176: 1093–1099,
1991.

[153412]

[153413]

[153414]14746.

[153415]Pirulli, D.; Puzzer, D.; Ferri, L.; Crovella, S.; Amoroso, A.;

[153416]Ferrettini, C.; Marangella, M.; Mazzola, G.; Florian, F.:
Molecular

[153417]analysis of hyperoxaluria type 1 in Italian patients reveals
eight

[153418]new mutations in the alanine:glyoxylate aminotransferase
gene. Hum.

[153419]Genet. 104: 523–525, 1999.

[153420]

[153421]

[153422]14747.

[153423]Watts, R. W. E.; Calne, R. Y.; Rolles, K.; Danpure, C. J.;
Morgan,

[153424]S. H.; Mansell, M. A.; Williams, R.; Purkiss, P.: Successful treatment

[153425]of primary hyperoxaluria type I by combined hepatic and renal transplantation. Lancet II:

[153426]474–475, 1987.

[153427]

[153428]

[153429]14748.

[153430]Cramer, S. D.; Ferree, P. M.; Lin, K.; Milliner, D. S.; Holmes,

[153431]R. P.: The gene encoding hydroxypyruvate reductase (GRHPR) is mutated

[153432]in patients with primary hyperoxaluria type II. Hum. Molec. Genet. 8:

[153433]2063–2069, 1999.

[153434]

[153435]

[153436]14749.

[153437]Natt, E.; Kida, K.; Odievre, M.; Di Rocco, M.; Scherer, G.: Point

[153438]mutations in the tyrosine aminotransferase gene in ty-

rosinemia type

[153439]II. Proc. Nat. Acad. Sci. 89: 9297–9301, 1992.

[153440]

[153441]

[153442]14750.

[153443]Johnson, W. G.: Metabolic interference and the +/- heterozygote:

[153444]a hypothetical form of simple inheritance which is neither dominant

[153445]nor recessive. Am. J. Hum. Genet. 32: 374–386, 1980.

[153446]

[153447]

[153448]14751.

[153449]Brenner, V.; Nyakatura, G.; Rosenthal, A.; Platzer, M.: Genomic

[153450]organization of two novel genes on human Xq28: compact head to head

[153451]arrangement of IDH-gamma and TRAP-delta is conserved in rat and mouse. Genomics 44:

[153452]8–14, 1997.

[153453]

[153454]

[153455]14752.

[153456]Achen, M. G.; Jeltsch, M.; Kukk, E.; Makinen, T.; Vitali, A.;
Wilks,

[153457]A. F.; Alitalo, K.; Stacker, S. A.: Vascular endothelial
growth factor

[153458]D (VEGF-D) is a ligand for the tyrosine kinases VEGF re-
ceptor 2 (Flk1)

[153459]and VEGF receptor 3 (Flt4). Proc. Nat. Acad. Sci. 95:
548-553, 1998.

[153460]

[153461]

[153462]14753.

[153463]Stacker, S. A.; Caesar, C.; Baldwin, M. E.; Thornton, G. E.;
Williams,

[153464]R. A.; Prevo, R.; Jackson, D. G.; Nishikawa, S.; Kubo, H.;
Achen,

[153465]M. G.: VEGF-D promotes the metastatic spread of tumor
cells via the

[153466]lymphatics. Nature Med. 7: 186-191, 2001.

[153467]

[153468]

[153469]14754.

[153470]Yamada, Y.; Nezu, J.; Shimane, M.; Hirata, Y.: Molecular
cloning

[153471]of a novel vascular endothelial growth factor, VEGF-D. Ge-
nomics 42:

[153472]483–488, 1997.

[153473]

[153474]

[153475]14755.

[153476]Davies, P. A.; Hanna, M. C.; Hales, T. G.; Kirkness, E. F.:
Insensitivity

[153477]to anaesthetic agents conferred by a class of GABA(A) re-
ceptor subunit. Nature 385:

[153478]820–823, 1997.

[153479]

[153480]

[153481]14756.

[153482]Sinkkonen, S. T.; Hanna, M. C.; Kirkness, E. F.; Korpi, E. R.:

[153483]GABA-A receptor epsilon and theta subunits display unusual structural

[153484]variation between species and are enriched in the rat locus ceruleus. J.

[153485]Neurosci. 20: 3588–3595, 2000.

[153486]

[153487]

[153488]14757.

[153489]Wilke, K.; Gaul, R.; Klauck, S. M.; Poustka, A.: A gene in human

[153490]chromosome band Xq28 (GABRE) defines a putative new subunit class

[153491]of the GABA(A) neurotransmitter receptor. Genomics 45: 1–10, 1997.

[153492]

[153493]

[153494]14758.

[153495]Gecz, J.; Baker, E.; Donnelly, A.; Ming, J. E.; McDonald-McGinn,

[153496]D. M.; Spinner, N. B.; Zackai, E. H.; Sutherland, G. R.; Mulley, J.

[153497]C.: Fibroblast growth factor homologous factor 2 (FHF2): gene structure,

[153498]expression and mapping to the Borjeson–Forssman–Lehmann syndrome region

[153499]in Xq26 delineated by a duplication breakpoint in a BFLS-like patient. Hum.

[153500]Genet. 104: 56–63, 1999.

[153501]

[153502]

[153503]14759.

[153504]Lovec, H.; Hartung, H.; Verdier, A.–S.; Mattei, M.–G.; Birnbaum,

[153505]D.; Goldfarb, M.; Coulier, F.: Assignment of FGF13 to human chromosome

[153506]band Xq21 by in situ hybridization. Cytogenet. Cell Genet. 76: 183–184,

[153507]1997.

[153508]

[153509]

[153510]14760.

[153511]Smallwood, P. M.; Munoz-Sanjuan, I.; Tong, P.; Macke, J. P.; Hendry,

[153512]S. H. C.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.; Nathans,

[153513]J.: Fibroblast growth factor (FGF) homologous factors: new members

[153514]of the FGF family implicated in nervous system development. Proc.

[153515]Nat. Acad. Sci. 93: 9850–9857, 1996.

[153516]

[153517]

[153518]14761.

[153519]Jones, M. H.; Furlong, R. A.; Burkin, H.; Chalmers, I. J.; Brown,

[153520]G. M.; Khwaja, O.; Affara, N. A.: The Drosophila developmental gene

[153521]fat facets has a human homologue in Xp11.4 which escapes X-inactivation

[153522]and has related sequences on Yq11.2. Hum. Molec. Genet. 5: 1695–1701,

[153523]1996.

[153524]

[153525]

[153526]14762.

[153527]Stoffers, D. A.; Zinkin, N. T.; Stanojevic, V.; Clarke, W. L.;

[153528]Habener, J. F.: Pancreatic agenesis attributable to a single
nucleotide

[153529]deletion in the human IPF1 gene coding sequence. Nature
Genet. 15:

[153530]106–110, 1997.

[153531]

[153532]

[153533]14763.

[153534]Theophilos, M. B.; Cox, D. W.; Mercer, J. F. B.: The toxic
milk

[153535]mouse is a murine model of Wilson disease. Hum. Molec.
Genet. 5:

[153536]1619–1624, 1996.

[153537]

[153538]

[153539]14764.

[153540]Natt, E.; Westphal, E.-M.; Toth-Fejel, S. E.; Magenis, R. E.;

[153541]Buist, N. R. M.; Rettenmeier, R.; Scherer, G.: Inherited and
de novo

[153542]deletion of the tyrosine aminotransferase gene locus at
16q22.1-q22.3

[153543]in a patient with tyrosinemia type II. Hum. Genet. 77:
352-358,

[153544]1987.

[153545]

[153546]

[153547]14765.

[153548]Pelet, B.; Antener, I.; Faggioni, R.; Spahr, A.; Gautier, E.:

[153549]Tyrosinemia without liver or renal damage with plantar
and palmar

[153550]keratosis and keratitis (hypertyrosinemia type II). Helv.
Paediat.

[153551]Acta 34: 177-183, 1979.

[153552]

[153553]

[153554]14766.

[153555]Rehak, A.; Selim, M. M.; Yadav, G.: Richner–Hanhart syndrome

[153556](tyrosinaemia–II) (report of four cases without ocular involvement). Brit.

[153557]J. Derm. 104: 469–475, 1981.

[153558]

[153559]

[153560]14767.

[153561]Rettenmeier, R.; Natt, E.; Zentgraf, H.; Scherer, G.: Isolation

[153562]and characterization of the human tyrosine aminotransferase gene. Nucleic

[153563]Acids Res. 18: 3853–3861, 1990.

[153564]

[153565]

[153566]14768.

[153567]Richner, H.: Hornhautaffektion bei Keratoma palmare et

plantare

[153568]hereditarium. Klin. Mbl. Augenheilk. 100: 580–588, 1938.

[153569]

[153570]

[153571]14769.

[153572]Schmid, W.; Muller, G.; Schutz, G.; Gluecksohn–Waelsch,
S.: Deletions

[153573]near the albino locus on chromosome 7 of the mouse af-
fect the level

[153574]of tyrosine aminotransferase mRNA. Proc. Nat. Acad. Sci.
82: 2866–2869,

[153575]1985.

[153576]

[153577]

[153578]14770.

[153579]Tallab, T.M.: Richner–Hanhart syndrome: importance of
early diagnosis

[153580]and early intervention. J. Am. Acad. Dermatol. 35:
857–859, 1996.

[153581]

[153582]

[153583]14771.

[153584]Ventura, G.; Biasini, G.; Petrozzi, M.: Cheratomia palmo-
plantare

[153585]dissipatum associato a lesioni corneali in due fratelli. Boll.
Oculist. 44:

[153586]497-510, 1965.

[153587]

[153588]

[153589]14772.

[153590]Waardenburg, P. J.; Franceschetti, A.; Klein, D.: Genetics
and

[153591]Ophthalmology. Springfield, Ill.: Charles C Thomas (pub.)
1:

[153592]1961. Pp. 515-517.

[153593]

[153594]

[153595]14773.

[153596]Westphal, E.-M.; Natt, E.; Grimm, T.; Odievre, M.; Scherer,

G.

[153597]: The human tyrosine aminotransferase gene: characterization of restriction

[153598]fragment length polymorphisms and haplotype analysis in a family with

[153599]tyrosinemia type II. Hum. Genet. 79: 260–264, 1988.

[153600]

[153601]

[153602]14774.

[153603]Perheentupa, J.: Autoimmune polyendocrinopathy–candidosis–ectodermal

[153604]dystrophy (APECED).In: Eriksson, A. W.; Forsius, H. R.; Nevanlinna,

[153605]H. R.; Workman, P. L.; Norio, R. K.: Population Structure and Genetic

[153606]Disorders. New York: Academic Press (pub.) 1980. Pp. 583–587.

[153607]

[153608]

[153609]14775.

[153610]Prader, A.: Personal Communication. Zurich, Switzerland

1972.

[153611]

[153612]

[153613]14776.

[153614]Ramsay, C.; Winqvist, O.; Puhakka, L.; Halonen, M.; Moro, A.;

[153615]Kampe, O.; Eskelin, P.; Peltö-Huikko, M.; Peltonen, L.: Aire deficient

[153616]mice develop multiple features of APECED phenotype and show altered

[153617]immune response. Hum. Molec. Genet. 11: 397–409, 2002.

[153618]

[153619]

[153620]14777.

[153621]Rinderle, C.; Christensen, H.–M.; Schweiger, S.; Lehrach, H.;

[153622]Yaspo, M.–L.: AIRE encodes a nuclear protein co-localizing with cytoskeletal

[153623]filaments: altered sub-cellular distribution of mutants lacking PHD

[153624]zinc fingers. Hum. Molec. Genet. 8: 277–290, 1999.

[153625]

[153626]

[153627]14778.

[153628]Rosatelli, M. C.; Meloni, A.; Meloni, A.; Devoto, M.; Cao, A.;

[153629]Scott, H. S.; Peterson, P.; Heino, M.; Krohn, K. J.; Nagamine, K.;

[153630]Kudoh, J.; Shimizu, N.; Antonarakis, S. E.: A common mutation in

[153631]Sardinian autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy

[153632]patients. Hum. Genet. 103: 428–434, 1998.

[153633]

[153634]

[153635]14779.

[153636]Saenger, P.: Primary hypoaldosteronism due to zona glomerulosa

[153637]defect. (Letter) New Eng. J. Med. 310: 1394 only, 1984.

[153638]

[153639]

[153640]14780.

[153641]Saenger, P.; Levine, L. S.; Irvine, W. J.; Gottesdiener, K.;

Rauh,

[153642]W.; Sonino, N.; Chow, D.; New, M. I.: Progressive adrenal

failure

[153643]in polyglandular autoimmune disease. J. Clin. Endocr.

Metab. 54:

[153644]863–868, 1982.

[153645]

[153646]

[153647]14781.

[153648]Scott, H. S.; Heino, M.; Peterson, P.; Mittaz, L.; Lalioti, M.

[153649]D.; Betterle, C.; Cohen, A.; Seri, M.; Lerone, M.; Romeo, G.;

Collin,

[153650]P.; Salo, M.; and 9 others: Common mutations in autoim-

mune polyendocrinopathy–candidiasis–ectodermal

[153651]dystrophy patients of different origins. Molec. Endocr. 12:

1112–1119,

[153652]1998.

[153653]

[153654]

[153655]14782.

[153656]Shapiro, M. S.; Zamir, R.; Weiss, E.; Radnay, J.; Shenkman, L.

[153657]: The polyglandular deficiency syndrome: a new variant in Persian

[153658]Jews. J. Endocr. Invest. 10: 1–7, 1987.

[153659]

[153660]

[153661]14783.

[153662]Spinner, M. W.; Blizzard, R. M.; Childs, B.: Clinical and genetic

[153663]heterogeneity in idiopathic Addison's disease and hypoparathyroidism. J.

[153664]Clin. Endocr. 28: 795–804, 1968.

[153665]

[153666]

[153667]14784.

[153668]Sweetnam, W. P.: Juvenile familial endocrinopathy. Lancet
I:

[153669]463–465, 1966.

[153670]

[153671]

[153672]14785.

[153673]Wang, C.–Y.; Davoodi–Semiromi, A.; Huang, W.; Connor,
E.; Shi,

[153674]J.–D.; She, J.–X.: Characterization of mutations in patients
with

[153675]autoimmune polyglandular syndrome type 1 (APS1). Hum.
Genet. 103:

[153676]681–685, 1998.

[153677]

[153678]

[153679]14786.

[153680]Whitaker, J. A.; Landing, B. H.; Esselborn, V. M.; Williams,
R.

[153681]R.: Syndrome of familial juvenile hypoadrenocorticism, hy-
poparathyroidism

[153682]and superficial moniliasis. J. Clin. Endocr. 16: 1374–1387,
1956.

[153683]

[153684]

[153685]14787.

[153686]Wirfalt, A.: Genetic heterogeneity in autoimmune polyg-
landular

[153687]failure. Acta Med. Scand. 210: 7–13, 1981.

[153688]

[153689]

[153690]14788.

[153691]Zlotogora, J.; Shapiro, M. S.: Polyglandular autoimmune
syndrome

[153692]type I among Iranian Jews. J. Med. Genet. 29: 824–826,
1992.

[153693]

[153694]

[153695]14789.

[153696]Hediger, M. A.: New view at C. Nature Med. 8: 445–446,

2002.

[153697]

[153698]

[153699]14790.

[153700]Sotiriou, S.; Gispert, S.; Cheng, J.; Wang, Y.; Chen, A.;

Hoogstraten-Miller,

[153701]S.; Miller, G. F.; Kwon, O.; Levine, M.; Guttentag, S. H.;

Nussbaum,

[153702]R. L.: Ascorbic-acid transporter Slc23a1 is essential for vitamin
tamin

[153703]C transport into the brain and for perinatal survival. Nature Med. 8:

[153704]514-517, 2002.

[153705]

[153706]

[153707]14791.

[153708]Nauseef, W. M.; Brigham, S.; Cogley, M.: Hereditary

myeloperoxidase

[153709]deficiency due to a missense mutation of arginine 569 to
tryptophan. J.

[153710]Biol. Chem. 269: 1212-1216, 1994.

[153711]

[153712]

[153713]14792.

[153714]Busard, B. L. S. M.; Renier, W. O.; Gabreels, F. J. M.; Jaspar,

[153715]H. H. J.; van Haelst, U. J. G.; Slooff, J. L.: Lafora's disease:

[153716]comparison of inclusion bodies in skin and in brain. Arch.

Neurol. 43:

[153717]296–299, 1986.

[153718]

[153719]

[153720]14793.

[153721]Busard, H. L. S. M.; Gabreels–Festen, A. A. W. M.; Renier,

W. O.;

[153722]Gabreels, F. J. M.; Stadhouders, A. M.: Axilla skin biopsy: a

reliable

[153723]test for the diagnosis of Lafora's disease. Ann. Neurol. 21:

599–601,

[153724]1987.

[153725]

[153726]

[153727]14794.

[153728]Fluharty, A. L.; Porter, M. T.; Hirsh, G. A.; Pevida, E.; Kihara,

[153729]H.: Metachromasia in fibroblasts from a patient with Lafora's disease.

[153730](Letter) Lancet II: 109–110, 1970.

[153731]

[153732]

[153733]14795.

[153734]Ganesh, S.; Agarwala, K. L.; Ueda, K.; Akagi, T.; Shoda, K.; Usui,

[153735]T.; Hashikawa, T.; Osada, H.; Delgado-Escueta, A. V.; Yamakawa, K.

[153736]: Laforin, defective in the progressive myoclonus epilepsy of Lafora

[153737]type, is a dual-specificity phosphatase associated with polyribosomes. Hum.

[153738]Molec. Genet. 9: 2251–2261, 2000.

[153739]

[153740]

[153741]14796.

[153742]Gomez–Garre, P.; Sanz, Y.; Rodriguez de Cordoba, S.; Ser–
ratos,

[153743]J. M.: Mutational spectrum of the EPM2A gene in progres–
sive myoclonus

[153744]epilepsy of Lafora: high degree of allelic heterogeneity
and prevalence

[153745]of deletions. Europ. J. Hum. Genet. 8: 946–954, 2000.

[153746]

[153747]

[153748]14797.

[153749]Harriman, D. G. F.; Millar, J. H. D.: Progressive familial my–
oclonic

[153750]epilepsy in 3 families: its clinical features and pathological
basis. Brain 78:

[153751]325–349, 1955.

[153752]

[153753]

[153754]14798.

- [153755]Janeway, R.; Ravens, J. R.; Pearce, L. A.; Odor, D. L.; Suzuki,
[153756]K.: Progressive myoclonus epilepsy with Lafora inclusion
bodies.
[153757]I. Clinical, genetic, histopathologic and biochemical as-
pects. Arch.
[153758]Neurol. 16: 565–582, 1967.
- [153759]
- [153760]
- [153761]14799.
- [153762]Lehesjoki, A.–E.; Koskiniemi, M.; Pandolfo, M.; Antonelli,
A.;
- [153763]Kyllerman, M.; Wahlstrom, J.; Nergardh, A.; Burmeister, M.;
Sistonen,
[153764]P.; Norio, R.; de la Chapelle, A.: Linkage studies in pro-
gressive
[153765]myoclonus epilepsy: Unverricht–Lundborg and Lafora's
diseases. Neurology 42:
[153766]1545–1550, 1992.
- [153767]
- [153768]

[153769]14800.

[153770]Maddox, L. O.; Descartes, M.; Collins, J.; Keating, J.;
Rosenfeld,

[153771]S.; Palmer, C.; Carroll, A. J.; Kuzniecky, R.: Identification of
[153772]a recombination event narrowing the Lafora disease gene
region. J.

[153773]Med. Genet. 34: 590–591, 1997.

[153774]

[153775]

[153776]14801.

[153777]Amadou, C.; Ribouchon, M. T.; Mattei, M. G.; Jenkins, N.
A.; Gilbert,

[153778]D. J.; Copeland, N. G.; Avoustin, P.; Pontarotti, P.: Local-
ization

[153779]of new genes and markers to the distal part of the human
major histocompatibility

[153780]complex (MHC) region and comparison with the mouse:
new insights into

[153781]the evolution of mammalian genomes. Genomics 26:
9–20, 1995.

[153782]

[153783]

[153784]14802.

[153785]Solinas-Toldo, S.; Lengauer, C.; Fries, R.: Comparative
genome

[153786]map of human and cattle. Genomics 27: 489–496, 1995.

[153787]

[153788]

[153789]14803.

[153790]Olives, B.; Martial, S.; Mattei, M.–G.; Matassi, G.; Rousse-
let,

[153791]G.; Ripoche, P.; Cartron, J.–P.; Bailly, P.: Molecular charac-
terization

[153792]of a new urea transporter in the human kidney. FEBS Lett.
386: 156–160,

[153793]1996.

[153794]

[153795]

[153796]14804.

[153797]Ranade, K.; Wu, K.–W.; Hwu, C.–M.; Ting, C.–T.; Pei, D.;

Pesich,

[153798]R.; Hebert, J.; Chen, Y.-D. I.; Pratt, R.; Olshen, R.; Masaki, K.;

[153799]Risch, N.; Cox, D. R.; Botstein, D.: Genetic variation in the human

[153800]urea transporter-2 is associated with variation in blood pressure. Hum.

[153801]Molec. Genet. 10: 2157-2164, 2001.

[153802]

[153803]

[153804]14805.

[153805]Ansel, K. M.; Ngo, V. N.; Hyman, P. L.; Luther, S. A.; Forster,

[153806]R.; Sedgwick, J. D.; Browning, J. L.; Lipp, M.; Cyster, J. G.: A

[153807]chemokine-driven positive feedback loop organizes lymphoid follicles. Nature 406:

[153808]309-314, 2000.

[153809]

[153810]

[153811]14806.

[153812]Galko, M. J.; Tessier-Lavigne, M.: Function of an axonal chemoattractant

[153813]modulated by metalloprotease activity. Science 289: 1365–1367, 2000.

[153814]

[153815]

[153816]14807.

[153817]Stein, E.; Tessier-Lavigne, M.: Hierarchical organization of guidance

[153818]receptors: silencing of netrin attraction by Slit through a Robo/DCC

[153819]receptor complex. Science 291: 1928–1938, 2001.

[153820]

[153821]

[153822]14808.

[153823]Stein, E.; Zou, Y.; Poo, M.; Tessier-Lavigne, M.: Binding of

[153824]DCC by netrin–1 to mediate axon guidance independent of adenosine

[153825]A2B receptor activation. Science 291: 1976–1982, 2001.

[153826]

[153827]

[153828]14809.

[153829]Lubahn, D. B.; Moyer, J. S.; Golding, T. S.; Couse, J. F.; Ko-
rach,

[153830]K. S.; Smithies, O.: Alteration of reproductive function but
not

[153831]prenatal sexual development after insertional disruption
of the mouse

[153832]estrogen receptor gene. Proc. Nat. Acad. Sci. 90:
11162–11166, 1993.

[153833]

[153834]

[153835]14810.

[153836]Hanna, I. H.; Dawling, S.; Roodi, N.; Guengerich, F. P.; Parl,

[153837]F. F.: Cytochrome P450 1B1 (CYP1B1) pharmacogenetics:
association

[153838]of polymorphisms with functional differences in estrogen
hydroxylation

[153839]activity. Cancer Res. 60: 3440–3444, 2000.

[153840]

[153841]

[153842]14811.

[153843]Moore, J. T.; McKee, D. D.; Slentz-Kesler, K.; Moore, L. B.;
Jones,

[153844]S. A.; Horne, E. L.; Su, J.-L.; Kliewer, S. A.; Lehmann, J. M.;
Willson,

[153845]T. M.: Cloning and characterization of human estrogen re-
ceptor beta

[153846]isoforms. Biochem. Biophys. Res. Commun. 247: 75–78,
1998.

[153847]

[153848]

[153849]14812.

[153850]Mosselman, S.; Polman, J.; Dijkema, R.: ER-beta: identifi-
cation

[153851]and characterization of a novel human estrogen receptor.
FEBS Lett. 392:

[153852]49–53, 1996.

[153853]

[153854]

[153855]14813.

[153856]Ogawa, S.; Emi, M.; Shiraki, M.; Hosoi, T.; Ouchi, Y.; Inoue,

[153857]S.: Association of estrogen receptor beta (ESR2) gene
polymorphism

[153858]with blood pressure. J. Hum. Genet. 45: 327–330, 2000.

[153859]

[153860]

[153861]14814.

[153862]Ogawa, S.; Inoue, S.; Watanabe, T.; Orimo, A.; Hosoi, T.;
Ouchi,

[153863]Y.; Muramatsu, M.: Molecular cloning and characterization
of human

[153864]estrogen receptor beta-cx: a potential inhibitor of estro-
gen action

[153865]in human. Nucleic Acids Res. 26: 3505–3512, 1998.

[153866]

[153867]

[153868]14815.

[153869]Osterlund, M. K.; Gustafsson, J. A.; Keller, E.; Hurd, Y. L.:

[153870]Estrogen receptor beta (ER-beta) messenger ribonucleic acid (mRNA)

[153871]expression within the human forebrain: distinct distribution pattern

[153872]to ER-alpha mRNA. J. Clin. Endocr. Metab. 85: 3840-3846, 2000.

[153873]

[153874]

[153875]14816.

[153876]Pasquali, D.; Rossi, V.; Esposito, D.; Abbondanza, C.; Puca, G.

[153877]A.; Bellastella, A.; Sinisi, A. A.: Loss of estrogen receptor beta

[153878]expression in malignant human prostate cells in primary cultures and

[153879]in prostate cancer tissues. J. Clin. Endocr. Metab. 86: 2051-2055,

[153880]2001.

[153881]

[153882]

[153883]14817.

[153884]Petersen, D. N.; Tkalcevic, G. T.; Koza-Taylor, P. H.; Turi, T.

[153885]G.; Brown, T. A.: Identification of estrogen receptor beta-2, a functional

[153886]variant of estrogen receptor beta expressed in normal rat tissues. Endocrinology 139:

[153887]1082-1092, 1998.

[153888]

[153889]

[153890]14818.

[153891]Rissman, E. F.; Heck, A. L.; Leonard, J. E.; Shupnik, M. A.;
Gustafsson,

[153892]J.-A.: Disruption of estrogen receptor beta gene impairs
spatial

[153893]learning in female mice. Proc. Nat. Acad. Sci. 99:
3996–4001, 2002.

[153894]

[153895]

[153896]14819.

[153897]Rosenkranz, K.; Hinney, A.; Ziegler, A.; Hermann, H.;
Fichter,

[153898]M.; Mayer, H.; Siegfried, W.; Young, J. K.; Remschmidt, H.;
Hebebrand,

[153899]J.: Systematic mutation screening of the estrogen receptor
beta gene

[153900]in probands of different weight extremes: identification of
several

[153901]genetic variants. J. Clin. Endocr. Metab. 83: 4524–4527,

1998.

[153902]

[153903]

[153904]14820.

[153905]Speirs, V.; Adams, I. P.; Walton, D. S.; Atkin, S. L.: Identifi-
cation

[153906]of wild-type and exon 5 deletion variants of estrogen re-
ceptor beta

[153907]in normal human mammary gland. J. Clin. Endocr. Metab.
85: 1601-1605,

[153908]2000.

[153909]

[153910]

[153911]14821.

[153912]Tsukamoto, K.; Inoue, S.; Hosoi, T.; Orimo, H.; Emi, M.:
Isolation

[153913]and radiation hybrid mapping of dinucleotide repeat poly-
morphism at

[153914]the human estrogen receptor beta locus. J. Hum. Genet.
43: 73-74,

[153915]1998.

[153916]

[153917]

[153918]14822.

[153919]Weihua, Z.; Saji, S.; Makinen, S.; Cheng, G.; Jensen, E. V.;
Warner,

[153920]M.; Gustafsson, J.-A.: Estrogen receptor (ER)-beta, a mod-
ulator of

[153921]ER-alpha in the uterus. Proc. Nat. Acad. Sci. 97:
5936-5941, 2000.

[153922]

[153923]

[153924]14823.

[153925]Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.;
Landen,

[153926]M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Poly-
morphisms

[153927]of the androgen receptor gene and the estrogen receptor
gene are associated

[153928]with androgen levels in women. J. Clin. Endocr. Metab. 86:
2562-2568,

[153929]2001.

[153930]

[153931]

[153932]14824.

[153933]Zhu, Y.; Bian, Z.; Lu, P.; Karas, R. H.; Bao, L.; Cox, D.; Hodgins,

[153934]J.; Shaul, P. W.; Thoren, P.; Smithies, O.; Gustafsson, J.-A.; Mendelsohn,

[153935]M. E.: Abnormal vascular function and hypertension in mice deficient

[153936]in estrogen receptor beta. Science 295: 505–508, 2002.

[153937]

[153938]

[153939]14825.

[153940]DesGroseillers, L.; Lemieux, N.: Localization of a human double-stranded

[153941]RNA-binding protein gene (STAU) to band 20q13.1 by fluorescence in

[153942]situ hybridization. Genomics 36: 527–529, 1996.

[153943]

[153944]

[153945]14826.

[153946]Marion, R. M.; Fortes, P.; Beloso, A.; Dotti, C.; Ortin, J.: A

[153947]human sequence homologue of staufen is an RNA-binding
protein that

[153948]is associated with polysomes and localizes to the rough
endoplasmic

[153949]reticulum. Molec. Cell. Biol. 19: 2212–2219, 1999.

[153950]

[153951]

[153952]14827.

[153953]Wickham, L.; Duchaine, T.; Luo, M.; Nabi, I. R.; DesGro-
seillers,

[153954]L.: Mammalian staufen is a double-stranded-RNA- and
tubulin-binding

[153955]protein which localizes to the rough endoplasmic reticu-
lum. Molec.

[153956]Cell. Biol. 19: 2220–2230, 1999.

[153957]

[153958]

[153959]14828.

[153960]Lee, J. E.; Hollenberg, S. M.; Snider, L.; Turner, D. L.; Lipnick,

[153961]N.; Weintraub, H.: Conversion of *Xenopus* ectoderm into neurons by

[153962]NeuroD, a basic helix-loop-helix protein. *Science* 268: 836–844,

[153963]1995.

[153964]

[153965]

[153966]14829.

[153967]Liu, M.; Pleasure, S. J.; Collins, A. E.; Noebels, J. L.; Naya,

[153968]F. J.; Tsai, M.-J.; Lowenstein, D. H.: Loss of BETA2/NeuroD leads

[153969]to malformation of the dentate gyrus and epilepsy. *Proc. Nat. Acad.*

[153970]*Sci.* 97: 865–870, 2000.

[153971]

[153972]

[153973]14830.

[153974]Malecki, M. T.; Jhala, U. S.; Antonellis, A.; Fields, L.; Doria,

[153975]A.; Orban, T.; Saad, M.; Warram, J. H.; Montminy, M.;
Krolewski, A.

[153976]S.: Mutations in NEUROD1 are associated with the devel-
opment of type

[153977]2 diabetes mellitus. Nature Genet. 23: 323–328, 1999.

[153978]

[153979]

[153980]14831.

[153981]Naya, F. J.; Huang, H.–P.; Qiu, Y.; Mutoh, H.; DeMayo, F. J.;
Leiter,

[153982]A. B.; Tsai, M.–J.: Diabetes, defective pancreatic morpho-
genesis,

[153983]and abnormal enteroendocrine differentiation in
BETA2/neurod–deficient

[153984]mice. Genes Dev. 11: 2323–2334, 1997.

[153985]

[153986]

[153987]14832.

[153988]Naya, F. J.; Stellrecht, C. M.; Tsai, M. J.: Tissue–specific
regulation

[153989]of the insulin gene by a novel basic helix-loop-helix tran-
scription

[153990]factor. Genes Dev. 9: 1009-1019, 1995.

[153991]

[153992]

[153993]14833.

[153994]Plasilova, M.; Stoilov, I.; Sarfarazi, M.; Kadasi, L.; Ferakova,

[153995]E.; Ferak, V.: Identification of a single ancestral CYP1B1
mutation

[153996]in Slovak Gypsies (Roms) affected with primary congenital
glaucoma. J.

[153997]Med. Genet. 36: 290-294, 1999.

[153998]

[153999]

[154000]14834.

[154001]Schwartzman, M. L.; Balazy, M.; Masferrer, J.; Abraham, N.
G.;

[154002]McGiff, J. C.; Murphy, R. C.: 12(R)-hydroxyicosatetraenoic
acid:

[154003]a cytochrome P450-dependent arachidonate metabolite

that inhibits

[154004]Na⁺,K⁺-ATPase in the cornea. Proc. Nat. Acad. Sci. 84:
8125-8129,

[154005]1987.

[154006]

[154007]

[154008]14835.

[154009]Stoilov, I.; Akarsu, A. N.; Alozie, I.; Child, A.; Barsoum-
Homsy,

[154010]M.; Turacli, M. E.; Or, M.; Lewis, R. A.; Ozdemir, N.; Brice,
G.;

[154011]Aktan, S. G.; Chevrette, L.; Coca-Prados, M.; Sarfarazi, M.:
Sequence

[154012]analysis and homology modeling suggest that primary
congenital glaucoma

[154013]on 2p21 results from mutations disrupting either the
hinge region

[154014]or the conserved core structures of cytochrome P4501B1.
Am. J. Hum.

[154015]Genet. 62: 573-584, 1998.

[154016]

[154017]

[154018]14836.

[154019]Stoilov, I.; Akarsu, A. N.; Sarfarazi, M.: Identification of
three

[154020]different truncating mutations in cytochrome P4501B1
(CYP1B1) as the

[154021]principal cause of primary congenital glaucoma
(buphthalmos) in families

[154022]linked to the GLC3A locus on chromosome 2p21. Hum.
Molec. Genet. 6:

[154023]641–647, 1997.

[154024]

[154025]

[154026]14837.

[154027]Sutter, T. R.; Tang, Y. M.; Hayes, C. L.; Wo, Y.–Y. P.; Jabs,
E.

[154028]W.; Li, X.; Yin, H.; Cody, C. W.; Greenlee, W. F.: Complete
cDNA

[154029]sequence of a human dioxin–inducible mRNA identifies a
new gene subfamily

[154030]of cytochrome P450 that maps to chromosome 2. J. Biol.

Chem. 269:

[154031]13092–13099, 1994.

[154032]

[154033]

[154034]14838.

[154035]Tang, Y. M.; Wo, Y.–Y. P.; Stewart, J.; Hawkins, A. L.; Griffin,
fin,

[154036]C. A.; Sutter, T. R.; Greenlee, W. F.: Isolation and characterization
terization

[154037]of the human cytochrome P450 CYP1B1 gene. J. Biol.

Chem. 271: 28324–28330,

[154038]1996.

[154039]

[154040]

[154041]14839.

[154042]Vincent, A.; Billingsley, G.; Priston, M.; Williams–Lyn, D.;
Sutherland,

[154043]J.; Glaser, T.; Oliver, E.; Walter, M. A.; Heathcote, G.; Levin,
A.;

[154044]Heon, E.: Phenotypic heterogeneity of CYP1B1: mutations

in a patient

[154045]with Peters' anomaly. J. Med. Genet. 38: 324–326, 2001.

[154046]

[154047]

[154048]14840.

[154049]Chen, H. T.; Bhandoola, A.; Difilippantonio, M. J.; Zhu, J.;
Brown,

[154050]M. J.; Tai, X.; Rogakou, E. P.; Brotz, T. M.; Bonner, W. M.;
Ried,

[154051]T.; Nussenzweig, A.: Response to RAG–mediated V(D)J
cleavage by NBS1

[154052]and gamma–H2AX. Science 290: 1962–1964, 2000.

[154053]

[154054]

[154055]14841.

[154056]Petersen, S.; Casellas, R.; Reina–San–Martin, B.; Chen, H.
T.;

[154057]Difilippantonio, M. J.; Wilson, P. C.; Hanitsch, L.; Celeste,
A.;

[154058]Muramatsu, M.; Pilch, D. R.; Redon, C.; Ried, T.; Bonner,

W. M.; Honjo,

[154059]T.; Nussenzweig, M. C.; Nussenzweig, A.: AID is required
to initiate

[154060]Nbs1/gamma-H2AX focus formation and mutations at
sites of class switching. Nature 414:

[154061]660–665, 2001.

[154062]

[154063]

[154064]14842.

[154065]Morasso, M. I.; Yonescu, R.; Griffin, C. A.; Sargent, T. D.:
Localization

[154066]of human DLX8 to chromosome 17q21.3–q22 by fluores-
cence in situ hybridization. Mammalian

[154067]Genome 8: 302–303, 1997.

[154068]

[154069]

[154070]14843.

[154071]Quinn, L. M.; Johnson, B. V.; Nicholl, J.; Sutherland, G. R.;
Kalionis,

[154072]B.: Isolation and identification of homeobox genes from

the human

[154073]placenta including a novel member of the Distal-less family, DLX4. Gene 187:

[154074]55–61, 1997.

[154075]

[154076]

[154077]14844.

[154078]Boddy, M. N.; Howe, K.; Etkin, L. D.; Solomon, E.; Freemont, P.

[154079]S.: PIC 1, a novel ubiquitin-like protein which interacts with the

[154080]PML component of a multiprotein complex that is disrupted in acute

[154081]promyelocytic leukaemia. Oncogene 13: 971–982, 1996.

[154082]

[154083]

[154084]14845.

[154085]Desterro, J. M. P.; Rodriguez, M. S.; Hay, R. T.: SUMO-1 modification

[154086]of I-kappa-B-alpha inhibits NF-kappa-B activation. Molec.

Cell 2:

[154087]233–239, 1998.

[154088]

[154089]

[154090]14846.

[154091]Lapenta, V.; Chiurazzi, P.; van der Spek, P.; Pizzuti, A.;
Hanaoka,

[154092]F.; Brahe, C.: SMT3A, a human homologue of the *S. cere-*
visiae SMT3

[154093]gene, maps to chromosome 21qter and defines a novel
gene family. *Genomics* 40:

[154094]362–366, 1997.

[154095]

[154096]

[154097]14847.

[154098]Mao, Y.; Sun, M.; Desai, S. D.; Liu, L. F.: SUMO–1 conjuga-
tion

[154099]to topoisomerase I: a possible repair response to topoiso-
merase-mediated

[154100]DNA damage. *Proc. Nat. Acad. Sci.* 97: 4046–4051, 2000.

[154101]

[154102]

[154103]14848.

[154104]Okura, T.; Gong, L.; Kamitani, T.; Wada, T.; Okura, I.; Wei, C.-F.;

[154105]Chang, H.-M.; Yeh, E. T. H.: Protection against Fas/
APO-1- and tumor

[154106]necrosis factor-mediated cell death by a novel protein,
sentrin. J.

[154107]Immun. 157: 4277-4281, 1996.

[154108]

[154109]

[154110]14849.

[154111]Shen, Z.; Pardington-Purtymun, P. E.; Comeaux, J. C.;
Moyzis, R.

[154112]K.; Chen, D. J.: UBL1, a human ubiquitin-like protein asso-
ciating

[154113]with human RAD51/RAD52 proteins. Genomics 36:
271-279, 1996.

[154114]

[154115]

[154116]14850.

[154117]Bhattacharjee, H.; Ho, Y.-S.; Rosen, B. P.: Genomic organization

[154118]and chromosomal localization of the Asna1 gene, a mouse homologue

[154119]of a bacterial arsenic-translocating ATPase gene. Gene 272: 291-299,

[154120]2001.

[154121]

[154122]

[154123]14851.

[154124]Kurdi-Haidar, B.; Aebi, S.; Heath, D.; Enns, R. E.; Naredi, P.;

[154125]Hom, D. K.; Howell, S. B.: Isolation of the ATP-binding human homolog

[154126]of the arsA component of the bacterial arsenite transporter. Genomics 36:

[154127]486-491, 1996.

[154128]

[154129]

[154130]14852.

[154131]Kurdi-Haidar, B.; Heath, D.; Aebi, S.; Howell, S. B.: Bio-
chemical

[154132]characterization of the human arsenite-stimulated ATPase
(hASNA-I). J.

[154133]Biol. Chem. 273: 22173-22176, 1998.

[154134]

[154135]

[154136]14853.

[154137]Kurdi-Haidar, B.; Heath, D.; Lennon, G.; Howell, S. B.:
Chromosomal

[154138]localization and genomic structure of the human arsenite-
stimulated

[154139]ATPase (hASNA-I). Somat. Cell Molec. Genet. 24: 307-311,
1998.

[154140]

[154141]

[154142]14854.

[154143]Kurdi-Haidar, B.; Hom, D. K.; Flittner, D. E.; Heath, D.;

Fink,

[154144]L.; Naredi, P.; Howell, S. B.: Dual cytoplasmic and nuclear
distribution

[154145]of the novel arsenite-stimulated human ATPase
(hASNA-I). J. Cell.

[154146]Biochem. 71: 1-10, 1998.

[154147]

[154148]

[154149]14855.

[154150]Bengtsson, E.; Neame, P. J.; Heinegard, D.; Sommarin, Y.:
The

[154151]primary structure of a basic leucine-rich repeat protein,
PRELP, found

[154152]in connective tissues. J. Biol. Chem. 270: 25639-25644,
1995.

[154153]

[154154]

[154155]14856.

[154156]Grover, J.; Chen, X.-N.; Korenberg, J. R.; Recklies, A. D.;
Roughley,

[154157]P. J.: The gene organization, chromosome location, and

expression

[154158]of a 55-kDa matrix protein (PRELP) of human articular cartilage. Genomics 38:

[154159]109–117, 1996.

[154160]

[154161]

[154162]14857.

[154163]Bigg, H. F.; Shi, Y. E.; Liu, Y. E.; Steffensen, B.; Overall, C.

[154164]M.: Specific, high affinity binding of tissue inhibitor of metalloproteinases–4

[154165](TIMP–4) to the COOH–terminal hemopexin–like domain of human gelatinase

[154166]A. J. Biol. Chem. 272: 15496–15500, 1997.

[154167]

[154168]

[154169]14858.

[154170]Greene, J.; Wang, M.; Liu, Y. E.; Raymond, L. A.; Rosen, C.; Shi,

[154171]Y. E.: Molecular cloning and characterization of human tissue inhibitor

[154172]of metalloproteinase 4. J. Biol. Chem. 271: 30375–30380,
1996.

[154173]

[154174]

[154175]14859.

[154176]Leco, K. J.; Apte, S. S.; Taniguchi, G. T.; Hawkes, S. P.;
Khokha,

[154177]R.; Schultz, G. A.; Edwards, D. R.: Murine tissue inhibitor
of metalloproteinases–4

[154178](Timp–4): cDNA isolation and expression in adult mouse
tissues. FEBS

[154179]Lett. 401: 213–217, 1997.

[154180]

[154181]

[154182]14860.

[154183]Olson, T. M; Hirohata, S.; Ye, J.; Leco, K.; Seldin, M. F.;
Apte,

[154184]S. S.: Cloning of the human tissue inhibitor of metallopro-
teinase–4

[154185]gene (TIMP4) and localization of the TIMP4 and Timp4
gene to human

[154186]chromosome 3p25 and mouse chromosome 6, respectively. Genomics 51:

[154187]148–151, 1998.

[154188]

[154189]

[154190]14861.

[154191]Wang, M.; Liu, Y. E.; Greene, J.; Sheng, S.; Fuchs, A.;
Rosen,

[154192]E. M.; Shi, Y. E.: Inhibition of tumor growth and metastasis
of human

[154193]breast cancer cells transfected with tissue inhibitor of
metalloproteinase

[154194]4. Oncogene 14: 2767–2774, 1997.

[154195]

[154196]

[154197]14862.

[154198]Shridhar, R.; Shridhar, V.; Rivard, S. Siegfried, J. M.;
Pietraszkiewicz,

[154199]H.; Ensley, J.; Pauley, R.; Grignon, D.; Sakr, W.; Miller, O. J.;

[154200]Smith, D. I.: Mutations in the arginine-rich protein gene,

in lung,

[154201]breast, and prostate cancers, and in squamous cell carcinoma of the

[154202]head and neck. Cancer Res. 56: 5576–5578, 1996.

[154203]

[154204]

[154205]14863.

[154206]Shridhar, V.; Rivard, S.; Shridhar, R.; Mullins, C.; Bostick, L.;

[154207]Sakr, W.; Grignon, D.; Miller, O. J.; Smith, D. I.: A gene from human

[154208]chromosomal band 3p21.1 encodes a highly conserved arginine-rich protein

[154209]and is mutated in renal cell carcinomas. Oncogene 12: 1931–1939,

[154210]1996.

[154211]

[154212]

[154213]14864.

[154214]Shridhar, V.; Rivard, S.; Wang, X.; Shridhar, R.; Paisley, C.;

[154215]Mullins, C.; Beirnat, L.; Dugan, M.; Sarkar, F.; Miller, O. J.;
Vaitkevicius,

[154216]V. K.; Smith, D. I.: Mutations in the arginine-rich protein
gene

[154217](ARP) in pancreatic cancer. Oncogene 14: 2213–2216,
1997.

[154218]

[154219]

[154220]14865.

[154221]Hsu, L. C.; Chang, W.-C.: Sequencing and expression of
the human

[154222]ALDH8 encoding a new member of the aldehyde dehydro-
genase family. Gene 174:

[154223]319–322, 1996.

[154224]

[154225]

[154226]14866.

[154227]Hsu, L. C.; Chang, W.-C.; Lin, S. W.; Yoshida, A.: Cloning
and

[154228]characterization of genes encoding four additional human
aldehyde

[154229]dehydrogenase isozymes. Adv. Exp. Med. Biol. 372:
159–168, 1995.

[154230]

[154231]

[154232]14867.

[154233]Ye, J.; Rawson, R. B.; Komuro, R.; Chen, X.; Dave, U. P.;
Prywes,

[154234]R.; Brown, M. S.; Goldstein, J. L.: ER stress induces cleav-
age of

[154235]membrane-bound ATF6 by the same proteases that pro-
cess SREBPs. Molec.

[154236]Cell 6: 1355–1364, 2000.

[154237]

[154238]

[154239]14868.

[154240]Kim, V. N.; Kataoka, N.; Dreyfuss, G.: Role of the non-
sense-mediated

[154241]decay factor hUpf3 in the splicing-dependent exon-exon
junction complex. Science 293:

[154242]1832–1836, 2001.

[154243]

[154244]

[154245]14869.

[154246]Lykke-Andersen, J.; Shu, M.-D.; Steitz, J. A.: Human Upf proteins

[154247]target an mRNA for nonsense-mediated decay when bound downstream of

[154248]a termination codon. Cell 103: 1121-1131, 2000.

[154249]

[154250]

[154251]14870.

[154252]Scott, A. F.: Personal Communication. Baltimore, Md. 2/9/2001.

[154253]

[154254]

[154255]14871.

[154256]Serin, G.; Gersappe, A.; Black, J. D.; Aronoff, R.; Maquat, L.

[154257]E.: Identification and characterization of human orthologues to Saccharomyces

[154258]cerevisiae Upf2 protein and Upf3 protein (Caenorhabditis elegans SMG-4). Molec.

[154259]Cell. Biol. 21: 209–223, 2001.

[154260]

[154261]

[154262]14872.

[154263]de Leeuw, B.; Balemans, M.; Geurts van Kessel, A.: A novel Kruppel-associated

[154264]box containing the SSX gene (SSX3) on the human X chromosome is not

[154265]implicated in t(X;18)-positive synovial sarcomas. Cytogenet. Cell

[154266]Genet. 73: 179–183, 1996.

[154267]

[154268]

[154269]14873.

[154270]Gure, A. O.; Tureci, O.; Sahin, U.; Tsang, S.; Scanlan, M. J.;

[154271]Jager, E.; Knuth, A.; Pfreundschuh, M.; Old, L. J.; Chen, Y.-T.:

[154272]SSX: a multigene family with several members transcribed

in normal

[154273]testis and human cancer. Int. J. Cancer 72: 965–971,
1997.

[154274]

[154275]

[154276]14874.

[154277]Kawai, A.; Woodruff, J.; Healey, J. H.; Brennan, M. F.; An-
tonescu,

[154278]C. R.; Ladanyi, M.: SYT–SSX gene fusion as a determinant
of morphology

[154279]and prognosis in synovial sarcoma. New Eng. J. Med. 338:
153–160,

[154280]1998.

[154281]

[154282]

[154283]14875.

[154284]Panagopoulos, I.; Mertens, F.; Isaksson, M.; Limon, J.;
Gustafson,

[154285]P.; Skytting, B.; Akerman, M.; Sciort, R.; Dal Cin, P.; Sam-
son, I.;

[154286]Illiszko, M.; Ryoe, J.; Debiec–Rychter, M.; Szadowska, A.;

Brosjo,

[154287]O.; Larsson, O.; Rydholm, A.; Mandahl, N. :Genes Chromo-
somes Cancer 31:

[154288]362–372, 2001.

[154289]

[154290]

[154291]14876.

[154292]Skytting, B.; Nilsson, G.; Brodin, B.; Xie, Y.; Lundeberg, J.;

[154293]Uhlen, M.; Larsson, O.: A novel fusion gene, SYT–SSX4, in
synovial

[154294]sarcoma. (Letter) J. Nat. Cancer Inst. 91: 974–975, 1999.

[154295]

[154296]

[154297]14877.

[154298]Yang, K.; Lui, W.–O.; Xie, Y.; Zhang, A.; Skytting, B.; Man-
dahl,

[154299]N.; Larsson, C.; Larsson, O.: Co-existence of SYT–SSX1
and SYT–SSX2

[154300]fusions in synovial sarcomas. Oncogene 21: 4181–4190,
2002.

[154301]

[154302]

[154303]14878.

[154304]Piccini, M.; Vitelli, F.; Seri, M.; Galletta, L. J. V.; Moran,

[154305]O.; Bulfone, A.; Banfi, S.; Pober, B.; Renieri, A.: KCNE1-like
gene

[154306]is deleted in AMME contiguous gene syndrome: identifica-
tion and characterization

[154307]of the human and mouse homologs. Genomics 60:
251-257, 1999.

[154308]

[154309]

[154310]14879.

[154311]Koda, T.; Kakinuma, M.: Molecular cloning of a cDNA en-
coding a

[154312]novel small GTP-binding protein. FEBS Lett. 328: 21-24,
1993.

[154313]

[154314]

[154315]14880.

[154316]Scott, A. F.: Personal Communication. Baltimore, Md.
5/14/2001.

[154317]

[154318]

[154319]14881.

[154320]Zheng, J. Y.; Koda, T.; Arimura, Y.; Kishi, M.; Kakinuma,
M.:

[154321]Structure and expression of the mouse S10 gene. Biochim.
Biophys.

[154322]Acta 1351: 47–50, 1997.

[154323]

[154324]

[154325]14882.

[154326]Tipnis, S. R.; Hooper, N. M.; Hyde, R.; Karran, E.; Christie,
G.;

[154327]Turner, A. J.: A human homolog of angiotensin–converting
enzyme:

[154328]cloning and functional expression as a captopril–insen–
sitive carboxypeptidase. J.

[154329]Biol. Chem. 275: 33238–33243, 2000.

[154330]

[154331]

[154332]14883.

[154333]Berta, P.; Hawkins, J. R.; Sinclair, A. H.; Taylor, A.; Griffiths,

[154334]B. L.; Goodfellow, P. N.; Fellous, M.: Genetic evidence
equating

[154335]SRY and the testis-determining factor. Nature 348:
448-450, 1990.

[154336]

[154337]

[154338]14884.

[154339]Disteche, C. M.; Casanova, M.; Saal, H.; Friedman, C.;
Sybert,

[154340]V.; Graham, J.; Thuline, H.; Page, D.; Fellous, M.: Small
deletions

[154341]of the short arm of the Y chromosome in 46,XY females.
Proc. Nat.

[154342]Acad. Sci. 83: 7841-7844, 1986.

[154343]

[154344]

[154345]14885.

[154346]De Plaen, E.; De Backer, O.; Arnaud, D.; Bonjean, B.;
Chomez, P.;

[154347]Martelange, V.; Avner, P.; Baldacci, P.; Babinet, C.; Hwang,
S.-Y.;

[154348]Knowles, B.; Boon, T.: A new family of mouse genes ho-
mologous to

[154349]the human MAGE genes. Genomics 55: 176–184, 1999.

[154350]

[154351]

[154352]14886.

[154353]Andersson, M.; Page, D. C.; de la Chapelle, A.: Chromo-
some Y-specific

[154354]DNA is transferred to the short arm of X chromosome in
human XX males. Science 233:

[154355]786–788, 1986.

[154356]

[154357]

[154358]14887.

[154359]Burgoyne, P. S.; Levy, E. R.; McLaren, A.: Spermatogenic

failure

[154360]in male mice lacking H-Y antigen. Nature 320: 170–172,
1986.

[154361]

[154362]

[154363]14888.

[154364]Ferguson-Smith, M. A.: X-Y chromosomal interchange in
the aetiology

[154365]of true hermaphroditism and of XX Klinefelter's syndrome.

Lancet II:

[154366]475–476, 1966.

[154367]

[154368]

[154369]14889.

[154370]Simpson, E.; Chandler, P.; Goulmy, E.; Distèche, C. M.;
Ferguson-Smith,

[154371]M. A.; Page, D. C.: Separation of the genetic loci for the H-
Y antigen

[154372]and for testis determination on human Y chromosome.

Nature 326:

[154373]876–878, 1987.

[154374]

[154375]

[154376]14890.

[154377]Tiepolo, L.; Zuffardi, O.: Localization of factors controlling

[154378]spermatogenesis in the nonfluorescent portion of the human Y chromosome

[154379]long arm. Hum. Genet. 34: 119–124, 1976.

[154380]

[154381]

[154382]14891.

[154383]Kennedy, B. P.; Diehl, R. E.; Boie, Y.; Adam, M.; Dixon, R. A.

[154384]: Gene characterization and promoter analysis of the human 5-lipoxygenase-activating

[154385]protein (FLAP). J. Biol. Chem. 266: 8511–8516, 1991.

[154386]

[154387]

[154388]14892.

[154389]Lam, B. K.; Penrose, J. F.; Freeman, G. J.; Austen, K. F.: Ex–

pression

[154390]cloning of a cDNA for human leukotriene C4 synthase, an
integral membrane

[154391]protein conjugating reduced glutathione to leukotriene
A4. Proc.

[154392]Nat. Acad. Sci. 91: 7663–7667, 1994.

[154393]

[154394]

[154395]14893.

[154396]Mayatepek, E.; Flock, B.: Leukotriene C4–synthesis defi-
ciency:

[154397]a new inborn error of metabolism linked to a fatal devel-
opmental syndrome. Lancet 352:

[154398]1514–1517, 1998.

[154399]

[154400]

[154401]14894.

[154402]Lynch, K. R.; O'Neill, G. P.; Liu, Q.; Im, D.-S.; Sawyer, N.;
Metters,

[154403]K. M.; Coulombe, N.; Abramovitz, M.; Figueroa, D. J.;

Zeng, Z.; Connolly,

[154404]B. M.; Bai, C.; Austin, C. P.; Chateauneuf, A.; Stocco, R.;
Greig,

[154405]G. M.; Kargman, S.; Hooks, S. B.; Hosfield, E.; Williams, D.
L., Jr.;

[154406]Ford-Hutchinson, A. W.; Caskey, C. T.; Evans, J. F.: Char-
acterization

[154407]of the human cysteinyl leukotriene CysLT-1 receptor. Na-
ture 399:

[154408]789-793, 1999.

[154409]

[154410]

[154411]14895.

[154412]Mayatepek, E.; Zelezny, R.; Lehmann, W. D.; Hammond, J.
W.; Hoffmann,

[154413]G. F.: Defects in the synthesis of cysteinyl leukotrienes: a
new

[154414]group of inborn errors of metabolism. J. Inherit. Metab.
Dis. 23:

[154415]404-408, 2000.

[154416]

[154417]

[154418]14896.

[154419]Pace-Asciak, C. R.; Klein, J.; Spielberg, S. P.: Human genetic
netic

[154420]defect in leukotriene C(4) synthesis. Biochem. Biophys.

Res. Commun. 140:

[154421]857-860, 1986.

[154422]

[154423]

[154424]14897.

[154425]Pace-Asciak, C. R.; Klein, J.; Spielberg, S. P.: Metabolism of

[154426]leukotriene A4 into C4 by human platelets. Biochim. Biophys. Acta 877:

[154427]68-74, 1986.

[154428]

[154429]

[154430]14898.

[154431]Penrose, J. F.; Gagnon, L.; Goppelt-Struebe, M.; Myers, P.;
Lam,

[154432]B. K.; Jack, R. M.; Austen, K. F.; Soberman, R. J.: Purification

[154433]of human leukotriene C(4) synthase: organization, nucleotide sequence,

[154434]and chromosomal localization to 5q35. Proc. Nat. Acad. Sci. 89:

[154435]11603–11606, 1992.

[154436]

[154437]

[154438]14899.

[154439]Penrose, J. F.; Spector, J.; Baldasaro, M.; Xu, K.; Boyce, J.;

[154440]Arm, J. P.; Austen, K. F.; Lam, B. K.: Molecular cloning of the gene

[154441]for human leukotriene C(4) synthase. J. Biol. Chem. 271: 11356–11361,

[154442]1996.

[154443]

[154444]

[154445]14900.

[154446]Aalfs, C. M.; van den Berg, H.; Barth, P. G.; Hennekam, R.

C. M.

[154447]: The Hoyeraal–Hreidarsson syndrome: the fourth case of
a separate

[154448]entity with prenatal growth retardation, progressive pan-
cytopenia

[154449]and cerebellar hypoplasia. *Europ. J. Pediat.* 154: 304–308,
1995.

[154450]

[154451]

[154452]14901.

[154453]Devriendt, K.; Matthijs, G.; Legius, E.; Schollen, E.; Block-
mans,

[154454]D.; van Geet, C.; Degreef, H.; Cassiman, J.–J.; Fryns, J.–P.:
Skewed

[154455]X–chromosome inactivation in female carriers of dysker-
atosis congenita. *Am.*

[154456]*J. Hum. Genet.* 60: 581–587, 1997.

[154457]

[154458]

[154459]14902.

[154460]Hassock, S.; Vetrie, D.; Giannelli, F.: Mapping and charac–

terization

[154461]of the X-linked dyskeratosis congenita (DKC) gene. Genomics 55:

[154462]21–27, 1999.

[154463]

[154464]

[154465]14903.

[154466]Heiss, N. S.; Knight, S. W.; Vulliamy, T. J.; Klauck, S. M.; Wiemann,

[154467]S.; Mason, P. J.; Poustka, A.; Dokal, I.: X-linked dyskeratosis congenita

[154468]is caused by mutations in a highly conserved gene with putative nucleolar

[154469]functions. Nature Genet. 19: 32–38, 1998.

[154470]

[154471]

[154472]14904.

[154473]Heiss, N. S.; Megarbane, A.; Klauck, S. M.; Kreuz, F. R.; Makhoul,

[154474]E.; Majewski, F.; Poustka, A.: One novel and two recurrent

missense

[154475]DKC1 mutations in patients with dyskeratosis congenita (DKC). Genet.

[154476]Counsel. 12: 129–136, 2001.

[154477]

[154478]

[154479]14905.

[154480]Abedinia, M.; Layfield, R.; Jones, S. M.; Nixon, P. F.; Mattick,

[154481]J. S.: Nucleotide and predicted amino acid sequence of a cDNA clone

[154482]encoding part of human transketolase. Biochem. Biophys. Res. Commun. 183:

[154483]1159–1166, 1992.

[154484]

[154485]

[154486]14906.

[154487]Coy, J. F.; Dubel, S.; Kioschis, P.; Thomas, K.; Micklem, G.; Delius,

[154488]H.; Poustka, A.: Molecular cloning of tissue-specific tran-

scripts

[154489]of a transketolase-related gene: implications for the evolution of

[154490]new vertebrate genes. Genomics 32: 309–316, 1996.

[154491]

[154492]

[154493]14907.

[154494]McCool, B. A.; Plonk, S. G.; Martin, P. R.; Singleton, C. K.:

[154495]Cloning of human transketolase cDNAs and comparison of the nucleotide

[154496]sequence of the coding region in Wernicke–Korsakoff and non–Wernicke–Korsakoff

[154497]individuals. J. Biol. Chem. 268: 1397–1404, 1993.

[154498]

[154499]

[154500]14908.

[154501]van der Maarel, S. M.; Scholten, I. H. J. M.; Huber, I.; Philippe,

[154502]C.; Suijkerbuijk, R. F.; Gilgenkrantz, S.; Kere, J.; Cremers, F. P.

- [154503]M.; Ropers, H.-H.: Cloning and characterization of
DXS6673E, a candidate
[154504]gene for X-linked mental retardation in Xq13.1. Hum.
Molec. Genet. 5:
[154505]887–897, 1996.
- [154506]
- [154507]
- [154508]14909.
- [154509]Jacobi, F. K.; Broghammer, M.; Pesch, K.; Zrenner, E.;
Berger,
[154510]W.; Meindl, A.; Pusch, C. M.: Physical mapping and exclu-
sion of GPR34
[154511]as the causative gene for congenital stationary night
blindness type
[154512]1. Hum. Genet. 107: 89–91, 2000.
- [154513]
- [154514]
- [154515]14910.
- [154516]Marchese, A.; Sawzdargo, M.; Nguyen, T.; Cheng, R.;
Heng, H. H.

[154517]Q.; Nowak, T.; Im, D-S.; Lynch, K. R.; George, S. R.;
O'Dowd, B. F.

[154518]: Discovery of three novel orphan G-protein-coupled re-
ceptors. Genomics 56:

[154519]12-21, 1999.

[154520]

[154521]

[154522]14911.

[154523]Schoneberg, T.; Schulz, A.; Grosse, R.; Schade, R.; Hen-
klein, P.;

[154524]Schultz, G.; Gudermann, T.: A novel subgroup of class I G-
protein-coupled

[154525]receptors. Biochim. Biophys. Acta 1446: 57-70, 1999.

[154526]

[154527]

[154528]14912.

[154529]Sanchis, D.; Fleury, C.; Chomiki, N.; Gubern, M.; Huang,
Q.; Neverova,

[154530]M.; Gregoire, F.; Easlick, J.; Raimbault, S.; Levi-Meyrueis,
C.; Miroux,

[154531]B.; Collins, S.; Seldin, M.; Richard, D.; Warden, C.; Bouil-
laud, F.;

[154532]Ricquier, D.: BMCP1, a novel mitochondrial carrier with
high expression

[154533]in the central nervous system of humans and rodents, and
respiration

[154534]uncoupling activity in recombinant yeast. J. Biol. Chem.
273: 34611–34615,

[154535]1998.

[154536]

[154537]

[154538]14913.

[154539]Blanco, P.; Sargent, C. A.; Boucher, C. A.; Mitchell, M.; Af-
fara,

[154540]N. A.: Conservation of PCDHX in mammals; expression of
X/Y genes

[154541]predominantly in brain. Mammalian Genome 11: 906–914,
2000.

[154542]

[154543]

[154544]14914.

[154545]Ciccodicola, A.; D'Esposito, M.; Esposito, T.; Gianfrancesco, F.;

[154546]Migliaccio, C.; Miano, M. G.; Matarazzo, M. R.; Vacca, M.; Franze,

[154547]A.; Cuccurese, M.; Cocchia, M.; Curci, A.; and 9 others: Differentially

[154548]regulated and evolved genes in the fully sequenced Xq/Yq pseudoautosomal

[154549]region. Hum. Molec. Genet. 9: 395–401, 2000.

[154550]

[154551]

[154552]14915.

[154553]Mumm, S.; Molini, B.; Terrell, J.; Srivastava, A.; Schlessinger,

[154554]D.: Evolutionary features of the 4-Mb Xq21.3 XY homology region revealed

[154555]by a map at 60-kb resolution. Genome Res. 7: 307–314, 1997.

[154556]

[154557]

[154558]14916.

[154559]Schwartz, A.; Chan, D. C.; Brown, L. G.; Alagappan, R.;
Pettay,

[154560]D.; Disteché, C.; McGillivray, B.; de la Chapelle, A.; Page,
D. C.

[154561]: Reconstructing hominid Y evolution: X-homologous
block, created

[154562]by X-Y transposition, was disrupted by Yp inversion
through LINE-LINE

[154563]recombination. Hum. Molec. Genet. 7: 1-11, 1998.

[154564]

[154565]

[154566]14917.

[154567]Tilford, C. A.; Kuroda-Kawaguchi, T.; Skaletsky, H.; Rozen,
S.;

[154568]Brown, L. G.; Rosenberg, M.; McPherson, J. D.; Wylie, K.;
Sekhon,

[154569]M.; Kucaba, T. A.; Waterston, R. H.; Page, D. C.: A physical
map

[154570]of the human Y chromosome. Nature 409: 943-945,
2001.

[154571]

[154572]

[154573]14918.

[154574]Batch, J. A.; Evans, B. A. J.; Hughes, I. A.; Patterson, M. N.

[154575]: Mutations of the androgen receptor gene identified in
perineal hypospadias. J.

[154576]Med. Genet. 30: 198–201, 1993.

[154577]

[154578]

[154579]14919.

[154580]Borkhardt, A.; Repp, R.; Haas, O. A.; Leis, T.; Harbott, J.;
Kreuder,

[154581]J.; Hammermann, J.; Henn, T.; Lampert, F.: Cloning and
characterization

[154582]of AFX, the gene that fuses to MLL in acute leukemias with
a t(X;11)(q13;q23). Oncogene 14:

[154583]195–202, 1997.

[154584]

[154585]

[154586]14920.

[154587]Chatterjee, A.; Faust, C. J.; Herman, G. E.: Genetic and physical

[154588]mapping of the biglycan gene on the mouse X chromosome. Mammalian

[154589]Genome 4: 33–36, 1993.

[154590]

[154591]

[154592]14921.

[154593]Das, S.; Metzenberg, A.; Pai, G. S.; Gitschier, J.: Mutational

[154594]analysis of the biglycan gene excludes it as a candidate for X-linked

[154595]dominant chondrodysplasia punctata, dyskeratosis congenita, and incontinentia

[154596]pigmenti. (Letter) Am. J. Hum. Genet. 54: 922–925, 1994.

[154597]

[154598]

[154599]14922.

[154600]Fisher, L. W.; Heegaard, A.–M.; Vetter, U.; Vogel, W.; Just, W.;

[154601]Termine, J. D.; Young, M. F.: Human biglycan gene: puta–

tive promoter,

[154602]intron–exon junctions, and chromosomal localization. J.

Biol. Chem. 266:

[154603]14371–14377, 1991.

[154604]

[154605]

[154606]14923.

[154607]Fisher, L. W.; Termine, J. D.; Young, M. F.: Deduced–pro–
tein sequence

[154608]of bone small proteoglycan I (biglycan) shows homology
with proteoglycan

[154609]II (decorin) and several nonconnective tissue proteins in a
variety

[154610]of species. J. Biol. Chem. 264: 4571–4576, 1989.

[154611]

[154612]

[154613]14924.

[154614]Geerkens, C.; Vetter, U.; Just, W.; Fedarko, N. S.; Fisher, L.

[154615]W.; Young, M. F.; Termine, J. D.; Gehron Robey, P.; Wohrle,
D.; Vogel,

[154616]W.: The X-chromosomal human biglycan gene BGN is
subject to X inactivation

[154617]but is transcribed like an X-Y homologous gene. Hum.
Genet. 96:

[154618]44–52, 1995.

[154619]

[154620]

[154621]14925.

[154622]McBride, O. W.; Fisher, L. W.; Young, M. F.: Localization of
PGI

[154623](biglycan, BGN) and PGII (decorin, DCN, PG-40) genes on
human chromosomes

[154624]Xq13-qter and 12q, respectively. Genomics 6: 219–225,
1990.

[154625]

[154626]

[154627]14926.

[154628]Traupe, H.; van den Ouweland, A. M. W.; van Oost, B. A.;
Vogel,

[154629]W.; Vetter, U.; Warren, S. T.; Rocchi, M.; Darlison, M. G.;
Ropers,

[154630]H.-H.: Fine mapping of the human biglycan (BGN) gene within the Xq28

[154631]region employing a hybrid cell panel. Genomics 13: 481–483, 1992.

[154632]

[154633]

[154634]14927.

[154635]Wegrowski, Y.; Pillarisetti, J.; Danielson, K. G.; Suzuki, S.;

[154636]Iozzo, R. V.: The murine biglycan: complete cDNA cloning, genomic

[154637]organization, promoter function, and expression. Genomics 30: 8–17,

[154638]1995.

[154639]

[154640]

[154641]14928.

[154642]Xu, T.; Bianco, P.; Fisher, L. W.; Longenecker, G.; Smith, E.;

[154643]Goldstein, S.; Bonadio, J.; Boskey, A.; Heegaard, A.-M.; Sommer, B.;

[154644]Satomura, K.; Dominguez, P.; Zhao, C.; Kulkarni, A. B.;

Robey, P.

[154645]G.; Young, M. F.: Targeted disruption of the biglycan gene leads

[154646]to an osteoporosis-like phenotype in mice. Nature Genet. 20: 78–82,

[154647]1998.

[154648]

[154649]

[154650]14929.

[154651]Brunialti, A. L. B.; Harding, C. O.; Wolff, J. A.; Guenet, J.–L.

[154652]: The mouse mutation sarcosinemia (sar) maps to chromosome 2 in a

[154653]region homologous to human 9q33–q34. Genomics 36: 182–184, 1996.

[154654]

[154655]

[154656]14930.

[154657]Yoshida, K.; Sugano, S.: Identification of a novel protocadherin

[154658]gene (PCDH11) on the human XY homology region in

Xq21.3. Genomics 62:

[154659]540–543, 1999.

[154660]

[154661]

[154662]14931.

[154663]de la Chapelle, A.; Sankila, E.–M.; Lindlof, M.; Aula, P.; Norio,
rio,

[154664]R.: Norrie disease caused by a gene deletion allowing carrier
detection

[154665]and prenatal diagnosis. Clin. Genet. 28: 317–320, 1985.

[154666]

[154667]

[154668]14932.

[154669]Sims, K. B.; de la Chapelle, A.; Norio, R.; Sankila, E.–M.;
Hsu,

[154670]Y.–P. P.; Rinehart, W. B.; Corey, T. J.; Ozelius, L.; Powell, J.
F.;

[154671]Bruns, G.; Gusella, J. F.; Murphy, D. L.; Breakefield, X. O.:
Monoamine

[154672]oxidase deficiency in males with an X chromosome dele–

tion. Neuron 2:

[154673]1069–1076, 1989.

[154674]

[154675]

[154676]14933.

[154677]Grino, P. B.; Griffin, J. E.; Cushard, W. G., Jr.; Wilson, J.

[154678]D.: A mutation of the androgen receptor associated with
partial androgen

[154679]resistance, familial gynecomastia, and fertility. J. Clin. En-
docr.

[154680]Metab. 66: 754–761, 1988.

[154681]

[154682]

[154683]14934.

[154684]Holterhus, P.–M.; Sinnecker, G. H. G.; Hiort, O.: Phenotypic

[154685]diversity and testosterone–induced normalization of mu-
tant L712F androgen

[154686]receptor function in a kindred with androgen insensitivity.
J. Clin.

[154687]Endocr. Metab. 85: 3245–3250, 2000.

[154688]

[154689]

[154690]14935.

[154691]Hughes, I. A.; Evans, B. A. J.: Complete androgen insensitivity

[154692]syndrome characterized by increased concentration of a normal androgen

[154693]receptor in genital skin fibroblasts. J. Clin. Endocr. Metab. 63:

[154694]309–315, 1986.

[154695]

[154696]

[154697]14936.

[154698]Imperato–McGinley, J.; Ip, N. Y.; Gautier, T.; Neuweiler, J.;

[154699]Gruenspan, H.; Liao, S.; Chang, C.; Balazs, I.: DNA linkage analysis

[154700]and studies of the androgen receptor gene in a large kindred with

[154701]complete androgen insensitivity. Am. J. Med. Genet. 36:

104–108,

[154702]1990.

[154703]

[154704]

[154705]14937.

[154706]Jukier, L.; Kaufman, M.; Pinsky, L.; Peterson, R. E.: Partial

[154707]androgen resistance associated with secondary

5- α -reductase deficiency:

[154708]identification of a novel qualitative androgen receptor defect and

[154709]clinical implications. J. Clin. Endocr. Metab. 59: 679–688, 1984.

[154710]

[154711]

[154712]14938.

[154713]Kaufman, M.; Pinsky, L.; Bowin, A.; Au, M. W. S.: Familial external

[154714]genital ambiguity due to a transformation defect of androgen-receptor

[154715]complexes that is expressed with

- 5-alpha-dihydrotestosterone and the
[154716]synthetic androgen methyltrienolone. Am. J. Med. Genet.
18: 493-507,
[154717]1984.
- [154718]
- [154719]
- [154720]14939.
- [154721]Kaufman, M.; Straisfeld, C.; Pinsky, L.: Male pseudo-
hermaphroditism
[154722]presumably due to target organ unresponsiveness to an-
drogens: deficient
[154723]5-alpha-dihydrotestosterone binding in cultured skin fi-
broblasts. J.
[154724]Clin. Invest. 58: 345-350, 1976.
- [154725]
- [154726]
- [154727]14940.
- [154728]Keenan, B. S.; Meyer, W. J., III; Hadjian, A. J.; Jones, H. W.;
- [154729]Migeon, C. J.: Syndrome of androgen insensitivity in man:
absence

[154730]of 5- α -dihydrotestosterone binding protein in skin fibroblasts. J.

[154731]Clin. Endocr. 38: 1143-1146, 1974.

[154732]

[154733]

[154734]14941.

[154735]Liao, S.; Witte, D.: Autoimmune anti-androgen-receptor antibodies

[154736]in human serum. Proc. Nat. Acad. Sci. 82: 8345-8348, 1985.

[154737]

[154738]

[154739]14942.

[154740]Lin, S.-Y.; Ohno, S.: The binding of androgen receptor to DNA

[154741]and RNA. Biochim. Biophys. Acta 654: 181-186, 1981.

[154742]

[154743]

[154744]14943.

[154745]Aradhya, S.; Bardaro, T.; Galgoczy, P.; Yamagata, T.; Esposito,

[154746]T.; Patlan, H.; Ciccodicola, A.; Munnich, A.; Kenwrick, S.; Platzer,

[154747]M.; D'Urso, M.; Nelson, D. L.: Multiple pathogenic and benign genomic

[154748]rearrangements occur at a 35 kb duplication involving the NEMO and

[154749]LAGE2 genes. Hum. Molec. Genet. 10: 2557–2567, 2001.

[154750]

[154751]

[154752]14944.

[154753]Aradhya, S.; Courtois, G.; Rajkovic, A.; Lewis, R. A.; Levy, M.;

[154754]Israel, A.; Nelson, D. L.: Atypical forms of incontinentia pigmenti

[154755]in male individuals result from mutations of a cytosine tract in exon

[154756]10 of NEMO (IKK–gamma). Am. J. Hum. Genet. 68: 765–771, 2001.

[154757]

[154758]

[154759]14945.

[154760]Aradhya, S.; Woffendin, H.; Jakins, T.; Bardaro, T.; Esposito,

[154761]T.; Smahi, A.; Shaw, C.; Levy, M.; Munnich, A.; D'Urso, M.; Lewis,

[154762]R. A.; Kenwrick, S.; Nelson, D. L.: A recurrent deletion in the ubiquitously

[154763]expressed NEMO (IKK-gamma) gene accounts for the vast majority of

[154764]incontinentia pigmenti mutations. Hum. Molec. Genet. 10: 2171-2179,

[154765]2001.

[154766]

[154767]

[154768]14946.

[154769]Doffinger, R.; Smahi, A.; Bessia, C.; Geissmann, F.; Feinberg,

[154770]J.; Durandy, A.; Bodemer, C.; Kenwrick, S.; Dupuis-Girod, S.; Blanche,

[154771]S.; Wood, P.; Rabia, S. H.; and 16 others: X-linked an-

hidrotic

[154772]ectodermal dysplasia with immunodeficiency is caused by
impaired NF-kappa-B

[154773]signaling. Nature Genet. 27: 277-285, 2001.

[154774]

[154775]

[154776]14947.

[154777]Jain, A.; Ma, C. A.; Liu, S.; Brown, M.; Cohen, J.; Strober, W.

[154778]: Specific missense mutations in NEMO result in hyper-IgM
syndrome

[154779]with hypohidrotic (sic) ectodermal dysplasia. Nature Im-
mun. 2: 223-228,

[154780]2001.

[154781]

[154782]

[154783]14948.

[154784]Jin, D. Y.; Jeang, K. T.: Isolation of full-length cDNA and
chromosomal

[154785]localization of human NF-kappaB modulator NEMO to
Xq28. J. Biomed.

[154786]Sci. 6: 115–120, 1999.

[154787]

[154788]

[154789]14949.

[154790]Kosaki, K.; Shimasaki, N.; Fukushima, H.; Hara, M.; Ogata, T.;

[154791]Matsuo, N.: Female patient showing hypohidrotic ectodermal dysplasia

[154792]and immunodeficiency (HED-ID). (Letter) Am. J. Hum. Genet. 69: 664–665,

[154793]2001.

[154794]

[154795]

[154796]14950.

[154797]Li, Q.; Van Antwerp, D.; Mercurio, F.; Lee, K.–F.; Verma, I. M.

[154798]: Severe liver degeneration in mice lacking the I-kappa-B kinase 2

[154799]gene. Science 284: 321–325, 1999.

[154800]

[154801]

[154802]14951.

[154803]Li, Y.; Kang, J.; Friedman, J.; Tarassishin, L.; Ye, J.; Kovalenko,

[154804]A.; Wallach, D.; Horwitz, M. S.: Identification of a cell protein

[154805](FIP-3) as a modulator of NF-kappa-B activity and as a target of an

[154806]adenovirus inhibitor of tumor necrosis factor alpha-induced apoptosis. Proc.

[154807]Nat. Acad. Sci. 96: 1042-1047, 1999.

[154808]

[154809]

[154810]14952.

[154811]Makris, C.; Godfrey, V. L.; Krahn-Senftleben, G.; Takahashi, T.;

[154812]Roberts, J. L.; Schwarz, T.; Feng, L.; Johnson, R. S.; Karin, M.:

[154813]Female mice heterozygous for IKK-gamma/NEMO deficiencies develop a

[154814]dermatopathy similar to the human X-linked disorder in-

continentia

[154815]pimenti. Molec. Cell 5: 969–979, 2000.

[154816]

[154817]

[154818]14953.

[154819]May, M. J.; D'Acquisto, F.; Madge, L. A.; Glockner, J.;
Poher,

[154820]J. S.; Ghosh, S.: Selective inhibition of NF–kappa–B activa-
tion by

[154821]a peptide that blocks the interaction of NEMO with the I–
kappa–B kinase

[154822]complex. Science 289: 1550–1554, 2000.

[154823]

[154824]

[154825]14954.

[154826]Roberts, J. L.; Morrow, B.; Vega–Rich, C.; Salafia, C. M.; Ni-
towsky,

[154827]H. M.: Incontinentia pigmenti in a newborn male infant
with DNA confirmation. Am.

[154828]J. Med. Genet. 75: 159–163, 1998.

[154829]

[154830]

[154831]14955.

[154832]Rothwarf, D. M.; Zandi, E.; Natoli, G.; Karin, M.: IKK-
gamma

[154833]is an essential regulatory subunit of the I-kappa-B kinase
complex. Nature 395:

[154834]297-300, 1998.

[154835]

[154836]

[154837]14956.

[154838]Rudolph, D.; Yeh, W.-C.; Wakeham, A.; Rudolph, B.; Nal-
lainathan,

[154839]D.; Potter, J.; Elia, A. J.; Mak, T. W.: Severe liver degenera-
tion

[154840]and lack of NF-kappa-B activation in NEMO/
IKK-gamma-deficient mice. Genes

[154841]Dev. 14: 854-862, 2000.

[154842]

[154843]

[154844]14957.

[154845]Schmidt-Supprian, M.; Bloch, W.; Courtois, G.; Addicks, K.;

Israel,

[154846]A.; Rajewsky, K.; Pasparakis, M.: NEMO/

IKK-gamma-deficient mice model

[154847]incontinentia pigmenti. Molec. Cell 5: 981-992, 2000.

[154848]

[154849]

[154850]14958.

[154851]The International Incontinentia Pigmenti Consortium: Ge-
nomic

[154852]rearrangement in NEMO impairs NF-kappa-B activation
and is a cause

[154853]of incontinentia pigmenti. Nature 405: 466-472, 2000.

[154854]

[154855]

[154856]14959.

[154857]The International IP Consortium: Survival of male patients

[154858]with incontinentia pigmenti carrying a lethal mutation can

be explained

[154859]by somatic mosaicism or Klinefelter syndrome. Am. J.

Hum. Genet. 69:

[154860]1210–1217, 2001.

[154861]

[154862]

[154863]14960.

[154864]Yamaoka, S.; Courtois, G.; Bessia, C.; Whiteside, S. T.;

Weil,

[154865]R.; Agou, F.; Kirk, H. E.; Kay, R. J.; Israel, A.: Complemen-

tation

[154866]cloning of NEMO, a component of the I-kappa-B kinase

complex essential

[154867]for NF-kappa-B activation. Cell 93: 1231–1240, 1998.

[154868]

[154869]

[154870]14961.

[154871]Zonana, J.; Elder, M. E.; Schneider, L. C.; Orlow, S. J.; Moss,

[154872]C.; Golabi, M.; Shapira, S. K.; Farndon, P. A.; Wara, D. W.;

Emmal,

[154873]S. A.; Ferguson, B. M.: A novel X-linked disorder of immune deficiency

[154874]and hypohidrotic ectodermal dysplasia is allelic to incontinentia

[154875]pigmenti and due to mutations in IKK-gamma (NEMO).

Am. J. Hum. Genet. 67:

[154876]1555–1562, 2000.

[154877]

[154878]

[154879]14962.

[154880]Bauer, M. F.; Gempel, K.; Reichert, A. S.; Rappold, G. A.; Lichtner,

[154881]P.; Gerbitz, K.-D.; Neupert, W.; Brunner, M.; Hofmann, S.: Genetic

[154882]and structural characterization of the human mitochondrial inner membrane

[154883]translocase. J. Molec. Biol. 289: 69–82, 1999.

[154884]

[154885]

[154886]14963.

[154887]Moller, D. E.; Xia, C. H.; Tang, W.; Zhu, A. X.; Jakubowski, M.

[154888]: Human rsk isoforms: cloning and characterization of tissue-specific

[154889]expression. Am. J. Physiol. 266: C351–C359, 1994.

[154890]

[154891]

[154892]14964.

[154893]Zhao, Y.; Bjorbaek, C.; Weremowicz, S.; Morton, C. C.; Moller,

[154894]D. E.: RSK3 encodes a novel pp90rsk isoform with a unique N-terminal

[154895]sequence: growth factor-stimulated kinase function and nuclear translocation. Molec.

[154896]Cell. Biol. 15: 4353–4363, 1995.

[154897]

[154898]

[154899]14965.

[154900]Au, H. C.; Seo, B. B.; Matsuno-Yagi, A.; Yagi, T.; Scheffler, I.

[154901]E.: The NDUFA1 gene product (MWFE protein) is essential
for activity

[154902]of complex I in mammalian mitochondria. Proc. Nat. Acad.
Sci. 96:

[154903]4354–4359, 1999.

[154904]

[154905]

[154906]14966.

[154907]Zhuchenko, O.; Wehnert, M.; Bailey, J.; Sun, Z. S.; Lee, C.
C.

[154908]: Isolation, mapping, and genomic structure of an X-
linked gene for

[154909]a subunit of human mitochondrial complex I. Genomics
37: 281–288,

[154910]1996.

[154911]

[154912]

[154913]14967.

[154914]Liu, Z.; Sun, C.; Olejniczak, E. T.; Meadows, R. P.; Betz, S.
F.;

[154915]Oost, T.; Herrmann, J.; Wu, J. C.; Fesik, S. W.: Structural
basis

[154916]for binding of Smac/DIABLO to the XIAP BIR3 domain. Na-
ture 408:

[154917]1004–1008, 2000.

[154918]

[154919]

[154920]14968.

[154921]Srinivasula, S. M.; Hegde, R.; Saleh, A.; Datta, P.; Shiozaki,
[154922]E.; Chai, J.; Lee, R.-A.; Robbins, P. D.; Fernandes-Alnemri,
T.; Shi,
[154923]Y.; Alnemri, E. S.: A conserved XIAP-interaction motif in
caspase-9
[154924]and Smac/DIABLO regulates caspase activity and apopto-
sis. Nature 410:
[154925]112-116, 2001.
[154926]
[154927]
[154928]14969.
[154929]Bomont, P.; Cavalier, L.; Blondeau, F.; Ben Hamida, C.; Be-
lal,
[154930]S.; Tazir, M.; Demir, E.; Topaloglu, H.; Korinthenberg, R.;
Tuysuz,
[154931]B.; Landrieu, P.; Hentati, F.; Koenig, M.: The gene encod-
ing gigaxonin,
[154932]a new member of the cytoskeletal BTB/kelch repeat fam-
ily, is mutated
[154933]in giant axonal neuropathy. Nature Genet. 26: 370-374,
2000.

[154934]

[154935]

[154936]14970.

[154937]Treiber-Held, S.; Budjarjo-Welim, H.; Riemann, D.; Richter, J.;

[154938]Kretzschmar, H. A.; Hanefeld, F.: Giant axonal neuropathy: a generalized

[154939]disorder of intermediate filaments with longitudinal grooves in the

[154940]hair. Neuropediatrics 25: 89–93, 1994.

[154941]

[154942]

[154943]14971.

[154944]Lieuallen, K.; Pennacchio, L. A.; Park, M.; Myers, R. M.; Lennon,

[154945]G. G.: Cystatin B-deficient mice have increased expression of apoptosis

[154946]and glial activation genes. Hum. Molec. Genet. 10: 1867–1871, 2001.

[154947]

[154948]

[154949]14972.

[154950]Mazarib, A.; Xiong, L.; Neufeld, M. Y.; Birnbaum, M.; Korczyn,

[154951]A. D.; Pandolfo, M.; Berkovic, S. F.: Unverricht–Lundborg disease

[154952]in a five-generation Arab family: instability of dodecamer repeats. *Neurology* 57:

[154953]1050–1054, 2001.

[154954]

[154955]

[154956]14973.

[154957]Pennacchio, L. A.; Bouley, D. M.; Higgins, K. M.; Scott, M. P.;

[154958]Noebels, J. L.; Myers, R. M.: Progressive ataxia, myoclonic epilepsy

[154959]and cerebellar apoptosis in cystatin B-deficient mice. *Nature Genet.* 20:

[154960]251–258, 1998.

[154961]

[154962]

[154963]14974.

[154964]Virtaneva, K.; D'Amato, E.; Miao, J.; Koskiniemi, M.; Norio,
R.;

[154965]Avanzini, G.; Franceschetti, S.; Michelucci, R.; Tassinari, C.
A.;

[154966]Omer, S.; Pennacchio, L. A.; Myers, R. M.; Dieguez-Lucena,
J. L.;

[154967]Krahe, R.; de la Chapelle, A.; Lehesjoki, A.-E.: Unstable
minisatellite

[154968]expansion causing recessively inherited myoclonus
epilepsy, EPM1. Nature

[154969]Genet. 15: 393-396, 1997.

[154970]

[154971]

[154972]14975.

[154973]Cannizzaro, L. A.; Chen, Y. Q.; Rafi, M. A.; Wenger, D. A.:
Regional

[154974]mapping of the human galactocerebrosidase gene (GALC)
to 14q31 by

[154975]in situ hybridization. Cytogenet. Cell Genet. 66: 244-245,

1994.

[154976]

[154977]

[154978]14976.

[154979]De Gasperi, R.; Sosa, M. A. G.; Sartorato, E. L.; Battistini, S.;

[154980]MacFarlane, H.; Gusella, J. F.; Krivit, W.; Kolodny, E. H.:
Molecular

[154981]heterogeneity of late-onset forms of globoid-cell
leukodystrophy. Am.

[154982]J. Hum. Genet. 59: 1233–1242, 1996.

[154983]

[154984]

[154985]14977.

[154986]Duchen, L. W.; Eicher, E. M.; Jacobs, J. M.; Scaravilli, F.;
Teixeira,

[154987]F.: Hereditary leucodystrophy in the mouse: the new mu-
tant twitcher. Brain 103:

[154988]695–710, 1980.

[154989]

[154990]

[154991]14978.

[154992]Minassian, B. A.; Lee, J. R.; Herbrick, J.-A.; Huizenga, J.;
Soder,

[154993]S.; Mungall, A. J.; Dunham, I.; Gardner, R.; Fong, C. G.;
Carpenter,

[154994]S.; Jardim, L.; Satishchandra, P.; Andermann, E.; Snead, O.
C., III;

[154995]Lopes-Cendes, I.; Tsui, L.-C.; Delgado-Escueta, A. V.;
Rouleau, G.

[154996]A.; Scherer, S. W.: Mutations in a gene encoding a novel
protein

[154997]tyrosine phosphatase cause progressive myoclonus
epilepsy. Nature

[154998]Genet. 20: 171-174, 1998.

[154999]

[155000]

[155001]14979.

[155002]Norio, R.; Koskiniemi, M.: Progressive myoclonus epilepsy:
genetic

[155003]and nosological aspects with special reference to 107

Finnish patients. Clin.

[155004]Genet. 15: 382–398, 1979.

[155005]

[155006]

[155007]14980.

[155008]Ortiz–Hidalgo, C.: The man behind Lafora's bodies. Am. J.
Surg.

[155009]Path. 10: 358–361, 1986.

[155010]

[155011]

[155012]14981.

[155013]Sainz, J.; Minassian, B. A.; Serratosa, J. M.; Gee, M. N.;
Sakamoto,

[155014]L. M.; Iranmanesh, R.; Bohlega, S.; Baumann, R. J.; Ryan,
S.; Sparkes,

[155015]R. S.; Delgado–Escueta, A. V.: Lafora progressive my-
oclonus epilepsy:

[155016]narrowing the chromosome 6q24 locus by recombinations
and homozygosities.

[155017](Letter) Am. J. Hum. Genet. 61: 1205–1209, 1997.

[155018]

[155019]

[155020]14982.

[155021]Sarlin, M. B.; Kloepper, H. W.; Mickle, W. A.; Heath, R. G.:

[155022]The detection of carriers in hereditary myoclonic epilepsy.

Acta

[155023]Genet. Med. Gemellol. 9: 466–471, 1960.

[155024]

[155025]

[155026]14983.

[155027]Schwarz, G. A.; Yanoff, M.: Lafora's disease, distinct clin-

ico-pathologic

[155028]form of Unverricht's syndrome. Arch. Neurol. 12:

172–188, 1965.

[155029]

[155030]

[155031]14984.

[155032]Serratosa, J. M.; Delgado-Escueta, A. V.; Posada, I.; Shih,

S.;

[155033]Drury, I.; Berciano, J.; Zabala, J. A.; Antunez, M. C.;
Sparkes, R.

[155034]S.: The gene for progressive myoclonus epilepsy of the
Lafora type

[155035]maps to chromosome 6q. Hum. Molec. Genet. 4:
1657–1663, 1995.

[155036]

[155037]

[155038]14985.

[155039]Serratosa, J. M.; Gomez–Garre, P.; Gallardo, M. E.; Anta, B.;

[155040]Beltran–Valero de Bernabe, D.; Lindhout, D.; Augustijn, P.
B.; Tassinari,

[155041]C. A.; Michelucci, R.; Malafosse, A.; Topcu, M.; Grid, D.;
Dravet,

[155042]C.; Berkovic, S. F.; Rodriguez de Cordoba, S.: A novel pro-
tein tyrosine

[155043]phosphatase gene is mutated in progressive myoclonus
epilepsy of the

[155044]Lafora type (EPM2). Hum. Molec. Genet. 8: 345–352,
1999.

[155045]

[155046]

[155047]14986.

[155048]Yanoff, M.; Schwarz, G. A.: Lafora's disease: a distinct genetically

[155049]determined form of Unverricht's syndrome. Genet. Hum. 14: 235-244,

[155050]1965.

[155051]

[155052]

[155053]14987.

[155054]Yokoi, S.; Austin, J.; Witmer, F.; Sakai, M.: Studies in myoclonus

[155055]epilepsy (Lafora body forms). I. Isolation and preliminary characterization

[155056]of Lafora bodies in two cases. Arch. Neurol. 19: 15-33, 1968.

[155057]

[155058]

[155059]14988.

[155060]Antonarakis, S.: Personal Communication. Geneva,

Switzerland

[155061]4/8/1997.

[155062]

[155063]

[155064]14989.

[155065]Lafreniere, R. G.; Rochefort, D. L.; Chretien, N.; Rommens, J.

[155066]M.; Cochius, J. I.; Kalviainen, R.; Nousiainen, U.; Patry, G.; Farrell,

[155067]K.; Soderfeldt, B.; Federico, A.; Hale, B. R.; Cossio, O. H.; Sorensen,

[155068]T.; Pouliot, M. A.; Kmiec, T.; Uldall, P.; Janszky, J.; Pranzatelli,

[155069]M. R.; Andermann, F.; Andermann, E.; Rouleau, G. A.: Unstable insertion

[155070]of the 5-prime flanking region of the cystatin B gene is the most

[155071]common mutation in progressive myoclonus epilepsy type 1, EPM1. Nature

[155072]Genet. 15: 298–302, 1997.

[155073]

[155074]

[155075]14990.

[155076]Lalioi, M. D.; Mirotsou, M.; Buresi, C.; Peitsch, M. C.;
Rossier,

[155077]C.; Ouazzani, R.; Baldy-Moulinier, M.; Bottani, A.;
Malafoffe, A.;

[155078]Antonarakis, S. E.: Identification of mutations in cystatin B,
the

[155079]gene responsible for the Unverricht-Lundborg type of
progressive myoclonus

[155080]epilepsy (EPM1). Am. J. Hum. Genet. 60: 342-351, 1997.

[155081]

[155082]

[155083]14991.

[155084]Lalioi, M. D.; Scott, H. S.; Buresi, C.; Rossier, C.; Bottani,

[155085]A.; Morris, M. A.; Malafoffe, A.; Antonarakis, S. E.: Dode-
camer repeat

[155086]expansion in cystatin B gene in progressive myoclonus
epilepsy. Nature 386:

[155087]847-851, 1997.

[155088]

[155089]

[155090]14992.

[155091]Carmel, R.: Gastric juice in congenital pernicious anemia
contains

[155092]no immunoreactive intrinsic factor molecule: study of
three kindreds

[155093]with variable ages at presentation, including a patient first
diagnosed

[155094]in adulthood. Am. J. Hum. Genet. 35: 67–77, 1983.

[155095]

[155096]

[155097]14993.

[155098]Lyerla, T. A.; Konola, J. T.; Skiba, M. C.; Raghavan, S.:
Galactocerebrosidase

[155099]activity in somatic cell hybrids derived from twitcher
mouse/control

[155100]human fibroblasts is associated with human chromosome
17. Am. J.

[155101]Hum. Genet. 44: 198–207, 1989.

[155102]

[155103]

[155104]14994.

[155105]Sakai, N.; Inui, K.; Fujii, N.; Fukushima, H.; Nishimoto, J.;

[155106]Yanagihara, I.; Isegawa, Y.; Iwamatsu, A.; Okada, S.:

Krabbe disease:

[155107]isolation and characterization of a full-length cDNA for
human galactocerebrosidase. Biochem.

[155108]Biophys. Res. Commun. 198: 485–491, 1994.

[155109]

[155110]

[155111]14995.

[155112]Sweet, H.: Twitcher (twi) is on chromosome 12. Mouse
Newsletter 75:

[155113]30, 1986.

[155114]

[155115]

[155116]14996.

[155117]Roume, J.; Genin, E.; Cormier-Daire, V.; Ma, H. W.;

Mehaye, B.;

[155118]Attie, T.; Razavi-Encha, F.; Fallet-Bianco, C.; Buenerd, A.; Clerget-Darpoux,

[155119]F.; Munnich, A.; Le Merrer, M.: A gene for Meckel syndrome maps to

[155120]chromosome 11q13. Am. J. Hum. Genet. 63: 1095-1101, 1998.

[155121]

[155122]

[155123]14997.

[155124]Fernandes, M.; Poirier, C.; Lespinasse, F.; Carle, G. F.: The

[155125]mouse homologs of human GIF, DDB1, and CFL1 genes are located on chromosome

[155126]19. Mammalian Genome 9: 339-342, 1998.

[155127]

[155128]

[155129]14998.

[155130]Hewitt, J. E.; Gordon, M. M.; Taggart, R. T.; Mohandas, T. K.;

[155131]Alpers, D. H.: Human gastric intrinsic factor: characteriza-

tion of

[155132]cDNA and genomic clones and localization to human
chromosome 11. Genomics 10:

[155133]432–440, 1991.

[155134]

[155135]

[155136]14999.

[155137]Katz, M.; Lee, S. K.; Cooper, B. A.: Vitamin B(12) malab-
sorption

[155138]due to biologically inert intrinsic factor. New Eng. J. Med.
287:

[155139]425–429, 1972.

[155140]

[155141]

[155142]15000.

[155143]Katz, M.; Mehlman, C. S.; Allen, R. H.: Isolation and char-
acterization

[155144]of an abnormal human intrinsic factor. J. Clin. Invest. 53:
1274–1283,

[155145]1974.

[155146]

[155147]

[155148]15001.

[155149]Levine, J. S.; Yang, Y.; Ducos, R.; Rosenberg, A. J.; Catron, P.

[155150]G.; Podell, E. R.; Allen, R. H.: Juvenile pernicious anemia (JPA)

[155151]due to an abnormal intrinsic factor (IF) that is markedly susceptible

[155152]to acid and proteolysis. (Abstract) Gastroenterology 80: 1210 only,

[155153]1981.

[155154]

[155155]

[155156]15002.

[155157]McNicholl, B.; Egan, B.: Congenital pernicious anemia: effects

[155158]on growth, brain, and absorption of B12. Pediatrics 42: 149-156,

[155159]1968.

[155160]

[155161]

[155162]15003.

[155163]Remacha, A. F.; Sambeat, M. A.; Barcelo, M. J.; Mones, J.;
Garcia-Die,

[155164]J.; Gimferrer, E.: Congenital intrinsic factor deficiency in a
Spanish

[155165]patient. Ann. Hemat. 64: 202-204, 1992.

[155166]

[155167]

[155168]15004.

[155169]Yang, Y. M.; Ducos, R.; Rosenberg, A. J.; Catrou, P. G.;
Levine,

[155170]J. S.; Podell, E. R.; Allen, R. H.: Cobalamin malabsorption in
three

[155171]siblings due to an abnormal intrinsic factor that is
markedly susceptible

[155172]to acid and proteolysis. J. Clin. Invest. 76: 2057-2065,
1985.

[155173]

[155174]

[155175]15005.

[155176]Carney, J. P.; Maser, R. S.; Olivares, H.; Davis, E. M.; Le
Beau,

[155177]M.; Yates, J. R., III; Hays, L.; Morgan, W. F.; Petrini, J. H. J.

[155178]: The hMre11/hRad50 protein complex and Nijmegen
breakage syndrome:

[155179]linkage of double-strand break repair to the cellular DNA
damage response. Cell 93:

[155180]477-486, 1998.

[155181]

[155182]

[155183]15006.

[155184]Mailly, F.; Palmen, J.; Muller, D. P. R.; Gibbs, T.; Lloyd, J.;

[155185]Brunzell, J.; Durrington, P.; Mitropoulos, K.; Betteridge, J.;
Watts,

[155186]G.; Lithell, H.; Angelico, F.; Humphries, S. E.; Talmud, P. J.:
Familial

[155187]lipoprotein lipase (LPL) deficiency: a catalogue of LPL gene
mutations

[155188]identified in 20 patients from the UK, Sweden, and Italy.

Hum. Mutat. 10:

[155189]465–473, 1997.

[155190]

[155191]

[155192]15007.

[155193]Normand, T.; Bergeron, J.; Fernandez–Margallo, T.;

Bharucha, A.;

[155194]Ven Murthy, M. R.; Julien, P.; Gagne, C.; Dionne, C.; De

Braekeleer,

[155195]M.; Ma, R.; Hayden, M. R.; Lupien, P. J.: Geographic distri-

bution

[155196]and genealogy of mutation 207 of the lipoprotein lipase

gene in the

[155197]French Canadian population of Quebec. Hum. Genet. 89:

671–675, 1992.

[155198]

[155199]

[155200]15008.

[155201]Monsalve, M. V.; Henderson, H.; Roederer, G.; Julien, P.;

Deeb,

[155202]S.; Kastelein, J. J. P.; Peritz, L.; Devlin, R.; Bruin, T.; Murthy,

[155203]M. R. V.; Gagne, C.; Davignon, J.; Lupien, P. J.; Brunzell, J.
D.;

[155204]Hayden, M. R.: A missense mutation at codon 188 of the
human lipoprotein

[155205]lipase gene is a frequent cause of lipoprotein lipase defi-
ciency in

[155206]persons of different ancestries. J. Clin. Invest. 86:
728-734, 1990.

[155207]

[155208]

[155209]15009.

[155210]Cobben, J. M.; van der Steege, G.; Grootscholten, P.; de
Visser,

[155211]M.; Scheffer, H.; Buys, C. H. C. M.: Deletions of the survival
motor

[155212]neuron gene in unaffected siblings of patients with spinal
muscular

[155213]atrophy. Am. J. Hum. Genet. 57: 805-808, 1995.

[155214]

[155215]

[155216]15010.

[155217]Hahnen, E.; Schonling, J.; Rudnik-Schoneborn, S.; Zerres, K.;

[155218]Wirth, B.: Hybrid survival motor neuron genes in patients with autosomal

[155219]recessive spinal muscular atrophy: new insights into molecular mechanisms

[155220]responsible for the disease. Am. J. Hum. Genet. 59: 1057–1065, 1996.

[155221]

[155222]

[155223]15011.

[155224]Lefebvre, S.; Burglen, L.; Reboullet, S.; Clermont, O.; Burlet,

[155225]P.; Viollet, L.; Benichou, B.; Cruaud, C.; Millasseau, P.; Zeviani,

[155226]M.; Le Paslier, D.; Frezal, J.; Cohen, D.; Weissenbach, J.; Munnich,

[155227]A.; Melki, J.: Identification and characterization of a spinal muscular

[155228]atrophy-determining gene. Cell 80: 155–165, 1995.

[155229]

[155230]

[155231]15012.

[155232]Matthijs, G.; Schollen, E.; Legius, E.; Devriendt, K.; Goe-
mans,

[155233]N.; Kayserili, H.; Apak, M. Y.; Cassiman, J.-J.: Unusual
molecular

[155234]findings in autosomal recessive spinal muscular atrophy.
J. Med.

[155235]Genet. 33: 469–474, 1996.

[155236]

[155237]

[155238]15013.

[155239]Handschug, K.; Sperling, S.; Yoon, S.-J. K.; Hennig, S.;
Clark,

[155240]A. J. L.; Huebner, A.: Triple A syndrome is caused by mu-
tations in

[155241]AAAS, a new WD-repeat protein gene. Hum. Molec. Genet.
10: 283–290,

[155242]2001.

[155243]

[155244]

[155245]15014.

[155246]Kaplan, J.; Gerber, S.; Larget-Piet, D.; Rozet, J.-M.; Dollfus,

[155247]H.; Dufier, J.-L.; Odent, S.; Postel-Vinay, A.; Janin, N.; Briard,

[155248]M.-L.; Frezal, J.; Munnich, A.: A gene for Stargardt's disease (fundus

[155249]flavimaculatus) maps to the short arm of chromosome 1.

Nature Genet. 5:

[155250]308-311, 1993.

[155251]

[155252]

[155253]15015.

[155254]Faiella, A.; Brunelli, S.; Granata, T.; D'Incerti, L.; Cardini,

[155255]R.; Lenti, C.; Battaglia, G.; Boncinelli, E.: A number of schizencephaly

[155256]patients including 2 brothers are heterozygous for germline mutations

[155257]in the homeobox gene EMX2. Europ. J. Hum. Genet. 5:

186–190, 1997.

[155258]

[155259]

[155260]15016.

[155261]Moyer, J. H.; Lee–Tischler, M. J.; Kwon, H.–Y.; Schrick, J. J.;

[155262]Avner, E. D.; Sweeney, W. E.; Godfrey, V. L.; Cacheiro, N. L.
A.;

[155263]Wilkinson, J. E.; Woychik, R. P.: Candidate gene associated
with

[155264]a mutation causing recessive polycystic kidney disease in
mice. Science 264:

[155265]1329–1333, 1994.

[155266]

[155267]

[155268]15017.

[155269]Bespalova, I. N.; Adkins, S.; Pranzatelli, M.; Burmeister, M.:

[155270]Novel cystatin B mutation and diagnostic PCR assay in an
Unverricht–Lundborg

[155271]progressive myoclonus epilepsy patient. Am. J. Med.
Genet. 74: 467–471,

[155272]1997.

[155273]

[155274]

[155275]15018.

[155276]Jarvinen, M.; Rinne, A.: Human spleen cysteine proteinase inhibitor:

[155277]purification, fractionation into isoelectric variants and some properties

[155278]of the variants. Biochim. Biophys. Acta 708: 210–217, 1982.

[155279]

[155280]

[155281]15019.

[155282]Jerala, R.; Trstenjak, M.; Lenarcic, B.; Turk, V.: Cloning a synthetic

[155283]gene for human stefin B and its expression in E. coli. FEBS Lett. 239:

[155284]41–44, 1988.

[155285]

[155286]

[155287]15020.

[155288]Lalioti, M. D.; Scott, H. S.; Antonarakis, S. E.: What is expanded

[155289]in progressive myoclonus epilepsy? (Letter) Nature Genet. 17: 17

[155290]only, 1997.

[155291]

[155292]

[155293]15021.

[155294]Lalioti, M. D.; Scott, H. S.; Genton, P.; Grid, D.; Ouazzani, R.;

[155295]M'Rabet, A.; Ibrahim, S.; Gouider, R.; Dravet, C.; Chkili, T.; Bottani,

[155296]A.; Buresi, C.; Malafosse, A.; Antonarakis, S. E.: A PCR amplification

[155297]method reveals instability of the dodecamer repeat in progressive

[155298]myoclonus epilepsy (EPM1) and no correlation between the size of the

[155299]repeat and age at onset. Am. J. Hum. Genet. 62: 842–847, 1998.

[155300]

[155301]

[155302]15022.

[155303]Pennacchio, L. A.; Lehesjoki, A.-E.; Stone, N. E.; Willour, V.

[155304]L.; Virtaneva, K.; Miao, J.; D'Amato, E.; Ramirez, L.; Faham, M.;

[155305]Koskiniemi, M.; Warrington, J. A.; Norio, R.; de la Chapelle, A.;

[155306]Cox, D. R.; Myers, R. M.: Mutations in the gene encoding cystatin

[155307]B in progressive myoclonus epilepsy (EPM1). Science 271: 1731-1733,

[155308]1996.

[155309]

[155310]

[155311]15023.

[155312]Pennacchio, L. A.; Myers, R. M.: Isolation and characterization

[155313]of the mouse cystatin B gene. Genome Res. 6: 1103-1109, 1996.

[155314]

[155315]

[155316]15024.

[155317]Pranzatelli, M. R.; Tate, E.; Huang, Y.; Haas, R. H.; Bodenstein,

[155318]J.; Ashwal, S.; Franz, D.: Neuropharmacology of progressive myoclonus

[155319]epilepsy: response to 5-hydroxy-L-tryptophan. *Epilepsia* 36: 783-791,

[155320]1995.

[155321]

[155322]

[155323]15025.

[155324]Turk, V.; Bode, W.: The cystatins: protein inhibitors of cysteine

[155325]proteinases. *FEBS Lett.* 285: 213-219, 1991.

[155326]

[155327]

[155328]15026.

- [155329]Chang, S. C.; Hoang, B.; Thomas, J. T.; Vukicevic, S.;
Luyten,
[155330]F. P.; Ryba, N. J. P.; Kozak, C. A.; Reddi, A. H.; Moos, M.,
Jr.:
[155331]Cartilage-derived morphogenetic proteins: new members
of the transforming
[155332]growth factor-beta superfamily predominantly expressed
in long bones
[155333]during human embryonic development. J. Biol. Chem. 269:
28227-28234,
[155334]1994.
[155335]
[155336]
[155337]15027.
[155338]Faiyaz-Ul-Haque, M.; Ahmad, W.; Wahab, A.; Haque, S.;
Azim, A.
[155339]C.; Zaidi, S. H. E.; Teebi, A. S.; Ahmad, M.; Cohn, D. H.;
Siddique,
[155340]T.; Tsui, L.-C.: Frameshift mutation in the cartilage-de-
rived morphogenetic
[155341]protein 1 (CDMP1) gene and severe acromesomelic chon-
drodysplasia resembling

[155342]Grebe-type chondrodysplasia. Am. J. Med. Genet. 111:
31-37, 2002.

[155343]

[155344]

[155345]15028.

[155346]Rousseau-Merck, M.-F.; Duro, D.; Berger, R.; Thiesen, H.
J.: Chromosomal

[155347]localization of two KOX zinc finger genes on chromosome
bands 7q21-q22. Ann.

[155348]Genet. 38: 81-84, 1995.

[155349]

[155350]

[155351]15029.

[155352]Tommerup, N.; Vissing, H.: Isolation and fine mapping of
16 novel

[155353]human zinc finger-encoding cDNAs identify putative can-
didate genes

[155354]for developmental and malignant disorders. Genomics 27:
259-264,

[155355]1995.

[155356]

[155357]

[155358]15030.

[155359]Canman, C. E.; Radany, E. H.; Parsels, L. A.; Davis, M. A.;
Lawrence,

[155360]T. S.; Maybaum, J.: Induction of resistance to fluo-
rodeoxyuridine

[155361]cytotoxicity and DNA damage in human tumor cells by
expression of

[155362]Escherichia coli deoxyuridinetriphosphatase. Cancer Res.
54: 2296-2298,

[155363]1994.

[155364]

[155365]

[155366]15031.

[155367]Canman, C. E.; Tang, H. Y.; Normolle, D. P.; Lawrence, T.
S.; Maybaum,

[155368]J.: Variations in patterns of DNA damage induced in hu-
man colorectal

[155369]tumor cells by 5-fluorodeoxyuridine: implications for
mechanisms of

[155370]resistance and cytotoxicity. Proc. Nat. Acad. Sci. 89:
10474–10478,
[155371]1992.

[155372]

[155373]

[155374]15032.

[155375]Chu, R.; Lin, Y.; Rao, M. S.; Reddy, J. K.: Cloning and identification
tification

[155376]of rat deoxyuridine triphosphatase as an inhibitor of peroxisome proliferator-activated
oxisome proliferator-activated

[155377]receptor alpha. J. Biol. Chem. 271: 27670–27676, 1996.

[155378]

[155379]

[155380]15033.

[155381]Cohen, D.; Heng, H. H. Q.; Shi, X.-M.; McIntosh, E. M.;
Tsui, L.-C.;

[155382]Pearlman, R. E.: Assignment of the human dUTPase gene
(DUT) to chromosome

[155383]15q15–q21.1 by fluorescence in situ hybridization. Genomics 40:

[155384]213–215, 1997.

[155385]

[155386]

[155387]15034.

[155388]el-Hajj, H. H.; Zhang, H.; Weiss, B.: Lethality of a dut
(deoxyuridine

[155389]triphosphatase) mutation in Escherichia coli. J. Bacteriol.
170:

[155390]1069–1075, 1988.

[155391]

[155392]

[155393]15035.

[155394]Ladner, R. D.; Caradonna, S. J.: The human dUTPase gene
encodes

[155395]both nuclear and mitochondrial isoforms: differential ex-
pression of

[155396]the isoforms and characterization of a cDNA encoding the
mitochondrial

[155397]species. J. Biol. Chem. 272: 19072–19080, 1997.

[155398]

[155399]

[155400]15036.

[155401]Ladner, R. D.; McNulty, D. E.; Carr, S. A.; Roberts, G. D.;
Caradonna,

[155402]S. J.: Characterization of distinct nuclear and mitochon-
drial forms

[155403]of human deoxyuridine triphosphate nucleotidohydrolase.
J. Biol.

[155404]Chem. 271: 7745–7751, 1996.

[155405]

[155406]

[155407]15037.

[155408]Lindahl, T.: DNA repair enzymes. Annu. Rev. Biochem. 51:
61–87,

[155409]1982.

[155410]

[155411]

[155412]15038.

[155413]McIntosh, E. M.; Ager, D. D.; Gadsden, M. H.; Haynes, R.

H.: Human

[155414]dUTP pyrophosphatase: cDNA sequence and potential biological importance

[155415]of the enzyme. Proc. Nat. Acad. Sci. 89: 8020–8024, 1992. Note:

[155416]Erratum: Proc. Nat. Acad. Sci.:90: 4328 only, 1993.

[155417]

[155418]

[155419]15039.

[155420]Boring, L.; Gosling, J.; Cleary, M.; Charo, I. F.: Decreased lesion

[155421]formation in CCR2 $-/-$ mice reveals a role for chemokines in the initiation

[155422]of atherosclerosis. Nature 394: 894–897, 1998.

[155423]

[155424]

[155425]15040.

[155426]Charo, I. F.; Myers, S. J.; Herman, A.; Franci, C.; Connolly, A.

[155427]J.; Coughlin, S. R.: Molecular cloning and functional ex–

pression

[155428]of two monocyte chemoattractant protein 1 receptors reveals alternative

[155429]splicing of the carboxyl-terminal tails. Proc. Nat. Acad. Sci. 91:

[155430]2752–2756, 1994.

[155431]

[155432]

[155433]15041.

[155434]Ullman, K. S.; Powers, M. A.; Forbes, D. J.: Nuclear export receptors:

[155435]from importin to exportin. Cell 90: 967–970, 1997.

[155436]

[155437]

[155438]15042.

[155439]Barrett, T. E.; Savva, R.; Panayotou, G.; Barlow, T.; Brown, T.;

[155440]Jiricny, J.; Pearl, L. H.: Crystal structure of a G:T/U mismatch-specific

[155441]DNA glycosylase: mismatch recognition by complemen-

tary-strand interactions. Cell 92:

[155442]117-129, 1998.

[155443]

[155444]

[155445]15043.

[155446]De Gregorio, L.; Gallinari, P.; Gariboldi, M.; Manenti, G.;
Pierotti,

[155447]M. A.; Jiricny, J.; Dragani, T. A.: Genetic mapping of
thymine DNA

[155448]glycosylase (Tdg) gene and of one pseudogene in the
mouse. Mammalian

[155449]Genome 7: 909-910, 1996.

[155450]

[155451]

[155452]15044.

[155453]Lindahl, T.: DNA repair enzymes. Ann. Rev. Biochem. 51:
61-87,

[155454]1982.

[155455]

[155456]

[155457]15045.

[155458]Neddermann, P.; Gallinari, P.; Lettieri, T.; Schmid, D.;

Truong,

[155459]O.; Hsuan, J. J.; Wiebauer, K.; Jiricny, J.: Cloning and expression

[155460]of human G/T mismatch-specific thymine-DNA glycosylase. J. Biol.

[155461]Chem. 271: 12767-12774, 1996.

[155462]

[155463]

[155464]15046.

[155465]Sard, L.; Tornielli, S.; Gallinari, P.; Minoletti, F.; Jiricny,

[155466]J.; Lettieri, T.; Pierotti, M. A.; Sozzi, G.; Radice, P.: Chromosomal

[155467]localizations and molecular analysis of TDG gene-related sequences. Genomics 44:

[155468]222-226, 1997.

[155469]

[155470]

[155471]15047.

[155472]DiMarco, S. P.; Glover, T. W.; Miller, D. E.; Reines, D.; Warren,
ren,

[155473]S. T.: Transcription elongation factor SII (TCEA) maps to
human chromosome

[155474]3p22-p21.3. Genomics 36: 185-188, 1996.

[155475]

[155476]

[155477]15048.

[155478]Park, H.; Baek, K.; Jeon, C.; Agarwal, K.; Yoo, O.: Character-
terization

[155479]of the gene encoding the human transcriptional elonga-
tion factor TFIIIS. Gene 139:

[155480]263-267, 1994.

[155481]

[155482]

[155483]15049.

[155484]Reines, D.: Nascent RNA cleavage by transcription elonga-
tion complexes.In:

[155485]Conaway, R. C.; Conaway, J. W. (eds.): Transcription:

Mechanisms

[155486]and Regulation. New York: Raven Press 1994. Pp.
263–278.

[155487]

[155488]

[155489]15050.

[155490]Thomas, M. J.; Platas, A. A.; Hawley, D. K.: Transcriptional
fidelity

[155491]and proofreading by RNA polymerase II. Cell 93: 627–637,
1998.

[155492]

[155493]

[155494]15051.

[155495]Chang, C.; da Silva, S. L.; Ideta, R.; Lee, Y.; Yeh, S.; Bur-
bach,

[155496]J. P. H.: Human and rat TR4 orphan receptors specify a
subclass of

[155497]the steroid receptor superfamily. Proc. Nat. Acad. Sci. 91:
6040–6044,

[155498]1994.

[155499]

[155500]

[155501]15052.

[155502]Chang, C.; Kokontis, J.; Acakpo-Satchivi, L.; Liao, S.;
Takeda,

[155503]H.; Chang, Y.: Molecular cloning of new human TR2 re-
ceptors: a class

[155504]of steroid receptor with multiple ligand-binding domains.
Biochem.

[155505]Biophys. Res. Commun. 165: 735-741, 1989.

[155506]

[155507]

[155508]15053.

[155509]Hirose, T.; Fujimoto, W.; Yamaai, T.; Kim, K. H.; Matsuura,
H.;

[155510]Jetten, A. M.: TAK1: Molecular cloning and characteriza-
tion of a

[155511]new member of the nuclear receptor superfamily. Molec.
Endocr. 1667-1680,

[155512]1994.

[155513]

[155514]

[155515]15054.

[155516]Yoshikawa, T.; DuPont, B. R.; Leach, R. J.; Detera-Wadleigh, S.

[155517]D.: New variants of the human and rat nuclear hormone receptor, TR4:

[155518]expression and chromosomal localization of the human gene. Genomics 35:

[155519]361–366, 1996.

[155520]

[155521]

[155522]15055.

[155523]Applequist, S. E.; Selg, M.; Raman, C.; Jack, H.–M.: Cloning and

[155524]characterization of HUPF1, a human homolog of the Saccharomyces cerevisiae

[155525]nonsense mRNA–reducing UPF1 protein. Nucleic Acids Res. 25: 814–821,

[155526]1997.

[155527]

[155528]

[155529]15056.

[155530]Medghalchi, S. M.; Frischmeyer, P. A.; Mendell, J. T.; Kelly, A.

[155531]G.; Lawler, A. M.; Dietz, H. C.: Rent1, a trans-effector of nonsense-mediated

[155532]mRNA decay, is essential for mammalian embryonic viability. Hum.

[155533]Molec. Genet. 10: 99–105, 2001.

[155534]

[155535]

[155536]15057.

[155537]Mendell, J. T.; ap Rhys, C. M. J.; Dietz, H. C.: Separable roles

[155538]for rent1/hUpf1 in altered splicing and decay of nonsense transcripts. Science 298:

[155539]419–371, 2002.

[155540]

[155541]

[155542]15058.

[155543]Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Dietz, H. C.

[155544]: Cloning and characterization of a human regulator of nonsense transcript

[155545]stability. (Abstract) Am. J. Hum. Genet. 59 (suppl.): A32 only,

[155546]1996.

[155547]

[155548]

[155549]15059.

[155550]Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Kendzior, R.

[155551]J., Jr.; Dietz, H. C.: Mammalian orthologues of a yeast regulator

[155552]of nonsense transcript stability. Proc. Nat. Acad. Sci. 93: 10928–10932,

[155553]1996.

[155554]

[155555]

[155556]15060.

[155557]Sun, X.; Perlick, H. A.; Dietz, H. C.; Maquat, L. E.: A mutated

[155558]human homologue to yeast Upf1 protein has a dominant-negative effect

[155559]on the decay of nonsense-containing mRNAs in mammalian cells. Proc.

[155560]Nat. Acad. Sci. 95: 10009–10014, 1998.

[155561]

[155562]

[155563]15061.

[155564]Eng, C.; Myers, S. M.; Kogon, M. D.; Sanicola, M.; Hession, C.;

[155565]Cate, R. L.; Mulligan, L. M.: Genomic structure and chromosomal localization

[155566]of the human GDNFR-alpha gene. Oncogene 16: 597–601, 1998.

[155567]

[155568]

[155569]15062.

[155570]GFR-alpha Nomenclature Committee: Nomenclature of GPI-linked

[155571]receptors for the GDNF ligand family. Neuron 19: 485
only, 1997.

[155572]

[155573]

[155574]15063.

[155575]Gorodinsky, A.; Zimonjic, D. B.; Popescu, N. C.; Milbrandt,
J.

[155576]: Assignment of the GDNF family receptor alpha-1
(GFRA1) to human

[155577]chromosome band 10q26 by in situ hybridization. Cyto-
genet. Cell Genet. 78:

[155578]289-290, 1997.

[155579]

[155580]

[155581]15064.

[155582]Jing, S.; Wen, D.; Yu, Y.; Holst, P. L.; Luo, Y.; Fang, M.;
Tamir,

[155583]R.; Antonio, L.; Hu, Z.; Cupples, R.; Louis, J.-C.; Hu, S.; Al-
trock,

[155584]B. W.; Fox, G. M.: GDNF-induced activation of the Ret pro-
tein tyrosine

[155585]kinase is mediated by GDNFR- α , a novel receptor for GDNF. Cell 85:

[155586]1113–1124, 1996.

[155587]

[155588]

[155589]15065.

[155590]Paratcha, G.; Ledda, F.; Baars, L.; Coulpier, M.; Besset, V.; Anders,

[155591]J.; Scott, R.; Ibanez, C. F.: Released GFR- α -1 potenti-ates downstream

[155592]signaling, neuronal survival, and differentiation via a novel mechanism

[155593]of recruitment of c-Ret to lipid rafts. Neuron 29: 171–184, 2001.

[155594]

[155595]

[155596]15066.

[155597]Puliti, A.; Cinti, R.; Seri, M.; Ceccherini, I.; Romeo, G.: As-signment

[155598]of mouse Gfra1, the homologue of a new human HSCR candidate gene,

[155599]to the telomeric region of mouse chromosome 19. Cyto-
genet. Cell Genet. 78:

[155600]291–294, 1997.

[155601]

[155602]

[155603]15067.

[155604]Shefelbine, S. E.; Khorana, S.; Schultz, P. N.; Huang, E.;
Thobe,

[155605]N.; Hu, Z. J.; Fox, G. M.; Jing, S.; Cote, G. J.; Gagel, R. F.:
Mutational

[155606]analysis of the GDNF/RET–GDNFR–alpha signaling com-
plex in a kindred

[155607]with vesicoureteral reflux. Hum. Genet. 102: 474–478,
1998.

[155608]

[155609]

[155610]15068.

[155611]Bardelli, A.; Longati, P.; Albero, D.; Goruppi, S.; Schneider,

[155612]C.; Ponzetto, C.; Comoglio, P. M.: HGF receptor associates
with the

[155613]anti-apoptotic protein BAG-1 and prevents cell death.

EMBO J. 15:

[155614]6205–6212, 1996.

[155615]

[155616]

[155617]15069.

[155618]Takayama, S.; Kochel, K.; Irie, S.; Inazawa, J.; Abe, T.; Sato,

[155619]T.; Druck, T.; Huebner, K.; Reed, J. C.: Cloning of cDNAs
encoding

[155620]the human BAG1 protein and localization of the human
BAG1 gene to

[155621]chromosome 9p12. Genomics 35: 494–498, 1996.

[155622]

[155623]

[155624]15070.

[155625]Takayama, S.; Sato, T.; Krajewski, S.; Kochel, K.; Irie, S.;
Millan,

[155626]J. A.; Reed, J. C.: Cloning and functional analysis of BAG-1:
a novel

[155627]Bcl-2-binding protein with anti-cell death activity. Cell

80: 279–284,

[155628]1995.

[155629]

[155630]

[155631]15071.

[155632]Wang, H.–G.; Takayama, S.; Rapp, U. R.; Reed, J. C.: Bcl–2
interacting

[155633]protein, BAG–1, binds to and activates the kinase Raf–1.
Proc. Nat.

[155634]Acad. Sci. 93: 7063–7068, 1996.

[155635]

[155636]

[155637]15072.

[155638]Yang, X.; Pater, A.; Tang, S.–C.: Cloning and characteriza-
tion

[155639]of the human BAG–1 gene promoter: upregulation by tu-
mor–derived p53

[155640]mutants. Oncogene 18: 4546–4553, 1999.

[155641]

[155642]

[155643]15073.

[155644]Chen, H.; Chrast, R.; Rossier, C.; Gos, A.; Antonarakis, S.
E.;

[155645]Kudoh, J.; Yamaki, A.; Shindoh, N.; Maeda, H.; Minoshima,
S.; Shimizu,

[155646]N.: Single-minded and Down syndrome? (Letter) Nature
Genet. 10:

[155647]9–10, 1995.

[155648]

[155649]

[155650]15074.

[155651]Chrast, R.; Scott, H. S.; Madani, R.; Huber, L.; Wolfer, D. P.;

[155652]Prinz, M.; Aguzzi, A.; Lipp, H.-P.; Antonarakis, S. E.: Mice
trisomic

[155653]for a bacterial artificial chromosome with the single-
minded 2 gene

[155654](Sim2) show phenotypes similar to some of those present
in the partial

[155655]trisomy 16 mouse models of Down syndrome. Hum.
Molec. Genet. 9:

[155656]1853–1864, 2000.

[155657]

[155658]

[155659]15075.

[155660]Dahmane, N.; Charron, G.; Lopes, C.; Yaspo, M.–L.; Maunoury, C.;

[155661]Decorte, L.; Sinet, P.–M.; Bloch, B.; Delabar, J.–M.: Down syndrome–critical

[155662]region contains a gene homologous to *Drosophila* sim expressed during

[155663]rat and human central nervous system development. Proc. Nat. Acad.

[155664]Sci. 92: 9191–9195, 1995.

[155665]

[155666]

[155667]15076.

[155668]Ema, M.; Ikegami, S.; Hosoya, T.; Mimura, J.; Ohtani, H.; Nakao,

[155669]K.; Inokuchi, K.; Katsuki, M.; Fujii–Kuriyama, Y.: Mild impairment

[155670]of learning and memory in mice overexpressing the
mSim2 gene located

[155671]on chromosome 16: an animal model of Down's syn-
drome. Hum. Molec.

[155672]Genet. 8: 1409–1415, 1999.

[155673]

[155674]

[155675]15077.

[155676]Moffett, P.; Dayo, M.; Reece, M.; McCormick, M. K.; Pel-
letier,

[155677]J.: Characterization of msim, a murine homologue of the
Drosophila

[155678]sim transcription factor. Genomics 35: 144–155, 1996.

[155679]

[155680]

[155681]15078.

[155682]Muenke, M.; Bone, L. J.; Mitchell, H. F.; Hart, I.; Walton, K.;

[155683]Hall-Johnson, K.; Ippel, E. F.; Dietz-Band, J.; Kvaloy, K.;

Fan, C.–M.;

[155684]Tessier-Lavigne, M.; Patterson, D.: Physical mapping of

the holoprosencephaly

[155685]critical region in 21q22.3, exclusion of SIM2 as a candidate gene

[155686]for holoprosencephaly, and mapping of SIM2 to a region of chromosome

[155687]21 important for Down syndrome. Am. J. Hum. Genet. 57: 1074–1079,

[155688]1995.

[155689]

[155690]

[155691]15079.

[155692]Yamaki, A.; Noda, S.; Kudoh, J.; Shindoh, N.; Maeda, H.; Minoshima,

[155693]S.; Kawasaki, K.; Shimizu, Y.; Shimizu, N.: The mammalian single-minded

[155694](SIM) gene: mouse cDNA structure and diencephalic expression indicate

[155695]a candidate gene for Down syndrome. Genomics 35: 136–143, 1996.

[155696]

[155697]

[155698]15080.

[155699]de Winter, J. P.; Leveille, F.; van Berkel, C. G. M.;

Rooimans,

[155700]M. A.; van der Weel, L.; Steltenpool, J.; Demuth, I.; Morgan,
N. V.;

[155701]Alon, N.; Bosnoyan-Collins, L.; Lightfoot, J.; Leegwater, P.
A.; Waisfisz,

[155702]Q.; Komatsu, K.; Arwert, F.; Pronk, J. C.; Mathew, C. G.;

Digweed,

[155703]M.; Buchwald, M.; Joenje, H.: Isolation of a cDNA repre-
senting the

[155704]Fanconi anemia complementation group E gene. Am. J.
Hum. Genet. 67:

[155705]1306-1308, 2000. Note: Erratum: Am. J. Hum. Genet. 67:
1365 only,

[155706]2000.

[155707]

[155708]

[155709]15081.

[155710]Sosa-Pineda, B.; Wigle, J. T.; Oliver, G.: Hepatocyte migra-
tion

[155711]during liver development requires Prox1. Nature Genet.
25: 254–255,
[155712]2000.

[155713]

[155714]

[155715]15082.

[155716]Wigle, J. T.; Chowdhury, K.; Gruss, P.; Oliver, G.: Prox1
function

[155717]is crucial for mouse lens–fibre elongation. Nature Genet.
21: 318–322,
[155718]1999.

[155719]

[155720]

[155721]15083.

[155722]Wigle, J. T.; Oliver, G.: Prox1 function is required for the
development

[155723]of the murine lymphatic system. Cell 98: 769–778, 1999.

[155724]

[155725]

[155726]15084.

[155727]Zinovieva, R. D.; Duncan, M. K.; Johnson, T. R.; Torres, R.;
Polymeropoulos,

[155728]M. H.; Tomarev, S. I.: Structure and chromosomal localization of

[155729]the human homeobox gene Prox 1. Genomics 35:
517–522, 1996.

[155730]

[155731]

[155732]15085.

[155733]Ladenburger, E. M.; Fackelmayer, F. O.; Hameister, H.;
Knippers,

[155734]R.: MCM4 and PRKDC, human genes encoding proteins
MCM4 and DNA–PKcs,

[155735]are close neighbours located on chromosome 8q12–q13.
Cytogenet. Cell

[155736]Genet. 77: 268–270, 1997.

[155737]

[155738]

[155739]15086.

[155740]Ma, Y.; Pannicke, U.; Schwarz, K.; Lieber, M. R.: Hairpin opening

[155741]and overhang processing by an Artemis/DNA-dependent protein kinase

[155742]complex in nonhomologous end joining and V(D)J recombination. Cell 108:

[155743]781–794, 2002.

[155744]

[155745]

[155746]15087.

[155747]Satoh, T.; Tsuruga, H.; Yabuta, N.; Ishidate, M., Jr.; Nojima,

[155748]H.: Assignment of the human CDC21 (MCM4) gene to chromosome 8q11.2. Genomics 46:

[155749]525–526, 1997.

[155750]

[155751]

[155752]15088.

[155753]Joenje, H.; Lo Ten Foe, J. R.; Oostra, A. B.; van Berkel, C. G.

[155754]M.; Rooimans, M. A.; Schroeder-Kurth, T.; Wegner, R.-D.;

Gille, J.

[155755]J. P.; Buchwald, M.; Arwert, F.: Classification of Fanconi anemia

[155756]patients by complementation analysis: evidence for a fifth genetic

[155757]subtype. Blood 86: 2156–2160, 1995.

[155758]

[155759]

[155760]15089.

[155761]Byk, T.; Dobransky, T.; Cifuentes–Diaz, C.; Sobel, A.: Identification

[155762]and molecular characterization of Unc–33–like phosphoprotein (Ulip),

[155763]a putative mammalian homolog of the axonal guidance-associated unc–33

[155764]gene product. J. Neurosci. 16: 688–701, 1996.

[155765]

[155766]

[155767]15090.

[155768]Choe, H.; Farzan, M.; Sun, Y.; Sullivan, N.; Rollins, B.;

Ponath,

[155769]P. D.; Wu, L.; Mackay, C. R.; LaRosa, G.; Newman, W.; Gerard, N.;

[155770]Gerard, C.; Sodroski, J.: The beta-chemokine receptors CCR3 and CCR5

[155771]facilitate infection by primary HIV-1 isolates. Cell 85: 1135-1148,

[155772]1996.

[155773]

[155774]

[155775]15091.

[155776]Diriong, S.; Lory, P.; Williams, M. E.; Ellis, S. B.; Harpold,

[155777]M. M.; Taviaux, S.: Chromosomal localization of the human genes for

[155778]alpha-1A, alpha-1B, and alpha-1E voltage-dependent Ca(2+) channel

[155779]subunits. Genomics 30: 605-609, 1995.

[155780]

[155781]

[155782]15092.

[155783]Ben-Porath, I.; Benvenisty, N.: Characterization of a tumor-associated

[155784]gene, a member of a novel family of genes encoding membrane glycoproteins. Gene 183:

[155785]69–75, 1996.

[155786]

[155787]

[155788]15093.

[155789]Hong, Y.; Ohishi, K.; Inoue, N.; Endo, Y.; Fujita, T.; Takeda,

[155790]J.; Kinoshita, T.: Structures and chromosomal localizations of the

[155791]glycosylphosphatidylinositol synthesis gene PIGC and its pseudogene

[155792]PIGCP1. Genomics 44: 347–349, 1997.

[155793]

[155794]

[155795]15094.

[155796]Gaetano, C; Matsuo, T.; Thiele, C. J.: Identification and characterization

[155797]of a retinoic acid-regulated human homologue of the

unc-33-like phosphoprotein

[155798]gene (hUlip) from neuroblastoma cells. J. Biol. Chem. 272:
12195-12201,

[155799]1997.

[155800]

[155801]

[155802]15095.

[155803]Matsuo, T.; Stauffer, J. K.; Walker, R. L.; Meltzer, P.; Thiele,

[155804]C. J.: Structure and promoter analysis of the human unc-
33-like phosphoprotein

[155805]gene: E-box required for maximal expression in neurob-
lastoma and myoblasts. J.

[155806]Biol. Chem. 275: 16560-16568, 2000.

[155807]

[155808]

[155809]15096.

[155810]Behm, F. G.; Smith, F. O.; Raimondi, S. C.; Pui, C.-H.;
Bernstein,

[155811]I. D.: Human homologue of the rat chondroitin sulfate
proteoglycan,

[155812]NG2, detected by monoclonal antibody 7.1, identifies
childhood acute

[155813]lymphoblastic leukemias with t(4;11)(q21;q23) or
t(11;19)(q23;p13)

[155814]and MLL gene rearrangements. Blood 87: 1134–1139,
1996.

[155815]

[155816]

[155817]15097.

[155818]Pluschke, G.; Vanek, M.; Evans, A.; Dittmar, T.; Schmid, P.;
Itin,

[155819]P.; Filardo, E. J.; Reisfeld, R. A.: Molecular cloning of a hu-
man

[155820]melanoma-associated chondroitin sulfate proteoglycan.
Proc. Nat.

[155821]Acad. Sci. 93: 9710–9715, 1996.

[155822]

[155823]

[155824]15098.

[155825]Rettig, W. J.; Dracopoli, N. C.; Goetzger, T. A.; Spengler, B.

[155826]A.; Biedler, J. L.; Oettgen, H. F.; Old, L. J.: Somatic cell genetic

[155827]analysis of human cell surface antigens: chromosomal assignments and

[155828]regulation of expression in rodent-human hybrid cells.
Proc. Nat.

[155829]Acad. Sci. 81: 6437-6441, 1984.

[155830]

[155831]

[155832]15099.

[155833]Rettig, W. J.; Real, F. X.; Spengler, B. A.; Biedler, J. L.; Old,

[155834]L. J.: Human melanoma proteoglycan: expression in hybrids controlled

[155835]by intrinsic and extrinsic signals. Science 231:
1281-1284, 1986.

[155836]

[155837]

[155838]15100.

[155839]Smith, F. O.; Rauch, C.; Williams, D. E.; March, C. J.;
Arthur,

[155840]D.; Hilden, J.; Lampkin, B. C.; Buckley, J. D.; Buckley, C. V.;
Woods,

[155841]W. G.; Dinndorf, P. A.; Sorensen, P.; Kersey, J.; Hammond,
D.; Bernstein,

[155842]I. D.: The human homologue of rat NG2, a chondroitin
sulfate proteoglycan,

[155843]is not expressed on the cell surface of normal hematopoi-
etic cells

[155844]but is expressed by acute myeloid leukemia blasts from
poor-prognosis

[155845]patients with abnormalities of chromosome band 11q23.

Blood 87:

[155846]1123-1133, 1996.

[155847]

[155848]

[155849]15101.

[155850]Gipp, J. J.; Bailey, H. H.; Mulcahy, R. T.: Cloning and se-
quencing

[155851]of the cDNA for the light subunit of human liver gamma-
glutamylcysteine

[155852]synthetase and relative mRNA levels for heavy and light
subunits in

[155853]human normal tissues. Biochem. Biophys. Res. Commun.
206: 584–589,
[155854]1995.

[155855]

[155856]

[155857]15102.

[155858]Huang; C.-S.; Anderson; M. E.; Meister, A.: Amino acid se-
quence

[155859]and function of the light subunit of rat kidney gamma-
glutamylcysteine

[155860]synthetase. J. Biol. Chem. 268: 20578–20583, 1993.

[155861]

[155862]

[155863]15103.

[155864]Sierra–Rivera, E.; Dasouki, M.; Summar, M. L.; Krishna-
mani, M.

[155865]R. S.; Meredith, M.; Rao, P. N.; Phillips, J. A., III; Freeman,
M.

[155866]L.: Assignment of the human gene (GLCLR) that encodes
the regulatory

[155867]subunit of gamma-glutamylcysteine synthetase to chromosome 1p21. Cytogenet.

[155868]Cell Genet. 72: 252–254, 1996.

[155869]

[155870]

[155871]15104.

[155872]Tsuchiya, K.; Mulcahy, R. T.; Reid, L. L.; Disteché, C. M.; Kavanagh,

[155873]T. J.: Mapping of the glutamate-cysteine ligase catalytic subunit

[155874]gene (GLCLC) to human chromosome 6p12 and mouse chromosome 9D-E and

[155875]of the regulatory subunit gene (GLCLR) to human chromosome 1p21–p22

[155876]and mouse chromosome 3H1–3. Genomics 30: 630–632, 1995.

[155877]

[155878]

[155879]15105.

[155880]Rozet, J.-M.; Gerber, S.; Perrault, I.; Calvas, P.; Souied, E.;

[155881]Chatelin, S.; Viegas-Pequignot, E.; Molina-Gomez, D.;
Munnich, A.;

[155882]Kaplan, J.: Structure and refinement of the physical map-
ping of the

[155883]gamma-glutamylcysteine ligase regulatory subunit
(GLCLR) gene to chromosome

[155884]1p22.1 within the critically deleted region of human ma-
lignant mesothelioma. Cytogenet.

[155885]Cell Genet. 82: 91-94, 1998.

[155886]

[155887]

[155888]15106.

[155889]Ellis, J. A.; Luzio, J. P.: Identification and characterization

[155890]of a novel protein (p137) which transcytoses bidirection-
ally in Caco-2

[155891]cells. J. Biol. Chem. 270: 20717-20723, 1995.

[155892]

[155893]

[155894]15107.

[155895]Gessler, M.; Klamt, B.; Tsaoussidou, S.; Ellis, J. A.; Luzio, J.

[155896]P.: The gene encoding the GPI-anchored membrane protein p137(GPI)

[155897](M11S1) maps to human chromosome 11p13 and is highly conserved in

[155898]the mouse. Genomics 32: 169-170, 1996.

[155899]

[155900]

[155901]15108.

[155902]Inoue, N.; Watanabe, R.; Takeda, J.; Kinoshita, T.: PI3-C, one

[155903]of the three human genes involved in the first step of glycosylphosphatidylinositol

[155904]biosynthesis is a homologue of *Saccharomyces cerevisiae* GPI2. Biochem.

[155905]Biophys. Res. Commun. 226: 193-199, 1996.

[155906]

[155907]

[155908]15109.

[155909]Ring, H. Z.; Vameghi-Meyers, V.; Wang, W.; Crabtree, G. R.; Francke,

[155910]U.: Five SWI/SNF-related, matrix-associated, actin-dependent regulator

[155911]of chromatin (SMARC) genes are dispersed in the human genome. Genomics 51:

[155912]140–143, 1998.

[155913]

[155914]

[155915]15110.

[155916]Wang, W.; Xue, Y.; Zhou, S.; Kuo, A.; Cairns, B. R.; Crabtree,

[155917]G. R.: Diversity and specialization of mammalian SWI/SNF complexes. Genes

[155918]Dev. 10: 2117–2130, 1996.

[155919]

[155920]

[155921]15111.

[155922]Kim, B.-T.; Kitagawa, H.; Tamura, J.; Saito, T.; Kusche-Gullberg,

[155923]M.; Lindahl, U.; Sugahara, K.: Human tumor suppressor EXT gene family

[155924]members EXTL1 and EXTL3 encode alpha-
1,4-N-acetylglucosaminyltransferases

[155925]that likely are involved in heparan sulfate/heparin biosynthesis. Proc.

[155926]Nat. Acad. Sci. 98: 7176–7181, 2001.

[155927]

[155928]

[155929]15112.

[155930]Wise, C. A.; Clines, G. A.; Massa, H.; Trask, B. J.; Lovett, M.

[155931]: Identification and localization of the gene for EXTL, a
third member

[155932]of the multiple exostoses gene family. Genome Res. 7:
10–16, 1997.

[155933]

[155934]

[155935]15113.

[155936]Moskow, J. J.; Bullrich, F.; Huebner, K.; Daar, I. O.; Buch-
berg,

[155937]A. M.: Meis1, a PBX1–related homeobox gene involved in
myeloid leukemia

[155938]in BXH–2 mice. Molec. Cell. Biol. 15: 5434–5443, 1995.

[155939]

[155940]

[155941]15114.

[155942]Steelman, S.; Moskow, J. J.; Muzynski, K.; North, C.; Druck, T.;

[155943]Montgomery, J. C.; Huebner, K.; Daar, I. O.; Buchberg, A. M.: Identification

[155944]of a conserved family of Meis1-related homeobox genes. Genome Res. 7:

[155945]142–156, 1997.

[155946]

[155947]

[155948]15115.

[155949]Thorsteinsdottir, U.; Kroon, E.; Jerome, L.; Blasi, F.; Sauvageau,

[155950]G.: Defining roles for HOX and MEIS1 genes in induction of acute

[155951]myeloid leukemia. Molec. Cell. Biol. 21: 224–234, 2001.

[155952]

[155953]

[155954]15116.

[155955]Capdevila, J.; Tsukui, T.; Esteban, C. R.; Zappavigna, V.;

Belmonte,

[155956]J. C. I.: Control of vertebrate limb outgrowth by the proximal factor

[155957]Meis2 and distal antagonism of BMPs by Gremlin. Molec. Cell 4: 839–849,

[155958]1999.

[155959]

[155960]

[155961]15117.

[155962]Nakamura, T.; Jenkins, N. A.; Copeland, N. G.: Identification

[155963]of a new family of Pbx-related homeobox genes. Oncogene 13: 2235–2242,

[155964]1996.

[155965]

[155966]

[155967]15118.

[155968]Smith, J. E.; Afonja, O.; Yee, H. T.; Inghirami, G.;

Takeshita,

[155969]K.: Chromosomal mapping to 15q14 and expression anal-

ysis of the human

[155970]MEIS2 homeobox gene. Mammalian Genome 8: 951–952,
1997.

[155971]

[155972]

[155973]15119.

[155974]Kobayashi, S.; Uemura, H.; Kohda, T.; Nagai, T.; Chinen,
Y.; Naritomi,

[155975]K.; Kinoshita, E.; Ohashi, H.; Imaizumi, K.; Tsukahara, M.;
Sugio,

[155976]Y.; Tonoki, H.; Kishino, T.; Tanaka, T.; Yamada, M.; Tsut-
sumi, O.;

[155977]Niikawa, N.; Kaneko-Ishino, T.; Ishino, F.: No evidence of
PEG1/MEST

[155978]gene mutations in Silver–Russell syndrome patients. Am. J.
Med. Genet. 104:

[155979]225–231, 2001.

[155980]

[155981]

[155982]15120.

[155983]Heighway, J.; Betticher, D. C.; Hoban, P. R.; Altermatt, H. J.;

[155984]Cowen, R.: Coamplification in tumors of KRAS2, type 2 inositol 1,4,5

[155985]triphosphate receptor gene, and a novel human gene, KRAG. Genomics 35:

[155986]207–214, 1996.

[155987]

[155988]

[155989]15121.

[155990]Scott, A. F.; Elizaga, A.; Morrell, J.; Bergen, A.; Penno, M. B.

[155991]: Characterization of a gene coamplified with Ki-ras in Y1 murine

[155992]adrenal carcinoma cells that codes for a putative membrane protein. Genomics 20:

[155993]227–230, 1994.

[155994]

[155995]

[155996]15122.

[155997]Monte, D.; Baert, J. L.; Defossez, P. A.; de Launoit, Y.; Stehelin,

[155998]D.: Molecular cloning and characterization of human ERM,

a new member

[155999]of the Ets family closely related to mouse PEA3 and ER81
transcription

[156000]factors. Oncogene 9: 1397–1406, 1994.

[156001]

[156002]

[156003]15123.

[156004]Monte, D.; Coutte, L.; Dewitte, F.; Defossez, P.–A.; Le Co-
niat,

[156005]M.; Stehelin, D.; Berger, R.; de Launoit, Y.: Genomic orga-
nization

[156006]of the human ERM (ETV5) gene, a PEA3 group member of
ETS transcription

[156007]factors. Genomics 35: 236–240, 1996.

[156008]

[156009]

[156010]15124.

[156011]Protopopova, M. V.; Vorobieva, N. V.; Protopopov, A. I.;
Gizatullin,

[156012]R. Z.; Kashuba, V. I.; Klein, G.; Zabarovsky, E. R.; Grapho–

datsky,

[156013]A. S.: Assignment of the ERM gene (ETV5) coding for the ets-related

[156014]protein to human chromosome band 3q28 by in situ hybridization. Cytogenet.

[156015]Cell Genet. 74: 220 only, 1996.

[156016]

[156017]

[156018]15125.

[156019]Bosher, J. M.; Williams, T.; Hurst, H. C.: The developmentally

[156020]regulated transcription factor AP-2 is involved in c-erbB-2 overexpression

[156021]in human mammary carcinoma. Proc. Nat. Acad. Sci. 92: 744-747, 1995.

[156022]

[156023]

[156024]15126.

[156025]Clements, J. L.; Lee, J. R.; Gross, B.; Yang, B.; Olson, J. D.;

[156026]Sandra, A.; Watson, S. P.; Lentz, S. R.; Koretzky, G. A.: Fe-

tal hemorrhage

[156027]and platelet dysfunction in SLP-76-deficient mice. J. Clin.

Invest. 103:

[156028]19-25, 1999.

[156029]

[156030]

[156031]15127.

[156032]Jackman, J. K.; Motto, D. G.; Sun, Q.; Tanemoto, M.; Turck,
C.

[156033]W.; Peltz, G. A.; Koretzky, G. A.; Findell, P. R.: Molecular
cloning

[156034]of SLP-76, a 76-kDa tyrosine phosphoprotein associated
with Grb2 in

[156035]T cells. J. Biol. Chem. 270: 7029-7032, 1995.

[156036]

[156037]

[156038]15128.

[156039]Motto, D. G.; Ross, S. E.; Wu, J.; Hendricks-Taylor, L. R.;
Koretzky,

[156040]G. A.: Implication of the GRB2-associated phosphoprotein

SLP-76 in

[156041]T cell receptor-mediated interleukin 2 production. J. Exp.
Med. 183:

[156042]1937-1943, 1996.

[156043]

[156044]

[156045]15129.

[156046]Pivniouk, V.; Tsitsikov, E.; Swinton, P.; Rathbun, G.; Alt, F.

[156047]W.; Geha, R. S.: Impaired viability and profound block in
thymocyte

[156048]development in mice lacking the adaptor protein SLP-76.

Cell 94:

[156049]229-238, 1998.

[156050]

[156051]

[156052]15130.

[156053]Combadiere, C.; Ahuja, S. K.; Murphy, P. M.: Cloning,
chromosomal

[156054]localization, and RNA expression of a human beta
chemokine receptor-like

[156055]gene. DNA Cell Biol. 14: 673–680, 1995.

[156056]

[156057]

[156058]15131.

[156059]Combadiere, C.; Ahuja, S. K.; Van Damme, J.; Tiffany, H.

L.; Gao,

[156060]J.–L.; Murphy, P. M.: Monocyte chemoattractant protein–3

is a functional

[156061]ligand for CC chemokine receptors 1 and 2B. J. Biol.

Chem. 270:

[156062]29671–29675, 1995.

[156063]

[156064]

[156065]15132.

[156066]Doranz, B. J.; Rucker, J.; Yi, Y.; Smyth, R. J.; Samson, M.;

Peiper,

[156067]S. C.; Parmentier, M.; Collman, R. G.; Doms, R. W.: A dual–

tropic

[156068]primary HIV–1 isolate that uses fusin and the beta–

chemokine receptors

[156069]CKR-5, CKR-3, and CKR-2b as fusion cofactors. Cell 85:
1149-1158,
[156070]1996.

[156071]

[156072]

[156073]15133.

[156074]Mummidi, S.; Ahuja, S. S.; Gonzalez, E.; Anderson, S. A.;
Santiago,

[156075]E. N.; Stephan, K. T.; Craig, F. E.; O'Connell, P.; Tryon, V.;
Clark,

[156076]R. A.; Dolan, M. J.; Ahuja, S. K.: Genealogy of the CCR5 lo-
cus and

[156077]chemokine system gene variants associated with altered
rates of HIV-1

[156078]disease progression. Nature Med. 4: 786-793, 1998.

[156079]

[156080]

[156081]15134.

[156082]Peters, W.; Dupuis, M.; Charo, I. F.: A mechanism for the
impaired

[156083]IFN- γ production in C-C chemokine receptor 2
(CCR2) knockout mice:

[156084]role of CCR2 in linking the innate and adaptive immune
responses. J.

[156085]Immun. 165: 7072–7077, 2000.

[156086]

[156087]

[156088]15135.

[156089]Peters, W.; Scott, H. M.; Chambers, H. F.; Flynn J. L.; Charo,

[156090]I. F.; Ernst, J. D.: Chemokine receptor 2 serves an early
and essential

[156091]role in resistance to *Mycobacterium tuberculosis*. Proc.
Nat. Acad.

[156092]Sci. 98: 7958–7963, 2001.

[156093]

[156094]

[156095]15136.

[156096]Samson, M.; Labbe, O.; Mollereau, C.; Vassart, G.; Par-
mentier,

[156097]M.: Molecular cloning and functional expression of a new

human CC-chemokine

[156098]receptor gene. Biochemistry 35: 3362–3367, 1996.

[156099]

[156100]

[156101]15137.

[156102]Samson, M.; Soularue, P.; Vassart, G.; Parmentier, M.: The
genes

[156103]encoding the human CC-chemokine receptors CC-CKR1
to CC-CKR5 (CMKBR1–CMKBR5)

[156104]are clustered in the p21.3–p24 region of chromosome 3.

Genomics 36:

[156105]522–526, 1996.

[156106]

[156107]

[156108]15138.

[156109]Sanders, S. K.; Crean, S. M.; Boxer, P. A.; Kellner, D.;
LaRosa,

[156110]G. J.; Hunt, S. W., III.: Functional differences between
monocyte

[156111]chemotactic protein–1 receptor A and monocyte chemo–

tactic protein-1

[156112]receptor B expressed in a Jurkat T cell. J. Immun. 165:
4877-4883,

[156113]2000.

[156114]

[156115]

[156116]15139.

[156117]Smith, M. W.; Dean, M.; Carrington, M.; Winkler, C.; Hutt-
ley,

[156118]G. A.; Lomb, D. A.; Goedert, J. J.; O'Brien, T. R.; Jacobson,
L. P.;

[156119]Kaslow, R.; Buchbinder, S.; Vittinghoff, E.; Vlahov, D.;
Hoots, K.;

[156120]Hilgartner, M. W.; Hemophilia Growth and Development
Study (HGDS);

[156121]Multicenter AIDS Cohort Study (MACS); Multicenter
Hemophilia Cohort

[156122]Study (MHCS); San Francisco City Cohort (SFCC); ALIVE
Study;

[156123]O'Brien, S. J.: Contrasting genetic influence of CCR2 and
CCR5 variants

[156124]on HIV-1 infection and disease progression. Science 277:

959-965,

[156125]1997.

[156126]

[156127]

[156128]15140.

[156129]Wong, L.-M.; Myers, S. J.; Tsou, C.-L.; Gosling, J.; Arai, H.;

[156130]Charo, I. F.: Organization and differential expression of
the human

[156131]monocyte chemoattractant protein 1 receptor gene: evi-
dence for the

[156132]role of the carboxyl-terminal tail in receptor trafficking. J.
Biol.

[156133]Chem. 272: 1038-1045, 1997.

[156134]

[156135]

[156136]15141.

[156137]Donald, L. J.; Wang, H. S.; Hamerton, J. L.: Assignment of
the

[156138]gene for cystathionase (CYS) to human chromosome 16.
(Abstract) Cytogenet.

[156139]Cell Genet. 32: 268 only, 1982.

[156140]

[156141]

[156142]15142.

[156143]Frimpter, G. W.: Cystathioninuria: nature of the defect.

Science 149:

[156144]1095–1096, 1965.

[156145]

[156146]

[156147]15143.

[156148]Frimpter, G. W.: Cystathioninuria, sulfite oxidase deficiency,

[156149]and 'beta-mercaptolactate-cysteine disulfiduria.In: Stanbury, J. B.;

[156150]Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic Basis of Inherited

[156151]Disease. New York: McGraw–Hill (pub.) (3rd ed.): 1972. Pp. 413–425.

[156152]

[156153]

[156154]15144.

[156155]Frimpter, G. W.; Haymovitz, A.; Horwith, M.: Cystathionin-
uria. New

[156156]Eng. J. Med. 268: 333–339, 1963.

[156157]

[156158]

[156159]15145.

[156160]Harris, H.; Penrose, L. S.; Thomas, D. H. H.: Cystathionin-
uria. Ann.

[156161]Hum. Genet. 23: 442–453, 1959.

[156162]

[156163]

[156164]15146.

[156165]Lu, Y.; O'Dowd, B. F.; Orrego, H.; Israel, Y.: Cloning and
nucleotide

[156166]sequence of human liver cDNA encoding for cystathionine
gamma-lyase. Biochem.

[156167]Biophys. Res. Commun. 189: 749–758, 1992.

[156168]

[156169]

[156170]15147.

[156171]Lyon, I. C. T.; Procopis, P. G.; Turner, B.: Cystathioninuria

[156172]in a well baby population. Acta Paediat. Scand. 60:
324–328, 1971.

[156173]

[156174]

[156175]15148.

[156176]Mongeau, J.–G.; Hilgartner, M.; Worthen, H. G.; Frimpter,
G. W.

[156177]: Cystathioninuria: study of an infant with normal mental–
ity, thrombocytopenia,

[156178]and renal calculi. J. Pediat. 69: 1113–1120, 1967.

[156179]

[156180]

[156181]15149.

[156182]Pascal, T. A.; Gaull, G. E.; Beratis, N. G.; Gillam, B. M.; Tal–
lan,

[156183]H. H.: Cystathionase deficiency: evidence for genetic het–

erogeneity

[156184]in primary cystathioninuria. *Pediat. Res.* 12: 125–133,
1978.

[156185]

[156186]

[156187]15150.

[156188]Perry, T. L.; Hardwick, D. F.; Hansen, S.; Love, D. L.; Is-
raels,

[156189]S.: Cystathioninuria in two healthy siblings. *New Eng. J.*
Med. 278:

[156190]590–592, 1968.

[156191]

[156192]

[156193]15151.

[156194]Schneiderman, L. J.: Latent cystathioninuria. *J. Med. Genet.*
4:

[156195]260–263, 1967.

[156196]

[156197]

[156198]15152.

[156199]Scott, C. R.; Dassell, S. W.; Clark, S. H.; Chiang-Teng, C.;
Swedberg,

[156200]K. R.: Cystathioninemia: a benign genetic condition. J. Pe-
diat. 76:

[156201]571-577, 1970.

[156202]

[156203]

[156204]15153.

[156205]Shaw, K. N. F.; Lieberman, E.; Koch, R.; Donnell, G. N.:
Cystathioninuria. Am.

[156206]J. Dis. Child. 113: 119-128, 1967.

[156207]

[156208]

[156209]15154.

[156210]Tada, K.; Yoshida, T.; Yokoyama, Y.; Sato, T.; Nakagawa,
H.; Arakawa,

[156211]T.: Cystathioninuria not associated with vitamin B6 de-
pendency: a

[156212]probably new type of cystathioninuria. Tohoku J. Exp.

Med. 95: 235–242,

[156213]1968.

[156214]

[156215]

[156216]15155.

[156217]Whelan, D. T.; Scriver, C. R.: Cystathioninuria and renal
iminoglycinuria

[156218]in a pedigree: a perspective on counseling. New Eng. J.

Med. 278:

[156219]924–927, 1968.

[156220]

[156221]

[156222]15156.

[156223]Haworth, J. C.; Gourley, B.; Hadorn, B.; Sumida, C.: Malab-
sorption

[156224]and growth failure due to intestinal enterokinase defi-
ciency. J.

[156225]Pediat. 78: 481–490, 1971.

[156226]

[156227]

[156228]15157.

[156229]Haworth, J. C.; Hadorn, B.; Gourley, B.; Prasad, A.;
Troesch, V.

[156230]: Intestinal enterokinase deficiency: occurrence in two sibs
and age

[156231]dependency of clinical expression. Arch. Dis. Child. 50:
277-282,

[156232]1975.

[156233]

[156234]

[156235]15158.

[156236]Holzinger, A.; Maier, E. M.; Buck, C.; Mayerhofer, P. U.;
Kappler,

[156237]M.; Haworth, J. C.; Moroz, S. P.; Hadorn, H.-B.; Sadler, J.
E.; Roscher,

[156238]A. A.: Mutations in the proenteropeptidase gene are the
molecular

[156239]cause of congenital enteropeptidase deficiency. Am. J.
Hum. Genet. 70:

[156240]20-25, 2002.

[156241]

[156242]

[156243]15159.

[156244]Joenje, H.: Fanconi anaemia complementation groups in
Germany

[156245]and The Netherlands. Hum. Genet. 97: 280–282, 1996.

[156246]

[156247]

[156248]15160.

[156249]Joenje, H.; Oostra, A. B.; Wijker, M.; di Summa, F. M.; van
Berkel,

[156250]C. G. M.; Rooimans, M. A.; Ebell, W.; van Weel, M.; Pronk, J.
C.;

[156251]Buchwald, M.; Arwert, F.: Evidence for at least eight Fan-
coni anemia

[156252]genes. Am. J. Hum. Genet. 61: 940–944, 1997.

[156253]

[156254]

[156255]15161.

[156256]Bongiovanni, A. M.: Disorders of adrenocortical steroid biogenesis.In:

[156257]Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic

[156258]Basis of Inherited Disease. New York: McGraw-Hill (pub.) (3rd

[156259]ed.): 1972. Pp. 857-885.

[156260]

[156261]

[156262]15162.

[156263]Bongiovanni, A. M.; Root, A. W.: The adrenogenital syndrome. New

[156264]Eng. J. Med. 268: 1283-1289; 1342-1351; 1391-1399, 1963.

[156265]

[156266]

[156267]15163.

[156268]Boudreaux, D.; Waisman, J.; Skinner, D. G.; Low, R.: Giant adrenal

[156269]myelolipoma and testicular interstitial cell tumor in a man with congenital

[156270]21-hydroxylase deficiency. Am. J. Surg. Path. 3: 109-123,
1979.

[156271]

[156272]

[156273]15164.

[156274]Carroll, M. C.; Campbell, R. D.; Porter, R. R.: Mapping of
steroid

[156275]21-hydroxylase genes adjacent to complement compo-
nent C4 genes in

[156276]HLA, the major histocompatibility complex in man. Proc.
Nat. Acad.

[156277]Sci. 82: 521-525, 1985.

[156278]

[156279]

[156280]15165.

[156281]Chaplin, D. D.; Galbraith, L. J.; Seidman, J. G.; White, P. C.;

[156282]Parker, K. L.: Nucleotide sequence analysis of murine
21-hydroxylase

[156283]gene: mutations affecting gene expression. Proc. Nat.
Acad. Sci. 83:

[156284]9601–9605, 1986.

[156285]

[156286]

[156287]15166.

[156288]Charmandari, E.; Hindmarsh, P. C.; Johnston, A.; Brook, C.
G.

[156289]D.: Congenital adrenal hyperplasia due to 21-hydroxylase
deficiency:

[156290]alterations in cortisol pharmacokinetics at puberty. J. Clin.
Endocr.

[156291]Metab. 86: 2701–2708, 2001.

[156292]

[156293]

[156294]15167.

[156295]Childs, B.; Grumbach, M. M.; Van Wyk, J. J.: Virilizing
adrenal

[156296]hyperplasia: genetic and hormonal studies. J. Clin. Invest.
35:

[156297]213–222, 1956.

[156298]

[156299]

[156300]15168.

[156301]Chiou, S.-H.; Hu, M.-C.; Chung, B.: A missense mutation
at ile172-to-asn

[156302]or arg356-to-trp causes steroid 21-hydroxylase defi-
ciency. J. Biol.

[156303]Chem. 265: 3549-3552, 1990.

[156304]

[156305]

[156306]15169.

[156307]Chrousos, G. P.; Loriaux, D. L.; Mann, D. L.; Cutler, G. B.,
Jr.

[156308]: Late-onset 21-hydroxylase deficiency mimicking idio-
pathic hirsutism

[156309]or polycystic ovarian disease: an allelic variant of congeni-
tal virilizing

[156310]adrenal hyperplasia with a milder enzymatic defect. Ann.
Intern.

[156311]Med. 96: 143-148, 1982.

[156312]

[156313]

[156314]15170.

[156315]Collier, S.; Tassabehji, M.; Strachan, T.: A de novo pathological

[156316]point mutation at the 21-hydroxylase locus: implications for gene

[156317]conversion in the human genome. Nature Genet. 3: 260-265, 1993.

[156318]

[156319]

[156320]15171.

[156321]Cooper, D. Y.; Levin, S.; Narasimhulu, S.; Rosenthal, O.; Estabrook,

[156322]R. W.: Photochemical action spectrum of the terminal oxidase of mixed

[156323]function oxidase systems. Science 147: 400-402, 1965.

[156324]

[156325]

[156326]15172.

[156327]Cornean, R. E.; Hindmarsh, P. C.; Brook, C. G. D.: Obesity

in

[156328]21-hydroxylase deficient patients. Arch. Dis. Child. 78:

261-263,

[156329]1998.

[156330]

[156331]

[156332]15173.

[156333]Couillin, P.; Kottler-Missonnier, M. L.; Grisard, M. C.; Hors,

[156334]J.; Feingold, J.; Boue, J.; Boue, A.: HLA-A, B, C, DR alleles

in

[156335]congenital adrenal hyperplasia. Hum. Genet. 53: 389-392,

1980.

[156336]

[156337]

[156338]15174.

[156339]Cutfield, R. G.; Bateman, J. M.; Odell, W. D.: Infertility

caused

[156340]by bilateral testicular masses secondary to congenital

adrenal hyperplasia

[156341](21-hydroxylase deficiency). Fertil. Steril. 40: 809-814,

1983.

[156342]

[156343]

[156344]15175.

[156345]Cutler, G. B., Jr.; Laue, L.: Congenital adrenal hyperplasia

[156346]due to 21-hydroxylase deficiency. New Eng. J. Med. 323:

1806-1813,

[156347]1990.

[156348]

[156349]

[156350]15176.

[156351]Dacou-Voutetakis, C.; Dracopoulou, M.: High incidence of
molecular

[156352]defects of the CYP21 gene in patients with premature
adrenarche. J.

[156353]Clin. Endocr. Metab. 84: 1570-1574, 1999.

[156354]

[156355]

[156356]15177.

[156357]Day, D. J.; Speiser, P. W.; Schulze, E.; Bettendorf, M.; Fitness,

[156358]J.; Barany, F.; White, P. C.: Identification of non-amplifying CYP21

[156359]genes when using PCR-based diagnosis of 21-hydroxylase deficiency

[156360]in congenital adrenal hyperplasia (CAH) affected pedigrees. Hum.

[156361]Molec. Genet. 5: 2039–2048, 1996.

[156362]

[156363]

[156364]15178.

[156365]Deneux, C.; Tardy, V.; Dib, A.; Mornet, E.; Billaud, L.; Charron,

[156366]D.; Morel, Y.; Kuttann, F.: Phenotype-genotype correlation in 56

[156367]women with nonclassical congenital adrenal hyperplasia due to 21-hydroxylase

[156368]deficiency. J. Clin. Endocr. Metab. 86: 207–213, 2001.

[156369]

[156370]

[156371]15179.

[156372]Donohoue, P. A.; Jospe, N.; Migeon, C. J.; Van Dop, C.:

Two distinct

[156373]areas of unequal crossingover within the steroid

21-hydroxylase genes

[156374]produce absence of CYP21B. Genomics 5: 397-406, 1989.

[156375]

[156376]

[156377]15180.

[156378]Dracopoulou-Vabouli, M.; Maniati-Christidi, M.; Dacou-

Voutetakis,

[156379]C.: The spectrum of molecular defects of the CYP21 gene

in the Hellenic

[156380]population: variable concordance between genotype and

phenotype in

[156381]the different forms of congenital adrenal hyperplasia. J.

Clin. Endocr.

[156382]Metab. 86: 2845-2848, 2001.

[156383]

[156384]

[156385]15181.

[156386]Duck, S. C.: Malignancy associated with congenital adrenal hyperplasia. J.

[156387]Pediat. 99: 423–424, 1981.

[156388]

[156389]

[156390]15182.

[156391]Dumic, M.; Brkljacic, L.; Mardesic, D.; Plavsic, V.; Lukenda,

[156392]M.; Kastelan, A.: 'Cryptic' form of congenital adrenal hyperplasia

[156393]due to 21-hydroxylase deficiency in the Yugoslav population. Acta

[156394]Endocr. 109: 386–392, 1985.

[156395]

[156396]

[156397]15183.

[156398]Dupont, B.; Smithwick, E. M.; Oberfield, S. E.; Lee, T. D.; Levine,

[156399]L. S.: Close genetic linkage between HLA and congenital

adrenal hyperplasia

[156400](21-hydroxylase deficiency). Lancet II: 1309-1312, 1977.

[156401]

[156402]

[156403]15184.

[156404]Ezquieta, B.; Oliver, A.; Gracia, R.; Gancedo, P. G.: Analysis

[156405]of steroid 21-hydroxylase gene mutations in the Spanish
population. Hum.

[156406]Genet. 96: 198-204, 1995.

[156407]

[156408]

[156409]15185.

[156410]Ferenczi, A.; Garami, M.; Kiss, E.; Pek, M.; Sasvari-Szekely,

[156411]M.; Barta, C.; Staub, M.; Solyom, J.; Fekete, G.: Screening
for mutations

[156412]of 21-hydroxylase gene in Hungarian patients with con-
genital adrenal

[156413]hyperplasia. J. Clin. Endocr. Metab. 84: 2369-2372, 1999.

[156414]

[156415]

[156416]15186.

[156417]Fitness, J.; Dixit, N.; Webster, D.; Torresani, T.; Pergolizzi,

[156418]R.; Speiser, P. W.; Day, D. J.: Genotyping of CYP21, linked
chromosome

[156419]6p markers, and a sex-specific gene in neonatal screening
for congenital

[156420]adrenal hyperplasia. J. Clin. Endocr. Metab. 84: 960–966,
1999.

[156421]

[156422]

[156423]15187.

[156424]Fleischnick, E.; Awdeh, Z. L.; Raum, D.; Granados, J.;
Alosco,

[156425]S. M.; Crigler, J. F., Jr.; Gerald, P. S.; Giles, C. M.; Yunis, E.

[156426]J.; Alper, C. A.: Extended MHC haplotypes in
21-hydroxylase-deficiency

[156427]congenital adrenal hyperplasia: shared genotypes in unre-
lated patients. Lancet I:

[156428]152–156, 1983.

[156429]

[156430]

[156431]15188.

[156432]Flori, J.; Tongio, M. M.; Kurtz, F.; Mayer, S.; Juif, J. G.:

[156433]Liaison entre le gene responsable de l'hyperplasie surre-
nalienne par

[156434]deficit en 21 hydroxylase et les marqueurs du bras court
du chromosome

[156435]6. J. Genet. Hum. 28: 185–193, 1980.

[156436]

[156437]

[156438]15189.

[156439]Forest, M. G.; Betuel, H.; David, M.: Prenatal treatment in
congenital

[156440]adrenal hyperplasia due to 21–hydroxylase deficiency:
up–date 88 of

[156441]the French multicentric study. Endocr. Res. 15: 277–301,
1989.

[156442]

[156443]

[156444]15190.

[156445]Galal, O. M.; Rudd, B. T.; Drayer, N. M.: Evaluation of deficiency

[156446]of 21-hydroxylation in patients with congenital adrenal hyperplasia. Arch.

[156447]Dis. Child. 43: 410-414, 1968.

[156448]

[156449]

[156450]15191.

[156451]Ghanem, N.; Lobaccaro, J. M.; Buresi, C.; Abbal, M.; Halaby, G.;

[156452]Sultan, C.; Lefranc, G.: Defective, deleted or converted CYP21B gene

[156453]and negative association with a rare restriction fragment length polymorphism

[156454]allele of the factor B gene in congenital adrenal hyperplasia. Hum.

[156455]Genet. 86: 117-125, 1990.

[156456]

[156457]

[156458]15192.

[156459]Rassool, F. V.; Le Beau, M. M.; Shen, M.-L.; Neilly, M. E.;

Espinosa,

[156460]R., III; Ong, S. T.; Boldog, F.; Drabkin, H.; McCarroll, R.;

McKeithan,

[156461]T. W.: Direct cloning of DNA sequences from the common

fragile site

[156462]region at chromosome band 3p14.2. Genomics 35:

109-117, 1996.

[156463]

[156464]

[156465]15193.

[156466]Rudduck, C.; Franzen, G.: A new heritable fragile site on

human

[156467]chromosome 3. Hereditas 98: 297-299, 1983.

[156468]

[156469]

[156470]15194.

[156471]Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association

[156472]of FHIT (fragile histidine triad), a candidate tumour sup-

pressor gene,

[156473]with the ubiquitin–conjugating enzyme hUBC9. *Biochem. J.*

352: 443–448,

[156474]2000.

[156475]

[156476]

[156477]15195.

[156478]Shiraishi, T.; Druck, T.; Mimori, K.; Flomenberg, J.; Berk,
L.;

[156479]Alder, H.; Miller, W.; Huebner, K.; Croce, C. M.: Sequence
conservation

[156480]at human and mouse orthologous common fragile re-
gions, FRA3B/FHIT

[156481]and Fra14A2/Fhit. *Proc. Nat. Acad. Sci.* 98: 5722–5727,
2001.

[156482]

[156483]

[156484]15196.

[156485]Sozzi, G.; Veronese, M. L.; Negrini, M.; Baffa, R.; Cotticelli,

[156486]M. G.; Inoue, H.; Tornielli, S.; Pilotti, S.; De Gregorio, L.;

Pastorino,

[156487]U.; Pierotti, M. A.; Ohta, M.; Huebner, K.; Croce, C. M.: The
FHIT

[156488]gene at 3p14.2 is abnormal in lung cancer. Cell 85:
17-26, 1996.

[156489]

[156490]

[156491]15197.

[156492]Stein, C. K.; Glover, T. W.; Palmer, J. L.; Glisson, B. S.: Di-
rect

[156493]correlation between FRA3B expression and cigarette
smoking. Genes

[156494]Chromosomes Cancer 34: 333-340, 2002.

[156495]

[156496]

[156497]15198.

[156498]Virgilio, L.; Shuster, M.; Gollin, S. M.; Veronese, M. L.;
Ohta,

[156499]M.; Huebner, K.; Croce, C. M.: FHIT gene alterations in
head and

[156500]neck squamous cell carcinomas. Proc. Nat. Acad. Sci. 93:

9770-9775,

[156501]1996.

[156502]

[156503]

[156504]15199.

[156505]Wegner, R.-D.: A new inducible fragile site on chromosome 3 (p14.2)

[156506]in human lymphocytes. Hum. Genet. 63: 297-298, 1983.

[156507]

[156508]

[156509]15200.

[156510]Wegner, R.-D.: Reply to the letter of A. Markkanen, S. Knuutila,

[156511]and A. de la Chapelle. (Letter) Hum. Genet. 65: 218 only, 1983.

[156512]

[156513]

[156514]15201.

[156515]Zanesi, N.; Fidanza, V.; Fong, L. Y.; Mancini, R.; Druck, T.;

[156516]Valtieri, M.; Rudiger, T.; McCue, P. A.; Croce, C. M.; Huebner, K.

[156517]: The tumor spectrum in FHIT-deficient mice. Proc. Nat. Acad. Sci. 98:

[156518]10250–10255, 2001.

[156519]

[156520]

[156521]15202.

[156522]Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayasi, Y.;

[156523]Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura, N.: Prediction

[156524]of the coding sequences of unidentified human genes. VI. The coding

[156525]sequences of 80 new genes (KIAA0201–KIAA0280) deduced by analysis

[156526]of cDNA clones from cell line KG-1 and brain. DNA Res. 3: 321–329,

[156527]1996.

[156528]

[156529]

[156530]15203.

[156531]Palfree, R. G. E.; Sadro, L. C.; Solomon, S.: The gene encoding

[156532]the human corticostatin HP-4 precursor contains a recent 86-base duplication

[156533]and is located on chromosome 8. Molec. Endocr. 7: 199-205, 1993.

[156534]

[156535]

[156536]15204.

[156537]Singh, A.; Bateman, A.; Zhu, Q. Z.; Shimasaki, S.; Esch, F.; Solomon,

[156538]S.: Structure of the novel human granulocyte peptide with anti-ACTH

[156539]activity. Biochem. Biophys. Res. Commun. 155: 524-529, 1988.

[156540]

[156541]

[156542]15205.

[156543]Lee, E.; Marcucci, M.; Daniell, L.; Pypaert, M.; Weisz, O. A.;

[156544]Ochoa, G.-C.; Farsad, K.; Wenk, M. R.; De Camilli, P.: Am-
phiphysin

[156545]2 (Bin1) and T-tubule biogenesis in muscle. Science 297:
1193-1196,

[156546]2002.

[156547]

[156548]

[156549]15206.

[156550]Negorev, D.; Riethman, H.; Wechsler-Reya, R.; Sakamuro,
D.; Prendergast,

[156551]G. C.; Simon, D.: The Bin1 gene localizes to human chro-
mosome 2q14

[156552]by PCR analysis of somatic cell hybrids and fluorescence
in situ hybridization. Genomics 33:

[156553]329-331, 1996.

[156554]

[156555]

[156556]15207.

[156557]Sakamuro, D.; Elliott, K. J.; Wechsler-Reya, R.; Prender-
gast, G.

[156558]C.: BIN1 is a novel MYC–interacting protein with features
of a tumour

[156559]suppressor. Nature Genet. 14: 69–77, 1996.

[156560]

[156561]

[156562]15208.

[156563]Ishizuka, T.; Ahmad, I.; Kita, K.; Sonoda, T.; Ishijima, S.;
Sawa,

[156564]K.; Suzuki, N.; Tatibana, M.: The human phosphoribo-
sylpyrophosphate

[156565]synthetase–associated protein 39 gene (PRPSAP1) is lo-
cated in the

[156566]chromosome region 17q24–q25. Genomics 33: 332–334,
1996.

[156567]

[156568]

[156569]15209.

[156570]Ishizuka, T.; Kita, K.; Sonoda, T.; Ishijima, S.; Sawa, K.;
Suzuki,

[156571]N.; Tatibana, M.: Cloning and sequencing of human com-
plementary DNA

[156572]for the phosphoribosylpyrophosphate synthetase-associated protein

[156573]39. Biochim. Biophys. Acta 1306: 27–30, 1996.

[156574]

[156575]

[156576]15210.

[156577]Kita, K.; Ishizuka, T.; Ishijima, S.; Sonoda, T.; Tatibana, M.

[156578]: A novel 39-kDa phosphoribosylpyrophosphate synthetase-associated

[156579]protein of rat liver: cloning, high sequence similarity to the catalytic

[156580]subunits, and a negative regulatory role. J. Biol. Chem. 269: 8334–8340,

[156581]1994.

[156582]

[156583]

[156584]15211.

[156585]Endo, Y.; Sato, Y.; Matsushita, M.; Fujita, T.: Cloning and characterization

[156586]of the human lectin P35 gene and its related gene. Ge-

nomics 36:

[156587]515–521, 1996.

[156588]

[156589]

[156590]15212.

[156591]Ichijo, H.; Hellman, U.; Wernstedt, C.; Gonez, L. J.; Claes–
son–Welsh,

[156592]L.; Heldin, C.–H.; Miyazono, K.: Molecular cloning and
characterization

[156593]of ficolin, a multimeric protein with fibrinogen– and colla–
gen–like

[156594]domains. J. Biol. Chem. 268: 14505–14513, 1993.

[156595]

[156596]

[156597]15213.

[156598]Ichijo, H.; Ronnstrand, L.; Miyagawa, K.; Ohashi, H.;
Heldin, C.–H.;

[156599]Miyazono, K.: Purification of transforming growth factor–
beta 1 binding

[156600]proteins from porcine uterus membranes. J. Biol. Chem.

266: 22459–22464,

[156601]1991.

[156602]

[156603]

[156604]15214.

[156605]Lu, J.; Tay, P. N.; Kon, O. L.; Reid, K. B. M.: Human ficolin:

[156606]cDNA cloning, demonstration of peripheral blood leuco-
cytes as the

[156607]major site of synthesis and assignment of the gene to
chromosome 9. Biochem.

[156608]J. 313: 473–478, 1996.

[156609]

[156610]

[156611]15215.

[156612]Matsushita, M.; Endo, Y.; Taira, S.; Sato, Y.; Fujita, T.;
Ichikawa,

[156613]N.; Nakata, M.; Mizuochi, T.: A novel human serum lectin
with collagen–

[156614]and fibrinogen–like domains that functions as an opsonin.
J. Biol.

[156615]Chem. 271: 2448–2454, 1996.

[156616]

[156617]

[156618]15216.

[156619]Bronstein, J. M.; Kozak, C. A.; Chen, X.–N.; Wu, S.; Dan-
ciger,

[156620]M.; Korenberg, J. R.; Farber, D. B.: Chromosomal localiza-
tion of

[156621]murine and human oligodendrocyte–specific protein
genes. Genomics 34:

[156622]255–257, 1996.

[156623]

[156624]

[156625]15217.

[156626]Wayne, S.; Robertson, N. G.; DeClau, F.; Chen, N.; Verho-
even, K.;

[156627]Prasad, S.; Tranebjarg, L.; Morton, C. C.; Ryan, A. F.; Van
Camp,

[156628]G.; Smith, R. J. H.: Mutations in the transcriptional activa-
tor EYA4

[156629]cause late-onset deafness at the DFNA10 locus. Hum.

Molec. Genet. 10:

[156630]195–200, 2001.

[156631]

[156632]

[156633]15218.

[156634]Kanai, Y.; Miura, K.; Uehara, T.; Amagai, M.; Takeda, O.;

Tanuma,

[156635]S.; Kurosawa, Y.: Natural occurrence of Nuc in the sera of
autoimmune-prone

[156636]MRL/lpr mice. Biochem. Biophys. Res. Commun. 196:
729–736, 1993.

[156637]

[156638]

[156639]15219.

[156640]Miura, K.; Hirai, M.; Kanai, Y.; Kurosawa, Y.: Organization
of

[156641]the human gene for nucleobindin (NUC) and its chromo-
somal assignment

[156642]to 19q13.2–q13.4. Genomics 34: 181–186, 1996.

[156643]

[156644]

[156645]15220.

[156646]Miura, K.; Titani, K.; Kurosawa, Y.; Kanai, Y.: Molecular cloning

[156647]of nucleobindin, a novel DNA-binding protein that contains both a

[156648]signal peptide and a leucine zipper structure. Biochem. Biophys.

[156649]Res. Commun. 187: 375–380, 1992.

[156650]

[156651]

[156652]15221.

[156653]Connelly, M. A.; Grady, R. C.; Mushinski, J. F.; Marcu, K. B.:

[156654]PANG, a gene encoding a neuronal glycoprotein, is ectopically activated

[156655]by intracisternal A-type particle long terminal repeats in murine

[156656]plasmacytomas. Proc. Nat. Acad. Sci. 91: 1337–1341, 1994.

[156657]

[156658]

[156659]15222.

[156660]Mock, B. A.; Connelly, M. A.; McBride, O. W.; Kozak, C. A.;
Marcu,

[156661]K. B.: Plasmacytoma-associated neuronal glycoprotein,
Pang, maps

[156662]to mouse chromosome 6 and human chromosome 3. Ge-
nomics 34: 226-228,

[156663]1996.

[156664]

[156665]

[156666]15223.

[156667]Gow, A.; Southwood, C. M.; Li, J. S.; Pariali, M.; Riordan, G.

[156668]P.; Brodie, S. E.; Danias, J.; Bronstein, J. M.; Kachar, B.;
Lazzarini,

[156669]R. A.: CNS myelin and Sertoli cell tight junction strands are
absent

[156670]in Osp/claudin-11 null mice. Cell 99: 649-659, 1999.

[156671]

[156672]

[156673]15224.

[156674]Noben-Trauth, K.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[156675]A.; Sonoda, G.; Testa, J. R.; Klempnauer, K.-H.: Mybl2 (Bmyb) maps

[156676]to mouse chromosome 2 and human chromosome 20q13.1. Genomics 35:

[156677]610-612, 1996.

[156678]

[156679]

[156680]15225.

[156681]Ando, A.; Kikuti, Y. Y.; Shigenari, A.; Kawata, H.; Okamoto, N.;

[156682]Shiina, T.; Chen, L.; Ikemura, T.; Abe, K.; Kimura, M.; Inoko, H.

[156683]: cDNA cloning of the human homologues of the mouse Ke4 and Ke6 genes

[156684]at the centromeric end of the human MHC region. Genomics 35: 600-602,

[156685]1996.

[156686]

[156687]

[156688]15226.

[156689]Aziz, N.; Maxwell, M. M.; Brenner, B. M.: Coordinate regulation

[156690]of 11-beta-HSD and Ke6 gene in cpk mouse: implications for steroid

[156691]metabolic defect in PKD. Am. J. Physiol. 267: F791-F797, 1994.

[156692]

[156693]

[156694]15227.

[156695]Aziz, N.; Maxwell, M. M.; St.-Jacques, B.; Brenner, B. M.: Downregulation

[156696]of Ke 6, a novel gene encoded within the major histocompatibility

[156697]complex, in murine polycystic kidney disease. Molec. Cell. Biol. 13:

[156698]1847-1853, 1993.

[156699]

[156700]

[156701]15228.

[156702]Kikuti, Y. Y.; Tamiya, G.; Ando, A.; Chen, L.; Kimura, M.;
Ferreira,

[156703]E.; Tsuji, K.; Trowsdale, J.; Inoko, H.: Physical mapping
220 kb

[156704]centromeric of the human MHC and DNA sequence analysis of the 43-kb

[156705]segment including the RING1, HKE6, and HKE4 genes. Genomics 42:

[156706]422–435, 1997.

[156707]

[156708]

[156709]15229.

[156710]Aziz, N.; Anderson, E.; Lee, G. Y.; Woo, D. D. L.: Arrested
testis

[156711]development in the cpk mouse may be the result of abnormal steroid

[156712]metabolism. Molec. Cell. Endocr. 171: 83–88, 2001.

[156713]

[156714]

[156715]15230.

[156716]Fomitcheva, J.; Baker, M. E.; Anderson, E.; Lee, G. Y.; Aziz, N.

[156717]: Characterization of Ke 6, a new 17-beta-hydroxysteroid dehydrogenase,

[156718]and its expression in gonadal tissues. J. Biol. Chem. 273: 22664-22671,

[156719]1998.

[156720]

[156721]

[156722]15231.

[156723]Khazak, V.; Estojak, J.; Cho, H.; Majors, J.; Sonoda, G.; Testa,

[156724]J. R.; Golemis, E. A.: Analysis of the interaction of the novel RNA

[156725]polymerase II (pol II) subunit hsRPB4 with its partner hsRPB7 and

[156726]with pol II. Molec. Cell. Biol. 18: 1935-1945, 1998.

[156727]

[156728]

[156729]15232.

[156730]Schoen, T. J.; Chandrasekharappa, S. C.; Guru, S. C.;
Mazuruk,

[156731]K.; Chader, G. J.; Rodriguez, I. R.: Human gene for the RNA
polymerase

[156732]II seventh subunit (hsRPB7): structure, expression and
chromosomal

[156733]localization. Biochim. Biophys. Acta 1353: 39–49, 1997.

[156734]

[156735]

[156736]15233.

[156737]Cheung, A. H.; Stewart, R. J.; Marsden, P. A.: Endothelial
Tie2/Tek

[156738]ligands angiopoietin–1 (ANGPT1) and angiopoietin–2
(ANGPT2): regional

[156739]localization of the human genes to 8q22.3–q23 and 8p23.
Genomics 48:

[156740]389–391, 1998.

[156741]

[156742]

[156743]15234.

[156744]Davis, S.; Aldrich, T. H.; Jones, P. F.; Acheson, A.; Comp-
ton,

[156745]D. L.; Jain, V.; Ryan, T. E.; Bruno, J.; Radziejewski, C.;
Maisonpierre,

[156746]P. C.; Yancopoulos, G. D.: Isolation of angiopoietin-1, a
ligand

[156747]for the TIE2 receptor, by secretion-trap expression
cloning. Cell 87:

[156748]1161-1169, 1996.

[156749]

[156750]

[156751]15235.

[156752]Folkman, J.; D'Amore, P. A.: Blood vessel formation: what
is its

[156753]molecular basis? Cell 87: 1153-1155, 1996.

[156754]

[156755]

[156756]15236.

[156757]Marziliano, N.; Crovella, S.; Audero, E.; Pecile, V.; Bus-
solino,

[156758]F.; Amoroso, A.; Garagna, S.: Genetic mapping of the
mouse homologue

[156759]of the human angiopoietin-1 gene (Agpt) to mouse chro-
mosome 9E2 by

[156760]in situ hybridization. Cytogenet. Cell Genet. 87: 199-200,
1999.

[156761]

[156762]

[156763]15237.

[156764]Sato, T. N.; Tozawa, Y.; Deutsch, U.; Wolburg-Buchholz,
K.; Fujiwara,

[156765]Y.; Gendron-Maguire, M.; Gridley, T.; Wolburg, H.; Risau,
W.; Qin,

[156766]Y.: Distinct roles of the receptor tyrosine kinases Tie-1
and Tie-2

[156767]in blood vessel formation. Nature 376: 70-73, 1995.

[156768]

[156769]

[156770]15238.

[156771]Suri, C.; Jones, P. F.; Patan, S.; Bartunkova, S.; Maison-
pierre,

[156772]P. C.; Davis, S.; Sato, T. N.; Yancopoulos, G. D.: Requisite
role

[156773]of angiopoietin-1, a ligand for the TIE2 receptor, during
embryonic

[156774]angiogenesis. Cell 87: 1171-1180, 1996.

[156775]

[156776]

[156777]15239.

[156778]Suri, C.; McClain, J.; Thurston, G.; McDonald, D. M.; Zhou,
H.;

[156779]Oldmixon, E. H.; Sato, T. N.; Yancopoulos, G. D.: Increased
vascularization

[156780]in mice overexpressing angiopoietin-1. Science 282:
468-471, 1998.

[156781]

[156782]

[156783]15240.

[156784]Valenzuela, D. M.; Griffiths, J. A.; Rojas, J.; Aldrich, T. H.;

[156785]Jones, P. F.; Zhou, H.; McClain, J.; Copeland, N. G.; Gilbert, D.

[156786]J.; Jenkins, N. A.; Huang, T.; Papadopoulos, N.; Maison-pierre, P.

[156787]C.; Davis, S.; Yancopoulos, G. D.: Angiopoietins 3 and 4: diverging

[156788]gene counterparts in mice and humans. Proc. Nat. Acad. Sci. 96:

[156789]1904–1909, 1999.

[156790]

[156791]

[156792]15241.

[156793]Chestkov, A. V.; Baka, I. D.; Kost, M. V.; Georgiev, G. P.; Buchman,

[156794]V. L.: The d4 gene family in the human genome. Genomics 36: 174–177,

[156795]1996.

[156796]

[156797]

[156798]15242.

[156799]Gabig, T. G.; Crean, C. D.; Klenk, A.; Long, H.; Copeland, N. G.;

[156800]Gilbert, D. J.; Jenkins, N. A.; Quincey, D.; Parente, F.; Lespinasse,

[156801]F.; Carle, G. F.; Gaudray, P.; and 13 others: Expression and chromosomal

[156802]localization of the Requiem gene. Mammalian Genome 9: 660–665, 1998.

[156803]

[156804]

[156805]15243.

[156806]Gabig, T. G.; Mantel, P. L.; Rosli, R.; Crean, C. D.: Requiem:

[156807]a novel zinc finger gene essential for apoptosis in myeloid cells. J.

[156808]Biol. Chem. 269: 29515–29519, 1994.

[156809]

[156810]

[156811]15244.

[156812]Smith, J. E., Jr.; Bollekens, J. A.; Inghirami, G.; Takeshita,

[156813]K.: Cloning and mapping of the MEIS1 gene, the human

homolog of a

[156814]murine leukemogenic gene. Genomics 43: 99–103, 1997.

[156815]

[156816]

[156817]15245.

[156818]Duperat, V. G.; Jacquelin, B.; Boisseau, P.; Arveiler, B.;
Nurden,

[156819]A. T.: Protease-activated receptor genes are clustered on
5q13. Blood 92:

[156820]25–31, 1998.

[156821]

[156822]

[156823]15246.

[156824]Ishihara, H.; Connolly, A. J.; Zeng, D.; Kahn, M. L.; Zheng,
Y.

[156825]W.; Timmons, C.; Tram, T.; Coughlin, S. R.: Protease-acti-
vated receptor

[156826]3 is a second thrombin receptor in humans. Nature 386:
502–506,

[156827]1997.

[156828]

[156829]

[156830]15247.

[156831]Kahn, M. L.; Zheng, Y. W.; Huang, W.; Bigornia, V.; Zeng, D.; Moff,

[156832]S.; Farese, R. V., Jr.; Tam, C.; Coughlin, S. R.: A dual
thrombin

[156833]receptor system for platelet activation. Nature 394:
690–694, 1998.

[156834]

[156835]

[156836]15248.

[156837]Nakanishi–Matsui, M.; Zheng, Y.–W.; Sulciner, D. J.; Weiss, E.

[156838]J.; Ludeman, M. J.; Coughlin, S. R.: PAR3 is a cofactor for
PAR4

[156839]activation by thrombin. Nature 404: 609–613, 2000.

[156840]

[156841]

[156842]15249.

[156843]Sambrano, G. R.; Weiss, E. J.; Zheng, Y.-W.; Huang, W.;
Coughlin,

[156844]S. R.: Role of thrombin signalling in platelets in haemosta-
sis and

[156845]thrombosis. Nature 413: 74–78, 2001.

[156846]

[156847]

[156848]15250.

[156849]Schmidt, V.; Wainer, J.; Bahou, W.: Identification and char-
acterization

[156850]of the human proteolytically activated receptor-3 (PAR-3)
gene within

[156851]the PAR gene cluster by positional mapping. (Abstract)
Blood (suppl.

[156852]1) 90: 284a only, 1997.

[156853]

[156854]

[156855]15251.

[156856]Paulweber, B.; Wiebusch, H.; Miesenboeck, G.; Funke, H.;
Assmann,

[156857]G.; Hoelzl, B.; Sippl, M. J.; Friedl, W.; Patsch, J. R.; Sand-
hofer,

[156858]F.: Molecular basis of lipoprotein lipase deficiency in two
Austrian

[156859]families with type I hyperlipoproteinemia. Atherosclerosis
86: 239–250,

[156860]1991.

[156861]

[156862]

[156863]15252.

[156864]Abe, M.; Suzuki, T.; Deguchi, T.: An improved method for
genotyping

[156865]of N-acetyltransferase polymorphism by polymerase chain
reaction. Jpn.

[156866]J. Hum. Genet. 38: 163–168, 1993.

[156867]

[156868]

[156869]15253.

[156870]Azad Khan, A. K.; Nurazzaman, M.; Truelove, S. C.: The
effect

[156871]of the acetylator phenotype on the metabolism of sul-
phasalazine in

[156872]man. J. Med. Genet. 20: 30–36, 1983.

[156873]

[156874]

[156875]15254.

[156876]Bell, D. A.; Taylor, J. A.; Butler, M. A.; Stephens, E. A.;
Wiest,

[156877]J.; Brubaker, L. H.; Kadlubar, F. F.; Lucier, G. W.: Geno-
type/phenotype

[156878]discordance for human arylamine N-acetyltransferase
(NAT2) reveals

[156879]a new slow-acetylator allele common in African-Amer-
icans. Carcinogenesis 14:

[156880]1689–1692, 1993.

[156881]

[156882]

[156883]15255.

[156884]Blum, M.; Grant, D.; Demierre, A.; Meyer, U. A.: N-
acetylation

[156885]pharmacogenetics: a gene deletion causes absence of arylamine N-acetyltransferase

[156886]in liver of slow acetylators rabbits. Proc. Nat. Acad. Sci. 86: 9554-9557,

[156887]1989.

[156888]

[156889]

[156890]15256.

[156891]Burrows, A. W.; Hockaday, T. D. R.; Mann, J. I.; Taylor, J. G.

[156892]: Diabetic dimorphism according to acetylator status. Brit. Med.

[156893]J. 1: 208-210, 1978.

[156894]

[156895]

[156896]15257.

[156897]Cascorbi, I.; Brockmoller, J.; Bauer, S.; Reum, T.; Roots, I.:

[156898]NAT2*12A (803A-G) codes for rapid arylamine N-acetylation in humans. Pharmacogenetics 6:

[156899]257-259, 1996.

[156900]

[156901]

[156902]15258.

[156903]Cascorbi, I.; Drakoulis, N.; Brockmoller, J.; Maurer, A.;
Sperling,

[156904]K.; Roots, I.: Arylamine N-acetyltransferase (NAT2) muta-
tions and

[156905]their allelic linkage in unrelated Caucasian individuals:
correlation

[156906]with phenotypic activity. Am. J. Hum. Genet. 57: 581–592,
1995.

[156907]

[156908]

[156909]15259.

[156910]Delomenie, C.; Sica, L.; Grant, D. M.; Krishnamoorthy, R.;
Dupret,

[156911]J.-M.: Genotyping of the polymorphic N-acetyltransferase
(NAT2*)

[156912]gene locus in two native African populations. Pharmaco-
genetics 6:

[156913]177–185, 1996.

[156914]

[156915]

[156916]15260.

[156917]Evans, D. A. P.: N-acetyltransferase. Pharm. Therap. 42:
157–234,

[156918]1989.

[156919]

[156920]

[156921]15261.

[156922]Evans, D. A. P.; Manley, K. A.; McKusick, V. A.: Genetic
control

[156923]of isoniazid metabolism in man. Brit. Med. J. 2: 485–491,
1960.

[156924]

[156925]

[156926]15262.

[156927]Evans, D. A. P.; White, T. A.: Human acetylation polymor-
phism. J.

[156928]Lab. Clin. Med. 63: 394–403, 1964.

[156929]

[156930]

[156931]15263.

[156932]Frymoyer, J. W.; Jacox, R. F.: Studies of genetically controlled

[156933]sulfadiazine acetylation in rabbit livers: possible identification

[156934]of the heterozygous trait. J. Lab. Clin. Med. 62: 905–909, 1963.

[156935]

[156936]

[156937]15264.

[156938]Glowinski, I. B.; Weber, W. W.: Biochemical characterization

[156939]of genetically variant aromatic amine N-acetyltransferases in A/J

[156940]and C57BL/6J mice. J. Biol. Chem. 257: 1431–1437, 1982.

[156941]

[156942]

[156943]15265.

[156944]Glowinski, I. B.; Weber, W. W.: Genetic regulation of aromatic

[156945]amine N-acetylation in inbred mice. J. Biol. Chem. 257:
1424-1430,

[156946]1982.

[156947]

[156948]

[156949]15266.

[156950]Grant, D. M.; Morike, K.; Eichelbaum, M.; Meyer, U. A.:
Acetylation

[156951]pharmacogenetics: the slow acetylator phenotype is
caused by decreased

[156952]or absent arylamine N-acetyltransferase in human liver. J.
Clin.

[156953]Invest. 85: 968-972, 1990.

[156954]

[156955]

[156956]15267.

[156957]Grant, D. M.; Tang, B. K.; Kalow, W.: A simple test for
acetylator

[156958]phenotype using caffeine. Brit. J. Clin. Pharm. 17:
459-464, 1984.

[156959]

[156960]

[156961]15268.

[156962]Hoo, J. J.; Hussein, L.; Goedde, H. W.: A simplified microchromethod

[156963]for the determination of the acetylator phenotype. J. Clin. Chem.

[156964]Clin. Biochem. 15: 329–331, 1977.

[156965]

[156966]

[156967]15269.

[156968]Ilett, K. F.; David, B. M.; Detchon, P.; Castleden, W. M.; Kwa,

[156969]R.: Acetylation phenotype in colorectal carcinoma. Cancer Res. 47:

[156970]1466–1469, 1987.

[156971]

[156972]

[156973]15270.

[156974]Iselius, L.; Evans, D. A. P.: Formal genetics of isoniazid metabolism

[156975]in man. Clin. Pharmacokinetics 8: 541–544, 1983.

[156976]

[156977]

[156978]15271.

[156979]Kukongviriyapan, V.; Lulitanond, V.; Areejitransorn, C.; Kongyingyose,

[156980]B.; Laupattarakasem, P.: N-acetyltransferase polymorphism in Thailand. Hum.

[156981]Hered. 34: 246–249, 1984.

[156982]

[156983]

[156984]15272.

[156985]Kutt, H.; Brennan, R.; Dehejia, H.; Verebely, K.: Diphenylhydantoin

[156986]intoxication: a complication of isoniazid therapy. Am. Rev. Resp.

[156987]Dis. 101: 377–384, 1970.

[156988]

[156989]

[156990]15273.

[156991]Lang, N. P.; Chu, D. Z. J.; Hunter C. F.; Kendall, D. C.;
Flammang,

[156992]T. J.; Kadlubar, F. F.: Role of aromatic amine acetyltrans-
ferase

[156993]in human colorectal cancer. Arch. Surg. 121: 1259–1261,
1986.

[156994]

[156995]

[156996]15274.

[156997]Lee, E. J. D.; Zhao, B.; Seow–Choen, F.: Relationship be-
tween

[156998]polymorphism of N–acetyltransferase gene and suscepti-
bility to colorectal

[156999]carcinoma in a Chinese population. Pharmacogenetics 8:
513–517,

[157000]1998.

[157001]

[157002]

[157003]15275.

[157004]Lin, H. J.; Han, C.-Y.; Lin, B. K.; Hardy, S.: Ethnic distribution

[157005]of slow acetylators mutations in the polymorphic N-acetyltransferase

[157006](NAT2) gene. Pharmacogenetics 4: 125-134, 1994.

[157007]

[157008]

[157009]15276.

[157010]Lin, H. J.; Han, C.-Y.; Lin, B. K.; Hardy, S.: Slow acetylators

[157011]mutations in the human polymorphic N-acetyltransferase gene in 786

[157012]Asians, blacks, Hispanics, and whites: application to metabolic epidemiology. Am.

[157013]J. Hum. Genet. 52: 827-834, 1993.

[157014]

[157015]

[157016]15277.

[157017]Mashimo, M.; Suzuki, T.; Abe, M.; Deguchi, T.: Molecular

genotyping

[157018]of N-acetylation polymorphism to predict phenotype.

Hum. Genet. 90:

[157019]139–143, 1992.

[157020]

[157021]

[157022]15278.

[157023]Mattano, S. S.; Weber, W. W.: Kinetics of arylamine N-acetyltransferase

[157024]in tissues from rapid and slow acetylators mice. Carcinogenesis 8:

[157025]133–137, 1987.

[157026]

[157027]

[157028]15279.

[157029]McLaren, E. H.; Burden, A. C.; Moorhead, P. J.: Acetylator phenotype

[157030]in diabetic neuropathy. Brit. Med. J. 2: 291–293, 1977.

[157031]

[157032]

[157033]15280.

[157034]McQueen, C. A.; Maslansky, C. J.; Glowinski, I. B.;
Crescenzi,

[157035]S. B.; Weber, W. W.; Williams, G. M.: Relationship between
the genetically

[157036]determined acetylator phenotype and DNA damage in-
duced by hydralazine

[157037]and 2-aminofluorene in cultured rabbit hepatocytes. Proc.
Nat. Acad.

[157038]Sci. 79: 1269-1272, 1982.

[157039]

[157040]

[157041]15281.

[157042]Giralt, M.; Park, E. A.; Gurney, A. L.; Liu, J.; Hakimi, P.;
Hanson,

[157043]R. W.: Identification of a thyroid hormone response ele-
ment in the

[157044]phosphoenolpyruvate carboxykinase (GTP) gene: evidence
for synergistic

[157045]interaction between thyroid hormone and cAMP cis-

regulatory elements. J.

[157046]Biol. Chem. 266: 21991–21996, 1991.

[157047]

[157048]

[157049]15282.

[157050]Pilz, A. J.; Willer, E.; Povey, S.; Abbott, C. M.: The genes
coding

[157051]for phosphoenolpyruvate carboxykinase–1 (PCK1) and
neuronal nicotinic

[157052]acetylcholine receptor alpha–4 subunit (CHRNA4) map to
human chromosome

[157053]20, extending the known region of homology with mouse
chromosome 2. Ann.

[157054]Hum. Genet. 56: 289–293, 1992.

[157055]

[157056]

[157057]15283.

[157058]Olswang, Y.; Cohen, H.; Papo, O.; Cassuto, H.; Croniger, C.
M.;

[157059]Hakimi, P.; Tilghman, S. M.; Hanson, R. W.; Reshef, L.: A

mutation

[157060]in the peroxisome proliferator-activated receptor gamma-binding site

[157061]in the gene for the cytosolic form of phosphoenolpyruvate carboxykinase

[157062]reduces adipose tissue size and fat content in mice. Proc. Nat. Acad.

[157063]Sci. 99: 625-630, 2002.

[157064]

[157065]

[157066]15284.

[157067]Stoffel, M.; Xiang, K.; Espinosa, R., III; Cox, N. J.; Le Beau,

[157068]M. M.; Bell, G. I.: cDNA sequence and localization of polymorphic

[157069]human cytosolic phosphoenolpyruvate carboxykinase gene (PCK1) to chromosome

[157070]20, band q13.31: PCK1 is not tightly linked to maturity-onset diabetes

[157071]of the young. Hum. Molec. Genet. 2: 1-4, 1993.

[157072]

[157073]

[157074]15285.

[157075]Ting, C.-N.; Burgess, D. L.; Chamberlain, J. S.; Keith, T. P.;

[157076]Falls, K.; Meisler, M. H.: Phosphoenolpyruvate carboxyki-
nase (GTP):

[157077]characterization of the human PCK1 gene and localization
distal to

[157078]MODY on chromosome 20. Genomics 16: 698–706, 1993.

[157079]

[157080]

[157081]15286.

[157082]Vidnes, J.; Sovik, O.: Gluconeogenesis in infancy and
childhood.

[157083]III. Deficiency of the extramitochondrial form of hepatic
phosphoenolpyruvate

[157084]carboxykinase in a case of persistent neonatal hypogly-
caemia. Acta

[157085]Paediat. Scand. 65: 301–312, 1976.

[157086]

[157087]

[157088]15287.

[157089]Yu, H.; Thun, R.; Chandrasekharappa, S.; Trent, J. M.;
Zhang, J.;

[157090]Meisler, M. H.: Human PCK1 encoding phosphoenolpyru-
vate carboxykinase

[157091]is located on chromosome 20q13.2. Genomics 15:
219–221, 1993.

[157092]

[157093]

[157094]15288.

[157095]Pepe, G.; Chimienti, G.; Resta, F.; Di Perna, V.; Tarricone,
C.;

[157096]Lovecchio, M.; Colacicco, A. M.; Capurso, A.: A new Italian
case

[157097]of lipoprotein lipase deficiency: a leu365-to-val change
resulting

[157098]in loss of enzyme activity. Biochem. Biophys. Res. Com-
mun. 199:

[157099]570–576, 1994.

[157100]

[157101]

[157102]15289.

[157103]Reymer, P. W. A.; Groenemeyer, B. E.; Gagne, E.; Miao, L.;
Appelman,

[157104]E. E. G.; Seidel, J. C.; Kromhout, D.; Bijvoet, S. M.; van de
Oever,

[157105]K.; Bruin, T.; Hayden, M. R.; Kastelein, J. J. P.: A frequently
occurring

[157106]mutation in the lipoprotein lipase gene (Asn291Ser) con-
tributes to

[157107]the expression of familial combined hyperlipidemia. Hum.
Molec. Genet. 4:

[157108]1543–1549, 1995.

[157109]

[157110]

[157111]15290.

[157112]Samuels, M. E.; Forbey, K. C.; Reid, J. E.; Abkevich, V.;
Bulka,

[157113]K.; Wardell, B. R.; Bowen, B. R.; Hopkins, P. N.; Hunt, S. C.;
Ballinger,

[157114]D. G.; Skolnick, M. H.; Wagner, S.: Identification of a com-
mon variant

[157115]in the lipoprotein lipase gene in a large Utah kindred as-
certained

[157116]for coronary heart disease: the -93G/D9N variant predisposes to low

[157117]HDL-C/high triglycerides. Clin. Genet. 59: 88-98, 2001.

[157118]

[157119]

[157120]15291.

[157121]Schreibman, P. H.; Arons, D. L.; Saudek, C. D.; Arky, R. A.:

[157122]Abnormal lipoprotein lipase in familial exogenous hypertriglyceridemia. J.

[157123]Clin. Invest. 52: 2074-2082, 1973.

[157124]

[157125]

[157126]15292.

[157127]Sprecher, D. L.; Kobayashi, J.; Rymaszewski, M.; Goldberg, I.

[157128]J.; Harris, B. V.; Bellet, P. S.; Ameis, D.; Yunker, R. L.; Black,

[157129]D. M.; Stein, E. A.; Schotz, M. C.; Wiginton, D. A.:

Trp64-to-nonsense

[157130]mutation in the lipoprotein lipase gene. J. Lipid Res. 33:

859-866,

[157131]1992.

[157132]

[157133]

[157134]15293.

[157135]Sternowsky, H. J.; Gaertner, U.; Stahnkel, N.; Kaukel, E.:
Juvenile

[157136]familial hypertriglyceridemia and growth retardation: clinical and

[157137]biochemical observations in three siblings. Europ. J. Pediatrics. 125:

[157138]59-70, 1977.

[157139]

[157140]

[157141]15294.

[157142]Takagi, A.; Ikeda, Y.; Tsutsumi, Z.; Shoji, T.; Yamamoto, A.:

[157143]Molecular studies on primary lipoprotein lipase (LPL) deficiency:

[157144]one base deletion (G-916) in exon 5 of LPL gene causes

no detectable

[157145]LPL protein due to the absence of LPL mRNA transcript. J.

Clin. Invest. 89:

[157146]581–591, 1992.

[157147]

[157148]

[157149]15295.

[157150]Weinstock, P. H.; Bisgaier, C. L.; Aalto-Setälä, K.; Radner, H.;

[157151]Ramakrishnan, R.; Levak-Frank, S.; Essenburg, A. D.; Zechner, R.;

[157152]Breslow, J. L.: Severe hypertriglyceridemia, reduced high density

[157153]lipoprotein, and neonatal death in lipoprotein lipase knockout mice:

[157154]mild hypertriglyceridemia with impaired very low density lipoprotein

[157155]clearance in heterozygotes. J. Clin. Invest. 96: 2555–2568, 1995.

[157156]

[157157]

[157158]15296.

[157159]Wessler, S.; Avioli, L. A.: Familial hyperlipoproteinemia.

J.A.M.A. 207:

[157160]929–937, 1969.

[157161]

[157162]

[157163]15297.

[157164]Wilson, D. E.; Edwards, C. Q.; Chan, I.–F.: Phenotypic heterogeneity

[157165]in the extended pedigree of a proband with lipoprotein lipase deficiency. Metabolism 32:

[157166]1107–1114, 1983.

[157167]

[157168]

[157169]15298.

[157170]Wilson, D. E.; Emi, M.; Iverius, P.–H.; Hata, A.; Wu, L. L.; Hillas,

[157171]E.; Williams, R. R.; Lalouel, J.–M.: Phenotypic expression of heterozygous

[157172]lipoprotein lipase deficiency in the extended pedigree of a

proband

[157173]homozygous for a missense mutation. J. Clin. Invest. 86:
735–750,

[157174]1990.

[157175]

[157176]

[157177]15299.

[157178]Wilson, D. E.; Hata, A.; Kwong, L. K.; Lingam, A.; Shuhua,
J.;

[157179]Ridinger, D. N.; Yeager, C.; Kaltenborn, K. C.; Iverius,
P.–H.; Lalouel,

[157180]J.–M.: Mutations in exon 3 of the lipoprotein lipase gene
segregating

[157181]in a family with hypertriglyceridemia, pancreatitis, and
non–insulin–dependent

[157182]diabetes. J. Clin. Invest. 92: 203–211, 1993.

[157183]

[157184]

[157185]15300.

[157186]Wion, K. L.; Kirchgessner, T. G.; Lusi, A. J.; Schotz, M. C.;

[157187]Lawn, R. M.: Human lipoprotein lipase complementary
DNA sequence. Science 235:

[157188]1638–1641, 1987.

[157189]

[157190]

[157191]15301.

[157192]Wittrup, H. H.; Tybjaerg-Hansen, A.; Abildgaard, S.; Stef-
fensen,

[157193]R.; Schnohr, P.; Nordestgaard, B. G.: A common substitu-
tion (asn291ser)

[157194]in lipoprotein lipase is associated with increased risk of
ischemic

[157195]heart disease. J. Clin. Invest. 99: 1606–1613, 1997.

[157196]

[157197]

[157198]15302.

[157199]Wittrup, H. H.; Tybjaerg-Hansen, A.; Steffensen, R.; Deeb,
S.

[157200]S.; Brunzell, J. D.; Jensen, G.; Nordestgaard, B. G.: Muta-
tions in

[157201]the lipoprotein lipase gene associated with ischemic heart disease

[157202]in men: the Copenhagen City Heart Study. Arterioscler. Thromb. Vasc.

[157203]Biol. 19: 1535–1540, 1999.

[157204]

[157205]

[157206]15303.

[157207]Wood, S.; Schertzer, M.; Hayden, M.; Ma, Y.: Support for founder

[157208]effect for two lipoprotein lipase (LPL) gene mutations in French Canadians

[157209]by analysis of GT microsatellites flanking the LPL gene. Hum. Genet. 91:

[157210]312–316, 1993.

[157211]

[157212]

[157213]15304.

[157214]McQueen, C. A.; Weber, W. W.: Characterization of human lymphocyte

[157215]N-acetyltransferase and its relationship to the isoniazid
acetylator

[157216]polymorphism. Biochem. Genet. 18: 889–904, 1980.

[157217]

[157218]

[157219]15305.

[157220]Nhachi, C. F. B.: Polymorphic acetylation of sulphamet-
hazine

[157221]in a Zimbabwe population. J. Med. Genet. 25: 29–31,
1988.

[157222]

[157223]

[157224]15306.

[157225]Ohsako, S.; Deguchi, T.: Cloning and expression of cDNAs
for

[157226]polymorphic and monomorphic arylamine N-
acetyltransferases from human

[157227]liver. J. Biol. Chem. 265: 4630–4634, 1990.

[157228]

[157229]

[157230]15307.

[157231]Parker, J. M.: Human variability in the metabolism of sul-
famethazine. Hum.

[157232]Hered. 19: 402–409, 1969.

[157233]

[157234]

[157235]15308.

[157236]Penketh, R. J. A.; Gibney, S. F. A.; Nurse, G. T.; Hopkinson,

[157237]D. A.: Acetylator phenotypes in Papua New Guinea. J. Med.
Genet. 20:

[157238]37–40, 1983.

[157239]

[157240]

[157241]15309.

[157242]Roberts–Thomson, I. C.; Ryan, P.; Khoo, K. K.; Hart, W. J.;
McMichael,

[157243]A. J.; Butler, R. N.: Diet, acetylator phenotype, and risk of
colorectal

[157244]neoplasia. Lancet 347: 1372–1374, 1996.

[157245]

[157246]

[157247]15310.

[157248]Roots, I.; Drakoulis, N.; Brockmoller, J.; Janicke, I.;
Cuprunov,

[157249]M.; Ritter, J.: Hydroxylation and acetylation phenotypes as
genetic

[157250]risk factors in certain malignancies.In: Kato, W.; Es-
tabrook, R. W.;

[157251]Cayen, M. N. (eds.): Xenobiotic Metabolism and Disposi-
tion. London:

[157252]Taylor and Francis 1989. Pp. 499–506.

[157253]

[157254]

[157255]15311.

[157256]Schloot, W.; Goedde, H. W.: Studies on the polymorphism
of isoniazid

[157257](INH) acetylation in rhesus monkeys (*Macaca mulatta*).
Acta Genet.

[157258]Statist. Med. 18: 394–398, 1968.

[157259]

[157260]

[157261]15312.

[157262]Schnakenberg, E.; Ehlers, C.; Feyerabend, W.; Werdin, R.;
Hubotter,

[157263]R.; Dreikorn, K.; Schloot, W.: Genotyping of the polymor-
phic N-acetyltransferase

[157264](NAT2) and loss of heterozygosity in bladder cancer pa-
tients. Clin.

[157265]Genet. 53: 396–402, 1998.

[157266]

[157267]

[157268]15313.

[157269]Sonnhaag, C.; Karlsson, E.; Hed, J.: Procainamide-induced
lupus

[157270]erythematosus-like syndrome in relation to acetylator
phenotype and

[157271]plasma levels of procainamide. Acta Med. Scand. 206:
245–251, 1979.

[157272]

[157273]

[157274]15314.

[157275]Sunahara, S.; Urano, M.; Ogawa, M.: Genetical and geographic

[157276]studies on isoniazid inactivation. Science 134:
1530–1531, 1961.

[157277]

[157278]

[157279]15315.

[157280]Tannen, R. H.; Weber, W. W.: Inheritance of acetylator
phenotype

[157281]in mice. J. Pharm. Exp. Therap. 213: 480–484, 1980.

[157282]

[157283]

[157284]15316.

[157285]Timbrell, J. A.; Wright, J. M.; Baillie, T. A.: Monoacetylhydrazine

[157286]as a metabolite of isoniazid in man. Clin. Pharm. Therap.
22: 602–609,

[157287]1977.

[157288]

[157289]

[157290]15317.

[157291]Vansant, J.; Woosley, R. L.; John, J. T.; Sergent, J. S.: Normal
mal

[157292]distribution of acetylation phenotypes in systemic lupus
erythematosus. Arthritis

[157293]Rheum. 21: 192–195, 1978.

[157294]

[157295]

[157296]15318.

[157297]Weber, W. W.: The Acetylator Genes and Drug Response.
New York:

[157298]Oxford Univ. Press (pub.) 1987.

[157299]

[157300]

[157301]15319.

[157302]Woosley, R. L.; Drayer, D. E.; Reidenberg, M. M.; Nies, A.
S.;

[157303]Carr, K.; Oates, J. A.: Effect of acetylator phenotype on the
rate

[157304]at which procainamide induces antinuclear antibodies and
the lupus

[157305]syndrome. New Eng. J. Med. 298: 1157–1159, 1978.

[157306]

[157307]

[157308]15320.

[157309]Pace, P.; Johnson, M.; Tan, W. M.; Mosedale, G.; Sng, C.;
Hoatlin,

[157310]M.; de Winter, J.; Joenje, H.; Gergely, F.; Patel, K. J.: FANCE:

[157311]the link between Fanconi anaemia complex assembly and
activity. EMBO

[157312]J. 21: 3414–3423, 2002.

[157313]

[157314]

[157315]15321.

[157316]Hodge, S. E.; Anderson, C. E.; Neiswanger, K.; Field, L. L.;
Spence,

[157317]M. A.; Sparkes, R. S.; Sparkes, M. C.; Crist, M.; Terasaki, P.

I.;

[157318]Rimoin, D. L.; Rotter, J. I.: Close genetic linkage between diabetes

[157319]mellitus and Kidd blood group. Lancet II: 893–895, 1981.

[157320]

[157321]

[157322]15322.

[157323]Faiyaz–Ul–Haque, M.; Ahmad, W.; Zaidi, S. H. E.; Haque, S.; Teebi,

[157324]A. S.; Ahmad, M.; Cohn, D. H.; Tsui, L. C.: Mutation in the cartilage–derived

[157325]morphogenetic protein–1 (CDMP1) gene in a kindred affected with fibular

[157326]hypoplasia and complex brachydactyly (DuPan syndrome). Clin. Genet. 61:

[157327]454–458, 2002.

[157328]

[157329]

[157330]15323.

[157331]Touchman, J. W.; Anikster, Y.; Dietrich, N. L.; Braden

Maduro,

[157332]V. V.; McDowell, G.; Shotelersuk, V.; Bouffard, G. G.; Beckstrom-Sternberg,

[157333]S. M.; Gahl, W. A.; Green, E. D.: The genomic region encompassing

[157334]the nephropathic cystinosis gene (CTNS): complete sequencing of a

[157335]200-kb segment and discovery of a novel gene within the common cystinosis-causing

[157336]deletion. *Genome Res.* 10: 165–173, 2000.

[157337]

[157338]

[157339]15324.

[157340]Shapiro, M. B.; Senapathy, P.: RNA splice junctions of different

[157341]classes of eukaryotes: sequence statistics and functional implications

[157342]in gene expression. *Nucleic Acids Res.* 15: 7155–7174, 1987.

[157343]

[157344]

[157345]15325.

[157346]Jobard, F.; Lefevre, C.; Karaduman, A.; Blanchet-Bardon, C.; Emre,

[157347]S.; Weissenbach, J.; Ozguc, M.; Lathrop, M.; Prud'homme, J.-F.; Fischer,

[157348]J.: Lipoxxygenase-3 (ALOXE3) and 12(R)-lipoxxygenase (ALOX12B) are

[157349]mutated in non-bullous congenital ichthyosiform erythroderma (NCIE)

[157350]linked to chromosome 17p13.1. Hum. Molec. Genet. 11: 107-113, 2002.

[157351]

[157352]

[157353]15326.

[157354]Varon, R.; Vissinga, C.; Platzer, M.; Cerosaletti, K. M.; Chrzanowska,

[157355]K. H.; Saar, K.; Beckmann, G.; Seemanova, E.; Cooper, P. R.; Nowak,

[157356]N. J.; Stumm, M.; Weemaes, C. M. R.; Gatti, R. A.; Wilson, R. K.;

[157357]Digweed, M.; Rosenthal, A.; Sperling, K.; Concannon, P.; Reis, A.

[157358]: Nibrin, a novel DNA double-strand break repair protein,
is mutated

[157359]in Nijmegen breakage syndrome. Cell 93: 467-476, 1998.

[157360]

[157361]

[157362]15327.

[157363]Cheung, P. P.; Cannizzaro, L. A.; Colman, R. W.: Chromo-
somal mapping

[157364]of human kininogen gene (KNG) to 3q26-qter. Cytogenet.
Cell Genet. 59:

[157365]24-26, 1992.

[157366]

[157367]

[157368]15328.

[157369]Cheung, P. P.; Kunapuli, S. P.; Scott, C. F.; Wachtfogel, Y.
T.;

[157370]Colman, R. W.: Genetic basis of total kininogen deficiency
in Williams'

[157371]trait. J. Biol. Chem. 268: 23361-23365, 1993.

[157372]

[157373]

[157374]15329.

[157375]Colman, R. W.; Bagdasarian, A.; Talamo, R. C.; Scott, C. F.;
Seavey,

[157376]M.; Guimaraes, J. A.; Pierce, J. V.; Kaplan, A. P.: Williams
trait:

[157377]human kininogen deficiency with diminished levels of
plasminogen proactivator

[157378]and prekallikrein associated with abnormalities of the
Hageman factor

[157379]dependent pathways. J. Clin. Invest. 56: 1650–1662,
1975.

[157380]

[157381]

[157382]15330.

[157383]Davie, E. W.: Personal Communication. Seattle, Wash.
11/26/1979.

[157384]

[157385]

[157386]15331.

[157387]Donaldson, V. H.; Kleniewski, J.; Saito, H.; Sayed, J. K.:

Prekallikrein

[157388]deficiency in a kindred with kininogen deficiency and

Fitzgerald trait

[157389]clotting defect: evidence that high molecular weight

kininogen and

[157390]prekallikrein exist as a complex in normal plasma. J. Clin.

Invest. 60:

[157391]571–583, 1977.

[157392]

[157393]

[157394]15332.

[157395]Fong, D.; Smith, D. I.; Hsieh, W.–T.: The human kininogen

gene

[157396](KNG) mapped to chromosome 3q26–qter by analysis of

somatic cell hybrids

[157397]using the polymerase chain reaction. Hum. Genet. 87:

189–192, 1991.

[157398]

[157399]

[157400]15333.

[157401]Hayashi, H.; Ishimaru, F.; Fujita, T.; Tsurumi, N.; Tsuda, T.;

[157402]Kimura, I.: Molecular genetic survey of five Japanese families with

[157403]high molecular weight kininogen deficiency. Blood 75: 1296–1304,

[157404]1990.

[157405]

[157406]

[157407]15334.

[157408]James, F. W.; Donaldson, V. H.: Decreased exercise tolerance and

[157409]hypertension in severe hereditary deficiency of plasma kininogens.

[157410](Letter) Lancet I: 889 only, 1981.

[157411]

[157412]

[157413]15335.

[157414]Kunapuli, S. P.: Personal Communication. Philadelphia, Pa.

[157415]12/21/1993.

[157416]

[157417]

[157418]15336.

[157419]Lacombe, M.-J.; Varet, B.; Levy, J.-P.: A hitherto undescribed

[157420]plasma factor acting at the contact phase of blood coagulation (Flaujeac

[157421]factor): case report and coagulation studies. Blood 46: 761-768,

[157422]1975.

[157423]

[157424]

[157425]15337.

[157426]Lefrere, J.-J.; Horellou, M.-H.; Gozin, D.; Conard, J.; Muller,

[157427]J.-Y.; Clark, M.; Soulier, J.-P.; Samama, M.: A new case of high

[157428]molecular weight kininogen inherited deficiency. Am. J. Hemat. 22:

[157429]415-419, 1986.

[157430]

[157431]

[157432]15338.

[157433]Saito, H.; Ratnoff, O. D.; Waldmann, R.; Abraham, J. P.:

Deficiency

[157434]of a hitherto unrecognized agent, Fitzgerald factor, participating

[157435]in surface-mediated reactions of clotting, fibrinolysis, generation

[157436]of kinins, and the property of diluted plasma enhancing vascular permeability

[157437](Pf-DIL). J. Clin. Invest. 55: 1082–1089, 1975.

[157438]

[157439]

[157440]15339.

[157441]Waldmann, R.; Abraham, J. P.; Rebuck, J. W.; Caldwell, J.;

Saito,

[157442]H.; Ratnoff, O. D.: Fitzgerald factor: a hitherto unrecognized coagulation

[157443]factor. Lancet I: 949–951, 1975.

[157444]

[157445]

[157446]15340.

[157447]Wuepper, K. D.; Miller, D. R.; Lacombe, M. J.: Flaujeac trait:

[157448]deficiency of human plasma kininogen. J. Clin. Invest. 56:

1663–1672,

[157449]1975.

[157450]

[157451]

[157452]15341.

[157453]ten Hove, T.; Corbaz, A.; Amitai, H.; Aloni, S.; Belzer, I.;

Graber,

[157454]P.; Drillenburger, P.; van Deventer, S. J.; Chvatchko, Y.; Te

Velde,

[157455]A. A.: Blockade of endogenous IL–18 ameliorates TNBS–

induced colitis

[157456]by decreasing local TNF–alpha production in mice. Gas-

troenterology 121:

[157457]1372–1379, 2001.

[157458]

[157459]

[157460]15342.

[157461]Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.;
Tanaka, A.;

[157462]Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding
sequences

[157463]of unidentified human genes. X. The complete sequences
of 100 new

[157464]cDNA clones from brain which can code for large proteins
in vitro. DNA

[157465]Res. 5: 169–176, 1998.

[157466]

[157467]

[157468]15343.

[157469]Kipreos, E. T.; Lander, L. E.; Wing, J. P.; He, W. W.; Hedge-
cock,

[157470]E. M.: cul-1 is required for cell cycle exit in *C. elegans* and
identifies

[157471]a novel gene family. Cell 85: 829–839, 1996.

[157472]

[157473]

[157474]15344.

[157475]Rasooly, R. S.: Personal Communication. Baltimore, Md.
9/29/1998.

[157476]

[157477]

[157478]15345.

[157479]Wang, P. J.; McCarrey, J. R.; Yang, F.; Page, D. C.: An abundance
dance

[157480]of X-linked genes expressed in spermatogonia. Nature
Genet. 27:

[157481]422-426, 2001.

[157482]

[157483]

[157484]15346.

[157485]Herold, A.; Suyama, M.; Rodrigues, J. P.; Braun, I. C.; Ku-
tay,

[157486]U.; Carmo-Fonseca, M.; Bork, P.; Izaurralde, E.: TAP (NXF1)
belongs

[157487]to a multigene family of putative RNA export factors with

a conserved

[157488]modular architecture. *Molec. Cell. Biol.* 20: 8996–9008, 2000.

[157489]

[157490]

[157491]15347.

[157492]Yang, J.; Bogerd, H. P.; Wang, P. J.; Page, D. C.; Cullen, B. R.

[157493]: Two closely related human nuclear export factors utilize entirely

[157494]distinct export pathways. *Molec. Cell* 8: 397–406, 2001.

[157495]

[157496]

[157497]15348.

[157498]Ikeda, M.; Ishida, O.; Hinoi, T.; Kishida, S.; Kikuchi, A.: Identification

[157499]and characterization of a novel protein interacting with Ral-binding

[157500]protein 1, a putative effector protein of Ral. *J. Biol. Chem.* 273:

[157501]814–821, 1998.

[157502]

[157503]

[157504]15349.

[157505]Koshiba, S.; Kigawa, T.; Iwahara, J.; Kikuchi, A.; Yokoyama, S.

[157506]: Solution structure of the Eps15 homology domain of a human POB1

[157507](partner of RalBP1). FEBS Lett. 442: 138–142, 1999.

[157508]

[157509]

[157510]15350.

[157511]Crackower, M. A.; Sarao, R.; Oudit, G. Y.; Yagil, C.; Kozier-
adzki,

[157512]I.; Scanga, S. E.; Oliveira-dos-Santos, A. J.; da Costa, J.;
Zhang,

[157513]L.; Pei, Y.; Scholey, J.; Ferrario, C. M.; Manoukian, A. S.;
Chappell,

[157514]M. C.; Backx, P. H.; Yagil, Y.; Penninger, J. M.: An-
giotensin-converting

[157515]enzyme 2 is an essential regulator of heart function. Na-
ture 417:

[157516]822–828, 2002.

[157517]

[157518]

[157519]15351.

[157520]Scanlan, M. J.; Gordan, J. D.; Williamson, B.; Stockert, E.;
Bander,

[157521]N. H.; Jongeneel, V.; Gure, A. O.; Jager, D.; Jager, E.;
Knuth, A.;

[157522]Chen, Y.–T.; Old, L. J.: Antigens recognized by autologous
antibody

[157523]in patients with renal–cell carcinoma. Int. J. Cancer 83:
456–464,

[157524]1999.

[157525]

[157526]

[157527]15352.

[157528]Yan, Z.; Fedorov, S. A.; Mumby, M. C.; Williams, R. S.:
PR48,

[157529]a novel regulatory subunit of protein phosphatase 2A, in–
teracts with

[157530]Cdc6 and modulates DNA replication in human cells.

Molec. Cell. Biol. 20:

[157531]1021–1029, 2000.

[157532]

[157533]

[157534]15353.

[157535]Gitzelmann, R.: Hereditary galactokinase deficiency, a
newly recognized

[157536]cause of juvenile cataracts. Pediat. Res. 1: 14–23, 1967.

[157537]

[157538]

[157539]15354.

[157540]Kalaydjieva, L.; Perez–Lezaun, A.; Angelicheva, D.;
Onengut, S.;

[157541]Dye, D.; Bosshard, N. U.; Jordanova, A.; Savov, A.;
Yanakiev, P.;

[157542]Kremensky, I.; Radeva, B.; Hallmayer, J.; Markov, A.; Ned–
kova, V.;

[157543]Tournev, I.; Aneva, L.; Gitzelmann, R.: A founder mutation
in the

[157544]GK1 gene is responsible for galactokinase deficiency in
Roma (Gypsies). Am.

[157545]J. Hum. Genet. 65: 1299–1307, 1999.

[157546]

[157547]

[157548]15355.

[157549]Stratakis, C. A.; Lin, J.–P.; Pras, E.; Rennert, O. M.; Bour-
dony,

[157550]C. J.; Chan, W.–Y.: Segregation of Allgrove (triple–A) syn-
drome in

[157551]Puerto Rican kindreds with chromosome 12 (12q13) poly-
morphic markers. Proc.

[157552]Assoc. Am. Phys. 109: 478–482, 1997.

[157553]

[157554]

[157555]15356.

[157556]Tullio–Pelet, A.; Salomon, R.; Hadj–Rabia, S.; Mugnier, C.;
de

[157557]Laet, M.–H.; Chaouachi, B.; Bakiri, F.; Brottier, P.; Cattolico,
L.;

[157558]Penet, C.; Begeot, M.; Naville, D.; Nicolino, M.; Chaussain, J.-L.;

[157559]Weissenbach, J.; Munnich, A.; Lyonnet, S.: Mutant WD-repeat protein

[157560]in triple-A syndrome. Nature Genet. 26: 332-335, 2000.

[157561]

[157562]

[157563]15357.

[157564]Camacho, J. A.; Obie, C.; Biery, B.; Goodman, B. K.; Hu, C.-A.;

[157565]Almashanu, S.; Steel, G.; Casey, R.; Lambert, M.; Mitchell, G. A.;

[157566]Valle, D.: Hyperornithinaemia-hyperammon-aemia-homocitrullinuria syndrome

[157567]is caused by mutations in a gene encoding a mitochondrial ornithine

[157568]transporter. Nature Genet. 22: 151-158, 1999.

[157569]

[157570]

[157571]15358.

[157572]Darnell, J. E., Jr.: Reflections on STAT3, STAT5, and STAT6
as

[157573]fat STATs. Proc. Nat. Acad. Sci. 93: 6221–6224, 1996.

[157574]

[157575]

[157576]15359.

[157577]Ghilardi, N.; Ziegler, S.; Wiestner, A.; Stoffel, R.; Heim, M.

[157578]H.; Skoda, R. C.: Defective STAT signaling by the leptin re-
ceptor

[157579]in diabetic mice. Proc. Nat. Acad. Sci. 93: 6231–6235,
1996.

[157580]

[157581]

[157582]15360.

[157583]Vaisse, C.; Halaas, J. L.; Horvath, C. M.; Darnell, J. E., Jr.;

[157584]Stoffel, M.; Friedman, J. M.: Leptin activation of Stat3 in
the hypothalamus

[157585]of wildtype and ob/ob mice but not in db/db mice. Nature
Genet. 14:

[157586]95–100, 1996.

[157587]

[157588]

[157589]15361.

[157590]Mohandas, T. K.; Chen, X.-N.; Rowe, L. B.; Birkenmeier, E. H.;

[157591]Fanning, A. S.; Anderson, J. M.; Korenberg, J. R.: Localiza-
tion of

[157592]the tight junction protein gene TJP1 to human chromo-
some 15q13, distal

[157593]to the Prader-Willi/Angelman region, and to mouse chro-
mosome 7. Genomics 30:

[157594]594-597, 1995.

[157595]

[157596]

[157597]15362.

[157598]Willott, E.; Balda, M. S.; Fanning, A. S.; Jameson, B.; Van
Itallie,

[157599]C.; Anderson, J. M.: The tight junction protein ZO-1 is ho-
mologous

[157600]to the Drosophila discs-large tumor suppressor protein of
septate

[157601]junctions. Proc. Nat. Acad. Sci. 90: 7834–7838, 1993.

[157602]

[157603]

[157604]15363.

[157605]Blancar, M. A.; Crossley, P. H.; Peters, K. G.; Steingrimsson, E.;

[157606]Copeland, N. G.; Jenkins, N. A.; Martin, G. R.; Rutter, W. J.:
Meso1,

[157607]a basic helix–loop–helix protein involved in mammalian
presomitic

[157608]mesoderm development. Proc. Nat. Acad. Sci. 92:
5870–5874, 1995.

[157609]

[157610]

[157611]15364.

[157612]Burgess, R.; Rawls, A.; Brown, D.; Bradley, A.; Olson, E. N.:

[157613]Requirement of the paraxis gene for somite formation and
musculoskeletal

[157614]patterning. Nature 384: 570–573, 1996.

[157615]

[157616]

[157617]15365.

[157618]Hidai, H.; Quertermous, E. E.; Espinosa, R., III; Le Beau, M.
M.;

[157619]Quertermous, T.: Genomic organization and chromosomal
localization

[157620]of the gene TCF15 encoding the early mesodermal basic
helix-loop-helix

[157621]factor bHLH-EC2. Genomics 30: 598-601, 1995.

[157622]

[157623]

[157624]15366.

[157625]Quertermous, E. E.; Hidai, H.; Blonar, M. A.; Quertermous,
T.:

[157626]Cloning and characterization of a basic helix-loop-helix
protein expressed

[157627]in early mesoderm and the developing somites. Proc. Nat.
Acad. Sci. 91:

[157628]7066-7070, 1994.

[157629]

[157630]

[157631]15367.

[157632]Quertermous, T.: Personal Communication. Nashville,
Tenn. 1/22/1996.

[157633]

[157634]

[157635]15368.

[157636]Clevidence, D. E.; Overdier, D. G.; Peterson, R. S.; Porcella,

[157637]A.; Ye, H.; Paulson, K. E.; Costa, R. H.: Members of the
HNF-3/forkhead

[157638]family of transcription factors exhibit distinct cellular ex-
pression

[157639]patterns in lung and regulate the surfactant protein B pro-
moter. Dev.

[157640]Biol. 166: 195-209, 1994.

[157641]

[157642]

[157643]15369.

[157644]Hellqvist, M.; Mahlapuu, M.; Blixt, A.; Enerback, S.; Carls-

son,

[157645]P.: The human forkhead protein FREAC-2 contains two functionally

[157646]redundant activation domains and interacts with TBP and TFIIB. J.

[157647]Biol. Chem. 273: 23335-23343, 1998.

[157648]

[157649]

[157650]15370.

[157651]Hellqvist, M.; Mahlapuu, M.; Samuelsson, L.; Enerback, S.; Carlsson,

[157652]P.: Differential activation of lung-specific genes by two forkhead

[157653]proteins, FREAC-1 and FREAC-2. J. Biol. Chem. 271: 4482-4490, 1996.

[157654]

[157655]

[157656]15371.

[157657]Larsson, C.; Hellqvist, M.; Pierrou, S.; White, I.; Enerback, S.;

[157658]Carlsson, P.: Chromosomal localization of six human

forkhead genes,

[157659]frec-1 (FKHL5), -3 (FKHL7), -4 (FKHL8), -5 (FKHL9), -6 (FKHL10),

[157660]and -8 (FKHL12). Genomics 30: 464-469, 1995.

[157661]

[157662]

[157663]15372.

[157664]Pierrou, S.; Hellqvist, M.; Samuelsson, L.; Enerback, S.; Carlsson,

[157665]P.: Cloning and characterization of seven human forkhead proteins:

[157666]binding site specificity and DNA bending. EMBO J. 13: 5002-5012,

[157667]1994.

[157668]

[157669]

[157670]15373.

[157671]Ernstsson, S.; Pierrou, S.; Hulander, M.; Cederberg, A.; Hellqvist,

[157672]M.; Carlsson, P.; Enerback, S.: Characterization of the hu-

man forkhead

[157673]gene FREAC-4. J. Biol. Chem. 271: 21094-21099, 1996.

[157674]

[157675]

[157676]15374.

[157677]Hulander, M.; Wurst, W.; Carlsson, P.; Enerback, S.: The
winged

[157678]helix transcription factor Fkh10 is required for normal de-
velopment

[157679]of the inner ear. Nature Genet. 20: 374-376, 1998.

[157680]

[157681]

[157682]15375.

[157683]De Smaele, E.; Zazzeroni, F.; Papa, S.; Nguyen, D. U.; Jin,
R.;

[157684]Jones, J.; Cong, R.; Franzoso, G.: Induction of
gadd45-beta by NF-kappa-B

[157685]downregulates pro-apoptotic JNK signaling. Nature 414:
308-313,

[157686]2001.

[157687]

[157688]

[157689]15376.

[157690]Dong, C.; Yang, D. D.; Tournier, C.; Whitmarsh, A. J.; Xu, J.;

[157691]Davis, R. J.; Flavell, R. A.: JNK is required for effector T-cell

[157692]function but not for T-cell activation. Nature 405: 91–94, 2000.

[157693]

[157694]

[157695]15377.

[157696]Gupta, S.; Barrett, T.; Whitmarsh, A. J.; Cavanagh, J.; Sluss,

[157697]H. K.; Derijard, B.; Davis, R. J.: Selective interaction of JNK protein

[157698]kinase isoforms with transcription factors. EMBO J. 15: 2760–2770,

[157699]1996.

[157700]

[157701]

[157702]15378.

[157703]Tournier, C.; Hess, P.; Yang, D. D.; Xu, J.; Turner, T. K.;
Nimnual,

[157704]A.; Bar-Sagi, D.; Jones, S. N.; Flavell, R. A.; Davis, R. J.: Re-
quirement

[157705]of JNK for stress-induced activation of the cytochrome c-
mediated

[157706]death pathway. Science 288: 870–874, 2000.

[157707]

[157708]

[157709]15379.

[157710]Daugherty, B. L.; Springer, M. S.: The beta-chemokine re-
ceptor

[157711]genes CCR1 (CMKBR1), CCR2 (CMKBR2), and CCR3
(CMKBR3) cluster within

[157712]285 kb on human chromosome 3p21. Genomics 41:
294–295, 1997.

[157713]

[157714]

[157715]15380.

[157716]Ayyanathan, K.; Naylor, S. L.; Kunapuli, S. P.: Structural characterization

[157717]and fine chromosomal mapping of the human P2Y(1) purinergic receptor

[157718]gene (P2RY1). Somat. Cell Molec. Genet. 22: 419–424, 1996.

[157719]

[157720]

[157721]15381.

[157722]Ayyanathan, K.; Webbs, T. E.; Sandhu, A. K.; Athwal, R. S.; Barnard,

[157723]E. A.; Kunapuli, S. P.: Cloning and chromosomal localization of the

[157724]human P2Y1 purinoceptor. Biochem. Biophys. Res. Commun. 218: 783–788,

[157725]1996.

[157726]

[157727]

[157728]15382.

[157729]Leon, C.; Hechler, B.; Freund, M.; Eckly, A.; Vial, C.; Ohlmann,

[157730]P.; Dierich, A.; LeMeur, M.; Cazenave, J.-P.; Gachet, C.:

Defective

[157731]platelet aggregation and increased resistance to thrombo-
sis in purinergic

[157732]P2Y1 receptor-null mice. J. Clin. Invest. 104: 1731-1737,
1999.

[157733]

[157734]

[157735]15383.

[157736]Leon, C.; Hechler, B.; Vial, C.; Leray, C.; Cazenave, J.-P.;
Gachet,

[157737]C.: The P2Y(1) receptor is an ADP receptor antagonized by
ATP and

[157738]expressed in platelets and megakaryoblastic cells. FEBS
Letts. 403:

[157739]26-30, 1997.

[157740]

[157741]

[157742]15384.

[157743]Leon, C.; Vial, C.; Cazenave, J.-P.; Gachet, C.: Cloning and
sequencing

[157744]of a human cDNA encoding endothelial P2Y1 purinoceptor. Gene 171:

[157745]295–297, 1996.

[157746]

[157747]

[157748]15385.

[157749]Matsuda, C.; Hayashi, Y. K.; Ogawa, M.; Aoki, M.; Murayama, K.;

[157750]Nishino, I.; Nonaka, I.; Arahata, K.; Brown, R. H., Jr.: The sarcolemmal

[157751]proteins dysferlin and caveolin-3 interact in skeletal muscle. Hum.

[157752]Molec. Genet. 10: 1761–1766, 2001.

[157753]

[157754]

[157755]15386.

[157756]Furlong, R. A.; Zhou, C. Y.; Ferguson-Smith, M. A.; Affara, N.

[157757]A.: Characterization of a kinesin-related gene ATSV, within the tuberous

[157758]sclerosis locus (TSC1) candidate region on chromosome
9q34. Genomics 33:
[157759]421–429, 1996.

[157760]

[157761]

[157762]15387.

[157763]Keller, M. P.; Seifried, B. A.; Rabin, B. A.; Chance, P. F.:
Mapping

[157764]of the kinesin–related gene ATSV to chromosome 2q37.
Hum. Genet. 104:

[157765]254–256, 1999.

[157766]

[157767]

[157768]15388.

[157769]Kikkawa, M.; Okada, Y.; Hirokawa, N.: 15–angstrom reso-
lution model

[157770]of the monomeric kinesin motor, KIF1A. Cell 100:
241–252, 2000.

[157771]

[157772]

[157773]15389.

[157774]Kikkawa, M.; Sablin, E. P.; Okada, Y.; Yajima, H.; Fletterick,

[157775]R. J.; Hirokawa, N.: Switch-based mechanism of kinesin
motors. Nature 411:

[157776]439–445, 2001.

[157777]

[157778]

[157779]15390.

[157780]Klopfenstein, D. R.; Tomishige, M.; Stuurman, N.; Vale, R.
D.:

[157781]Role of phosphatidylinositol(4,5)bisphosphate organiza-
tion in membrane

[157782]transport by the Unc104 kinesin motor. Cell 109:
347–358, 2002.

[157783]

[157784]

[157785]15391.

[157786]Okada, Y.; Yamazaki, H.; Sekine-Aizawa, Y.; Hirokawa, N.:
The

[157787]neuron-specific kinesin superfamily protein KIF1A is a

unique monomeric

[157788]motor for anterograde axonal transport of synaptic vesicle precursors. Cell 81:

[157789]769–780, 1995.

[157790]

[157791]

[157792]15392.

[157793]Yonekawa, Y.; Harada, A.; Okada, Y.; Funakoshi, T.; Kanai, Y.;

[157794]Takei, Y.; Terada, S.; Noda, T.; Hirokawa, N.: Defect in synaptic

[157795]vesicle precursor transport and neuronal cell death in KIF1A motor

[157796]protein-deficient mice. J. Cell Biol. 141: 431–441, 1998.

[157797]

[157798]

[157799]15393.

[157800]Kikuta, Y.; Sogawa, K.; Haniu, M.; Kinoshita, M.; Kusunose, E.;

[157801]Nojima, Y.; Yamamoto, S.; Ichihara, K.; Kusunose, M.; Fu–

jii-Kuriyama,

[157802]Y.: A novel species of cytochrome P-450 (P-450ib) specific for the

[157803]small intestine of rabbits: cDNA cloning and its expression in COS

[157804]cells. J. Biol. Chem. 266: 17821-17825, 1991.

[157805]

[157806]

[157807]15394.

[157808]Ma, J.; Ramachandran, S.; Fiedorek, F. T., Jr.; Zeldin, D. C.:

[157809]Mapping of the CYP2J cytochrome P450 genes to human chromosome 1 and

[157810]mouse chromosome 4. Genomics 49: 152-155, 1998.

[157811]

[157812]

[157813]15395.

[157814]Wu, S.; Moomaw, C. R.; Tomer, K. B.; Falck, J. R.; Zeldin, D. C.

[157815]: Molecular cloning and expression of CYP2J2, a human cytochrome P450

[157816]arachidonic acid epoxxygenase highly expressed in heart. J. Biol.

[157817]Chem. 271: 3460–3468, 1996.

[157818]

[157819]

[157820]15396.

[157821]de Sauvage, F. J.; Camerato, T. R.; Goeddel, D. V.: Primary structure

[157822]and functional expression of the human receptor for Escherichia coli

[157823]heat-stable enterotoxin. J. Biol. Chem. 266: 17912–17918, 1991.

[157824]

[157825]

[157826]15397.

[157827]Mann, E. A.; Swenson, E. S.; Copeland, N. G.; Gilbert, D. J.; Jenkins,

[157828]N. A.; Taguchi, T.; Testa, J. R.; Giannella, R. A.: Localization

[157829]of the guanylyl cyclase C gene to mouse chromosome 6 and human chromosome

[157830]12p12. Genomics 34: 265–267, 1996.

[157831]

[157832]

[157833]15398.

[157834]Singh, S.; Singh, G.; Heim, J. M.; Gerzer, R.: Isolation and
expression

[157835]of a guanylate cyclase–coupled heat stable enterotoxin re-
ceptor cDNA

[157836]from a human colonic cell line. Biochem. Biophys. Res.
Commun. 179:

[157837]1455–1463, 1991.

[157838]

[157839]

[157840]15399.

[157841]Chiang, P.–W.; Wang, S.; Smithivas, P.; Song, W.–J.; Ra-
mamoorthy,

[157842]S.; Hillman, J.; Puett, S.; Van Keuren, M. L.; Crombez, E.;
Kumar,

[157843]A.; Glover, T. W.; Miller, D. E.; Tsai, C.–H.; Blackburn, C.
C.; Chen,

[157844]X.-N.; Sun, Z.; Cheng, J.-F.; Korenberg, J. R.; Kurnit, D. M.:
Identification
[157845]and analysis of the human and murine putative chromatin
structure
[157846]regulator SUPT6H and Supt6h. Genomics 34: 328–333,
1996.
[157847]
[157848]
[157849]15400.
[157850]Segre, J. A.; Nemhauser, J. L.; Taylor, B. A.; Nadeau, J. H.;
Lander,
[157851]E. S.: Positional cloning of the nude locus: genetic, physi-
cal and
[157852]transcription maps of the region and mutations in the
mouse and rat. Genomics 28:
[157853]549–559, 1995.
[157854]
[157855]
[157856]15401.
[157857]Han, J.; Knops, J. F.; Longshore, J. W.; King, P. H.: Local-
ization

[157858]of human elav-like neuronal protein 1 (Hel-N1) on chromosome 9p21

[157859]by chromosome microdissection polymerase chain reaction and fluorescence

[157860]in situ hybridization. Genomics 36: 189–191, 1996.

[157861]

[157862]

[157863]15402.

[157864]King, P. H.: Hel-N2: a novel isoform of Hel-N1 which is conserved

[157865]in rat neural tissue and produced in early embryogenesis.

Gene 151:

[157866]261–265, 1994.

[157867]

[157868]

[157869]15403.

[157870]King, P. H.; Levine, T. D.; Fremeau, R. T., Jr.; Keene, J. D.:

[157871]Mammalian homologs of Drosophila ELAV localized to a neuronal subset

[157872]can bind in vitro to the 3' UTR of mRNA encoding the Id

transcriptional

[157873]repressor. J. Neurosci. 14: 1943–52, 1994.

[157874]

[157875]

[157876]15404.

[157877]Abel, K. J.; Brody, L. C.; Valdes, J. M.; Erdos, M. R.; McKin-
ley,

[157878]D. R.; Castilla, L. H.; Merajver, S. D.; Couch, F. J.; Fried-
man, L.

[157879]S.; Ostermeyer, E. A.; Lynch, E. D.; King, M.-C.; Welcsh, P.
L.; Osborne-Lawrence,

[157880]S.; Spillman, M.; Bowcock, A. M.; Collins, F. S.; Weber, B.
L.: Characterization

[157881]of EZH1, a human homolog of Drosophila enhancer of
zeste near BRCA1. Genomics 37:

[157882]161–171, 1996.

[157883]

[157884]

[157885]15405.

[157886]Akazawa, K.; Yamane, S.; Shiota, H.; Naito, E.: A case of

retinoblastoma

[157887]associated with Rieger's anomaly and 13q deletion. Jpn. J.

Ophthal. 25:

[157888]321–325, 1981.

[157889]

[157890]

[157891]15406.

[157892]Stathacopoulos, R. A.; Bateman, J. B.; Sparkes, R. S.; Hep-
ler,

[157893]R. S.: The Rieger syndrome and a chromosome 13 dele-
tion. J. Pediat.

[157894]Ophthal. Strabismus 24: 198–203, 1987.

[157895]

[157896]

[157897]15407.

[157898]Ferbus, D.; Le Chalony, C.; Prosperi, M.–T.; Muleris, M.;
Vincent–Salomon,

[157899]A.; Goubin, G.: Identification, nuclear localization, and
binding

[157900]activities of OZF, a human protein solely composed of zinc

finger

[157901]motifs. Europ. J. Biochem. 236: 991–995, 1996.

[157902]

[157903]

[157904]15408.

[157905]Le Chalony, C.; Apiou, F.; Pibouin, L.; Dutrillaux, B.;
Goubin,

[157906]G.: Constitutive amplification of a zinc finger protein gene
in cattle. DNA

[157907]Cell Biol. 15: 83–88, 1996.

[157908]

[157909]

[157910]15409.

[157911]Le Chalony, C.; Prosperi, M.–T.; Haluza, R.; Apiou, F.;
Dutrillaux,

[157912]B.; Goubin, G.: The OZF gene encodes a protein consisting
essentially

[157913]of zinc–finger motifs. J. Molec. Biol. 236: 399–404, 1994.

[157914]

[157915]

[157916]15410.

[157917]Dell'Angelica, E. C.; Ohno, H.; Ooi, C. E.; Rabinovich, E.;
Roche,

[157918]K. W.; Bonifacino, J. S.: AP-3: an adaptor-like protein
complex with

[157919]ubiquitous expression. EMBO J. 16: 917-928, 1997.

[157920]

[157921]

[157922]15411.

[157923]Watanabe, T. K.; Shimizu, F.; Nagata, M.; Takaichi, A.; Fu-
jiwara,

[157924]T.; Nakamura, Y.; Takahashi, E.; Hirai, Y.: Cloning, expres-
sion pattern

[157925]and mapping to 12p13.2-p13.1 of CLAPS3, a gene encod-
ing a novel clathrin-adaptor

[157926]small chain. Cytogenet. Cell Genet. 73: 214-217, 1996.

[157927]

[157928]

[157929]15412.

[157930]Angrist, M.; Wells, D. E.; Chakravarti, A.; Pandey, A.: Chromosomal

[157931]localization of the mouse Src-like adapter protein (Slap) gene and

[157932]its putative human homolog SLA. Genomics 30: 623–625, 1995.

[157933]

[157934]

[157935]15413.

[157936]Holland, S. J.; Liao, X. C.; Mendenhall, M. K.; Zhou, X.; Pardo,

[157937]J.; Chu, P.; Spencer, C.; Fu, A.; Sheng, N.; Yu, P.; Pali, E.; Nagin,

[157938]A.; and 14 others: Functional cloning of Src-like adapter protein-2

[157939](SLAP-2), a novel inhibitor of antigen receptor signaling. J. Exp.

[157940]Med. 194: 1263–1276, 2001.

[157941]

[157942]

[157943]15414.

[157944]Kratchmarova, I.; Sosinowski, T.; Weiss, A.; Witter, K.; Vincenz,

[157945]C.; Pandey, A.: Characterization of promoter region and genomic structure

[157946]of the murine and human genes encoding Src like adapter protein. Gene 262:

[157947]267–273, 2001.

[157948]

[157949]

[157950]15415.

[157951]Meijerink, P. H. S.; Yanakiev, P.; Zorn, I.; Grierson, A. J.; Bikker,

[157952]H.; Dye, D.; Kalaydjieva, L.; Baas, F.: The gene for the human Src-like

[157953]adaptor protein (hSLAP) is located within the 64-kb intron of the

[157954]thyroglobulin gene. Europ. J. Biochem. 254: 297–303, 1998.

[157955]

[157956]

[157957]15416.

[157958]Sosinowski, T.; Pandey, A.; Dixit, V. M.; Weiss, A.: Src-like
[157959]adaptor protein (SLAP) is a negative regulator of T cell re-
ceptor

[157960]signaling. J. Exp. Med. 191: 463–474, 2000.

[157961]

[157962]

[157963]15417.

[157964]Bailey, S. M.; Cornforth, M. N.; Kurimasa, A.; Chen, D. J.;
Goodwin,

[157965]E. H.: Strand-specific postreplicative processing of mam-
malian telomeres. *Science* 293:

[157966]2462–2465, 2001.

[157967]

[157968]

[157969]15418.

[157970]Maniatis, T.: A ubiquitin ligase complex essential for the
NF- κ B,

[157971]Wnt/Wingless, and Hedgehog signaling pathways. *Genes*
Dev. 13: 505–510,

[157972]1999.

[157973]

[157974]

[157975]15419.

[157976]Winston, J. T.; Strack, P.; Beer-Romero, P.; Chu, C. Y.;
Elledge,

[157977]S. J.; Harper, J. W.: The SCF(β -TRCP)-ubiquitin ligase
complex

[157978]associates specifically with phosphorylated destruction motifs in

[157979]I-kappa-B-alpha and beta-catenin and stimulates I-kappa-B-alpha ubiquitination

[157980]in vitro. Genes Dev. 13: 270-283, 1999.

[157981]

[157982]

[157983]15420.

[157984]Ahouse, J. J.; Hagerman, C. L.; Mittal, P.; Gilbert, D. J.; Copeland,

[157985]N. G.; Jenkins, N. A.; Simister, N. E.: Mouse MHC class I-like Fc

[157986]receptor encoded outside the MHC. J. Immun. 151: 6076-6088, 1993.

[157987]

[157988]

[157989]15421.

[157990]Junghans, R. P.; Anderson, C. L.: The protection receptor for

[157991]IgG catabolism is the beta-2-microglobulin-containing neonatal intestinal

[157992]transport receptor. Proc. Nat. Acad. Sci. 93: 5512–5516,
1996.

[157993]

[157994]

[157995]15422.

[157996]Junghans, R. P.; Ebralidze, A.; Tiwari, B.: Does (CUG)_n re-
peat

[157997]in DMPK mRNA 'paint' chromosome 19 to suppress distant
genes to create

[157998]the diverse phenotype of myotonic dystrophy? A new hy-
pothesis of long-range

[157999]cis autosomal inactivation. Neurogenetics 3: 59–67, 2001.

[158000]

[158001]

[158002]15423.

[158003]Kandil, E.; Egashira, M.; Miyosi, O.; Niikawa, N.; Ishibashi,
T.;

[158004]Kasahara, M.: The human gene encoding the heavy chain
of the major

[158005]histocompatibility complex class I-like Fc receptor
(FCGRT) maps to

[158006]19q13.3. Cytogenet. Cell Genet. 73: 97–98, 1996.

[158007]

[158008]

[158009]15424.

[158010]Story, C. M.; Mikulska, J. E.; Simister, N. E.: A major histocompatibility

[158011]complex class I-like Fc receptor cloned from human placenta: possible

[158012]role in transfer of immunoglobulin G from mother to fetus. J. Exp.

[158013]Med. 180: 2377–2381, 1994.

[158014]

[158015]

[158016]15425.

[158017]Waldmann, T. A.; Strober, W.: Metabolism of immunoglobulins. Prog.

[158018]Allergy 13: 1–110, 1969.

[158019]

[158020]

[158021]15426.

[158022]Hart, T. C.; Price, J. A.; Bobby, P. L.; Pettenati, M. J.; Shashi,

[158023]V.; Von Kap Herr, C.; Van Dyke, T. E.: Cytogenetic assignment and

[158024]physical mapping of the human DGKE gene to chromosome 17q22. Genomics 56:

[158025]233–235, 1999.

[158026]

[158027]

[158028]15427.

[158029]Tang, W.; Bunting, M.; Zimmerman, G. A.; McIntyre, T. M.; Prescott,

[158030]S. M.: Molecular cloning of a novel human diacylglycerol kinase highly

[158031]selective for arachidonate-containing substrates. J. Biol. Chem. 271:

[158032]10237–10241, 1996.

[158033]

[158034]

[158035]15428.

[158036]Beddow, A. L.; Richards, S. A.; Orem, N. R.; Macara, I. G.:

The

[158037]Ran/TC4 GTPase-binding domain: identification by expression cloning

[158038]and characterization of a conserved sequence motif. Proc. Nat. Acad.

[158039]Sci. 92: 3328-3332, 1995.

[158040]

[158041]

[158042]15429.

[158043]Fauser, S.; Aslanukov, A.; Roepman, R.; Ferreira, P. A.: Genomic

[158044]organization, expression, and localization of murine Ran-binding protein

[158045]2 (RanBP2) gene. Mammalian Genome 12: 406-415, 2001.

[158046]

[158047]

[158048]15430.

[158049]Krebber, H.; Bastians, H.; Hoheisel, J.; Lichter, P.; Ponstingl,

[158050]H.; Joos, S.: Localization of the gene encoding the Ran-binding protein

[158051]RanBP2 to human chromosome 2q11-q13 by fluorescence in situ hybridization. Genomics 43:

[158052]247-248, 1997.

[158053]

[158054]

[158055]15431.

[158056]Pichler, A.; Gast, A.; Seeler, J. S.; Dejean, A.; Melchior, F.

[158057]: The nucleoporin RanBP2 has SUMO1 E3 ligase activity. Cell 108:

[158058]109-120, 2002.

[158059]

[158060]

[158061]15432.

[158062]Wu, J.; Matunis, M. J.; Kraemer, D.; Blobel, G.; Coutavas, E.:

[158063]Nup358, a cytoplasmically exposed nucleoporin with peptide repeats,

[158064]Ran-GTP binding sites, zinc fingers, a cyclophilin A ho-

mologous domain,

[158065]and a leucine-rich region. J. Biol. Chem. 270:

14209–14213, 1995.

[158066]

[158067]

[158068]15433.

[158069]Yokoyama, N.; Hayashi, N.; Seki, T.; Pante, N.; Ohba, T.;

Nishii,

[158070]K.; Kuma, K.; Hayashida, T.; Miyata, T.; Abei, U.; Fukui, M.;

Nishimoto,

[158071]T.: A giant nucleopore protein that binds Ran/TC4. Nature

376:

[158072]184–188, 1995.

[158073]

[158074]

[158075]15434.

[158076]Dhar, S. K.; Yoshida, K.; Machida, Y.; Khaira, P.; Chaud-

huri, B.;

[158077]Wohlschlegel, J. A.; Leffak, M.; Yates, J.; Dutta, A.: Replica-

tion

[158078]from oriP of Epstein-Barr virus requires human ORC and is

inhibited

[158079]by geminin. Cell 106: 287–296, 2001.

[158080]

[158081]

[158082]15435.

[158083]Gavin, K. A.; Hidaka, M.; Stillman, B.: Conserved initiator
proteins

[158084]in eukaryotes. Science 270: 1667–1671, 1995.

[158085]

[158086]

[158087]15436.

[158088]Ohtani, K.; DeGregori, J.; Leone, G.; Herendeen, D. R.;
Kelly,

[158089]T. J.; Nevins, J. R.: Expression of the HsOrc1 gene, a hu-
man ORC1

[158090]homolog, is regulated by cell proliferation via the E2F
transcription

[158091]factor. Molec. Cell. Biol. 16: 6977–6984, 1996.

[158092]

[158093]

[158094]15437.

[158095]Takahara, K.; Bong, M.; Brevard, R.; Eddy, R. L.; Haley, L.
L.;

[158096]Sait, S. J.; Shows, T. B.; Hoffman, G. G.; Greenspan, D. S.:
Mouse

[158097]and human homologues of the yeast origin of replication
recognition

[158098]complex subunit ORC2 and chromosomal localization of
the cognate human

[158099]gene ORC2L. Genomics 31: 119–122, 1996.

[158100]

[158101]

[158102]15438.

[158103]Chang, A. C.–M.; Janosi, J.; Hulsbeek, M.; de Jong, D.; Jef-
frey,

[158104]K. J.; Noble, J. R.; Reddel, R. R.: A novel human cDNA
highly homologous

[158105]to the fish hormone stanniocalcin. Molec. Cell. Endocr.
112: 241–247,

[158106]1995.

[158107]

[158108]

[158109]15439.

[158110]Chang, A. C.-M.; Jeffrey, K. J.; Tokutake, Y.; Shimamoto, A.; Neumann,

[158111]A. A.; Dunham, M. A.; Cha, J.; Sugawara, M.; Furuichi, Y.; Reddel,

[158112]R. R.: Human stanniocalcin (STC): genomic structure, chromosomal

[158113]localization, and the presence of CAG trinucleotide repeats. Genomics 47:

[158114]393–398, 1998.

[158115]

[158116]

[158117]15440.

[158118]Jellinek, D. A.; Chang, A. C.; Larsen, M. R.; Wang, X.; Robinson,

[158119]P. J.; Reddel, R. R.: Stanniocalcin 1 and 2 are secreted as phosphoproteins

[158120]from human fibrosarcoma cells. Biochem. J. 350: 453–461, 2000.

[158121]

[158122]

[158123]15441.

[158124]Olsen, H. S.; Cepeda, M. A.; Zhang, Q.-Q.; Rosen, C. A.;
Vozzolo,

[158125]B. L.; Wagner, G. F.: Human stanniocalcin: a possible hor-
monal regulator

[158126]of mineral metabolism. Proc. Nat. Acad. Sci. 93:
1792-1796, 1996.

[158127]

[158128]

[158129]15442.

[158130]Varghese, R.; Wong, C. K. C.; Deol, H.; Wagner, G. F.; Di-
Mattia,

[158131]G. E.: Comparative analysis of mammalian stanniocalcin
genes. Endocrinology 139:

[158132]4714-4725, 1998.

[158133]

[158134]

[158135]15443.

[158136]Wagner, G. F.; Guiraudon, C. C.; Milliken, C.; Copp, D. H.:
Immunological

[158137]and biological evidence for a stanniocalcin-like hormone
in human

[158138]kidney. Proc. Nat. Acad. Sci. 92: 1871–1875, 1995.

[158139]

[158140]

[158141]15444.

[158142]Bartsch, O.; Wagner, A.; Hinkel, G. K.; Lichtner, P.; Murken,
J.;

[158143]Schuffenhauer, S.: No evidence for chromosomal mi-
crodeletions at

[158144]the second DiGeorge syndrome locus on 10p near
D10S585. (Letter) Am.

[158145]J. Med. Genet. 83: 425–426, 1999.

[158146]

[158147]

[158148]15445.

[158149]Daw, S. C. M.; Taylor, C.; Kraman, M.; Call, K.; Mao, J.;
Schuffenhauer,

[158150]S.; Meitinger, T.; Lipson, T.; Goodship, J.; Scambler, P.: A
common

[158151]region of 10p deleted in DiGeorge and velocardiofacial
syndromes. Nature

[158152]Genet. 13: 458–461, 1996.

[158153]

[158154]

[158155]15446.

[158156]Lichtner, P.; Konig, R.; Hasegawa, T.; Van Esch, H.;
Meitinger,

[158157]T.; Schuffenhauer, S.: An HDR (hypoparathyroidism, deaf-
ness, renal

[158158]dysplasia) syndrome locus maps distal to the DiGeorge
syndrome region

[158159]on 10p13/14. J. Med. Genet. 37: 33–37, 2000.

[158160]

[158161]

[158162]15447.

[158163]Chen, C.–Y.; Gherzi, R.; Ong, S.–E.; Chan, E. L.; Raijmak-
ers, R.;

[158164]Pruijn, G. J. M.; Stoecklin, G.; Moroni, C.; Mann, M.; Karin, M.:

[158165]AU binding proteins recruit the exosome to degrade ARE-containing

[158166]mRNAs. Cell 107: 451–464, 2001.

[158167]

[158168]

[158169]15448.

[158170]Dangel, A. W.; Shen, L.; Mendoza, A. R.; Wu, L.-c.; Yu, C. Y.:

[158171]Human helicase gene SKI2W in the HLA class III region exhibits striking

[158172]structural similarities to the yeast antiviral gene SKI2 and to the

[158173]human gene KIAA0052: emergence of a new gene family. Nucleic Acids

[158174]Res. 23: 2120–2126, 1995.

[158175]

[158176]

[158177]15449.

[158178]Lee, S.-G.; Lee, I.; Park, S. H.; Kang, C.; Song, K.: Identifi-
cation

[158179]and characterization of a human cDNA homologous to
yeast SKI2. Genomics 25:

[158180]660–666, 1995.

[158181]

[158182]

[158183]15450.

[158184]Shen, L.; Wu, L. C.; Sanlioglu, S.; Chen, R.; Mendoza, A. R.;
Dangel,

[158185]A. W.; Carroll, M. C.; Zipf, W. B.; Yu, C. Y.: Structure and
genetics

[158186]of the partially duplicated gene RP located immediately
upstream of

[158187]the complement C4A and the C4B genes in the HLA class
III region:

[158188]molecular cloning, exon–intron structure, composite
retroposon, and

[158189]breakpoint of gene duplication. J. Biol. Chem. 269:
8466–8476, 1994.

[158190]

[158191]

[158192]15451.

[158193]Yang, Z.; Shen, L.; Dangel, A. W.; Wu, L.-C.; Yu, C. Y.: Four

[158194]ubiquitously expressed genes, RD (D6S45)–SKI2W
(SKIV2L)–DOM3Z–RP1

[158195](D6S60E), are present between complement component
genes factor B

[158196]and C4 in the class III region of the HLA. Genomics 53:
338–347,

[158197]1998.

[158198]

[158199]

[158200]15452.

[158201]Doyle, K.; Zhang, Y.; Baer, R.; Bina, M.: Distinguishable
patterns

[158202]of protein–DNA interactions involving complexes of basic
helix–loop–helix

[158203]proteins. J. Biol. Chem. 269: 12099–12105, 1994.

[158204]

[158205]

[158206]15453.

[158207]Hu, J.-S.; Olson, E. N.; Kingston, R. E.: HEB, a helix-loop-helix

[158208]protein related to E2A and ITF2 that can modulate the DNA-binding

[158209]ability of myogenic regulatory factors. Molec. Cell. Biol. 12: 1031-1042,

[158210]1992.

[158211]

[158212]

[158213]15454.

[158214]Sawada, S.; Littman, D. R.: A heterodimer of HEB and an E12-related

[158215]protein interacts with the CD4 enhancer and regulates its activity

[158216]in T-cell lines. Molec. Cell. Biol. 13: 5620-5628, 1993.

[158217]

[158218]

[158219]15455.

[158220]Zhang, Y.; Babin, J.; Feldhaus, A. L.; Singh, H.; Sharp, P. A.;

[158221]Bina, M.: HTF4: a new human helix-loop-helix protein.

Nucleic Acids

[158222]Res. 19: 4555 only, 1991.

[158223]

[158224]

[158225]15456.

[158226]Zhang, Y.; Bina, M.: The nucleotide sequence of the human transcription

[158227]factor HTF4a cDNA. DNA Sequence 2: 397-403, 1992.

[158228]

[158229]

[158230]15457.

[158231]Zhang, Y.; Flejter, W. L.; Barcroft, C. L.; Riviere, M.; Szpirer,

[158232]J.; Szpirer, C.; Bina, M.: Localization of the human HTF4 transcription

[158233]factors 4 gene (TCF12) to chromosome 15q21. Cytogenet. Cell Genet. 68:

[158234]235-238, 1995.

[158235]

[158236]

[158237]15458.

[158238]Dubrulle, J.; McGrew, M. J.; Pourquie, O.: FGF signaling controls

[158239]somite boundary position and regulates segmentation clock control

[158240]of spatiotemporal Hox gene activation. Cell 106: 219–232, 2001.

[158241]

[158242]

[158243]15459.

[158244]Fukuchi–Shimogori, T.; Grove, E. A.: Neocortex patterning by the

[158245]secreted signaling molecule FGF8. Science 294: 1071–1074, 2001.

[158246]

[158247]

[158248]15460.

[158249]Gemel, J.; Gorry, M.; Ehrlich, G. D.; MacArthur, C. A.: Structure

[158250]and sequence of human FGF8. Genomics 35: 253–257,
1996.

[158251]

[158252]

[158253]15461.

[158254]Rowen, L.; Young, J.; Birditt, B.; Kaur, A.; Madan, A.;
Philipps,

[158255]D. L.; Qin, S.; Minx, P.; Wilson, R. K.; Hood, L.; Graveley, B.
R.

[158256]: Analysis of the human neurexin genes: alternative splic-
ing and the

[158257]generation of protein diversity. Genomics 79: 587–597,
2002.

[158258]

[158259]

[158260]15462.

[158261]Tabuchi, K.; Sudhof, T. C.: Structure and evolution of
neurexin

[158262]genes: insight into the mechanism of alternative splicing.
Genomics 79:

[158263]849–859, 2002.

[158264]

[158265]

[158266]15463.

[158267]Ullrich, B.; Ushkaryov, Y. A.; Sudhof, T. C.: Cartography of
neurexins:

[158268]more than 1000 isoforms generated by alternative splicing and expressed

[158269]in distinct subsets of neurons. Neuron 14: 497–507,
1995.

[158270]

[158271]

[158272]15464.

[158273]Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T.
C.

[158274]: Neurexins: synaptic cell surface proteins related to the
alpha-latrotoxin

[158275]receptor and laminin. Science 257: 50–56, 1992.

[158276]

[158277]

[158278]15465.

[158279]Bergman, L.; Silins, G.; Grimmond, S.; Hummerich, H.;
Stewart,

[158280]C.; Little, P.; Hayward, N.: A 500-kb sequence-ready cos-
mid contig

[158281]and transcript map of the MEN1 region on 11q13. Ge-
nomics 55: 49–56,

[158282]1999.

[158283]

[158284]

[158285]15466.

[158286]Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T.
C.:

[158287]Neurexins: synaptic cell surface proteins related to the al-
pha-latrotoxin

[158288]receptor and laminin. Science 257: 50–56, 1992.

[158289]

[158290]

[158291]15467.

[158292]Scheiffele, P.; Fan, J.; Choih, J.; Fetter, R.; Serafini, T.:

[158293]Neurologin expressed in nonneuronal cells triggers presy-

naptic development

[158294]in contacting axons. Cell 101: 657–669, 2000.

[158295]

[158296]

[158297]15468.

[158298]Abderrahmani, A.; Steinmann, M.; Plaisance, V.; Niederhauser, G.;

[158299]Haefliger, J.–A.; Mooser, V.; Bonny, C.; Nicod, P.; Waeber, G.: The

[158300]transcriptional repressor REST determines the cell-specific expression

[158301]of the human MAPK8IP1 gene encoding IB1 (JIP–1). Molec. Cell. Biol. 21:

[158302]7256–7627, 2001.

[158303]

[158304]

[158305]15469.

[158306]Chen, Z.–F.; Paquette, A. J.; Anderson, D. J.: NRSF/REST is required

[158307]in vivo for repression of multiple neuronal target genes

during embryogenesis. Nature

[158308]Genet. 20: 136–142, 1998.

[158309]

[158310]

[158311]15470.

[158312]Palm, K.; Metsis, M.; Timmusk, T.: Neuron-specific splicing of

[158313]zinc finger transcription factor REST/NRSF/XBR is frequent in neuroblastomas

[158314]and conserved in human, mouse and rat. Molec. Brain Res. 72: 30–39,

[158315]1999.

[158316]

[158317]

[158318]15471.

[158319]Schoenherr, C. J.; Anderson, D. J.: The neuron-restrictive silencer

[158320]factor (NRSF): a coordinate repressor of multiple neuron-specific

[158321]genes. Science 267: 1360–1363, 1995.

[158322]

[158323]

[158324]15472.

[158325]Thiel, G.; Lietz, M.; Cramer, M.: Biological activity and
modulator

[158326]structure of RE-1-silencing transcription factor (REST), a
repressor

[158327]of neuronal genes. J. Biol. Chem. 273: 26891-26899,
1998.

[158328]

[158329]

[158330]15473.

[158331]Cavaloc, Y.; Popielarz, M.; Fuchs, J.-P.; Gattoni, R.;
Stevenin,

[158332]J.: Characterization and cloning of the human splicing fac-
tor 9G8:

[158333]a novel 35 kDa factor of the serine/arginine protein fam-
ily. EMBO

[158334]J. 13: 2639-2649, 1994.

[158335]

[158336]

[158337]15474.

[158338]Popielarz, M.; Cavaloc, Y.; Mattei, M.-G.; Gattoni, R.;
Stevenin,

[158339]J.: The gene encoding human splicing factor 9G8: struc-
ture, chromosomal

[158340]localization, and expression of alternatively processed
transcripts. J.

[158341]Biol. Chem. 270: 17830-17835, 1995.

[158342]

[158343]

[158344]15475.

[158345]Chiang, C.-M.; Roeder, R. G.: Cloning of an intrinsic hu-
man TFIID

[158346]subunit that interacts with multiple transcriptional activa-
tors. Science 267:

[158347]531-536, 1995.

[158348]

[158349]

[158350]15476.

[158351]Wu, Q.; Zhang, T.; Cheng, J.-F.; Kim, Y.; Grimwood, J.;
Schmutz,

[158352]J.; Dickson, M.; Noonan, J. P.; Zhang, M. Q.; Myers, R. M.;
Maniatis,

[158353]T.: Comparative DNA sequence analysis of mouse and hu-
man protocadherin

[158354]gene clusters. Genome Res. 11: 389–404, 2001.

[158355]

[158356]

[158357]15477.

[158358]Aoki, K.; Inazawa, J.; Takahashi, T.; Nakahara, K.; Kasai,
M.:

[158359]Genomic structure and chromosomal localization of the
gene encoding

[158360]translin, a recombination hotspot binding protein. Ge-
nomics 43:

[158361]237–241, 1997.

[158362]

[158363]

[158364]15478.

[158365]Aoki, K.; Nakahara, K.; Ikegawa, C.; Seto, M.; Takahashi, T.; Minowada,

[158366]J.; Strominger, J. L.; Maziarz, R. T.; Kasai, M.: Nuclear proteins

[158367]binding to a novel target sequence within the recombination hotspot

[158368]regions of Bcl-2 and the immunoglobulin D(H) gene family. Oncogene 9:

[158369]1109-1115, 1994.

[158370]

[158371]

[158372]15479.

[158373]Blair, I. P.; Gibson, R. R.; Bennett, C. L.; Chance, P. F.: Search

[158374]for genes involved in Joubert syndrome: evidence that one or more

[158375]major loci are yet to be identified and exclusion of candidate genes

[158376]EN1, EN2, FGF8, and BARHL1. Am. J. Med. Genet. 107: 190-196, 2002.

[158377]

[158378]

[158379]15480.

[158380]Berge, K. E.; Tian, H.; Graf, G. A.; Yu, L.; Grishin, N. V.;
Schultz,

[158381]J.; Kwiterovich, P.; Shan, B.; Barnes, R.; Hobbs, H. H.: Ac-
cumulation

[158382]of dietary cholesterol in sitosterolemia caused by muta-
tions in adjacent

[158383]ABC transporters. Science 290: 1771–1775, 2000.

[158384]

[158385]

[158386]15481.

[158387]Bodzioch, M.; Orso, E.; Klucken, J.; Langmann, T.;
Bottcher, A.;

[158388]Diederich, W.; Drobnik, W.; Barlage, S.; Buchler, C.;
Porsch-Ozcurumez,

[158389]M.; Kaminski, W. E.; Hahmann, H. W.; Oette, K.; Rothe, G.;
Aslanidis,

[158390]C.; Lackner, K. J.; Schmitz, G.: The gene encoding ATP-
binding cassette

[158391]transporter 1 is mutated in Tangier disease. Nature Genet.

22: 347–351,

[158392]1999.

[158393]

[158394]

[158395]15482.

[158396]Brooks–Wilson, A.; Marcil, M.; Clee, S. M.; Zhang, L.–H.;

Roomp,

[158397]K.; van Dam, M.; Yu, L.; Brewer, C.; Collins, J. A.; Mol–

huizen, H.

[158398]O. F.; Loubser, O.; Ouelette, B. F. F.; and 14 others: Muta–

tions

[158399]in ABC1 in Tangier disease and familial high–density

lipoprotein deficiency. Nature

[158400]Genet. 22: 336–345, 1999.

[158401]

[158402]

[158403]15483.

[158404]Asamoah, A.; Wilson, A. F.; Elston, R. C.; Dalferes, E., Jr.;

Berenson,

[158405]G. S.: Segregation and linkage analyses of dopamine–

beta-hydroxylase

[158406]activity in a six-generation pedigree. Am. J. Med. Genet.
27: 613-621,

[158407]1987.

[158408]

[158409]

[158410]15484.

[158411]Biaggioni, I.; Goldstein, D. S.; Atkinson, T.; Robertson, D.:

[158412]Dopamine-beta-hydroxylase deficiency in humans. Neu-
rology 40: 370-373,

[158413]1990.

[158414]

[158415]

[158416]15485.

[158417]Biaggioni, I.; Robertson, D.: Endogenous restoration of
noradrenaline

[158418]by precursor therapy in dopamine-beta-hydroxylase defi-
ciency. Lancet II:

[158419]1170-1172, 1987.

[158420]

[158421]

[158422]15486.

[158423]Craig, S. P.; Buckle, V. J.; Lamouroux, A.; Mallet, J.; Craig,

[158424]I. W.: Localization of the human dopamine beta hydroxy-
lase (DBH)

[158425]gene to chromosome 9q34. Cytogenet. Cell Genet. 48:
48-50, 1988.

[158426]

[158427]

[158428]15487.

[158429]Dunnette, J.; Weinshilboum, R.: Human serum dopamine
beta-hydroxylase:

[158430]correlation of enzymatic activity with immunoreactive
protein in genetically

[158431]defined samples. Am. J. Hum. Genet. 28: 155-166, 1976.

[158432]

[158433]

[158434]15488.

[158435]Dunnette, J.; Weinshilboum, R.: Inheritance of low immunoreactive

[158436]human plasma dopamine-beta-hydroxylase: radioimmunoassay studies. J.

[158437]Clin. Invest. 60: 1080-1087, 1977.

[158438]

[158439]

[158440]15489.

[158441]Elston, R. C.; Namboodiri, K. K.; Hames, C. G.: Segregation and

[158442]linkage analysis of dopamine-beta-hydroxylase activity. Hum. Hered. 29:

[158443]284-292, 1979.

[158444]

[158445]

[158446]15490.

[158447]Gershon, E. S.; Goldin, L. R.: Segregation and linkage studies

[158448]of plasma dopamine-beta-hydroxylase (DBH), erythrocyte catechol-O-methyltransferase

[158449](COMT) and platelet monoamine oxidase (MAO): possible linkage between

[158450]the ABO locus and a gene controlling DBH activity.

(Abstract) Am.

[158451]J. Hum. Genet. 33: 136A only, 1981.

[158452]

[158453]

[158454]15491.

[158455]Goldin, L. R.; Gershon, E. S.; Lake, C. R.; Murphy, D. L.; McGinniss,

[158456]M.; Sparkes, R. S.: Segregation and linkage studies of plasma dopamine-beta-hydroxylase

[158457](DBH), erythrocyte catechol-O-methyltransferase (COMT), and platelet

[158458]monoamine oxidase (MAO): possible linkage between the ABO locus and

[158459]a gene controlling DBH activity. Am. J. Hum. Genet. 34: 250-262,

[158460]1982.

[158461]

[158462]

[158463]15492.

[158464]Joh, T. H.; Baetge, E. E.; Reis, D. J.: Evidence for the existence
tence

[158465]of a single gene or linked genes coding for catecholamine
biosynthetic

[158466]enzymes. Trans. Assoc. Am. Phys. 96: 38–43, 1983.

[158467]

[158468]

[158469]15493.

[158470]Joh, T. H.; Baetge, E. E.; Ross, M. E.; Albert, V. R.; Moon, H.

[158471]M.; Reis, D. J.: Existence of catecholamine biosynthetic
enzyme gene

[158472]family. (Abstract) Clin. Res. 31: 528 only, 1983.

[158473]

[158474]

[158475]15494.

[158476]Joh, T. H.; Baetge, E. E.; Ross, M. E.; Reis, D. J.: Biochem–
istry

[158477]and molecular biology of catecholamine neurons: a single

gene or gene

[158478]family hypothesis. Clin. Exp. Hypertension 6A: 11–21,
1984.

[158479]

[158480]

[158481]15495.

[158482]Kobayashi, K.; Kurosawa, Y.; Fujita, K.; Nagatsu, T.: Hu-
man dopamine

[158483]beta-hydroxylase gene: two mRNA types having different
3-prime-terminal

[158484]regions are produced through alternative polyadenylation.
Nucleic

[158485]Acids Res. 17: 1089–1102, 1989.

[158486]

[158487]

[158488]15496.

[158489]Lamouroux, A.; Vigny, A.; Faucon Biguet, N.; Darmon, M.
C.; Franck,

[158490]R.; Henry, J.–P.; Mallet, J.: The primary structure of human
dopamine-beta-hydroxylase:

[158491]insights into the relationship between the soluble and the

membrane-bound

[158492]forms of the enzyme. EMBO J. 6: 3931–3937, 1987.

[158493]

[158494]

[158495]15497.

[158496]Lea, R. A.; Dohy, A.; Jordan, K.; Quinlan, S.; Brimage, P. J.;

[158497]Griffiths, L. R.: Evidence for allelic association of the
dopamine

[158498]beta-hydroxylase gene (DBH) with susceptibility to typical
migraine. Neurogenetics 3:

[158499]35–40, 2000.

[158500]

[158501]

[158502]15498.

[158503]Mathias, C. J.; Bannister, R. B.; Cortelli, P.; Heslop, K.; Po-
lak,

[158504]J. M.; Raimbach, S.; Springall, D. R.; Watson, L.: Clinical,
autonomic

[158505]and therapeutic observations in two siblings with postural
hypotension

[158506]and sympathetic failure due to an inability to synthesize
noradrenaline

[158507]from dopamine because of a deficiency of dopamine beta
hydroxylase. Quart.

[158508]J. Med. 75: 617–633, 1990.

[158509]

[158510]

[158511]15499.

[158512]McKinney, E. F.; Walton, R. T.; Yudkin, P.; Fuller, A.; Hal-
dar,

[158513]N. A.; Mant, D.; Murphy, M.; Welsh, K. I.; Marshall, S. E.:
Association

[158514]between polymorphisms in dopamine metabolic enzymes
and tobacco consumption

[158515]in smokers. Pharmacogenetics 10: 483–491, 2000.

[158516]

[158517]

[158518]15500.

[158519]O'Malley, K. L.; Mauron, A.; Raese, J.; Barchas, J. D.; Kedes,

[158520]L.: Genes for catecholamine biosynthesis: cloning by ex–

pression and

[158521]identification of the cDNA for rat dopamine beta-hydroxylase. Proc.

[158522]Nat. Acad. Sci. 80: 2161–2165, 1983.

[158523]

[158524]

[158525]15501.

[158526]Ogihara, T.; Nugent, C. A., Jr.; Shen, S.-W.; Goldfein, S.:
Serum

[158527]dopamine-beta-hydroxylase activity in parents and children. J. Lab.

[158528]Clin. Med. 85: 566–573, 1975.

[158529]

[158530]

[158531]15502.

[158532]Perry, S. E.; Summar, M. L.; Phillips, J. A., III; Robertson,

[158533]D.: Linkage analysis of the human dopamine beta-hydroxylase gene. Genomics 10:

[158534]493–495, 1991.

[158535]

[158536]

[158537]15503.

[158538]Robertson, D.; Haile, V.; Perry, S. E.; Robertson, R. M.;
Phillips,

[158539]J. A., III; Biaggioni, I.: Dopamine beta-hydroxylase defi-
ciency:

[158540]a genetic disorder of cardiovascular regulation. Hyperten-
sion 18:

[158541]1-8, 1991.

[158542]

[158543]

[158544]15504.

[158545]Ross, S. B.; Wetterberg, L.; Myrhed, M.: Genetic control of
plasma

[158546]dopamine-beta-hydroxylase. Life Sci. 12: 529-532, 1973.

[158547]

[158548]

[158549]15505.

[158550]Schanberg, S. M.; Stone, R. A.; Kirshner, N.; Gunnells, J. C.;

[158551]Robinson, R. R.: Plasma dopamine beta-hydroxylase: a
possible aid

[158552]in the study and evaluation of hypertension. Science 183:
523-525,

[158553]1974.

[158554]

[158555]

[158556]15506.

[158557]Marlar, R. A.; Griffin, J. H.: Deficiency of protein C inhibitor

[158558]in combined factor V/VIII deficiency disease. J. Clin. In-
vest. 66:

[158559]1186-1189, 1980.

[158560]

[158561]

[158562]15507.

[158563]Meijers, J. C.; Chung, D. W.: Organization of the gene cod-
ing

[158564]for human protein C inhibitor (plasminogen activator in-

hibitor-3):

[158565]assignment of the gene to chromosome 14. J. Biol. Chem.
266: 15028-15034,
[158566]1991.

[158567]

[158568]

[158569]15508.

[158570]Nichols, W. C.; Seligsohn, U.; Zivelin, A.; Terry, V. H.;
Arnold,

[158571]N. D.; Siemieniak, D. R.; Kaufman, R. J.; Ginsburg, D.:
Linkage of

[158572]combined factors V and VIII deficiency to chromosome
18q by homozygosity

[158573]mapping. J. Clin. Invest. 99: 596-601, 1997.

[158574]

[158575]

[158576]15509.

[158577]Suzuki, K.; Deyashiki, Y.; Nishioka, J.; Kurachi, K.; Akira,
M.;

[158578]Yamamoto, S.; Hashimoto, S.: Characterization of a cDNA

for human

[158579]protein C inhibitor: a new member of the plasma serine
protease inhibitor

[158580]superfamily. J. Biol. Chem. 262: 611–616, 1987.

[158581]

[158582]

[158583]15510.

[158584]Cathelineau, L.; Dinh, D. P.; Briand, P.; Kamoun, P.: Studies

[158585]on complementation in argininosuccinate synthetase and
argininosuccinate

[158586]lyase deficiencies in human fibroblasts. Hum. Genet. 57:
282–284,

[158587]1981.

[158588]

[158589]

[158590]15511.

[158591]Kromberg, J. G. R.; Castle, D.; Zwane, E. M.; Jenkins, T.:
Albinism

[158592]and skin cancer in southern Africa. Clin. Genet. 36:
43–52, 1989.

[158593]

[158594]

[158595]15512.

[158596]El Kahloun, A.; Chauvel, B.; Mauvieux, V.; Dorval, I.;
Jouanolle,

[158597]A.-M.; Gicquel, I.; Le Gall, J.-Y.; David, V.: Localization of
seven

[158598]new genes around the HLA-A locus. Hum. Molec. Genet.
2: 55-60, 1993.

[158599]

[158600]

[158601]15513.

[158602]Beattie, E. C.; Stellwagen, D.; Morishita, W.; Bresnahan, J.
C.;

[158603]Ha, B. K.; Von Zastrow, M.; Beattie, M. S.; Malenka, R. C.:
Control

[158604]of synaptic strength by glial TNF-alpha. Science 295:
2282-2285,

[158605]2002.

[158606]

[158607]

[158608]15514.

[158609]Beutler, B.; Krochin, N.; Milsark, I. W.; Luedke, C.; Cerami, A.

[158610]: Control of cachectin (tumor necrosis factor) synthesis: mechanisms

[158611]of endotoxin resistance. Science 232: 977–980, 1986.

[158612]

[158613]

[158614]15515.

[158615]Brenner, D. A.; O'Hara, M.; Angel, P.; Chojkier, M.; Karin, M.

[158616]: Prolonged activation of JUN and collagenase genes by tumour necrosis

[158617]factor- α . Nature 337: 661–663, 1989.

[158618]

[158619]

[158620]15516.

[158621]Broudy, V. C.; Kaushansky, K.; Segal, G. M.; Harlan, J. M.;

Adamson,

[158622]J. W.: Tumor necrosis factor type alpha stimulates human endothelial

[158623]cells to produce granulocyte/macrophage colony-stimulating factor. Proc.

[158624]Nat. Acad. Sci. 83: 7467–7471, 1986.

[158625]

[158626]

[158627]15517.

[158628]Bruce, A. J.; Boling, W.; Kindy, M. S.; Peschon, J.; Kraemer, P.

[158629]J.; Carpenter, M. K.; Holtsberg, F. W.; Mattson, M. P.: Altered neuronal

[158630]and microglial responses to excitotoxic and ischemic brain injury

[158631]in mice lacking TNF receptors. Nature Med. 2: 788–794, 1996.

[158632]

[158633]

[158634]15518.

[158635]Cabrera, M.; Shaw, M. A.; Sharples, C.; Williams, H.;

Castes, M.;

[158636]Convit, J.; Blackwell, J. M.: Polymorphism in tumor necrosis factor

[158637]genes associated with mucocutaneous leishmaniasis. J. Exp. Med. 182:

[158638]1259–1264, 1995.

[158639]

[158640]

[158641]15519.

[158642]Conway, D. J.; Holland, M. J.; Bailey, R. L.; Campbell, A. E.;

[158643]Mahdi, O. S.; Jennings, R.; Mbeni, E.; Mabey, D. C.: Scarring trachoma

[158644]is associated with polymorphism in the tumor necrosis factor alpha

[158645](TNF-alpha) gene promoter and with elevated TNF-alpha levels in tear

[158646]fluid. Infect. Immun. 65: 1003–1006, 1997.

[158647]

[158648]

[158649]15520.

[158650]Davis, J. M.; Narachi, M. A.; Alton, N. K.; Arakawa, T.:

Structure

[158651]of human tumor necrosis factor alpha derived from re-
combinant DNA. Biochemistry 26:

[158652]1322-1326, 1987.

[158653]

[158654]

[158655]15521.

[158656]Escobar-Morreale, H. F.; Calvo, R. M.; Sancho, J.; San Mil-
lan,

[158657]J. L.: TNF-alpha and hyperandrogenism: a clinical, bio-
chemical, and

[158658]molecular genetic study. J. Clin. Endocr. Metab. 86:
3761-3767,

[158659]2001.

[158660]

[158661]

[158662]15522.

[158663]Gorman, J. D.; Sack, K. E.; Davis, J. C., Jr.: Treatment of
ankylosing

[158664]spondylitis by inhibition of tumor necrosis factor- α .

New Eng.

[158665]J. Med. 346: 1349–1356, 2002.

[158666]

[158667]

[158668]15523.

[158669]Herrmann, S.–M.; Ricard, S.; Nicaud, V.; Mallet, C.;

Arveiler,

[158670]D.; Evans, A.; Ruidavets, J.–B.; Luc, G.; Bara, L.; Parra,

H.–J.;

[158671]Poirier, O.; Cambien, F.: Polymorphisms of the tumour
necrosis factor- α

[158672]gene, coronary heart disease and obesity. Europ. J. Clin.
Invest. 28:

[158673]59–66, 1998.

[158674]

[158675]

[158676]15524.

[158677]Inoko, H.; Trowsdale, J.: Linkage of TNF genes to the HLA–
B locus. Nucleic

[158678]Acids Res. 15: 8957–8962, 1987.

[158679]

[158680]

[158681]15525.

[158682]Knight, J. C.; Udalova, I.; Hill, A. V. S.; Greenwood, B. M.;

[158683]Peshu, N.; Marsh, K.; Kwiatkowski, D.: A polymorphism
that affects

[158684]OCT–1 binding to the TNF promoter region is associated
with severe

[158685]malaria. Nature Genet. 22: 145–150, 1999.

[158686]

[158687]

[158688]15526.

[158689]Koss, K.; Satsangi, J.; Fanning, G. C.; Welsh, K. I.; Jewell,

[158690]D. P.: Cytokine (TNF–alpha, LT–alpha, and IL–10) poly-
morphisms in

[158691]inflammatory bowel diseases and normal controls: differ-
ential effects

[158692]on production and allele frequencies. Genes Immun. 1:

185–190, 2000.

[158693]

[158694]

[158695]15527.

[158696]Marino, M. W.; Dunn, A.; Grail, D.; Inglese, M.; Noguchi, Y.;

[158697]Richards, E.; Jungbluth, A.; Wada, H.; Moore, M.; Williamson, B.;

[158698]Basu, S.; Old, L. J.: Characterization of tumor necrosis factor-deficient

[158699]mice. Proc. Nat. Acad. Sci. 94: 8093–8098, 1997.

[158700]

[158701]

[158702]15528.

[158703]McCusker, S. M.; Curran, M. D.; Dynan, K. B.; McCullagh, C. D.;

[158704]Urquhart, D. D.; Middleton, D.; Patterson, C. C.; McIlroy, S. P.;

[158705]Passmore, A. P.: Association between polymorphism in regulatory region

[158706]of gene encoding tumour necrosis factor- α and risk

of Alzheimer's

[158707]disease and vascular dementia: a case-control study.

Lancet 357:

[158708]436-439, 2001.

[158709]

[158710]

[158711]15529.

[158712]Remaley, A. T.; Rust, S.; Rosier, M.; Knapper, C.; Naudin,
L.;

[158713]Broccardo, C.; Peterson, K. M.; Koch, C.; Arnould, I.;
Prades, C.;

[158714]Duverger, N.; Funke, H.; Assman, G.; Dinger, M.; Dean, M.;
Chimini,

[158715]G.; Santamarina-Fojo, S.; Fredrickson, D. S.; Deneffe, P.
Brewer,

[158716]H. B., Jr.: Human ATP-binding cassette transporter 1
(ABC1): genomic

[158717]organization and identification of the genetic defect in the
original

[158718]Tangier disease kindred. Proc. Nat. Acad. Sci. 96:
12685-12690,

[158719]1999.

[158720]

[158721]

[158722]15530.

[158723]Rust, S.; Rosier, M.; Funke, H.; Real, J.; Amoura, Z.; Piette,

[158724]J.-C.; Deleuze, J.-F.; Brewer, H. B.; Duverger, N.; Deneffe,
P.; Assmann,

[158725]G.: Tangier disease is caused by mutations in the gene
encoding ATP-binding

[158726]cassette transporter 1. Nature Genet. 22: 352–355, 1999.

[158727]

[158728]

[158729]15531.

[158730]Young, S. G.; Fielding, C. J.: The ABCs of cholesterol ef-
flux. Nature

[158731]Genet. 22: 316–318, 1999.

[158732]

[158733]

[158734]15532.

[158735]Buchwald, M.: Complementation groups: one or more per

gene? Nature

[158736]Genet. 11: 228–230, 1995.

[158737]

[158738]

[158739]15533.

[158740]Stambolian, D.; Ai, Y.; Sidjanin, D.; Nesburn, K.; Sathe, G.;

[158741]Rosenberg, M.; Bergsma, D. J.: Cloning of the galactokinase cDNA

[158742]and identification of mutations in two families with cataracts. Nature

[158743]Genet. 10: 307–312, 1995.

[158744]

[158745]

[158746]15534.

[158747]Coppola, G.; De Michele, G.; Cavalcanti, F.; Pianese, L.; Perretti,

[158748]A.; Santoro, L.; Vita, G.; Toscano, A.; Amboni, M.; Grimaldi, G.;

[158749]Salvatore, E.; Caruso, G.; Filla, A.: Why do some Friedreich's ataxia

[158750]patients retain tendon reflexes? A clinical, neurophysio-
logical and

[158751]molecular study. J. Neurol. 246: 353–357, 1999.

[158752]

[158753]

[158754]15535.

[158755]Delatycki, M. B.; Knight, M.; Koenig, M.; Cossee, M.;
Williamson,

[158756]R.; Forrest, S. M.: G130V, a common FRDA point mutation,
appears

[158757]to have arisen from a common founder. Hum. Genet. 105:
343–346,

[158758]1999.

[158759]

[158760]

[158761]15536.

[158762]Delatycki, M. B.; Paris, D. B. B. P.; Gardner, R. J. M.; Nichol-
son,

[158763]G. A.; Nassif, N.; Storey, E.; MacMillan, J. C.; Collins, V.;
Williamson,

[158764]R.; Forrest, S. M.: Clinical and genetic study of Friedreich
ataxia

[158765]in an Australian population. Am. J. Med. Genet. 87:
168–174, 1999.

[158766]

[158767]

[158768]15537.

[158769]Durr, A.; Cossee, M.; Agid, Y.; Campuzano, V.; Mignard,
C.; Penet,

[158770]C.; Mandel, J.–L.; Brice, A.; Koenig, M.: Clinical and genetic
abnormalities

[158771]in patients with Friedreich's ataxia. New Eng. J. Med. 335:
1169–1175,

[158772]1996.

[158773]

[158774]

[158775]15538.

[158776]Filla, A.; De Michele, G.; Cavalcanti, F.; Pianese, L.; Monti-
celli,

[158777]A.; Campanella, G.; Coccozza, S.: The relationship between
trinucleotide

[158778](GAA) repeat length and clinical features in Friedreich
ataxia. Am.

[158779]J. Hum. Genet. 59: 554–560, 1996.

[158780]

[158781]

[158782]15539.

[158783]Fujita, R.; Hanauer, A.; Vincent, A.; Mandel, J.–L.; Koenig,
M.

[158784]: Physical mapping of two loci (D9S5 and D9S15) tightly
linked to

[158785]Friedreich ataxia locus (FRDA) and identification of nearby
CpG islands

[158786]by pulse–field gel electrophoresis. Genomics 10:
915–920, 1991.

[158787]

[158788]

[158789]15540.

[158790]Gray, J. V.; Johnson, K. J.: Waiting for frataxin. Nature
Genet. 16:

[158791]323–325, 1997.

[158792]

[158793]

[158794]15541.

[158795]Victoria, T.; Rafi, M. A.; Wenger, D. A.: Cloning of the canine
nine

[158796]GALC cDNA and identification of the mutation causing
globoid cell

[158797]leukodystrophy in West Highland White and Cairn terriers.
Genomics 33:

[158798]457–462, 1996.

[158799]

[158800]

[158801]15542.

[158802]Wenger, D. A.; Rafi, M. A.; Luzi, P.: Molecular genetics of
Krabbe

[158803]disease (globoid cell leukodystrophy): diagnostic and clinical
implications. Hum.

[158804]Mutat. 10: 268–279, 1997.

[158805]

[158806]

[158807]15543.

[158808]Zlotogora, J.; Chakraborty, S.; Knowlton, R. G.; Wenger, D. A.

[158809]: Krabbe disease locus mapped to chromosome 14 by genetic linkage. Am.

[158810]J. Hum. Genet. 47: 37–44, 1990.

[158811]

[158812]

[158813]15544.

[158814]Zlotogora, J.; Regev, R.; Zeigler, M.; Iancu, T. C.; Bach, G.

[158815]: Krabbe disease: increased incidence in a highly inbred community. Am.

[158816]J. Med. Genet. 21: 765–770, 1985.

[158817]

[158818]

[158819]15545.

[158820]Rushton, A. R.; Dawson, G.: Genetic linkage studies of the human

[158821]glycosphingolipid beta-galactosidases. Biochem. Genet.

15: 1071–1082,

[158822]1977.

[158823]

[158824]

[158825]15546.

[158826]Balciunaite, G.; Keller, M. P.; Balciunaite, E.; Piali, L.;

Zuklys,

[158827]S.; Mathieu, Y. D.; Gill, J.; Boyd, R.; Sussman, D. J.; Hollander,

[158828]G. A.: Wnt glycoproteins regulate the expression of FoxN1, the gene

[158829]defective in nude mice. Nature Immun. 15Oct, 2002. Note: Advance

[158830]Electronic Publication.

[158831]

[158832]

[158833]15547.

[158834]Frank, J.; Pignata, C.; Panteleyev, A. A.; Prowse, D. M.; Baden,

[158835]H.; Weiner, L.; Gaetaniello, L.; Ahmad, W.; Pozzi, N.; Caer-

halmi-Friedman,

[158836]P. B.; Aita, V. M.; Uyttendaele, H.; Gordon, D.; Ott, J.; Brissette,

[158837]J. L.; Christiano, A. M.: Exposing the human nude phenotype. Nature 398:

[158838]473–474, 1999.

[158839]

[158840]

[158841]15548.

[158842]Nehls, M.; Pfeifer, D.; Schorpp, M.; Hedrich, H.; Boehm, T.:
New

[158843]member of the winged-helix protein family disrupted in
mouse and rat

[158844]nude mutations. Nature 372: 103–107, 1994.

[158845]

[158846]

[158847]15549.

[158848]Pignata, C.; Fiore, M.; Guzzetta, V.; Castaldo, A.; Sebastio,
G.;

[158849]Porta, F.; Guarino, A.: Congenital alopecia and nail dystro-

phy associated

[158850]with severe functional T-cell immunodeficiency in two
sibs. Am. J.

[158851]Med. Genet. 65: 167–170, 1996.

[158852]

[158853]

[158854]15550.

[158855]Schorpp, M.; Hofmann, M.; Dear, T. N.; Boehm, T.: Char-
acterization

[158856]of mouse and human nude genes. Immunogenetics 46:
509–515, 1997.

[158857]

[158858]

[158859]15551.

[158860]Segre, J. A.; Nemhauser, J. L.; Taylor, B. A.; Nadeau, J. H.;
Lander,

[158861]E. S.: Positional cloning of the nude locus: genetic, physi-
cal, and

[158862]transcription maps of the region and mutations in the
mouse and rat. Genomics 28:

[158863]549–559, 1995.

[158864]

[158865]

[158866]15552.

[158867]Delpire, E.; Lu, J.; England, R.; Dull, C.; Thorne, T.: Deafness

[158868]and imbalance associated with inactivation of the secretory Na-K-2Cl

[158869]co-transporter. Nature Genet. 22: 192-195, 1999.

[158870]

[158871]

[158872]15553.

[158873]Evans, R. L.; Park, K.; Turner, R. J.; Watson, G. E.; Nguyen, H.-V.;

[158874]Dennett, M. R.; Hand, A. R.; Flagella, M.; Shull, G. E.; Melvin, J.

[158875]E.: Severe impairment of salivation in Na⁺/K⁺/2Cl⁻ co-transporter

[158876](NKCC1)-deficient mice. J. Biol. Chem. 275: 26720-26726, 2000.

[158877]

[158878]

[158879]15554.

[158880]Payne, J. A.; Xu, J.-C.; Haas, M.; Lytle, C. Y.; Ward, D.; Forbush,

[158881]B., III: Primary structure, functional expression, and chromosomal

[158882]localization of the bumetanide-sensitive Na-K-Cl cotransporter in

[158883]human colon. J. Biol. Chem. 270: 17977-17985, 1995.

[158884]

[158885]

[158886]15555.

[158887]Quaggin, S. E.; Payne, J. A.; Forbush, B., III; Igarashi, P.:

[158888]Localization of the renal Na-K-Cl cotransporter gene (Slc12a1) on

[158889]mouse chromosome 2. Mammalian Genome 6: 557-561, 1995.

[158890]

[158891]

[158892]15556.

[158893]Xu, J.-C.; Lytle, C.; Zhu, T. T.; Payne, J. A.; Benz, E., Jr.;

[158894]Forbush, B., III: Molecular cloning and functional expression of

[158895]the bumetanide-sensitive Na-K-Cl cotransporter. Proc. Nat. Acad.

[158896]Sci. 91: 2201-2205, 1994.

[158897]

[158898]

[158899]15557.

[158900]Daniels, S. E.; Bhattacharrya, S.; James, A.; Leaves, N. I.; Young,

[158901]A.; Hill, M. R.; Faux, J. A.; Ryan, G. F.; le Souef, P. N.; Lathrop,

[158902]G. M.; Musk, A. W.; Cookson, W. O. C. M.: A genome-wide search for

[158903]quantitative trait loci underlying asthma. Nature 383: 247-250,

[158904]1996.

[158905]

[158906]

[158907]15558.

[158908]MacDonald, S. M.; Paznekas, W. A.; Jabs, E. W.: Chromosomal localization

[158909]of tumor protein, translationally-controlled 1 (TPT1) encoding the

[158910]human histamine releasing factor (HRF) to 13q12-q14.

Cytogenet. Cell

[158911]Genet. (in-press), 1999.

[158912]

[158913]

[158914]15559.

[158915]MacDonald, S. M.; Rafnar, T.; Langdon, J.; Lichtenstein, L. M.

[158916]: Molecular identification of an IgE-dependent histamine-releasing

[158917]factor. Science 269: 688-690, 1995.

[158918]

[158919]

[158920]15560.

[158921]Zhang, K.; Saxon, A.; Max, E. E.: Two unusual forms of hu-

man immunoglobulin

[158922]E encoded by alternative RNA splicing of the epsilon heavy chain membrane

[158923]exons. J. Exp. Med. 176: 233–243, 1992.

[158924]

[158925]

[158926]15561.

[158927]Simister, N. E.; Mostov, K. E.: An Fc receptor structurally related

[158928]to MHC class I antigens. Nature 337: 184–187, 1989.

[158929]

[158930]

[158931]15562.

[158932]Sedlacek, Z.; Konecki, D. S.; Korn, B.; Klauck, S. M.; Poustka,

[158933]A.: Evolutionary conservation and genomic organization of XAP-4,

[158934]an Xq28 located gene coding for a human rab GDP-dissociation inhibitor

[158935](GDI). Mammalian Genome 5: 633–639, 1995.

[158936]

[158937]

[158938]15563.

[158939]Sedlacek, Z.; Munstermann, E.; Mincheva, A.; Lichter, P.;
Poustka,

[158940]A.: The human rab GDI beta gene with long retroposon-
rich introns

[158941]maps to 10p15 and its pseudogene to 7p11-p13. Mam-
malian Genome 9:

[158942]78-80, 1998.

[158943]

[158944]

[158945]15564.

[158946]Desseyn, J.-L.; Aubert, J.-P.; van Seuninghen, I.; Porchet, N.;

[158947]Laine, A.: Genomic organization of the 3-prime region of
the human

[158948]mucin gene MUC5B. J. Biol. Chem. 272: 16873-16883,
1997.

[158949]

[158950]

[158951]15565.

[158952]Desseyn, J.-L.; Guyonnet-Duperat, V.; Porchet, N.; Aubert, J.-P.;

[158953]Laine, A.: Human mucin gene MUC5B, the 10.7-kb large central exon

[158954]encodes various alternate subdomains resulting in a super-repeat:

[158955]structural evidence for a 11p15.5 gene family. J. Biol. Chem. 272:

[158956]3168–3178, 1997.

[158957]

[158958]

[158959]15566.

[158960]Gipson, I. K.; Moccia, R.; Spurr-Michaud, S.; Argueso, P.; Gargiulo,

[158961]A. R.; Hill, J. A., III; Offner, G. D.; Keutmann, H. T.: The amount

[158962]of MUC5B in cervical mucus peaks at midcycle. J. Clin. Endocr. Metab. 86:

[158963]594–600, 2001.

[158964]

[158965]

[158966]15567.

[158967]Keates, A. C.; Nunes, D. P.; Afdhal, N. H.; Troxler, R. F.;
Offner,

[158968]G. D.: Molecular cloning of a major human gall bladder
mucin: complete

[158969]C-terminal sequence and genomic organization of
MUC5B. Biochem. J. 324:

[158970]295–303, 1997.

[158971]

[158972]

[158973]15568.

[158974]Troxler, R. F.; Offner, G. D.; Zhang, F.; Iontcheva, I.; Op-
penheim,

[158975]F. G.: Molecular cloning of a novel high molecular weight
mucin (MG1)

[158976]from human sublingual gland. Biochem. Biophys. Res.
Commun. 217:

[158977]1112–1119, 1995.

[158978]

[158979]

[158980]15569.

[158981]Birck, C.; Poch, O.; Romier, C.; Ruff, M.; Mengus, G.; Lavi-
gne,

[158982]A.-C.; Davidson, I.; Moras, D.: Human TAFII28 and TAFII18
interact

[158983]through a histone fold encoded by atypical evolutionary conserved

[158984]motifs also found in the SPT3 family. Cell 94: 239–249, 1998.

[158985]

[158986]

[158987]15570.

[158988]Mengus, G.; May, M.; Jacq, X.; Staub, A.; Tora, L.; Chambon, P.;

[158989]Davidson, I.: Cloning and characterization of hTAFII18, hTAFII20

[158990]and hTAFII28: three subunits of the human transcription factor TFIID. EMBO

[158991]J. 14: 1520–1531, 1995.

[158992]

[158993]

[158994]15571.

[158995]Waisfisz, Q.; Saar, K.; Morgan, N. V.; Altay, C.; Leegwater, P.

[158996]A.; de Winter, J. P.; Komatsu, K.; Evans, G. R.; Wegner,

R.-D.; Reis,

[158997]A.; Joenje, H.; Arwert, F.; Mathew, C. G.; Pronk, J. C.; Dig-
weed,

[158998]M.: The Fanconi anemia group E gene, FANCE, maps to
chromosome 6p. Am.

[158999]J. Hum. Genet. 64: 1400–1405, 1999.

[159000]

[159001]

[159002]15572.

[159003]Wegner, R.-D.; Henrichs, I.; Joenje, H.; Schroeder-Kurth,
T.:

[159004]Fanconi anemia complementation group E: clinical and cy-
togenetic data

[159005]of the first patient. Clin. Genet. 50: 479–482, 1996.

[159006]

[159007]

[159008]15573.

[159009]Green, P.; Lipman, D.; Hillier, L.; Waterston, R.; States, D.;

[159010]Claverie, J.-M.: Ancient conserved regions in new gene se-
quences

[159011]and the protein databases. Science 259: 1711–1716,
1993.

[159012]

[159013]

[159014]15574.

[159015]Schwartz, F.; Eisenman, R.; Knoll, J.; Gessler, M.; Bruns, G.:

[159016]cDNA sequence, genomic organization, and evolutionary
conservation

[159017]of a novel gene from the WAGR region. Genomics 29:
526–532, 1995.

[159018]

[159019]

[159020]15575.

[159021]Schwartz, F.; Neve, R.; Eisenman, R.; Gessler, M.; Bruns,
G.:

[159022]A WAGR region gene between PAX–6 and FSHB expressed
in fetal brain. Hum.

[159023]Genet. 94: 658–664, 1994.

[159024]

[159025]

[159026]15576.

[159027]Krishnan, B. R.; Jamry, I.; Chaplin, D. D.: Feature mapping
of

[159028]the HLA class I region: localization of the POU5F1 and
TCF19 genes. Genomics 30:

[159029]53–58, 1995.

[159030]

[159031]

[159032]15577.

[159033]Ku, D.–H.; Chang, C.; Koniecki, J.; Cannizzaro, L. A.;
Boghosian–Sell,

[159034]L.; Alder, H.; Baserga, R.: A new growth–regulated com–
plementary

[159035]DNA with the sequence of a putative trans–activating fac–
tor. Cell

[159036]Growth Differ. 2: 179–186, 1991.

[159037]

[159038]

[159039]15578.

[159040]Joseph, R. E.; Walker, J.; Norris, F. A.: Assignment of the

inositol

[159041]polyphosphate 4-phosphatase type I gene (INPP4A) to human chromosome

[159042]band 2q11.2 by in situ hybridization. Cytogenet. Cell Genet. 87:

[159043]276-277, 1999.

[159044]

[159045]

[159046]15579.

[159047]Norris, F. A.; Auethavekiat, V.; Majerus, P. W.: The isolation

[159048]and characterization of cDNA encoding human and rat brain inositol

[159049]polyphosphate 4-phosphatase. J. Biol. Chem. 270: 16128-16133, 1995.

[159050]

[159051]

[159052]15580.

[159053]van den Maagdenberg, A. M. J. M.; van den Hurk, H. H.; Olde Weghuis,

[159054]D.; Wieringa, B.; Geurts van Kessel, A.; Hendriks, W. J. A.

J.: Assignment

[159055]of the human protein tyrosine phosphatase epsilon
(PTPRE) gene to

[159056]chromosome 10q26 by fluorescence in situ hybridization.

Genomics 30:

[159057]128–129, 1995.

[159058]

[159059]

[159060]15581.

[159061]Kim, D. S.; Jung, H.–H.; Park, S.–H.; Chin, H.: Isolation and

[159062]characterization of the 5–prime–upstream region of the
human N–type

[159063]calcium channel alpha–1B subunit gene: chromosomal lo-
calization and

[159064]promoter analysis. J. Biol. Chem. 272: 5098–5104, 1997.

[159065]

[159066]

[159067]15582.

[159068]Williams, M. E.; Brust, P. F.; Feldman, D. H.; Patthi, S.;
Simerson,

- [159069]S.; Maroufi, A.; McCue, A. F.; Velicelebi, G.; Ellis, S. B.; Harpold,
[159070]M. M.: Structure and functional expression of an omega-conotoxin-sensitive
[159071]human N-type calcium channel. Science 257: 389–395, 1992.
- [159072]
- [159073]
- [159074]15583.
- [159075]Yamazaki, K.; Oki, T.; Tanaka, I.: Locations of human genes for
[159076]alpha(1A), alpha(1B), and alpha(1E) calcium channels determined by
[159077]radiation hybrid mapping. J. Hered. 89: 269–271, 1998.
- [159078]
- [159079]
- [159080]15584.
- [159081]Azim, A. C.; Knoll, J. H. M.; Marfatia, S. M.; Peel, D. J.; Bryant,
[159082]P. J.; Chishti, A. H.: DLG1: chromosome location of the closest human

[159083]homologue of the Drosophila discs large tumor suppressor gene. Genomics 30:

[159084]613–616, 1995.

[159085]

[159086]

[159087]15585.

[159088]Burgess, D. L.; Rafael, J. A.; Meisler, M. H.; Chamberlain, J.

[159089]S.: Dlgh1, a mouse homolog of the Drosophila discs-large gene, is

[159090]located on chromosome 16. Mammalian Genome 7:
623–624, 1996.

[159091]

[159092]

[159093]15586.

[159094]Hanada, T.; Lin, L.; Chandy, K. G.; Oh, S. S.; Chishti, A. H.:

[159095]Human homologue of the Drosophila discs large tumor suppressor binds

[159096]to p56lck tyrosine kinase and Shaker type Kv1.3 potassium channel

[159097]in T lymphocytes. J. Biol. Chem. 272: 26899–26904,

1997.

[159098]

[159099]

[159100]15587.

[159101]Lue, R. A.; Marfatia, S. M.; Branton, D.; Chishti, A. H.:

Cloning

[159102]and characterization of hdlg: the human homologue of the

Drosophila

[159103]discs large tumor suppressor binds to protein 4.1. Proc.

Nat. Acad.

[159104]Sci. 91: 9818–9822, 1994.

[159105]

[159106]

[159107]15588.

[159108]Mori, K.; Iwao, K.; Miyoshi, Y.; Nakagawara, A.; Kofu, K.;

Akiyama,

[159109]T.; Arita, N.; Hayakawa, T.; Nakamura, Y.: Identification of

brain-specific

[159110]splicing variants of the hDLG1 gene and altered splicing in

neuroblastoma

[159111]cell lines. J. Hum. Genet. 43: 123–127, 1998.

[159112]

[159113]

[159114]15589.

[159115]Peters, L. L.; Ciciotte, S. L.; Lin, L.; Chishti, A. H.: The
mouse

[159116]homolog of the Drosophila discs large tumor suppressor
gene maps to

[159117]chromosome 16. Mammalian Genome 7: 619–620, 1996.

[159118]

[159119]

[159120]15590.

[159121]Bertaux, F.; Sharp, A. H.; Ross, C. A.; Lehrach, H.; Bates, G.

[159122]P.; Wanker, E.: HAP1–huntingtin interactions do not con-
tribute to

[159123]the molecular pathology in Huntington's disease trans-
genic mice. FEBS

[159124]Lett. 426: 229–232, 1998.

[159125]

[159126]

[159127]15591.

[159128]Engelender, S.; Sharp, A. H.; Colomer, V.; Tokito, M. K.;
Lanahan,

[159129]A.; Worley, P.; Holzbaur, E. L. F.; Ross, C. A.: Huntingtin-
associated

[159130]protein 1 (HAP1) interacts with the p150(Glued) subunit of
dynactin. Hum.

[159131]Molec. Genet. 6: 2205–2212, 1997.

[159132]

[159133]

[159134]15592.

[159135]Li, S.–H.; Hosseini, S. H.; Gutekunst, C.–A.; Hersch, S. M.;
Ferrante,

[159136]R. J.; Li, X.–J.: A human HAP1 homologue: cloning, expres-
sion, and

[159137]interaction with huntingtin. J. Biol. Chem. 273:
19220–19227, 1998.

[159138]

[159139]

[159140]15593.

[159141]Li, X.-J.; Li, S.-H.; Sharp, A. H.; Nucifora, F. C., Jr.;

Schilling,

[159142]G.; Lanahan, A.; Worley, P.; Snyder, S. H.; Ross, C. A.: A

huntingtin-associated

[159143]protein enriched in brain with implications for pathology.

Nature 378:

[159144]398–402, 1995.

[159145]

[159146]

[159147]15594.

[159148]Nasir, J.; Duan, K.; Nichol, K.; Engelender, S.; Ashworth, R.;

[159149]Colomer, V.; Thomas, S.; Distèche, C. M.; Hayden, M. R.;

Ross, C.

[159150]A.: Gene structure and map location of the murine homolog of the

[159151]Huntington-associated protein, Hap1. Mammalian

Genome 9: 565–570,

[159152]1998.

[159153]

[159154]

[159155]15595.

[159156]Nasir, J.; Lafuente, M.-J.; Duan, K.; Colomer, V.; Engelen-
der,

[159157]S.; Ingersoll, R.; Margolis, R. L.; Ross, C. A.; Hayden, M. R.:
Human

[159158]huntingtin-associated protein (HAP-1) gene: genomic or-
ganisation and

[159159]an intragenic polymorphism. Gene 254: 181-187, 2000.

[159160]

[159161]

[159162]15596.

[159163]Broccoli, D.; Chong, L.; Oelmann, S.; Fernald, A. A.;
Marziliano,

[159164]N.; van Steensel, B.; Kipling, D.; Le Beau, M. M.; de Lange,
T.:

[159165]Comparison of the human and mouse genes encoding the
telomeric protein,

[159166]TRF1: chromosomal localization, expression and con-
served protein domains. Hum.

[159167]Molec. Genet. 6: 69-76, 1997.

[159168]

[159169]

[159170]15597.

[159171]Chong, L.; van Steensel, B.; Broccoli, D.; Erdjument-Bromage, H.;

[159172]Hanish, J.; Tempst, P.; de Lange, T.: A human telomeric protein. Science 270:

[159173]1663–1667, 1995.

[159174]

[159175]

[159176]15598.

[159177]Fairall, L.; Chapman, L.; Moss, H.; de Lange, T.; Rhodes, D.:

[159178]Structure of the TRFH dimerization domain of the human telomeric proteins

[159179]TRF1 and TRF2. Molec. Cell 8: 351–361, 2001.

[159180]

[159181]

[159182]15599.

[159183]Griffith, J. D.; Comeau, L.; Rosenfield, S.; Stansel, R. M.;

Bianchi,

[159184]A.; Moss, H.; de Lange, T.: Mammalian telomeres end in a
large duplex

[159185]loop. Cell 97: 503–514, 1999.

[159186]

[159187]

[159188]15600.

[159189]Kim, S.; Kaminker, P.; Campisi, J.: TIN2, a new regulator of
telomere

[159190]length in human cells. Nature Genet. 23: 405–412, 1999.

[159191]

[159192]

[159193]15601.

[159194]Marcand, S.; Gilson, E.; Shore, D.: A protein–counting
mechanism

[159195]for telomere length regulation in yeast. Science 275:
986–990, 1997.

[159196]

[159197]

[159198]15602.

[159199]Okabe, J.; Eguchi, A.; Masago, A.; Hayakawa, T.; Nakanishi, M.

[159200]: TRF1 is a critical trans-acting factor required for de novo telomere

[159201]formation in human cells. Hum. Molec. Genet. 9: 2639–2650, 2000.

[159202]

[159203]

[159204]15603.

[159205]Young, A. C.; Chavez, M.; Giambernardi, T. A.; Mattern, V.; McGill,

[159206]J. R.; Harris, J. M.; Sarosdy, M. F.; Patel, P.; Sakaguchi, A. Y.

[159207]: Organization and expression of human telomere repeat binding factor

[159208]genes. Somat. Cell Molec. Genet. 23: 275–286, 1997.

[159209]

[159210]

[159211]15604.

[159212]Zakian, V. A.: Telomeres: beginning to understand the

end. Science 270:

[159213]1601–1607, 1995.

[159214]

[159215]

[159216]15605.

[159217]Brotsky, G.; Otterson, G. A.; Parry, B. B.; Hart, I.; Patterson,
son,

[159218]D.; Kaye, F. J.: Localization of STCH to human chromosome 21q11.1. Genomics 30:

[159219]627–628, 1995.

[159220]

[159221]

[159222]15606.

[159223]Otterson, G. A.; Flynn, G. C.; Kratzke, R. A.; Coxon, A.;
Johnston,

[159224]P. G.; Kaye, F. J.: Stch encodes the 'ATPase core' of a mitochondrial

[159225]stress70 protein. EMBO J. 13: 1216–1225, 1994.

[159226]

[159227]

[159228]15607.

[159229]Reeves, R. H.; Rue, E.; Yu, J.; Kao, F.-T.: Stch maps to
mouse

[159230]chromosome 16, extending the conserved synteny with
human chromosome

[159231]21. Genomics 49: 156–157, 1998.

[159232]

[159233]

[159234]15608.

[159235]Nielsen, P. J.; Rochelle, J. M.; Seldin, M. F.: The functional

[159236]genes for protein synthesis initiation factor 4A1 and 4A2
map to

[159237]mouse chromosomes 11 and 16. Mammalian Genome 4:
185–186, 1993.

[159238]

[159239]

[159240]15609.

[159241]Nielsen, P. J.; Trachsel, H.: The mouse protein synthesis

initiation

[159242]factor 4A gene family includes two related functional
genes which

[159243]are differentially expressed. EMBO J. 7: 2097–2105, 1988.

[159244]

[159245]

[159246]15610.

[159247]Sudo, K.; Takahashi, E.; Nakamura, Y.: Isolation and map-
ping of

[159248]the human EIF4A2 gene homologous to the murine pro-
tein synthesis initiation

[159249]factor 4A-II gene Eif4a2. Cytogenet. Cell Genet. 71:
385–388, 1995.

[159250]

[159251]

[159252]15611.

[159253]Gladyshev, V. N.; Jeang, K.-T.; Stadtman, T. C.: Selenocys-
teine,

[159254]identified as the penultimate C-terminal residue in human
T-cell thioredoxin

[159255]reductase, corresponds to TGA in the human placental

gene. Proc.

[159256]Nat. Acad. Sci. 93: 6146–6151, 1996.

[159257]

[159258]

[159259]15612.

[159260]Zhang, K.; Lindsberg, P. J.; Tatlisumak, T.; Kaste, M.;
Olsen,

[159261]H. S.; Andersson, L. C.: Stanniocalcin: a molecular guard
of neurons

[159262]during cerebral ischemia. Proc. Nat. Acad. Sci. 97:
3637–3642, 2000.

[159263]

[159264]

[159265]15613.

[159266]Gillett, G. T.; Fox, M. F.; Rowe, P. S. N.; Casimir, C. M.;
Povey,

[159267]S.: Mapping of human non-muscle type cofilin (CFL1) to
chromosome

[159268]11q13 and muscle-type cofilin (CFL2) to chromosome 14.
Ann. Hum.

[159269]Genet. 60: 201–211, 1996.

[159270]

[159271]

[159272]15614.

[159273]Ogawa, K.; Tashima, M.; Yumoto, Y.; Okuda, T.; Sawada, H.; Okuma,

[159274]M.; Maruyama, Y.: Coding sequence of human placenta cofilin cDNA. Nuc.

[159275]Acids Res. 18: 7169 only, 1990.

[159276]

[159277]

[159278]15615.

[159279]Ono, S.; Minami, N.; Abe, H.; Obinata, T.: Characterization of

[159280]a novel cofilin isoform that is predominantly expressed in mammalian

[159281]skeletal muscle. J. Biol. Chem. 269: 15280–15286, 1994.

[159282]

[159283]

[159284]15616.

[159285]Georgakopoulos, T.; Thireos, G.: Two distinct yeast transcriptional

[159286]activators require the function of the GCN5 protein to promote normal

[159287]levels of transcription. EMBO J. 11: 4145–4152, 1992.

[159288]

[159289]

[159290]15617.

[159291]Inoue, M.; Isomura, M.; Ikegawa, S.; Fujiwara, T.; Shin, S.; Moriya,

[159292]H.; Nakamura, Y.: Isolation and characterization of a human cDNA

[159293]clone (GCN5L1) homologous to GCN5, a yeast transcription factor. Cytogenet.

[159294]Cell Genet. 73: 134–136, 1996.

[159295]

[159296]

[159297]15618.

[159298]Emahazion, T.; Beskow, A.; Gyllenstein, U.; Brookes, A. J.: Intron

[159299]based radiation hybrid mapping of 15 complex I genes of
the human

[159300]electron transport chain. Cytogenet. Cell Genet. 82:
115–119, 1998.

[159301]

[159302]

[159303]15619.

[159304]Gu, J. Z.; Lin, X.; Wells, D. E.: The human B22 subunit of
the

[159305]NADH–ubiquinone oxidoreductase maps to the region of
chromosome 8

[159306]involved in branchio–oto–renal syndrome. Genomics 35:
6–10, 1996.

[159307]

[159308]

[159309]15620.

[159310]Lin, X.; Wells, D. E.; Kimberling, W. J.; Kumar, S.: Human
NDUFB9

[159311]gene: genomic organization and a possible candidate
gene associated

[159312]with deafness disorder mapped to chromosome 8q13.

Hum. Hered. 49:

[159313]75–80, 1999.

[159314]

[159315]

[159316]15621.

[159317]Wilkinson, K. D.; Tashayev, V. L.; O'Connor, L. B.; Larsen, C.

[159318]N.; Kasperek, E.; Pickart, C. M.: Metabolism of the polyubiquitin

[159319]degradation signal: structure, mechanism, and role of isopeptidase

[159320]T. Biochemistry 34: 14535–14546, 1995.

[159321]

[159322]

[159323]15622.

[159324]Byrd, P. J.; Cooper, P. R.; Stankovic, T.; Kullar, H. S.; Watts,

[159325]G. D. J.; Robinson, P. J.; Taylor, M. R.: A gene transcribed from

[159326]the bidirectional ATM promoter coding for a serine rich protein: amino

[159327]acid sequence, structure and expression studies. Hum.

Molec. Genet. 5:

[159328]1785–1791, 1996.

[159329]

[159330]

[159331]15623.

[159332]Chen, X.; Yang, L.; Udar, N.; Liang, T.; Uhrhammer, N.; Xu,
S.;

[159333]Bay, J.–O.; Wang, Z.; Dandakar, S.; Chiplunkar, S.; Klisak,
I.; Telatar,

[159334]M.; Yang, H.; Concannon, P.; Gatti, R. A.: CAND3: a ubiqu-
itously

[159335]expressed gene immediately adjacent and in opposite
transcriptional

[159336]orientation to the ATM gene at 11q23.1. Mammalian
Genome 8: 129–133,

[159337]1997.

[159338]

[159339]

[159340]15624.

[159341]Imai, T.: Personal Communication. Chiba, Japan

9/12/1996.

[159342]

[159343]

[159344]15625.

[159345]Imai, T.; Sugawara, T.; Nishiyama, A.; Shimada, R.; Ohki,
R.; Seki,

[159346]N.; Sagara, M.; Ito, H.; Yamauchi, M.; Hori, T.: The struc-
ture and

[159347]organization of the human NPAT gene. Genomics 42:
388–392, 1997.

[159348]

[159349]

[159350]15626.

[159351]Imai, T.; Yamauchi, M.; Seki, N.; Sugawara, T.; Saito, T.;
Matsuda,

[159352]Y.; Ito, H.; Nagase, T.; Nomura, N.; Hori, T.: Identification
and

[159353]characterization of a new gene physically linked to the
ATM gene. Genome

[159354]Res. 6: 439–447, 1996.

[159355]

[159356]

[159357]15627.

[159358]Borowsky, B.; Hoffman, B. J.: Analysis of a gene encoding
two

[159359]glycine transporter variants reveals alternative promoter
usage and

[159360]a novel gene structure. J. Biol. Chem. 273: 29077–29085,
1998.

[159361]

[159362]

[159363]15628.

[159364]Borowsky, B.; Mezey, E.; Hoffman, B. J.: Two glycine trans-
porter

[159365]variants with distinct localization in the CNS and periph-
eral tissues

[159366]are encoded by a common gene. Neuron 10: 851–863,
1993.

[159367]

[159368]

[159369]15629.

[159370]Jones, E. M. C.; Fernald, A.; Bell, G. I.; Le Beau, M. M.: Assignment
segment

[159371]of SLC6A9 to human chromosome band 1p33 by in situ
hybridization. Cytogenet.

[159372]Cell Genet. 71: 211, 1995.

[159373]

[159374]

[159375]15630.

[159376]Kim, K.-M.; Kingsmore, S. F.; Han, H.; Yang-Feng, T. L.;
Godinot,

[159377]N.; Seldin, M. F.; Caron, M. G.; Giros, B.: Cloning of the
human

[159378]glycine transporter type 1: molecular and pharmacological
characterization

[159379]of novel isoform variants and chromosomal localization of
the gene

[159380]in the human and mouse genomes. Molec. Pharmacol. 45:
608-617, 1994.

[159381]

[159382]

[159383]15631.

[159384]Freeman, G. J.; Gribben, J. G.; Boussiotis, V. A.; Ng, J. W.;
Restivo,

[159385]V. A.; Lombard, L. A.; Gray, G. S.; Nadler, L. M. Cloning of
B7-2:

[159386]a CTLA-4 counter-receptor that costimulates human T
cell proliferation. Science 262:

[159387]909-911, 1993.

[159388]

[159389]

[159390]15632.

[159391]Jeannin, P.; Magistrelli, G.; Aubry, J.-P.; Caron, G.;
Gauchat,

[159392]J.-F.; Renno, T.; Herbault, N.; Goetsch, L.; Blaecke, A.; Di-
etrich,

[159393]P.-Y.; Bonnefoy, J.-Y.; Delneste, Y.: Soluble CD86 is a cos-
timulatory

[159394]molecule for human T lymphocytes. Immunity 13:
303-312, 2000.

[159395]

[159396]

[159397]15633.

[159398]Jellis, C. L.; Wang, S. S.; Rennert, P.; Borriello, F.; Sharpe,

[159399]A. H.; Green, N. R.; Gray, G. S.: Genomic organization of
the gene

[159400]coding for the costimulatory human B-lymphocyte antigen
B7-2 (CD86). Immunogenetics 42:

[159401]85-89, 1995.

[159402]

[159403]

[159404]15634.

[159405]Arai, Y.; Hosoda, F.; Kobayashi, H.; Arai, K.; Hayashi, Y.;
Kamada,

[159406]N.; Kaneko, Y.; Ohki, M.: The inv(11)(p15q22) chromo-
some translocation

[159407]of de novo and therapy-related myeloid malignancies re-
sults in fusion

[159408]of the nucleoporin gene, NUP98, with the putative RNA
helicase gene,

[159409]DDX10. Blood 89: 3936-3944, 1997.

[159410]

[159411]

[159412]15635.

[159413]Dash, A. B.; Williams, I. R.; Kutok, J. L.; Tomasson, M. H.;

Anastasiadou,

[159414]E.; Lindahl, K.; Li, S.; Van Etten, R. A.; Borrow, J.; Housman,

D.;

[159415]Druker, B.; Gilliland, D. G.: A murine model of CML blast
crisis

[159416]induced by cooperation between BCR/ABL and
NUP98/HOXA9. Proc. Nat.

[159417]Acad. Sci. 99: 7622–7627, 2002.

[159418]

[159419]

[159420]15636.

[159421]Enninga, J.; Levy, D. E.; Blobel, G.; Fontoura, B. M. A.: Role

[159422]of nucleoporin induction in releasing an mRNA nuclear
export block. Science 295:

[159423]1523–1525, 2002.

[159424]

[159425]

[159426]15637.

[159427]Fontoura, B. M.; Blobel, G.; Matunis, M. J.: A conserved
biogenesis

[159428]pathway for nucleoporins: proteolytic processing of a
186-kilodalton

[159429]precursor generates Nup98 and the novel nucleoporin,
Nup96. J. Cell

[159430]Biol. 144: 1097–1112, 1999.

[159431]

[159432]

[159433]15638.

[159434]Hodel, A. E.; Hodel, M. R.; Griffis, E. R.; Hennig, K. A.; Rat-
ner,

[159435]G. A.; Xu, S.; Powers, M. A.: The three-dimensional struc-
ture of

[159436]the autoproteolytic, nuclear pore-targeting domain of the
human nucleoporin

[159437]Nup98. Molec. Cell 10: 347–358, 2002.

[159438]

[159439]

[159440]15639.

[159441]Jaju, R. J.; Fidler, C.; Haas, O. A.; Strickson, A. J.; Watkins,

[159442]F.; Clark, K.; Cross, N. C. P.; Cheng, J.-F.; Aplan, P. D.;

Kearney,

[159443]L.; Boultonwood, J.; Wainscoat, J. S.: A novel gene, NSD1, is

fused

[159444]to NUP98 in the t(5;11)(q35;p15.5) in de novo childhood

acute myeloid

[159445]leukemia. Blood 98: 1264–1267, 2001.

[159446]

[159447]

[159448]15640.

[159449]Jaju, R. J.; Haas, C. A.; Neat, M.; et al: A new recurrent

translocation,

[159450]t(5;11)(q35p15.5), associated with del(5q) in childhood

acute myeloid

[159451]leukemia. Blood 94: 773–780, 1999.

[159452]

[159453]

[159454]15641.

[159455]Mizuno, T.; Kaibuchi, K.; Yamamoto, T.; Kawamura, M.;
Sakoda,

[159456]T.; Fujioka, H.; Matsuura, Y.; Takai, Y.: A stimulatory GDP/
GTP exchange

[159457]protein for smg p21 is active on the post-translationally
processed

[159458]form of c-Ki-ras p21 and rhoA p21. Proc. Nat. Acad. Sci.
88: 6442-6446,

[159459]1991.

[159460]

[159461]

[159462]15642.

[159463]Nakamura, T.; Largaespada, D. A.; Lee, M. P.; Johnson, L.
A.;

[159464]Ohyashiki, K.; Toyama, K.; Chen, S. J.; Willman, C. L.;
Chen, I.-M.;

[159465]Feinberg, A. P.; Jenkins, N. A.; Copeland, N. G.; Shaugh-
nessy, J.

[159466]D., Jr.: Fusion of the nucleoporin gene NUP98 to HOXA9
by the chromosome

[159467]translocation t(7;11)(p15;p15) in human myeloid
leukaemia. Nature

[159468]Genet. 12: 154–158, 1996.

[159469]

[159470]

[159471]15643.

[159472]Radu, A.; Moore, M. S.; Blobel, G.: The peptide repeat domain

[159473]of nucleoporin Nup98 functions as a docking site in transport across

[159474]the nuclear pore complex. Cell 81: 215–222, 1995.

[159475]

[159476]

[159477]15644.

[159478]Rosati, R.; La Starza, R.; Veronese, A.; Aventin, A.; Schwienbacher,

[159479]C.; Vallespi, T.; Negrini, M.; Martelli, M. F.; Mecucci, C.: NUP98

[159480]is fused to the NSD3 gene in acute myeloid leukemia associated with

[159481]t(8;11)(p11.2;p15). Blood 99: 3857–3860, 2002.

[159482]

[159483]

[159484]15645.

[159485]von Kobbe, C.; van Deursen, J. M. A.; Rodrigues, J. P.; Sitterlin,

[159486]D.; Bachi, A.; Wu, X.; Wilm, M.; Carmo-Fonseca, M.; Izauralde, E.

[159487]: Vesicular stomatitis virus matrix protein inhibits host cell gene

[159488]expression by targeting the nucleoporin Nup98. Molec. Cell 6: 1243–1252,

[159489]2000.

[159490]

[159491]

[159492]15646.

[159493]Cheng, Y.; Austin, S. C.; Rocca, B.; Koller, B. H.; Coffman, T.

[159494]M.; Grosser, T.; Lawson, J. A.; FitzGerald, G. A.: Role of prostacyclin

[159495]in the cardiovascular response to thromboxane A₂. Science 296: 539–541,

[159496]2002.

[159497]

[159498]

[159499]15647.

[159500]Fuse, I.; Mito, M.; Hattori, A.; Higuchi, W.; Shibata, A.;
Ushikubi,

[159501]F.; Okuma, M.; Yahata, K.: Defective signal transduction
induced

[159502]by thromboxane A₂ in a patient with a mild bleeding dis-
order: impaired

[159503]phospholipase C activation despite normal phospholipase
A₂ activation. Blood 81:

[159504]994–1000, 1993.

[159505]

[159506]

[159507]15648.

[159508]Hirata, M.; Hayashi, Y.; Ushikubi, F.; Yokota, Y.; Kageyama,
R.;

[159509]Nakanishi, S.; Narumiya, S.: Cloning and expression of
cDNA for a

[159510]human thromboxane A₂ receptor. Nature 349: 617–620,
1991.

[159511]

[159512]

[159513]15649.

[159514]Hirata, T.; Kakizuka, A.; Ushikubi, F.; Fuse, I.; Okuma, M.;
Narumiya,

[159515]S.: Arg60-to-leu mutation of the human thromboxane A2
receptor in

[159516]a dominantly inherited bleeding disorder. J. Clin. Invest.
94: 1662-1667,

[159517]1994.

[159518]

[159519]

[159520]15650.

[159521]Watanabe, O.; Maruyama, I.; Arimura, K.; Kitajima, I.;
Arimura,

[159522]H.; Hanatani, M.; Matsuo, K.; Arisato, T.; Osame, M.: Over-
production

[159523]of vascular endothelial growth factor/vascular permeabil-
ity factor

[159524]is causative in Crow-Fukase (POEMS) syndrome. Muscle
Nerve 21: 1390-1397,

[159525]1998.

[159526]

[159527]

[159528]15651.

[159529]Wei, M.-H.; Popescu, N. C.; Lerman, M. I.; Merrill, M. J.; Zimonjic,

[159530]D. B.: Localization of the human vascular endothelial growth factor

[159531]gene, VEGF, at chromosome 6p12. Hum. Genet. 97: 794–797, 1996.

[159532]

[159533]

[159534]15652.

[159535]Wong, A. K.; Alfert, M.; Castrillon, D. H.; Shen, Q.; Holash,

[159536]J.; Yancopoulos, G. D.; Chin, L.: Excessive tumor-elaborated VEGF

[159537]and its neutralization define a lethal paraneoplastic syndrome. Proc.

[159538]Nat. Acad. Sci. 98: 7481–7486, 2001.

[159539]

[159540]

[159541]15653.

[159542]Wulff, C.; Wilson, H.; Lague, P.; Duncan, W. C.; Armstrong, D.

[159543]G.; Fraser, H. M.: Angiogenesis in the human corpus luteum: localization

[159544]and changes in angiopoietins, Tie-2, and vascular endothelial growth

[159545]factor messenger ribonucleic acid. J. Clin. Endocr. Metab. 85: 4302-4309,

[159546]2000.

[159547]

[159548]

[159549]15654.

[159550]Ylikorkala, A.; Rossi, D. J.; Korsisaari, N.; Luukko, K.; Alitalo,

[159551]K.; Henkemeyer, M.; Makela, T. P.: Vascular abnormalities and deregulation

[159552]of VEGF in Lkb1-deficient mice. Science 293: 1323-1326, 2001.

[159553]

[159554]

[159555]15655.

[159556]Bodner, M.; Fridkin, M.; Gozes, I.: Coding sequences for
vasoactive

[159557]intestinal peptide and PHM-27 peptide are located on two
adjacent

[159558]exons in the human genome. Proc. Nat. Acad. Sci. 82:
3548-3551,

[159559]1985.

[159560]

[159561]

[159562]15656.

[159563]Delgado, M.; Abad, C.; Martinez, C.; Leceta, J.; Gomariz, R.
P.

[159564]: Vasoactive intestinal peptide prevents experimental
arthritis by

[159565]downregulating both autoimmune and inflammatory com-
ponents of the

[159566]disease. Nature Med. 7: 563-568, 2001.

[159567]

[159568]

[159569]15657.

[159570]Gotoh, E.; Yamagami, T.; Yamamoto, H.; Okamoto, H.:

Chromosomal

[159571]assignment of human VIP/PHM-27 gene to 6q26-q27 re-
gion by spot blot

[159572]hybridization and in situ hybridization. Biochem. Int. 17:
555-562,

[159573]1988.

[159574]

[159575]

[159576]15658.

[159577]Gozes, I.; Avidor, R.; Yahav, Y.; Katznelson, D.; Croce, C.
M.;

[159578]Huebner, K.: The gene encoding vasoactive intestinal pep-
tide is located

[159579]on human chromosome 6p21-6qter. Hum. Genet. 75:
41-44, 1987.

[159580]

[159581]

[159582]15659.

[159583]Gozes, I.; Nakai, H.; Byers, M.; Avidor, R.; Weinstein, Y.;
Shani,

[159584]Y.; Shows, T. B.: Sequential expression in the nervous sys-
tem of

[159585]C-MYB and VIP genes, located in human chromosomal re-
gion 6q24. Somat.

[159586]Cell Molec. Genet. 13: 305-313, 1987.

[159587]

[159588]

[159589]15660.

[159590]Heinz-Erian, P.; Dey, R. D.; Flux, M.; Said, S. I.: Deficient

[159591]vasoactive intestinal peptide innervation in sweat glands
of cystic

[159592]fibrosis patients. Science 229: 1407-1408, 1985.

[159593]

[159594]

[159595]15661.

[159596]Itoh, N.; Obata, K.; Yanaihara, N.; Okamoto, H.: Human

preprovasoactive

[159597]intestinal polypeptide contains a novel PHI-27-like peptide, PHM-27. Nature 304:

[159598]547-549, 1983.

[159599]

[159600]

[159601]15662.

[159602]Linder, S.; Barkhem, T.; Norberg, A.; Persson, H.; Schalling, M.;

[159603]Hokfelt, T.; Magnusson, G.: Structure and expression of the gene

[159604]encoding the vasoactive intestinal peptide precursor. Proc. Nat.

[159605]Acad. Sci. 84: 605-609, 1987.

[159606]

[159607]

[159608]15663.

[159609]Omary, M. B.; Kagnoff, M. F.: Identification of nuclear receptors

[159610]for VIP on a human colonic adenocarcinoma cell line. Sci-

ence 238:

[159611]1578–1581, 1987.

[159612]

[159613]

[159614]15664.

[159615]Couvineau, A.; Rouyer–Fessard, C.; Darmoul, D.; Maoret, J.–J.;

[159616]Carrero, I.; Ogier–Denis, E.; Laburthe, M.: Human intestinal VIP

[159617]receptor: cloning and functional expression of two cDNA encoding proteins

[159618]with different N–terminal domains. Biochem. Biophys. Res. Commun. 200:

[159619]769–776, 1994.

[159620]

[159621]

[159622]15665.

[159623]Hashimoto, H.; Nishino, A.; Shintani, N.; Hagihara, N.; Copeland,

[159624]N. G.; Jenkins, N. A.; Yamamoto, K.; Matsuda, T.; Ishihara,

T.; Nagata,

[159625]S.; Baba, A.: Genomic organization and chromosomal location of the

[159626]mouse vasoactive intestinal polypeptide 1 (VPAC-1) receptor. Genomics 58:

[159627]90-93, 1999.

[159628]

[159629]

[159630]15666.

[159631]Sreedharan, S. P.; Huang, J.-X.; Cheung, M.-C.; Goetzl, E. J.:

[159632]Structure, expression, and chromosomal localization of the type I

[159633]human vasoactive intestinal peptide receptor gene. Proc. Nat. Acad.

[159634]Sci. 92: 2939-2943, 1995.

[159635]

[159636]

[159637]15667.

[159638]Sreedharan, S. P.; Patel, D. R.; Huang, J.-X.; Goetzl, E. J.:

[159639]Cloning and functional expression of a human neuroen-
docrine vasoactive

[159640]intestinal peptide receptor. Biochem. Biophys. Res. Com-
mun. 193:

[159641]546–553, 1993.

[159642]

[159643]

[159644]15668.

[159645]Sreedharan, S. P.; Robichon, A.; Peterson, K. E.; Goetzl, E.
J.

[159646]: Cloning and expression of the human vasoactive intesti-
nal peptide

[159647]receptor. Proc. Nat. Acad. Sci. 88: 4986–4990, 1991.

[159648]

[159649]

[159650]15669.

[159651]Ueno, S.; Maruki, Y.; Nakamura, M.; Tomemori, Y.; Kamae,
K.; Tanabe,

[159652]H.; Yamashita, Y.; Matsuda, S.; Kaneko, S.; Sano, A.: The
gene encoding

[159653]a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. Nature

[159654]Genet. 28: 121-122, 2001.

[159655]

[159656]

[159657]15670.

[159658]Hirata, T.; Ushikubi, F.; Kakizuka, A.; Okuma, M.; Narumiya, S.

[159659]: Two thromboxane A(2) receptor isoforms in human platelets: opposite

[159660]coupling to adenylyl cyclase with different sensitivity to arg60-to-leu

[159661]mutation. J. Clin. Invest. 97: 949-956, 1996.

[159662]

[159663]

[159664]15671.

[159665]Nusing, R. M.; Hirata, M.; Kakizuka, A.; Eki, T.; Ozawa, K.; Narumiya,

[159666]S.: Characterization and chromosomal mapping of the human thromboxane

[159667]A2 receptor gene. J. Biol. Chem. 268: 25253–25259,
1993.

[159668]

[159669]

[159670]15672.

[159671]Schwengel, D. A.; Nouri, N.; Meyers, D. A.; Levitt, R. C.:
Linkage

[159672]mapping of the human thromboxane A2 receptor
(TBXA2R) to chromosome

[159673]19p13.3 using transcribed 3–prime untranslated DNA se-
quence polymorphisms. Genomics 18:

[159674]212–215, 1993.

[159675]

[159676]

[159677]15673.

[159678]Thomas, D. W.; Mannon, R. B.; Mannon, P. J.; Latour, A.;
Oliver,

[159679]J. A.; Hoffman, M.; Smithies, O.; Koller, B. H.; Coffman, T.
M.:

[159680]Coagulation defects and altered hemodynamic responses
in mice lacking

[159681]receptors for thromboxane A(2). J. Clin. Invest. 102:

1994–2001,

[159682]1998.

[159683]

[159684]

[159685]15674.

[159686]Unoki, M.; Furuta, S.; Onouchi, Y.; Watanabe, O.; Doi, S.;

Fujiwara,

[159687]H.; Miyatake, A.; Fujita, K.; Tamari, M.; Nakamura, Y.: As-

sociation

[159688]studies of 33 single nucleotide polymorphisms (SNPs) in

29 candidate

[159689]genes for bronchial asthma: positive association a T924C

polymorphism

[159690]in the thromboxane A2 receptor gene. Hum. Genet. 106:

440–446, 2000.

[159691]

[159692]

[159693]15675.

[159694]Ushikubi, F.; Nakajima, M.; Hirata, M.; Okuma, M.; Fuji-

wara, M.;

[159695]Narumiya, S.: Purification of the thromboxane

A₂/prostaglandin H₂

[159696]receptor from human blood platelets.. J. Biol. Chem. 264:

16496–16501,

[159697]1989.

[159698]

[159699]

[159700]15676.

[159701]Ushikubi, F.; Okuma, M.; Kanaji, K.; Sugiyama, T.;

Ogorochi, T.;

[159702]Narumiya, S.; Uchino, H.: Hemorrhagic thrombocytopathy

with platelet

[159703]thromboxane A₂ abnormality: defective signal transduc-

tion with normal

[159704]binding activity. Thromb. Haemost. 57: 158–164, 1987.

[159705]

[159706]

[159707]15677.

[159708]Ades, E. W.; Zwerner, R. K.; Acton, R. T.; Balch, C. M.: Iso-

lation

[159709]and partial characterisation of the human homologue of
Thy-1. J.

[159710]Exp. Med. 151: 400-406, 1980.

[159711]

[159712]

[159713]15678.

[159714]Bonewald, L.; Ades, E. W.; Tung, E.; Marchalonis, J. J.;
Wang,

[159715]A. C.: Biochemical characterization of human Thy-1. J. Im-
munogenet. 11:

[159716]283-296, 1984.

[159717]

[159718]

[159719]15679.

[159720]Gatti, R. A.; Lathrop, G. M.; Salser, W.; Silver, J.; Lalouel,

[159721]J. M.; White, R.: Location of Thy-1 with respect to a pri-
mary linkage

[159722]map of chromosome 11q. (Abstract) Cytogenet. Cell
Genet. 46: 618

[159723]only, 1987.

[159724]

[159725]

[159726]15680.

[159727]Gatti, R. A.; Shaked, R.; Wei, S.; Koyama, M.; Salser, W.;
Silver,

[159728]J.: DNA polymorphism in the human Thy-1 gene. Hum.
Immun. 22:

[159729]145-150, 1988.

[159730]

[159731]

[159732]15681.

[159733]Greenspan, R. J.; O'Brien, M. C.: Genetic evidence for the
role

[159734]of Thy-1 in neurite outgrowth in the mouse. J. Neuro-
genet. 5: 25-36,

[159735]1989.

[159736]

[159737]

[159738]15682.

[159739]Grzeschik, K.-H.; Kazazian, H. H.: Report of the committee
on

[159740]the genetic constitution of chromosomes 10, 11, and 12.
Cytogenet.

[159741]Cell Genet. 40: 179–205, 1985.

[159742]

[159743]

[159744]15683.

[159745]Letarte–Muirhead, M.; Barclay, A. N.; Williams, A. F.: Purifi-
cation

[159746]of the Thy–1 molecule, a major cell surface glycoprotein
of rat thymocytes.

[159747]Biochem. J. 151: 685–697, 1975.

[159748]

[159749]

[159750]15684.

[159751]Mansour, M. H.; Negm, H. I.; Cooper, E. L.: Thy–1 evolu-
tion. Dev.

[159752]Comp. Immun. 11: 3–15, 1987.

[159753]

[159754]

[159755]15685.

[159756]McKenzie, J. L.; Fabre, J. W.: Human Thy-1: unusual localization

[159757]and possible functional significance in lymphoid tissues. J. Immun. 126:

[159758]843-850, 1981.

[159759]

[159760]

[159761]15686.

[159762]Morris, R.: Thy-1 in developing nervous tissue. Dev. Neurosci. 7:

[159763]133-160, 1985.

[159764]

[159765]

[159766]15687.

[159767]Raff, M. C.: Surface antigenic markers for distinguishing T and

[159768]B lymphocytes in mice. Transplant. Rev. 6: 52-80, 1971.

[159769]

[159770]

[159771]15688.

[159772]Rettig, W. J.; Dracopoli, N. C.; Chesa, P. G.; Spengler, B. A.;

[159773]Beresford, H. R.; Davies, P.; Biedler, J. L.; Old, L. J.: Role of

[159774]human chromosome 11 in determining surface antigenic
phenotype of

[159775]normal and malignant cells: somatic cell genetic analysis
of eight

[159776]antigens, including putative human Thy-1. J. Exp. Med.
162: 1603-1619,

[159777]1985.

[159778]

[159779]

[159780]15689.

[159781]Rettig, W. J.; Dracopoli, N. C.; Silver, J.; Old, L. J.: Human

[159782]THY-1: regional mapping on chromosome 11 and com-
parison with other

[159783]chromosome 11-encoded cell surface glycoproteins.

(Abstract) Cytogenet.

[159784]Cell Genet. 40: 731 only, 1985.

[159785]

[159786]

[159787]15690.

[159788]Rettig, W. J.; Dracopoli, N. C.; Spengler, B. A.; Biedler, J.

[159789]L.; Old, L. J.: Somatic cell genetic analysis of human cell
surface

[159790]antigens, including putative human Thy-1: eight distinct
antigenic

[159791]systems controlled by chromosome 11. (Abstract) Cyto-
genet. Cell

[159792]Genet. 40: 732 only, 1985.

[159793]

[159794]

[159795]15691.

[159796]Seki, T.; Spurr, N.; Obata, F.; Goyert, S.; Goodfellow, P.;
Silver,

[159797]J.: The human Thy-1 gene: structure and chromosomal
location. Proc.

[159798]Nat. Acad. Sci. 82: 6657–6661, 1985.

[159799]

[159800]

[159801]15692.

[159802]Tse, A. G. D.; Barclay, A. N.; Watts, A.; Williams, A. F.: A

[159803]glycophospholipid tail at the carboxyl terminus of the
Thy-1 glycoprotein

[159804]of neurons and thymocytes. Science 230: 1003–1008,
1985.

[159805]

[159806]

[159807]15693.

[159808]Chen, Z.; Friedrich, G. A.; Soriano, P.: Transcriptional en-
hancer

[159809]factor 1 disruption by a retroviral gene trap leads to heart
defects

[159810]and embryonic lethality in mice. Genes Dev. 8:
2293–2301, 1994.

[159811]

[159812]

[159813]15694.

[159814]Aso, T.; Tsai, P.; Kawaguchi, T.; Menninger, J. C.; Kitajima, S.;

[159815]Yasukochi, Y.; Ward, D. C.; Weissman, S. M.: Assignment of the human

[159816]GTF2F1 gene to chromosome 19p13.3. Genomics 16: 252–253, 1993.

[159817]

[159818]

[159819]15695.

[159820]Aso, T.; Vasavada, H. A.; Kawaguchi, T.; Germino, F. J.; Ganguly,

[159821]S.; Kitajima, S.; Weissman, S. M.; Yasukochi, Y.: Characterization

[159822]of cDNA for the large subunit of the transcription initiation factor

[159823]TFIIF. Nature 355: 461–464, 1992.

[159824]

[159825]

[159826]15696.

[159827]Finkelstein, A.; Kostrub, C. F.; Li, J.; Chavez, D. P.; Wang, B.

[159828]Q.; Fang, S. M.; Greenblatt, J.; Burton, Z. F.: A cDNA encoding RAP74,

[159829]a general initiation factor for transcription by RNA polymerase II. Nature 355:

[159830]464–467, 1992.

[159831]

[159832]

[159833]15697.

[159834]Joliot, V.; Demma, M.; Prywes, R.: Interaction with RAP74 subunit

[159835]of TFIIF is required for transcriptional activation by serum response

[159836]factor. Nature 373: 632–635, 1995.

[159837]

[159838]

[159839]15698.

[159840]Sopta, M.; Burton, Z. F.; Greenblatt, J.: Structure and associated

[159841]DNA-helicase activity of a general transcription initiation factor

[159842]that binds to RNA polymerase II. Nature 341: 410–414, 1989.

[159843]

[159844]

[159845]15699.

[159846]Field, S. J.; Tsai, F.–Y.; Kuo, F.; Zubiaga, A. M.; Kaelin, W.

[159847]G., Jr.; Livingston, D. M.; Orkin, S. H.; Greenberg, M. E.: E2F–1

[159848]functions in mice to promote apoptosis and suppress proliferation. Cell 85:

[159849]549–561, 1996.

[159850]

[159851]

[159852]15700.

[159853]Helin, K.; Lees, J. A.; Vidal, M.; Dyson, N.; Harlow, E.; Fataey,

[159854]A.: A cDNA encoding a pRB-binding protein with properties of the

[159855]transcription factor E2F. Cell 70: 337–350, 1992.

[159856]

[159857]

[159858]15701.

[159859]Irwin, M.; Marin, M. C.; Phillips, A. C.; Seelan, R. S.; Smith,

[159860]D. I.; Liu, W.; Flores, E. R.; Tsai, K. Y.; Jacks, T.; Vousden,
K.

[159861]H.; Kaelin, W. G., Jr.: Role for the p53 homologue p73 in
E2F–1–induced

[159862]apoptosis. Nature 407: 645–648, 2000.

[159863]

[159864]

[159865]15702.

[159866]Jacks, T.; Fazeli, A.; Schmitt, E. M.; Bronson, R. T.; Goodell,

[159867]M. A.; Weinberg, R. A.: Effects of an Rb mutation in the
mouse. Nature 359:

[159868]295–300, 1992.

[159869]

[159870]

[159871]15703.

[159872]Lees, J. A.; Saito, M.; Valentine, M.; Look, T.; Harlow, E.;
Dyson,

[159873]N.; Helin, K.: The retinoblastoma protein binds to a family
of E2F

[159874]transcription factors. Molec. Cell. Biol. 13: 7813–7825,
1993.

[159875]

[159876]

[159877]15704.

[159878]Leone, G.; Sears, R.; Huang, E.; Rempel, R.; Nuckolls, F.;
Park,

[159879]C.–H.; Giangrande, P.; Wu, L.; Saavedra, H. I.; Field, S. J.;
Thompson,

[159880]M. A.; Yang, H.; Fujiwara, Y.; Greenberg, M. E.; Orkin, S.;
Smith,

[159881]C.; Nevins, J. R.: Myc requires distinct E2F activities to in-
duce

[159882]S phase and apoptosis. Molec. Cell 8: 105–113, 2001.

[159883]

[159884]

[159885]15705.

[159886]Lissy, N. A.; Davis, P. K.; Irwin, M.; Kaelin, W. G.; Dowdy, S.

[159887]F.: A common E2F-1 and p73 pathway mediates cell death
induced by

[159888]TCR activation. Nature 407: 642-645, 2000.

[159889]

[159890]

[159891]15706.

[159892]Neuman, E.; Sellers, W. R.; McNeil, J. A.; Lawrence, J. B.;
Kaelin,

[159893]W. G., Jr.: Structure and partial genomic sequence of the
human E2F1

[159894]gene. Gene 173: 163-169, 1996.

[159895]

[159896]

[159897]15707.

[159898]Nevins, J. R.: The Rb/E2F pathway and cancer. Hum.
Molec. Genet. 10:

[159899]699-703, 2001.

[159900]

[159901]

[159902]15708.

[159903]Nevins, J. R.: E2F: a link between the Rb tumor suppressor protein

[159904]and viral oncoproteins. Science 258: 424–429, 1992.

[159905]

[159906]

[159907]15709.

[159908]Ohtani, K.; DeGregori, J.; Nevins, J. R.: Regulation of the cyclin

[159909]E gene by transcription factor E2F1. Proc. Nat. Acad. Sci. 92: 12146–12150,

[159910]1995.

[159911]

[159912]

[159913]15710.

[159914]Arden, K. C.; Boutin, J.–M.; Djiane, J.; Kelly, P. A.; Cavenee,

[159915]W. K.: The receptors for prolactin and growth hormone are

localized

[159916]in the same region of human chromosome 5. Cytogenet.

Cell Genet. 53:

[159917]161–165, 1990.

[159918]

[159919]

[159920]15711.

[159921]Arden, K. C.; Cavenee, W. K.; Boutin, J.–M.; Kelly, P. A.:

The

[159922]genes encoding the receptors for prolactin and growth
hormone map

[159923]to human chromosome 5. (Abstract) Am. J. Hum. Genet.

45 (suppl.):

[159924]A129 only, 1989.

[159925]

[159926]

[159927]15712.

[159928]Boutin, J.–M.; Edery, M.; Shirota, M.; Jolicoeur, C.; Lesueur,

[159929]L.; Ali, S.; Gould, D.; Djiane, J.; Kelly, P. A.: Identification

[159930]of a cDNA encoding a long form of prolactin receptor in
human hepatoma

[159931]and breast cancer cells. Molec. Endocr. 3: 1455–1461,
1989.

[159932]

[159933]

[159934]15713.

[159935]Cunningham, B. C.; Bass, S.; Fuh, G.; Wells, J. A.: Zinc me-
diation

[159936]of the binding of human growth hormone to the human
prolactin receptor. Science 250:

[159937]1709–1712, 1990.

[159938]

[159939]

[159940]15714.

[159941]Glasow, A.; Horn, L.–C.; Taymans, S. E.; Stratakis, C. A.;
Kelly,

[159942]P. A.; Kohler, U.; Gillespie, J.; Vonderhaar, B. K.; Bornstein,
S.

[159943]R.: Mutational analysis of the PRL receptor gene in human
breast

[159944]tumors with differential PRL receptor protein expression.
J. Clin.

[159945]Endocr. Metab. 86: 3826–3832, 2001.

[159946]

[159947]

[159948]15715.

[159949]Hu, Z.-Z.; Zhuang, L.; Meng, J.; Leondires, M.; Dufau, M.
L.:

[159950]The human prolactin receptor gene structure and alterna-
tive promoter

[159951]utilization: the generic promoter hPIII and a novel human
promoter

[159952]hP(N). J. Clin. Endocr. Metab. 84: 1153–1156, 1999.

[159953]

[159954]

[159955]15716.

[159956]Ormandy, C. J.; Camus, A.; Barra, J.; Damotte, D.; Lucas,
B.; Buteau,

[159957]H.; Edery, M.; Brousse, N.; Babinet, C.; Binart, N.; Kelly, P.
A.

[159958]: Null mutation of the prolactin receptor gene produces multiple reproductive

[159959]defects in the mouse. *Genes Dev.* 11: 167–178, 1997.

[159960]

[159961]

[159962]15717.

[159963]Perrot–Applanat, M.; Gualillo, O.; Pezet, A.; Vincent, V.; Edery,

[159964]M.; Kelly, P. A.: Dominant negative and cooperative effects of mutant

[159965]forms of prolactin receptor. *Molec. Endocr.* 11: 1020–1032, 1997.

[159966]

[159967]

[159968]15718.

[159969]Tan, F.; Morris, P. W.; Skidgel, R. A.; Erdos, E. G.: Sequencing

[159970]and cloning of human prolylcarboxypeptidase (angiotensinase C): similarity

[159971]to both serine carboxypeptidase and prolylendopeptidase families. *J.*

[159972]Biol. Chem. 268: 16631–16638, 1993.

[159973]

[159974]

[159975]15719.

[159976]Watson, B., Jr.; Nowak, N. J.; Myracle, A. D.; Shows, T. B.;
Warnock,

[159977]D. G.: The human angiotensinase C gene (HUMPCP) maps
to 11q14 within

[159978]700 kb of D11S901: a candidate gene for essential hyper-
tension. Genomics 44:

[159979]365–367, 1997.

[159980]

[159981]

[159982]15720.

[159983]Cheng, S. Y.; Gong, Q. H.; Parkinson, C.; Robinson, E. A.;
Appella,

[159984]E.; Merlino, G. T.; Pastan, I.: The nucleotide sequence of a
human

[159985]cellular thyroid hormone-binding protein present in en-
doplasmic reticulum. J.

[159986]Biol. Chem. 262: 11221–11227, 1987.

[159987]

[159988]

[159989]15721.

[159990]Koivu, J.; Myllyla, R.; Halaakoski, T.; Pihlajaniemi, T.; Tasa-
nen,

[159991]K.; Kivirikko, K. I.: A single polypeptide acts both as the
beta

[159992]subunit of prolyl 4-hydroxylase and as a protein disul-
fide-isomerase. J.

[159993]Biol. Chem. 262: 6447–6449, 1987.

[159994]

[159995]

[159996]15722.

[159997]Morris, J. I.; Varandani, P. T.: Characterization of a cDNA
for

[159998]human glutathione–insulin transhydrogenase
(protein–disulfide isomerase/oxidoreductase). Biochim.

[159999]Biophys. Acta 949: 169–180, 1988.

[160000]

[160001]

[160002]15723.

[160003]Noiva, R.; Lennarz, W. J.: Protein disulfide isomerase: a
multifunctional

[160004]protein resident in the lumen of the endoplasmic reticu-
lum. J. Biol.

[160005]Chem. 267: 3553–3556, 1992.

[160006]

[160007]

[160008]15724.

[160009]Pajunen, L.; Hoyhtya, M.; Tryggvason, K.; Kivirikko, K. I.;
Myllyla,

[160010]R.: Species-specific antibodies in the assignment of the
gene for

[160011]the beta-subunit of human prolyl 4-hydroxylase.

(Abstract) Cytogenet.

[160012]Cell Genet. 40: 719 only, 1985.

[160013]

[160014]

[160015]15725.

[160016]Pajunen, L.; Jones, T. A.; Goddard, A.; Sheer, D.; Solomon,

E.;

[160017]Pihlajaniemi, T.; Kivirikko, K. I.: Regional assignment of the human

[160018]gene coding for a multifunctional polypeptide (P4HB) acting as the

[160019]beta-subunit of prolyl 4-hydroxylase and the enzyme protein disulfide

[160020]isomerase to 17q25. Cytogenet. Cell Genet. 56: 165-168, 1991.

[160021]

[160022]

[160023]15726.

[160024]Pajunen, L.; Myllyla, R.; Helaakoski, T.; Pihlajaniemi, T.; Tasanen,

[160025]K.; Hoyhtya, M.; Tryggvason, K.; Solomon, E.; Kivirikko, K. I.: Assignment

[160026]of the gene coding for both the beta-subunit of prolyl 4-hydroxylase

[160027]and protein disulphide isomerase to human chromosome region 17q23-25.

[160028](Abstract) Cytogenet. Cell Genet. 46: 674 only, 1987.

[160029]

[160030]

[160031]15727.

[160032]Pajunen, L.; Myllyla, R.; Helaakoski, T.; Pihlajaniemi, T.;

Tasanen,

[160033]K.; Hoyhtya, M.; Tryggvason, K.; Solomon, E.; Kivirikko, K.

I.: Assignment

[160034]of the gene coding for both the beta-subunit of prolyl

4-hydroxylase

[160035]and the enzyme disulfide isomerase to human chromo-

some region 17p11-qter. Cytogenet.

[160036]Cell Genet. 47: 37-41, 1988.

[160037]

[160038]

[160039]15728.

[160040]Pihlajaniemi, T.; Helaakoski, T.; Tasanen, K.; Myllyla, R.;

Huhtala,

[160041]M.-L.; Koivu, J.; Kivirikko, K. I.: Molecular cloning of the

beta-subunit

[160042]of human prolyl 4-hydroxylase: this subunit and protein

disulphide

[160043]isomerase are products of the same gene. EMBO J. 6:

643–649, 1987.

[160044]

[160045]

[160046]15729.

[160047]Popescu, N. C.; Cheng, S.; Pastan, I.: Chromosomal localization

[160048]of the gene for a human thyroid hormone-binding protein. Am. J. Hum.

[160049]Genet. 42: 560–564, 1988.

[160050]

[160051]

[160052]15730.

[160053]Tasanen, K.; Parkkonen, T.; Chow, L. T.; Kivirikko, K. I.; Pihlajaniemi,

[160054]T.: Characterization of the human gene for a polypeptide that acts

[160055]both as the beta-subunit of prolyl 4-hydroxylase and as protein disulfide

[160056]isomerase. J. Biol. Chem. 263: 16218–16224, 1988.

[160057]

[160058]

[160059]15731.

[160060]Kelly, A.; Powis, S. H.; Glynne, R.; Radley, E.; Beck, S.;
Trowsdale,

[160061]J.: Second proteasome-related gene in the human MHC
class II region. Nature 353:

[160062]667-668, 1991.

[160063]

[160064]

[160065]15732.

[160066]Martinez, C. K.; Monaco, J. J.: Homology of proteasome
subunits

[160067]to a major histocompatibility complex-linked LMP gene.
Nature 353:

[160068]664-667, 1991.

[160069]

[160070]

[160071]15733.

[160072]Petes, T. D.: Meiotic recombination hot spots and cold

spots. Nature

[160073]Rev. Genet. 2: 360–369, 2001.

[160074]

[160075]

[160076]15734.

[160077]Van Kaer, L.; Ashton–Rickardt, P. G.; Eichelberger, M.;
Gaczynska,

[160078]M.; Nagashima, K.; Rock, K. L.; Goldberg, A. L.; Doherty, P.
C.; Tonegawa,

[160079]S.: Altered peptidase and viral–specific T cell response in
LMP2

[160080]mutant mice. Immunity 1: 533–541, 1994.

[160081]

[160082]

[160083]15735.

[160084]Zhou, P.; Zanelli, E.; Smart, M.; David, C.: Genomic organi-
zation

[160085]and tissue expression of mouse proteasome gene Lmp–2.
Genomics 16:

[160086]664–668, 1993.

[160087]

[160088]

[160089]15736.

[160090]Bodmer, J. G.; Marsh, S. G. E.; Albert, E. D.; Bodmer, W. F.;
Dupont,

[160091]B.; Erlich, H. A.; Mach, B.; Mayr, W. R.; Parham, P.;
Sasazuki, T.;

[160092]Schreuder, G. M. T.; Strominger, J. L.; Svejgaard, A.;
Terasaki, P.

[160093]I.: Nomenclature for factors of the HLA system, 1991. Tis-
sue Antigens 39:

[160094]161–173, 1992.

[160095]

[160096]

[160097]15737.

[160098]Deng, G. Y.; Muir, A.; Maclaren, N. K.; She, J.-X.: Associa-
tion

[160099]of LMP2 and LMP7 genes within the major histocompati-
bility complex

[160100]with insulin-dependent diabetes mellitus: population and
family studies. Am.

[160101]J. Hum. Genet. 56: 528–534, 1995.

[160102]

[160103]

[160104]15738.

[160105]Fehling, H. J.; Swat, W.; Laplace, C.; Kuhn, R.; Rajewsky, K.;

[160106]Muller, U.; von Boehmer, H.: MHC class I expression in
mice lacking

[160107]the proteasome subunit LMP-7. Science 265: 1234–1237,
1994.

[160108]

[160109]

[160110]15739.

[160111]Doolittle, R. F.; Hunkapiller, M. W.; Hood, L. E.; Devare, S.

[160112]G.; Robbins, K. C.; Aaronson, S. A.; Antoniades, H. N.:
Simian sarcoma

[160113]virus onc gene, v-sis, is derived from the gene (or genes)
encoding

[160114]a platelet-derived growth factor. Science 221: 275–277,
1983.

[160115]

[160116]

[160117]15740.

[160118]Frolova, L. Y.; Sudomoina, M. A.; Grigorieva, A. Y.; Zinovieva,

[160119]O. L.; Kisselev, L. L.: Cloning and nucleotide sequence of the structural

[160120]gene encoding for human tryptophanyl-tRNA synthetase. Gene 109:

[160121]291-296, 1991.

[160122]

[160123]

[160124]15741.

[160125]Vassart, G.: Personal Communication. Brussels, Belgium 1/15/1992.

[160126]

[160127]

[160128]15742.

[160129]Wenger, G. D.: Personal Communication. Columbus, Ohio 8/3/1993.

[160130]

[160131]

[160132]15743.

[160133]Gardner, T. L.; Elston, D. M.; Wotowic, P. J.: A familial dermatofibrosarcoma

[160134]protuberans. J. Am. Acad. Derm. 39: 504–505, 1998.

[160135]

[160136]

[160137]15744.

[160138]Groffen, J.; Heisterkamp, N.; Stephenson, J. R.; Geurts van Kessel,

[160139]A.; de Klein, A.; Grosveld, G.; Bootsma, D.: c-sis is translocated

[160140]from chromosome 22 to chromosome 9 in chronic myelocytic leukemia. J.

[160141]Exp. Med. 158: 9–15, 1983.

[160142]

[160143]

[160144]15745.

[160145]Hermansson, M.; Nister, M.; Betsholtz, C.; Heldin, C.-H.;

Westermarck,

[160146]B.; Funa, K.: Endothelial cell hyperplasia in human
glioblastoma:

[160147]coexpression of mRNA for platelet-derived growth factor
(PDGF) B chain

[160148]and PDGF receptor suggests autocrine growth stimulation.

Proc. Nat.

[160149]Acad. Sci. 85: 7748–7752, 1988.

[160150]

[160151]

[160152]15746.

[160153]Josephs, S. F.; Dalla-Favera, R.; Gelmann, E. P.; Gallo, R.
C.;

[160154]Wong-Staal, F.: 5-prime viral and human cellular se-
quences corresponding

[160155]to the transforming gene of simian sarcoma virus. Science
219: 503–505,

[160156]1983.

[160157]

[160158]

[160159]15747.

[160160]Josephs, S. F.; Guo, C.; Ratner, L.; Wong-Staal, F.: Human
proto-oncogene

[160161]nucleotide sequences corresponding to the transforming
region of simian

[160162]sarcoma virus. Science 223: 487-491, 1984.

[160163]

[160164]

[160165]15748.

[160166]Josephs, S. F.; Ratner, L.; Clarke, M. F.; Westin, E. H.; Reitz,

[160167]M. S.; Wong-Staal, F.: Transforming potential of human c-
sis nucleotide

[160168]sequences encoding platelet-derived growth factor. Sci-
ence 225:

[160169]636-639, 1984.

[160170]

[160171]

[160172]15749.

[160173]Kelly, J. D.; Raines, E. W.; Ross, R.; Murray, M. J.: The B
chain

[160174]of PDGF alone is sufficient for mitogenesis. EMBO J. 4:

3399–3405,

[160175]1985.

[160176]

[160177]

[160178]15750.

[160179]Kiuru–Kuhlefelt, S.; El–Rifai, W.; Fanburg–Smith, J.; Kere, J.;

[160180]Miettinen, M.; Knuutila, S.: Concomitant DNA copy number amplification

[160181]at 17q and 22q in dermatofibrosarcoma protuberans. Cytogenet. Cell

[160182]Genet. 92: 192–195, 2001.

[160183]

[160184]

[160185]15751.

[160186]Kozak, C. A.; Sears, J. F.; Hoggan, M. D.: Genetic mapping of

[160187]the mouse proto–oncogene c–sis to chromosome 15. Science 221: 867–869,

[160188]1983.

[160189]

[160190]

[160191]15752.

[160192]Lindahl, P.; Johansson, B. R.; Leveen, P.; Betsholtz, C.: Pericyte

[160193]loss and microaneurysm formation in PDGF-B-deficient mice. Science 277:

[160194]242–245, 1997.

[160195]

[160196]

[160197]15753.

[160198]Owen, A. J.; Pantazis, P.; Antoniades, H. N.: Simian sarcoma

[160199]virus-transformed cells secrete a mitogen identical to platelet-derived

[160200]growth factor. Science 225: 54–56, 1984.

[160201]

[160202]

[160203]15754.

[160204]Rao, C. D.; Igarashi, H.; Chiu, I.-M.; Robbins, K. C.; Aaronson,

[160205]S. A.: Structure and sequence of the human c-sis/platelet-derived

[160206]growth factor 2 (SIS/PDGF2) transcriptional unit. Proc. Nat. Acad.

[160207]Sci. 83: 2392-2396, 1986.

[160208]

[160209]

[160210]15755.

[160211]Robbins, K. C.; Antoniades, H. N.; Devare, S. G.; Hunkapiller,

[160212]M. W.; Aaronson, S. A.: Structural and immunological similarities

[160213]between simian sarcoma virus gene product(s) and human platelet-derived

[160214]growth factor. Nature 305: 605-608, 1983.

[160215]

[160216]

[160217]15756.

[160218]Robbins, K. C.; Devare, S. G.; Reddy, E. P.; Aaronson, S. A.:

[160219]In vivo identification of the transforming gene product of
simian

[160220]sarcoma virus. Science 218: 1131–1133, 1982.

[160221]

[160222]

[160223]15757.

[160224]Simon, M.–P.; Navarro, M.; Roux, D.; Pouyssegur, J.: Struc-
tural

[160225]and functional analysis of a chimeric protein
COL1A1–PDGFB generated

[160226]by the translocation t(17;22)(q22;q13.1) in dermatofi-
brosarcoma protuberans

[160227](DP). Oncogene 20: 2965–2975, 2001.

[160228]

[160229]

[160230]15758.

[160231]Smidt, M.; Kirsch, I.; Ratner, L.: Deletion of Alu sequences

[160232]in the fifth c-sis intron in individuals with meningiomas. J.

Clin.

[160233]Invest. 86: 1151–1157, 1990.

[160234]

[160235]

[160236]15759.

[160237]Turc–Carel, C.; Philip, I.; Berger, M. P.; Philip, T.; Lenoir,

[160238]G. M.: Chromosomal translocations in Ewing's sarcoma.

(Letter) New

[160239]Eng. J. Med. 309: 497–498, 1983.

[160240]

[160241]

[160242]15760.

[160243]Shimizu, A.; O'Brien, K. P.; Sjoblom, T.; Pietras, K.; Buch–

dunger,

[160244]E.; Collins, V. P.; Heldin, C.–H.; Dumanski, J. P.; Ostman,

A.: The

[160245]dermatofibrosarcoma protuberans–associated collagen

type I–alpha–1/platelet–derived

[160246]growth factor (PDGF) beta–chain fusion gene generates a

transforming

[160247]protein that is processed to functional PDGF-BB. Cancer
Res. 59:

[160248]3719–3723, 1999.

[160249]

[160250]

[160251]15761.

[160252]Simon, M.-P.; Pedeutour, F.; Sirvent, N.; Grosgeorge, J.;
Minoletti,

[160253]F.; Coindre, J.-M.; Terrier-Lacombe, M.-J.; Mandahl, N.;
Craver, R.

[160254]D.; Blin, N.; Sozzi, G.; Turc-Carel, C.; O'Brien, K. P.; Kedra,
D.;

[160255]Fransson, I.; Guilbaud, C.; Dumanski, J. P.: Deregulation of
the

[160256]platelet-derived growth factor B-chain gene via fusion
with collagen

[160257]gene COL1A1 in dermatofibrosarcoma protuberans and
giant-cell fibroblastoma. Nature

[160258]Genet. 15: 95–98, 1997.

[160259]

[160260]

[160261]15762.

[160262]Waterfield, M. D.; Scrace, G. T.; Whittle, N.; Stroobant, P.;

[160263]Johnsson, A.; Wasteson, A.; Westermarck, B.; Heldin, C.-H.;
Huang,

[160264]J. S.; Deuel, T. F.: Platelet-derived growth factor is struc-
turally

[160265]related to the putative transforming protein p28(sis) of
simian sarcoma

[160266]virus. Nature 304: 35-39, 1983.

[160267]

[160268]

[160269]15763.

[160270]Graphodatsky, A.; Frolova, L.; Biltueva, L.; Eremina, V.;
Lushnikova,

[160271]T.; Sudomoina, M.; Zinovieva, O.; Kisselev, L.: Localization
of the

[160272]tryptophanyl tRNA synthetase gene (WARS) on human and
bovine chromosomes

[160273]by in situ hybridization. Mammalian Genome 4: 183-184,
1993.

[160274]

[160275]

[160276]15764.

[160277]Jensen, L. L.; Nielsen, M. M.; Justesen, J.; Hansen, L. L.:

Assignment

[160278]of human NADH dehydrogenase (ubiquinone) 1 beta sub-complex 3 (NDUFB3)

[160279]and of its four pseudogenes to human chromosomes 2q31.3, 1p13.3-p13.1,

[160280]9q32-q34.1, 14q22.3-q23.1 and 14q32.2 by radiation hybrid mapping. Cytogenet.

[160281]Cell Genet. 93: 147-150, 2001.

[160282]

[160283]

[160284]15765.

[160285]Otani, A.; Slike, B. M.; Dorrell, M. I.; Hood, J.; Kinder, K.;

[160286]Ewalt, K. L.; Cheresch, D.; Schimmel, P.; Friedlander, M.: A fragment

[160287]of human TrpRS as a potent antagonist of ocular angiogenesis. Proc.

[160288]Nat. Acad. Sci. 99: 178-183, 2002.

[160289]

[160290]

[160291]15766.

[160292]Shimizu, N.; Kucherlapati, R. S.; Ruddle, F. H.: Assignment
of

[160293]a human gene for tryptophanyl-tRNA synthetase to chromosome 14 using

[160294]human-mouse somatic cell hybrids. Somat. Cell Genet. 2:
345-357,

[160295]1976.

[160296]

[160297]

[160298]15767.

[160299]Tolstrup, A. B.; Bejder, A.; Fleckner, J.; Justesen, J.: Transcriptional

[160300]regulation of the interferon-gamma-inducible tryptophanyl-tRNA synthetase

[160301]includes alternative splicing. J. Biol. Chem. 270: 397-403,
1995.

[160302]

[160303]

[160304]15768.

[160305]Turpaev, K. T.; Zakhariev, V. M.; Sokolova, I. V.;

Narovlyansky,

[160306]A. N.; Amchenkova, A. M.; Justesen, J.; Frolova, L. Y.: Al-

ternative

[160307]processing of the tryptophanyl-tRNA synthetase mRNA

from interferon-treated

[160308]human cells. Europ. J. Biochem. 240: 732-737, 1996.

[160309]

[160310]

[160311]15769.

[160312]Wakasugi, K.; Slike, B. M.; Hood, J.; Otani, A.; Ewalt, K. L.;

[160313]Friedlander, M.; Cheresh, D. A.; Schimmel, P.: A human

aminoacyl-tRNA

[160314]synthetase as a regulator of angiogenesis. Proc. Nat. Acad.

Sci. 99:

[160315]173-177, 2002.

[160316]

[160317]

[160318]15770.

[160319]Miller, J. S.; Moxley, G.; Schwartz, L. B.: Cloning and characterization

[160320]of a second complementary DNA for human tryptase. J. Clin. Invest. 86:

[160321]864–870, 1990.

[160322]

[160323]

[160324]15771.

[160325]Miller, J. S.; Westin, E. H.; Schwartz, L. B.: Cloning and characterization

[160326]of complementary DNA for human tryptase. J. Clin. Invest. 84: 1188–1195,

[160327]1989.

[160328]

[160329]

[160330]15772.

[160331]Vanderslice, P.; Ballinger, S. M.; Tam, E. K.; Goldstein, S. M.;

[160332]Craik, C. S.; Caughey, G. H.: Human mast cell tryptase:

multiple

[160333]cDNAs and genes reveal a multigene serine protease family. Proc.

[160334]Nat. Acad. Sci. 87: 3811–3815, 1990.

[160335]

[160336]

[160337]15773.

[160338]Mach, B.; Steimle, V.; Reith, W.: MHC class II-deficient combined

[160339]immunodeficiency: a disease of gene regulation. Immun. Rev. 138:

[160340]207–221, 1994.

[160341]

[160342]

[160343]15774.

[160344]Babcock, M.; de Silva, D.; Oaks, R.; Davis–Kaplan, S.; Jiralspong,

[160345]S.; Montermini, L.; Pandolfo, M.; Kaplan, J.: Regulation of mitochondrial

[160346]iron accumulation by Yfh1p, a putative homolog of

frataxin. Science 276:

[160347]1709–1712, 1997.

[160348]

[160349]

[160350]15775.

[160351]Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: The GAA
triplet-repeat

[160352]expansion in Friedreich ataxia interferes with transcription
and may

[160353]be associated with an unusual DNA structure. Am. J. Hum.
Genet. 62:

[160354]111–121, 1998.

[160355]

[160356]

[160357]15776.

[160358]Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: Atypical
Friedreich

[160359]ataxia caused by compound heterozygosity for a novel
missense mutation

[160360]and the GAA triplet-repeat expansion. (Letter) Am. J. Hum.

Genet. 60:

[160361]1251–1256, 1997.

[160362]

[160363]

[160364]15777.

[160365]Cavadini, P.; Gellera, C.; Patel, P. I.; Isaya, G.: Human
frataxin

[160366]maintains mitochondrial iron homeostasis in *Saccha-
romyces cerevisiae*. Hum.

[160367]Molec. Genet. 9: 2523–2530, 2000.

[160368]

[160369]

[160370]15778.

[160371]Sukhatme, V. P.; Vollmer, A. C.; Erikson, J.; Isobe, M.;
Croce,

[160372]C.; Parnes, J. R.: Gene for the human T cell differentiation
antigen

[160373]Leu-2/T8 is closely linked to the kappa light chain locus
on chromosome

[160374]2. J. Exp. Med. 161: 429–434, 1985.

[160375]

[160376]

[160377]15779.

[160378]Traver, D.; Akashi, K.; Manz, M.; Merad, M.; Miyamoto, T.;
Engleman,

[160379]E. G.; Weissman, I. L.: Development of CD8–alpha–positive
dendritic

[160380]cells from a common myeloid progenitor. Science 290:
2152–2154,

[160381]2000.

[160382]

[160383]

[160384]15780.

[160385]Weichhold, G. M.; Huber, C.; Parnes, J. R.; Zachau, H. G.:
The

[160386]CD8–alpha locus is located on the telomere side of the
immunoglobulin–kappa

[160387]locus at a distance of 2 Mb. Genomics 16: 512–514, 1993.

[160388]

[160389]

[160390]15781.

[160391]Balk, S. P.; Burke, S.; Polischuk, J. E.; Frantz, M. E.; Yang,

[160392]L.; Porcelli, S.; Colgan, S. P.; Blumberg, R. S.: Beta-

2-microglobulin-independent

[160393]MHC class Ib molecule expressed by human intestinal ep-

ithelium. Science 265:

[160394]259-262, 1994.

[160395]

[160396]

[160397]15782.

[160398]Benlagha, K.; Kyin, T.; Beavis, A.; Teyton, L.; Bendelac, A.:

[160399]A thymic precursor to NK T cell lineage. Science 296:

553-555, 2002.

[160400]

[160401]

[160402]15783.

[160403]Bradbury, A.; Milstein, C.; Kozak, C. A.: Chromosomal lo-

calization

[160404]of Cd1d genes in the mouse. Somat. Cell Molec. Genet.

17: 93–96,

[160405]1991.

[160406]

[160407]

[160408]15784.

[160409]Honey, K.; Benlagha, K.; Beers, C.; Forbush, K.; Teyton, L.;
Kleijmeer,

[160410]M. J.; Rudensky, A. Y.; Bendelac, A.: Thymocyte expression
of cathepsin

[160411]L is essential for NKT cell development. Nature Immun.
7Oct, 2002.

[160412]Note: Advance Electronic Publication.

[160413]

[160414]

[160415]15785.

[160416]MacDonald, H. R.: T before NK. Science 296: 481–482,
2002.

[160417]

[160418]

[160419]15786.

[160420]Pellicci, D. G.; Hammond, K. J. L.; Uldrich, A. P.; Baxter, A.

[160421]G.; Smyth, M. J.; Godfrey, D. I.: A natural killer T (NKT) cell
developmental

[160422]pathway involving a thymus-dependent NK1.1(-)CD4(+)
CD1d-dependent

[160423]precursor stage. J. Exp. Med. 195: 835–844, 2002.

[160424]

[160425]

[160426]15787.

[160427]Prigozy, T. I.; Naidenko, O.; Qasba, P.; Elewaut, D.;
Brossay,

[160428]L.; Khurana, A.; Natori, T.; Koezuka, Y.; Kulkarni, A.; Kro-
nenberg,

[160429]M.: Glycolipid antigen processing for presentation by
CD1d molecules. Science 291:

[160430]664–667, 2001.

[160431]

[160432]

[160433]15788.

[160434]Shi, F.-D.; Flodstrom, M.; Balasa, B.; Kim, S. H.; Van Gunst,
[160435]K.; Strominger, J. L.; Wilson, S. B.; Sarvetnick, N.: Germ line
deletion
[160436]of the CD1 locus exacerbates diabetes in the NOD mouse.
Proc. Nat.
[160437]Acad. Sci. 98: 6777–6782, 2001.

[160438]

[160439]

[160440]15789.

[160441]Hashimoto, K.; Hirai, M.; Kurosawa, Y.: A gene outside the
human
[160442]MHC related to classical HLA class I genes. Science 269:
693–695,
[160443]1995.

[160444]

[160445]

[160446]15790.

[160447]Shilo, B. Z.; Weinberg, R. A.: DNA sequences homologous
to vertebrate
[160448]oncogenes are conserved in *Drosophila melanogaster*.

Proc. Nat. Acad.

[160449]Sci. 78: 6789–6792, 1981.

[160450]

[160451]

[160452]15791.

[160453]Takahashi, E.; Hori, T.; O'Connell, P.; Leppert, M.; White, R.

[160454]: Mapping of the MYC gene to band 8q24.12–q24.13 by

R–banding and

[160455]distal to fra(8)(q24.11), FRA8E, by fluorescence in situ hy–

bridization. Cytogenet.

[160456]Cell Genet. 57: 109–111, 1991.

[160457]

[160458]

[160459]15792.

[160460]Taub, R.; Kirsch, I.; Morton, C.; Lenoir, G.; Swan, D.; Tron–

ick,

[160461]S.; Aaronson, S.; Leder, P.: Translocation of the c–myc

gene into

[160462]the immunoglobulin heavy chain locus in human Burkitt

lymphoma and

[160463]murine plasmacytoma cells. Proc. Nat. Acad. Sci. 79:
7837–7841,
[160464]1982.

[160465]

[160466]

[160467]15793.

[160468]Trumpf, A.; Refaeli, Y.; Oskarsson, T.; Gasser, S.; Murphy,
M.;

[160469]Martin, G. R.; Bishop, J. M.: c-Myc regulates mammalian
body size

[160470]by controlling cell number but not cell size. Nature 414:
768–773,

[160471]2001.

[160472]

[160473]

[160474]15794.

[160475]Vafa, O.; Wade, M.; Kern, S.; Beeche, M.; Pandita, T. K.;
Hampton,

[160476]G. M.; Wahl, G. M.: c-Myc can induce DNA damage, in-
crease reactive

[160477]oxygen species, and mitigate p53 function: a mechanism
for oncogene-induced

[160478]genetic instability. *Molec. Cell* 9: 1031–1044, 2002.

[160479]

[160480]

[160481]15795.

[160482]Watt, R.; Nishikura, K.; Sorrentino, J.; ar-Rushdi, A.; Croce,

[160483]C. M.; Rovera, G.: The structure and nucleotide sequence
of the 5-prime

[160484]end of the human c-myc oncogene. *Proc. Nat. Acad. Sci.*
80: 6307–6311,

[160485]1983.

[160486]

[160487]

[160488]15796.

[160489]Watt, R.; Stanton, L. W.; Marcu, K. B.; Gallo, R. C.; Croce, C.

[160490]M.; Rovera, G.: Nucleotide sequence of cloned cDNA of
human c-myc

[160491]oncogene. *Nature* 303: 725–728, 1983.

[160492]

[160493]

[160494]15797.

[160495]Azarnia, R.; Reddy, S.; Kmiecik, T. E.; Shalloway, D.;
Loewenstein,

[160496]W. R.: The cellular src gene product regulates junctional
cell-to-cell

[160497]communication. Science 239: 398–401, 1988.

[160498]

[160499]

[160500]15798.

[160501]Czernilofsky, A. P.; Levinson, A. D.; Varmus, H. E.; Bishop,
J.

[160502]M.; Tischer, E.; Goodman, H.: Correction to the nucleotide
sequence

[160503]of the src gene of Rous sarcoma virus. Nature 301:
736–738, 1983.

[160504]

[160505]

[160506]15799.

[160507]Gibbs, C. P.; Tanaka, A.; Anderson, S. K.; Radul, J.; Baar, J.;

[160508]Ridgway, A.; Kung, H.-J.; Fujita, D. J.: Isolation and structural

[160509]mapping of a human c-src gene homologous to the transforming gene

[160510](v-src) of Rous sarcoma virus. J. Virol. 53: 19-24, 1985.

[160511]

[160512]

[160513]15800.

[160514]Irby, R. B.; Mao, W.; Coppola, D.; Kang, J.; Loubeau, J. M.;
Trudeau,

[160515]W.; Karl, R.; Fujita, D. J.; Jove, R.; Yeatman, T. J.: Activating

[160516]SRC mutation in a subset of advanced human colon cancers. Nature

[160517]Genet. 21: 187-190, 1999.

[160518]

[160519]

[160520]15801.

[160521]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J.

D.:

[160522]c-src is consistently conserved in the chromosomal deletion (20q)

[160523]observed in myeloid disorders. Proc. Nat. Acad. Sci. 82: 6692-6696,

[160524]1985.

[160525]

[160526]

[160527]15802.

[160528]Lowe, C.; Yoneda, T.; Boyce, B. F.; Chen, H.; Mundy, G. R.; Soriano,

[160529]P.: Osteopetrosis in Src-deficient mice is due to an autonomous defect

[160530]of osteoclasts. Proc. Nat. Acad. Sci. 90: 4485-4489, 1993.

[160531]

[160532]

[160533]15803.

[160534]Morris, C. M.; Honeybone, L. M.; Hollings, P. E.; Fitzgerald,

[160535]P. H.: Localization of the SRC oncogene to chromosome band 20q11.2

[160536]and loss of this gene with deletion (20q) in two leukemic patients. Blood 74:

[160537]1768–1773, 1989.

[160538]

[160539]

[160540]15804.

[160541]Sakaguchi, A. Y.; Mohandas, T.; Naylor, S. L.: A human c-src

[160542]gene resides on the proximal long arm of chromosome 20 (cen–q13.1). Cancer

[160543]Genet. Cytogenet. 18: 123–129, 1985.

[160544]

[160545]

[160546]15805.

[160547]Sakaguchi, A. Y.; Naylor, S. L.; Weinberg, R. A.; Shows, T. B.

[160548]: Organization of human proto–oncogenes. (Abstract) Am. J. Hum. Genet. 34:

[160549]175A, 1982.

[160550]

[160551]

[160552]15806.

[160553]Sakaguchi, A. Y.; Zabel, B. U.; Grzeschik, K. H.; Law, M. L.;

[160554]Naylor, S. L.: Human proto-oncogene assignments.

(Abstract) Cytogenet.

[160555]Cell Genet. 37: 572–573, 1984.

[160556]

[160557]

[160558]15807.

[160559]Wong, B. R.; Besser, D.; Kim, N.; Arron, J. R.; Vologodskaya,

[160560]M.; Hanafusa, H.; Choi, Y.: TRANCE, a TNF family member,
activates

[160561]Akt/PKB through a signaling complex involving TRAF6 and
c-Src. Molec.

[160562]Cell 4: 1041–1049, 1999.

[160563]

[160564]

[160565]15808.

[160566]Mangelsdorf, D. J.; Borgmeyer, U.; Heyman, R. A.; Zhou, J.

Y.;

[160567]Ong, E. S.; Oro, A. E.; Kakizuka, A.; Evans, R. M.: Charac-
terization

[160568]of three RXR genes that mediate the action of 9-cis
retinoic acid. Genes

[160569]Dev. 6: 329-344, 1992.

[160570]

[160571]

[160572]15809.

[160573]Attard-Montalto, S.; Evans, N.; Sherwood, R. A.: Carote-
naemia

[160574]with low vitamin A levels and low retinol-binding protein.
J. Inherit.

[160575]Metab. Dis. 15: 929-930, 1992.

[160576]

[160577]

[160578]15810.

[160579]Chainani, M.; Sampsell, B.; Elliott, R. W.: Localization of
the

[160580]gene for plasma retinol binding protein to the distal half

of mouse

[160581]chromosome 19. Genomics 9: 376–379, 1991.

[160582]

[160583]

[160584]15811.

[160585]Matsuo, T.: Familial retinol-binding-protein deficiency?

(Letter) Lancet II:

[160586]910 only, 1987.

[160587]

[160588]

[160589]15812.

[160590]Matsuo, T.; Matsuo, N.; Shiraga, F.; Koide, N.: Familial
retinol-binding-protein

[160591]deficiency. (Letter) Lancet II: 402–403, 1987.

[160592]

[160593]

[160594]15813.

[160595]Morgan, F. F.; Canfield, R. E.; Goodman, D. S.: The partial
structure

[160596]of human plasma prealbumin and retinol-binding protein.

Biochim.

[160597]Biophys. Acta 236: 798–801, 1971.

[160598]

[160599]

[160600]15814.

[160601]Rask, L.; Anundi, H.; Fohlman, J.; Peterson, P. A.: The
complete

[160602]amino acid sequence of human serum retinol-binding
protein. Upsala

[160603]J. Med. Sci. 92: 115–146, 1987.

[160604]

[160605]

[160606]15815.

[160607]Rask, L.; Vahlquist, A.; Peterson, P. A.: Studies on two
physiological

[160608]forms of the human retinol-binding protein differing in
vitamin A

[160609]and arginine content. J. Biol. Chem. 246: 6638–6646,
1971.

[160610]

[160611]

[160612]15816.

[160613]Rocchi, M.; Covone, A.; Romeo, G.; Faraonio, R.; Colantuoni, V.

[160614]: Regional mapping of RBP4 to 10q23–q24 and RBP1 to 3q21–q22 in man. Somat.

[160615]Cell Molec. Genet. 15: 185–190, 1989.

[160616]

[160617]

[160618]15817.

[160619]Seeliger, M. W.; Biesalski, H. K.; Wissinger, B.; Gollnick, H.;

[160620]Gielen, S.; Frank, J.; Beck, S.; Zrenner, E.: Phenotype in retinol

[160621]deficiency due to a hereditary defect in retinol binding protein synthesis. Invest.

[160622]Ophthal. Vis. Sci. 40: 3–11, 1999.

[160623]

[160624]

[160625]15818.

[160626]Thuluvath, P. J.: Familial retinol-binding-protein deficiency?

[160627](Letter) Lancet II: 910 only, 1987.

[160628]

[160629]

[160630]15819.

[160631]Colantuoni, V.; Cortese, R.; Nilsson, M.; Lundvall, J.; Bavik,

[160632]C. O.; Eriksson, U.; Peterson, P. A.; Sundelin, J.: Cloning and sequencing

[160633]of a full length cDNA corresponding to human cellular retinol-binding

[160634]protein. Biochem. Biophys. Res. Commun. 130: 431-439, 1985.

[160635]

[160636]

[160637]15820.

[160638]Colantuoni, V.; Rocchi, M.; Roncuzzi, L.; Romeo, G.: Mapping of

[160639]human cellular retinol-binding protein to chromosome 3.

Cytogenet.

[160640]Cell Genet. 43: 221–222, 1986.

[160641]

[160642]

[160643]15821.

[160644]De Baere, E.; Speleman, F.; Van Roy, N.; De Paepe, A.;
Messiaen,

[160645]L.: Assignment of the cellular retinol-binding protein 1
gene (RBP1)

[160646]and of the coatmer beta-prime subunit gene (COPB2) to
human chromosome

[160647]band 3q23 by in situ hybridization. Cytogenet. Cell Genet.
82: 226–227,

[160648]1998.

[160649]

[160650]

[160651]15822.

[160652]Nilsson, M. H. L.; Spurr, N. K.; Lundvall, J.; Rask, L.; Peter-
son,

[160653]P. A.: Human cellular retinol-binding protein gene organi-

zation and

[160654]chromosomal location. *Europ. J. Biochem.* 173: 35–44,
1988.

[160655]

[160656]

[160657]15823.

[160658]Carson, N. L.; Simpson, N. E.: A physical map of 13 mark-
ers on

[160659]chromosome 10 from dosage studies on abnormal cell
lines. (Abstract) *Cytogenet.*

[160660]*Cell Genet.* 51: 974–975, 1989.

[160661]

[160662]

[160663]15824.

[160664]Chin, K. S.; Mathew, C. G. P.; Fong, S. L.; Bridges, C. D. B.;

[160665]Ponder, B. A. J.: Styl RFLP recognised by a human IRBP
cDNA localised

[160666]to chromosome 10. *Nucleic Acids Res.* 16: 1645 only,
1988.

[160667]

[160668]

[160669]15825.

[160670]Danciger, M.; Kozak, C. A.; Nickerson, J.; Redmond, T. M.;

Farber,

[160671]D. B.: Localization of the gene for interphotoreceptor
retinoid-binding

[160672]protein to mouse chromosome 14 near Np-1. Genomics
8: 727-731, 1990.

[160673]

[160674]

[160675]15826.

[160676]Farrer, L. A.; Castiglione, C. M.; Kidd, J. R.; Myers, S.; Car-

son,

[160677]N.; Simpson, N. E.; Kidd, K. K.: A linkage group of five
DNA markers

[160678]on human chromosome 10. Genomics 3: 72-77, 1988.

[160679]

[160680]

[160681]15827.

[160682]Fong, S.-L.; Liou, G. I.; Landers, R. A.; Alvarez, R. A.; Gon-

zalez-Fernandez,

[160683]F.; Glazebrook, P. A.; Lam, D. M. K.; Bridges, C. D. B.:

Characterization,

[160684]localization, and biosynthesis of an interstitial retinol-

binding

[160685]glycoprotein in the human eye. J. Neurochem. 42:

1667-1676, 1984.

[160686]

[160687]

[160688]15828.

[160689]Liou, G. I.; Fong, S.-L.; Beattie, W. G.; Cook, R. G.; Leone,
J.;

[160690]Landers, R. A.; Alvarez, R. A.; Wang, C.; Li, Y.; Bridges, C.
D. B.

[160691]: Bovine interstitial retinol-binding protein
(IRBP)--isolation and

[160692]sequence analysis of cDNA clones, characterization and in
vitro translation

[160693]of mRNA. Vision Res. 26: 1645-1653, 1986.

[160694]

[160695]

[160696]15829.

[160697]Liou, G. I.; Fong, S.-L.; Gosden, J.; vanTuinen, P.; Ledbet-
ter,

[160698]D. H.; Christie, S.; Rout, D.; Bhattacharya, S.; Cook, R. G.;
Li,

[160699]Y.; Wang, C.; Bridges, C. D. B.: Human interstitial retinol-
binding

[160700]protein (IRBP): cloning, partial sequence, and chromoso-
mal localization. Somat.

[160701]Cell Molec. Genet. 13: 315-323, 1987.

[160702]

[160703]

[160704]15830.

[160705]Liou, G. I.; Li, Y.; Wang, C.; Fong, S.-L.; Bhattacharya, F. S.;

[160706]Bridges, C. D. B.: Bgl II RFLP recognized by a human IRBP
cDNA localized

[160707]to chromosome 10. Nucleic Acids Res. 15: 3196 only,
1987.

[160708]

[160709]

[160710]15831.

[160711]McGuire, W.; Hill, A. V. S.; Allsopp, C. E. M.; Greenwood, B.

[160712]M.; Kwiatkowski, D.: Variation in the TNF- α promoter
region associated

[160713]with susceptibility to cerebral malaria. Nature 371:
508-511, 1994.

[160714]

[160715]

[160716]15832.

[160717]Benbrook, D.; Pfahl, M.: A novel thyroid hormone receptor
encoded

[160718]by a cDNA clone from a human testis library. Science 238:
788-791,

[160719]1987.

[160720]

[160721]

[160722]15833.

[160723]Bernal, J.; Refetoff, S.; DeGroot, L. J.: Abnormalities of tri-
iodothyronine

[160724]binding to lymphocyte and fibroblast nuclei from a patient

with peripheral

[160725]tissue resistance to thyroid hormone action. J. Clin. Endocr. Metab. 47:

[160726]1266–1272, 1978.

[160727]

[160728]

[160729]15834.

[160730]Dayton, A. I.; Selden, J. R.; Laws, G.; Dorney, D. J.; Finan, J.;

[160731]Tripputi, P.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.; Croce, C.

[160732]M.: A human c-erbA oncogene homologue is closely proximal to the

[160733]chromosome 17 breakpoint in acute promyelocytic leukemia. Proc. Nat.

[160734]Acad. Sci. 81: 4495–4499, 1984.

[160735]

[160736]

[160737]15835.

[160738]Debuire, B.; Henry, C.; Benaissa, M.; Biserte, G.; Claverie, J.

[160739]M.; Saule, S.; Martin, P.; Stehelin, D.: Sequencing the erbA
gene

[160740]of avian erythroblastosis virus reveals a new type of onco-
gene. Science 224:

[160741]1456–1459, 1984.

[160742]

[160743]

[160744]15836.

[160745]Ferro, M. T.; San Roman, C.: Constitutional t(15;17). Can-
cer

[160746]Genet. Cytogenet. 4: 89–91, 1981.

[160747]

[160748]

[160749]15837.

[160750]Gullberg, H.; Rudling, M.; Forrest, D.; Angelin, B.;
Vennstrom,

[160751]B.: Thyroid hormone receptor beta-deficient mice show
complete loss

[160752]of the normal cholesterol 7-alpha-hydroxylase (CYP7A)
response to

[160753]thyroid hormone but display enhanced resistance to dietary cholesterol. Molec.

[160754]Endocr. 14: 1739–1749, 2000.

[160755]

[160756]

[160757]15838.

[160758]Ichikawa, K.; Hughes, I. A.; Horwitz, A. L.; DeGroot, L. J.:
Characterization

[160759]of nuclear thyroid hormone receptors of cultured skin fibroblasts

[160760]from patients with resistance to thyroid hormone.

Metabolism 36:

[160761]392–399, 1987.

[160762]

[160763]

[160764]15839.

[160765]Iskaros, J.; Pickard, M.; Evans, I.; Sinha, A.; Hardiman, P.;
Ekins,

[160766]R.: Thyroid hormone receptor gene expression in first
trimester human

[160767]fetal brain. J. Clin. Endocr. Metab. 85: 2620–2623, 2000.

[160768]

[160769]

[160770]15840.

[160771]Jansson, M.; Philipson, L.; Vennstrom, B.: Isolation and
characterization

[160772]of multiple human genes homologous to the oncogenes of
avian erythroblastosis

[160773]virus. EMBO J. 2: 561–565, 1983.

[160774]

[160775]

[160776]15841.

[160777]Jhanwar, S. C.; Chaganti, R. S. K.; Croce, C. M.: Germ–line
chromosomal

[160778]localization of human c–erb–A oncogene. Somat. Cell
Molec. Genet. 11:

[160779]99–102, 1985.

[160780]

[160781]

[160782]15842.

[160783]Kaneshige, M.; Suzuki, H.; Kaneshige, K.; Cheng, J.; Wimbrow,

[160784]H.; Barlow, C.; Willingham, M. C.; Cheng, S.: A targeted dominant

[160785]negative mutation of the thyroid hormone alpha-1 receptor causes increased

[160786]mortality, infertility, and dwarfism in mice. Proc. Nat. Acad. Sci. 98:

[160787]15095-15100, 2001.

[160788]

[160789]

[160790]15843.

[160791]Laudet, V.; Begue, A.; Henry-Duthoit, C.; Joubel, A.; Martin,

[160792]P.; Stehelin, D.; Saule, S.: Genomic organization of the human thyroid

[160793]hormone receptor alpha (c-erbA-1) gene. Nucleic Acids Res. 19: 1105-1112,

[160794]1991.

[160795]

[160796]

[160797]15844.

[160798]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J.
D.;

[160799]Oren, M.: Translocation of the p53 gene in t(15;17) in
acute promyelocytic

[160800]leukaemia. Nature 316: 826–828, 1985.

[160801]

[160802]

[160803]15845.

[160804]Mathieu–Mahul, D.; Xu, D. Q.; Saule, S.; Lidereau, R.; Gal-
ibert,

[160805]F.; Berger, R.; Mauchauffe, M.; Larsen, C. J.: An EcoRI re-
striction

[160806]fragment length polymorphism (RFLP) in the human c-erb
A locus. Hum.

[160807]Genet. 71: 41–44, 1985.

[160808]

[160809]

[160810]15846.

[160811]McCabe, C. J.; Gittoes, N. J.; Sheppard, M. C.; Franklyn, J.

A.

[160812]: Thyroid receptor alpha-1 and alpha-2 mutations in non-functioning

[160813]pituitary tumors. J. Clin. Endocr. Metab. 84: 649-653, 1999.

[160814]

[160815]

[160816]15847.

[160817]Menezes-Ferreira, M. M.; Eil, C.; Wortsman, J.; Weintraub,

B.

[160818]D.: Decreased nuclear uptake of [125-I]triiodo-L-thyronine in fibroblasts

[160819]from patients with peripheral thyroid hormone resistance. J. Clin.

[160820]Endocr. Metab. 59: 1081-1087, 1984.

[160821]

[160822]

[160823]15848.

[160824]Mitelman, F.; Manolov, G.; Manolova, Y.; Billstrom, R.; Heim,

[160825]S.; Kristoffersson, U.; Mandahl, N.; Ferro, M. T.; San Roman, C.:

[160826]High resolution chromosome analysis of constitutional and acquired

[160827]t(15;17) maps c-erbA to subband 17q11.2. Cancer Genet. Cytogenet. 22:

[160828]95–98, 1986.

[160829]

[160830]

[160831]15849.

[160832]Miyajima, N.; Horiuchi, R.; Shibuya, Y.; Fukushige, S.; Matsubara,

[160833]K.; Toyoshima, K.; Yamamoto, T.: Two erbA homologs encoding proteins

[160834]with different T(3) binding capacities are transcribed from opposite

[160835]DNA strands of the same genetic locus. Cell 57: 31–39, 1989.

[160836]

[160837]

[160838]15850.

[160839]Nagaya, T.; Nomura, Y.; Fujieda, M.; Seo, H.: Heterodimer-
ization

[160840]preferences of thyroid hormone receptor alpha isoforms.
Biochem.

[160841]Biophys. Res. Commun. 226: 426–430, 1996.

[160842]

[160843]

[160844]15851.

[160845]Nakai, A.; Seino, S.; Sakurai, A.; Szilak, I.; Bell, G. I.; DeG-
root,

[160846]L. J.: Characterization of a thyroid hormone receptor ex-
pressed in

[160847]human kidney and other tissues. Proc. Nat. Acad. Sci. 85:
2781–2785,

[160848]1988.

[160849]

[160850]

[160851]15852.

[160852]Ng, L.; Rusch, A.; Amma, L. L.; Nordstrom, K.; Erway, L. C.;
Vennstrom,

[160853]B.; Forrest, D.: Suppression of the deafness and thyroid dysfunction

[160854]in Thrb-null mice by an independent mutation in the Thra thyroid hormone

[160855]receptor gene. Hum. Molec. Genet. 10: 2701–2708, 2001.

[160856]

[160857]

[160858]15853.

[160859]Puzianowska-Kuznicka, M.; Krystyniak, A.; Madej, A.; Cheng, S.-Y.;

[160860]Nauman, J.: Functionally impaired TR mutants are present in thyroid

[160861]papillary cancer. J. Clin. Endocr. Metab. 87: 1120–1128, 2002.

[160862]

[160863]

[160864]15854.

[160865]Refetoff, S.; DeGroot, L. J.; Benard, B.; DeWind, L. T.: Studies

[160866]of a sibship with apparent hereditary resistance to the intracellular

[160867]action of thyroid hormone. Metabolism 21: 723–756,
1972.

[160868]

[160869]

[160870]15855.

[160871]Rider, S. H.; Bailey, C. J.; Voss, R.; Sheer, D.; Hiorns, L. R.;

[160872]Solomon, E.: RFLP for the human erb-A1 gene. Nucleic
Acids Res. 15:

[160873]863 only, 1987.

[160874]

[160875]

[160876]15856.

[160877]Sakurai, A.; Nakai, A.; DeGroot, L. J.: Expression of three
forms

[160878]of thyroid hormone receptor in human tissues. Molec. En-
docr. 3:

[160879]392–399, 1989.

[160880]

[160881]

[160882]15857.

[160883]Cerretti, D. P.; Lyman, S. D.; Kozlosky, C. J.; Copeland, N. G.;

[160884]Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Kirstein, M. N.; Shapiro,

[160885]D. N.; Morris, S. W.: The genes encoding the Eph-related receptor

[160886]tyrosine kinase ligands LERK-1 (EPLG1, Epl1), LERK-3 (EPLG3, Epl3),

[160887]and LERK-4 (EPLG4, Epl4) are clustered on human chromosome 1 and mouse

[160888]chromosome 3. Genomics 33: 277-282, 1996.

[160889]

[160890]

[160891]15858.

[160892]Holzman, L. B.; Marks, R. M.; Dixit, V. M.: A novel immediate-early

[160893]response gene of endothelium is induced by cytokines and encodes a

[160894]secreted protein. Molec. Cell. Biol. 10: 5830-5838, 1990.

[160895]

[160896]

[160897]15859.

[160898]Pandey, A.; Lindberg, R. A.; Dixit, V. M.: Receptor orphans
find

[160899]a family. Curr. Biol. 5: 986–989, 1995.

[160900]

[160901]

[160902]15860.

[160903]Bunz, F.; Dutriaux, A.; Lengauer, C.; Waldman, T.; Zhou,
S.; Brown,

[160904]J. P.; Sedivy, J. M.; Kinzler, K. W.; Vogelstein, B.: Require-
ment

[160905]for p53 and p21 to sustain G2 arrest after DNA damage.
Science 282:

[160906]1497–1501, 1998.

[160907]

[160908]

[160909]15861.

[160910]Colucci-Guyon, E.; Portier, M.–M.; Dunia, I.; Paulin, D.;

Pournin,

[160911]S.; Babinet, C.: Mice lacking vimentin develop and reproduce without

[160912]an obvious phenotype. Cell 79: 679–694, 1994.

[160913]

[160914]

[160915]15862.

[160916]Ferrari, S.; Battini, R.; Kaczmarek, L.; Rittling, S.; Calabretta,

[160917]B.; de Riel, J. K.; Philiponis, V.; Weil, J.–F.; Baserga, R.: Coding

[160918]sequence and growth regulation of the human vimentin gene. Molec.

[160919]Cell. Biol. 6: 3614–3620, 1986.

[160920]

[160921]

[160922]15863.

[160923]Gieser, L.; Swaroop, A.: Expressed sequence tags and chromosomal

[160924]localization of cDNA clones from a subtracted retinal pig–

ment epithelium

[160925]library. Genomics 13: 873–876, 1992.

[160926]

[160927]

[160928]15864.

[160929]Lilienbaum, A.; Legagneux, V.; Portier, M.–M.; Dellagi, K.;
Paulin,

[160930]D.: Vimentin gene: expression in human lymphocytes and
in Burkitt's

[160931]lymphoma cells. EMBO J. 5: 2809–2814, 1986.

[160932]

[160933]

[160934]15865.

[160935]Marcus, E. M.; Smith, B. A.; Telenius, H.; Landsvater, R. M.;
Buys,

[160936]C. H. C. M.; Ferrari, S.; Ponder, B. A. J.; Mathew, C. G. P.:
Bcll

[160937]RFLP for the human vimentin gene. Nucleic Acids Res. 16:
9068 only,

[160938]1988.

[160939]

[160940]

[160941]15866.

[160942]Mathew, C. G.; Wakeling, W.; Jones, E.; Easton, D.; Fisher, R.;

[160943]Strong, C.; Smith, B.; Chin, K.; Little, P.; Nakamura, Y.; Shows,

[160944]T. B.; Jones, C.; Goodfellow, P. J.; Povey, S.; Ponder, B. A. J.:

[160945]Regional localization of polymorphic markers on chromosome 10 by physical

[160946]and genetic mapping. Ann. Hum. Genet. 54: 121–129, 1990.

[160947]

[160948]

[160949]15867.

[160950]Perreau, J.; Lilienbaum, A.; Vasseur, M.; Paulin, D.: Nucleotide

[160951]sequence of the human vimentin gene and regulation of its transcription

[160952]in tissues and cultured cells. Gene 62: 7–16, 1988.

[160953]

[160954]

[160955]15868.

[160956]Gimona, M.; Small, J. V.; Moeremans, M.; Van Damme, J.;
Puype,

[160957]M.; Vandekerckhove, J.: Porcine vinculin and metavinculin
differ

[160958]by a 68-residue insert located close to the carboxy-ter-
minal part

[160959]of the molecule. EMBO J. 7: 2329-2334, 1988.

[160960]

[160961]

[160962]15869.

[160963]Burglen, L.; Amiel, J.; Viollet, L.; Lefebvre, S.; Burlet, P.;

[160964]Clermont, O.; Raclin, V.; Landrieu, P.; Verloes, A.;
Munnich, A.;

[160965]Melki, J.: Survival motor neuron gene deletion in the
arthrogryposis

[160966]multiplex congenita-spinal muscular atrophy association.
J. Clin.

[160967]Invest. 98: 1130–1132, 1996.

[160968]

[160969]

[160970]15870.

[160971]Gabreels, B. A. T. F.; Swaab, D. F.; de Kleijn, D. P. V.; Seidah,

[160972]N. G.; Van de Loo, J.–W.; Van de Ven, W. J. M.; Martens, G. J. M.;

[160973]van Leeuwen, F. W.: Attenuation of the polypeptide 7B2, prohormone

[160974]convertase PC2, and vasopressin in the hypothalamus of some Prader–Willi

[160975]patients: indications for a processing defect. J. Clin. Endocr. Metab. 83:

[160976]591–599, 1998.

[160977]

[160978]

[160979]15871.

[160980]Martens, G. J. M.: Cloning and sequence analysis of human pituitary

[160981]cDNA encoding the novel polypeptide 7B2. FEBS Lett. 234:
160–164,
[160982]1988.

[160983]

[160984]

[160985]15872.

[160986]Mattei, M. G.; Mbikay, M.; Sylla, B. S.; Lenoir, G.; Mattei, J.
[160987]F.; Seidah, N. G.; Chretien, M.: Assignment of the gene for
neuroendocrine
[160988]protein 7B2 (SGNE1 locus) to mouse chromosome region
2[E3–F3] and
[160989]to human chromosome region 15q11–q15. Genomics 6:
436–440, 1990.

[160990]

[160991]

[160992]15873.

[160993]Mbikay, M.; Seidah, N. G.; Chretien, M.: Neuroendocrine
secretory
[160994]protein 7B2: structure, expression and functions.
Biochem. J. 357:

[160995]329–342, 2001.

[160996]

[160997]

[160998]15874.

[160999]Roebroek, A. J. M.; Dehaen, M. R. M.; van Bokhoven, A.;
Martens,

[161000]G. J. M.; Marynen, P.; van den Berghe, H.; Van de Ven, W. J.
M.:

[161001]Regional mapping of the human gene encoding the novel
pituitary polypeptide

[161002]7B2 to chromosome 15q13–q14 by in situ hybridization.
Cytogenet.

[161003]Cell Genet. 50: 158–160, 1989.

[161004]

[161005]

[161006]15875.

[161007]Westphal, C. H.; Muller, L.; Zhou, A.; Zhu, X.; Bonner–Weir,
S.;

[161008]Schambelan, M.; Steiner, D. F.; Lindberg, I.; Leder, P.: The
neuroendocrine

[161009]protein 7B2 is required for peptide hormone processing in vivo and

[161010]provides a novel mechanism for pituitary Cushing's disease. Cell 96:

[161011]689–700, 1999.

[161012]

[161013]

[161014]15876.

[161015]Bell, K.; Hopper, K. E.; McKenzie, H. A.; Murphy, W. H.; Shaw,

[161016]D. C.: A comparison of bovine alpha-lactalbumin A and B of Droughtmaster. Biochim.

[161017]Biophys. Acta 214: 437–444, 1970.

[161018]

[161019]

[161020]15877.

[161021]Bell, K.; McKenzie, H. A.; Murphy, W. H.; Shaw, D. C.: Beta-lactoglobulin

[161022](Droughtmaster): a unique protein variant. Biochim. Biophys. Acta 214:

[161023]427–436, 1970.

[161024]

[161025]

[161026]15878.

[161027]Julkunen, M.; Seppala, M.; Janne, O. A.: Complete amino acid sequence

[161028]of human placental protein 14: a progesterone-regulated uterine protein

[161029]homologous to beta-lactoglobulins. Proc. Nat. Acad. Sci. 85: 8845–8849,

[161030]1988.

[161031]

[161032]

[161033]15879.

[161034]Morris, H. R.; Dell, A.; Easton, R. L.; Panico, M.; Koistinen,

[161035]H.; Koistinen, R.; Oehninger, S.; Patankar, M. S.; Seppala,

M.; Clark,

[161036]G. F.: Gender-specific glycosylation of human glycodelin affects

[161037]its contraceptive activity. J. Biol. Chem. 271: 32159–32167, 1996.

[161038]

[161039]

[161040]15880.

[161041]Song, M.; Ramaswamy, S.; Ramachandran, S.; Flowers, L. C.; Horowitz,

[161042]I. R.; Rock, J. A.; Parthasarathy, S.: Angiogenic role for glycodelin

[161043]in tumorigenesis. Proc. Nat. Acad. Sci. 98: 9265–9270, 2001.

[161044]

[161045]

[161046]15881.

[161047]Van Cong, N.; Vaisse, C.; Gross, M.–S.; Slim, R.; Milgrom, E.;

[161048]Bernheim, A.: The human placental protein 14 (PP14) gene is localized

[161049]on chromosome 9q34. Hum. Genet. 86: 515–518, 1991.

[161050]

[161051]

[161052]15882.

[161053]Lee, F. S.; Fox, E. A.; Zhou, H.–M.; Strydom, D. J.; Vallee, B.

[161054]L.: Primary structure of human placental ribonuclease inhibitor. Biochemistry 27:

[161055]8545–8553, 1988.

[161056]

[161057]

[161058]15883.

[161059]Weremowicz, S.; Fox, E. A.; Morton, C. C.; Vallee, B. L.: The

[161060]placental ribonuclease inhibitor (RNH) gene is located on chromosome

[161061]subband 11p15.5. Genomics 8: 717–721, 1990.

[161062]

[161063]

[161064]15884.

[161065]Zneimer, S. M.; Crawford, D.; Schneider, N. R.; Beutler, B.:

Mapping

[161066]of the human ribonuclease inhibitor gene (RNH) to chromosome 11p15

[161067]by in situ hybridization. Genomics 8: 175–178, 1990.

[161068]

[161069]

[161070]15885.

[161071]Coughlin, P.; Nicholl, J.; Sun, J.; Salem, H.; Bird, P.; Sutherland,

[161072]G. R.: Chromosomal mapping of the human proteinase inhibitor 6 (PI6)

[161073]gene to 6p25 by fluorescence in situ hybridization. Genomics 26:

[161074]431–433, 1995.

[161075]

[161076]

[161077]15886.

[161078]Coughlin, P.; Sun, J.; Cerruti, L.; Salem, H. H.; Bird, P.:

Cloning

[161079]and molecular characterization of a human intracellular
serine proteinase

[161080]inhibitor. Proc. Nat. Acad. Sci. 90: 9417–9421, 1993.

[161081]

[161082]

[161083]15887.

[161084]Siu, G.; Wurster, A. L.; Duncan, D. D.; Soliman, T. M.;
Hedrick,

[161085]S. M.: A transcriptional silencer controls the developmen-
tal expression

[161086]of the CD4 gene. EMBO J. 13: 3570–3579, 1994.

[161087]

[161088]

[161089]15888.

[161090]Zhang, K.; Westberg, J. A.; Paetau, A.; von Boguslawsky,
K.; Lindsberg,

[161091]P.; Erlander, M.; Guo, H.; Su, J.; Olsen, H. S.; Andersson, L.
C.

[161092]: High expression of stanniocalcin in differentiated brain
neurons. Am.

[161093]J. Path. 153: 439–445, 1998.

[161094]

[161095]

[161096]15889.

[161097]Shpakovski, G. V.; Acker, J.; Wintzerith, M.; Lacroix, J. F.;
Thuriaux,

[161098]P.; Vigneron, M.: Four subunits that are shared by the
three classes

[161099]of RNA polymerase are functionally interchangeable be-
tween Homo sapiens

[161100]and Saccharomyces cerevisiae. Molec. Cell. Biol. 15:
4702–4710,

[161101]1995.

[161102]

[161103]

[161104]15890.

[161105]Hamilton, S. E.; Hurley, J. B.: A phosphodiesterase in-
hibitor

[161106]specific to a subset of bovine retinal cones. J. Biol. Chem.
265:

[161107]11259–11264, 1990.

[161108]

[161109]

[161110]15891.

[161111]Shimizu–Matsumoto, A.; Itoh, K.; Inazawa, J.; Nishida, K.;
Matsumoto,

[161112]Y.; Kinoshita, S.; Matsubara, K.; Okubo, K.: Isolation and
chromosomal

[161113]localization of the human cone cGMP phosphodiesterase
gamma cDNA (PDE6H). Genomics 32:

[161114]121–124, 1996.

[161115]

[161116]

[161117]15892.

[161118]Chen, J.; Engle, S. J.; Seilhamer, J. J.; Tischfield, J. A.:
Cloning

[161119]and recombinant expression of a novel human low molec-
ular weight Ca^{2+} –dependent

[161120]phospholipase A2. J. Biol. Chem. 269: 2365–2368, 1994.

[161121]

[161122]

[161123]15893.

[161124]Tischfield, J. A.; Xia, Y.-R.; Shih, D. M.; Klisak, I.; Chen, J.;

[161125]Engle, S. J.; Siakotos, A. N.; Winstead, M. V.; Seilhamer, J. J.;

[161126]Allamand, V.; Gyapay, G.; Lysis, A. J.: Low molecular weight, calcium-dependent

[161127]phospholipase A(2) genes are linked and map to homologous chromosome

[161128]regions in mouse and human. Genomics 32: 328–333, 1996.

[161129]

[161130]

[161131]15894.

[161132]Alexopoulou, L.; Thomas, V.; Schnare, M.; Lobet, Y.; Anguita, J.;

[161133]Schoen, R. T.; Medzhitov, R.; Fikrig, E.; Flavell, R. A.: Hyporesponsiveness

[161134]to vaccination with *Borrelia burgdorferi* OspA in humans and in TLR1–

[161135]and TLR2–deficient mice. Nature Med. 8: 878–884, 2002.

[161136]

[161137]

[161138]15895.

[161139]Gay, N. J.; Keith, F. J.: *Drosophila* Toll and IL-1 receptor.

(Letter) *Nature* 351:

[161140]355–356, 1991.

[161141]

[161142]

[161143]15896.

[161144]Muzio, M.; Bosisio, D.; Polentarutti, N.; D'amico, G.; Stop-
pacciaro,

[161145]A.; Mancinelli, R.; van't Veer, C.; Penton-Rol, G.; Ruco, L.
P.; Allavena,

[161146]P.; Mantovani, A.: Differential expression and regulation of
Toll-like

[161147]receptors (TLR) in human leukocytes: selective expression
of TLR3

[161148]in dendritic cells. *J. Immun.* 164: 5998–6004, 2000.

[161149]

[161150]

[161151]15897.

[161152]Rock, F. L.; Hardiman, G.; Timans, J. C.; Kastelein, R. A.;

Bazan,

[161153]J. F.: A family of human receptors structurally related to

Drosophila

[161154]Toll. *Proc. Nat. Acad. Sci.* 95: 588–593, 1998.

[161155]

[161156]

[161157]15898.

[161158]Taguchi, T.; Mitcham, J. L.; Dower, S. K.; Sims, J. E.; Testa,

[161159]J. R.: Chromosomal localization of TIL, a gene encoding a
protein

[161160]related to the *Drosophila* transmembrane receptor Toll, to
human chromosome

[161161]4p14. *Genomics* 32: 486–488, 1996.

[161162]

[161163]

[161164]15899.

[161165]Xu, Y.; Tao, X.; Shen, B.; Horng, T.; Medzhitov, R.; Manley,

J.

[161166]L.; Tong, L.: Structural basis for signal transduction by the
Toll/interleukin-1

[161167]receptor domains. Nature 408: 111-115, 2000.

[161168]

[161169]

[161170]15900.

[161171]Perrimon, N.; Mahowald, A. P.: Multiple functions of seg-
ment polarity

[161172]genes in Drosophila. Dev. Biol. 119: 587-600, 1987.

[161173]

[161174]

[161175]15901.

[161176]Pizzuti, A.; Novelli, G.; Mari, A.; Ratti, A.; Colosimo, A.;
Amati,

[161177]F.; Penso, D.; Sangiuolo, F.; Calabrese, G.; Palka, G.; Silani,
V.;

[161178]Gennarelli, M.; Mingarelli, R.; Scarlato, G.; Scambler, P.;
Dallapiccola,

[161179]B.: Human homologue sequences to the Drosophila di-

shevelled segment-polarity

[161180]gene are deleted in the DiGeorge syndrome. Am. J. Hum. Genet. 58:

[161181]722–729, 1996.

[161182]

[161183]

[161184]15902.

[161185]Brown, E. J.; Albers, M. W.; Shin, T. B.; Ichikawa, K.; Keith,

[161186]C. T.; Lane, W. S.; Schreiber, S. L.: A mammalian protein targeted

[161187]by G1-arresting rapamycin-receptor complex. Nature 369: 756–758,

[161188]1994.

[161189]

[161190]

[161191]15903.

[161192]Dennis, P. B.; Jaeschke, A.; Saitoh, M.; Fowler, B.; Kozma, S.

[161193]C.; Thomas, G.: Mammalian TOR: a homeostatic ATP sensor. Science 294:

[161194]1102–1105, 2001.

[161195]

[161196]

[161197]15904.

[161198]Fang, Y.; Vilella–Bach, M.; Bachmann, R.; Flanigan, A.;
Chen, J.

[161199]: Phosphatidic acid–mediated mitogenic activation of
mTOR signaling. Science 294:

[161200]1942–1945, 2001.

[161201]

[161202]

[161203]15905.

[161204]Hara, K.; Maruki, Y.; Long, X.; Yoshino, K.; Oshiro, N.; Hi–
dayat,

[161205]S.; Tokunaga, C.; Avruch, J.; Yonezawa, K.: Raptor, a bind–
ing partner

[161206]of target of rapamycin, mediates TOR action. Cell 110:
177–189,

[161207]2002.

[161208]

[161209]

[161210]15906.

[161211]Kim, D.-H.; Sarbassov, D. D.; Ali, S. M.; King, J. E.; Latek, R.

[161212]R.; Erdjument-Bromage, H.; Tempst, P.; Sabatini, D. M.:
mTOR interacts

[161213]with raptor to form a nutrient-sensitive complex that signals to the

[161214]cell growth machinery. Cell 110: 163–175, 2002.

[161215]

[161216]

[161217]15907.

[161218]Lench, N. J.; Macadam, R.; Markham, A. F.: The human
gene encoding

[161219]FKBP–rapamycin associated protein (FRAP) maps to chromosomal band

[161220]1p36.2. Hum. Genet. 99: 547–549, 1997.

[161221]

[161222]

[161223]15908.

[161224]Moore, P. A.; Rosen, C. A.; Carter, K. C.: Assignment of the human

[161225]FKBP12–rapamycin–associated protein (FRAP) gene to chromosome 1p36

[161226]by fluorescence in situ hybridization. Genomics 33: 331–332, 1996.

[161227]

[161228]

[161229]15909.

[161230]Onyango, P.; Lubyova, B.; Gardellin, P.; Kurzbauer, R.; Weith,

[161231]A.: Molecular cloning and expression analysis of five novel genes

[161232]in chromosome 1p36. Genomics 50: 187–198, 1998.

[161233]

[161234]

[161235]15910.

[161236]Sabatini, D. M.; Erdjument–Bromage, H.; Lui, M.; Tempst, P.; Snyder,

[161237]S. H.: RAFT1: a mammalian protein that binds to FKBP12 in a rapamycin–dependent

[161238]fashion and is homologous to yeast TORs. Cell 78: 35–43,
1994.

[161239]

[161240]

[161241]15911.

[161242]Barbier, M.; Attoub, S.; Calvez, R.; Laffargue, M.; Jarry, A.;

[161243]Mareel, M.; Altruda, F.; Gespach, C.; Wu, D.; Lu, B.; Hirsch,
E.;

[161244]Wymann, M. P.: Weakening link to colorectal cancer? Na-
ture 413:

[161245]796 only, 2001.

[161246]

[161247]

[161248]15912.

[161249]Crackower, M. A.; Oudit, G. Y.; Kozieradzki, I.; Sarao, R.;
Sun,

[161250]H.; Sasaki, T.; Hirsch, E.; Suzuki, A.; Shioi, T.; Irie–Sasaki,
J.;

[161251]Sah, R.; Cheng, H.–Y. M.; and 13 others: Regulation of
myocardial

[161252]contractility and cell size by distinct PI3K–PTEN signaling pathways. Cell 110:

[161253]737–749, 2002.

[161254]

[161255]

[161256]15913.

[161257]Hirsch, E.; Katanaev, V. L.; Garlanda, C.; Azzolino, O.; Pirola,

[161258]L.; Silengo, L.; Sozzani, S.; Mantovani, A.; Altruda, F.; Wymann,

[161259]M. P.: Central role for G protein–coupled phosphoinositide 3–kinase

[161260]gamma in inflammation. Science 287: 1049–1053, 2000.

[161261]

[161262]

[161263]15914.

[161264]Jiang, K.; Zhong, B.; Gilvary, D. L.; Corliss, B. C. Hong–Geller,

[161265]E.; Wei, S; Djeu, J. Y.: Pivotal role of phosphoinositide–3 kinase

[161266]in regulation of cytotoxicity in natural killer cells. Nature
Immun. 1:

[161267]419–425, 2000.

[161268]

[161269]

[161270]15915.

[161271]Kikuta, Y.; Kato, M.; Yamashita, Y.; Miyauchi, Y.; Tanaka,
K.;

[161272]Kamada, N.; Kusunose, M.: Human leukotriene B4 omega-
hydroxylase

[161273](CYP4F3) gene: molecular cloning and chromosomal local-
ization. DNA

[161274]Cell Biol. 17: 221–230, 1998.

[161275]

[161276]

[161277]15916.

[161278]Kikuta, Y.; Kusunose, E.; Endo, K.; Yamamoto, S.; Sogawa,
K.; Fujii–Kuriyama,

[161279]Y.; Kusunose, M.: A novel form of cytochrome P–450 fam-
ily 4 in human

[161280]polymorphonuclear leukocytes: cDNA cloning and expression of leukotriene

[161281]B4 omega-hydroxylase. J. Biol. Chem. 268: 9376–9380, 1993.

[161282]

[161283]

[161284]15917.

[161285]Klingensmith, J.; Nusse, R.; Perrimon, N.: The Drosophila segment

[161286]polarity gene dishevelled encodes a novel protein required for response

[161287]to wingless signal. Genes Dev. 8: 118–130, 1994.

[161288]

[161289]

[161290]15918.

[161291]Holmes, S. E.; Riazzi, M. A.; Gong, W.; McDermid, H. E.; Sellinger,

[161292]B. T.; Hua, A.; Chen, F.; Wang, Z.; Zhang, G.; Roe, B.; Gonzalez,

[161293]I.; McDonald-McGinn, D. M.; Zackai, E.; Emanuel, B. S.; Budarf, M.

[161294]L.: Disruption of the clathrin heavy chain-like gene
(CLTCL) associated

[161295]with features of DGS/VCFS: a balanced (21;22)(p12;q11)
translocation. Hum.

[161296]Molec. Genet. 6: 357–367, 1997.

[161297]

[161298]

[161299]15919.

[161300]Kedra, D.; Peyrard, M.; Fransson, I.; Collins, J. E.; Dunham,
I.;

[161301]Roe, B. A.; Dumanski, J. P.: Characterization of a second
human clathrin

[161302]heavy chain polypeptide gene (CLH-22) from chromosome
22q11. Hum.

[161303]Molec. Genet. 5: 625–631, 1996.

[161304]

[161305]

[161306]15920.

[161307]Long, K. R.; Trofatter, J. A.; Ramesh, V.; McCormick, M. K.;
Buckler,

[161308]A. J.: Cloning and characterization of a novel human
clathrin heavy

[161309]chain gene (CLTCL). Genomics 35: 466–472, 1996.

[161310]

[161311]

[161312]15921.

[161313]Lagenaur, C.; Kunemund, V.; Fischer, G.; Fushiki, S.;
Schachner,

[161314]M.: Monoclonal M6 antibody interferes with neurite extension of cultured

[161315]neurons. J. Neurobiol. 23: 71–88, 1992.

[161316]

[161317]

[161318]15922.

[161319]Shimizu, F.; Watanabe, T. K.; Fujiwara, T.; Takahashi, E.;
Nakamura,

[161320]Y.; Maekawa, H.: Isolation and mapping of the human glycoprotein

[161321]M6 gene (GPM6A) to 4q33–to–q34. Cytogenet. Cell Genet.
74: 138–139,

[161322]1996.

[161323]

[161324]

[161325]15923.

[161326]Bui, T. D.; Beier, D. R.; Jonssen, M.; Smith, K.; Dorrington, S.

[161327]M.; Kaklamanis, L.; Kearney, L.; Regan, R.; Sussman, D. J.; Harris,

[161328]A. L.: cDNA cloning of a human dishevelled DVL-3 gene, mapping to

[161329]3q27, and expression in human breast and colon carcinomas. Biochem.

[161330]Biophys. Res. Commun. 239: 510-516, 1997.

[161331]

[161332]

[161333]15924.

[161334]Cheng, Y.-S. E.; Patterson, C. E.; Staeheli, P.: Interferon-induced

[161335]guanylate-binding proteins lack an N(T)KXD consensus motif and bind

[161336]GMP in addition to GDP and GTP. Molec. Cell. Biol. 11:
4717–4725,
[161337]1991.

[161338]

[161339]

[161340]15925.

[161341]Kumar, S.; Li, Q.; Dua, A.; Ying, Y.–K.; Bagchi, M. K.;
Bagchi,

[161342]I. C.: Messenger ribonucleic acid encoding interferon–in–
ducible guanylate

[161343]binding protein 1 is induced in human endometrium
within the putative

[161344]window of implantation. J. Clin. Endocr. Metab. 86:
2420–2427, 2001.

[161345]

[161346]

[161347]15926.

[161348]Prochazka, M.; Staeheli, P.; Holmes, R. S.; Haller, O.: Inter–
feron–induced

[161349]guanylate–binding proteins: mapping of the murine Gbp–
1 locus to chromosome

[161350]3. Virology 145: 273–279, 1985.

[161351]

[161352]

[161353]15927.

[161354]Strehlow, I.; Lohmann–Matthes, M. L.; Decker, T.: The interferon–inducible

[161355]GBP1 gene: structure and mapping to human chromosome 1. Gene 144:

[161356]295–299, 1994.

[161357]

[161358]

[161359]15928.

[161360]Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek,

[161361]F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.; Copeland, N.

[161362]G.: Murine chromosomal location of four hepatocyte–enriched transcription

[161363]factors: HNF–3–alpha, HNF3–beta, HNF–3–gamma, and HFN–4. Genomics 13:

[161364]264–268, 1992.

[161365]

[161366]

[161367]15929.

[161368]Chartier, F. L.; Bossu, J.-P.; Laudet, V.; Fruchart, J.-C.:

Cloning

[161369]and sequencing of cDNAs encoding the human hepatocyte
nuclear factor

[161370]4 indicate the presence of two isoforms in human liver.

Gene 147:

[161371]269-272, 1994.

[161372]

[161373]

[161374]15930.

[161375]Eeckhoute, J.; Formstecher, P.; Laine, B.: Maturity-onset

diabetes

[161376]of the young type 1 (MODY1)-associated mutations R154X
and E276Q in

[161377]hepatocyte nuclear factor 4-alpha (HNF4-alpha) gene im-
pair recruitment

[161378]of p300, a key transcriptional coactivator. Molec. Endocr.
15: 1200-1210,

[161379]2001.

[161380]

[161381]

[161382]15931.

[161383]Furuta, H.; Iwasaki, N.; Oda, N.; Hinokio, Y.; Horikawa, Y.;
Yamagata,

[161384]K.; Yano, N.; Sugahiro, J.; Ogata, M.; Ohgawara, H.; Omori,
Y.; Iwamoto,

[161385]Y.; Bell, G. I.: Organization and partial sequence of the
hepatocyte

[161386]nuclear factor-4-alpha/MODY1 gene and identification of
a missense

[161387]mutation, R127W, in a Japanese family with MODY. Dia-
betes 46: 1652-1657,

[161388]1997.

[161389]

[161390]

[161391]15932.

[161392]Hani, E. H.; Suaud, L.; Boutin, P.; Chevre, J.-C.; Durand, E.;

[161393]Philippi, A.; Demenais, F.; Vionnet, N.; Furuta, H.; Velho,

G.; Bell,

[161394]G. I.; Laine, B.; Froguel, P.: A missense mutation in hepatocyte

[161395]nuclear factor-4-alpha, resulting in a reduced transactivation activity,

[161396]in human late-onset non-insulin-dependent diabetes mellitus. J. Clin.

[161397]Invest. 101: 521-526, 1998.

[161398]

[161399]

[161400]15933.

[161401]Lausen, J.; Thomas, H.; Lemm, I.; Bulman, M.; Borgschulze, M.;

[161402]Lingott, A.; Hattersley, A. T.; Ryffel, G. U.: Naturally occurring ring

[161403]mutations in the human HNF4-alpha gene impair the function of the

[161404]transcription factor to a varying degree. Nucleic Acids Res. 28:

[161405]430-437, 2000.

[161406]

[161407]

[161408]15934.

[161409]Li, J.; Ning, G.; Duncan, S. A.: Mammalian hepatocyte differentiation

[161410]requires the transcription factor HNF-4-alpha. Genes Dev. 14: 464-474,

[161411]2000.

[161412]

[161413]

[161414]15935.

[161415]Stoffel, M.; Duncan, S. A.: The maturity-onset diabetes of the

[161416]young (MODY1) transcription factor HNF4-alpha regulates expression

[161417]of genes required for glucose transport and metabolism. Proc. Nat.

[161418]Acad. Sci. 94: 13209-13214, 1997.

[161419]

[161420]

[161421]15936.

[161422]Yamagata, K.; Furuta, H.; Oda, N.; Kaisaki, P. J.; Menzel, S.;

[161423]Cox, N. J.; Fajans, S. S.; Signorini, S.; Stoffel, M.; Bell, G. I.

[161424]: Mutations in the hepatocyte nuclear factor-4-alpha gene
in maturity-onset

[161425]diabetes of the young (MODY1). Nature 384: 458-460,
1996.

[161426]

[161427]

[161428]15937.

[161429]Zouali, H.; Hani, E. H.; Philippi, A.; Vionnet, N.; Beckmann,

[161430]J. S.; Demenais, F.; Froguel, P.: A susceptibility locus for
early-onset

[161431]non-insulin dependent (type 2) diabetes mellitus maps to
chromosome

[161432]20q, proximal to the phosphoenolpyruvate carboxykinase
gene. Hum.

[161433]Molec. Genet. 6: 1401-1408, 1997.

[161434]

[161435]

[161436]15938.

- [161437]Frolova, L.; Le Goff, X.; Rasmussen, H. H.; Cheperegin, S.;
Drugeon,
[161438]G.; Kress, M.; Arman, I.; Haenni, A.–L.; Celis, J. E.; Philippe,
M.;
- [161439]Justesen, J.; Kisselev, L.: A highly conserved eukaryotic
protein
[161440]family possessing properties of polypeptide chain release
factor. Nature 372:
[161441]701–703, 1994.
- [161442]
- [161443]
- [161444]15939.
- [161445]Guenet, L.; Henry, C.; Toutain, B.; Dubourg, C.; Le Gall, J.
Y.;
- [161446]David, V.; Le Treut, A.: Eukaryotic translation termination
factor
[161447]gene (ETF1/eRF1) maps at D5S500 in a commonly deleted
region of chromosome
[161448]5q31 in malignant myeloid diseases. Cytogenet. Cell
Genet. 88: 82–86,
[161449]2000.

[161450]

[161451]

[161452]15940.

[161453]Hansen, L. L.; Jakobsen, C. G.; Justesen, J.: Assignment of
the

[161454]human translation termination factor 1 (ETF1) to 5q31.1
and of the

[161455]proximal marker D5S1995 by radiation hybrid mapping.
Cytogenet. Cell

[161456]Genet. 87: 256–257, 1999.

[161457]

[161458]

[161459]15941.

[161460]Le Goff, X.; Philippe, M.; Jean-Jean, O.: Overexpression of
human

[161461]release factor 1 alone has an antisuppressor effect in hu-
man cells. Molec.

[161462]Cell Biol. 17: 3164–3172, 1997.

[161463]

[161464]

[161465]15942.

[161466]Song, H.; Mugnier, P.; Das, A. K.; Webb, H. M.; Evans, D.

R.; Tuite,

[161467]M. F.; Hemmings, B. A.; Barford, D.: The crystal structure
of human

[161468]eukaryotic release factor eRF1--mechanism of stop codon
recognition

[161469]and peptidyl-tRNA hydrolysis. Cell 100: 311-321, 2000.

[161470]

[161471]

[161472]15943.

[161473]DiDonato, C. J.; Chen, X.-N.; Noya, D.; Korenberg, J. R.;

Nadeau,

[161474]J. H.; Simard, L. R.: Cloning, characterization, and copy
number

[161475]of the murine survival motor neuron gene: homolog of the
spinal muscular

[161476]atrophy-determining gene. Genome Res. 7: 339-352,
1997.

[161477]

[161478]

[161479]15944.

[161480]DiDonato, C. J.; Morgan, K.; Carpten, J. D.; Fuerst, P.; In-
graham,

[161481]S. E.; Prescott, G.; McPherson, J. D.; Wirth, B.; Zerres, K.;
Hurko,

[161482]O.; Wasmuth, J. J.; Mendell, J. R.; Burghes, A. H. M.;
Simard, L.

[161483]R.: Association between Ag1-CA alleles and severity of
autosomal

[161484]recessive proximal spinal muscular atrophy. Am. J. Hum.
Genet. 55:

[161485]1218-1229, 1994.

[161486]

[161487]

[161488]15945.

[161489]Feldkotter, M.; Schwarzer, V.; Wirth, R.; Wienker, T. F.;
Wirth,

[161490]B.: Quantitative analyses of SMN1 and SMN2 based on
real-time lightCycler

[161491]PCR: fast and highly reliable carrier testing and prediction
of severity

[161492]of spinal muscular atrophy. Am. J. Hum. Genet. 70:
358–368, 2002.

[161493]

[161494]

[161495]15946.

[161496]Friesen, W. J.; Massenet, S.; Paushkin, S.; Wyce, A.; Drey-
fuss,

[161497]G.: SMN, the product of the spinal muscular atrophy gene,
binds preferentially

[161498]to dimethylarginine-containing protein targets. Molec.
Cell 7: 1111–1117,

[161499]2001.

[161500]

[161501]

[161502]15947.

[161503]Frugier, T.; Tiziano, F. D.; Cifuentes–Diaz, C.; Miniou, P.;
Roblot,

[161504]N.; Dierich, A.; Le Meur, M.; Melki, J.: Nuclear targeting
defect

[161505]of SMN lacking the C-terminus in a mouse model of spinal
muscular

[161506]atrophy. Hum. Molec. Genet. 9: 849–858, 2000.

[161507]

[161508]

[161509]15948.

[161510]Gambardella, A.; Mazzei, R.; Toscano, A.; Annesi, G.;
Pasqua,

[161511]A.; Annesi, F.; Quattrone, F.; Oliveri, R. L.; Valentino, P.;
Bono,

[161512]F.; Aguglia, U.; Zappia, M.; Vita, G.; Quattrone, A.: Spinal
muscular

[161513]atrophy due to an isolated deletion of exon 8 of the
telomeric survival

[161514]motor neuron gene. Ann. Neurol. 44: 836–839, 1998.

[161515]

[161516]

[161517]15949.

[161518]Hahnen, E.; Schonling, J.; Rudnik–Schoneborn, S.;
Raschke, H.;

[161519]Zerres, K.; Wirth, B.: Missense mutations in exon 6 of the
survival

[161520]motor neuron gene in patients with spinal muscular atrophy (SMA). Hum.

[161521]Molec. Genet. 6: 821–825, 1997.

[161522]

[161523]

[161524]15950.

[161525]Hannus, S.; Buhler, D.; Romano, M.; Seraphin, B.; Fischer, U.

[161526]: The Schizosaccharomyces pombe protein Yab8p and a novel factor,

[161527]Yip1p, share structural and functional similarity with the spinal

[161528]muscular atrophy-associated proteins SMN and SIP1. Hum. Molec. Genet. 9:

[161529]663–674, 2000.

[161530]

[161531]

[161532]15951.

[161533]Hsieh-Li, H. M.; Chang, J.-G.; Jong, Y.-J.; Wu, M.-H.; Wang, N.

[161534]M.; Tsai, C. H.; Li, H.: A mouse model for spinal muscular atrophy. Nature

[161535]Genet. 24: 66–70, 2000.

[161536]

[161537]

[161538]15952.

[161539]Jablonka, S.; Bandilla, M.; Wiese, S.; Buhler, D.; Wirth, B.;

[161540]Sendtner, M.; Fischer, U.: Co-regulation of survival of motor neuron

[161541](SMN) protein and its interactor SIP1 during development and in spinal

[161542]muscular atrophy. Hum. Molec. Genet. 10: 497–505, 2001.

[161543]

[161544]

[161545]15953.

[161546]Liu, Q.; Dreyfuss, G.: A novel nuclear structure containing the

[161547]survival of motor neurons protein. EMBO J. 15: 3555–3565, 1996.

[161548]

[161549]

[161550]15954.

[161551]Liu, Q.; Fischer, U.; Wang, F.; Dreyfuss, G.: The spinal
muscular

[161552]atrophy disease gene product, SMN, and its associated
protein SIP1

[161553]are in a complex with spliceosomal snRNP proteins. Cell
90: 1013–1021,

[161554]1997.

[161555]

[161556]

[161557]15955.

[161558]Lorson, C. L.; Androphy, E. J.: The domain encoded by
exon 2

[161559]of the survival motor neuron protein mediates nucleic acid
binding. Hum.

[161560]Molec. Genet. 7: 1269–1275, 1998.

[161561]

[161562]

[161563]15956.

[161564]Lorson, C. L.; Hahnen, E.; Androphy, E. J.; Wirth, B.: A single
gle

[161565]nucleotide in the SMN gene regulates splicing and is responsible for

[161566]spinal muscular atrophy. Proc. Nat. Acad. Sci. 96:
6307–6311, 1999.

[161567]

[161568]

[161569]15957.

[161570]Lorson, C. L.; Strasswimmer, J.; Yao, J.–M.; Baleja, J. D.;
Hahnen,

[161571]E.; Wirth, B.; Le, T.; Burghes, A. H. M.; Androphy, E. J.: SMN
oligomerization

[161572]defect correlates with spinal muscular atrophy severity.
Nature Genet. 19:

[161573]63–66, 1998.

[161574]

[161575]

[161576]15958.

[161577]McAndrew, P. E.; Parsons, D. W.; Simard, L. R.; Rochette, C.;

[161578]Ray, P. N.; Mendell, J. R.; Prior, T. W.; Burghes, A. H.: Identification

[161579]of proximal spinal muscular atrophy carriers and patients by analysis

[161580]of SMNT and SMNC gene copy number. Am. J. Hum. Genet. 60: 1411–1422,

[161581]1997.

[161582]

[161583]

[161584]15959.

[161585]Meister, G.; Buhler, D.; Laggerbauer, B.; Zobawa, M.; Lottspeich,

[161586]F.; Fischer, U.: Characterization of a nuclear 20S complex containing

[161587]the survival of motor neurons (SMN) protein and a specific subset

[161588]of spliceosomal Sm proteins. Hum. Molec. Genet. 9: 1977–1986, 2000.

[161589]

[161590]

[161591]15960.

[161592]Mohaghegh, P.; Rodrigues, N. R.; Owen, N.; Ponting, C. P.;

Le,

[161593]T. T.; Burghes, A. H. M.; Davies, K. E.: Analysis of mutations in

[161594]the tudor domain of the survival motor neuron protein SMN. Europ.

[161595]J. Hum. Genet. 7: 519–525, 1999.

[161596]

[161597]

[161598]15961.

[161599]Monani, U. R.; Lorson, C. L.; Parsons, D. W.; Prior, T. W.;

Androphy,

[161600]E. J.; Burghes, A. H. M.; McPherson, J. D.: A single nucleotide difference

[161601]that alters splicing patterns distinguishes the SMA gene SMN1 from

[161602]the copy gene SMN2. Hum. Molec. Genet. 8: 1177–1183, 1999.

[161603]

[161604]

[161605]15962.

[161606]Mourelatos, Z.; Abel, L.; Yong, J.; Kataoka, N.; Dreyfuss, G.

[161607]: SMN interacts with a novel family of hnRNP and spliceosomal proteins. EMBO

[161608]J. 20: 5443–5452, 2001.

[161609]

[161610]

[161611]15963.

[161612]Owen, N.; Doe, C. L.; Mellor, J.; Davies, K. E.: Characterization

[161613]of the Schizosaccharomyces pombe orthologue of the human survival

[161614]motor neuron (SMN) protein. Hum. Molec. Genet. 9: 675–684, 2000.

[161615]

[161616]

[161617]15964.

[161618]Pagliardini, S.; Giavazzi, A.; Setola, V.; Lizier, C.; Di Luca,

[161619]M.; DeBiasi, S.; Battaglia, G.: Subcellular localization and axonal

[161620]transport of the survival motor neuron (SMN) protein in the developing

[161621]rat spinal cord. Hum. Molec. Genet. 9: 47–56, 2000.

[161622]

[161623]

[161624]15965.

[161625]Parsons, D. W.; McAndrew, P. E.; Iannaccone, S. T.; Mendell, J.

[161626]R.; Burghes, A. H. M.; Prior, T. W.: Intragenic telSMN mutations:

[161627]frequency, distribution, evidence of a founder effect, and modification

[161628]of the spinal muscular atrophy phenotype by cenSMN copy number. Am.

[161629]J. Hum. Genet. 63: 1712–1723, 1998.

[161630]

[161631]

[161632]15966.

[161633]Parsons, D. W.; McAndrew, P. E.; Monani, U. R.; Mendell, J. R.;

[161634]Burghes, A. H. M.; Prior, T. W.: An 11 base pair duplication in exon

[161635]6 of the SMN gene produces a type I spinal muscular atrophy (SMA)

[161636]phenotype: further evidence for SMN as the primary SMA-determining

[161637]gene. Hum. Molec. Genet. 5: 1727–1732, 1996.

[161638]

[161639]

[161640]15967.

[161641]Lewandoski, M.; Sun, X.; Martin, G. R.: Fgf8 signalling from the

[161642]AER is essential for normal limb development. Nature Genet. 26:

[161643]460–463, 2000.

[161644]

[161645]

[161646]15968.

[161647]Lorenzi, M. V.; Long, J. E.; Miki, T.; Aaronson, S. A.: Ex-
pression

[161648]cloning, developmental expression and chromosomal lo-
calization of

[161649]fibroblast growth factor-8. Oncogene 10: 2051-2055,
1995.

[161650]

[161651]

[161652]15969.

[161653]Meyers, E. N.; Martin, G. R.: Differences in left-right axis

[161654]pathways in mouse and chick: functions of FGF8 and SHH.
Science 285:

[161655]403-406, 1999.

[161656]

[161657]

[161658]15970.

[161659]Moon, A. M.; Capecchi, M. R.: Fgf8 is required for out-
growth

[161660]and patterning of the limbs. Nature Genet. 26: 455-459,
2000.

[161661]

[161662]

[161663]15971.

[161664]Payson, R. A.; Wu, J.; Liu, Y.; Chiu, I.-M.: The human FGF-8

[161665]gene localizes on chromosome 10q24 and is subjected to induction by

[161666]androgen in breast cancer cells. Oncogene 13: 47-53, 1996.

[161667]

[161668]

[161669]15972.

[161670]Streit, A.; Berliner, A. J.; Papanayotou, C.; Sirulnik, A.; Stern,

[161671]C. D.: Initiation of neural induction by FGF signalling before gastrulation. Nature 406:

[161672]74-78, 2000.

[161673]

[161674]

[161675]15973.

[161676]Sun, X.; Meyers, E. N.; Lewandoski, M.; Martin, G. R.: Tar-
geted

[161677]disruption of Fgf8 causes failure of cell migration in the
gastrulating

[161678]mouse embryo. Genes Dev. 13: 1834–1846, 1999.

[161679]

[161680]

[161681]15974.

[161682]White, R. A.; Dowler, L. L.; Angeloni, S. V.; Pasztor, L. M.;

[161683]MacArthur, C. A.: Assignment of FGF8 to human chromo-
some 10q25–q26:

[161684]mutations in FGF8 may be responsible for some types of
acrocephalosyndactyly

[161685]linked to this region. Genomics 30: 109–111, 1995.

[161686]

[161687]

[161688]15975.

[161689]Yoshiura, K.; Leysens, N. J.; Chang, J.; Ward, D.; Murray, J.

[161690]C.; Muenke, M.: Genomic structure, sequence, and map–

ping of human

[161691]FGF8 with no evidence for its role in craniosynostosis/
limb defect

[161692]syndromes. Am. J. Med. Genet. 72: 354–362, 1997.

[161693]

[161694]

[161695]15976.

[161696]Zammit, C.; Coope, R.; Gomm, J. J.; Shousha, S.; Johnston,
C.

[161697]L.; Coombes, R. C.: Fibroblast growth factor 8 is ex-
pressed at higher

[161698]levels in lactating human breast and in breast cancer. Brit.
J. Cancer 86:

[161699]1097–1103, 2002.

[161700]

[161701]

[161702]15977.

[161703]Cao, H.; Mok, A.; Miskie, B.; Hegele, R. A.: Single-nu-
cleotide

[161704]polymorphisms of the proprotein convertase subtilisin/

kexin type 5

[161705](PCSK5) gene. J. Hum. Genet. 46: 730–732, 2001.

[161706]

[161707]

[161708]15978.

[161709]van de Loo, J.–W. H. P.; Creemers, J. W. M.; Kas, K.; Roe–
broek,

[161710]A. J. M.; Van de Ven, W. J. M.: Assignment of the human
proprotein

[161711]convertase gene PCSK5 to chromosome 9q21.3 Cyto–
genet. Cell Genet. 75:

[161712]227–229, 1996.

[161713]

[161714]

[161715]15979.

[161716]Aramburu, J.; Garcia–Cozar, F.; Raghavan, A.; Okamura,
H.; Rao,

[161717]A.; Hogan, P. G.: Selective inhibition of NFAT activation by
a peptide

[161718]spanning the calcineurin targeting site of NFAT. Molec.

Cell 1:

[161719]627–637, 1998.

[161720]

[161721]

[161722]15980.

[161723]Castigli, E.; Pahwa, R.; Good, R. A.; Geha, R. S.; Chatila, T.

[161724]A.: Molecular basis of a multiple lymphokine deficiency in
a patient

[161725]with severe combined immunodeficiency. Proc. Nat. Acad.
Sci. 90:

[161726]4728–4732, 1993.

[161727]

[161728]

[161729]15981.

[161730]Chatila, T.; Castigli, E.; Pahwa, R.; Pahwa, S.; Chirmule, N.;

[161731]Oyaizu, N.; Good, R. A.; Geha, R. S.: Primary combined im-
munodeficiency

[161732]resulting from defective transcription of multiple T-cell
lymphokine

[161733]genes. Proc. Nat. Acad. Sci. 87: 10033–10037, 1990.

[161734]

[161735]

[161736]15982.

[161737]Horsley, V.; Pavlath, G. K.: NFAT: ubiquitous regulator of
cell

[161738]differentiation and adaptation. J. Cell Biol. 156: 771–774,
2002.

[161739]

[161740]

[161741]15983.

[161742]Jauliac, S.; Lopez–Rodriguez, C.; Shaw, L. M.; Brown, L. F.;
Rao,

[161743]A.; Toker, A.: The role of NFAT transcription factors in in-
tegrin–mediated

[161744]carcinoma invasion. Nature Cell Biol. 4: 540–544, 2002.

[161745]

[161746]

[161747]15984.

[161748]Li, X.; Ho, S. N.; Luna, J.; Giacalone, J.; Thomas, D. J.; Timmerman,

[161749]L. A.; Crabtree, G. R.; Francke, U.: Cloning and chromosomal localization

[161750]of the human and murine genes for the T-cell transcription factors

[161751]NFATc and NFATp. Cytogenet. Cell Genet. 68: 185–191, 1995.

[161752]

[161753]

[161754]15985.

[161755]Northrop, J. P.; Ho, S. N.; Chen, L.; Thomas, D. J.; Timmerman,

[161756]L. A.; Nolan, G. P.; Admon, A.; Crabtree, G. R.: NF-AT components

[161757]define a family of transcription factors targeted in T-cell activation. Nature 369:

[161758]497–502, 1994.

[161759]

[161760]

[161761]15986.

[161762]Okamura, H.; Aramburu, J.; Garcia-Rodriguez, C.; Viola, J. P. B.;

[161763]Raghavan, A.; Tahiliani, M.; Zhang, X.; Qin, J.; Hogan, P. G.; Rao,

[161764]A.: Concerted dephosphorylation of the transcription factor NFAT1

[161765]induces a conformational switch that regulates transcriptional activity. Molec.

[161766]Cell 6: 539–550, 2000.

[161767]

[161768]

[161769]15987.

[161770]Pahwa, R.; Chatila, T.; Pahwa, S.; Paradise, C.; Day, N. K.; Geha,

[161771]R.; Schwartz, S. A.; Slade, H.; Oyaizu, N.; Good, R. A.: Recombinant

[161772]interleukin 2 therapy in severe combined immunodeficiency disease. Proc.

[161773]Nat. Acad. Sci. 86: 5069–5073, 1989.

[161774]

[161775]

[161776]15988.

[161777]Park, J.; Takeuchi, A.; Sharma, S.: Characterization of a
new

[161778]isoform of the NFAT (nuclear factor of activated T cells)
gene family

[161779]member NFATc. J. Biol. Chem. 271: 20914–20921, 1996.

[161780]

[161781]

[161782]15989.

[161783]Peng, S. L.; Gerth, A. J.; Ranger, A. M.; Glimcher, L. H.:
NFATc1

[161784]and NFATc2 together control both T and B cell activation
and differentiation. Immunity 14:

[161785]13–20, 2001.

[161786]

[161787]

[161788]15990.

[161789]Aoki, K.; Suzuki, K.; Sugano, T.; Tasaka, T.; Nakahara, K.;
Kuge,

[161790]O.; Omori, A.; Kasai, M.: A novel gene, 'Translin,' encodes

a recombination

[161791]hotspot binding protein associated with chromosomal
translocations. Nature

[161792]Genet. 10: 167–174, 1995.

[161793]

[161794]

[161795]15991.

[161796]Badge, R. M.; Yardley, J.; Jeffreys, A. J.; Armour, J. A. L.:

[161797]Crossover breakpoint mapping identifies a subtelomeric
hotspot for

[161798]male meiotic recombination. Hum. Molec. Genet. 9:
1239–1244, 2000.

[161799]

[161800]

[161801]15992.

[161802]Hosaka, T.; Kanoe, H.; Nakayama, T.; Murakami, H.; Ya-
mamoto, H.;

[161803]Nakamata, T.; Tsuboyama, T.; Oka, M.; Kasai, M.; Sasaki,
M. S.; Nakamura,

[161804]T.; Toguchida, J.: Translin binds to the sequences adjacent

to the

[161805]breakpoints of the TLS and CHOP genes in liposarcomas
with translocation

[161806]t(12;16). Oncogene 19: 5821–5825, 2000.

[161807]

[161808]

[161809]15993.

[161810]Kasai, M.; Aoki, K.; Matsuo, Y.; Minowada, J.; Maziarz, R.
T.;

[161811]Strominger, J. L.: Recombination hotspot associated fac-
tors specifically

[161812]recognize novel target sequences at the site of interchro-
mosomal rearrangements

[161813]in T-ALL patients with t(8;14)(q24;q11)and
(t(1;14)(p32;q11). Int.

[161814]Immun. 6: 1017–1025, 1994.

[161815]

[161816]

[161817]15994.

[161818]Arceci, R. J.; King, A. A.; Simon, M. C.; Orkin, S. H.; Wilson,

[161819]D. B.: Mouse GATA-4: a retinoic acid-inducible GATA-binding transcription

[161820]factor expressed in endodermally derived tissues and heart. Molec.

[161821]Cell. Biol. 13: 2235-2246, 1993.

[161822]

[161823]

[161824]15995.

[161825]Cirillo, L. A.; Lin, F. R.; Cuesta, I.; Friedman, D.; Jarnik, M.;

[161826]Zaret, K. S.: Opening of compacted chromatin by early developmental

[161827]transcription factors HNF3(FoxA) and GATA-4. Molec. Cell
9: 279-289,

[161828]2002.

[161829]

[161830]

[161831]15996.

[161832]Crispino, J. D.; Lodish, M. B.; Thurberg, B. L.; Litovsky, S.
H.;

[161833]Collins, T.; Molkentin, J. D.; Orkin, S. H.: Proper coronary

vascular

[161834]development and heart morphogenesis depend on interaction of GATA-4

[161835]with FOG cofactors. Genes Dev. 15: 839-844, 2001.

[161836]

[161837]

[161838]15997.

[161839]Hasegawa, K.; Lee, S. J.; Jobe, S. M.; Markham, B. E.; Kitsis,

[161840]R. N.: cis-acting sequences that mediate induction of beta-myosin

[161841]heavy chain gene expression during left ventricular hypertrophy due

[161842]to aortic constriction. Circulation 96: 3943-3953, 1997.

[161843]

[161844]

[161845]15998.

[161846]Huang, W.-Y.; Cukerman, E.; Liew, C.-C.: Identification of a GATA

[161847]motif in the cardiac alpha-myosin heavy-chain-encoding gene and isolation

[161848]of a human GATA-4 cDNA. Gene 155: 219-223, 1995.

[161849]

[161850]

[161851]15999.

[161852]Huang, W.-Y.; Heng, H. H. Q.; Liew, C.-C.: Assignment of
the human

[161853]GATA4 gene to 8p23.1-p22 using fluorescence in situ hy-
bridization

[161854]analysis. Cytogenet. Cell Genet. 72: 217-218, 1996.

[161855]

[161856]

[161857]16000.

[161858]Kennedy, S. J.; Teebi, A. S.; Adatia, I.; Teshima, I.: Inher-
ited

[161859]duplication, dup(8)(p23.1p23.1)Pat, in a father and
daughter with

[161860]congenital heart defects. (Letter) Am. J. Med. Genet. 104:
79-80,

[161861]2001.

[161862]

[161863]

[161864]16001.

[161865]Ketola, I.; Pentikainen, V.; Vaskivuo, T.; Ilvesmaki, V.;

Herva,

[161866]R.; Dunkel, L.; Tapanainen, J. S.; Toppari, J.; Heikinheimo,

M.:

[161867]Expression of transcription factor GATA-4 during human
testicular

[161868]development and disease. J. Clin. Endocr. Metab. 85:

3925-3931,

[161869]2000.

[161870]

[161871]

[161872]16002.

[161873]Laitinen, M. P. E.; Anttonen, M.; Ketola, I.; Wilson, D. B.;

Ritvos,

[161874]O.; Butzow, R.; Heikinheimo, M.: Transcription factors

GATA-4 and

[161875]GATA-6 and a GATA family cofactor, FOG-2, are ex-
pressed in human ovary

[161876]and sex cord-derived ovarian tumors. J. Clin. Endocr.

Metab. 85:

[161877]3476–3483, 2000.

[161878]

[161879]

[161880]16003.

[161881]Molkentin, J. D.; Kalvakolanu, D. V.; Markham, B. E.: Tran-
scription

[161882]factor GATA-4 regulates cardiac muscle-specific expres-
sion of the

[161883]alpha-myosin heavy-chain gene. Molec. Cell. Biol. 14:
4947–4957,

[161884]1994.

[161885]

[161886]

[161887]16004.

[161888]Pehlivan, T.; Pober, B. R.; Brueckner, M.; Garrett, S.;
Slaugh,

[161889]R.; Van Rheeden, R.; Wilson, D. B.; Watson, M. S.; Hing, A.
V.: GATA4

[161890]haploinsufficiency in patients with interstitial deletion of

chromosome

[161891]region 8p23.1 and congenital heart disease. Am. J. Med.

Genet. 83:

[161892]201–206, 1999.

[161893]

[161894]

[161895]16005.

[161896]White, R. A.; Dowler, L. L.; Pasztor, L. M.; Gatson, L. L.;
Adkison,

[161897]L. R.; Angeloni, S. V.; Wilson, D. B.: Assignment of the
transcription

[161898]factor GATA4 gene to human chromosome 8 and mouse
chromosome 14: Gata4

[161899]is a candidate gene for Ds (disorganization). Genomics 27:
20–26,

[161900]1995.

[161901]

[161902]

[161903]16006.

[161904]Brenneman, M. A.; Wagener, B. M.; Miller, C. A.; Allen, C.;

Nickoloff,

[161905]J. A.: XRCC3 controls the fidelity of homologous recombination: roles

[161906]for XRCC3 in late stages of recombination. Molec. Cell 10: 387–395,

[161907]2002.

[161908]

[161909]

[161910]16007.

[161911]Liu, N.; Lamerdin, J. E.; Tebbs, R. S.; Schild, D.; Tucker, J.

[161912]D.; Shen, M. R.; Brookman, K. W.; Siciliano, M. J.; Walter, C. A.;

[161913]Fan, W.; Narayana, L. S.; Zhou, Z.-Q.; Adamson, A. W.; Sorensen, K.

[161914]J.; Chen, D. J.; Jones, N. J.; Thompson, L. H.: XRCC2 and XRCC3,

[161915]new human Rad51–family members, promote chromosome stability and protect

[161916]against DNA cross–links and other damages. Molec. Cell 1: 783–793,

[161917]1998.

[161918]

[161919]

[161920]16008.

[161921]Tebbs, R. S.; Zhao, Y.; Tucker, J. D.; Scheerer, J. B.; Siciliano,

[161922]M. J.; Hwang, M.; Liu, N.; Legerski, R. J.; Thompson, L. H.:
Correction

[161923]of chromosomal instability and sensitivity to diverse mutagens by

[161924]a cloned cDNA of the XRCC3 DNA repair gene. Proc. Nat. Acad. Sci. 92:

[161925]6354–6358, 1995.

[161926]

[161927]

[161928]16009.

[161929]Winsey, S. L.; Haldar, N. A.; Marsh, H. P.; Bunce, M.; Marshall,

[161930]S. E.; Harris, A. L.; Wojnarowska, F.; Welsh, K. I.: A variant within

[161931]the DNA repair gene XRCC3 is associated with the development of melanoma

[161932]skin cancer. Cancer Res. 60: 5612–5616, 2000.

[161933]

[161934]

[161935]16010.

[161936]Hayashida, S.; Yamasaki, K.; Asada, Y.; Soeda, E.; Niikawa, N.;

[161937]Kishino, T.: Construction of a physical and transcript map flanking

[161938]the imprinted MEST/PEG1 region at 7q32. Genomics 66: 221–225, 2000.

[161939]

[161940]

[161941]16011.

[161942]de Wind, N.; Dekker, M.; Claij, N.; Jansen, L.; van Klink, Y.;

[161943]Radman, M.; Riggins, G.; van der Valk, M.; van't Wout, K.; te Riele,

[161944]H.: HNPCC–like cancer predisposition in mice through simultaneous

[161945]loss of Msh3 and Msh6 mismatch–repair protein functions. Nature Genet. 23:

[161946]359–362, 1999.

[161947]

[161948]

[161949]16012.

[161950]Farrelly, D.; Brown, K. S.; Tieman, A.; Ren, J.; Lira, S. A.;
Hagan,

[161951]D.; Gregg, R.; Mookhtiar, K. A.; Hariharan, N.: Mice mutant
for glucokinase

[161952]regulatory protein exhibit decreased liver glucokinase: a
sequestration

[161953]mechanism in metabolic regulation. Proc. Nat. Acad. Sci.
96: 14511–14516,

[161954]1999.

[161955]

[161956]

[161957]16013.

[161958]Braun–Dullaеus, R. C.; Mann, M. J.; Ziegler, A.; von der
Leyen,

[161959]H. E.; Dzaу, V. J.: A novel role for the cyclin–dependent ki-
nase

[161960]inhibitor p27(Kip1) in angiotensin II–stimulated vascular smooth muscle

[161961]cell hypertrophy. J. Clin. Invest. 104: 815–823, 1999.

[161962]

[161963]

[161964]16014.

[161965]Carrano, A. C.; Eytan, E.; Hershko, A.; Pagano, M.: SKP2 is required

[161966]for ubiquitin–mediated degradation of the CDK inhibitor p27. Nature

[161967]Cell Biol. 1: 193–199, 1999.

[161968]

[161969]

[161970]16015.

[161971]Di Cristofano, A.; De Acetis, M.; Koff, A.; Cordon–Cardo, C.; Pandolfi,

[161972]P. P.: Pten and p27(KIP1) cooperate in prostate cancer tumor suppression

[161973]in the mouse. Nature Genet. 27: 222–224, 2001.

[161974]

[161975]

[161976]16016.

[161977]Fero, M. L.; Rivkin, M.; Tasch, M.; Porter, P.; Carow, C. E.;

Firpo,

[161978]E.; Polyak, K.; Tsai, L.-H.; Broudy, V.; Perlmutter, R. M.;

Kaushansky,

[161979]K.; Roberts, J. M.: A syndrome of multiorgan hyperplasia
with features

[161980]of gigantism, tumorigenesis, and female sterility in
p27(Kip1)-deficient

[161981]mice. Cell 85: 733-744, 1996.

[161982]

[161983]

[161984]16017.

[161985]Albertella, M. R.; Campbell, R. D.: Characterization of a
novel

[161986]gene in the human major histocompatibility complex that
encodes a

[161987]potential new member of the I kappa B family of proteins.

Hum. Molec.

[161988]Genet. 3: 793-799, 1994.

[161989]

[161990]

[161991]16018.

[161992]Buyse, I. M.; Shao, G.; Huang, S.: The retinoblastoma protein

[161993]binds to RIZ, a zinc-finger protein that shares an epitope with the

[161994]adenovirus E1A protein. Proc. Nat. Acad. Sci. 92: 4467-4471, 1995.

[161995]

[161996]

[161997]16019.

[161998]Buyse, I. M.; Takahashi, E.; Huang, S.: Physical mapping of the

[161999]retinoblastoma interacting zinc finger gene RIZ to D1S228 on chromosome

[162000]1p36. Genomics 34: 119-121, 1996.

[162001]

[162002]

[162003]16020.

[162004]Chadwick, R. B.; Jiang, G.-L.; Bennington, G. A.; Yuan, B.;
Johnson,

[162005]C. K.; Stevens, M. W.; Niemann, T. H.; Peltomaki, P.;
Huang, S.; de

[162006]la Chapelle, A.: Candidate tumor suppressor RIZ is fre-
quently involved

[162007]in colorectal carcinogenesis. Proc. Nat. Acad. Sci. 97:
2662-2667,

[162008]2000.

[162009]

[162010]

[162011]16021.

[162012]Mock, B. A.; Coleman, M. P.; Huang, S.: Riz maps to distal
chromosome

[162013]4 near genes involved in tumorigenesis and nerve degen-
eration. Mammalian

[162014]Genome 7: 637 only, 1996.

[162015]

[162016]

[162017]16022.

[162018]Poetsch, M.; Dittberner, T.; Woenckhaus, C.: Frameshift mutations

[162019]of RIZ, but no point mutations in RIZ1 exons in malignant melanomas

[162020]with deletions in 1p36. Oncogene 21: 3038–3042, 2002.

[162021]

[162022]

[162023]16023.

[162024]Malek, N. P.; Sundberg, H.; McGrew, S.; Nakayama, K.; Kyriakidis,

[162025]T. R.; Roberts, J. M.: A mouse knock-in model exposes sequential

[162026]proteolytic pathways that regulate p27(Kip1) in G1 and S phase. Nature 413:

[162027]323–327, 2001.

[162028]

[162029]

[162030]16024.

[162031]Martin, E.; Cacheux, V.; Cave, H.; Lapierre, J. M.; Le\Paslier,

[162032]D.; Grandchamp, B.: Localization of the CDKN4/p27(Kip1) gene to human

[162033]chromosome 12p12.3. Hum. Genet. 96: 668–670, 1995.

[162034]

[162035]

[162036]16025.

[162037]Mitsuhashi, T.; Aoki, Y.; Eksioglu, Y. Z.; Takahashi, T.; Bhide,

[162038]P. G.; Reeves, S. A.; Caviness, V. S., Jr.: Overexpression of p27(Kip1)

[162039]lengthens the G1 phase in a mouse model that targets inducible gene

[162040]expression to central nervous system progenitor cells.

Proc. Nat.

[162041]Acad. Sci. 98: 6435–6440, 2001.

[162042]

[162043]

[162044]16026.

[162045]Polyak, K.; Lee, M.–H.; Erdjument–Bromage, H.; Koff, A.;

Roberts,

[162046]J. M.; Tempst, P.; Massague, J.: Cloning of p27(Kip1), a cy–

clin–dependent

[162047]kinase inhibitor and a potential mediator of extracellular

antimitogenic

[162048]signals. Cell 78: 59–66, 1994.

[162049]

[162050]

[162051]16027.

[162052]Saito, T.; Seki, N.; Hattori, A.; Hayashi, A.; Abe, M.; Araki,

[162053]R.; Fujimori, A.; Fukumura, R.; Kozuma, S.; Matsuda, Y.:

Structure,

[162054]expression profile, and chromosomal location of a mouse

gene homologous

[162055]to human DNA-PK(cs) interacting protein (KIP) gene.

Mammalian Genome 10:

[162056]315–317, 1999.

[162057]

[162058]

[162059]16028.

[162060]Sherr, C. J.; Roberts, J. M.: Inhibitors of mammalian G1 cyclin-dependent

[162061]kinases. Genes Dev. 9: 1149–1163, 1995.

[162062]

[162063]

[162064]16029.

[162065]Toyoshima, H.; Hunter, T.: p27, a novel inhibitor of G1 cyclin-Cdk

[162066]protein kinase activity, is related to p21. Cell 78: 67–74, 1994.

[162067]

[162068]

[162069]16030.

[162070]Hudson, A. W.; Birnbaum, M. J.: Identification of a non-neuronal

[162071]isoform of synaptotagmin. Proc. Nat. Acad. Sci. 92: 5895–5899, 1995.

[162072]

[162073]

[162074]16031.

[162075]Warner, J. P.; Leek, J. P.; Intody, S.; Markham, A. F.; Bonthron,

[162076]D. T.: Human glucokinase regulatory protein (GCKR): cDNA and genomic

[162077]cloning, complete primary structure, and chromosomal localization. Mammalian

[162078]Genome 6: 532–536, 1995.

[162079]

[162080]

[162081]16032.

[162082]Chen, C.–C.; Akopian, A. N.; Sivilotti, L.; Colquhoun, D.; Burnstock,

[162083]G.; Wood, J. N.: A P2X purinoceptor expressed by a subset
of sensory

[162084]neurons. Nature 377: 428–430, 1995.

[162085]

[162086]

[162087]16033.

[162088]Cockayne, D. A.; Hamilton, S. G.; Zhu, Q.–M.; Dunn, P. M.;
Zhong,

[162089]Y.; Novakovic, S.; Malmberg, A. B.; Cain, G.; Berson, A.;
Kassotakis,

[162090]L.; Hedley, L.; Lachnit, W. G.; Burnstock, G.; McMahon, S.
B.; Ford,

[162091]A. P. D. W.: Urinary bladder hyporeflexia and reduced
pain–related

[162092]behaviour in P2X3–deficient mice. Nature 407:
1011–1015, 2000.

[162093]

[162094]

[162095]16034.

[162096]Vaxillaire, M.; Vionnet, N.; Vigouroux, C.; Sun, F.; Es–
pinosa,

[162097]R., III; LeBeau, M. M.; Stoffel, M.; Lehto, M.; Beckmann, J.
S.; Detheux,

[162098]M.; Passa, P.; Cohen, D.; Van Schaftingen, E.; Velho, G.;
Bell, G.

[162099]I.; Froguel, P.: Search for a third susceptibility gene for
maturity-onset

[162100]diabetes of the young: studies with eleven candidate
genes. Diabetes 43:

[162101]389–395, 1994.

[162102]

[162103]

[162104]16035.

[162105]Cook, S. P.; McCleskey, E. W.: ATP, pain and a full bladder.
Nature 407:

[162106]951–952, 2000.

[162107]

[162108]

[162109]16036.

[162110]Garcia-Guzman, M.; Stuhmer, W.; Soto, F.: Molecular char-
acterization

[162111]and pharmacological properties of the human P2X3
purinoceptor. Molec.

[162112]Brain Res. 47 –59–66, 1997.

[162113]

[162114]

[162115]16037.

[162116]Kennedy, C.; Leff, P.: Painful connection for ATP. Nature
377:

[162117]385–386, 1995.

[162118]

[162119]

[162120]16038.

[162121]Lewis, C.; Neldhart, S.; Holy, C.; North, R. A.; Buell, G.;
Surprenant,

[162122]A.: Coexpression of P2X2 and P2X3 receptor subunits can
account for

[162123]ATP-gated currents in sensory neurons. Nature 377:
432–435, 1995.

[162124]

[162125]

[162126]16039.

[162127]Souslova, V.; Cesare, P.; Ding, Y.; Akopian, A. N.; Stanfa, L.;

[162128]Suzuki, R.; Carpenter, K.; Dickenson, A.; Boyce, S.; Hill, R.; Nebenius–Oosthuizen,

[162129]D.; Smith, A. J. H.; Kidd, E. J.; Wood, J. N.: Warm–coding deficits

[162130]and aberrant inflammatory pain in mice lacking P2X3 receptors. Nature 407:

[162131]1015–1017, 2000.

[162132]

[162133]

[162134]16040.

[162135]Souslova, V.; Ravenall, S.; Fox, M.; Wells, D.; Wood, J. N.; Akopian,

[162136]A. N.: Structure and chromosomal mapping of the mouse P2X(3) gene. Gene 195:

[162137]101–111, 1997.

[162138]

[162139]

[162140]16041.

[162141]Khakh, B. S.; Zhou, X.; Sydes, J.; Galligan, J. J.; Lester, H.

[162142]A.: State-dependent cross-inhibition between transmitter-gated cation

[162143]channels. *Nature* 406: 405–410, 2000.

[162144]

[162145]

[162146]16042.

[162147]Buell, G.; Lewis, C.; Collo, G.; North, R. A.; Surprenant, A.:

[162148]An antagonist-insensitive P2X receptor expressed in epithelia and

[162149]brain. *EMBO J.* 15: 55–62, 1996.

[162150]

[162151]

[162152]16043.

[162153]Buell, G.; Lewis, C.; Collo, G.; North, R. A.; Surprenant, A.:

[162154]Cloning and expression of a family of ATP-gated ion channels. (Abstract) Soc.

[162155]*Neurosci.* 21: 1262, 1995.

[162156]

[162157]

[162158]16044.

[162159]Garcia-Guzman, M.; Soto, F.; Gomez-Hernandez, J. M.;
Lund, P. E.;

[162160]Stuhmer, W.: Characterization of the recombinant human
P2X4 receptor

[162161]reveals pharmacological differences to the rat homologue.
Molec.

[162162]Pharm. 51: 109–118, 1997.

[162163]

[162164]

[162165]16045.

[162166]Chen, J. D.; Evans, R. M.: A transcriptional co-repressor
that

[162167]interacts with nuclear hormone receptors. Nature 377:
454–457, 1995.

[162168]

[162169]

[162170]16046.

[162171]Fischle, W.; Dequiedt, F.; Hendzel, M. J.; Guenther, M. G.; Lazar,

[162172]M. A.; Voelter, W.; Verdin, E.: Enzymatic activity associated with

[162173]class II HDACs is dependent on a multiprotein complex containing HDAC3

[162174]and SMRT/N-CoR. Molec. Cell 9: 45–57, 2002.

[162175]

[162176]

[162177]16047.

[162178]Horlein, A. J.; Naar, A. M.; Heinzl, T.; Torchia, J.; Gloss, B.;

[162179]Kurokawa, R.; Ryan, A.; Kamel, Y.; Soderstrom, M.; Glass, C. K.; Rosenfeld,

[162180]M. G.: Ligand-independent repression by the thyroid hormone receptor

[162181]mediated by a nuclear receptor co-repressor. Nature 377: 397–403,

[162182]1995.

[162183]

[162184]

[162185]16048.

[162186]Hu, X.; Lazar, M. A.: The CoRNR motif controls the recruitment

[162187]of corepressors by nuclear hormone receptors. Nature
402: 93–96,

[162188]1999.

[162189]

[162190]

[162191]16049.

[162192]Jiang, Q.; Galiegue–Zouitina, S.; Roumier, C.; Hildebrand,
M. P.;

[162193]Thomas, S.; Coignet, L. J.: Genomic organization and re–
fined mapping

[162194]of the human nuclear corepressor 2 (NCOR2)/ silencing
mediator of

[162195]retinoid and thyroid hormone receptor (SMRT) gene on
chromosome 12q24.3. Cytogenet.

[162196]Cell Genet. 92: 217–220, 2001.

[162197]

[162198]

[162199]16050.

[162200]Ordentlich, P.; Downes, M.; Xie, W.; Genin, A.; Spinner, N. B.;

[162201]Evans, R. M.: Unique forms of human and mouse nuclear receptor corepressor

[162202]SMRT. Proc. Nat. Acad. Sci. 96: 2639–2644, 1999.

[162203]

[162204]

[162205]16051.

[162206]Colvin, J. S.; Green, R. P.; Schmahl, J.; Capel, B.; Ornitz, D.

[162207]M.: Male-to-female sex reversal in mice lacking fibroblast growth

[162208]factor 9. Cell 104: 875–889, 2001.

[162209]

[162210]

[162211]16052.

[162212]Mattei, M.-G.; Penault-Llorca, F.; Coulier, F.; Birnbaum, D.:

[162213]The human FGF9 gene maps to chromosomal region 13q11–q12. Genomics 29:

[162214]811–812, 1995.

[162215]

[162216]

[162217]16053.

[162218]Miyamoto, M.; Naruo, K.–I.; Seko, C.; Matsumoto, S.;
Kondo, T.;

[162219]Kurokawa, T.: Molecular cloning of a novel cytokine cDNA
encoding

[162220]the ninth member of the fibroblast growth factor family,
which has

[162221]a unique secretion property. Molec. Cell. Biol. 13:
4251–4259, 1993.

[162222]

[162223]

[162224]16054.

[162225]Garcia, J. G.; Lazar, V.; Gilbert–McClain, L. I.; Gallagher, P.

[162226]J.; Verin, A. D.: Myosin light chain kinase in endothelium:
molecular

[162227]cloning and regulation. Am. J. Resp. Cell Molec. Biol. 16:
489–494,

[162228]1997.

[162229]

[162230]

[162231]16055.

[162232]Giorgi, D.; Brand-Arpon, V.; Rouquier, S.: The functional
myosin

[162233]light chain kinase (MYLK) gene localizes with marker
D3S3552 on human

[162234]chromosome 3q21 in a greater than 5-Mb yeast artificial
chromosome

[162235]region and is not linked to olfactory receptor genes. Cyto-
genet.

[162236]Cell Genet. 92: 85-88, 2001.

[162237]

[162238]

[162239]16056.

[162240]Lazar, V.; Garcia, J. G. N.: A single human myosin light
chain

[162241]kinase gene (MLCK; MYLK) transcribes multiple nonmuscle
isoforms. Genomics 57:

[162242]256–267, 1999.

[162243]

[162244]

[162245]16057.

[162246]Potier, M.–C.; Chelot, E.; Pekarsky, Y.; Gardiner, K.;
Rossier,

[162247]J.; Turnell, W. G.: The human myosin light chain kinase
(MLCK) from

[162248]hippocampus: cloning, sequencing, expression, and local-
ization to

[162249]3cen–q21. Genomics 29: 562–570, 1995.

[162250]

[162251]

[162252]16058.

[162253]Walker, L. A.; MacDonald, J. A.; Liu, X.; Nakamoto, R. K.;
Haystead,

[162254]T. A. J.; Somlyo, A. V.; Somlyo, A. P.: Site–specific phos-
phorylation

[162255]and point mutations of telokin modulate its
Ca(2+)–desensitizing effect

[162256]in smooth muscle. J. Biol. Chem. 276: 24519–24524,
2001.

[162257]

[162258]

[162259]16059.

[162260]Watterson, D. M.; Schavocky, J. P.; Guo, L.; Weiss, C.;
Chlenski,

[162261]A.; Shirinsky, V. P.; Van Eldik, L. J.; Haiech, J.: Analysis of
the

[162262]kinase-related protein gene found at human chromosome
3q21 in a multi-gene

[162263]cluster: organization, expression, alternative splicing, and
polymorphic

[162264]marker. J. Cell. Biochem. 75: 481–491, 1999.

[162265]

[162266]

[162267]16060.

[162268]Allcock, R. J. N.; Christiansen, F. T.; Price, P.: The central

[162269]MHC gene IKBL carries a structural polymorphism that is
associated

[162270]with HLA-A3,B7,DR15. Immunogenetics 49: 660-665,
1999.

[162271]

[162272]

[162273]16061.

[162274]Druck, T.; Gu, Y.; Prabhala. G.; Cannizzaro, L. A.; Park,
S.-H.;

[162275]Huebner, K.; Keen, J. H.: Chromosome localization of hu-
man genes

[162276]for clathrin adaptor polypeptides AP2-beta and AP50 and
the clathrin-binding

[162277]protein, VCP. Genomics 30: 94-97, 1995.

[162278]

[162279]

[162280]16062.

[162281]Egerton, M.; Ashe, O. R.; Chen, D.; Druker, B. J.: Burgess,
W.

[162282]H.: Samelson, L. E.:VCP, the mammalian homolog of
cdc48, is tyrosine

[162283]phosphorylated in response to T cell antigen receptor ac-
tivation. EMBO

[162284]J. 11: 3533–3540, 1992.

[162285]

[162286]

[162287]16063.

[162288]Hoyle, J.; Tan, K. H.; Fisher, E. M. C.: Mapping the valosin-containing

[162289]protein (VCP) gene on human chromosome 9 and mouse chromosome 4, and

[162290]a likely pseudogene on the mouse X chromosome. Mammalian Genome 8:

[162291]778–780, 1997.

[162292]

[162293]

[162294]16064.

[162295]Koller, K. J.; Brownstein, M. J.: Use of a cDNA clone to identify

[162296]a supposed precursor protein containing valosin. Nature 325: 542–545,

[162297]1987.

[162298]

[162299]

[162300]16065.

[162301]Pleasure, I. T.; Black, M. M.; Keen, J. H.: Valosin-containing

[162302]protein, VCP, is a ubiquitous clathrin-binding protein. Nature 365:

[162303]459–462, 1993.

[162304]

[162305]

[162306]16066.

[162307]Ye, Y.; Meyer, H. H.; Rapoport, T. A.: The AAA ATPase Cdc48/p97

[162308]and its partners transport proteins from the ER into the cytosol. Nature 414:

[162309]652–656, 2001.

[162310]

[162311]

[162312]16067.

[162313]Liu, Q.; Feng, Y.; Forgac, M.: Activity and in vitro reassembly

[162314]of the coated vesicle (H⁺)-ATPase requires the 50-kDa subunit of the

[162315]clathrin assembly complex AP-2. J. Biol. Chem. 269: 31592-31597,

[162316]1994.

[162317]

[162318]

[162319]16068.

[162320]Thurieu, C.; Brosius, J.; Burne, C.; Jolles, P.; Keen, J. H.;

[162321]Mattaliano, R. J.; Chow, E. P.; Ramachandran, K. L.; Kirchhausen,

[162322]T.: Molecular cloning and complete amino acid sequence of AP50, an

[162323]assembly protein associated with clathrin-coated vesicles.

DNA 7:

[162324]663-669, 1988.

[162325]

[162326]

[162327]16069.

[162328]Ponnambalam, S.; Robinson, M. S.; Jackson, A. P.; Peiperl,

L.;

[162329]Parham, P.: Conservation and diversity in families of coated vesicle

[162330]adaptins. J. Biol. Chem. 265: 4814–4820, 1990.

[162331]

[162332]

[162333]16070.

[162334]Robinson, M. S.: Cloning of cDNAs encoding two related 100-kD

[162335]coated vesicle proteins (alpha-adaptins). J. Cell Biol. 108: 833–42,

[162336]1989.

[162337]

[162338]

[162339]16071.

[162340]Waelter, S.; Scherzinger, E.; Hasenbank, R.; Nordhoff, E.; Lurz,

[162341]R.; Goehler, H.; Gauss, C.; Sathasivam, K.; Bates, G. P.; Lehrach,

[162342]H.; Wanker, E. E.: The huntingtin interacting protein HIP1

is a clathrin

[162343]and alpha-adaptin-binding protein involved in receptor-mediated endocytosis. Hum.

[162344]Molec. Genet. 10: 1807–1817, 2001.

[162345]

[162346]

[162347]16072.

[162348]Kaneko-Ishino, T.; Kuroiwa, Y.; Miyoshi, N.; Kohda, T.; Suzuki,

[162349]R.; Yokoyama, M.; Viville, S.; Barton, S. C.; Ishino, F.; Surani,

[162350]M. A.: Peg1/Mest imprinted gene on chromosome 6 identified by cDNA

[162351]subtraction hybridization. Nature Genet. 11: 52–59, 1995.

[162352]

[162353]

[162354]16073.

[162355]Kobayashi, S.; Kohda, T.; Miyoshi, N.; Kuroiwa, Y.; Aisaka, K.;

[162356]Tsutsumi, O.; Kaneko-Ishino, T.; Ishino, F.: Human

PEG1/MEST, an

[162357]imprinted gene on chromosome 7. Hum. Molec. Genet. 6:
781–786, 1997.

[162358]

[162359]

[162360]16074.

[162361]Kosaki, K.; Kosaki, R.; Craigen, W. J.; Matsuo, N.: Isoform-
specific

[162362]imprinting of the human PEG1/MEST gene. (Letter) Am. J.
Hum. Genet. 66:

[162363]309–312, 2000.

[162364]

[162365]

[162366]16075.

[162367]Kotzot, D.; Schmitt, S.; Bernasconi, F.; Robinson, W. P.;
Lurie,

[162368]I. W.; Ilyina, H.; Mehes, K.; Hamel, B. C. J.; Otten, B. J.;
Hergersberg,

[162369]M.; Werder, E.; Shoenle, E.; Schinzel, A.: Uniparental dis-
omy 7 in

[162370]Silver–Russell syndrome and primordial growth retarda-

tion. Hum. Molec.

[162371]Genet. 4: 583–587, 1995.

[162372]

[162373]

[162374]16076.

[162375]Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Keverne,

[162376]E. B.; Surani, M. A.: Abnormal maternal behaviour and
growth retardation

[162377]associated with loss of the imprinted gene Mest. Nature
Genet. 20:

[162378]163–169, 1998.

[162379]

[162380]

[162381]16077.

[162382]Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Surani, M.

[162383]A.: Genomic structure and parent-of-origin-specific
methylation of

[162384]Peg1. Hum. Molec. Genet. 6: 1907–1915, 1997.

[162385]

[162386]

[162387]16078.

[162388]Seki, T.; Tada, S.; Katada, T.; Enomoto, T.: Cloning of a
cDNA

[162389]encoding a novel importin- α homologue, Qip1: discrimination of

[162390]Qip1 and Rch1 from hSrp1 by their ability to interact with
DNA helicase

[162391]Q1/RecQL. Biochem. Biophys. Res. Commun. 234: 48–53,
1997.

[162392]

[162393]

[162394]16079.

[162395]Ayala-Madrigal, M. L.; Doerr, S.; Ramirez-Duenas, M. L.;
Hansmann,

[162396]I.: Assignment of karyopherin α 1 (KPNA1) to human
chromosome

[162397]band 3q21 by in situ hybridization. Cytogenet. Cell Genet.
90: 58–59,

[162398]2000.

[162399]

[162400]

[162401]16080.

[162402]Conti, E.; Uy, M.; Leighton, L.; Blobel, G.; Kuriyan, J.: Crys-
tallographic

[162403]analysis of the recognition of a nuclear localization signal
by the

[162404]nuclear import factor karyopherin alpha. Cell 94:
193-204, 1998.

[162405]

[162406]

[162407]16081.

[162408]Cortes, P.; Ye, Z.-S.; Baltimore, D.: RAG-1 interacts with
the

[162409]repeated amino acid motif of the human homologue of
the yeast protein

[162410]SRP1. Proc. Nat. Acad. Sci. 91: 7633-7637, 1994.

[162411]

[162412]

[162413]16082.

[162414]Chen, H.; Antonarakis, S. E.: Localization of a human ho-

molog

[162415]of the mouse Tiam-1 gene to chromosome 21q22.1. Genomics 30: 123-127,

[162416]1995.

[162417]

[162418]

[162419]16083.

[162420]Habets, G. G. M.; Scholtes, E. H. M.; Zuydgeest, D.; van der Kammen,

[162421]R. A.; Stam, J. C.; Berns, A.; Collard, J. G.: Identification of

[162422]an invasion-inducing gene, Tiam-1, that encodes a protein with homology

[162423]to GDP-GTP exchangers for Rho-like proteins. Cell 77: 537-549, 1994.

[162424]

[162425]

[162426]16084.

[162427]Habets, G. G. M.; van der Kammen, R. A.; Jenkins, N. A.; Gilbert,

[162428]D. J.; Copeland, N. G.; Hagemeijer, A.; Collard, J. G.: The

invasion-inducing

[162429]TIAM1 gene maps to human chromosome band 21q22
and mouse chromosome

[162430]16. Cytogenet. Cell. Genet. 70: 48–51, 1995.

[162431]

[162432]

[162433]16085.

[162434]Habets, G. G. M.; van der Kammen, R. A.; Stam, J. C.;
Michiels,

[162435]F.; Collard, J. G.: Sequence of the human invasion-induc-
ing TIAM1

[162436]gene, its conservation in evolution and its expression in
tumor cell

[162437]lines of different tissue origin. Oncogene 10: 1371–1376,
1995.

[162438]

[162439]

[162440]16086.

[162441]Malliri, A.; van der Kammen, R. A.; Clark, K.; van der Valk,
M.;

[162442]Michiels, F.; Collard, J. G.: Mice deficient in the Rac activa-

tor

[162443]Tiam1 are resistant to Ras-induced skin tumours. Nature
417: 867–871,

[162444]2002.

[162445]

[162446]

[162447]16087.

[162448]Koch, A.; Juan, T. S.-C.; Jenkins, N. A.; Gilbert, D. J.;
Copeland,

[162449]N. G.; McNiece, I. K.; Fletcher, F. A.: cDNA cloning and
chromosomal

[162450]mapping of mouse fast skeletal muscle troponin T. Mam-
malian Genome 8:

[162451]346–348, 1997.

[162452]

[162453]

[162454]16088.

[162455]Mao, C.; Baumgartner, A. P.; Jha, P. K.; Huang, T. H.-M.;
Sarkar,

[162456]S.: Assignment of the human fast skeletal troponin T gene

(TNNT3)

[162457]to chromosome 11p15.5: evidence for the presence of
11pter in a monochromosome

[162458]9 somatic cell hybrid in NIGMS mapping panel 2. Ge-
nomics 31: 385-388,

[162459]1996.

[162460]

[162461]

[162462]16089.

[162463]Wu, Q.-L.; Jha, P. K.; Raychowdhury, M. K.; Du, Y.; Leavis,
P.

[162464]C.; Sarkar, S.: Isolation and characterization of human fast
skeletal

[162465]beta troponin T cDNA: comparative sequence analysis of
isoforms and

[162466]insight into the evolution of members of a multigene fam-
ily. DNA

[162467]Cell Biol. 13: 217-233, 1994.

[162468]

[162469]

[162470]16090.

[162471]Ghetti, A.; Pinol-Roma, S.; Michael, W. M.; Morandi, C.;
Dreyfuss,

[162472]G.: HNRNP I, the polypyrimidine tract-binding protein:
distinct nuclear

[162473]localization and association with hnRNAs. Nucleic Acids
Res. 20:

[162474]3671–3678, 1992.

[162475]

[162476]

[162477]16091.

[162478]Raimondi, E.; Romanelli, M. G.; Moralli, D.; Gamberi, C.;
Russo,

[162479]M. P.; Morandi, C.: Assignment of the human gene encod-
ing heterogeneous

[162480]nuclear RNA ribonucleoprotein I (PTB) to chromosome
14q23–q24.1. Genomics 27:

[162481]553–555, 1995.

[162482]

[162483]

[162484]16092.

[162485]Romanelli, M. G.; Lorenzi, P.; Morandi, C.: Organization of the

[162486]human gene encoding heterogeneous nuclear ribonucleo-protein type I

[162487](hnRNP I) and characterization of hnRNP I related pseudo-gene. Gene 255:

[162488]267–272, 2000.

[162489]

[162490]

[162491]16093.

[162492]Hoogenraad, C. C.; Koekkoek, B.; Akhmanova, A.; Krugers, H.; Dortland,

[162493]B.; Miedema, M.; van Alphen, A.; Kistler, W. M.; Jaegle, M.; Koutsourakis,

[162494]M.; Van Camp, N.; Verhoye, M.; van der Linden, A.; Kaverina, I.; Grosveld,

[162495]F.; De Zeeuw, C. I.; Galjart, N.: Targeted mutation of Cyln2 in the

[162496]Williams syndrome critical region links CLIP-115 haploinsufficiency

[162497]to neurodevelopmental abnormalities in mice. Nature Genet. 32: 116–127,

[162498]2002. Note: Erratum: Nature Genet. 32: 331 only, 2002.

[162499]

[162500]

[162501]16094.

[162502]Gollob, M. H.; Green, M. S.; Roberts, R.: A gene responsible for

[162503]familial Wolff–Parkinson–White syndrome. (Letter) New Eng. J. Med. 345:

[162504]1063–1064, 2001.

[162505]

[162506]

[162507]16095.

[162508]L'Allemand, D.; Tardy, V.; Gruters, A.; Schnabel, D.; Krude, H.;

[162509]Morel, Y.: How a patient homozygous for a 30–kb deletion of the C4–CYP

[162510]21 genomic region can have a nonclassic form of 21–hydroxylase deficiency. J.

[162511]Clin. Endocr. Metab. 85: 4562–4567, 2000.

[162512]

[162513]

[162514]16096.

[162515]Lako, M.; Ramsden, S.; Campbell, R. D.; Strachan, T.: Mutation
tation

[162516]screening in British 21-hydroxylase deficiency families
and development

[162517]of novel microsatellite based approaches to prenatal diagnosis. J.

[162518]Med. Genet. 36: 119–124, 1999.

[162519]

[162520]

[162521]16097.

[162522]Lau, I. F.; Soardi, F. C.; Lemos–Marini, S. H. V.; Guerra, G.,

[162523]Jr.; Baptista, M. T. M.; De Mello, M. P.: H28+C insertion in
the

[162524]CYP21 gene: a novel frameshift mutation in a Brazilian patient with

[162525]the classical form of 21-hydroxylase deficiency. J. Clin.
Endocr.

[162526]Metab. 86: 5877–5880, 2001.

[162527]

[162528]

[162529]16098.

[162530]Laue, L.; Merke, D. P.; Jones, J. V.; Barnes, K. M.; Hill, S.;

[162531]Cutler, G. B.: A preliminary study of flutamide, testolactone, and

[162532]reduced hydrocortisone dose in the treatment of congenital adrenal

[162533]hyperplasia. J. Clin. Endocr. Metab. 81: 3535–3539, 1996.

[162534]

[162535]

[162536]16099.

[162537]Layrisse, A.; White, C.; Gunczler, P.; Gafaro Valera, L.;
Arias,

[162538]S.; Yunis, E. J.; Alper, C. A.; Awdeh, Z. L.: Sharing of MHC
haplotypes

[162539]among apparently unrelated patients with congenital
adrenal hyperplasia

[162540]due to 21-hydroxylase deficiency. Immunogenetics 25:
99–103, 1987.

[162541]

[162542]

[162543]16100.

[162544]Lee, H.-H.; Chao, H.-T.; Ng, H.-T.; Choo, K.-B.: Direct
molecular

[162545]diagnosis of CYP21 mutations in congenital adrenal hy-
perplasia. J.

[162546]Med. Genet. 33: 371-375, 1996.

[162547]

[162548]

[162549]16101.

[162550]Levine, L. S.; Dupont, B.; Lorenzen, F.; Pang, S.; Pollack,
M.;

[162551]Oberfield, S.; Kohn, B.; Lerner, A.; Cacciari, E.; Mantero, F.;
Cassio,

[162552]A.; Scaroni, C.; Chiumello, G.; Rondanini, G. F.; Gargantini,
L.;

[162553]Giovannelli, G.; Viridis, R.; Bartolotta, E.; Migliori, C.; Pin-
tor,

[162554]C.; Tato, L.; Barboni, F.; New, M. I.: Cryptic
21-hydroxylase deficiency

[162555]in families of patients with classical congenital adrenal hyperplasia. J.

[162556]Clin. Endocr. Metab. 51: 1316–1324, 1980.

[162557]

[162558]

[162559]16102.

[162560]Levine, L. S.; Dupont, B.; Lorenzen, F.; Pang, S.; Pollack, M.;

[162561]Oberfield, S. E.; Kohn, B.; Lerner, A.; Cacciari, E.; Mantero, F.;

[162562]Cassio, A.; Scaroni, C.; Chiumello, G.; Rondanini, G. F.; Gargantini,

[162563]L.; Giovannelli, G.; Viridis, R.; Bartolotta, E.; Migliori, C.; Pintor,

[162564]C.; Tato, L.; Barboni, F.; New, M. I.: Genetic and hormonal characterization

[162565]of cryptic 21–hydroxylase deficiency. J. Clin. Endocr. Metab. 53:

[162566]1193–1198, 1981.

[162567]

[162568]

[162569]16103.

[162570]Levine, L. S.; Pang, S.; Dupont, B.; Pollack, M.; Lorenzen, F.;

[162571]New, M. I.: Detection of heterozygote of 21-hydroxylase deficiency.

[162572](Letter) Lancet I: 603-604, 1980.

[162573]

[162574]

[162575]16104.

[162576]Levine, L. S.; Zachmann, M.; New, M. I.; Prader, A.; Pollack,

[162577]M. S.; O'Neill, G. J.; Yang, S. Y.; Oberfield, S. E.; Dupont, B.:

[162578]Genetic mapping of the 21-hydroxylase-deficiency gene within the HLA

[162579]linkage group. New Eng. J. Med. 299: 911-915, 1978.

[162580]

[162581]

[162582]16105.

[162583]Levo, A.; Partanen, J.: Mutation-haplotype analysis of steroid

[162584]21-hydroxylase (CYP21) deficiency in Finland. Implications for the

[162585]population history of defective alleles. Hum. Genet. 99: 488-497,

[162586]1997.

[162587]

[162588]

[162589]16106.

[162590]Lewis, V. G.; Money, J.; Epstein, R.: Concordance of verbal and

[162591]nonverbal ability in the adrenogenital syndrome. Johns Hopkins Med.

[162592]J. 122: 192-195, 1968.

[162593]

[162594]

[162595]16107.

[162596]Libber, S. M.; Migeon, C. J.; Bias, W. B.: Ascertainment of 21-hydroxylase

[162597]deficiency in individuals with HLA-B14 haplotype. J. Clin. Endocr.

[162598]Metab. 60: 727–730, 1985.

[162599]

[162600]

[162601]16108.

[162602]Lo, J. C.; Schwitzgebel, V. M.; Tyrrell, J. B.; Fitzgerald, P.

[162603]A.; Kaplan, S. L.; Conte, F. A.; Grumbach, M. M.: Normal
female infants

[162604]born of mothers with classic congenital adrenal hyperpla-
sia due to

[162605]21-hydroxylase deficiency. J. Clin. Endocr. Metab. 84:
930–936,

[162606]1999.

[162607]

[162608]

[162609]16109.

[162610]Lopez-Gutierrez, A. U.; Riba, L.; Ordonez-Sanchez, M. L.;
Ramirez-Jimenez,

[162611]S.; Cerrillo-Hinojosa, M.; Tusie-Luna, M. T.: Uniparental
disomy

[162612]for chromosome 6 results in steroid 21-hydroxylase defi-

ciency: evidence

[162613]of different genetic mechanisms involved in the production of the

[162614]disease. J. Med. Genet. 35: 1014–1019, 1998.

[162615]

[162616]

[162617]16110.

[162618]Lorenzen, F.; Pang, S.; New, M. I.; Dupont, B.; Pollack, M.;
Chow,

[162619]D. M.; Levine, L. S.: Hormonal phenotype and HLA-genotype in families

[162620]of patients with congenital adrenal hyperplasia
(21-hydroxylase deficiency). Pediat.

[162621]Res. 13: 1356–1360, 1979.

[162622]

[162623]

[162624]16111.

[162625]Lorenzen, F.; Pang, S.; New, M. I.; Oberfield, S.; Dupont,
B.;

[162626]Chow, D.; Schneider, B.; Levine, L.: Studies of the C-21

and C-19

[162627]steroids and HLA genotyping in siblings and parents of patients with

[162628]congenital adrenal hyperplasia due to 21-hydroxylase deficiency. J.

[162629]Clin. Endocr. Metab. 50: 572-577, 1980.

[162630]

[162631]

[162632]16112.

[162633]Lubani, M. M.; Issa, A.-R. A.; Bushnaq, R.; Al-Saleh, Q. A.; Dudin,

[162634]K. I.; Reavey, P. C.; El-Khalifa, M. Y.; Manandhar, D. S.; Abdul Al,

[162635]Y. K.; Ismail, E. A.; Teebi, A. S.: Prevalence of congenital adrenal

[162636]hyperplasia in Kuwait. Europ. J. Pediat. 149: 391-392, 1990.

[162637]

[162638]

[162639]16113.

[162640]Matteson, K. J.; Phillips, J. A., III; Miller, W. L.; Chung,

B.-C.;

[162641]Orlando, P. J.; Frisch, H.; Ferrandez, A.; Burr, I. M.:

P450XXI (steroid

[162642]21-hydroxylase) gene deletions are not found in family studies of

[162643]congenital adrenal hyperplasia. Proc. Nat. Acad. Sci. 84: 5858-5862,

[162644]1987.

[162645]

[162646]

[162647]16114.

[162648]McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba,

[162649]Canada 2/26/1988.

[162650]

[162651]

[162652]16115.

[162653]McGuire, L. S.; Omenn, G. S.: Congenital adrenal hyperplasia.

[162654]I. Family studies of IQ. Behav. Genet. 5: 165-173, 1975.

[162655]

[162656]

[162657]16116.

[162658]Merkatz, I. R.; New, M. I.; Peterson, R. E.; Seaman, M. P.:

Prenatal

[162659]diagnosis of adrenogenital syndrome by amniocentesis. J.

Pediat. 75:

[162660]977–982, 1969.

[162661]

[162662]

[162663]16117.

[162664]Merke, D. P.; Chrousos, G. P.; Eisenhofer, G.; Weise, M.;

Keil,

[162665]M. F.; Rogol, A. D.; Van Wyk, J. J.; Bornstein, S. R.:

Adrenomedullary

[162666]dysplasia and hypofunction in patients with classic

21-hydroxylase

[162667]deficiency. New Eng. J. Med. 343: 1362–1368, 2000.

[162668]

[162669]

[162670]16118.

[162671]Merke, D. P.; Keil, M. F.; Jones, J. V.; Fields, J.; Hill, S.;

[162672]Cutler, G. B., Jr.: Flutamide, testolactone, and reduced hydrocortisone

[162673]dose maintain normal growth velocity and bone maturation despite elevated

[162674]androgen levels in children with congenital adrenal hyperplasia. J.

[162675]Clin. Endocr. Metab. 85: 1114–1120, 2000.

[162676]

[162677]

[162678]16119.

[162679]Meyer–Bahlburg, H. F. L.: What causes low rates of childbearing

[162680]in congenital adrenal hyperplasia? J. Clin. Endocr. Metab. 84: 1844–1847,

[162681]1999.

[162682]

[162683]

[162684]16120.

[162685]Miller, W. L.: Gene conversions, deletions and polymorphisms

[162686]in congenital adrenal hyperplasia. Am. J. Hum. Genet. 42: 4–7, 1988.

[162687]

[162688]

[162689]16121.

[162690]Miller, W. L.: Personal Communication. San Francisco, Calif.

[162691]1/3/1996.

[162692]

[162693]

[162694]16122.

[162695]Bannerman, R. M.; Edwards, J. A.; Kreimer–Birnbaum, M.; McFarland,

[162696]E.; Russell, E. S.: Hereditary microcytic anaemia in the mouse; studies

[162697]in iron distribution and metabolism. Brit. J. Haemat. 23: 235–245,

[162698]1972.

[162699]

[162700]

[162701]16123.

[162702]Vincent, J. B.; Herbrick, J.-A.; Gurling, H. M. D.; Bolton, P.

[162703]F.; Roberts, W.; Scherer, S. W.: Identification of a novel
gene on

[162704]chromosome 7q31 that is interrupted by a translocation
breakpoint

[162705]in an autistic individual. Am. J. Hum. Genet. 67: 510–514,
2000.

[162706]

[162707]

[162708]16124.

[162709]Barichard, F.; Joulin, V.; Henry, I.; Garel, M.-C.; Valentin,
C.;

[162710]Rosa, R.; Cohen-Solal, M.; Junien, C.: Chromosomal as-
signment of

[162711]the human 2,3-bisphosphoglycerate mutase gene (BPGM)
to region 7q34–7q22. Hum.

[162712]Genet. 77: 283–285, 1987.

[162713]

[162714]

[162715]16125.

[162716]Bowdler, A. J.; Prankerd, T. A. J.: Studies in congenital
non-spherocytic

[162717]haemolytic anaemias with specific enzyme defects. Acta
Haemat. 31:

[162718]65–78, 1964.

[162719]

[162720]

[162721]16126.

[162722]Chen, S.–H.; Anderson, J. E.; Giblett, E. R.:
2,3-diphosphoglycerate

[162723]mutase: its demonstration by electrophoresis and the de-
tection of

[162724]a genetic variant. Biochem. Genet. 5: 481–486, 1971.

[162725]

[162726]

[162727]16127.

[162728]Galacteros, F.; Rosa, R.; Prehu, M. O.; Najean, Y.; Calvin, M.

[162729]C.: Deficit en diphosphoglycerate mutase: nouveaux cas associes a

[162730]une polyglobulie. Nouv. Rev. Franc. Hemat. 26: 69–74, 1984.

[162731]

[162732]

[162733]16128.

[162734]Joulin, V.; Barichard, F.; Henry, I.; Garel, M. C.; Valentin, C.;

[162735]Rosa, R.; Cohen–Solal, M.; Junien, C.: Chromosomal assignment of

[162736]the human 2,3–bisphosphoglycerate mutase gene (BPGM) to region 7q22–7q34.

[162737](Abstract) Cytogenet. Cell Genet. 46: 635 only, 1987.

[162738]

[162739]

[162740]16129.

[162741]Joulin, V.; Garel, M.–C.; Le Boulch, P.; Valentin, C.; Rosa, R.;

[162742]Rosa, J.; Cohen-Solal, M.: Isolation and characterization of
the

[162743]human 2,3-bisphosphoglycerate mutase gene. J. Biol.
Chem. 263: 15785-15790,

[162744]1988.

[162745]

[162746]

[162747]16130.

[162748]Joulin, V.; Peduzzi, J.; Romeo, P.-H.; Rosa, R.; Valentin, C.;

[162749]Dubart, A.; Lapeyre, B.; Blouquit, Y.; Garel, M.-C.;
Goossens, M.;

[162750]Rosa, J.; Cohen-Solal, M.: Molecular cloning and sequenc-
ing of the

[162751]human erythrocyte 2,3-bisphosphoglycerate mutase
cDNA: revised amino

[162752]acid sequence. EMBO J. 5: 2275-2283, 1986.

[162753]

[162754]

[162755]16131.

[162756]Labie, D.; Leroux, J.-P.; Najman, A.; Reyrolle, C.: Familial

diphosphoglycerate

[162757]mutase deficiency: influence on the oxygen affinity curves
of hemoglobin. FEBS

[162758]Lett. 9: 37–40, 1970.

[162759]

[162760]

[162761]16132.

[162762]Lemarchandel, V.; Joulin, V.; Valentin, C.; Rosa, R.; Galac-
teros,

[162763]F.; Rosa, J.; Cohen–Solal, M.: Compound heterozygosity in
a complete

[162764]erythrocyte bisphosphoglycerate mutase deficiency. Blood
80: 2643–2649,

[162765]1992.

[162766]

[162767]

[162768]16133.

[162769]Rosa, R.; Audit, I.; Rosa, J.: Diphosphoglycerate mutase
and

[162770]2,3–diphosphoglycerate phosphatase activities of red

cells: comparative

[162771]electrophoretic study. Biochem. Biophys. Res. Commun.
51: 536–542,

[162772]1973.

[162773]

[162774]

[162775]16134.

[162776]Rosa, R.; Blouquit, Y.; Calvin, M.–C.; Prome, D.; Prome,
J.–C.;

[162777]Rosa, J.: Isolation, characterization, and structure of a
mutant

[162778]89 arg–to–cys bisphosphoglycerate mutase: implication of
the active

[162779]site in the mutation. J. Biol. Chem. 264: 7837–7843,
1989.

[162780]

[162781]

[162782]16135.

[162783]Rosa, R.; Prehu, M.–O.; Beuzard, Y.; Rosa, J.: The first case

[162784]of a complete deficiency of diphosphoglycerate mutase in

human erythrocytes. J.

[162785]Clin. Invest. 62: 907–915, 1978.

[162786]

[162787]

[162788]16136.

[162789]Sasaki, R.; Ikura, K.; Sugimoto, E.; Chiba, H.: Purification

[162790]of bisphosphoglyceromutase, 2,3–bisphosphoglycerate
phosphatase and

[162791]phosphoglyceromutase from human erythrocytes. Europ.
J. Biochem. 50:

[162792]581–593, 1975.

[162793]

[162794]

[162795]16137.

[162796]Schroter, W.: Kongenitale nichtsphaerocytaere haemolytis-
che Anaemie

[162797]bei 2,3–Diphosphoglyceratmutasemangel der Erythrocyten
im fruehen

[162798]Saeuglingsalter. Klin. Wschr. 43: 1147–1153, 1965.

[162799]

[162800]

[162801]16138.

[162802]Scott, E. M.; Wright, R. C.: An alternate method for
demonstration

[162803]of bisphosphoglyceromutase (DPGM) on starch gels. Am.
J. Hum. Genet. 34:

[162804]1013–1015, 1982.

[162805]

[162806]

[162807]16139.

[162808]Yanagawa, S.; Hitomi, K.; Sasaki, R.; Chiba, H.: Isolation
and

[162809]characterization of cDNA encoding rabbit reticulocyte
2,3–bisphosphoglycerate

[162810]synthase. Gene 44: 185–191, 1986.

[162811]

[162812]

[162813]16140.

[162814]Cali, J. J.; Hsieh, C.-L.; Francke, U.; Russell, D. W.: Muta-
tions

[162815]in the bile acid biosynthetic enzyme sterol 27-hydroxylase
underlie

[162816]cerebrotendinous xanthomatosis. J. Biol. Chem. 266:
7779-7783, 1991.

[162817]

[162818]

[162819]16141.

[162820]Cali, J. J.; Russell, D. W.: Characterization of human sterol

[162821]27-hydroxylase: a mitochondrial cytochrome P-450 that
catalyzes multiple

[162822]oxidation reactions in bile acid biosynthesis. J. Biol. Chem.
266:

[162823]7774-7778, 1991.

[162824]

[162825]

[162826]16142.

[162827]Jaspers, M.; Zhang, Z.; Marynen, P.; Vekemans, S.; Aly, M.
S.;

[162828]Cuppens, H.; Hillicker, C.; Cassiman, J.-J.: Localization of
the

[162829]genes encoding the alpha-2 and alpha-4 subunits of the
human VLA-receptors

[162830]to chromosome 5q23-31 and 2q31-32 respectively.

(Abstract) Cytogenet.

[162831]Cell Genet. 58: 1870 only, 1991.

[162832]

[162833]

[162834]16143.

[162835]Kaplan, C.; Morel-Kopp, M. C.; Kroll, H.; Kiefel, V.;
Schlegel,

[162836]N.; Chesnel, N.; Mueller-Eckhardt, C.: HPA-5b (Br-a)
neonatal alloimmune

[162837]thrombocytopenia: clinical and immunological analysis of
39 cases. Brit.

[162838]J. Haemat. 78: 425-429, 1991.

[162839]

[162840]

[162841]16144.

[162842]Kiefel, V.; Santoso, S.; Katzmann, B.; Mueller-Eckhardt, C.:

A

[162843]new platelet-specific alloantigen Br(a): report of 4 cases
with neonatal

[162844]alloimmune thrombocytopenia. Vox Sang. 54: 101-106,
1988.

[162845]

[162846]

[162847]16145.

[162848]Kiefel, V.; Shechter, Y.; Atias, D.; Kroll, H.; Santoso, S.;
Mueller-Eckhardt,

[162849]C.: Neonatal alloimmune thrombocytopenia due to anti-
Br(b) (HPA-5a):

[162850]report of three cases in two families. Vox Sang. 60:
244-245, 1991.

[162851]

[162852]

[162853]16146.

[162854]Kritzik, M.; Savage, B.; Nugent, D. J.; Santoso, S.; Ruggeri,
Z.

[162855]M.; Kunicki, T. J.: Nucleotide polymorphisms in the alpha-

2 gene

[162856]define multiple alleles that are associated with differences
in platelet

[162857]alpha-2/beta-1 density. Blood 92: 2382-2388, 1998.

[162858]

[162859]

[162860]16147.

[162861]Kunicki, T. J.; Kritzik, M.; Annis, D. S.; Nugent, D. J.:
Hereditary

[162862]variation in platelet integrin alpha-2-beta-1 density is as-
sociated

[162863]with two silent polymorphisms in the alpha-2 gene coding
sequence. Blood 89:

[162864]1939-1943, 1997.

[162865]

[162866]

[162867]16148.

[162868]Mueller-Eckhardt, C.; Kiefel, V.; Grubert, A.; Kroll, H.;
Weisheit,

[162869]M.; Schmidt, S.; Mueller-Eckhardt, G.; Santoso, S.: 348

cases of

[162870]suspected neonatal alloimmune thrombocytopenia. Lancet
I: 363–366,
[162871]1989.

[162872]

[162873]

[162874]16149.

[162875]Nieuwenhuis, H. K.; Akkerman, J. W. N.; Houdijk, W. P.;
Sixma,

[162876]J. J.: Human blood platelets showing no response to colla-
gen fail

[162877]to express surface glycoprotein Ia. Nature 318: 470–472,
1985.

[162878]

[162879]

[162880]16150.

[162881]Santoso, S.; Amrhein, J.; Hofmann, H. A.; Sachs, U. J. H.;
Walka,

[162882]M. M.; Kroll, H.; Kiefel, V.: A point mutation thr799met on
the alpha-2

[162883]integrin leads to the formation of new human platelet al-

loantigen

[162884]Sit(a) and affects collagen-induced aggregation. Blood 94:
4103-4111,

[162885]1999.

[162886]

[162887]

[162888]16151.

[162889]Santoso, S.; Kalb, R.; Walka, M.; Kiefel, V.; Mueller-Eck-
hardt,

[162890]C.; Newman, P. J.: The human platelet alloantigens Br(a)
and Br(b)

[162891]are associated with a single amino acid polymorphism on
glycoprotein

[162892]Ia (integrin subunit alpha-2). J. Clin. Invest. 92:
2427-2432, 1993.

[162893]

[162894]

[162895]16152.

[162896]Santoso, S.; Kunicki, T. J.; Kroll, H.; Haberbosch, W.;
Gardemann,

[162897]A.: Association of the platelet glycoprotein Ia C807T gene

polymorphism

[162898]with nonfatal myocardial infarction in younger patients.

Blood 93:

[162899]2449–2453, 1999.

[162900]

[162901]

[162902]16153.

[162903]Takada, Y.; Hemler, M. E.: The primary structure of the
VLA-2/collagen

[162904]receptor alpha-2 subunit (platelet GPIa): homology to
other integrins

[162905]and the presence of a possible collagen-binding domain.

J. Cell Biol. 109:

[162906]397–407, 1989.

[162907]

[162908]

[162909]16154.

[162910]von Beckerath, N.; Koch, W.; Mehilli, J.; Bottiger, C.;

Schomig,

[162911]A.; Kastrati, A.: Glycoprotein Ia gene C807T polymorphism

and risk

[162912]for major adverse cardiac events within the first 30 days
after coronary

[162913]artery stenting. Blood 95: 3297–3301, 2000.

[162914]

[162915]

[162916]16155.

[162917]Fernandez–Ruiz, E.; Pardo–Manuel de Villena, F.; Rubio, M.
A.;

[162918]Corbi, A. L.; Rodriguez de Cordoba, S.; Sanchez–Madrid,
F.: Mapping

[162919]of the human VLA–alpha–4 gene to chromosome
2q31–q32. Europ. J. Immun. 22:

[162920]587–590, 1992.

[162921]

[162922]

[162923]16156.

[162924]Roseblatt, M.; Vuillet–Gaugler, M. H.; Leroy, C.;
Coulombel, L.

[162925]: Coexpression of two fibronectin receptors, VLA–4 and

VLA-5, by immature

[162926]human erythroblastic precursor cells. J. Clin. Invest. 87:
6-11,

[162927]1991.

[162928]

[162929]

[162930]16157.

[162931]Zhang, Z.; Vekemans, S.; Aly, M. S.; Jaspers, M.; Marynen,
P.;

[162932]Cassiman, J.-J.: The gene for the alpha-4 subunit of the
VLA-4 integrin

[162933]maps to chromosome 2q31-32. Blood 78: 2396-2399,
1991.

[162934]

[162935]

[162936]16158.

[162937]Hiesberger, T.; Trommsdorff, M.; Howell, B. W.; Goffinet,
A.; Mumby,

[162938]M. C.; Cooper, J. A.; Herz, J.: Direct binding of reelin to
VLDL

[162939]receptor and apoE receptor 2 induces tyrosine phosphory-

lation of disabled-1

[162940]and modulates tau phosphorylation. Neuron 24: 481–489,
1999.

[162941]

[162942]

[162943]16159.

[162944]Oka, K.; Tzung, K.-W.; Sullivan, M.; Lindsay, E.; Baldini, A.;

[162945]Chan, L.: Human very-low-density lipoprotein receptor
complementary

[162946]DNA and deduced amino acid sequence and localization
of its gene (VLDLR)

[162947]to chromosome band 9q24 by fluorescence in situ hy-
bridization. Genomics 20:

[162948]298–300, 1994.

[162949]

[162950]

[162951]16160.

[162952]Okuizumi, K.; Onodera, O.; Namba, Y.; Ikeda, K.; Ya-
mamoto, T.;

[162953]Seki, K.; Ueki, A.; Nanko, S.; Tanaka, H.; Takahashi, H.;

Oyanagi,

[162954]K.; Mizusawa, H.; Kanazawa, I.; Tsuji, S.: Genetic association of

[162955]the very low density lipoprotein (VLDL) receptor gene with sporadic

[162956]Alzheimer's disease. Nature Genet. 11: 207–209, 1995.

[162957]

[162958]

[162959]16161.

[162960]Gafvels, M. E.; Paavola, L. G.; Boyd, C. O.; Nolan, P. M.; Wittmaack,

[162961]F.; Chawla, A.; Lazar, M. A.; Bucan, M.; Angelin, B.; Strauss, J.

[162962]F., III: Cloning of a complementary deoxyribonucleic acid encoding

[162963]the murine homolog of the very low density lipoprotein/apolipoprotein-E

[162964]receptor: expression pattern and assignment of the gene to mouse chromosome

[162965]19. Endocrinology 135: 387–394, 1994.

[162966]

[162967]

[162968]16162.

[162969]Sakai, J.; Hoshino, A.; Takahashi, S.; Miura, Y.; Ishii, H.;
Suzuki,

[162970]H.; Kawarabayasi, Y.; Yamamoto, T.: Structure, chromo-
some location,

[162971]and expression of the human very low density lipoprotein
receptor

[162972]gene. J. Biol. Chem. 269: 2173–2182, 1994.

[162973]

[162974]

[162975]16163.

[162976]Trommsdorff, M.; Gotthardt, M.; Hiesberger, T.; Shelton,
J.; Stockinger,

[162977]W.; Nimpf, J.; Hammer, R. E.; Richardson, J. A.; Herz, J.:
Reeler/Disabled-like

[162978]disruption of neuronal migration in knockout mice lacking
the VLDL

[162979]receptor and ApoE receptor 2. Cell 97: 689–701, 1999.

[162980]

[162981]

[162982]16164.

[162983]Huebner, K.; Druck, T.; Croce, C. M.; Thiesen, H. J.:

Twenty-seven

[162984]nonoverlapping zinc finger cDNAs from human T cells

map to nine different

[162985]chromosomes with apparent clustering. Am. J. Hum.

Genet. 48: 726-740,

[162986]1991.

[162987]

[162988]

[162989]16165.

[162990]Rousseau-Merck, M.-F.; Hillion, J.; Jonveaux, P.; Couillin,

P.;

[162991]Seite, P.; Thiesen, H.-J.; Berger, R.: Chromosomal localiza-

tion of

[162992]9 KOX zinc finger genes: physical linkages suggest clus-

tering of KOX

[162993]genes on chromosomes 12, 16, and 19. Hum. Genet. 92:

583-587, 1993.

[162994]

[162995]

[162996]16166.

[162997]Bray, P.; Lichter, P.; Thiesen, H.-J.; Ward, D. C.; Dawid, I. B.

[162998]: Characterization and mapping of human genes encoding
zinc finger

[162999]proteins. Proc. Nat. Acad. Sci. 88: 9563–9567, 1991.

[163000]

[163001]

[163002]16167.

[163003]Lehmann, O. J.; El-ashry, M. F.; Ebenezer, N. D.; Ocaka, L.;
Francis,

[163004]P. J.; Wilkie, S. E.; Patel, R. J.; Ficker, L.; Jordan, T.; Khaw,

[163005]P. T.; Bhattacharya, S. S.: A novel keratocan mutation
causing autosomal

[163006]recessive cornea plana. Invest. Ophthalm. Vis. Sci. 42:
3118–3122,

[163007]2001.

[163008]

[163009]

[163010]16168.

[163011]Gahlmann, R.; Kedes, L.: Cloning, structural analysis, and
expression

[163012]of the human fast twitch skeletal muscle troponin C gene.
J. Biol.

[163013]Chem. 265: 12520–12528, 1990.

[163014]

[163015]

[163016]16169.

[163017]Gahlmann, R.; Wade, R.; Gunning, P.; Kedes, L.: Differen-
tial expression

[163018]of slow and fast skeletal muscle troponin C: slow skeletal
muscle

[163019]troponin C is expressed in human fibroblasts. J. Molec.
Biol. 201:

[163020]379–391, 1988.

[163021]

[163022]

[163023]16170.

[163024]Townsend, P. J.; Yacoub, M. H.; Barton, P. J. R.: Assign-

ment of

[163025]the human fast skeletal muscle troponin C gene (TNNC2)
between D20S721

[163026]and GCT10F11 on chromosome 20 by somatic cell hybrid
analysis. Ann.

[163027]Hum. Genet. 61: 457–459, 1997.

[163028]

[163029]

[163030]16171.

[163031]Bermingham, N.; Hernandez, D.; Balfour, A.; Gilmour, F.;
Martin,

[163032]J. E.; Fisher, E. M. C.: Mapping TNNC1, the gene that en-
codes cardiac

[163033]troponin I in the human and mouse. Genomics 30:
620–622, 1995.

[163034]

[163035]

[163036]16172.

[163037]Roher, A.; Lieska, N.; Spitz, W.: The amino acid sequence
of human

[163038]cardiac troponin–C. Muscle Nerve 9: 73–77, 1986.

[163039]

[163040]

[163041]16173.

[163042]Romero-Herrera, A. E.; Castillo, O.; Lehmann, H.: Human
skeletal

[163043]muscle proteins: the primary structure of troponin C. J.
Molec. Evol. 8:

[163044]251-270, 1976.

[163045]

[163046]

[163047]16174.

[163048]Schreier, T.; Kedes, L.; Gahlmann, R.: Cloning, structural
analysis,

[163049]and expression of the human slow twitch skeletal muscle/
cardiac troponin

[163050]C gene. J. Biol. Chem. 265: 21247-21253, 1990.

[163051]

[163052]

[163053]16175.

[163054]Song, W.-J.; Van Keuren, M. L.; Drabkin, H. A.; Cypser, J. R.;

[163055]Gemmell, R. M.; Kurnit, D. M.: Assignment of the human slow twitch

[163056]skeletal muscle/cardiac troponin C gene (TNNC1) to human chromosome

[163057]3p21.3-3p14.3 using somatic cell hybrids. Cytogenet. Cell Genet. 75:

[163058]36-37, 1996.

[163059]

[163060]

[163061]16176.

[163062]Song, W. J.: Personal Communication. Ann Arbor, Mich.
1/17/1997.

[163063]

[163064]

[163065]16177.

[163066]Townsend, P. J.; Yacoub, M. H.; Barton, P. J. R.: Assign-
ment of

[163067]the human cardiac/slow skeletal muscle troponin C gene
(TNNC1) between

[163068]D3S3118 and GCT4B10 on the short arm of chromosome
3 by somatic cell

[163069]hybrid analysis. Ann. Hum. Genet. 61: 375–377, 1997.

[163070]

[163071]

[163072]16178.

[163073]Haldane, J. B. S.; Sprunt, A. D.; Haldane, N. M.: Reduplica-

tion

[163074]in mice. J. Genet. 5: 133–135, 1915.

[163075]

[163076]

[163077]16179.

[163078]Lyon, M. F.; King, T. R.; Gondo, Y.; Gardner, J. M.; Nakatsu,

[163079]Y.; Eicher, E. M.; Brilliant, M. H.: Genetic and molecular
analysis

[163080]of recessive alleles at the pink-eyed dilution (p) locus of
the mouse. Proc.

[163081]Nat. Acad. Sci. 89: 6968–6972, 1992.

[163082]

[163083]

[163084]16180.

[163085]Stohl, W.; Kunkel, H. G.: Heterogeneity in expression of
the

[163086]T4 epitope in black individuals. Scand. J. Immun. 20:
273–278, 1984.

[163087]

[163088]

[163089]16181.

[163090]Tishkoff, S. A.; Dietzsch, E.; Speed, W.; Pakstis, A. J.; Kidd,

[163091]J. R.; Cheung, K.; Bonne-Tamir, B.; Santachiara-Benere-
cetti, A. S.;

[163092]Moral, P.; Krings, M.; Paabo, S.; Watson, E.; Risch, N.;
Jenkins,

[163093]T.; Kidd, K. K.: Global patterns of linkage disequilibrium at
the

[163094]CD4 locus and modern human origins. Science 271:
1380-1387, 1996.

[163095]

[163096]

[163097]16182.

[163098]van Dongen, J. J. M.; Wolvers-Tettero, I. L. M.; Versnel, M.
A.;

[163099]Westerveld, A.; Geurts van Kessel, A. H. M.: Assignment of
the genes

[163100]coding for the T-cell antigens CD7 (Tp41), CD5 (T1) and
CD4 (T4) to

[163101]human chromosome 17, 11, 12 respectively. (Abstract)

Cytogenet. Cell

[163102]Genet. 40: 767, 1985.

[163103]

[163104]

[163105]16183.

[163106]Zou, Y.-R.; Sunshine, M.-J.; Taniuchi, I.; Hatam, F.; Killeen,

[163107]N.; Littman, D. R.: Epigenetic silencing of CD4 in T cells
committed

[163108]to the cytotoxic lineage. Nature Genet. 332-336, 2001.

[163109]

[163110]

[163111]16184.

[163112]DiLella, A. G.: Chromosomal assignment of the human im-
munophilin

[163113]FKBP-12 gene. Biochem. Biophys. Res. Commun. 179:
1427-1433, 1991.

[163114]

[163115]

[163116]16185.

[163117]DiLella, A. G.; Hawkins, A.; Craig, R. J.; Schreiber, S. L.;
Griffin,

[163118]C. A.: Chromosomal band assignments of the genes en-
coding human FKBP12

[163119]and FKBP13. Biochem. Biophys. Res. Commun. 189:
819–823, 1992.

[163120]

[163121]

[163122]16186.

[163123]Goebel, M. G.: The peptidyl–prolyl isomerase,
FK506–binding protein,

[163124]is most likely the 12 kd endogenous inhibitor 2 of protein
kinase

[163125]C. (Letter) Cell 64: 1051–1052, 1991.

[163126]

[163127]

[163128]16187.

[163129]Jin, Y.–J.; Albers, M. W.; Lane, W. S.; Bierer, B. E.; Schreiber,

[163130]S. L.; Burakoff, S. J.: Molecular cloning of a membrane–as–
sociated

[163131]human FK506- and rapamycin-binding protein, FKBP-13.

Proc. Nat. Acad.

[163132]Sci. 88: 6677-6681, 1991.

[163133]

[163134]

[163135]16188.

[163136]Maki, N.; Sekiguchi, F.; Nishimaki, J.; Miwa, K.; Hayano, T.;
Takahashi,

[163137]N.; Suzuki, M.: Complementary DNA encoding the human
T-cell FK506-binding

[163138]protein, a peptidylprolyl cis-trans isomerase distinct from
cyclophilin. Proc.

[163139]Nat. Acad. Sci. 87: 5440-5443, 1990.

[163140]

[163141]

[163142]16189.

[163143]Peattie, D. A.; Hsaio, K.; Benasutti, M.; Lippke, J. A.: Three

[163144]distinct messenger RNAs can encode the human immuno-
suppressant-binding

[163145]protein FKBP12. Gene 150: 251-257, 1994.

[163146]

[163147]

[163148]16190.

[163149]Shou, W.; Aghdasi, B.; Armstrong, D. L.; Guo, Q.; Bao, S.;
Charng,

[163150]M.-J.; Mathews, L. M.; Schneider, M. D.; Hamilton, S. L.;
Matzuk,

[163151]M. M.: Cardiac defects and altered ryanodine receptor
function in

[163152]mice lacking FKBP12. Nature 391: 489–492, 1998.

[163153]

[163154]

[163155]16191.

[163156]Standaert, R. F.; Galat, A.; Verdine, G. L.; Schreiber, S. L.:

[163157]Molecular cloning and overexpression of the human
FK506–binding protein

[163158]FKBP. Nature 346: 671–674, 1990.

[163159]

[163160]

[163161]16192.

[163162]Wang, T.; Donahoe, P. K.; Zervos, A. S.: Specific interaction
[163163]of type I receptors of the TGF-beta family with the im-
munophilin FKBP-12. Science 265:
[163164]674-676, 1994.

[163165]

[163166]

[163167]16193.

[163168]Jin, Y. J.; Albers, M. W.; Lane, W. S.; Bierer, B. E.; Schreiber,
[163169]S. L.; Burakoff, S. J.: Molecular cloning of a membrane-as-
sociated
[163170]human FK506- and rapamycin-binding protein, FKBP-13.
Proc. Nat. Acad.
[163171]Sci. 88: 6677-6681, 1991.

[163172]

[163173]

[163174]16194.

[163175]Howard, H. C.; Mount, D. B.; Rochefort, D.; Byun, N.;
Dupre, N.;

[163176]Lu, J.; Fan, X.; Song, L.; Riviere, J.-B.; Prevost, C.; Horst, J.;
[163177]Simonati, A.; and 12 others: The K-Cl cotransporter KCC3
is mutant
[163178]in a severe peripheral neuropathy associated with agene-
sis of the
[163179]corpus callosum. Nature Genet. 7Oct, 2002. Note: Ad-
vance Electronic
[163180]Publication.

[163181]

[163182]

[163183]16195.

[163184]Haire, R. N.; Litman, G. W.: The murine form of TXK, a
novel TEC

[163185]kinase expressed in thymus maps to chromosome 5.

Mammalian Genome 6:

[163186]476-480, 1995.

[163187]

[163188]

[163189]16196.

[163190]Haire, R. N.; Ohta, Y.; Lewis, J. E.; Fu, S. M.; Kroisel, P.; Lit-

man,

[163191]G. W.: TXK, a novel human tyrosine kinase expressed in T cells shares

[163192]sequence identity with Tec family kinases and maps to 4p12. Hum.

[163193]Molec. Genet. 3: 897–901, 1994.

[163194]

[163195]

[163196]16197.

[163197]Schneider, H.; Schwartzberg, P. L.; Rudd, C. E.: Resting lymphocyte

[163198]kinase (Rlk/Txk) phosphorylates the YVKM motif and regulates PI 3-kinase

[163199]binding to T-cell antigen CTLA-4. Biochem. Biophys. Res. Commun. 252:

[163200]14–19, 1998.

[163201]

[163202]

[163203]16198.

[163204]Takeba, Y.; Nagafuchi, H.; Takeno, M.; Kashiwakura, J.;

Suzuki,

[163205]N.: Txk, a member of nonreceptor tyrosine kinase of Tec family, acts

[163206]as a Th1 cell-specific transcription factor and regulates IFN-gamma

[163207]gene transcription. J. Immun. 168: 2365-2370, 2002.

[163208]

[163209]

[163210]16199.

[163211]Gasdaska, J. R.; Gasdaska, P. Y.; Gallegos, A.; Powis, G.: Human

[163212]thioredoxin reductase gene localization to chromosomal position 12q23-q24.1

[163213]and mRNA distribution in human tissue. Genomics 37: 257-259, 1996.

[163214]

[163215]

[163216]16200.

[163217]Gasdaska, P. Y.; Gasdaska, J. R.; Cochran, S.; Powis, G.: Cloning

[163218]and sequencing of a human thioredoxin reductase. FEBS

Lett. 373:

[163219]5–9, 1995.

[163220]

[163221]

[163222]16201.

[163223]Svaren, J.; Apel, E. D.; Simburger, K. S.; Jenkins, N. A.;
Gilbert,

[163224]D. J.; Copeland, N. A.; Milbrandt, J.: The Nab2 and Stat6
genes share

[163225]a common transcription termination region. Genomics 41:
33–39, 1997.

[163226]

[163227]

[163228]16202.

[163229]Weichenrieder, O.; Wild, K.; Strub, K.; Cusack, S.: Structure

[163230]and assembly of the Alu domain of the mammalian signal
recognition

[163231]particle. Nature 408: 167–173, 2000.

[163232]

[163233]

[163234]16203.

[163235]Nichols, R. C.; Blinder, J.; Pai, S. I.; Ge, Q.; Targoff, I. N.;

[163236]Plotz, P. H.; Liu, P.: Assignment of two human autoantigen
genes:

[163237]isoleucyl-tRNA synthetase locates to 9q21 and lysyl-tRNA
synthetase

[163238]locates to 16q23-q24. Genomics 36: 210-213, 1996.

[163239]

[163240]

[163241]16204.

[163242]Nichols, R. C.; Raben, N.; Boerkoel, C. F.; Plotz, P. H.: Hu-
man

[163243]isoleucyl-tRNA synthetase: sequence of the cDNA, alter-
native mRNA

[163244]splicing, and the characteristics of an unusually long C-
terminal

[163245]extension. Gene 155: 299-304, 1995.

[163246]

[163247]

[163248]16205.

[163249]Dejgaard, K.; Leffers, H.; Rasmussen, H. H.; Madsen, P.;
Kruse,

[163250]T. A.; Gesser, B.; Nielsen, H.; Celis, J. E.: Identification,
molecular

[163251]cloning, expression and chromosome mapping of a family
of transformation

[163252]upregulated hnRNP-K proteins derived by alternative
splicing. J.

[163253]Molec. Biol. 236: 33-48, 1994.

[163254]

[163255]

[163256]16206.

[163257]Tommerup, N.; Leffers, H.: Assignment of human KH-
box-containing

[163258]genes by in situ hybridization: HNRNPK maps to
9q21.32-q21.33, PCBP1

[163259]to 2p12-p13, and PCBP2 to 12q13.12-q13.13, distal to
FRA12A. Genomics 32:

[163260]297-298, 1996.

[163261]

[163262]

[163263]16207.

[163264]Agarwal, A. K.; Monder, C.; Eckstein, B.; White, P. C.:

Cloning

[163265]and expression of rat cDNA encoding corticosteroid

11-beta-dehydrogenase. J.

[163266]Biol. Chem. 264: 18939-18943, 1989.

[163267]

[163268]

[163269]16208.

[163270]Agarwal, A. K.; Rogerson, F. M.; Mune, T.; White, P. C.:

Gene

[163271]structure and chromosomal localization of the human

HSD11K gene encoding

[163272]the kidney (type 2) isozyme of 11-beta-hydroxysteroid

dehydrogenase. Genomics 29:

[163273]195-199, 1995.

[163274]

[163275]

[163276]16209.

[163277]Lakshmi, V.; Monder, C.: Purification and characterization
of

[163278]the corticosteroid 11-beta-dehydrogenase component of
the rat liver

[163279]11-beta-hydroxysteroid dehydrogenase complex. En-
docrinology 123:

[163280]2390-2398, 1988.

[163281]

[163282]

[163283]16210.

[163284]Kotelevtsev, Y.; Holmes, M. C.; Burchell, A.; Houston, P.
M.; Schmoll,

[163285]D.; Jamieson, P.; Best, R.; Brown, R.; Edwards, C. R. W.;
Seckl, J.

[163286]R.; Mullins, J. J.: 11-beta-hydroxysteroid dehydrogenase
type 1 knockout

[163287]mice show attenuated glucocorticoid-inducible responses
and resist

[163288]hyperglycemia on obesity or stress. Proc. Nat. Acad. Sci.
94: 14924-14929,

[163289]1997.

[163290]

[163291]

[163292]16211.

[163293]Masuzaki, H.; Paterson, J.; Shinyama, H.; Morton, N. M.;
Mullins,

[163294]J. J.; Seckl, J. R.; Flier, J. S.: A transgenic model of visceral

[163295]obesity and the metabolic syndrome. Science 294:
2166–2170, 2001.

[163296]

[163297]

[163298]16212.

[163299]Morton, N. M.; Holmes, M. C.; Fievet, C.; Staels, B.;
Tailleux,

[163300]A.; Mullins, J. J.; Seck, J. R.: Improved lipid and lipoprotein
profile,

[163301]hepatic insulin sensitivity, and glucose tolerance in
11-beta-hydroxysteroid

[163302]dehydrogenase type 1 null mice. J. Biol. Chem. 276:
41293–41300,

[163303]2001.

[163304]

[163305]

[163306]16213.

[163307]Ricketts, M. L.; Verhaeg, J. M.; Bujalska, I.; Howie, A. J.;
Rainey,

[163308]W. E.; Stewart, P. M.: Immunohistochemical localization of
type 1

[163309]11-beta-hydroxysteroid dehydrogenase in human tissues.
J.Clin. Endocr.

[163310]Metab. 83: 1325-1335, 1998.

[163311]

[163312]

[163313]16214.

[163314]Alessi, D. R.; Smythe, C.; Keyse, S. M.: The human CL100
gene

[163315]encodes a tyr/thr-protein phosphatase which potently and
specifically

[163316]inactivates MAP kinase and suppresses its activation by
oncogenic

[163317]ras in Xenopus oocyte extracts. Oncogene 8: 2015-2020,
1993.

[163318]

[163319]

[163320]16215.

[163321]Brondello, J.-M.; Pouyssegur, J.; McKenzie, F. R.: Reduced
MAP

[163322]kinase phosphatase-1 degradation after
p42/p44(MAPK)-dependent phosphorylation. Science 286:

[163323]2514-2517, 1999.

[163324]

[163325]

[163326]16216.

[163327]Emslie, E. A.; Jones, T. A.; Sheer, D.; Keyse, S. M.: The
CL100

[163328]gene, which encodes a dual specificity (tyr/thr) MAP ki-
nase phosphatase,

[163329]is highly conserved and maps to human chromosome
5q34. Hum. Genet. 93:

[163330]513-516, 1994.

[163331]

[163332]

[163333]16217.

[163334]Keyse, S. M.; Emslie, E. A.: Oxidative stress and heat shock
induce

[163335]a human gene encoding a protein-tyrosine phosphatase.
Nature 359:

[163336]644-647, 1992.

[163337]

[163338]

[163339]16218.

[163340]Martell, K. J.; Kwak, S.; Hakes, D. J.; Dixon, J. E.; Trent, J.

[163341]M.: Chromosomal localization of four human VH1-like
protein-tyrosine

[163342]phosphatases. Genomics 22: 462-464, 1994.

[163343]

[163344]

[163345]16219.

[163346]Chrast, R.; Chen, H.; Morris, M. A.; Antonarakis, S. E.:
Mapping

[163347]of the human transcription factor GABPA (E4TF1-60) gene

to chromosome

[163348]21. Genomics 28: 119–122, 1995.

[163349]

[163350]

[163351]16220.

[163352]Gugneja, S.; Virbasius, J. V.; Scarpulla, R. C.: Four structurally

[163353]distinct, non-DNA-binding subunits of human nuclear respiratory factor

[163354]2 share a conserved transcriptional activation domain. Molec. Cell.

[163355]Biol. 15: 102–111, 1995.

[163356]

[163357]

[163358]16221.

[163359]Guo, A.; Nie, F.; Wong-Riley, M.: Human nuclear respiratory factor

[163360]2-alpha subunit cDNA: isolation, subcloning, sequencing, and in situ

[163361]hybridization of transcripts in normal and monocularly

deprived macaque

[163362]visual system. J. Comp. Neurol. 417: 221–232, 2000.

[163363]

[163364]

[163365]16222.

[163366]Sawada, J.; Goto, M.; Watanabe, H.; Handa, H.; Yoshida, M.
C.:

[163367]Regional mapping of two subunits of transcription factor
E4TF1 to

[163368]human chromosome. Jpn. J. Cancer Res. 86: 10–12, 1995.

[163369]

[163370]

[163371]16223.

[163372]Virbasius, J. V.; Scarpulla, R. C.: Activation of the human
mitochondrial

[163373]transcription factor A gene by nuclear respiratory factors:
a potential

[163374]regulatory link between nuclear and mitochondrial gene
expression

[163375]in organelle biogenesis. Proc. Nat. Acad. Sci. 91:

1309–1313, 1994.

[163376]

[163377]

[163378]16224.

[163379]Watanabe, H.; Sawada, J.; Yano, K.–I.; Yamaguchi, K.; Goto, M.;

[163380]Handa, H.: cDNA cloning of transcription factor E4TF1 subunits with

[163381]Ets and notch motifs. Molec. Cell. Biol. 13: 1385–1391, 1993.

[163382]

[163383]

[163384]16225.

[163385]Dixon, J.; Loftus, S. K.; Gladwin, A. J.; Scambler, P. J.; Wasmuth,

[163386]J. J.; Dixon, M. J.: Cloning of the human heparan sulfate–N–deacetylase/N–sulfotransferase

[163387]gene from the Treacher Collins syndrome candidate region at 5q32–q33.1. Genomics 26:

[163388]239–244, 1995.

[163389]

[163390]

[163391]16226.

[163392]Gladwin, A. J.; Dixon, J.; Loftus, S. K.; Wasmuth, J. J.;

Dixon,

[163393]M. J.: Genomic organization of the human heparan sul-

fate-N-deacetylase/N-sulfotransferase

[163394]gene: exclusion from a causative role in the pathogenesis

of Treacher

[163395]Collins syndrome. Genomics 32: 471-473, 1996.

[163396]

[163397]

[163398]16227.

[163399]Hashimoto, Y.; Orellana, A.; Gil, G.; Hirschberg, C. B.:

Molecular

[163400]cloning and expression of rat liver N-heparan sulfate sul-

fotransferase. J.

[163401]Biol. Chem. 267: 15744-15750, 1995.

[163402]

[163403]

[163404]16228.

[163405]Chen, H.; Antonarakis, S. E.: Localisation of a human homologue

[163406]of the *Drosophila* *mnb* and rat *Dyrk* genes to chromosome 21q22.2. Hum.

[163407]Genet. 99: 262–265, 1997.

[163408]

[163409]

[163410]16229.

[163411]Patil, N.; Cox, D. R.; Bhat, D.; Faham, M.; Myers, R. M.; Peterson,

[163412]A. S.: A potassium channel mutation in weaver mice implicates membrane

[163413]excitability in granule cell differentiation. Nature Genet. 11:

[163414]126–129, 1995.

[163415]

[163416]

[163417]16230.

[163418]Shindoh, N.; Kudoh, J.; Maeda, H.; Yamaki, A.; Minoshima, S.; Shimizu,

[163419]Y.; Shimizu, N.: Cloning of a human homolog of the
Drosophila minibrain/rat

[163420]Dyrk gene from 'the Down syndrome critical region' of
chromosome 21. Biochem.

[163421]Biophys. Res. Commun. 225: 92–99, 1996.

[163422]

[163423]

[163424]16231.

[163425]Smith, D. J.; Stevens, M. E.; Sudanagunta, S. P.; Bronson, R.
T.;

[163426]Makhinson, M.; Watabe, A. M.; O'Dell, T. J.; Fung, J.; Weier,
H.–U.

[163427]G.; Cheng, J.–F.; Rubin, E. M.: Functional screening of 2
Mb of human

[163428]chromosome 21q22.2 in transgenic mice implicates mini–
brain in learning

[163429]defects associated with Down syndrome. Nature Genet.
16: 28–36,

[163430]1997.

[163431]

[163432]

[163433]16232.

[163434]Song, W.-J.; Chung, S.-H.; Kurnit, D. M.: The murine Dyrk protein

[163435]maps to chromosome 16, localizes to the nucleus, and can form multimers. Biochem.

[163436]Biophys. Res. Commun. 231: 640–644, 1997.

[163437]

[163438]

[163439]16233.

[163440]Song, W.-J.; Sternberg, L. R.; Kasten-Sportes, C.; Van Keuren,

[163441]M. L.; Chung, S.-H.; Slack, A. C.; Miller, D. E.; Glover, T. W.; Chiang,

[163442]P.-W.; Lou, L.; Kurnit, D. M.: Isolation of human and murine homologues

[163443]of the Drosophila minibrain gene: human homologue maps to 21q22.2

[163444]in the Down syndrome 'critical region.' Genomics 38: 331–339, 1996.

[163445]

[163446]

[163447]16234.

[163448]Tejedor, F.; Zhu, X. R.; Kaltenbach, E.; Ackermann, A.;

Baumann,

[163449]A.; Canal, I.; Heisenberg, M.; Fischbach, K. F.; Pongs, O.:

Minibrain:

[163450]a new protein kinase family involved in postembryonic

neurogenesis

[163451]in *Drosophila*. *Neuron* 14: 287–301, 1995.

[163452]

[163453]

[163454]16235.

[163455]Niethammer, D.; Dieterle, U.; Kleihauer, E.; Wildfeuer, A.;

Haferkamp,

[163456]O.; Hitzig, W. H.: An inherited defect in granulocyte func-

tion: impaired

[163457]chemotaxis, phagocytosis and intracellular killing of mi-

croorganisms. *Helv.*

[163458]*Paediat. Acta* 30: 537–541, 1976.

[163459]

[163460]

[163461]16236.

[163462]Ostermann, G.; Weber, K. S. C.; Zerneck, A.; Schroder, A.;
Weber,

[163463]C.: JAM-1 is a ligand of the beta-2 integrin LFA-1 in-
volved in transendothelial

[163464]migration of leukocytes. Nature Immun. 3: 151-158,
2002.

[163465]

[163466]

[163467]16237.

[163468]Rivera-Matos, I. R.; Rakita, R. M.; Mariscalco, M. M.; Elder,

[163469]F. F. B.; Dreyer, S. A.; Cleary, T. G.: Leukocyte adhesion
deficiency

[163470]mimicking Hirschsprung disease. J. Pediat. 127: 755-757,
1995.

[163471]

[163472]

[163473]16238.

[163474]Rosmarin, A. G.; Caprio, D.; Levy, R.; Simkevich, C.: CD18
(beta-2

[163475]leukocyte integrin) promoter requires PU.1 transcription factor for

[163476]myeloid activity. Proc. Nat. Acad. Sci. 92: 801–805, 1995.

[163477]

[163478]

[163479]16239.

[163480]Ross, G. D.: Clinical and laboratory features of patients with

[163481]an inherited deficiency of neutrophil membrane complement receptor

[163482]type 3 (CR3) and the related membrane antigens LFA-1 and p150,95. J.

[163483]Clin. Immun. 6: 107–113, 1986.

[163484]

[163485]

[163486]16240.

[163487]Shuster, D. E.; Kehrli, M. E., Jr.; Ackermann, M. R.; Gilbert,

[163488]R. O.: Identification and prevalence of a genetic defect that causes

[163489]leukocyte adhesion deficiency in Holstein cattle. Proc. Nat.

Acad.

[163490]Sci. 89: 9225–9229, 1992.

[163491]

[163492]

[163493]16241.

[163494]Sligh, J. E., Jr.; Anderson, D. C.; Beaudet, A. L.: A mutation

[163495]in the initiation codon of the CD18 gene in a patient with
the moderate

[163496]phenotype of leukocyte adhesion deficiency. (Abstract)
Am. J. Hum.

[163497]Genet. 45 (suppl.): A219, 1989.

[163498]

[163499]

[163500]16242.

[163501]Solomon, E.; Palmer, R. W.; Hing, S.; Law, S. K. A.: Regional

[163502]localization of CD18, the beta-subunit of the cell surface
adhesion

[163503]molecule LFA-1, on human chromosome 21 by in situ hybridization. Ann.

[163504]Hum. Genet. 52: 123–128, 1988.

[163505]

[163506]

[163507]16243.

[163508]Springer, T. A.; Miller, L. J.; Anderson, D. C.: p150,95, the

[163509]third member of the Mac–1, LFA–1 human leukocyte ad–
hesion glycoprotein

[163510]family. J. Immun. 136: 240–245, 1986.

[163511]

[163512]

[163513]16244.

[163514]Springer, T. A.; Thompson, W. S.; Miller, L. J.; Schmalstieg,

[163515]F. C.; Anderson, D. C.: Inherited deficiency of the Mac–1,
LFA–1,

[163516]p150,95 glycoprotein family and its molecular basis. J.

Exp. Med. 160:

[163517]1901–1918, 1984.

[163518]

[163519]

[163520]16245.

[163521]Ross, G. D.; Thompson, R. A.; Walport, M. J.; Springer, T. A.;

[163522]Watson, J. V.; Ward, R. H. R.; Lida, J.; Newman, S. L.; Harrison,

[163523]R. A.; Lachmann, P. J.: Characterization of patients with an increased

[163524]susceptibility to bacterial infections and a genetic deficiency of

[163525]leukocyte membrane complement receptor type three (CR3) and the related

[163526]membrane antigen LFA-1. Blood 66: 882-890, 1985.

[163527]

[163528]

[163529]16246.

[163530]Suomalainen, H. A.; Gahmberg, C. G.; Patarroyo, M.; Beatty, P.

[163531]G.; Schroder, J.: Genetic assignment of GP90, leukocyte adhesion

[163532]glycoprotein to human chromosome 21. Somat. Cell Molec. Genet. 12:

[163533]297–302, 1986.

[163534]

[163535]

[163536]16247.

[163537]Suomalainen, H. A.; Gahmberg, C. G.; Patarroyo, M.;
Schroder,

[163538]J.: GP90 (Leu–CAM antigen) is coded for by genes on
chromosome 21.

[163539](Abstract) Cytogenet. Cell Genet. 40: 755, 1985.

[163540]

[163541]

[163542]16248.

[163543]Taylor, G. M.; Williams, A.; D'Souza, S. W.; Fergusson, W.
D.;

[163544]Donnai, D.; Fennell, J.; Harris, R.: The expression of CD18
is increased

[163545]on trisomy 21 (Down syndrome) lymphoblastoid cells.
Clin. Exp. Immun. 71:

[163546]324–328, 1988.

[163547]

[163548]

[163549]16249.

[163550]Todd, R. F., III; Freyer, D. R.: The CD11/CD18 leukocyte
glycoprotein

[163551]deficiency. Hemat. Oncol. Clin. North Am. 2: 13–31, 1988.

[163552]

[163553]

[163554]16250.

[163555]van der Meer, J. W. M.; van Zwet, T. L.; van Furth, R.: New
familial

[163556]defect in microbicidal function of polymorphonuclear leu-
cocytes. Lancet II:

[163557]630–632, 1975.

[163558]

[163559]

[163560]16251.

[163561]Vazquez–Torres, A.; Jones–Carson, J.; Baumler, A. J.;
Falkow,

[163562]S.; Valdivia, R.; Brown, W.; Le, M.; Berggren, R.; Parks, W.

T.; Fang,

[163563]F. C.: Extraintestinal dissemination of Salmonella by
CD18-expressing

[163564]phagocytes. Nature 401: 804–808, 1999.

[163565]

[163566]

[163567]16252.

[163568]Vedder, N. B.; Winn, R. K.; Rice, C. L.; Chi, E. Y.; Arfors,
K.–E.;

[163569]Harlan, J. M.: A monoclonal antibody to the adherence–
promoting leukocyte

[163570]glycoprotein, CD18, reduces organ injury and improves
survival from

[163571]hemorrhagic shock and resuscitation in rabbits. J. Clin. In–
vest. 81:

[163572]939–944, 1988.

[163573]

[163574]

[163575]16253.

[163576]Wardlaw, A. J.; Hibbs, M. L.; Stacker, S. A.; Springer, T. A.

[163577]: Distinct mutations in two patients with leukocyte adhesion deficiency

[163578]and their functional correlates. J. Exp. Med. 172: 335–345, 1990.

[163579]

[163580]

[163581]16254.

[163582]Weening, R. S.; Roos, D.; Weemaes, C. M. R.; Homan-Muller, J.

[163583]W. T.; van Schaik, M. L. J.: Defective initiation of the metabolic

[163584]stimulation in phagocytizing granulocytes: a new congenital defect. J.

[163585]Lab. Clin. Med. 88: 757–768, 1976.

[163586]

[163587]

[163588]16255.

[163589]Weitzman, J. B.; Wells, C. E.; Wright, A. H.; Clark, P. A.; Law,

[163590]S. K. A.: The gene organisation of the human beta-2 integrin subunit

[163591](CD18). FEBS Lett. 294: 97–103, 1991.

[163592]

[163593]

[163594]16256.

[163595]Wilson, J. M.; Ping, A. J.; Krauss, J. C.; Mayo–Bond, L.;
Rogers,

[163596]C. E.; Anderson, D. C.; Todd, R. F., III: Correction of
CD18–deficient

[163597]lymphocytes by retrovirus–mediated gene transfer. Sci–
ence 248: 1413–1416,

[163598]1990.

[163599]

[163600]

[163601]16257.

[163602]Wilson, R. W.; Ballantyne, C. M.; Smith, C. W.; Mont–
gomery, C.;

[163603]Bradley, A.; O'Brien, W. E.; Beaudet, A. L.: Gene targeting
yields

[163604]a CD18–mutant mouse for study of inflammation. J. Im–
mun. 151: 1571–1578,

[163605]1993.

[163606]

[163607]

[163608]16258.

[163609]Eichwald, E. J.; Silmser, C. R.: Skin. Transplant. Bull. 2:

[163610]148–149, 1955.

[163611]

[163612]

[163613]16259.

[163614]Gasser, D. L.; Silvers, W. K.: Genetics and immunology of
sex-linked

[163615]antigens. Adv. Immun. 15: 215–247, 1972.

[163616]

[163617]

[163618]16260.

[163619]Kent–First, M. G.; Maffitt, M.; Muallem, A.; Brisco, P.;
Shultz,

[163620]J.; Ekenberg, S.; Agulnik, A. I.; Agulnik, I.; Shramm, D.;

Bavister,

[163621]B.; Abdul-Mawgood, A.; VandeBerg, J.: Gene sequence and evolutionary

[163622]conservation of human SMCY. (Letter) Nature Genet. 14: 128–129,

[163623]1996.

[163624]

[163625]

[163626]16261.

[163627]Scott, D. M.; Ehrmann, I. E.; Ellis, P. S.; Bishop, C. E.; Agul-nik,

[163628]A. I.; Simpson, E.; Mitchell, M. J.: Identification of a mouse male-specific

[163629]transplantation antigen, H-Y. Nature 376: 695–698, 1995.

[163630]

[163631]

[163632]16262.

[163633]Wachtel, S. S.; Koo, G. C.; Breg, W. R.; Elias, S.; Boyse, E.

[163634]A.; Miller, O. J.: Expression of H-Y antigen in human males with

[163635]two Y chromosomes. New Eng. J. Med. 293: 1070–1072,
1975.

[163636]

[163637]

[163638]16263.

[163639]Wachtel, S. S.; Koo, G. C.; Zuckerman, E. E.; Hammerling,
U.;

[163640]Scheid, M. P.; Boyse, E. A.: Serological crossreactivity be-
tween

[163641]H–Y (male) antigens of mouse and man. Proc. Nat. Acad.
Sci. 71:

[163642]1215–1218, 1974.

[163643]

[163644]

[163645]16264.

[163646]Wang, W.; Meadows, L. R.; den Haan, J. M. M.; Sherman, N.
E.;

[163647]Chen, Y.; Blokland, E.; Shabanowitz, J.; Agulnik, A. I.; Hen-
drickson,

[163648]R. C.; Bishop, C. E.; Hunt, D. F.; Goulmy, E.; Engelhard, V.
H.:

[163649]Human H-Y: a male-specific histocompatibility antigen
derived from

[163650]the SMCY protein. Science 269: 1588–1590, 1995.

[163651]

[163652]

[163653]16265.

[163654]Redmond, T. M.; Yu, S.; Lee, E.; Bok, D.; Hamasaki, D.;
Chen,

[163655]N.; Goletz, P.; Ma, J.-X.; Crouch, R. K.; Pfeifer, K.: Rpe65 is
necessary

[163656]for production of 11-cis-vitamin A in the retinal visual cy-
cle. Nature

[163657]Genet. 20: 344–351, 1998.

[163658]

[163659]

[163660]16266.

[163661]Seeliger, M. W.; Grimm, C.; Stahlberg, F.; Friedburg, C.;
Jaissle,

[163662]G.; Zrenner, E.; Guo, H.; Reme, C. E.; Humphries, P.; Hof-
mann, F.;

[163663]Biel, M.; Fariss, R. N.; Redmond, T. M.; Wenzel, A.: New
views on

[163664]RPE65 deficiency: the rod system is the source of vision in
a mouse

[163665]model of Leber congenital amaurosis. Nature Genet. 29:
70–74, 2001.

[163666]

[163667]

[163668]16267.

[163669]Thompson, D. A.; Gyurus, P.; Fleischer, L.; Bingham, E. L.;
McHenry,

[163670]C. L.; Apfelstedt–Sylla, E.; Zrenner, E.; Lorenz, B.; Richards,
J.

[163671]E.; Jacobson, S. G.; Sieving, P. A.; Gal, A.: Genetics and
phenotypes

[163672]of RPE65 mutations in inherited retinal degeneration. In–
vest. Ophthal.

[163673]Vis. Sci. 41: 4293–4299, 2000.

[163674]

[163675]

[163676]16268.

[163677]Thompson, D. A.; McHenry, C. L.; Li, Y.; Richards, J. E.;
Othman,

[163678]M. I.; Schwinger, E.; Vollrath, D.; Jacobson, S. G.; Gal, A.:
Retinal

[163679]dystrophy due to paternal isodisomy for chromosome 1 or
chromosome

[163680]2, with homoallelism for mutations in RPE65 or MERTK,
respectively. Am.

[163681]J. Hum. Genet. 70: 224–229, 2002.

[163682]

[163683]

[163684]16269.

[163685]Wrigstad, A.; Narfstrom, K.; Nilsson, S. E.: Slowly progres-
sive

[163686]changes of the retina and retinal pigment epithelium in
Briard dogs

[163687]with hereditary retinal dystrophy: a morphological study.
Doc. Ophthal. 87:

[163688]337–354, 1994.

[163689]

[163690]

[163691]16270.

[163692]Bogenmann, E.; Lochrie, M. A.; Simon, M. I.: Cone cell-specific

[163693]genes expressed in retinoblastoma. Science 240: 76–78, 1988.

[163694]

[163695]

[163696]16271.

[163697]Bonaiti-Pellie, C.; Briard-Guillemot, M. L.: Segregation analysis

[163698]in hereditary retinoblastoma. Hum. Genet. 57: 411–419, 1981.

[163699]

[163700]

[163701]16272.

[163702]Harbitz, I.; Chowdhary, B.; Thomsen, P. D.; Davies, W.; Kaufmann,

[163703]U.; Kran, S.; Gustavsson, I.; Christensen, K.; Hauge, J. G.: Assignment

[163704]of the porcine calcium release channel gene, a candidate for the malignant

[163705]hyperthermia locus, to the 6p11-q21 segment of chromosome 6. Genomics 8:

[163706]243-248, 1990.

[163707]

[163708]

[163709]16273.

[163710]Hunter, A. G. W.; Thompson, M. W.: Acromesomelic dwarfism: description

[163711]of a patient and comparison with previously reported cases. Hum.

[163712]Genet. 34: 107-113, 1976.

[163713]

[163714]

[163715]16274.

[163716]Harbitz, I.; Kristensen, T.; Bosnes, M.; Kran, S.; Davies, W.

[163717]: DNA sequence of the skeletal muscle calcium release channel cDNA

[163718]and verification of the arg615-to-cys615 mutation, associated with

[163719]porcine malignant hyperthermia, in Norwegian Landrace

pigs. Animal

[163720]Genet. 23: 395–402, 1992.

[163721]

[163722]

[163723]16275.

[163724]Keating, K. E.; Quane, K. A.; Manning, B. M.; Lehane, M.;
Hartung,

[163725]E.; Censier, K.; Urwyler, A.; Klausnitzer, M.; Muller, C. R.;
Heffron,

[163726]J. J. A.; McCarthy, T. V.: Detection of a novel RYR1 muta-
tion in

[163727]four malignant hyperthermia pedigrees. Hum. Molec.
Genet. 3: 1855–1858,

[163728]1994.

[163729]

[163730]

[163731]16276.

[163732]Lynch, P. J.; Tong, J.; Lehane, M.; Mallet, A.; Giblin, L.; Hef-
fron,

[163733]J. J. A.; Vaughan, P.; Zafra, G.; MacLennan, D. H.; Mc-

Carthy, T. V.

[163734]: A mutation in the transmembrane/luminal domain of the ryanodine

[163735]receptor is associated with abnormal Ca^{2+} release channel function

[163736]and severe central core disease. Proc. Nat. Acad. Sci. 96: 4164–4169,

[163737]1999.

[163738]

[163739]

[163740]16277.

[163741]MacKenzie, A. E.; Korneluk, R. G.; Duff, C.; Worton, R. G.; MacLennan,

[163742]D. H.: Location of the human ryanodine receptor gene with respect

[163743]to an established human chromosome 19 linkage group.(Abstract) Am.

[163744]J. Hum. Genet. 45 (suppl.): A149, 1989.

[163745]

[163746]

[163747]16278.

[163748]MacKenzie, A. E.; Korneluk, R. G.; Zorzato, F.; Fujii, J.;
Phillips,

[163749]M.; Iles, D.; Wieringa, B.; Leblond, S.; Bailly, J.; Willard, H.
F.;

[163750]Duff, C.; Worton, R. G.; MacLennan, D. H.: The human
ryanodine receptor

[163751]gene: its mapping to 19q13.1, placement in a chromo-
some 19 linkage

[163752]group, and exclusion as the gene causing myotonic dys-
trophy. Am.

[163753]J. Hum. Genet. 46: 1082–1089, 1990.

[163754]

[163755]

[163756]16279.

[163757]MacLennan, D. H.: Personal Communication. Toronto, On-
tario,

[163758]Canada 7/5/1991.

[163759]

[163760]

[163761]16280.

- [163762]Manning, B. M.; Quane, K. A.; Lynch, P. J.; Urwyler, A.;
Tegazzin,
[163763]V.; Krivosic-Horber, R.; Censier, K.; Comi, G.; Adnet, P.;
Wolz, W.;
- [163764]Lunardi, J.; Muller, C. R.; McCarthy, T. V.: Novel mutations
at a
[163765]CpG dinucleotide in the ryanodine receptor in malignant
hyperthermia. Hum.
[163766]Mutat. 11: 45–50, 1998.
- [163767]
- [163768]
- [163769]16281.
- [163770]Mattei, M. G.; Giannini, G.; Moscatelli, F.; Sorrentino, V.:
- [163771]Chromosomal localization of murine ryanodine receptor
genes RYR1,
[163772]RYR2, and RYR3 by in situ hybridization. Genomics 22:
202–204, 1994.
- [163773]
- [163774]
- [163775]16282.

[163776]McCarthy, T. V.; Quane, K. A.; Lynch, P. J.: Ryanodine receptor

[163777]mutations in malignant hyperthermia and central core disease. Hum.

[163778]Mutat. 15: 410–417, 2000.

[163779]

[163780]

[163781]16283.

[163782]Monnier, N.; Romero, N. B.; Lerale, J.; Landrieu, P.; Nivoche,

[163783]Y.; Fardeau, M.; Lunardi, J.: Familial and sporadic forms of central

[163784]core disease are associated with mutations in the C-terminal domain

[163785]of the skeletal muscle ryanodine receptor. Hum. Molec. Genet. 10:

[163786]2581–2592, 2001.

[163787]

[163788]

[163789]16284.

[163790]MacLennan, D. H.; Zorzato, F.; Fujii, J.; Otsu, K.; Phillips,

[163791]M.; Lai, F. A.; Meissner, G.; Green, N. M.; Willard, H. F.;

Britt,

[163792]B. A.; Worton, R. G.; Korneluk, R. G.: Cloning and localization of

[163793]the human calcium release channel (ryanodine receptor) gene to the

[163794]proximal long arm (cen-q13.2) of human chromosome 19.(Abstract) Am.

[163795]J. Hum. Genet. 45 (suppl.): A205, 1989.

[163796]

[163797]

[163798]16285.

[163799]Manning, B. M.; Quane, K. A.; Ording, H.; Urwyler, A.; Tegazzin,

[163800]V.; Lehane, M.; O'Halloran, J.; Hartung, E.; Giblin, L. M.; Lynch,

[163801]P. J.; Vaughan, P.; Censier, K.; Bendixen, D.; Comi, G.; Heytens,

[163802]L.; Monsieurs, K.; Fagerlund, T.; Wolz, W.; Heffron, J. J. A.; Muller,

[163803]C. R.; McCarthy, T. V.: Identification of novel mutations in the

[163804]ryanodine-receptor gene (RYR1) in malignant hyperthermia: genotype-phenotype

[163805]correlation. Am. J. Hum. Genet. 62: 599-609, 1998.

[163806]

[163807]

[163808]16286.

[163809]Monnier, N.; Romero, N. B.; Lerale, J.; Nivoche, Y.; Qi, D.; MacLennan,

[163810]D. H.; Fardeau, M.; Lunardi, J.: An autosomal dominant congenital

[163811]myopathy with cores and rods is associated with a neo-mutation in the

[163812]RYR1 gene encoding the skeletal muscle ryanodine receptor. Hum. Molec.

[163813]Genet. 9: 2599-2608, 2000.

[163814]

[163815]

[163816]16287.

[163817]Monsieurs, K. G.; Van Broekhoven, C.; Martin, J.-J.; Van
Hoof,

[163818]V. O.; Heytens, L.: Gly341arg mutation indicating malignant hyperthermia

[163819]susceptibility: specific cause of chronically elevated serum
creatine

[163820]kinase activity. J. Neurol. Sci. 154: 62–65, 1998.

[163821]

[163822]

[163823]16288.

[163824]O'Brien, R. O.; Taske, N. L.; Hansbro, P. M.; Matthaei, K. I.;

[163825]Hogan, S. P.; Denborough, M. A.; Foster, P. S.: Exclusion of
defects

[163826]in the skeletal muscle specific regions of the DHPR alpha-
1 subunit

[163827]as frequent causes of malignant hyperthermia. J. Med.
Genet. 32:

[163828]913–914, 1995.

[163829]

[163830]

[163831]16289.

[163832]Otsu, K.; Khanna, V. K.; Archibald, A. L.; MacLennan, D. H.:

[163833]Cosegregation of porcine malignant hyperthermia and a
probable causal

[163834]mutation in the skeletal muscle ryanodine receptor gene
in backcross

[163835]families. Genomics 11: 744–750, 1991.

[163836]

[163837]

[163838]16290.

[163839]Otsu, K.; Nishida, K.; Kimura, Y.; Kuzuya, T.; Hori, M.; Ka-
mada,

[163840]T.; Tada, M.: The point mutation arg615-to-cys in the
Ca(2+) release

[163841]channel of skeletal sarcoplasmic reticulum is responsible
for hypersensitivity

[163842]to caffeine and halothane in malignant hyperthermia. J.
Biol. Chem. 269:

[163843]9413–9415, 1994.

[163844]

[163845]

[163846]16291.

[163847]Otsu, K.; Phillips, M. S.; Khanna, V. K.; de Leon, S.;

MacLennan,

[163848]D. H.: Refinement of diagnostic assays for a probable

causal mutation

[163849]for porcine and human malignant hyperthermia. Genomics

13: 835–837,

[163850]1992.

[163851]

[163852]

[163853]16292.

[163854]Phillips, M. S.; Fujii, J.; Khanna, V. K.; DeLeon, S.; Yoko-

bata,

[163855]K.; De Jong, P. J.; MacLennan, D. H.: The structural organi-

zation

[163856]of the human skeletal muscle ryanodine receptor (RYR1)

gene. Genomics 34:

[163857]24–41, 1996.

[163858]

[163859]

[163860]16293.

[163861]Quane, K. A.; Healy, J. M. S.; Keating, K. E.; Manning, B. M.;

[163862]Couch, F. J.; Palmucci, L. M.; Doriguzzi, C.; Fagerlund, T.
H.; Berg,

[163863]K.; Ording, H.; Bendixen, D.; Mortier, W.; Linz, U.; Muller,
C. R.;

[163864]McCarthy, T. V.: Mutations in the ryanodine receptor gene
in central

[163865]core disease and malignant hyperthermia. Nature Genet.
5: 51–55,

[163866]1993.

[163867]

[163868]

[163869]16294.

[163870]Richter, M.; Schleithoff, L.; Deufel, T.; Lehmann–Horn, F.;
Herrmann–Frank,

[163871]A.: Functional characterization of a distinct ryanodine re-
ceptor

[163872]mutation in human malignant hyperthermia–susceptible
muscle. J. Biol.

[163873]Chem. 272: 5256–5260, 1997.

[163874]

[163875]

[163876]16295.

[163877]Robinson, R. L.; Brooks, C.; Brown, S. L.; Ellis, F. R.; Hall,
sall,

[163878]P. J.; Quinnell, R. J.; Shaw, M.–A.; Hopkins, P. M.: RYR1
mutations

[163879]causing central core disease are associated with more se–
vere malignant

[163880]hyperthermia in vitro contracture test phenotypes. Hum.
Mutat. 20:

[163881]88–97, 2002.

[163882]

[163883]

[163884]16296.

[163885]Sambuughin, N.; McWilliams, S.; de Bantel, A.; Sivakumar,
K.;

[163886]Nelson, T. E.: Single–amino–acid deletion in the RYR1
gene, associated

[163887]with malignant hyperthermia susceptibility and unusual contraction

[163888]phenotype. Am. J. Hum. Genet. 69: 204–208, 2001.

[163889]

[163890]

[163891]16297.

[163892]Oesterling, J. E.; Chan, D. W.; Epstein, J. I.; Kimball, A. W.,

[163893]Jr.; Bruzek, D. J.; Rock, R. C.; Brendler, C. B.; Walsh, P. C.:

Prostate

[163894]specific antigen in the preoperative and postoperative evaluation

[163895]of localized prostatic cancer treated with radical prostate-ctomy. J.

[163896]Urol. 139: 766–772, 1988.

[163897]

[163898]

[163899]16298.

[163900]Riegman, P. H. J.; Vlietstra, R. J.; van der Korput, J. A. G.

M.;

[163901]Brinkmann, A. O.; Trapman, J.: The promoter of the

prostate-specific

[163902]antigen gene contains a functional androgen responsive
element. Molec.

[163903]Endocr. 5: 1921-1930, 1991.

[163904]

[163905]

[163906]16299.

[163907]van den Elsen, P.; Bruns, G.; Gerhard, D. S.; Pravtcheva, D.;

[163908]Jones, C.; Housman, D.; Ruddle, F. A.; Orkin, S.; Terhorst,
C.: Assignment

[163909]of the gene coding for the T3-delta subunit of the
T3--T-cell receptor

[163910]complex to the long arm of human chromosome 11 and
to mouse chromosome

[163911]9. Proc. Nat. Acad. Sci. 82: 2920-2924, 1985.

[163912]

[163913]

[163914]16300.

[163915]van Rijs, J.; Giguere, V.; Hurst, J.; van Agthoven, T.; Geurts

[163916]van Kessel, A.; Goyert, S.; Grosveld, F.: Chromosomal localization

[163917]of the human Thy-1 gene. Proc. Nat. Acad. Sci. 82: 5832-5835, 1985.

[163918]

[163919]

[163920]16301.

[163921]Williams, A. F.; Gagnon, J.: Neuronal cell Thy-1 glycoprotein:

[163922]homology with immunoglobulin. Science 216: 696-703, 1982.

[163923]

[163924]

[163925]16302.

[163926]Johansson, M.; Karlsson, A.: Cloning of the cDNA and chromosome

[163927]localization of the gene for human thymidine kinase 2. J. Biol. Chem. 272:

[163928]8454-8458, 1997.

[163929]

[163930]

[163931]16303.

[163932]Mandel, H.; Szargel, R.; Labay, V.; Elpeleg, O.; Saada, A.;

Shalata,

[163933]A.; Anbinder, Y.; Berkowitz, D.; Hartman, C.; Barak, M.;

Eriksson,

[163934]S.; Cohen, N.: The deoxyguanosine kinase gene is mutated in individuals

[163935]with depleted hepatocerebral mitochondrial DNA. Nature Genet. 29:

[163936]337–341, 2001. Note: Erratum: Nature Genet. 29: 491 only, 2001.

[163937]

[163938]

[163939]16304.

[163940]Saada, A.; Shaag, A.; Mandel, H.; Nevo, Y.; Eriksson, S.;

Elpeleg,

[163941]O.: Mutant mitochondrial thymidine kinase in mitochondrial DNA depletion

[163942]myopathy. Nature Genet. 29: 342–344, 2001.

[163943]

[163944]

[163945]16305.

[163946]Wang, L.; Munch-Petersen, B.; Herrstrom Sjoberg, A.; Hellman, U.;

[163947]Bergman, T.; Jornvall, H.; Eriksson, S.: Human thymidine kinase 2:

[163948]molecular cloning and characterisation of the enzyme activity with

[163949]antiviral and cytostatic nucleoside substrates. FEBS Lett. 443:

[163950]170-174, 1999.

[163951]

[163952]

[163953]16306.

[163954]Willecke, K.; Reuber, T.; Kucherlapati, R. S.; Ruddle, F. H.:

[163955]Human mitochondrial thymidine kinase is coded for by a gene on chromosome

[163956]16 of the nucleus. Somat. Cell Genet. 3: 237-245, 1977.

[163957]

[163958]

[163959]16307.

[163960]Chang, J. G.; Scarpa, A.; Eddy, R. L.; Byers, M. G.; Harris, A.

[163961]S.; Morrow, J. S.; Watkins, P.; Shows, T. B.; Forget, B. G.:

Cloning

[163962]of a portion of the chromosomal gene and cDNA for hu-

man beta-fodrin,

[163963]the nonerythroid form of beta-spectrin. Genomics 17:

287-293, 1993.

[163964]

[163965]

[163966]16308.

[163967]Watkins, P. C.; Eddy, R.; Forget, B. G.; Chang, J. G.;

Rochelle,

[163968]R.; Shows, T. B.: Assignment of a non-erythroid spectrin

gene to

[163969]human chromosome 2. (Abstract) Am. J. Hum. Genet. 43:

A161, 1988.

[163970]

[163971]

[163972]16309.

[163973]Miceli–Richard, C.; Lesage, S.; Rybojad, M.; Prieur, A.–M.;
Manouvrier–Hanu,

[163974]S.; Hafner, R.; Chamaillard, M.; Zouali, H.; Thomas, G.;
Hugot, J.–P.

[163975]: CARD15 mutations in Blau syndrome. Nature Genet. 29:
19–20, 2001.

[163976]

[163977]

[163978]16310.

[163979]Cong, F.; Spencer, S.; Cote, J.–F.; Wu, Y.; Tremblay, M. L.;
Lasky,

[163980]L. A.; Goff, S. P.: Cytoskeletal protein PSTPIP1 directs the
PEST–type

[163981]protein tyrosine phosphatase to the c–Abl kinase to medi–
ate Abl dephosphorylation. Molec.

[163982]Cell 6: 1413–1423, 2000.

[163983]

[163984]

[163985]16311.

[163986]Han, L. Wong, D.; Dhaka, A.; Afar, D.; White, M.; Xie, W.;
Herschman,

[163987]H.; Witte, O.; Colicelli, J.: Protein binding and signaling properties

[163988]of RIN1 suggest a unique effector function. Proc. Nat. Acad. Sci. 94:

[163989]4954–4959, 1997.

[163990]

[163991]

[163992]16312.

[163993]Jhanwar, S. C.; Neel, B. G.; Hayward, W. S.; Chaganti, R. S. K.

[163994]: Localization of the cellular oncogenes ABL, SIS, and FES on human

[163995]germ–line chromosomes. Cytogenet. Cell Genet. 38: 73–75, 1984.

[163996]

[163997]

[163998]16313.

[163999]Blank, V.; Andrews, N. C.: The Maf transcription factors: regulators

[164000]of differentiation. Trends Biochem. Sci. 22: 437–441, 1997.

[164001]

[164002]

[164003]16314.

[164004]Jamieson, R. V.; Perveen, R.; Kerr, B.; Carette, M.; Yardley, J.;

[164005]Heon, E.; Wirth, M. G.; van Heyningen, V.; Donnai, D.; Munier, F.;

[164006]Black, G. C. M.: Domain disruption and mutation of the bZIP transcription

[164007]factor, MAF, associated with cataract, ocular anterior segment dysgenesis

[164008]and coloboma. Hum. Molec. Genet. 11: 33–42, 2002.

[164009]

[164010]

[164011]16315.

[164012]Kim, J. I.; Li, T.; Ho, I. C.; Grusby, M. J.; Glimcher, L. H.:

[164013]Requirement for the c-Maf transcription factor in crystallin gene

[164014]regulation and lens development. Proc. Nat. Acad. Sci. 96: 3781–3785,

[164015]1999.

[164016]

[164017]

[164018]16316.

[164019]Nishizawa, M.; Kataoka, K.; Goto, N.; Fujiwara, K. T.;
Kawai, S.

[164020]: v-maf, a viral oncogene that encodes a 'leucine zipper'
motif. Proc.

[164021]Nat. Acad. Sci. 86: 7711-7715, 1989.

[164022]

[164023]

[164024]16317.

[164025]Yoshida, M. C.; Nishizawa, M.; Kataoka, K.; Goto, N.; Fuji-
wara,

[164026]K. T.; Kawai, S.: Localization of the human MAF pro-
tooncogene on

[164027]chromosome 16 to bands q22-q23. (Abstract) Cytogenet.
Cell Genet. 58:

[164028]2003 only, 1991.

[164029]

[164030]

[164031]16318.

[164032]Bao, S.; Tibbetts, R. S.; Brumbaugh, K. M.; Fang, Y.;

Richardson,

[164033]D. A.; Ali, A.; Chen, S. M.; Abraham, R. T.; Wang, X.-F.:

ATR/ATM-mediated

[164034]phosphorylation of human Rad17 is required for genotoxic stress responses. Nature 411:

[164035]969–974, 2001.

[164036]

[164037]

[164038]16319.

[164039]Barbosa, M. D. F. S.; Barrat, F. J.; Tchernev, V. T.; Nguyen, Q.

[164040]A.; Mishra, V. S.; Colman, S. D.; Pastural, E.; Dufourcq-Lagelouse,

[164041]R.; Fischer, A.; Holcombe, R. F.; Wallace, M. R.; Brandt, S. J.; de

[164042]Saint Basile, G.; Kingsmore S. F.: Identification of mutations in

[164043]two major mRNA isoforms of the Chediak-Higashi syn-

drome gene in human

[164044]and mouse. Hum. Molec. Genet. 6: 1091–1098, 1997.

[164045]

[164046]

[164047]16320.

[164048]Barrat, F. J.; Auloge, L.; Pastural, E.; Dufourcq Lagelouse,
R.;

[164049]Vilmer, E.; Cant, A. J.; Weissenbach, J.; Le Paslier, D.; Fis-
cher,

[164050]A.; de Saint Basile, G.: Genetic and physical mapping of
the Chediak–Higashi

[164051]syndrome on chromosome 1q42–43. Am. J. Hum. Genet.
59: 625–632,

[164052]1996.

[164053]

[164054]

[164055]16321.

[164056]Dufourcq–Lagelouse, R.; Lambert, N.; Duval, M.; Viot, G.;
Vilmer,

[164057]E.; Fischer, A.; Prieur, M.; de Saint Basile, G.: Chediak–Hi–

gashi

[164058]syndrome associated with maternal uniparental isodisomy
of chromosome

[164059]1. Europ. J. Hum. Genet. 7: 633–637, 1999.

[164060]

[164061]

[164062]16322.

[164063]Faigle, W.; Raposo, G.; Tenza, D.; Pinet, V.; Vogt, A. B.;
Kropshofer,

[164064]H.; Fischer, A.; de Saint-Basile, G.; Amigorena, S.: Defi-
cient peptide

[164065]loading and MHC class II endosomal sorting in a human
genetic immunodeficiency

[164066]disease: the Chediak–Higashi syndrome. J. Cell Biol. 141:
1121–1134,

[164067]1998.

[164068]

[164069]

[164070]16323.

[164071]Karim, M. A.; Suzuki, K.; Fukai, K.; Oh, J.; Nagle, D. L.;

Moore,

[164072]K. J.; Barbosa, E.; Falik-Borenstein, T.; Filipovich, A.;

Ishida,

[164073]Y. Kivrikko, S.; Klein, C.; and 8 others: Apparent geno-

type-phenotype

[164074]correlation in childhood, adolescent, and adult Chediak-

Higashi syndrome. Am.

[164075]J. Med. Genet. 108: 16-22, 2002.

[164076]

[164077]

[164078]16324.

[164079]Kunieda, T.; Ide, H.; Nakagiri, M.; Yoneda, K.; Konfortov, B.;

[164080]Ogawa, H.: Localization of the locus responsible for Che-diak-Higashi

[164081]syndrome in cattle to bovine chromosome 28. Anim. Genet. 31: 87-90,

[164082]2000.

[164083]

[164084]

[164085]16325.

[164086]Akama, T. O.; Nishida, K.; Nakayama, J.; Watanabe, H.; Ozaki, K.;

[164087]Nakamura, T.; Dota, A.; Kawasaki, S.; Inoue, Y.; Maeda, N.; Yamamoto,

[164088]S.; Fujiwara, T.; Thonar, E. J.-M. A.; Shimomura, Y.; Kinoshita, S.;

[164089]Tanigami, A.; Fukuda, M. N.: Macular corneal dystrophy type I and

[164090]type II are caused by distinct mutations in a new sulpho-transferase

[164091]gene. Nature Genet. 26: 237–241, 2000.

[164092]

[164093]

[164094]16326.

[164095]Enomoto, A.; Kimura, H.; Chairoungdua, A.; Shigeta, Y.;
Jutabha,

[164096]P.; Cha, S. H.; Hosoyamada, M.; Takeda, M.; Sekine, T.;
Igarashi,

[164097]T.; Matsuo, H.; Kikuchi, Y.; Oda, T.; Ichida, K.; Hosoya, T.;
Shimokata,

[164098]K.; Niwa, T.; Kanai, Y.; Endou, H.: Molecular identification
of a

[164099]renal urate–anion exchanger that regulates blood urate
levels. Nature 417:

[164100]447–452, 2002.

[164101]

[164102]

[164103]16327.

[164104]Goedert, M.; Hasegawa, J.; Craxton, M.; Leversha, M. A.;
Clegg,

[164105]S.: Assignment of the human stress-activated protein kinase-3 gene

[164106](SAPK3) to chromosome 22q13.3 by fluorescence in situ hybridization. Genomics 41:

[164107]501-502, 1997.

[164108]

[164109]

[164110]16328.

[164111]Yang, H.; Jeffrey, P. D.; Miller, J.; Kinnucan, E.; Sun, Y.; Thoma,

[164112]N. H.; Zheng, N.; Chen, P.-L.; Lee, W.-H.; Pavletich, N. P.: BRCA2

[164113]function in DNA binding and recombination from a BRCA2-DSS1-ssDNA

[164114]structure. Science 297: 1837-1848, 2002.

[164115]

[164116]

[164117]16329.

[164118]Koettnitz, K.; Kappel, B.; Baumruker, T.; Hauber, J.; Bevec, D.

[164119]: The genomic structure encoding human initiation factor

eIF-5A. Gene 144:

[164120]249-252, 1994.

[164121]

[164122]

[164123]16330.

[164124]Koettnitz, K.; Wohl, T.; Kappel, B.; Lottspeich, F.; Hauber,
J.;

[164125]Bevec, D.: Identification of a new member of the human
eIF-5A gene

[164126]family. Gene 159: 283-284, 1995.

[164127]

[164128]

[164129]16331.

[164130]Steinkasserer, A.; Jones, T.; Sheer, D.; Koettnitz, K.;
Hauber,

[164131]J.; Bevec, D.: The eukaryotic cofactor for the human im-
munodeficiency

[164132]virus type 1 (HIV-1) rev protein, eIF-5A, maps to chromo-
some 17p12-p13:

[164133]three eIF-5A pseudogenes map to 10q23.3, 17q25, and
19q13.2. Genomics 25:

[164134]749-752, 1995.

[164135]

[164136]

[164137]16332.

[164138]Hartley, D. A.; Preiss, A.; Artavanis-Tsakonas, S.: A de-
duced

[164139]gene product from the Drosophila neurogenic locus, en-
hancer of split,

[164140]shows homology to mammalian G-protein beta subunit.
Cell 55: 785-795,

[164141]1988.

[164142]

[164143]

[164144]16333.

[164145]Hou, E. W.; Li, S.-L.: Genomic organization and chromo-
some localization

[164146]to band 19p13.3 of the human AES gene: gene product
exhibits strong

[164147]similarity to the N-terminal domain of Drosophila enhancer of split

[164148]Groucho protein. DNA Cell Biol. 17: 911–913, 1998.

[164149]

[164150]

[164151]16334.

[164152]Miyasaka, H.; Choudhury, B. K.; Hou, E. W.; Li, S. S.-L.:
Molecular

[164153]cloning and expression of mouse and human cDNA encoding AES and ESG

[164154]proteins with strong similarity to Drosophila enhancer of split groucho

[164155]protein. Europ. J. Biochem. 216: 343–352, 1993.

[164156]

[164157]

[164158]16335.

[164159]Imai, Y.; Kurokawa, M.; Tanaka, K.; Friedman, A. D.;
Ogawa, S.;

[164160]Mitani, K.; Yazaki, Y.; Hirai, H.: TLE, the human homolog of groucho,

[164161]interacts with AML1 and acts as a repressor of
AML1–induced transactivation. Biochem.

[164162]Biophys. Res. Commun. 252: 582–589, 1998.

[164163]

[164164]

[164165]16336.

[164166]Liu, Y.; Dehni, G.; Purcell, K. J.; Sokolow, J.; Carcangiu, M.

[164167]L.; Artavanis–Tsakonas, S.; Stifani, S.: Epithelial expression
and

[164168]chromosomal location of human TLE genes: implications
for Notch signaling

[164169]and neoplasia. Genomics 31: 58–64, 1996.

[164170]

[164171]

[164172]16337.

[164173]Stifani, S.; Blaumueller, C. M.; Redhead, N. J.; Hill, R. E.;
Artavanis–Tsakonas,

[164174]S.: Human homologs of a Drosophila enhancer of split
gene product

[164175]define a novel family of nuclear proteins. Nature Genet. 2:

119–127,

[164176]1992.

[164177]

[164178]

[164179]16338.

[164180]de Bruijn, D. R. H.; Baats, E.; Zechner, U.; de Leeuw, B.;
Balemans,

[164181]M.; Olde Weghuis, D.; Hirning–Folz, U.; Geurts van Kessel,
A.: Isolation

[164182]and characterization of the mouse homolog of SYT, a
gene implicated

[164183]in the development of human synovial sarcomas. Onco–
gene 13: 643–648,

[164184]1996.

[164185]

[164186]

[164187]16339.

[164188]de Bruijn, D. R. H.; Kater–Baats, E.; Eleveld, M.; Merkx, G.;
van

[164189]Kessel, A. G.: Mapping and characterization of the mouse

and human

[164190]SS18 genes, two human SS18-like genes and a mouse
Ss18 pseudogene. Cytogenet.

[164191]Cell Genet. 92: 310-319, 2001.

[164192]

[164193]

[164194]16340.

[164195]Thaete, C.; Brett, D.; Monaghan, P.; Whitehouse, S.; Ren-
nie, G.;

[164196]Rayner, E.; Cooper, C. S.; Goodwin, G.: Functional domains
of the

[164197]SYT and SYT-SSX synovial sarcoma translocation proteins
and co-localization

[164198]with the SNF protein BRM in the nucleus. Hum. Molec.
Genet. 8: 585-591,

[164199]1999.

[164200]

[164201]

[164202]16341.

[164203]Ge, Q.; Trieu, E. P.; Targoff, I. N.: Primary structure and

functional

[164204]expression of human glycyl-tRNA synthetase, an autoantigen in myositis.

[164205]J. Biol. Chem. 269: 28790-28797, 1994.

[164206]

[164207]

[164208]16342.

[164209]Nichols, R. C.; Pai, S. I.; Ge, Q.; Targoff, I. N.; Plotz, P. H.;

[164210]Liu, P.: Localization of two human autoantigen genes by PCR screening

[164211]and in situ hybridization--Glycyl-tRNA synthetase locates to 7p15

[164212]and alanyl-tRNA synthetase locates to 16q22. Genomics 30: 131-132,

[164213]1995.

[164214]

[164215]

[164216]16343.

[164217]Shiba, K.; Schimmel, P.; Motegi, H.; Noda, T.: Human glycyl-tRNA

[164218]synthetase: wide divergence of primary structure from
bacterial counterpart

[164219]and species-specific aminoacylation. J. Biol. Chem. 269:
30049–30055,

[164220]1994.

[164221]

[164222]

[164223]16344.

[164224]Williams, J.; Osvath, S.; Khong, T. F.; Pearse, M.; Power, D.:

[164225]Cloning, sequencing and bacterial expression of human
glycine tRNA

[164226]synthetase. Nucleic Acids Res. 23: 1307–1310, 1995.

[164227]

[164228]

[164229]16345.

[164230]Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek,

[164231]F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.;
Copeland, N.

[164232]G.: Murine chromosomal location of four hepatocyte-en-

riched transcription

[164233]factors: HNF-3-alpha, HNF-3-beta, HNF-3-gamma, and
HFN-4. Genomics 13:

[164234]264-268, 1992.

[164235]

[164236]

[164237]16346.

[164238]Deleuze, J. F.; Dhorne, S.; Hazan, J.; Borghi, E.; Raynaud,
N.;

[164239]Pollet, N.; Meunier-Rotival, M.; Deschatrette, J.; Alagille,
D.; Hadchouel,

[164240]M.: Deleted chromosome 20 from a patient with Alagille
syndrome isolated

[164241]in a cell hybrid through leucine transport selection: study
of three

[164242]candidate genes. Mammalian Genome 5: 663-669, 1994.

[164243]

[164244]

[164245]16347.

[164246]Kaestner, K. H.; Hiemisch, H.; Luckow, B.; Schutz, G.: The

HNF-3

[164247]gene family of transcription factors in mice: gene structure, cDNA

[164248]sequence, and mRNA distribution. Genomics 20: 377-385, 1994.

[164249]

[164250]

[164251]16348.

[164252]Mincheva, A.; Lichter, P.; Schutz, G.; Kaestner, K. H.: Assignment

[164253]of the human genes for hepatocyte nuclear factor 3-alpha, -beta, and

[164254]-gamma (HNF3A, HNF3B, HNF3G) to 14q12-q13, 20p11, and 19q13.2-q13.4. Genomics 39:

[164255]417-419, 1997.

[164256]

[164257]

[164258]16349.

[164259]Adams, R. H.; Porras, A.; Alonso, G.; Jones, M.; Vintersten, K.;

[164260]Panelli, S.; Valladares, A.; Perez, L.; Klein, R.; Nebreda, A.

R.

[164261]: Essential role of p38–alpha MAP kinase in placental but not embryonic

[164262]cardiovascular development. Molec. Cell 6: 109–116, 2000.

[164263]

[164264]

[164265]16350.

[164266]Ge, B.; Gram, H.; Di Padova, F.; Huang, B.; New, L.; Ulevitch,

[164267]R. J.; Luo, Y.; Han, J.: MAPKK–independent activation of p38–alpha

[164268]mediated by TAB1–dependent autophosphorylation of p38–alpha. Science 295:

[164269]1291–1294, 2002.

[164270]

[164271]

[164272]16351.

[164273]Han, J.; Lee, J.–D.; Bibbs, L.; Ulevitch, R. J.: A MAP kinase

[164274]targeted by endotoxin and hyperosmolarity in mammalian

cells. Science 265:

[164275]808–811, 1994.

[164276]

[164277]

[164278]16352.

[164279]Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.;
Garsin,

[164280]D. A.; Inoue, H.; Tanaka-Hino, M.; Hisamoto, N.; Mat-
sumoto, K.; Tan,

[164281]M.-W.; Ausubel, F. M.: A conserved p38 MAP kinase path-
way in *Caenorhabditis*

[164282]*elegans* innate immunity. Science 297: 623–626, 2002.

[164283]

[164284]

[164285]16353.

[164286]Kumar, S.; McLaughlin, M. M.; McDonnell, P. C.; Lee, J. C.;
Livi,

[164287]G. P.; Young, P. R.: Human mitogen-activated protein ki-
nase CSBP1,

[164288]but not CSBP2, complements a *hog1* deletion in yeast. J.

Biol. Chem. 270:

[164289]29043–29046, 1995.

[164290]

[164291]

[164292]16354.

[164293]Lee, J. C.; Laydon, J. T.; McDonnell, P. C.; Gallagher, T. F.;

[164294]Kumar, S.; Green, D.; McNulty, D.; Blumenthal, M. J.; Heys,
J. R.;

[164295]Landvatter, S. W.; Stickler, J. E.; McLaughlin, M. M.;
Siemens, I.

[164296]R.; Fisher, S. M.; Livi, G. P.; White, J. R.; Adams, J. L.;
Young,

[164297]P. R.: A protein kinase involved in the regulation of in-
flammatory

[164298]cytokine biosynthesis. Nature 372: 739–746, 1994.

[164299]

[164300]

[164301]16355.

[164302]Liao, P.; Georgakopoulos, D.; Kovacs, A.; Zheng, M.;
Lerner, D.;

[164303]Pu, H.; Saffitz, J.; Chien, K.; Xiao, R.-P.; Kass, D. A.; Wang, Y.

[164304]: The in vivo role of p38 MAP kinases in cardiac remodeling and restrictive

[164305]cardiomyopathy. Proc. Nat. Acad. Sci. 98: 12283–12288, 2001.

[164306]

[164307]

[164308]16356.

[164309]Maizels, E. T.; Mukherjee, A.; Sithanandam, G.; Peters, C. A.;

[164310]Cottom, J.; Mayo, K. E.; Hunzicker-Dunn, M.: Developmental regulation

[164311]of mitogen-activated protein kinase-activated kinases-2 and -3 (MAPKAPK-2/-3)

[164312]in vivo during corpus luteum formation in the rat. Molec. Endocr. 15:

[164313]716–733, 2001.

[164314]

[164315]

[164316]16357.

[164317]McDonnell, P. C.; DiLella, A. G.; Lee, J. C.; Young, P. R.:

Localization

[164318]of the human stress responsive MAP kinase-like CSAIDs
binding protein

[164319](CSBP) gene to chromosome 6p21.3/21.2. Genomics 29:
301–302, 1995.

[164320]

[164321]

[164322]16358.

[164323]New, L.; Jiang, Y.; Zhao, M.; Liu, K.; Zhu, W.; Flood, L. J.;

[164324]Kato, Y.; Parry, G. C. N.; Han, J.: PRAK, a novel protein ki-
nase

[164325]regulated by the p38 MAP kinase. EMBO J. 17: 3372–3384,
1998.

[164326]

[164327]

[164328]16359.

[164329]Ni, H.; Wang, X. S.; Diener, K.; Yao, Z.: MAPKAPK5, a novel
mitogen-activate

[164330]d protein kinase (MAPK)–activated protein kinase, is a
substrate of

[164331]the extracellular-regulated kinase (ERK) and p38 kinase.

Biochem.

[164332]Biophys. Res. Commun. 243: 492–496, 1998.

[164333]

[164334]

[164335]16360.

[164336]Takekawa, M.; Maeda, T.; Saito, H.: Protein phosphatase
2C-alpha

[164337]inhibits the human stress-responsive p38 and JNK MAPK
pathways. EMBO

[164338]J. 17: 4744–4752, 1998.

[164339]

[164340]

[164341]16361.

[164342]Tamura, K.; Sudo, T.; Senftleben, U.; Dadak, A. M.; John-
son, R.;

[164343]Karin, M.: Requirement for p38-alpha in erythropoietin
expression:

[164344]a role for stress kinases in erythropoiesis. Cell 102:
221–231,

[164345]2000.

[164346]

[164347]

[164348]16362.

[164349]Haber, N.; Stengel, D.; Defer, N.; Roeckel, N.; Mattei,
M.-G.;

[164350]Hanoune, J.: Chromosomal mapping of human adenylyl
cyclase genes

[164351]type III, type V and type VI. Hum. Genet. 94: 69–73, 1994.

[164352]

[164353]

[164354]16363.

[164355]Sinnarajah, S.; Dessauer, C. W.; Srikumar, D.; Chen, J.;
Yuen,

[164356]J.; Yilma, S.; Dennis, J. C.; Morrison, E. E.; Vodyanoy, V.;
Kehrl,

[164357]J. H.: RGS2 regulates signal transduction in olfactory neu-
rons by

[164358]attenuating activation of adenylyl cyclase III. Nature 409:
1051–1055,

[164359]2001.

[164360]

[164361]

[164362]16364.

[164363]Pellizzoni, L.; Kataoka, N.; Charroux, B.; Dreyfuss, G.: A
novel

[164364]function for SMN, the spinal muscular atrophy gene prod-
uct, in pre-mRNA

[164365]splicing. Cell 95: 615-624, 1998.

[164366]

[164367]

[164368]16365.

[164369]Brown, K. A.; Leek, J. P.; Lench, N. J.; Moynihan, L. M.;
Markham,

[164370]A. F.; Mueller, R. F.: Human sequences homologous to the
gene for

[164371]the cochlear protein Ocp-II do not map to currently known
non-syndromic

[164372]hearing loss loci. Ann. Hum. Genet. 60: 385-389, 1996.

[164373]

[164374]

[164375]16366.

[164376]Chen, H.; Thalmann, I.; Adams, J. C.; Avraham, K. B.;
Copeland,

[164377]N. G.; Jenkins, N. A.; Beier, D. R.; Corey, D. P.; Thalmann,
R.; Duyk,

[164378]G. M.: cDNA cloning, tissue distribution, and chromoso-
mal localization

[164379]of Ocp2, a gene encoding a putative transcription-associ-
ated factor

[164380]predominantly expressed in the auditory organs. Ge-
nomics 27: 389-398,

[164381]1995.

[164382]

[164383]

[164384]16367.

[164385]Demetrick, D. J.; Zhang, H.; Beach, D. H.: Chromosomal
mapping

[164386]of the genes for the human CDK2/cyclin A-associated
proteins p19 (SKP1A

[164387]and SKP1B) and p45 (SKP2). Cytogenet. Cell Genet. 73:

104–107, 1996.

[164388]

[164389]

[164390]16368.

[164391]Liang, Y.; Chen, H.; Asher, J. H., Jr.; Chang, C.–C.; Friedman,
man,

[164392]T. B.: Human inner ear OCP2 cDNA maps to 5q22–5q35.2
with related

[164393]sequences on chromosomes 4p16.2–4p14, 5p13–5q22,
7pter–q22, 10 and

[164394]12p13–12qter. Gene 184: 163–167, 1997.

[164395]

[164396]

[164397]16369.

[164398]Sowden, J.; Morrison, K.; Schofield, J.; Putt, W.; Edwards, Y.

[164399]: A novel cDNA with homology to an RNA polymerase II
elongation factors

[164400]maps to human chromosome 5q31 (TCEB1L) and to
mouse chromosome 11

[164401](Tceb1l). Genomics 29: 145–151, 1995.

[164402]

[164403]

[164404]16370.

[164405]Thalmann, I.; Rosenthal, H. L.; Moore, B. W.; Thalmann, R.:

Organ

[164406]of Corti-specific polypeptides: OCP-I and OCP-II. Arch.

Oto-Rhino-Laryngol. 226:

[164407]123-128, 1980.

[164408]

[164409]

[164410]16371.

[164411]Thalmann, I.; Takahashi, K.; Varghese, J.; Comegys, T. H.;

Thalmann,

[164412]R.: Biochemical features of major organ of Corti proteins

(OCP-I

[164413]and OCP-II) including partial amino acid sequence. Laryn-

goscope 100:

[164414]99-105, 1990.

[164415]

[164416]

[164417]16372.

[164418]Zhang, H.; Kobayashi, R.; Galaktionov, K.; Beach, D.:

p19Skp1

[164419]and p45Skp2 are essential elements of the cyclin A-CDK2
S phase kinase. Cell 82:

[164420]915-925, 1995.

[164421]

[164422]

[164423]16373.

[164424]Anand, A.; Chada, K.: In vivo modulation of Hmgic reduces
obesity. Nature

[164425]Genet. 24: 377-380, 2000.

[164426]

[164427]

[164428]16374.

[164429]Arlotta, P.; Tai, A. K.-F.; Manfioletti, G.; Clifford, C.; Jay,

[164430]G.; Ono, S. J.: Transgenic mice expressing a truncated
form of the

[164431]high mobility group I-C protein develop adiposity and an

abnormally

[164432]high prevalence of lipomas. J. Biol. Chem. 275:
14394–14400, 2000.

[164433]

[164434]

[164435]16375.

[164436]Ashar, H. R.; Cherath, L.; Przybysz, K. M.; Chada, K.: Ge-
nomic

[164437]characterization of human HMGIC, a member of the ac-
cessory transcription

[164438]factor family found at translocation breakpoints in lipo-
mas. Genomics 31:

[164439]207–214, 1996.

[164440]

[164441]

[164442]16376.

[164443]Ashar, H. R.; Schoenberg Fejzo, M.; Tkachenko, A.; Zhou,
X.; Fletcher,

[164444]J. A.; Weremowicz, S.; Morton, C. C.; Chada, K.: Disruption
of the

[164445]architectural factor HMGI-C: DNA-binding AT hook motifs

fused in lipomas

[164446]to distinct transcriptional regulatory domains. Cell 82:

57–65,

[164447]1995.

[164448]

[164449]

[164450]16377.

[164451]Chau, K.–Y.; Patel, U. A.; Lee, K.–L. D.; Lam, H.–Y. P.;

Crane–Robinson,

[164452]C.: The gene for the human architectural transcription

factor HMGI–C

[164453]consists of five exons each coding for a distinct functional

element. Nucleic

[164454]Acids Res. 23: 4262–4266, 1995.

[164455]

[164456]

[164457]16378.

[164458]Danforth, E., Jr.: Failure of adipocyte differentiation causes

[164459]type II diabetes mellitus. Nature Genet. 26: 13 only, 2000.

[164460]

[164461]

[164462]16379.

[164463]Friedmann, M.; Holth, L. T.; Zoghbi, H. Y.; Reeves, R.: Or-
ganization,

[164464]inducible-expression and chromosome localization of the
human HMG-I(Y)

[164465]nonhistone protein gene. Nucleic Acids Res. 21:
4259-4267, 1993.

[164466]

[164467]

[164468]16380.

[164469]Ishwad, C. S.; Shriver, M. D.; Lassige, D. M.; Ferrell, R. E.:

[164470]The high mobility group I-C gene (HMGI-C): polymor-
phism and genetic

[164471]localization. Hum. Genet. 99: 103-105, 1997.

[164472]

[164473]

[164474]16381.

[164475]Kazmierczak, B.; Dal Cin, P.; Wanschura, S.; Bartnitzke, S.;
Van

[164476]den Berghe, H.; Bullerdiek, J.: Cloning and molecular characterization

[164477]of part of a new gene fused to HMGIC in mesenchymal tumors. Am. J.

[164478]Path. 152: 431–435, 1998.

[164479]

[164480]

[164481]16382.

[164482]Kazmierczak, B.; Pohnke, Y.; Bullerdiek, J.: Fusion transcripts

[164483]between the HMGIC gene and RTVL–H–related sequences in mesenchymal

[164484]tumors without cytogenetic aberrations. Genomics 38: 223–226, 1996.

[164485]

[164486]

[164487]16383.

[164488]Manfioletti, G.; Rustighi, A.; Mantovani, F.; Goodwin, G. H.;

[164489]Giancotti, V.: Isolation and characterization of the gene coding

[164490]for murine high-mobility-group protein HMGI-C. Gene 167: 249-253,

[164491]1995.

[164492]

[164493]

[164494]16384.

[164495]Mine, N.; Kurose, K.; Nagai, H.; Doi, D.; Ota, Y.; Yoneyama, K.;

[164496]Konishi, H.; Araki, T.; Emi, M.: Gene fusion involving HMGIC is a

[164497]frequent aberration in uterine leiomyomas. J. Hum. Genet. 46: 408-412,

[164498]2001.

[164499]

[164500]

[164501]16385.

[164502]Nucci, M. R.; Weremowicz, S.; Neskey, D. M.; Sornberger, K.; Tallini,

[164503]G.; Morton, C. C.; Quade, B. J.: Chromosomal translocation
t(8;12)

[164504]induces aberrant HMGIC expression in aggressive angiosarcoma of the

[164505]vulva. Genes Chromosomes Cancer 32: 172–176, 2001.

[164506]

[164507]

[164508]16386.

[164509]Rodrigues, N. R.; Owen, N.; Talbot, K.; Ignatius, J.;
Dubowitz,

[164510]V.; Davies, K. E.: Deletions in the survival motor neuron
gene on

[164511]5q13 in autosomal recessive spinal muscular atrophy.
Hum. Molec.

[164512]Genet. 4: 631–634, 1995.

[164513]

[164514]

[164515]16387.

[164516]Rossoll, W.; Kroning, A.-K.; Ohndorf, U.-M.; Steegborn, C.;
Jablonka,

[164517]S.; Sendtner, M.: Specific interaction of Smn, the spinal muscular

[164518]atrophy determining gene product, with hnRNP-R and gry-rbp/hnRNP-Q:

[164519]a role for Smn in RNA processing in motor axons? Hum. Molec. Genet. 11:

[164520]93–105, 2002.

[164521]

[164522]

[164523]16388.

[164524]Simard, L. R.; Rochette, C.; Semionov, A.; Morgan, K.; Vanasse,

[164525]M.: SMN(T) and NAIP mutations in Canadian families with spinal muscular

[164526]atrophy (SMA): genotype/phenotype correlations with disease severity. Am.

[164527]J. Med. Genet. 72: 51–58, 1997.

[164528]

[164529]

[164530]16389.

[164531]Sossi, V.; Giuli, A.; Vitali, T.; Tiziano, F.; Mirabella, M.;

[164532]Antonelli, A.; Neri, G.; Brahe, C.: Premature termination mutations

[164533]in exon 3 of the SMN1 gene are associated with exon skipping and a

[164534]relatively mild SMA phenotype. *Europ. J. Hum. Genet.* 9: 113–120,

[164535]2001.

[164536]

[164537]

[164538]16390.

[164539]Stewart, H.; Wallace, A.; McGaughran, J.; Mountford, R.; Kingston,

[164540]H.: Molecular diagnosis of spinal muscular atrophy. *Arch. Dis. Child.* 78:

[164541]531–535, 1998.

[164542]

[164543]

[164544]16391.

[164545]Talbot, K.; Ponting, C. P.; Theodosiou, A. M.; Rodrigues, N.

R.;

[164546]Surtees, R.; Mountford, R.; Davies, K. E.: Missense mutation clustering

[164547]in the survival motor neuron gene: a role for a conserved tyrosine

[164548]and glycine rich region of the protein in RNA metabolism?

Hum. Molec.

[164549]Genet. 6: 497–500, 1997.

[164550]

[164551]

[164552]16392.

[164553]van der Steege, G.; Grootschloten, P. M.; Cobben, J. M.; Zappata,

[164554]S.; Scheffer, H.; den Dunnen, J. T.; van Ommen, G.–J. B.; Brahe, C.;

[164555]Buys, C. H. C. M.: Apparent gene conversions involving the SMN gene

[164556]in the region of the spinal muscular atrophy locus on chromosome 5. Am.

[164557]J. Hum. Genet. 59: 834–838, 1996.

[164558]

[164559]

[164560]16393.

[164561]Viollet, L.; Bertrand, S.; Bueno Brunialti, A. L.; Lefebvre,

[164562]S.; Burlet, P.; Clermont, O.; Cruaud, C.; Guenet, J.-L.;

Munnich,

[164563]A.; Melki, J.: cDNA isolation, expression, and chromosomal localization

[164564]of the mouse survival motor neuron gene (Smn). *Genomics* 40: 185–188,

[164565]1997.

[164566]

[164567]

[164568]16394.

[164569]Wang, C. H.; Xu, J.; Carter, T. A.; Ross, B. M.; Dominski, M.

[164570]K.; Bellcross, C. A.; Penchaszadeh, G. K.; Munsat, T. L.;

Gilliam,

[164571]T. C.: Characterization of survival motor neuron (SMN) gene deletions

[164572]in asymptomatic carriers of spinal muscular atrophy. *Hum. Molec.*

[164573]Genet. 5: 359–365, 1996.

[164574]

[164575]

[164576]16395.

[164577]Wang, J.; Dreyfuss, G.: A cell system with targeted disruption
tion

[164578]of the SMN gene: functional conservation of the SMN protein and dependence

[164579]of Gemin2 on SMN. J. Biol. Chem. 276: 9599–9605, 2001.

[164580]

[164581]

[164582]16396.

[164583]Wirth, B.: An update of the mutation spectrum of the survival
vival

[164584]motor neuron gene (SMN1) in autosomal recessive spinal
muscular atrophy

[164585](SMA). Hum. Mutat. 15: 228–237, 2000.

[164586]

[164587]

[164588]16397.

[164589]Wirth, B.; Hahnen, E.; Morgan, K.; DiDonato, C. J.; Dadze, A.;

[164590]Rudnik-Schoneborn, S.; Simard, L. R.; Zerres, K.; Burghes, A. H. M.

[164591]: Allelic association and deletions in autosomal recessive proximal

[164592]spinal muscular atrophy: association of marker genotype with disease

[164593]severity and candidate cDNAs. Hum. Molec. Genet. 4: 1273-1284, 1995.

[164594]

[164595]

[164596]16398.

[164597]Wirth, B.; Herz, M.; Wetter, A.; Moskau, S.; Hahnen, E.; Rudnik-Schoneborn,

[164598]S.; Wienker, T.; Zerres, K.: Quantitative analysis of survival motor

[164599]neuron copies: identification of subtle SMN1 mutations in patients

[164600]with spinal muscular atrophy, genotype-phenotype correlation, and

[164601]implications for genetic counseling. Am. J. Hum. Genet.
64: 1340–1356,
[164602]1999.

[164603]

[164604]

[164605]16399.

[164606]Young, P. J.; Man, N.; Lorson, C. L.; Le, T. T.; Androphy, E.

[164607]J.; Burghes, A. H. M.; Morris, G. E.: The exon 2b region of
the spinal

[164608]muscular atrophy protein, SMN, is involved in self-
association and

[164609]SIP1 binding. Hum. Molec. Genet. 9: 2869–2877, 2000.

[164610]

[164611]

[164612]16400.

[164613]David, C.; Solimena, M.; De Camilli, P.: Autoimmunity in
stiff-man

[164614]syndrome with breast cancer is targeted to the C-terminal
region of

[164615]human amphiphysin, a protein similar to the yeast pro-

teins, Rvs167

[164616]and Rvs161. FEBS Lett. 351: 73–79, 1994.

[164617]

[164618]

[164619]16401.

[164620]De Camilli, P.; Thomas, A.; Cofiell, R.; Folli, F.; Lichte, B.;

[164621]Piccolo, G.; Meinck, H.–M.; Austoni, M.; Fassetta, G.; Bot-
tazzo, G.;

[164622]Bates, D.; Cartlidge, N.; Solimena, M.; Kilimann, M. W.: The
synaptic

[164623]vesicle-associated protein amphiphysin is the 128-kD au-
toantigen of

[164624]stiff-man syndrome with breast cancer. J. Exp. Med. 178:
2219–2223,

[164625]1993.

[164626]

[164627]

[164628]16402.

[164629]Jenkins, N. A.; Gilbert, D. J.; Yamamoto, R.; Kilimann, M.
W.;

[164630]Copeland, N. G.: Amphiphysin (Amph) maps to the proximal region of

[164631]mouse chromosome 13. Genomics 28: 363–365, 1995.

[164632]

[164633]

[164634]16403.

[164635]Lichte, B.; Veh, R. W.; Meyer, H. E.; Kilimann, M. W.: Amphiphysin,

[164636]a novel protein associated with synaptic vesicles. EMBO J. 11: 2521–2530,

[164637]1992.

[164638]

[164639]

[164640]16404.

[164641]Yamamoto, R.; Li, X.; Francke, U.; Kilimann, M. W.: Genetic mapping

[164642]of the human amphiphysin gene (AMPH) at 7p14–p13 excludes its involvement

[164643]in retinitis pigmentosa 9 or dominant cystoid macular dystrophy. (Letter) Am.

[164644]J. Hum. Genet. 57: 970–972, 1995.

[164645]

[164646]

[164647]16405.

[164648]Yamamoto, R.; Li, X.; Winter, S.; Francke, U.; Kilimann, M.
W.

[164649]: Primary structure of human amphiphysin, the dominant
autoantigen

[164650]of paraneoplastic Stiff–Man syndrome, and mapping of its
gene (AMPH)

[164651]to chromosome 7p13–p14. Hum. Molec. Genet. 4:
265–268, 1995.

[164652]

[164653]

[164654]16406.

[164655]Kingsmore, S. F.; Suh, D.; Seldin, M. F.: Genetic mapping
of the

[164656]glycine receptor alpha–3 subunit on mouse chromosome
8. Mammalian

[164657]Genome 5: 831–832, 1994.

[164658]

[164659]

[164660]16407.

[164661]Kuhse, J.; Schmieden, V.; Betz, H.: Identification and functional

[164662]expression of a novel ligand binding subunit of the inhibitory glycine

[164663]receptor. J. Biol. Chem. 265: 22317–22320, 1990.

[164664]

[164665]

[164666]16408.

[164667]Nikolic, Z.; Laube, B.; Weber, R. G.; Lichter, P.; Kioschis, P.;

[164668]Poustka, A.; Mulhardt, C.; Becker, C.–M.: The human glycine receptor

[164669]subunit alpha-3: GLRA3 gene structure, chromosomal localization, and

[164670]functional characterization of alternative transcripts. J. Biol.

[164671]Chem. 273: 19708–19714, 1998.

[164672]

[164673]

[164674]16409.

[164675]Birkenmeier, E. H.; Rowe, L. B.; Crossman, M. W.; Gordon, J. I.

[164676]: Ileal lipid-binding protein (IILBP) gene maps to mouse chromosome

[164677]11. Mammalian Genome 5: 805-806, 1994.

[164678]

[164679]

[164680]16410.

[164681]Oelkers, P.; Dawson, P. A.: Cloning and chromosomal localization

[164682]of the human ileal lipid-binding protein. Biochim. Biophys. Acta 1257:

[164683]199-202, 1995.

[164684]

[164685]

[164686]16411.

[164687]Johnson, D. W.; Berg, J. N.; Baldwin, M. A.; Gallione, C. J.;

Marondel,

[164688]I.; Yoon, S.-J.; Stenzel, T. T.; Speer, M.; Pericak-Vance, M. A.;

[164689]Diamond, A.; Guttmacher, A. E.; Jackson, C. E.; Attisano, L.; Kucherlapati,

[164690]R.; Porteous, M. E. M.; Marchuk, D. A.: Mutations in the activin

[164691]receptor-like kinase 1 gene in hereditary haemorrhagic telangiectasia

[164692]type 2. Nature Genet. 13: 189-195, 1996.

[164693]

[164694]

[164695]16412.

[164696]Seimiya, H.; Sawabe, T.; Inazawa, J.; Tsuruo, T.: Cloning, expression

[164697]and chromosomal localization of a novel gene for protein tyrosine

[164698]phosphatase (PTP-U2) induced by various differentiation-inducing agents. Oncogene 10:

[164699]1731-1738, 1995.

[164700]

[164701]

[164702]16413.

[164703]Wiggins, R. C.; Wiggins, J. E.; Goyal, M.; Wharram, B. L.;
Thomas,

[164704]P. E.: Molecular cloning of cDNAs encoding human
GLEPP1, a membrane

[164705]protein tyrosine phosphatase: characterization of the
GLEPP1 protein

[164706]distribution in human kidney and assignment of the
GLEPP1 gene to

[164707]human chromosome 12p12–p13. Genomics 27: 174–181,
1995.

[164708]

[164709]

[164710]16414.

[164711]Borsani, G.; Rugarli, E. I.; Taglialatela, M.; Wong, C.; Bal-
labio,

[164712]A.: Characterization of a human and murine gene (CLCN3)
sharing similarities

[164713]to voltage-gated chloride channels and to a yeast integral
membrane

[164714]protein. Genomics 27: 131–141, 1995.

[164715]

[164716]

[164717]16415.

[164718]Mills, K. A.; Mathews, K. D.; Scherpbier–Heddema, T.; Buetow, K.

[164719]H.; Baldini, A.; Ballabio, A.; Borsani, G.: Genetic and physical

[164720]mapping of a voltage–dependent chloride channel gene to human 4q32

[164721]and to mouse 8. Genomics 36: 374–376, 1996.

[164722]

[164723]

[164724]16416.

[164725]Stobrawa, S. M.; Breiderhoff, T.; Takamori, S.; Engel, D.; Schweizer,

[164726]M.; Zdebik, A. A.; Bosl, M. R.; Ruether, K.; Jahn, H.; Draguhn, A.;

[164727]Jahn, R.; Jentsch, T. J.: Disruption of ClC–3, a chloride channel

[164728]expressed on synaptic vesicles, leads to a loss of the hippocampus. Neuron 29:

[164729]185–196, 2001.

[164730]

[164731]

[164732]16417.

[164733]Taine, L.; Coupry, I.; Boisseau, P.; Saura, R.; Lacombe, D.; Arveiler,

[164734]B.: Refined localisation of the voltage-gated chloride channel, CLCN3,

[164735]to 4q33. Hum. Genet. 102: 178–181, 1998.

[164736]

[164737]

[164738]16418.

[164739]Yoshikawa, M.; Uchida, S.; Ezaki, J.; Rai, T.; Hayama, A.; Kobayashi,

[164740]K.; Kida, Y.; Noda, M.; Koike, M.; Uchiyama, Y.; Marumo, F.; Kominami,

[164741]E.; Sasaki, S.: CLC–3 deficiency leads to phenotypes similar to human

[164742]neuronal ceroid lipofuscinosis. Genes Cells 7: 597–605,
2002.

[164743]

[164744]

[164745]16419.

[164746]Pagliuca, A.; Bartoli, P. C.; Saccone, S.; Valle, G. D.; Lania,

[164747]L.: Molecular cloning of ID4, a novel dominant negative
helix–loop–helix

[164748]human gene on chromosome 6p21.3–p22. Genomics 27:
200–203, 1995.

[164749]

[164750]

[164751]16420.

[164752]Rigolet, M.; Rich, T.; Gross–Morand, M.–S.; Molina–Gomes,
D.; Viegas–Pequignot,

[164753]E.; Junien, C.: cDNA cloning, tissue distribution and chro-
mosomal

[164754]localization of the human ID4 gene. DNA Res. 5: 309–313,
1998.

[164755]

[164756]

[164757]16421.

[164758]Jia, S.; VanDusen, W. J.; Diehl, R. E.; Kohl, N. E.; Dixon, R.

[164759]A. F.; Elliston, K. O.; Stern, A. M.; Friedman, P. A.: cDNA
cloning

[164760]and expression of bovine aspartyl (asparaginy) beta-
hydroxylase. J.

[164761]Biol. Chem. 267: 14322–14327, 1992.

[164762]

[164763]

[164764]16422.

[164765]Korioth, F.; Gieffers, C.; Frey, J.: Cloning and characteriza-
tion

[164766]of the human gene encoding aspartyl beta-hydroxylase.
Gene 150:

[164767]395–399, 1994.

[164768]

[164769]

[164770]16423.

- [164771]Lavaissiere, L.; Jia, S.; Nishiyama, M.; de la Monte, S.;
Stern,
[164772]A. M.; Wands, J. R.; Friedman, P. A.: Overexpression of hu-
man aspartyl(asparaginyl)-beta-hydroxylase
[164773]in hepatocellular carcinoma and cholangiocarcinoma. J.
Clin. Invest. 98:
[164774]1313-1323, 1996.
- [164775]
- [164776]
- [164777]16424.
- [164778]Lim, K. Y.; Hong, C.-S.; Kim, D. H.: cDNA cloning and
characterization
[164779]of human cardiac junctin. Gene 255: 35-42, 2000.
- [164780]
- [164781]
- [164782]16425.
- [164783]Scott, A. F.: Personal Communication. Baltimore, Md.
2/19/2001.
- [164784]
- [164785]

[164786]16426.

[164787]Treves, S.; Feriotto, G.; Moccagatta, L.; Gambari, R.;

Zorzato,

[164788]F.: Molecular cloning, expression, functional characteriza-
tion, chromosomal

[164789]localization, and gene structure of junctate, a novel inte-
gral calcium

[164790]binding protein of sarco(endo)plasmic reticulum mem-
brane. J. Biol.

[164791]Chem. 275: 39555–39568, 2000.

[164792]

[164793]

[164794]16427.

[164795]Wetzel, G. T.; Ding, S.; Chen, F.: Molecular cloning of
junctin

[164796]from human and developing rabbit heart. Molec. Genet.
Metab. 69:

[164797]252–258, 2000.

[164798]

[164799]

[164800]16428.

[164801]Zhou, P.; Sun, L. J.; Dotsch, V.; Wagner, G.; Verdine, G. L.:

[164802]Solution structure of the core NFATC1/DNA complex. Cell
92: 687–696,

[164803]1998.

[164804]

[164805]

[164806]16429.

[164807]McCaffrey, P. G.; Luo, C.; Kerppola, T. K.; Jain, J.; Badalian,

[164808]T. M.; Ho, A. M.; Burgeon, E.; Lane, W. S.; Lambert, J. N.;
Curran,

[164809]T.; Verdine, G. L.; Rao, A.; Hogan, P. G.: Isolation of the
cyclosporin-sensitive

[164810]T cell transcription factor NFATp. Science 262: 750–754,
1993.

[164811]

[164812]

[164813]16430.

[164814]Rengarajan, J.; Tang, B.; Glimcher, L. H.: NFATc2 and

NFATc3 regulate

[164815]TH2 differentiation and modulate TCR-responsiveness of naive TH cells. Nature

[164816]Immun. 3: 48–54, 2002.

[164817]

[164818]

[164819]16431.

[164820]Chan, K.; Han, X.–D.; Kan, Y. W.: An important function of Nrf2

[164821]in combating oxidative stress: detoxification of acetaminophen. Proc.

[164822]Nat. Acad. Sci. 98: 4611–4616, 2001.

[164823]

[164824]

[164825]16432.

[164826]Baud, V.; Chisoe, S. L.; Viegas–Pequignot, E.; Diriong, S.; N'Guyen,

[164827]V. C.; Roe, B. A.; Lipinski, M.: EMR1, an unusual member in the family

[164828]of hormone receptors with seven transmembrane seg–

ments. Genomics 26:

[164829]334–344, 1995.

[164830]

[164831]

[164832]16433.

[164833]McKnight, A. J.; Macfarlane, A. J.; Seldin, M. F.; Gordon, S.:

[164834]Chromosome mapping of the Emr1 gene. Mammalian
Genome 8: 946–947,

[164835]1997.

[164836]

[164837]

[164838]16434.

[164839]Atanasoski, S.; Toldo, S. S.; Malipiero, U.; Schreiber, E.;
Fries,

[164840]R.; Fontana, A.: Isolation of the human genomic brain–
2/N–Oct 3 gene

[164841](POUF3) and assignment to chromosome 6q16. Genomics
26: 272–280,

[164842]1995.

[164843]

[164844]

[164845]16435.

[164846]Schreiber, E.; Tobler, A.; Malipiero, U.; Schaffner, W.;
Fontana,

[164847]A.: cDNA cloning of human N-Oct 3, a nervous-system
specific POU

[164848]domain transcription factor binding to the octamer DNA
motif. Nucleic

[164849]Acids Res. 21: 253–258, 1993.

[164850]

[164851]

[164852]16436.

[164853]Xia, Y.–R.; Andersen, B.; Mehrabian, M.; Diep, A. T.; War-
den, C.

[164854]H.; Mohandas, T.; McEvilly, R. J.; Rosenfeld, M. G.; Lusi, A.
J.

[164855]: Chromosomal organization of mammalian POU domain
factors. Genomics 18:

[164856]126–130, 1993.

[164857]

[164858]

[164859]16437.

[164860]Brass, N.; Heckel, D.; Sahin, U.; Pfreundschuh, M.;
Sybrecht, G.

[164861]W.; Meese, E.: Translation initiation factor eIF-4gamma is
encoded

[164862]by an amplified gene and induces an immune response in
squamous cell

[164863]lung carcinoma. Hum. Molec. Genet. 6: 33-39, 1997.

[164864]

[164865]

[164866]16438.

[164867]Gradi, A.; Imataka, H.; Svitkin, Y. V.; Rom, E.; Raught, B.;
Morino,

[164868]S.; Sonenberg, N.: A novel functional human eukaryotic
translation

[164869]initiation factor 4G. Molec. Cell. Biol. 18: 334-342, 1998.

[164870]

[164871]

[164872]16439.

[164873]Imataka, H.; Gradi, A.; Sonenberg, N.: A newly identified
N-terminal

[164874]amino acid sequence of human eIF4G binds
poly(A)-binding protein and

[164875]functions in poly(A)-dependent translation. EMBO J. 17:
7480-7489,

[164876]1998.

[164877]

[164878]

[164879]16440.

[164880]Imataka, H.; Sonenberg, N.: Human eukaryotic translation
initiation

[164881]factor 4G (eIF4G) possesses two separate and independent
binding sites

[164882]for eIF4A. Molec. Cell. Biol. 17: 6940-6947, 1997.

[164883]

[164884]

[164885]16441.

[164886]Yan, R.; Rhoads, R. E.: Human protein synthesis initiation

factor

[164887]eIF-4-gamma is encoded by a single gene (EIF4G) that maps to chromosome

[164888]3q27-qter. Genomics 26: 394-398, 1995.

[164889]

[164890]

[164891]16442.

[164892]Yan, R.; Rychlik, W.; Etchison, D.; Rhoads, R. E.: Amino acid

[164893]sequence of the human protein synthesis initiation factor eIF-4-gamma. J.

[164894]Biol. Chem. 267: 23226-23231, 1992.

[164895]

[164896]

[164897]16443.

[164898]Corrigall, A. V.; Hift, R. J.; Hancock, V.; Meissner, D.; Davids,

[164899]L.; Kirsch, R. E.; Meissner, P. N.: Identification and characterisation

[164900]of a deletion (537delAT) in the protoporphyrinogen oxi-

dase gene in

[164901]a South African variegate porphyria family. Hum. Mutat.
12: 403–407,

[164902]1998.

[164903]

[164904]

[164905]16444.

[164906]Aso, T.; Haque, D.; Fukudome, K.; Brower, C. S.; Conaway,
J. W.;

[164907]Conaway, R. C.: A human cDNA encoding the 110–kDa A
subunit of RNA

[164908]polymerase II transcription factor elongin. Gene 168:
277–278, 1996.

[164909]

[164910]

[164911]16445.

[164912]Aso, T.; Lane, W. S.; Conaway, J. W.; Conaway, R. C.: Elon-
gin

[164913](SIII): a multisubunit regulator of elongation by RNA poly-
merase II. Science 269:

[164914]1439–1443, 1995.

[164915]

[164916]

[164917]16446.

[164918]Aso, T.; Mokady, N.; Haque, D.; Conaway, R. C.; Conaway, J. W.

[164919]: Assignment of a human gene encoding the 110-kDa subunit of general

[164920]transcription factor elongin (SIII) to chromosome 1p36.1.

Genomics 30:

[164921]393–394, 1995.

[164922]

[164923]

[164924]16447.

[164925]Vikkula, M.; Boon, L. M.; Carraway, K. L., III; Calvert, J. T.;

[164926]Diamonti, A. J.; Goumnerov, B.; Pasyk, K. A.; Marchuk, D. A.; Warman,

[164927]M. L.; Cantley, L. C.; Mulliken, J. B.; Olsen, B. R.: Vascular dysmorphogenesis

[164928]caused by an activating mutation in the receptor tyrosine kinase TIE2. Cell 87:

[164929]1181–1190, 1996.

[164930]

[164931]

[164932]16448.

[164933]Iwata, T.; Kogame, K.; Toki, T.; Yokoyama, M.; Yamamoto, M.; Ito,

[164934]E.: Structure and chromosome mapping of the human small maf genes

[164935]MAFG and MAFK. Cytogenet. Cell Genet. 82: 88–90, 1998.

[164936]

[164937]

[164938]16449.

[164939]Motohashi, H.; Katsuoka, F.; Shavit, J. A.; Engel, J. D.; Yamamoto,

[164940]M.: Positive or negative MARE-dependent transcriptional regulation

[164941]is determined by the abundance of small Maf proteins.

Cell 103:

[164942]865–875, 2000.

[164943]

[164944]

[164945]16450.

[164946]Peters, L. L.; Eicher, E. M.: The ubiquitous subunit of the
globin

[164947]enhancer-binding protein NF-E2 (Nfe2u) maps to mouse
chromosome 5. Genomics 22:

[164948]490-491, 1994.

[164949]

[164950]

[164951]16451.

[164952]Shavit, J. A.; Motohashi, H.; Onodera, K.; Akasaka, J.; Ya-
mamoto,

[164953]M.; Engel, J. D.: Impaired megakaryopoiesis and behavioral
defects

[164954]in mafG-null mutant mice. Genes Dev. 12: 2164-2174,
1998.

[164955]

[164956]

[164957]16452.

[164958]Toki, T.; Itoh, J.; Kitazawa, J.; Arai, K.; Hatakeyama, K.;

Akasaka,

[164959]J.; Igarashi, K.; Nomura, N.; Yokoyama, M.; Yamamoto, M.;
Ito, E.

[164960]: Human small Maf proteins form heterodimers with CNC
family transcription

[164961]factors and recognize the NF-E2 motif. Oncogene 14:
1901–1910, 1997.

[164962]

[164963]

[164964]16453.

[164965]Yang, B.; He, B.; Abdel-Halim, S. M.; Tibell, A.; Brendel, M.
D.;

[164966]Bretzel, R. G.; Efendic, S.; Hillert, J.: Molecular cloning of a

[164967]full-length cDNA for human type 3 adenylyl cyclase and
its expression

[164968]in human islets. Biochem. Biophys. Res. Commun. 254:
548–551, 1999.

[164969]

[164970]

[164971]16454.

[164972]Arden, K. C.; Viars, C. S.; Weiss, S.; Argentin, S.; Nemer, M.

[164973]: Localization of the human B-type natriuretic peptide precursor (NPPB)

[164974]gene to chromosome 1p36. Genomics 26: 385–389, 1995.

[164975]

[164976]

[164977]16455.

[164978]Jordan, C. E.; White, R. I., Jr.; Fischer, K. C.; Neill, C.; Dorst,

[164979]J. P.: The scoliosis of congenital heart disease. Am. Heart J. 84:

[164980]463–469, 1972.

[164981]

[164982]

[164983]16456.

[164984]Ogawa, Y.; Itoh, H.; Yoshitake, Y.; Inoue, M.; Yoshimasa, T.; Serikawa,

[164985]T.; Nakao, K.: Molecular cloning and chromosomal assignment of the

[164986]mouse C-type natriuretic peptide (CNP) gene (Nppc): comparison with

[164987]the human CNP gene (NPPC). Genomics 24: 383–387,
1994.

[164988]

[164989]

[164990]16457.

[164991]Reckles, L. N.; Peterson, H. A.; Weidman, W. H.; Bianco, A.
J.,

[164992]Jr. The association of scoliosis and congenital heart dis-
ease. J.

[164993]Bone Joint Surg. 57A: 449–455, 1975.

[164994]

[164995]

[164996]16458.

[164997]Steinhelper, M. E.: Structure, expression, and genomic
mapping

[164998]of the mouse natriuretic peptide type-B gene. Circ. Res.
72: 984–992,

[164999]1993.

[165000]

[165001]

[165002]16459.

[165003]Tamura, N.; Ogawa, Y.; Chusho, H.; Nakamura, K.; Nakao, K.; Suda,

[165004]M.; Kasahara, M.; Hashimoto, R.; Katsuura, G.; Mukoyama, M.; Itoh,

[165005]H.; Saito, Y.; Tanaka, I.; Otani, H.; Katsuki, M.; Nakao, K.:
Cardiac

[165006]fibrosis in mice lacking brain natriuretic peptide. Proc.
Nat. Acad.

[165007]Sci. 97: 4239–4244, 2000.

[165008]

[165009]

[165010]16460.

[165011]Chawengsaksophak, K.; James, R.; Hammond, V. E.; Kont-
gen, F.; Beck,

[165012]F.: Homeosis and intestinal tumours in Cdx2 mutant mice.
Nature 386:

[165013]84–87, 1997.

[165014]

[165015]

[165016]16461.

[165017]Drummond, F.; Putt, W.; Fox, M.; Edwards, Y. H.: Cloning
and chromosome

[165018]assignment of the human CDX2 gene. Ann. Hum. Genet.
61: 393–400,

[165019]1997.

[165020]

[165021]

[165022]16462.

[165023]German, M. S.; Wang, J.; Fernald, A. A.; Espinosa, R., III; Le

[165024]Beau, M. M.; Bell, G. I.: Localization of the genes encoding
two

[165025]transcription factors, LMX1 and CDX3, regulating insulin
gene expression

[165026]to human chromosomes 1 and 13. Genomics 24:
403–404, 1994.

[165027]

[165028]

[165029]16463.

[165030]Woodford–Richens, K. L.; Halford, S.; Rowan, A.; Bevan, S.;

Aaltonen,

[165031]L. A.; Wasan, H.; Bicknell, D.; Bodmer, W. F.; Houlston, R.
S.; Tomlinson,

[165032]I. P. M.: CDX2 mutations do not account for juvenile poly-
posis or

[165033]Peutz–Jeghers syndrome and occur infrequently in spo-
radic colorectal

[165034]cancers. Brit. J. Cancer 84: 1314–1316, 2001.

[165035]

[165036]

[165037]16464.

[165038]Dubbink, H. J.; Verkaik, N. S.; Faber, P. W.; Trapman, J.;
Schroder,

[165039]F. H.; Romijn, J. C.: Tissue-specific and androgen-regu-
lated expression

[165040]of human prostate-specific transglutaminase. Biochem. J.
315: 901–908,

[165041]1996.

[165042]

[165043]

[165044]16465.

[165045]Gentile, V.; Grant, F. J.; Porta, R.; Baldini, A.: Localization
[165046]of the human prostate transglutaminase (type IV) gene
(TGM4) to chromosome
[165047]3p21.33–p22 by fluorescence in situ hybridization. Ge-
nomics 27:
[165048]219–220, 1995.

[165049]

[165050]

[165051]16466.

[165052]Grant, F. J.; Taylor, D. A.; Sheppard, P. O.; Mathewes, S. L.;

[165053]Lint, W.; Vanaja, E.; Bishop, P. D.; O'Hara, P. J.: Molecular
cloning

[165054]and characterization of a novel transglutaminase cDNA
from a human

[165055]prostate cDNA library. Biochem. Biophys. Res. Commun.
203: 1117–1123,

[165056]1994.

[165057]

[165058]

[165059]16467.

[165060]Barker, H. M.; Brewis, N. D.; Street, A. J.; Spurr, N. K.; Cohen,

[165061]P. T. W.: Three genes for protein phosphatase 1 map to different

[165062]human chromosomes: sequence, expression and gene localisation of protein

[165063]serine/threonine phosphatase 1 beta (PPP1CB). Biochim. Biophys. Acta 1220:

[165064]212–218, 1994.

[165065]

[165066]

[165067]16468.

[165068]Saadat, M.; Kakinoki, Y.; Mizuno, Y.; Kikuchi, K.; Yoshida, M.

[165069]C.: Chromosomal localization of human, rat, and mouse protein phosphatase

[165070]type 1 beta catalytic subunit genes (PPP1CB) by fluorescence in situ

[165071]hybridization. Jpn. J. Genet. 69: 697–700, 1994.

[165072]

[165073]

[165074]16469.

[165075]King, L. S.; Nielsen, S.; Agre, P.: Aquaporins and the respiratory

[165076]system: advice for a lung investigator. J. Clin. Invest. 105: 15–16,

[165077]2000.

[165078]

[165079]

[165080]16470.

[165081]Lee, M. D.; Bhakta, K. Y.; Raina, S.; Yonescu, R.; Griffin, C.

[165082]A.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Preston, G. M.;

[165083]Agre, P.: The human aquaporin–5 gene: molecular characterization

[165084]and chromosomal localization. J. Biol. Chem. 271: 8599–8604, 1996.

[165085]

[165086]

[165087]16471.

[165088]Ma, T.; Fukuda, N.; Song, Y.; Matthay, M. A.; Verkman, A.

S.:

[165089]Lung fluid transport in aquaporin-5 knockout mice. J.

Clin. Invest. 105:

[165090]93-100, 2000.

[165091]

[165092]

[165093]16472.

[165094]Raina, S.; Preston, G. M.; Guggino, W. B.; Agre, P.: Molecular
lar

[165095]cloning and characterization of an aquaporin cDNA from
salivary, lacrimal,

[165096]and respiratory tissues. J. Biol. Chem. 270: 1908-1912,
1995.

[165097]

[165098]

[165099]16473.

[165100]Padilla, C. A.; Bajalica, S.; Lagercrantz, J.; Holmgren, A.:

The

[165101]gene for human glutaredoxin (GLRX) is localized to human
chromosome

[165102]5q14. Genomics 32: 455–457, 1996.

[165103]

[165104]

[165105]16474.

[165106]Padilla, C. A.; Martinez–Galisteo, E.; Barcena, J. A.; Spyrou,

[165107]G.; Holmgren, A.: Purification from placenta, amino acid
sequence,

[165108]structure comparisons and cDNA cloning of human
glutaredoxin. Europ.

[165109]J. Biochem. 227: 27–34, 1995.

[165110]

[165111]

[165112]16475.

[165113]Raghavachari, N.; Krysan, K.; Xing K.; Lou, M. F.: Regula-
tion

[165114]of thioltransferase expression in human lens epithelial
cells. Invest.

[165115]Ophthalmol. Vis. Sci. 42: 1002–1008, 2001.

[165116]

[165117]

[165118]16476.

[165119]Liang, B. T.; Jacobson, K. A.: A physiological role of the
adenosine

[165120]A₃ receptor: sustained cardioprotection. Proc. Nat. Acad.
Sci. 95:

[165121]6995–6999, 1998.

[165122]

[165123]

[165124]16477.

[165125]Monitto, C. L.; Levitt, R. C.; DiSilvestre, D.; Holroyd, K. J.

[165126]: Localization of the A(3) adenosine receptor gene

(ADORA3) to human

[165127]chromosome 1p. Genomics 26: 637–638, 1995.

[165128]

[165129]

[165130]16478.

[165131]Sajjadi, F. G.; Firestein, G. S.: cDNA cloning and sequence
analysis

[165132]of the human A3 adenosine receptor. Biochim. Biophys.
Acta 1179:

[165133]105–107, 1993.

[165134]

[165135]

[165136]16479.

[165137]Zhao, Z.; Ravid, S.; Ravid, K.: Chromosomal mapping of
the mouse

[165138]A3 adenosine receptor gene, Adora3. Genomics 30:
118–119, 1995.

[165139]

[165140]

[165141]16480.

[165142]Zhou, Q.-Y.; Li, C.; Olah, M. E.; Johnson, R. A.; Stiles, G. L.;

[165143]Civelli, O.: Molecular cloning and characterization of an
adenosine

[165144]receptor: the A3 adenosine receptor. Proc. Nat. Acad. Sci.
89: 7432–7436,

[165145]1992.

[165146]

[165147]

[165148]16481.

[165149]Corset, V.; Nguyen-Ba-Charvet, K. T.; Forcet, C.; Moyse,
E.; Chedotal,

[165150]A.; Mehlen, P.: Netrin–1–mediated axon outgrowth and
cAMP production

[165151]requires interaction with adenosine A2b receptor. Nature
407: 747–750,

[165152]2000.

[165153]

[165154]

[165155]16482.

[165156]Mino, R. P.; Spoerri, P. E.; Caballero, S.; Player, D.; Belar-
dinelli,
[165157]L.; Biaggioni, I.; Grant, M. B.: Adenosine receptor antago-
nists and
[165158]retinal neovascularization in vivo. Invest. Ophthalm. Vis. Sci.
42:
[165159]3320–3324, 2001.

[165160]

[165161]

[165162]16483.

[165163]Stein, E.; Zou, Y.; Poo, M.; Tessier–Lavigne, M.: Binding of
DCC
[165164]by netrin–1 to mediate axon guidance independent of
adenosine A2B
[165165]receptor activation. Science 291: 1976–1982, 2001.

[165166]

[165167]

[165168]16484.

[165169]Townsend–Nicholson, A.; Baker, E.; Sutherland, G. R.;
Schofield,

[165170]P. R.: Localization of the adenosine A2b receptor subtype gene (ADORA2B)

[165171]to chromosome 17p11.2–p12 by FISH and PCR screening of somatic cell

[165172]hybrids. Genomics 25: 605–607, 1995.

[165173]

[165174]

[165175]16485.

[165176]von Gall, C.; Garabette, M. L.; Kell, C. A.; Frenzel, S.; Dehghani,

[165177]F.; Schumm–Draeger, P.–M.; Weaver, D. R.; Korf, H.–W.; Hastings, M.

[165178]H.; Stehle, J. H.: Rhythmic gene expression in pituitary depends

[165179]on heterologous sensitization by the neurohormone melatonin. Nature

[165180]Neurosci. 5: 234–238, 2002.

[165181]

[165182]

[165183]16486.

[165184]Baier, G.; Telford, D.; Giampa, L.; Coggeshall, K. M.; Baier–
Bitterlich,

[165185]G.; Isakov, N.; Altman, A.: Molecular cloning and charac–
terization

[165186]of PKC–theta, a novel member of the protein (PKC) gene
family expressed

[165187]predominantly in hematopoietic cells. J. Biol. Chem. 268:
4997–5004,

[165188]1993.

[165189]

[165190]

[165191]16487.

[165192]Becq, F.; Hamon, Y.; Bajetto, A.; Gola, M.; Verrier, B.; Chi–
mini,

[165193]G.: ABC1, an ATP binding cassette transporter required for
phagocytosis

[165194]of apoptotic cells, generates a regulated anion flux after
expression

[165195]in *Xenopus laevis* oocytes. J. Biol. Chem. 272: 2695–2699,
1997.

[165196]

[165197]

[165198]16488.

[165199]Decottignies, A.; Goffeau, A.: Complete inventory of the
yeast

[165200]ABC proteins. Nature Genet. 15: 137–145, 1997.

[165201]

[165202]

[165203]16489.

[165204]Guo, Z.; Inazu, A.; Yu, W.; Suzumura, T.; Okamoto, M.;
Nohara,

[165205]A.; Higashikata, T.; Sano, R.; Wakasugi, K.; Hayakawa, T.;
Yoshida,

[165206]K.; Suehiro, T.; Schmitz, G.; Mabuchi, H.: Double deletions
and missense

[165207]mutations in the first nucleotide-binding fold of the ATP-
binding

[165208]cassette transporter A1 (ABCA1) gene in Japanese patients
with Tangier

[165209]disease. J. Hum. Genet. 47: 325–329, 2002.

[165210]

[165211]

[165212]16490.

[165213]Hong, S. H.; Rhyne, J.; Zeller, K.; Miller, M.: Novel ABCA1 compound

[165214]variant associated with HDL cholesterol deficiency. Biochim. Biophys.

[165215]Acta 1587: 60–64, 2002.

[165216]

[165217]

[165218]16491.

[165219]Huang, W.; Moriyama, K.; Koga, T.; Hua, H.; Ageta, M.; Kawabata,

[165220]S.; Mawatari, K.; Imamura, T.; Eto, T.; Kawamura, M.; Teramoto, T.;

[165221]Sasaki, J.: Novel mutations in ABCA1 gene in Japanese patients with

[165222]Tangier disease and familial high density lipoprotein deficiency with

[165223]coronary heart disease. Biochim. Biophys. Acta 1537: 71–78, 2001.

[165224]

[165225]

[165226]16492.

[165227]Ishii, J.; Nagano, M.; Kujiraoka, T.; Ishihara, M.; Egashira,

[165228]T.; Takada, D.; Tsuji, M.; Hattori, H.; Emi, M.: Clinical variant
ant

[165229]of Tangier disease in Japan: mutation of the ABCA1 gene
in hypoalphalipoproteinemia

[165230]with corneal lipidosis. J. Hum. Genet. 47: 366–369, 2002.

[165231]

[165232]

[165233]16493.

[165234]Jennings, M. W.; Jones, R. W.; Wood, W. G.; Weatherall, D.
J.

[165235]: Analysis of an inversion within the human beta globin
gene cluster. Nucleic

[165236]Acids Res. 13: 2897–2906, 1985.

[165237]

[165238]

[165239]16494.

- [165240]Kulozik, A. E.; Bellan-Koch, A.; Kohne, E.; Kleihauer, E.: A
[165241]deletion/inversion rearrangement of the beta-globin gene
cluster in
[165242]a Turkish family with delta-beta(0)-thalassemia interme-
dia. Blood 79:
[165243]2455-2459, 1992.
- [165244]
- [165245]
- [165246]16495.
- [165247]Langmann, T.; Klucken, J.; Reil, M.; Liebisch, G.; Luciani,
M.-F.;
- [165248]Chimini, G.; Kaminski, W. E.; Schmitz, G.: Molecular
cloning of the
[165249]human ATP-binding cassette transporter 1 (hABC1): evi-
dence for sterol-dependent
[165250]regulation in macrophages. Biochem. Biophys. Res. Com-
mun. 257: 29-33,
[165251]1999.
- [165252]
- [165253]

[165254]16496.

[165255]Lawn, R. M.; Wade, D. P.; Garvin, M. R.; Wang, X.;

Schwartz, K.;

[165256]Porter, J. G.; Seilhamer, J. J.; Vaughan, A. M.; Oram, J. F.:

The

[165257]Tangier disease gene product ABC1 controls the cellular

apolipoprotein-mediated

[165258]lipid removal pathway. J. Clin. Invest. 104: R25-R31,

1999.

[165259]

[165260]

[165261]16497.

[165262]Luciani, M. F.; Denizot, F.; Savary, S.; Mattei, M. G.; Chi-

mini,

[165263]G.: Cloning of two novel ABC transporters mapping on

human chromosome

[165264]9. Genomics 21: 150-159, 1994.

[165265]

[165266]

[165267]16498.

[165268]Lapicka-Bodzioch, K.; Bodzioch, M.; Krull, M.; Kielar, D.;
Probst,

[165269]M.; Kiec, B.; Andrikovics, H.; Bottcher, A.; Hubacek, J.;
Aslanidis,

[165270]C.; Suttorp, N.; Schmitz, G.: Homogeneous assay based on
52 primer

[165271]sets to scan for mutations of the ABCA1 gene and its ap-
plication in

[165272]genetic analysis of a new patient with familial high-
density lipoprotein

[165273]deficiency syndrome. Biochim. Biophys. Acta 1537: 42-48,
2001.

[165274]

[165275]

[165276]16499.

[165277]Marcil, M.; Boucher, B.; Krimbou, L.; Solymoss, B. C.; Davi-
gnon,

[165278]J.; Frohlich, J.; Genest, J., Jr.: Severe familial HDL deficiency

[165279]in French-Canadian kindreds: clinical, biochemical, and
molecular

[165280]characterization. Arterioscler. Thromb. Vasc. Biol. 15:
1015-1024,

[165281]1995.

[165282]

[165283]

[165284]16500.

[165285]Marcil, M.; Yu, L.; Krimbou, L.; Boucher, B.; Oram, J. F.;

Cohn,

[165286]J. S.; Genest, J., Jr.: Cellular cholesterol transport and efflux

[165287]in fibroblasts are abnormal in subjects with familial HDL deficiency. Arterioscler.

[165288]Thromb. Vasc. Biol. 19: 159–169, 1999.

[165289]

[165290]

[165291]16501.

[165292]McNeish, J.; Aiello, R. J.; Guyot, D.; Turi, T.; Gabel, C.;

Aldinger,

[165293]C.; Hoppe, K. L.; Roach, M. L.; Royer, L. J.; de Wet, J.; Brocardo,

[165294]C.; Chimini, G.; Francone, O. L.: High density lipoprotein deficiency

[165295]and foam cell accumulation in mice with targeted disruption of ATP-binding

[165296]cassette transporter-1. Proc. Nat. Acad. Sci. 97: 4245-4250, 2000.

[165297]

[165298]

[165299]16502.

[165300]Pullinger, C. R.; Hakamata, H.; Duchateau, P. N.; Eng, C.; Aouizerat,

[165301]B. E.; Cho, M. H.; Fielding, C. J.; Kane, J. P.: Analysis of hABC1

[165302]gene 5-prime end: additional peptide sequence, promoter region, and

[165303]four polymorphisms. Biochem. Biophys. Res. Commun. 271: 451-455,

[165304]2000.

[165305]

[165306]

[165307]16503.

[165308]Petit, M. M. R.; Schoenmakers, E. F. P. M.; Huysmans, C.; Geurts,

[165309]J. M. W.; Mandahl, N.; Van de Ven, W. J. M.: LHFP, a novel
translocation

[165310]partner gene of HMGIC in a lipoma, is a member of a new
family of

[165311]LHFP-like genes. Genomics 57: 438–441, 1999.

[165312]

[165313]

[165314]16504.

[165315]Nishita, Y.; Yoshida, I.; Sado, T.; Takagi, N.: Genomic im-
printing

[165316]and chromosomal localization of the human MEST gene.
Genomics 36:

[165317]539–542, 1996.

[165318]

[165319]

[165320]16505.

[165321]Kohen, R.; Metcalf, M. A.; Khan, N.; Druck, T.; Huebner, K.;
Lachowicz,

[165322]J. E.; Meltzer, H. Y.; Sibley, D. R.; Roth, B. L.; Hamblin, M.
W.

[165323]: Cloning, characterization, and chromosomal localization
of a human

[165324]5-HT-6 serotonin receptor. J. Neurochem. 66: 47–56,
1996.

[165325]

[165326]

[165327]16506.

[165328]Frank, J.; Aita, V. M.; Ahmad, W.; Lam, H.; Wolff, C.; Chris-
tiano,

[165329]A. M.: Identification of a founder mutation in the proto-
porphyrinogen

[165330]oxidase gene in variegate porphyria patients from Chile.
Hum. Hered. 51:

[165331]160–168, 2001.

[165332]

[165333]

[165334]16507.

[165335]Frank, J.; Jugert, F. K.; Breitkopf, C.; Goerz, G.; Merk, H. F.;

[165336]Christiano, A. M.: Recurrent missense mutation in the
protoporphyrinogen

[165337]oxidase gene underlies variegate porphyria. Am. J. Med.

Genet. 79:

[165338]22–26, 1998.

[165339]

[165340]

[165341]16508.

[165342]Orso, E.; Broccardo, C.; Kaminski, W. E.; Bottcher, A.;

Liebisch,

[165343]G.; Drobnik, W.; Gotz, A.; Chambenoit, O.; Diederich, W.;

Langmann,

[165344]T.; Spruss, T.; Luciani, M.–F.; Rothe, G.; Lackner, K. J.; Chi–
mini,

[165345]G.; Schmitz, G.: Transport of lipids from Golgi to plasma
membrane

[165346]is defective in Tangier disease patients and Abc1–deficient
mice. Nature

[165347]Genet. 24: 192–196, 2000.

[165348]

[165349]

[165350]16509.

[165351]Hift, R. J.; Meissner, P. N.; Corrigall, A. V.; Ziman, M. R.;
Petersen,

[165352]L. A.; Meissner, D. M.; Davidson, B. P.; Sutherland, J.; Dai-
ley, H.

[165353]A.; Kirsch, R. E.: Variegate porphyria in South Africa,
1688–1996:

[165354]new developments in an old disease. S. Afr. Med. J. 87:
722–731,

[165355]1997.

[165356]

[165357]

[165358]16510.

[165359]Hift, R. J.; Meissner, P. N.; Todd, G.; Kirby, P.; Bilsland, D.;

[165360]Collins, P.; Ferguson, J.; Moore, M. R.: Homozygous varie-
gate porphyria:

[165361]an evolving clinical syndrome. Postgrad. Med. J. 69:
781–786, 1993.

[165362]

[165363]

[165364]16511.

[165365]Lam, H.; Dragan, L.; Tsou, H. C.; Merk, H.; Peacocke, M.;
Goerz,

[165366]G.; Sassa, S.; Poh-Fitzpatrick, M.; Bickers, D. R.; Chris-
tiano, A.

[165367]M.: Molecular basis of variegate porphyria: a de novo in-
sertion mutation

[165368]in the protoporphyrinogen oxidase gene. Hum. Genet. 99:
126-129,

[165369]1997.

[165370]

[165371]

[165372]16512.

[165373]Puy, H.; Robreau, A.-M.; Rosipal, R.; Nordmann, Y.; Dey-
bach, J.-C.

[165374]: Protoporphyrinogen oxidase: complete genomic se-
quence and polymorphisms

[165375]in the human gene. Biochem. Biophys. Res. Comm. 226:
226-230, 1996.

[165376]

[165377]

[165378]16513.

[165379]Roberts, A. G.; Puy, H.; Dailey, T. A.; Morgan, R. R.; What-
ley,

[165380]S. D.; Dailey, H. A.; Martasek, P.; Nordmann, Y.; Deybach,
J.-C.;

[165381]Elder, G. H.: Molecular characterization of homozygous
variegate

[165382]porphyria. Hum. Molec. Genet. 7: 1921–1925, 1998.

[165383]

[165384]

[165385]16514.

[165386]Honda, H.; Inazawa, J.; Nishida, J.; Yazaki, Y.; Hirai, H.:
Molecular

[165387]cloning, characterization, and chromosomal localization
of a novel

[165388]protein-tyrosine phosphatase, HPTP eta. Blood 84:
4186–4194, 1994.

[165389]

[165390]

[165391]16515.

[165392]Ostman, A.; Yang, Q.; Tonks, N. K.: Expression of DEP-1, a
receptor-like

[165393]protein-tyrosine-phosphatase, is enhanced with increasing cell density. Proc.

[165394]Nat. Acad. Sci. 91: 9680–9684, 1994.

[165395]

[165396]

[165397]16516.

[165398]Katoh, M.; Hirai, M.; Sugimura, T.; Terada, M.: Cloning and characterization

[165399]of MST, a novel (putative) serine/threonine kinase with SH3 domain. Oncogene 10:

[165400]1447–1451, 1995.

[165401]

[165402]

[165403]16517.

[165404]Morgan, H. D.; Sutherland, H. G. E.; Martin, D. I. K.; Whitelaw,

[165405]E.: Epigenetic inheritance at the agouti locus in the mouse. Nature

[165406]Genet. 23: 314–318, 1999.

[165407]

[165408]

[165409]16518.

[165410]Jacoby, A. S.; Webb, G. C.; Liu, M. L.; Kofler, B.; Hort, Y. J.;

[165411]Fathi, Z.; Bottema, C. D. K.; Shine, J.; Iismaa, T. P.: Structural
tural

[165412]organization of the mouse and human GALR1 galanin receptor genes (Galnr

[165413]and GALNR) and chromosomal localization of the mouse
gene. Genomics 45:

[165414]496–508, 1997.

[165415]

[165416]

[165417]16519.

[165418]Tomoda, T.; Kurashige, T.; Moriki, T.; Yamamoto, H.; Fujimoto,
moto,

[165419]S.; Taniguchi, T.: Enhanced expression of
poly(ADP-ribose) synthetase

[165420]gene in malignant lymphoma. Am. J. Hemat. 37: 223–227,
1991.

[165421]

[165422]

[165423]16520.

[165424]Vasquez, K. M.; Marburger, K.; Intody, Z.; Wilson, J. H.:

Manipulating

[165425]the mammalian genome by homologous recombination.

Proc. Nat. Acad.

[165426]Sci. 98: 8403–8410, 2001.

[165427]

[165428]

[165429]16521.

[165430]Yu, S.–W.; Wang, H.; Poitras, M. F.; Coombs, C.; Bowers, W.

J.;

[165431]Federoff, H. J.; Poirier, G. G.; Dawson, T. M.; Dawson, V. L.:

Mediation

[165432]of poly(ADP–ribose) polymerase–1–dependent cell death

by apoptosis–inducing

[165433]factor. Science 297: 259–263, 2002.

[165434]

[165435]

[165436]16522.

[165437]Zabel, B. U.; Herzog, H.; Schneider, R.; Auer, B.; Hirsch-Kauffmann,

[165438]M.; Schweiger, M.: Chromosomal sublocalization of the gene for human

[165439]poly (ADP-ribose) polymerase (NAD⁺ ADP-ribosyltransferase) at 1q41-42.

[165440](Abstract) Cytogenet. Cell Genet. 51: 1115, 1989.

[165441]

[165442]

[165443]16523.

[165444]Hood, L.; Kronenberg, M.; Hunkapiller, T.: T cell antigen receptors

[165445]and the immunoglobulin supergene family. Cell 40: 225-229, 1985.

[165446]

[165447]

[165448]16524.

[165449]de Villiers, J. N. P.; Hillermann, R.; Loubser, L.; Kotze, M. J.

[165450]: Spectrum of mutations in the HFE gene implicated in haemochromatosis

[165451]and porphyria. Hum. Molec. Genet. 8: 1517–1522, 1999.

[165452]

[165453]

[165454]16525.

[165455]Deybach, J.–C.; Puy, H.; Robreau, A.–M.; Lamoril, J.; Da
Silva,

[165456]V.; Grandchamp, B.; Nordmann, Y.: Mutations in the pro-
toporphyrinogen

[165457]oxidase gene in patients with variegate porphyria. Hum.
Molec. Genet. 5:

[165458]407–410, 1996.

[165459]

[165460]

[165461]16526.

[165462]Meissner, P. N.; Dailey, T. A.; Hift, R. J.; Ziman, M.; Corri-
gall,

[165463]A. V.; Roberts, A. G.; Meissner, D. M.; Kirsch, R. E.; Dailey,
H.

[165464]A.: A R59W mutation in human protoporphyrinogen oxi-
dase results in

[165465]decreased enzyme activity and is prevalent in South Africans with

[165466]variegate porphyria. *Nature Genet.* 13: 95–97, 1996.

[165467]

[165468]

[165469]16527.

[165470]Kalman, K.; Nguyen, A.; Tseng–Crank, J.; Dukes, I. D.; Chandy,

[165471]G.; Hustad, C. M.; Copeland, N. G.; Jenkins, N. A.; Mohren–weiser,

[165472]H.; Brandriff, B.; Cahalan, M.; Gutman, G. A.; Chandy, K. G.: Genomic

[165473]organization, chromosomal localization, tissue distribution, and biophysical

[165474]characterization of a novel mammalian Shaker–related voltage–gated

[165475]potassium channel, Kv1.7. *J. Biol. Chem.* 273: 5851–5857, 1998.

[165476]

[165477]

[165478]16528.

[165479]Stewart, S. L.; Rosenberg, H.; Fletcher, J. E.: Failure to
identify

[165480]the ryanodine receptor G1021A mutation in a large North
American population

[165481]with malignant hyperthermia. Clin. Genet. 54: 358–361,
1998.

[165482]

[165483]

[165484]16529.

[165485]Orkwiszewski, K. G.; Tedesco, T. A.; Mellman, W. J.; Croce,
C.

[165486]M.: Linkage relationship between the genes for thymidine
kinase and

[165487]galactokinase in different primates. Somat. Cell Genet. 2:
21–26,

[165488]1976.

[165489]

[165490]

[165491]16530.

[165492]Dunham, I.; Collins, J.; Wadey, R.; Scambler, P.: Possible
role

[165493]for COMT in psychosis associated with velo-cardio-facial syndrome.

[165494](Letter) Lancet 340: 1361-1362, 1992.

[165495]

[165496]

[165497]16531.

[165498]Pulkka, A.; Ihalainen, R.; Suorsa, A.; Riviere, M.; Szpirer, J.;

[165499]Pajunen, A.: Structures and chromosomal localizations of two rat

[165500]genes encoding S-adenosylmethionine decarboxylase.

Genomics 16:

[165501]342-349, 1993.

[165502]

[165503]

[165504]16532.

[165505]Radford, D. M.; Eddy, R.; Haley, L.; Henry, W. M.; Pegg, A. E.;

[165506]Pajunen, A.; Shows, T. B.: Gene sequences coding for S-adenosylmethionine

[165507]decarboxylase are present on human chromosome 6 and

the X and are

[165508]not amplified in colon neoplasia. Cytogenet. Cell Genet.

49: 285–288,

[165509]1989.

[165510]

[165511]

[165512]16533.

[165513]Grignani, F.; De Matteis, S.; Nervi, C.; Tomassoni, L.; Gelmetti,

[165514]V.; Cioce, M.; Fanelli, M.; Ruthardt, M.; Ferrara, F. F.; Zamir, I.;

[165515]Seiser, C.; Grignani, F.; Lazar, M. A.; Minucci, S.; Pelicci, P. G.

[165516]: Fusion proteins of the retinoic acid receptor–alpha recruit histone

[165517]deacetylase in promyelocytic leukaemia. Nature 391: 815–818, 1998.

[165518]

[165519]

[165520]16534.

[165521]Lin, R. J.; Nagy, L.; Inoue, S.; Shao, W.; Miller, W. H., Jr.;

[165522]Evans, R. M.: Role of the histone deacetylase complex in acute promyelocytic

[165523]leukaemia. Nature 391: 811–814, 1998.

[165524]

[165525]

[165526]16535.

[165527]Mendez, R.; Hake, L. E.; Andresson, T.; Littlepage, L. E.; Ruderman,

[165528]J. V.; Richter, J. D.: Phosphorylation of CPE binding factor by Eg2

[165529]regulates translation of c-mos mRNA. Nature 404: 302–307, 2000.

[165530]

[165531]

[165532]16536.

[165533]Ahrendt, S. A.; Decker, P. A.; Alawi, E. A.; Zhu, Y.; Sanchez–Cespedes,

[165534]M.; Yang, S. C.; Haasler, G. B.; Kajdacsy–Balla, A.; De–meure, M. J.;

[165535]Sidransky, D.: Cigarette smoking is strongly associated with mutation

[165536]of the K-ras gene in patients with primary adenocarcinoma of the lung. Cancer 92:

[165537]1525-1530, 2001.

[165538]

[165539]

[165540]16537.

[165541]Almoguera, C.; Shibata, D.; Forrester, K.; Martin, J.; Arnheim,

[165542]N.; Perucho, M.: Most human carcinomas of the exocrine pancreas contain

[165543]mutant c-K-ras genes. Cell 53: 549-554, 1988.

[165544]

[165545]

[165546]16538.

[165547]Andreyev, H. J. N.; Tilsed, J. V. T.; Cunningham, D.; Sampson,

[165548]S. A.; Norman, A. R.; Schneider, H. J.; Clarke, P. A.: K-ras mutations

[165549]in patients with early colorectal cancers. Gut 41: 323-329, 1997.

[165550]

[165551]

[165552]16539.

[165553]Bollag, G.; Adler, F.; elMasry, N.; McCabe, P. C.; Connor, E.,

[165554]Jr.; Thompson, P.; McCormick, F.; Shannon, K.: Biochemical characterization

[165555]of a novel KRAS insertion mutation from a human leukemia. J. Biol.

[165556]Chem. 271: 32491–32494, 1996.

[165557]

[165558]

[165559]16540.

[165560]Burmer, G. C.; Loeb, L. A.: Mutations in the KRAS2 oncogene during

[165561]progressive stages of human colon carcinoma. Proc. Nat. Acad. Sci. 86:

[165562]2403–2407, 1989.

[165563]

[165564]

[165565]16541.

[165566]Capon, D. J.; Seeburg, P. H.; McGrath, J. P.; Hayflick, J. S.;

[165567]Edman, U.; Levinson, A. D.; Goeddel, D. V.: Activation of
Ki-ras2

[165568]gene in human colon and lung carcinomas by two different
point mutations. Nature 304:

[165569]507-513, 1983.

[165570]

[165571]

[165572]16542.

[165573]Cubilla, A. L.; Fitzgerald, P. J. :Cancer Res. 36:
2690-2698,

[165574]1976.

[165575]

[165576]

[165577]16543.

[165578]Der, C. J.; Cooper, G. M.: Altered gene products are associated
with

[165579]activation of cellular ras-k genes in human lung and

colon carcinomas. Cell 32:

[165580]201–208, 1983.

[165581]

[165582]

[165583]16544.

[165584]Feig, L. A.; Bast, R. C., Jr.; Knapp, R. C.; Cooper, G. M.: So-
matic

[165585]activation of ras-K gene in a human ovarian carcinoma.

Science 223:

[165586]698–701, 1984.

[165587]

[165588]

[165589]16545.

[165590]Grimmond, S. M.; Raghavan, D.; Russell, P. J.: Detection of
a

[165591]rare point mutation in Ki-ras of a human bladder cancer
xenograft

[165592]by polymerase chain reaction and direct sequencing. Urol.

Res. 20:

[165593]121–126, 1992.

[165594]

[165595]

[165596]16546.

[165597]Hayashi, N.; Sugai, S.; Ito, I.; Nakamori, S.; Ogawa, M.;
Nakamura,

[165598]Y.: Ethnic difference in the pattern of K-ras oncogene mu-
tations

[165599]in human colorectal cancers. Hum. Mutat. 8: 258-261,
1996.

[165600]

[165601]

[165602]16547.

[165603]Tegazzin, V.; Accorsi, A.; Gritti, G.; Arcelli, L.; Di Giovanni,

[165604]A.: MH fulminant reaction after eight anaesthetics: a case
report. Minerva

[165605]Anesthesiol. 60: 217-219, 1994.

[165606]

[165607]

[165608]16548.

[165609]Tilgen, N.; Zorzato, F.; Halliger-Keller, B.; Muntoni, F.;

Sewry,

[165610]C.; Palumucci, L. M.; Schneider, C.; Hauser, E.; Lehmann-

Horn, F.;

[165611]Muller, C. R.; Treves, S.: Identification of four novel mutations

[165612]in the C-terminal membrane spanning domain of the ryanodine receptor

[165613]1: association with central core disease and alteration of calcium

[165614]homeostasis. Hum. Molec. Genet. 10: 2879-2887, 2001.

[165615]

[165616]

[165617]16549.

[165618]Tobin, J. R.; Jason, D. R.; Challa, V. R.; Nelson, T. E.; Sam-
buughin,

[165619]N.: Malignant hyperthermia and apparent heat stroke.
(Letter) JAMA 286:

[165620]168-169, 2001.

[165621]

[165622]

[165623]16550.

[165624]Wang, S.-Q.; Song, L.-S.; Lakatta, E. G.; Cheng, H.: Ca(2+) signalling

[165625]between single L-type Ca(2+) channels and ryanodine receptors in heart

[165626]cells. Nature 410: 592–596, 2001.

[165627]

[165628]

[165629]16551.

[165630]Zhang, Y.; Chen, H. S.; Khanna, V. K.; De Leon, S.; Phillips,

[165631]M. S.; Schappert, K.; Britt, B. A.; Brownell, A. K. W.;

MacLennan,

[165632]D. H.: A mutation in the human ryanodine receptor gene associated

[165633]with central core disease. Nature Genet. 5: 46–50, 1993.

[165634]

[165635]

[165636]16552.

[165637]Zorzato, F.; Fujii, J.; Otsu, K.; Phillips, M.; Green, N. M.;

[165638]Lai, F. A.; Meissner, G.; MacLennan, D. H.: Molecular cloning of

[165639]cDNA encoding human and rabbit forms of the Ca(2+) release channel

[165640](ryanodine receptor) of skeletal muscle sarcoplasmic reticulum. J.

[165641]Biol. Chem. 265: 2244–2256, 1990.

[165642]

[165643]

[165644]16553.

[165645]Marx, S. O.; Reiken, S.; Hisamatsu, Y.; Jayaraman, T.; Burkhoff,

[165646]D.; Rosemlit, N.; Marks, A. R.: PKA phosphorylation dissociates

[165647]FKBP12.6 from the calcium release channel (ryanodine receptor): defective

[165648]regulation in failing hearts. Cell 101: 365–376, 2000.

[165649]

[165650]

[165651]16554.

[165652]Mattei, M. G.; Giannini, G.; Moscatelli, F.; Sorrentino, V.:
Chromosomal

[165653]localization of murine ryanodine receptor genes RYR1,
RYR2, and RYR3

[165654]by in situ hybridization. Genomics 22: 202–204, 1994.

[165655]

[165656]

[165657]16555.

[165658]Bertocchini, F.; Ovitt, C. E.; Conti, A.; Barone, V.; Scholer,

[165659]H. R.; Bottinelli, R.; Reggiani, C.; Sorrentino, V.: Require-
ment

[165660]for the ryanodine receptor type 3 for efficient contraction
in neonatal

[165661]skeletal muscles. EMBO J. 16: 6956–6963, 1997.

[165662]

[165663]

[165664]16556.

[165665]Futatsugi, A.; Kato, K.; Ogura, H.; Li, S.–T.; Nagata, E.;
Kuwajima,

[165666]G.; Tanaka, K.; Itohara, S.; Mikoshiba, K.: Facilitation of

NMDAR-independent

[165667]LTP and spatial learning in mutant mice lacking ryanodine receptor

[165668]type 3. Neuron 24: 701–713, 1999.

[165669]

[165670]

[165671]16557.

[165672]Hakamata, Y.; Nakai, J.; Takeshima, H.; Imoto, K.: Primary structure

[165673]and distribution of a novel ryanodine receptor/calcium release channel

[165674]from rabbit brain. FEBS Lett. 312: 229–235, 1992.

[165675]

[165676]

[165677]16558.

[165678]Leeb, T.; Brenig, B.: cDNA cloning and sequencing of the human

[165679]ryanodine receptor type 3 (RYR3) reveals a novel alternative splice

[165680]site in the RYR3 gene. FEBS Lett. 423: 367–370, 1998.

[165681]

[165682]

[165683]16559.

[165684]Sorrentino, V.; Giannini, G.; Malzac, P.; Mattei, M. G.: Localization

[165685]of a novel ryanodine receptor gene (RYR3) to human chromosome 15q14-q15

[165686]by in situ hybridization. Genomics 18: 163-165, 1993.

[165687]

[165688]

[165689]16560.

[165690]Takeshima, H.; Ikemoto, T.; Nishi, M.; Nishiyama, N.; Shimuta,

[165691]M.; Sugitani, Y.; Kuno, J.; Saito, I.; Saito, H.; Endo, M.; Iino,

[165692]M.; Noda, T.: Generation and characterization of mutant mice lacking

[165693]ryanodine receptor type 3. J. Biol. Chem. 271: 19649-19652, 1996.

[165694]

[165695]

[165696]16561.

[165697]Bigler, R. D.; Bushkin, Y.; Chiorazzi, N.: S152 (CD27): a
modulating

[165698]disulfide-linked T cell activation antigen. J. Immun. 141:
21–28,

[165699]1988.

[165700]

[165701]

[165702]16562.

[165703]Hendriks, J.; Gravestien, L. A.; Tesselaar, K.; van Lier, R. A.

[165704]W.; Schumacher, T. N. M.; Borst, J.: CD27 is required for
generation

[165705]and long-term maintenance of T cell immunity. Nature
Immun. 1: 433–440,

[165706]2000.

[165707]

[165708]

[165709]16563.

[165710]Prasad, K. V. S.; Ao, Z.; Yoon, Y.; Wu, M. X.; Rizk, M.;

Jacquot,

[165711]S.; Schlossman, S. F.: CD27, a member of the tumor
necrosis factor

[165712]receptor family, induces apoptosis and binds to Siva, a
proapoptotic

[165713]protein. Proc. Nat. Acad. Sci. 94: 6346–6351, 1997.

[165714]

[165715]

[165716]16564.

[165717]Arnaiz–Villena, A.; Perez–Aciego, P.; Ballestin, C.; Sotelo,
T.;

[165718]Perez–Seoane, C.; Martin–Villa, J. M.; Regueiro, J. R.: Bio–
chemical

[165719]basis of a novel T lymphocyte receptor immunodeficiency
by immunohistochemistry:

[165720]a possible CD3–gamma abnormality. Lab. Invest. 64:
675–681, 1991.

[165721]

[165722]

[165723]16565.

[165724]Arnaiz–Villena, A.; Timon, M.; Corell, A.; Perez–Aciego, P.;

Martin-Villa,

[165725]J. M.; Regueiro, J. R.: Primary immunodeficiency caused by mutations

[165726]in the gene encoding the CD3-gamma subunit of the T-lymphocyte receptor. New

[165727]Eng. J. Med. 327: 529-533, 1992.

[165728]

[165729]

[165730]16566.

[165731]Das, S.; Cotter, F. E.; Gibbons, B.; Dhut, S.; Young, B. D.: CD3G

[165732]is within 200 kb of the leukemic t(4;11) translocation breakpoint. Genes

[165733]Chromosomes Cancer 3: 44-47, 1991.

[165734]

[165735]

[165736]16567.

[165737]Evans, G. A.; Lewis, K. A.; Lawless, G. M.: Molecular organization

[165738]of the human CD3 gene family on chromosome 11q23.

Immunogenetics 28:

[165739]365–373, 1988.

[165740]

[165741]

[165742]16568.

[165743]Krissansen, G. W.; Gorman, P. A.; Kozak, C. A.; Spurr, N.
K.; Sheer,

[165744]D.; Goodfellow, P. N.; Crumpton, M. J.: Chromosomal lo-
cations of

[165745]the gene coding for the CD3 (T3) gamma-subunit of the
human and mouse

[165746]CD3/T-cell antigen receptor complexes. Immunogenetics
26: 258–266,

[165747]1987.

[165748]

[165749]

[165750]16569.

[165751]Krissansen, G. W.; Owen, M. J.; Verbi, W.; Crumpton, M. J.:
Primary

[165752]structure of the T3 gamma subunit of the T3/T cell anti-

gen receptor

[165753]complex deduced from cDNA sequences: evolution of the
T3 gamma and

[165754]delta subunits. EMBO J. 5: 1799–1808, 1986.

[165755]

[165756]

[165757]16570.

[165758]Saito, H.; Koyama, T.; Georgopoulos, K.; Clevers, H.;
Haser, W.

[165759]G.; LeBien, T.; Tonegawa, S.; Terhorst, C.: Close linkage of
the

[165760]mouse and human CD3 gamma- and delta-chain genes
suggests that their

[165761]transcription is controlled by common regulatory ele-
ments. Proc.

[165762]Nat. Acad. Sci. 84: 9131–9134, 1987.

[165763]

[165764]

[165765]16571.

[165766]Tunnacliffe, A.; Buluwela, L.; Rabbitts, T. H.: Physical link-

age

[165767]of three CD3 genes on human chromosome 11. EMBO J. 6:
2953–2957,

[165768]1987.

[165769]

[165770]

[165771]16572.

[165772]Tunnacliffe, A.; McGuire, R. S.: A physical linkage group in
human

[165773]chromosome band 11q23 covering a region implicated in
leukocyte neoplasia. Genomics 8:

[165774]447–453, 1990.

[165775]

[165776]

[165777]16573.

[165778]Tunnacliffe, A.; Olsson, C.; Buluwela, L.; Rabbitts, T. H.:
Organization

[165779]of the human CD3 locus on chromosome 11. Europ. J. Im-
mun. 18: 1639–1642,

[165780]1988.

[165781]

[165782]

[165783]16574.

[165784]Yunis, J. J.; Jones, C.; Madden, M. T.; Lu, D.; Mayer, M. G.:

[165785]Gene order, amplification, and rearrangement of chromosome band 11q23

[165786]in hematologic malignancies. Genomics 5: 84–90, 1989.

[165787]

[165788]

[165789]16575.

[165790]Park, S.–H.; Bendelac, A.: CD1–restricted T–cell responses and

[165791]microbial infection. Nature 406: 788–792, 2000.

[165792]

[165793]

[165794]16576.

[165795]Beckman, E. M.; Porcelli, S. A.; Morita, C. T.; Behar, S. M.; Furlong,

[165796]S. T.; Brenner, M. B.: Recognition of a lipid antigen by

CD1-restricted

[165797]alpha-beta(+) T cells. Nature 372: 691-694, 1994.

[165798]

[165799]

[165800]16577.

[165801]Sieling, P. A.; Chatterjee, D.; Porcelli, S. A.; Prigozy, T. I.;

[165802]Mazzaccaro, R. J.; Soriano, T.; Bloom, B. R.; Brenner, M. B.;

Kronenberg,

[165803]M.; Brennan, P. J.; Modlin, R. L.: CD1-restricted T cell

recognition

[165804]of microbial lipoglycan antigens. Science 269: 227-230,
1995.

[165805]

[165806]

[165807]16578.

[165808]Sieling, P. A.; Jullien, D.; Dahlem, M.; Tedder, T. F.; Rea, T.

[165809]H.; Modlin, R. L.; Porcelli, S. A.: CD1 expression by den-
dritic cells

[165810]in human leprosy lesions: correlation with effective host
immunity. J.

[165811]Immun. 162: 1851–1858, 1999.

[165812]

[165813]

[165814]16579.

[165815]Albertson, D. G.; Fishpool, R.; Sherrington, P.; Nacheva, E.;
Milstein,

[165816]C.: Sensitive and high resolution in situ hybridization to
human

[165817]chromosomes using biotin labelled probes: assignment of
the human

[165818]thymocyte CD1 antigen genes to chromosome 1. EMBO J.
7: 2801–2805,

[165819]1988.

[165820]

[165821]

[165822]16580.

[165823]Balk, S. P.; Bleicher, P. A.; Terhorst, C.: Isolation and char-
acterization

[165824]of a cDNA and gene coding for a fourth CD1 molecule.
Proc. Nat. Acad.

[165825]Sci. 86: 252–256, 1989.

[165826]

[165827]

[165828]16581.

[165829]Calabi, F.; Milstein, C.: A novel family of human major histocompatibility

[165830]complex-related genes not mapping to chromosome 6.
Nature 323: 540–543,

[165831]1986.

[165832]

[165833]

[165834]16582.

[165835]Nishimura, K.; Taketani, S.; Inokuchi, H.: Cloning of a human

[165836]cDNA for protoporphyrinogen oxidase by complementation in vivo of

[165837]a hemG mutant of Escherichia coli. J. Biol. Chem. 270:
8076–8080,

[165838]1995.

[165839]

[165840]

[165841]16583.

[165842]Roberts, A. G.; Whatley, S. D.; Daniels, J.; Holmans, P.;

Fenton,

[165843]I.; Owen, M. J.; Thompson, P.; Long, C.; Elder, G. H.: Partial
characterization

[165844]and assignment of the gene for protoporphyrinogen oxi-
dase and variegate

[165845]porphyria to human chromosome 1q23. Hum. Molec.
Genet. 4: 2387-2390,

[165846]1995.

[165847]

[165848]

[165849]16584.

[165850]Taketani, S.; Inazawa, J.; Abe, T.; Furukawa, T.; Kohno, H.;
Tokunaga,

[165851]R.; Nishimura, K.; Inokuchi, H.: The human protopor-
phyrinogen oxidase

[165852]gene (PPOX): organization and location to chromosome 1.
Genomics 29:

[165853]698-703, 1995.

[165854]

[165855]

[165856]16585.

[165857]Warnich, L.; Kotze, M. J.; Groenewald, I. M.; Groenewald, J.
Z.;

[165858]van Brakel, M. G.; van Heerden, C. J.; de Villiers, J. N. P.;
van

[165859]de Ven, W. J. M.; Schoenmakers, E. F. P. M.; Taketani, S.;
Retief,

[165860]A. E.: Identification of three mutations and associated
haplotypes

[165861]in the protoporphyrinogen oxidase gene in South African
families with

[165862]variegate porphyria. Hum. Molec. Genet. 5: 981–984,
1996.

[165863]

[165864]

[165865]16586.

[165866]Haas, M.; Ward, D. C.; Lee, J.; Roses, A. D.; Clarke, V.;
D'Eustachio,

[165867]P.; Lau, D.; Vega–Saenz de Miera, E.; Rudy, B.: Localization

of Shaw-related

[165868]K(+) channel genes on mouse and human chromosomes.

Mammalian Genome 4:

[165869]711–715, 1993.

[165870]

[165871]

[165872]16587.

[165873]Albrecht, B.; Weber, K.; Pongs, O.: Characterization of a
voltage-activated

[165874]K-channel gene cluster on human chromosome 12p13.

Receptors Channels 3:

[165875]213–220, 1995.

[165876]

[165877]

[165878]16588.

[165879]Grupe, A.; Schroter, K. H.; Ruppertsberg, J. P.; Stocker, M.;
Drewes,

[165880]T.; Beckh, S.; Pongs, O.: Cloning and expression of a hu-
man voltage-gated

[165881]potassium channel: a novel member of the RCK potassium

channel family. EMBO

[165882]J. 9: 1749–1756, 1990.

[165883]

[165884]

[165885]16589.

[165886]Klocke, R.; Roberds, S. L.; Tamkun, M. M.; Gronemeier, M.;
Augustin,

[165887]A.; Albrecht, B.; Pongs, O.; Jockusch, H.: Chromosomal
mapping in

[165888]the mouse of eight K(+)-channel genes representing the
four Shaker-like

[165889]subfamilies Shaker, Shab, Shaw, and Shal. Genomics 18:
568–574,

[165890]1993.

[165891]

[165892]

[165893]16590.

[165894]Grissmer, S.; Ghanshani, S.; Dethlefs, B.; McPherson, J. D.;
Wasmuth,

[165895]J. J.; Gutman, G. A.; Cahalan, M. D.; Chandy, K. G.: The

Shaw-related

[165896]potassium channel gene, Kv3.1, on human chromosome 11, encodes the

[165897]type 1 K⁺ channel in T cells. J. Biol. Chem. 267: 20971–20979, 1992.

[165898]

[165899]

[165900]16591.

[165901]Ried, T.; Rudy, B.; Vega-Saenz de Miera, E.; Lau, D.; Ward, D.

[165902]C.; Sen, K.: Localization of a highly conserved human potassium channel

[165903]gene (NGK2–KV4; KCNC1) to chromosome 11p15. Genomics 15: 405–411,

[165904]1993.

[165905]

[165906]

[165907]16592.

[165908]Stubbs, L.; Rinchik, E. M.; Goldberg, E.; Rudy, B.; Handel, M.

[165909]A.; Johnson, D.: Clustering of six human 11p15 gene ho-

mologs within

[165910]a 500-kb interval of proximal mouse chromosome 7. Genomics 24: 324–332,

[165911]1994.

[165912]

[165913]

[165914]16593.

[165915]Curran, M. E.; Landes, G. M.; Keating, M. T.: Molecular cloning,

[165916]characterization, and genomic localization of a human potassium channel

[165917]gene. Genomics 12: 729–737, 1992.

[165918]

[165919]

[165920]16594.

[165921]Al-Chalabi, A.; Andersen, P. M.; Nilsson, P.; Chioza, B.; Andersson,

[165922]J. L.; Russ, C.; Shaw, C. E.; Powell, J. F.; Leigh, P. N.: Deletions

[165923]of the heavy neurofilament subunit tail in amyotrophic

lateral sclerosis. Hum.

[165924]Molec. Genet. 8: 157–164, 1999.

[165925]

[165926]

[165927]16595.

[165928]Collard, J.–F.; Cote, F.; Julien, J.–P.: Defective axonal
transport

[165929]in a transgenic mouse model of amyotrophic lateral sclerosis. Nature 375:

[165930]61–64, 1995.

[165931]

[165932]

[165933]16596.

[165934]Figlewicz, D. A.; Krizus, A.; Martinoli, M. G.; Meiningner, V.;

[165935]Dib, M.; Rouleau, G. A.; Julien, J.–P.: Variants of the heavy
neurofilament

[165936]subunit are associated with the development of amy-
otrophic lateral

[165937]sclerosis. Hum. Molec. Genet. 3: 1757–1761, 1994.

[165938]

[165939]

[165940]16597.

[165941]Lees, J. F.; Shneidman, P. S.; Skuntz, S. F.; Carden, M. J.;
Lazzarini,

[165942]R. A.: The structure and organization of the human heavy
neurofilament

[165943]subunit (NF-H) and the gene encoding it. EMBO J. 7:
1947-1955, 1988.

[165944]

[165945]

[165946]16598.

[165947]Mattei, M.-G.; Dautigny, A.; Pham-Dinh, D.; Passage, E.;
Mattei,

[165948]J.-F.; Jolles, P.: The gene encoding the large human neu-
rofilament

[165949]subunit (NF-H) maps to the q121-q131 region on human
chromosome 22. Hum.

[165950]Genet. 80: 293-295, 1988.

[165951]

[165952]

[165953]16599.

[165954]Rooke, K.; Figlewicz, D. A.; Han, F.; Rouleau, G. A.: Analysis
sis

[165955]of the KSP repeat of the neurofilament heavy subunit in
familial amyotrophic

[165956]lateral sclerosis. Neurology 46: 789–790, 1996.

[165957]

[165958]

[165959]16600.

[165960]Rouleau, G. A.; Merel, P.; Lutchman, M.; Sanson, M.; Zuc-
man, J.;

[165961]Marineau, C.; Hoang-Xuan, K.; Demczuk, S.; Desmaze, C.;
Plougastel,

[165962]B.; Pulst, S. M.; Lenoir, G.; Bijlsma, E.; Fashold, R.; Duman-
ski,

[165963]J.; de Jong, P.; Parry, D.; Eldrige, R.; Aurias, A.; Delattre,
O.;

[165964]Thomas, G.: Alteration in a new gene encoding a putative
membrane-organizing

[165965]protein causes neuro-fibromatosis type 2. Nature 363:

515–521, 1993.

[165966]

[165967]

[165968]16601.

[165969]Tomkins, J.; Usher, P.; Slade, J. Y.; Ince, P. G.; Curtis, A.;

[165970]Bushby, K.; Shaw, P. J.: Novel insertion in the KSP region of
the

[165971]neurofilament heavy gene in amyotrophic lateral sclerosis.

Neuroreport 9:

[165972]3967–3970, 1998.

[165973]

[165974]

[165975]16602.

[165976]Vechio, J. D.; Bruijn, L. I.; Xu, Z.; Brown, R. H., Jr.; Cleve-
land,

[165977]D. W.: Sequence variants in human neurofilament pro-
teins: absence

[165978]of linkage to familial amyotrophic lateral sclerosis. Ann.
Neurol. 40:

[165979]603–610, 1996.

[165980]

[165981]

[165982]16603.

[165983]Watson, C. J.; Gaunt, L.; Evans, G.; Patel, K.; Harris, R.;
Strachan,

[165984]T.: A disease-associated germline deletion maps the type
2 neurofibromatosis

[165985](NF2) gene between the Ewing sarcoma region and the
leukaemia inhibitory

[165986]factor locus. Hum. Molec. Genet. 2: 701–704, 1993.

[165987]

[165988]

[165989]16604.

[165990]Levy, E.; Liem, R. K. H.; D'Eustachio, P.; Cowan, N. J.:
Structure

[165991]and evolutionary origin of the gene encoding NF-M, the
middle-molecular-mass

[165992]neurofilament protein. Europ. J. Biochem. 166: 71–77,
1987.

[165993]

[165994]

[165995]16605.

[165996]Mersiyanova, I. V.; Perepelov, A. V.; Polyakov, A. V.; Sitnikov,

[165997]V. F.; Dadali, E. L.; Oparin, R. B.; Petrin, A. N.; Evgrafov, O. V.

[165998]: A new variant of Charcot–Marie–Tooth disease type 2 is probably

[165999]the result of a mutation in the neurofilament–light gene. Am. J.

[166000]Hum. Genet. 67: 37–46, 2000.

[166001]

[166002]

[166003]16606.

[166004]Myers, M. W.; Lazzarini, R. A.; Lee, V. M.–Y.; Schlaepfer, W. W.;

[166005]Nelson, D. L.: The human mid–size neurofilament subunit: a repeated

[166006]protein sequence and the relationship of its gene to the intermediate

[166007]filament gene family. EMBO J. 6: 1617–1626, 1987.

[166008]

[166009]

[166010]16607.

[166011]Barker, P. E.; Besmer, P.; Ruddle, F. H.: Human c-kit oncogene

[166012]on human chromosome 4. (Abstract) Am. J. Hum. Genet. 37: A143, 1985.

[166013]

[166014]

[166015]16608.

[166016]Beghini, A.; Larizza, L.; Cairoli, R.; Morra, E.: c-kit activating

[166017]mutations and mast cell proliferation in human leukemia. (Letter) Blood 92:

[166018]701–702, 1998.

[166019]

[166020]

[166021]16609.

[166022]Beghini, A.; Tibiletti, M.; Roversi, G.; Chiaravalli, A.; Serio,

[166023]G.; Capella, C.; Larizza, L.: Germline mutation in the jux-
tamembrane

[166024]domain of the KIT gene in a family with gastrointestinal
stromal tumors

[166025]and urticaria pigmentosa. Cancer 92: 657–662, 2001.

[166026]

[166027]

[166028]16610.

[166029]Blume-Jensen, P.; Jiang, G.; Hyman, R.; Lee, K.-F.;
O'Gorman, S.;

[166030]Hunter, T.: Kit/stem cell factor receptor-induced activa-
tion of phosphatidylinositol

[166031]3-prime-kinase is essential for male fertility. Nature
Genet. 24:

[166032]157–162, 2000.

[166033]

[166034]

[166035]16611.

[166036]Bologna, J. L.; Pawelek, J. M.: Biology of hypopigmenta-
tion. J.

[166037]Am. Acad. Derm. 19: 217–255, 1988.

[166038]

[166039]

[166040]16612.

[166041]Brannan, C. I.; Lyman, S. D.; Williams, D. E.; Eisenman, J.;
Anderson,

[166042]D. M.; Cosman, D.; Bedell, M. A.; Jenkins, N. A.; Copeland,
N. G.

[166043]: Steel–Dickie mutation encodes a c–kit ligand lacking
transmembrane

[166044]and cytoplasmic domains. Proc. Nat. Acad. Sci. 88:
4671–4674, 1991.

[166045]

[166046]

[166047]16613.

[166048]Chabot, B.; Stephenson, D. A.; Chapman, V. M.; Besmer,
P.; Bernstein,

[166049]A.: The proto–oncogene c–kit encoding a transmembrane
tyrosine kinase

[166050]receptor maps to the mouse W locus. Nature 335: 88–89,
1988.

[166051]

[166052]

[166053]16614.

[166054]d'Auriol, L.; Mattei, M.-G.; Andre, C.; Galibert, F.: Local-
ization

[166055]of the human c-kit protooncogene on the q11-q12 region
of chromosome

[166056]4. Hum. Genet. 78: 374-376, 1988.

[166057]

[166058]

[166059]16615.

[166060]De Miguel, M. P.; Cheng, L.; Holland, E. C.; Federspiel, M.
J.;

[166061]Donovan, P. J.: Dissection of the c-Kit signaling pathway
in mouse

[166062]primordial germ cells by retroviral-mediated gene trans-
fer. Proc.

[166063]Nat. Acad. Sci. 99: 10458-10463, 2002.

[166064]

[166065]

[166066]16616.

[166067]Dubreuil, P.; Forrester, L.; Rottapel, R.; Reedijk, M.; Fujita,

[166068]J.; Bernstein, A.: The c-fms gene complements the mitogenic defect

[166069]in mast cells derived from mutant W mice but not mi (microphthalmia)

[166070]mice. Proc. Nat. Acad. Sci. 88: 2341–2345, 1991.

[166071]

[166072]

[166073]16617.

[166074]el-Omar, M.; Davies, J.; Gupta, S.; Ross, H.; Thompson, R.: Leiomyosarcoma

[166075]in leiomyomatosis of the small intestine. Postgrad. Med. J. 70:

[166076]661–664, 1994.

[166077]

[166078]

[166079]16618.

[166080]Fleischman, R. A.: Human piebald trait resulting from a

dominant

[166081]negative mutant allele of the c-kit membrane receptor gene. J. Clin.

[166082]Invest. 89: 1713–1717, 1992.

[166083]

[166084]

[166085]16619.

[166086]Fleischman, R. A.; Saltman, D. L.; Stastny, V.; Zneimer, S.:

[166087]Deletion of the c-kit protooncogene in the human developmental defect

[166088]piebald trait. Proc. Nat. Acad. Sci. 88: 10885–10889, 1991.

[166089]

[166090]

[166091]16620.

[166092]Fritsche-Polanz, R.; Jordan, J.-H.; Feix, A.; Sperr, W. R.; Sunder-Plassmann,

[166093]G.; Valent, P.; Fodinger, M.: Mutation analysis of C-KIT in patients

[166094]with myelodysplastic syndromes without mastocytosis and

cases of systemic

[166095]mastocytosis. Brit. J. Haemat. 113: 357–364, 2001.

[166096]

[166097]

[166098]16621.

[166099]Geissler, E. N.; Ryan, M. A.; Housman, D. E.: The dominant-white
spotting (W) locus of the mouse encodes the c-kit proto-

[166100]oncogene. Cell 55:
185–192, 1988.

[166101]185–192, 1988.

[166102]

[166103]

[166104]16622.

[166105]Giebel, L. B.; Spritz, R. A.: Mutation of the KIT (mast/stem

[166106]cell growth factor receptor) protooncogene in human
piebaldism. Proc.

[166107]Nat. Acad. Sci. 88: 8696–8699, 1991.

[166108]

[166109]

[166110]16623.

[166111]Hirota, S.; Isozaki, K.; Moriyama, Y.; Hashimoto, K.;
Nishida,

[166112]T.; Ishiguro, S.; Kawano, K.; Hanada, M.; Kurata, A.;
Takeda, M.;

[166113]Tunio, G. M.; Matsuzawa, Y.; Kanakura, Y.; Shinomura, Y.;
Kitamura,

[166114]Y.: Gain-of-function mutations of c-kit in human gas-
trointestinal

[166115]stromal tumors. Science 279: 577–580, 1998.

[166116]

[166117]

[166118]16624.

[166119]Huizinga, J. D.; Thuneberg, L.; Kluppel, M.; Malysz, J.;
Mikkelsen,

[166120]H. B.; Bernstein, A.: W/kit gene required for interstitial
cells

[166121]of Cajal and for intestinal pacemaker activity. Nature 373:
347–349,

[166122]1995.

[166123]

[166124]

[166125]16625.

[166126]Furitsu, T.; Tsujimura, T.; Tono, T.; Ikeda, H.; Kitayama,
H.;

[166127]Koshimizu, U.; Sugahara, H.; Butterfield, J. H.; Ashman, L.
K.; Kanayama,

[166128]Y.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Identifica-
tion of

[166129]mutations in the coding sequence of the proto-oncogene

c-kit in a

[166130]human mast cell leukemia cell line causing ligand-independent activation

[166131]of c-kit product. J. Clin. Invest. 92: 1736–1744, 1993.

[166132]

[166133]

[166134]16626.

[166135]Ingram, D. A.; Yang, F.-C.; Travers, J. B.; Wenning, M. J.; Hiatt,

[166136]K.; New, S.; Hood, A.; Shannon, K.; Williams, D. A.; Clapp, D. W.

[166137]: Genetic and biochemical evidence that haploinsufficiency of the

[166138]Nf1 tumor suppressor gene modulates melanocyte and mast cell fates

[166139]in vivo. J. Exp. Med. 191: 181–187, 2000.

[166140]

[166141]

[166142]16627.

[166143]Isozaki, K.; Terris, B.; Belghiti, J.; Schiffmann, S.; Hirota,

[166144]S.; Vanderwinden, J.-M.: Germline-activating mutation in the kinase

[166145]domain of KIT gene in familial gastrointestinal stromal tumors. Am.

[166146]J. Path. 157: 1581-1585, 2000.

[166147]

[166148]

[166149]16628.

[166150]Joensuu, H.; Roberts, P. J.; Sarlomo-Rikala, M.; Andersson, L.

[166151]C.; Tervahartiala, P.; Tuveson, D.; Silberman, S. L.; Capdeville,

[166152]R.; Dimitrijevic, S.; Druker, B.; Demetri, G. D.: Effect of the tyrosine

[166153]kinase inhibitor STI571 in a patient with a metastatic gastrointestinal

[166154]stromal tumor. New Eng. J. Med. 344: 1052-1056, 2001.

[166155]

[166156]

[166157]16629.

[166158]Johansson Moller, M.; Chaudhary, R.; Hellmen, E.; Hoy-
heim, B.;

[166159]Chowdhary, B.; Andersson, L.: Pigs with the dominant
white coat color

[166160]phenotype carry a duplication of the KIT gene encoding
the mast/stem

[166161]cell growth factor receptor. Mammalian Genome 7:
822–830, 1996.

[166162]

[166163]

[166164]16630.

[166165]Lasota, J.; Jasinski, M.; Sarlomo–Rikala, M.; Miettinen, M.:

[166166]Mutations in exon 11 of c-kit occur preferentially in ma-
lignant versus

[166167]benign gastrointestinal stromal tumors and do not occur
in leiomyomas

[166168]or leiomyosarcomas. Am. J. Path. 154: 53–60, 1999.

[166169]

[166170]

[166171]16631.

[166172]Longley, B. J.; Tyrrell, L.; Lu, S.-Z.; Ma, Y.-S.; Langley, K.;

[166173]Ding, T.; Duffy, T.; Jacobs, P.; Tang, L. H.; Modlin, I.: So-
matic

[166174]c-KIT activating mutation in urticaria pigmentosa and ag-
gressive mastocytosis:

[166175]establishment of clonality in a human mast cell neoplasm.
Nature

[166176]Genet. 12: 312-314, 1996.

[166177]

[166178]

[166179]16632.

[166180]Longley, B. J., Jr.; Metcalfe, D. D.; Tharp, M.; Wang, X.;
Tyrrell,

[166181]L.; Lu, S.-Z.; Heitjan, D.; Ma, Y.: Activating and dominant
inactivating

[166182]c-KIT catalytic domain mutations in distinct clinical forms
of human

[166183]mastocytosis. Proc. Nat. Acad. Sci. 96: 1609-1614, 1999.

[166184]

[166185]

[166186]16633.

[166187]Marklund, S.; Kijas, J.; Rodriguez–Martinez, H.;

Ronnstrand, L.;

[166188]Funa, K.; Moller, M.; Lange, D.; Edfors–Lilja, I.; Andersson,

L.:

[166189]Molecular basis for the dominant white phenotype in the
domestic pig. Genome

[166190]Res. 8: 826–833, 1998.

[166191]

[166192]

[166193]16634.

[166194]Minokoshi, Y.; Kim, Y.–B.; Peroni, O. D.; Fryer, L. G. D.;

Muller,

[166195]C.; Carling, D.; Kahn, B. B.: Leptin stimulates fatty–acid
oxidation

[166196]by activating AMP–activated protein kinase. Nature 415:
339–343,

[166197]2002.

[166198]

[166199]

[166200]16635.

[166201]Kiefer, M. C.; Tucker, J. E.; Joh, R.; Landsberg, K. E.; Saltman,

[166202]D.; Barr, P. J.: Identification of a second human subtilisin-like

[166203]protease gene in the fes/fps region of chromosome 15.

DNA Cell Biol. 10:

[166204]757–769, 1991.

[166205]

[166206]

[166207]16636.

[166208]Amiel, J.; Audollent, S.; Joly, D.; Dureau, P.; Salomon, R.;
Tellier,

[166209]A.–L.; Auge, J.; Bouissou, F.; Antignac, C.; Gubler, M.–C.;
Eccles,

[166210]M. R.; Munnich, A.; Vekemans, M.; Lyonnet, S.; Attie–
Bitach, T.:

[166211]PAX2 mutations in renal–coloboma syndrome: mutational
hotspot and

[166212]germline mosaicism. Europ. J. Hum. Genet. 8: 820–826,
2000.

[166213]

[166214]

[166215]16637.

[166216]Dehbi, M.; Ghahremani, M.; Lechner, M.; Dressler, G.; Pelletier,

[166217]J.: The paired-box transcription factor, PAX2, positively modulates

[166218]expression of the Wilms' tumor suppressor gene. *Oncogene* 13: 447-453,

[166219]1996.

[166220]

[166221]

[166222]16638.

[166223]Devriendt, K.; Matthijs, G.; Van Damme, B.; Van Caesbroeck, D.;

[166224]Eccles, M.; Vanrenterghem, Y.; Fryns, J.-P.; Leys, A.: Missense mutation

[166225]and hexanucleotide duplication in the PAX2 gene in two unrelated families

[166226]with renal-coloboma syndrome (MIM 120330). *Hum. Genet.* 103: 149-153,

[166227]1998.

[166228]

[166229]

[166230]16639.

[166231]Dressler, G. R.; Deutsch, U.; Chowdhury, K.; Nornes, H. O.;

Gruss,

[166232]P.: Pax2, a new murine paired-box-containing gene and
its expression

[166233]in the developing excretory system. Development 109:
787-795, 1990.

[166234]

[166235]

[166236]16640.

[166237]Ford, B.; Rupps, R.; Lirenman, D.; Van Allen, M. I.; Far-

quharson,

[166238]D.; Lyons, C.; Friedman, J. M.: Renal-coloboma syndrome:
prenatal

[166239]detection and clinical spectrum in a large family. Am. J.
Med. Genet. 99:

[166240]137-141, 2001.

[166241]

[166242]

[166243]16641.

[166244]Keller, S. A.; Jones, J. M.; Boyle, A.; Barrow, L. L.; Killen,

[166245]P. D.; Green, D. G.; Kapousta, N. V.; Hitchcock, P. F.;

Swank, R.

[166246]T.; Meisler, M. H.: Kidney and retinal defects (Krd), a

transgene-induced

[166247]mutation with a deletion of mouse chromosome 19 that

includes the

[166248]Pax2 locus. Genomics 23: 309–320, 1994.

[166249]

[166250]

[166251]16642.

[166252]Narahara, K.; Baker, E.; Ito, S.; Yokoyama, Y.; Yu, S.; He-

witt,

[166253]D.; Sutherland, G. R.; Eccles, M. R.; Richards, R. I.: Locali-

sation

[166254]of a 10q breakpoint within the PAX2 gene in a patient with

a de novo

[166255]t(10;13) translocation and optic nerve coloboma–renal disease. J.

[166256]Med. Genet. 34: 213–216, 1997.

[166257]

[166258]

[166259]16643.

[166260]Pilz, A. J.; Povey, S.; Gruss, P.; Abbott, C. M.: Mapping of the

[166261]human homologs of the murine paired–box–containing genes. Mammalian

[166262]Genome 4: 78–82, 1993.

[166263]

[166264]

[166265]16644.

[166266]Favor, J.; Sandulache, R.; Neuhauser–Klaus, A.; Pretsch, W.; Chatterjee,

[166267]B.; Senft, E.; Wurst, W.; Blanquet, V.; Grimes, P.; Sporle, R.; Schughart,

[166268]K.: The mouse Pax2(1Neu) mutation is identical to a human PAX2 mutation

[166269]in a family with renal–coloboma syndrome and results in
developmental

[166270]defects of the brain, ear, eye, and kidney. Proc. Nat. Acad.
Sci. 93:

[166271]13870–13875, 1996.

[166272]

[166273]

[166274]16645.

[166275]Liou, G. I.; Ma, D.–P.; Yang, Y.–W.; Geng, L.; Zhu, C.;
Baehr,

[166276]W.: Human interstitial retinoid–binding protein: gene
structure and

[166277]primary sequence. J. Biol. Chem. 264: 8200–8206, 1989.

[166278]

[166279]

[166280]16646.

[166281]Nakamura, Y.; Lathrop, M.; Bragg, T.; Leppert, M.;
O'Connell,

[166282]P.; Jones, C.; Lalouel, J. M.; White, R.: An extended genetic
linkage

[166283]map of markers for human chromosome 10. Genomics 3:
389–392, 1988.

[166284]

[166285]

[166286]16647.

[166287]Sap, J.; Munoz, A.; Damm, K.; Goldberg, Y.; Ghysdael, J.;
Leutz,

[166288]A.; Beug, H.; Vennstrom, B.: The c-erb-A protein is a
high-affinity

[166289]receptor for thyroid hormone. Nature 324: 635–640,
1986.

[166290]

[166291]

[166292]16648.

[166293]Bayliss, W.; Starling, E. H.: The mechanism of pancreatic
secretion. J.

[166294]Physiol. (London) 28: 325–353, 1902.

[166295]

[166296]

[166297]16649.

[166298]Chow, B. K.-C.: Molecular cloning and functional characterization

[166299]of a human secretin receptor. Biochem. Biophys. Res. Commun. 212:

[166300]204-211, 1995.

[166301]

[166302]

[166303]16650.

[166304]Ishihara, T.; Nakamura, S.; Kaziro, Y.; Takahashi, T.; Takahashi,

[166305]K.; Nagata, S.: Molecular cloning and expression of a cDNA encoding

[166306]the secretin receptor. EMBO J. 10: 1635-1641, 1991.

[166307]

[166308]

[166309]16651.

[166310]Mark, H. F. L.; Chow, B. K.-C.: Localization of the gene encoding

[166311]the secretin receptor, SCTR, on human chromosome 2q14.1 by fluorescence

[166312]in situ hybridization and chromosome morphometry. Ge-
nomics 29: 817–818,
[166313]1995.

[166314]

[166315]

[166316]16652.

[166317]Peel, A. L.; Rao, R. V.; Cottrell, B. A.; Hayden, M. R.;
Ellerby,

[166318]L. M.; Bredesen, D. E.: Double-stranded RNA-dependent
protein kinase,

[166319]PKR, binds preferentially to Huntington's disease (HD)
transcripts

[166320]and is activated in HD tissue. Hum. Molec. Genet. 10:
1531–1538,

[166321]2001.

[166322]

[166323]

[166324]16653.

[166325]Squire, J.; Meurs, E. F.; Chong, K. L.; McMillan, N. A. J.;
Hovanessian,

[166326]A. G.; Williams, B. R. G.: Localization of the human interferon-induced,

[166327]ds-RNA activated p68 kinase gene (PRKR) to chromosome 2p21-p22. Genomics 16:

[166328]768-770, 1993.

[166329]

[166330]

[166331]16654.

[166332]Taylor, D. R.; Shi, S. T.; Romano, P. R.; Barber, G. N.; Lal, M.

[166333]M. C.: Inhibition of the interferon-inducible protein kinase PKR

[166334]by HCV E2 protein. Science 285: 107-110, 1999.

[166335]

[166336]

[166337]16655.

[166338]Brott, B. K.; Alessandrini, A.; Largaespada, D. A.; Copeland, N.

[166339]G.; Jenkins, N. A.; Crews, C. M.; Erikson, R. L.: MEK2 is a kinase

[166340]related to MEK1 and is differentially expressed in murine tissues. Cell

[166341]Growth Differ. 4: 921–929, 1993.

[166342]

[166343]

[166344]16656.

[166345]Favata, M. F.; Horiuchi, K. Y.; Manos, E. J.; Daulerio, A. J.;

[166346]Stradley, D. A.; Feeser, W. S.; Van Dyk, D. E.; Pitts, W. J.;
Earl,

[166347]R. A.; Hobbs, F.; Copeland, R. A.; Magolda, R. L.; Scherle,
P. A.;

[166348]Trzaskos, J. M.: Identification of a novel inhibitor of mitogen-activated

[166349]protein kinase kinase. J. Biol. Chem. 273: 18623–18632,
1998.

[166350]

[166351]

[166352]16657.

[166353]Meloche, S.; Gopalbhai, K.; Beatty, B. G.; Scherer, S. W.;
Pellerin,

[166354]J.: Chromosome mapping of the human genes encoding the MAP kinase

[166355]kinase MEK1 (MAP2K1) to 15q21 and MEK2 (MAP2K2) to 7q32. Cytogenet.

[166356]Cell Genet. 88: 249–252, 2000.

[166357]

[166358]

[166359]16658.

[166360]Orth, K.; Palmer, L. E.; Bao, Z. Q.; Stewart, S.; Rudolph, A. E.;

[166361]Bliska, J. B.; Dixon, J. E.: Inhibition of the mitogen-activated

[166362]protein kinase kinase superfamily by a Yersinia effector. Science 285:

[166363]1920–1923, 1999.

[166364]

[166365]

[166366]16659.

[166367]Pleschka, S.; Wolff, T.; Ehrhardt, C.; Hobom, G.; Planz, O.; Rapp,

[166368]U. R.; Ludwig, S.: Influenza virus propagation is impaired
by inhibition

[166369]of the Raf/MEK/ERK signalling cascade. Nature Cell Biol. 3:
301–305,

[166370]2001.

[166371]

[166372]

[166373]16660.

[166374]Rampoldi, L.; Zimbello, R.; Bortoluzzi, S.; Tiso, N.; Valle,
G.;

[166375]Lanfranchi, G.; Danieli, G. A.: Chromosomal localization of
four

[166376]MAPK signaling cascade genes: MEK1, MEK3, MEK4 and
MEKK5. Cytogenet.

[166377]Cell Genet. 78: 301–303, 1997.

[166378]

[166379]

[166380]16661.

[166381]Seger, R.; Krebs, E. G.: The MAPK signaling cascade. FASEB
J. 9:

[166382]726–735, 1995.

[166383]

[166384]

[166385]16662.

[166386]Zheng, C. F.; Guan, K. L.: Cloning and characterization of
two

[166387]distinct human extracellular signal-regulated kinase acti-
vator kinases,

[166388]MEK1 and MEK2. J. Biol. Chem. 268: 11435–11439, 1993.

[166389]

[166390]

[166391]16663.

[166392]Bunnell, B.; Heath, L. S.; Adams, D. E.; Lahti, J. M.; Kidd, V.

[166393]J.: Elevated expression of a p58 protein kinase leads to
changes

[166394]in the CHO cell cycle. Proc. Nat. Acad. Sci. 87: 7467–7471,
1990.

[166395]

[166396]

[166397]16664.

[166398]Cornelis, S.; Bruynooghe, Y.; Denecker, G.; Van Huffel, S.;

Tinton,

[166399]S.; Beyaert, R.: Identification and characterization of a
novel cell

[166400]cycle-regulated internal ribosome entry site. Molec. Cell 5:
597-605,

[166401]2000.

[166402]

[166403]

[166404]16665.

[166405]Eipers, P. G.; Barnoski, B. L.; Han, J.; Carroll, A. J.; Kidd,

[166406]V. J.: Localization of the expressed human p58 protein ki-
nase chromosomal

[166407]gene to chromosome 1p36 and a highly related sequence
to chromosome

[166408]15. Genomics 11: 621-629, 1991.

[166409]

[166410]

[166411]16666.

[166412]Eipers, P. G.; Lahti, J. M.; Kidd, V. J.: Structure and expression

[166413]of the human p58(cdk-1) protein kinase chromosomal gene. Genomics 13:

[166414]613-621, 1992.

[166415]

[166416]

[166417]16667.

[166418]Gururajan, R.; Lahti, J. M.; Grenet, J.; Easton, J.; Gruber, I.;

[166419]Ambros, P. F.; Kidd, V. J.: Duplication of a genomic region containing

[166420]the Cdc2L1-2 and MMP21-22 genes on human chromosome 1p36.3 and their

[166421]linkage to D1Z2. Genome Res. 8: 929-939, 1998.

[166422]

[166423]

[166424]16668.

[166425]Lahti, J. M.; Valentine, M.; Xiang, J.; Jones, B.; Amann, J.; Grenet,

[166426]J.; Richmond, G.; Look, A. T.; Kidd, V. J.: Alterations in the

PITSLRE

[166427]protein kinase gene complex on chromosome 1p36 in
childhood neuroblastoma. Nature

[166428]Genet. 7: 370–375, 1994.

[166429]

[166430]

[166431]16669.

[166432]White, P. S.; Maris, J. M.; Beltinger, C.; Sulman, E.; Mar-
shall,

[166433]H. N.; Fujimori, M.; Kaufman, B. A.; Biegel, J. A.; Allen, C.;
Hilliard,

[166434]C.; Valentine, M. B.; Look, A. T.; Enomoto, H.; Sakiyama,
S.; Brodeur,

[166435]G. M.: A region of consistent deletion in neuroblastoma
maps within

[166436]human chromosome 1p36.2–36.3. Proc. Nat. Acad. Sci.
92: 5520–5524,

[166437]1995.

[166438]

[166439]

[166440]16670.

[166441]Xiang, J.; Lahti, J. M.; Grenet, J.; Easton, J.; Kidd, V. J.:

[166442]Molecular cloning and expression of alternatively spliced
PITSLRE

[166443]protein kinase isoforms. J. Biol. Chem. 269:
15786–15794, 1994.

[166444]

[166445]

[166446]16671.

[166447]Collu, R.; Tang, J.; Castagne, J.; Lagace, G.; Masson, N.;
Huot,

[166448]C.; Deal, C.; Delvin, E.; Faccenda, E.; Eidne, K. A.; Van
Vliet, G.

[166449]: A novel mechanism for isolated central hypothyroidism:
inactivating

[166450]mutations in the thyrotropin–releasing hormone receptor
gene. J.

[166451]Clin. Endocr. Metab. 82: 1361–1365, 1997.

[166452]

[166453]

[166454]16672.

[166455]Amin, R. S.; Wert, S. E.; Baughman, R. P.; Tomashevski, J.
F.,

[166456]Jr.; Noguee, L. M.; Brody, A. S.; Hull, W. M.; Whitsett, J. A.:
Surfactant

[166457]protein deficiency in familial interstitial lung disease. J.
Pediat. 139:

[166458]85–92, 2001.

[166459]

[166460]

[166461]16673.

[166462]Glasser, S. W.; Burhans, M. S.; Korfhagen, T. R.; Na, C.–L.;
Sly,

[166463]P. D.; Ross, G. F.; Ikegami, M.; Whitsett, J. A.: Altered sta-
bility

[166464]of pulmonary surfactant in SP–C–deficient mice. Proc. Nat.
Acad.

[166465]Sci. 98: 6366–6371, 2001.

[166466]

[166467]

[166468]16674.

[166469]Keller, D. M.; Zeng, X.; Wang, Y.; Zhang, Q. H.; Kapoor, M.;
Shu,

[166470]H.; Goodman, R.; Lozano, G.; Zhao, Y.; Lu, H.: A DNA
damage-induced

[166471]p53 serine 392 kinase complex contains CK2, hSpt16, and
SSRP1. Molec.

[166472]Cell 283–292, 2001.

[166473]

[166474]

[166475]16675.

[166476]Maki, R. G.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Srivas-
tava,

[166477]P. K.: Localization of genes for the stress-inducible tumor
rejection

[166478]antigen gp96. (Abstract) Cytogenet. Cell Genet. 58: 1978
only, 1991.

[166479]

[166480]

[166481]16676.

[166482]Maki, R. G.; Old, L. J.; Srivastava, P. K.: Human homologue
of

[166483]murine tumor rejection antigen gp96: 5-prime-regulatory
and coding

[166484]regions and relationship to stress-induced proteins. Proc.
Nat. Acad.

[166485]Sci. 87: 5658–5662, 1990.

[166486]

[166487]

[166488]16677.

[166489]Schild, H.; Rammensee, H.–G.: gp96: the immune system's
Swiss

[166490]army knife. Nature Immun. 1: 100–101, 2000.

[166491]

[166492]

[166493]16678.

[166494]Lanske, B.; Karaplis, A. C.; Lee, K.; Luz, A.; Vortkamp, A.;
Pirro,

[166495]A.; Karperien, M.; Defize, L. H. K.; Ho, C.; Mulligan, R. C.;
Abou-Samra,

[166496]A.–B.; Juppner, H.; Segre, G. V.; Kronenberg, H. M.: PTH/
PTHrP receptor

[166497]in early development and Indian hedgehog–regulated
bone growth. Science 273:

[166498]663–666, 1996.

[166499]

[166500]

[166501]16679.

[166502]Albright, F.: Case records of the Massachusetts General
Hospital

[166503](case 27461). New Eng. J. Med. 225: 789–791, 1941.

[166504]

[166505]

[166506]16680.

[166507]Bakre, M. M.; Zhu, Y.; Yin, H.; Burton, D. W.; Terkeltaub,
R.;

[166508]Deftos, L. J.; Varner, J. A.: Parathyroid hormone–related
peptide

[166509]is a naturally occurring, protein kinase A–dependent an–
giogenesis

[166510]inhibitor. Nature Med. 8: 995–1003, 2002.

[166511]

[166512]

[166513]16681.

[166514]Broadus, A. E.; Mangin, M.; Ikeda, K.; Insogna, K. L.; Weir, E.

[166515]C.; Burtis, W. J.; Stewart, A. F.: Humoral hypercalcemia of cancer:

[166516]identification of a novel parathyroid hormone-like peptide. New Eng.

[166517]J. Med. 319: 556–563, 1988.

[166518]

[166519]

[166520]16682.

[166521]Hammonds, R. G., Jr.; McKay, P.; Winslow, G. A.; Diefenbach-Jagger,

[166522]H.; Grill, V.; Glatz, J.; Rodda, C. P.; Moseley, J. M.; Wood, W. I.;

[166523]Martin, T. J.: Purification and characterization of recombinant human

[166524]parathyroid hormone-related protein. J. Biol. Chem. 264: 14806–14811,

[166525]1989.

[166526]

[166527]

[166528]16683.

[166529]Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.;
Yasuda,

[166530]T.; Banville, D.; Goltzman, D.: Gene for parathyroid hor-
mone-like

[166531]peptide is on mouse chromosome 6. Cytogenet. Cell
Genet. 53: 80-82,

[166532]1990.

[166533]

[166534]

[166535]16684.

[166536]Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.;
Yasuda,

[166537]T.; Banville, D.; Goltzman, D.: Gene for parathyroid hor-
mone-like

[166538]peptide is on mouse chromosome 6. (Abstract) Cytogenet.
Cell Genet. 51:

[166539]1003 only, 1989.

[166540]

[166541]

[166542]16685.

[166543]Kamps, M. P.; Look, A. T.; Baltimore, D.: The human
t(1:19) translocation

[166544]in pre-B ALL produces multiple nuclear E2A-Pbx1 fusion
proteins with

[166545]differing transforming potentials. Genes Dev. 5: 358–368,
1991.

[166546]

[166547]

[166548]16686.

[166549]Kim, S. K.; Selleri, L.; Lee, J. S.; Zhang, A. Y.; Gu, X.; Ja-
cobs,

[166550]Y.; Cleary, M. L.: Pbx1 inactivation disrupts pancreas de-
velopment

[166551]and in *lpl1*-deficient mice promotes diabetes mellitus.
Nature Genet. 30:

[166552]430–435, 2002.

[166553]

[166554]

[166555]16687.

[166556]Mercader, N.; Leonardo, E.; Azpiazu, N.; Serrano, A.;
Morata, G.;

[166557]Martinez-A, C.; Torres, M.: Conserved regulation of proximal
modistal

[166558]limb axis development by Meis1/Hth. Nature 402:
425-429, 1999.

[166559]

[166560]

[166561]16688.

[166562]Monica, K.; Galili, N.; Nourse, J.; Saltman, D.; Cleary, M. L.

[166563]: PBX2 and PBX3, new homeobox genes with extensive homology to the

[166564]human proto-oncogene PBX1. Molec. Cell. Biol. 11:
6149-6157, 1991.

[166565]

[166566]

[166567]16689.

[166568]Aguado, B.; Campbell, R. D.: The novel gene G17, located

in the

[166569]human major histocompatibility complex, encodes PBX2, a
homeodomain-containing

[166570]protein. Genomics 25: 650–659, 1995.

[166571]

[166572]

[166573]16690.

[166574]Katsanis, N.; Fitzgibbon, J.; Fisher, E. M. C.: Paralogy map-
ping:

[166575]identification of a region in the human MHC triplicated
onto human

[166576]chromosomes 1 and 9 allows the prediction and isolation
of novel PBX

[166577]and NOTCH loci. Genomics 35: 101–108, 1996.

[166578]

[166579]

[166580]16691.

[166581]Hourvitz, A.; Widger, A. E.; Filho, F. L. T.; Chang, R. J.;
Adashi,

[166582]E. Y.; Erickson, G. F.: Pregnancy-associated plasma pro-

tein-A gene

[166583]expression in human ovaries is restricted to healthy follicles and

[166584]corpora lutea. J. Clin. Endocr. Metab. 85: 4916–4919, 2000.

[166585]

[166586]

[166587]16692.

[166588]Lawrence, J. B.; Oxvig, C.; Overgaard, M. T.; Sottrup-Jensen, L.;

[166589]Gleich, G. J.; Hays, L. G.; Yates, J. R., III; Conover, C. A.: The

[166590]insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease

[166591]secreted by human fibroblasts is pregnancy-associated plasma protein-A. Proc.

[166592]Nat. Acad. Sci. 96: 3149–3153, 1999.

[166593]

[166594]

[166595]16693.

[166596]Silahtaroglu, A. N.; Tumer, Z.; Kristensen, T.; Sottrup-

Jensen,

[166597]L.; Tommerup, N.: Assignment of the human gene for pregnancy-associated

[166598]plasma protein A (PAPPA) to 9q33.1 by fluorescence in situ hybridization

[166599]to mitotic and meiotic chromosomes. Cytogenet. Cell Genet. 62: 214–216,

[166600]1993.

[166601]

[166602]

[166603]16694.

[166604]Smith, G. C. S.; Stenhouse, E. J.; Crossley, J. A.; Aitken, D.

[166605]A.; Cameron, A. D.; Connor, J. M.: Early pregnancy levels of pregnancy-associated

[166606]plasma protein A and the risk of intrauterine growth restriction,

[166607]premature birth, preeclampsia, and stillbirth. J. Clin. Endocr. Metab. 87:

[166608]1762–1767, 2002.

[166609]

[166610]

[166611]16695.

[166612]Smith, G. C. S.; Stenhouse, E. J.; Crossley, J. A.; Aitken, D.

[166613]A.; Cameron, A. D.; Connor, J. M.: Early-pregnancy origins
of low

[166614]birth weight. *Nature* 417: 916 only, 2002.

[166615]

[166616]

[166617]16696.

[166618]Eisenhofer, G.; Walther, M. M.; Huynh, T.-T.; Li, S.-T.;
Bornstein,

[166619]S. R.; Vortmeyer, A.; Mannelli, M.; Goldstein, D. S.; Line-
han, W.

[166620]M.; Lenders, J. W. M.; Pacak, K.: Pheochromocytomas in
von Hippel-Lindau

[166621]syndrome and multiple endocrine neoplasia type 2 display
distinct

[166622]biochemical and clinical phenotypes. *J. Clin. Endocr.*
Metab. 86:

[166623]1999-2008, 2001.

[166624]

[166625]

[166626]16697.

[166627]Glasser, S. W.; Korfhagen, T. R.; Weaver, T. E.; Clark, J. C.;

[166628]Pilot-Matias, T.; Meuth, J.; Fox, J. L.; Whitsett, J. A.: cDNA,
deduced

[166629]polypeptide structure and chromosomal assignment of
human pulmonary

[166630]surfactant proteolipid, SPL(pVal). J. Biol. Chem. 263: 9–12,
1988.

[166631]

[166632]

[166633]16698.

[166634]Zhang, S.-H.; Liu, J.; Kobayashi, R.; Tonks, N. K.: Identifi-
cation

[166635]of the cell cycle regulator VCP (p97/CDC48) as a substrate
of the

[166636]band 4.1-related protein-tyrosine phosphatase PTPH1. J.
Biol. Chem. 274:

[166637]17806–17812, 1999.

[166638]

[166639]

[166640]16699.

[166641]Lorimer, D. D.; Benya, R. V.: Cloning and quantification of
galanin-1

[166642]receptor expression by mucosal cells lining the human
gastrointestinal

[166643]tract. Biochem. Biophys. Res. Commun. 222: 379-385,
1996.

[166644]

[166645]

[166646]16700.

[166647]Nicholl, J.; Kofler, B.; Sutherland, G. R.; Shine, J.; Iismaa,

[166648]T. P.: Assignment of the gene encoding human galanin re-
ceptor (GALNR)

[166649]to 18q23 by in situ hybridization. Genomics 30: 629-630,
1995.

[166650]

[166651]

[166652]16701.

[166653]Parker, E. M.; Izzarelli, D. G.; Nowak, H. P.; Mahle, C. D.;

Iben,

[166654]L. G.; Wang, J.; Goldstein, M. E.: Cloning and characterization of

[166655]the rat GALR1 galanin receptor from Rin14B insulinoma cells. Molec.

[166656]Brain Res. 34: 179–189, 1995.

[166657]

[166658]

[166659]16702.

[166660]Simoneaux, D. K.; Leach, R. J.; O'Connell, P.: Galanin receptor

[166661]1 gene (Galnr1) is tightly linked to the myelin basic protein gene

[166662]on chromosome 18 in mouse. Mammalian Genome 8: 875 only, 1997.

[166663]

[166664]

[166665]16703.

[166666]Walli, R.; Schafer, H.; Morys-Wortmann, C.; Paetzold, G.; Nustede,

[166667]R.; Schmidt, W. E.: Identification and biochemical charac-

terization

[166668]of the human brain galanin receptor. J. Molec. Endocr. 13:
347–356,

[166669]1994.

[166670]

[166671]

[166672]16704.

[166673]Janowski, B. A.; Grogan, M. J.; Jones, S. A.; Wisely, G. B.;
Kliwer,

[166674]S. A.; Corey, E. J.; Mangelsdorf, D. J.: Structural require-
ments

[166675]of ligands for the oxysterol liver X receptors LXR-alpha
and LXR-beta. Proc.

[166676]Nat. Acad. Sci. 96: 266–271, 1999.

[166677]

[166678]

[166679]16705.

[166680]Le Beau, M. M.; Song, C.; Davis, E. M.; Hiipakka, R. A.;
Kokontis,

[166681]J. M.; Liao, S.: Assignment of the human ubiquitous recep-

tor gene

[166682](UNR) to 19q13.3 using fluorescence in situ hybridization.

Genomics 26:

[166683]166–168, 1995.

[166684]

[166685]

[166686]16706.

[166687]Repa, J. J.; Berge, K. E.; Pomajzl, C.; Richardson, J. A.;

Hobbs,

[166688]H.; Mangelsdorf, D. J.: Regulation of ATP-binding cassette

sterol

[166689]transporters ABCG5 and ABCG8 by the liver X receptors

alpha and beta. J.

[166690]Biol. Chem. 277: 18793–18800, 2002.

[166691]

[166692]

[166693]16707.

[166694]Shinar, D. M.; Endo, N.; Rutledge, S. J.; Vogel, R.; Rodan,

G.

[166695]A.; Schmidt, A.: NER, a new member of the gene family

encoding the

[166696]human steroid hormone nuclear receptor. Gene 147:
273–276, 1994.

[166697]

[166698]

[166699]16708.

[166700]Song, C.; Hiipakka, R. A.; Kokontis, J. M.; Liao, S.: Ubiqui-
tous

[166701]receptor: structures, immunocytochemical localization,
and modulation

[166702]of gene activation by receptors for retinoic acids and thy-
roid hormones. Ann.

[166703]N.Y. Acad. Sci. 761: 38–49, 1995.

[166704]

[166705]

[166706]16709.

[166707]Tangirala, R. K.; Bischoff, E. D.; Joseph, S. B.; Wagner, B.
L.;

[166708]Walczak, R.; Laffitte, B. A.; Daige, C. L.; Thomas, D.; Hey-
man, R.

[166709]A.; Mangelsdorf, D. J.; Wang, X.; Lusi, A. J.; Tontonoz, P.;

Schulman,

[166710]I. G.: Identification of macrophage liver X receptors as inhibitors

[166711]of atherosclerosis. Proc. Nat. Acad. Sci. 99: 11896–11901, 2002.

[166712]

[166713]

[166714]16710.

[166715]Futterer, A.; Kruppa, G.; Kramer, B.; Lemke, H.; Kronke, M.: Molecular

[166716]cloning and characterization of human kinectin. Molec. Biol. Cell 6:

[166717]161–170, 1995.

[166718]

[166719]

[166720]16711.

[166721]Plitz, T.; Pfeffer, K.: Intact lysosome transport and phagosome

[166722]function despite kinectin deficiency. Molec. Cell. Biol. 21: 6044–6055,

[166723]2001.

[166724]

[166725]

[166726]16712.

[166727]Print, C. G.; Leung, E.; Harrison, J. E. B.; Watson, J. D.;
Krissansen,

[166728]G. W.: Cloning of a gene encoding a human leukocyte
protein characterised

[166729]by extensive heptad repeats. Gene 144: 221–228, 1994.

[166730]

[166731]

[166732]16713.

[166733]Print, C. G.; Morris, C. M.; Spurr, N. K.; Rooke, L.; Kris-
sansen,

[166734]G. W.: The CG–1 gene, a member of the kinectin and ES/
130 family,

[166735]maps to human chromosome band 14q22. Immunogenet-
ics 43: 227–229,

[166736]1996.

[166737]

[166738]

[166739]16714.

[166740]Rao, P. N.; Yu, H.; Hodge, R.; Pettenati, M. J.; Sheetz, M. P.

[166741]: Assignment of the human kinectin gene (KTN1), encoding a kinesin-binding

[166742]protein, to chromosome 14 band q22.1 by in situ hybridization. Cytogenet.

[166743]Cell Genet. 79: 196–197, 1997.

[166744]

[166745]

[166746]16715.

[166747]Yu, H.; Nicchitta, C. V.; Kumar, J.; Becker, M.; Toyoshima, I.;

[166748]Sheetz, M. P.: Characterization of kinectin, a kinesin-binding protein:

[166749]primary sequence and N-terminal topogenic signal analysis. Molec.

[166750]Biol. Cell 6: 171–183, 1995.

[166751]

[166752]

[166753]16716.

[166754]Gordenin, D. A.; Kunkel, T. A.; Resnick, M. A.: Repeat expansion--all

[166755]in a flap? *Nature Genet.* 16: 116–118, 1997.

[166756]

[166757]

[166758]16717.

[166759]Greene, A. L.; Snipe, J. R.; Gordenin, D. A.; Resnick, M. A.:

[166760]Functional analysis of human FEN1 in *Saccharomyces cerevisiae* and

[166761]its role in genome stability. *Hum. Molec. Genet.* 8: 2263–2273, 1999.

[166762]

[166763]

[166764]16718.

[166765]Augousseau, S.; Jouk, S.; Jalbert, P.; Prieur, M.: DiGeorge syndrome

[166766]and 22q11 rearrangements. (Letter) *Hum. Genet.* 74: 206 only, 1986.

[166767]

[166768]

[166769]16719.

[166770]Van Esch, H.; Groenen, P.; Daw, S.; Poffyn, A.; Holvoet, M.;
Scambler,

[166771]P.; Fryns, J.-P.; Van de Ven, W.; Devriendt, K.: Partial DiGe-
orge

[166772]syndrome in two patients with a 10p rearrangement. Clin.
Genet. 55:

[166773]269–276, 1999.

[166774]

[166775]

[166776]16720.

[166777]Wadey, R.; Daw, S.; Taylor, C.; Atif, U.; Kamath, S.; Halford,

[166778]S.; O'Donnell, H.; Wilson, D.; Goodship, J.; Burn, J.; Scam-
bler, P.

[166779]: Isolation of a gene encoding an integral membrane pro-
tein from the

[166780]vicinity of a balanced translocation breakpoint associated
with DiGeorge

[166781]syndrome. Hum. Molec. Genet. 4: 1027–1033, 1995.

[166782]

[166783]

[166784]16721.

[166785]Onuchic, L. F.; Schrick, J. J.; Ma, J.; Hudson, T.; Guay-
Woodford,

[166786]L. M.; Zerres, K.; Woychik, R. P.; Reeders, S. T.: Sequence
analysis

[166787]of the human hTg737 gene and its polymorphic sites in
patients with

[166788]autosomal recessive polycystic kidney disease. Mam-
malian Genome 6:

[166789]805–808, 1995.

[166790]

[166791]

[166792]16722.

[166793]Schrick, J. J.; Onuchic, L. F.; Reeders, S. T.; Korenberg, J.;

[166794]Chen, X.–N.; Moyer, J. H.; Wilkinson, J. E.; Woychik, R. P.:
Characterization

[166795]of the human homologue of the mouse Tg737 candidate
polycystic kidney

[166796]disease gene. Hum. Molec. Genet. 4: 559–567, 1995.

[166797]

[166798]

[166799]16723.

[166800]Armstrong, J. A.; Bieker, J. J.; Emerson, B. M.: A SWI/
SNF-related

[166801]chromatin remodeling complex, E-RC1, is required for
tissue-specific

[166802]transcriptional regulation by EKLF in vitro. Cell 95:
93-104, 1998.

[166803]

[166804]

[166805]16724.

[166806]Bieker, J. J.: Isolation, genomic structure, and expression
of

[166807]human erythroid Kruppel-like factor (EKLF). DNA Cell Biol.
15: 347-352,

[166808]1996.

[166809]

[166810]

[166811]16725.

[166812]Nuez, B.; Michalovich, D.; Bygrave, A.; Ploemacher, R.;
Grosveld,

[166813]F.: Defective haematopoiesis in fetal liver resulting from
inactivation

[166814]of the EKLF gene. Nature 375: 316–318, 1995.

[166815]

[166816]

[166817]16726.

[166818]Perkins, A. C.; Sharpe, A. H.; Orkin, S. H.: Lethal beta-
thalassaemia

[166819]in mice lacking the erythroid CACCC–transcription factor
EKLF. Nature 375:

[166820]318–322, 1995.

[166821]

[166822]

[166823]16727.

[166824]van Ree, J. H.; Roskrow, M. A.; Becher, A. M.; McNall, R.;
Valentine,

[166825]V. A.; Jane, S. M.; Cunningham, J. M.: The human ery-
throid–specific

[166826]transcription factor EKLF localizes to chromosome

19p13.12–p13.13. Genomics 39:

[166827]393–395, 1997.

[166828]

[166829]

[166830]16728.

[166831]Wijgerde, M.; Gribnau, J.; Trimborn, T.; Nuez, B.; Philipsen, S.;

[166832]Grosveld, F.; Fraser, P.: The role of EKLF in human beta-globin gene

[166833]competition. Genes Dev. 10: 2894–2902, 1996.

[166834]

[166835]

[166836]16729.

[166837]Contractor, A.; Rogers, C.; Maron, C.; Henkemeyer, M.; Swanson,

[166838]G. T.; Heinemann, S. F.: Trans-synaptic Eph receptor-ephrin signaling

[166839]in hippocampal mossy fiber LTP. Science 296: 1864–1869, 2002.

[166840]

[166841]

[166842]16730.

[166843]Tang, X. X.; Biegel, J. A.; Nycum, L. M.; Yoshioka, A.;
Brodeur,

[166844]G. M.; Pleasure, D. E.; Ikegaki, N.: cDNA cloning, molecular
characterization,

[166845]and chromosomal localization of NET (EPHT2), a human
EPH-related receptor

[166846]protein-tyrosine kinase gene preferentially expressed in
brain. :Genomics 29:

[166847]426-437, 1995.

[166848]

[166849]

[166850]16731.

[166851]Kamp, C.; Hirschmann, P.; Voss, H.; Huellen, K.; Vogt, P.
H.:

[166852]Two long homologous retroviral sequence blocks in prox-
imal Yq11 cause

[166853]AZFa microdeletions as a result of intrachromosomal re-
combination

[166854]events. Hum. Molec. Genet. 9: 2563–2572, 2000.

[166855]

[166856]

[166857]16732.

[166858]Krausz, C.; Meyts, E. R.–D.; Frydelund–Larson, L.; Quintana–Murci,

[166859]L.; McElreavey, K.; Skakkebaek, N. E.: Double–blind Y chromosome

[166860]microdeletion analysis in men with known sperm parameters and reproductive

[166861]hormone profiles: microdeletions are specific for spermatogenic failure. J.

[166862]Clin. Endocr. Metab. 86: 2638–2642, 2001.

[166863]

[166864]

[166865]16733.

[166866]Simpson, E.; Chandler, P.; Goulmy, E.; Ma, K.; Hargreave, T. B.;

[166867]Chandley, A. C.: Loss of the 'azoospermia factor' (AZF) on Yq in

[166868]man is not associated with loss of HYA. Hum. Molec.

Genet. 2: 469–471,

[166869]1993.

[166870]

[166871]

[166872]16734.

[166873]Sun, C.; Skaletsky, H.; Rozen, S.; Gromoll, J.; Nieschlag, E.;

[166874]Oates, R.; Page, D. C.: Deletion of azoospermia factor a
(AZFa) region

[166875]of human Y chromosome caused by recombination be-
tween HERV15 proviruses. Hum.

[166876]Molec. Genet. 9: 2291–2296, 2000.

[166877]

[166878]

[166879]16735.

[166880]Krausz, C.; Quintana–Murci, L.; Barbaux, S.; Siffroi, J.–P.;
Rouba,

[166881]H.; Delafontaine, D.; Souleyreau–Therville, N.; Arvis, G.;
Antoine,

[166882]J. M.; Erdei, E.; Taar, J. P.; Tar, A.; Jeandidier, E.; Plessis,

G.;

[166883]Bourgeron, T.; Dadoune, J.-P.; Fellous, M.; McElreavey, K.:

A high

[166884]frequency of Y chromosome deletions in males with nonidiopathic infertility. J.

[166885]Clin. Endocr. Metab. 84: 3606–3612, 1999.

[166886]

[166887]

[166888]16736.

[166889]Ma, K.; Inglis, J. D.; Sharkey, A.; Bickmore, W. A.; Hill, R.

[166890]E.; Prosser, E. J.; Speed, R. M.; Thomson, E. J.; Jobling, M.; Taylor,

[166891]K.; Wolfe, J.; Cooke, H. J.; Hargreave, T. B.; Chandley, A.

C.: A

[166892]Y chromosome gene family with RNA-binding protein homology: candidates

[166893]for the azoospermia factor AZF controlling human spermatogenesis. Cell 75:

[166894]1287–1295, 1993.

[166895]

[166896]

[166897]16737.

[166898]Vergnaud, G.; Page, D. C.; Simmler, M. C.; Brown, L.;

Rouyer,

[166899]F.; Noel, B.; Botstein, D.; de la Chapelle, A.; Weissenbach,

J.:

[166900]A deletion map of the human Y chromosome based on

DNA hybridization. Am.

[166901]J. Hum. Genet. 38: 109–124, 1986.

[166902]

[166903]

[166904]16738.

[166905]Vogt, P.; Chandley, A. C.; Hargreave, T. B.; Keil, R.; Ma, K.;

[166906]Sharkey, A.: Microdeletions in interval 6 of the Y chromosome of

[166907]males with idiopathic sterility point to disruption of AZF, a human

[166908]spermatogenesis gene. Hum. Genet. 89: 491–496, 1992.

[166909]

[166910]

[166911]16739.

[166912]Vogt, P. H.; Edelman, A.; Kirsch, S.; Henegariu, O.;

Hirschmann,

[166913]P.; Kieseewetter, F.; Kohn, F. M.; Schill, W. B.; Farah, S.;

Ramos,

[166914]C.; Hartmann, M.; Hartschuh, W.; Meschede, D.; Behre, H.

M.; Castel,

[166915]A.; Nieschlag, E.; Weidner, W.; Grone, H.-J.; Jung, A.; En-

gel, W.;

[166916]Haidl, G.: Human Y chromosome azoospermia factors

(AZF) mapped to

[166917]different subregions in Yq11. Hum. Molec. Genet. 5:

933-943, 1996.

[166918]

[166919]

[166920]16740.

[166921]Agulnik, A. I.; Bishop, C. E.; Lerner, J. L.; Agulnik, S. I.;

Solovyev,

[166922]V. V.: Analysis of mutation rates in the SMCY/SMCX genes

shows that

[166923]mammalian evolution is male driven. Mammalian Genome

8: 134-138,

[166924]1997.

[166925]

[166926]

[166927]16741.

[166928]Hawkins, J. R.: Mutational analysis of SRY in XY females.
Hum.

[166929]Mutat. 2: 347–350, 1993.

[166930]

[166931]

[166932]16742.

[166933]Hawkins, J. R.; Taylor, A.; Berta, P.; Levilliers, J.; Van der

[166934]Auwera, B.; Goodfellow, P. N.: Mutational analysis of SRY:
nonsense

[166935]and missense mutations in XY sex reversal. Hum. Genet.
88: 471–474,

[166936]1992.

[166937]

[166938]

[166939]16743.

[166940]Rohen, C.; Caselitz, J.; Stern, C.; Wanschura, S.; Schoen–

makers,

[166941]E. F.; Van de Ven, W. J.; Barnitzke, S.; Bullerdiek, J.: A
hamartoma

[166942]of the breast with an aberration of 12q mapped to the
MAR region by

[166943]fluorescence in situ hybridization. Genes Chromosomes
Cancer 84:

[166944]82–84, 1995.

[166945]

[166946]

[166947]16744.

[166948]Schoenmakers, E. F. P. M.; Huysmans, C.; Van de Ven, W. J.
M.

[166949]: Allelic knockout of novel splice variants of human re-
combination

[166950]repair gene RAD51B in t(12;14) uterine leiomyomas. Can-
cer Res. 59:

[166951]19–23, 1999.

[166952]

[166953]

[166954]16745.

[166955]Zhou, X.; Benson, K. F.; Ashar, H. R.; Chada, K.: Mutation responsible

[166956]for the mouse pygmy phenotype in the developmentally regulated factor

[166957]HMGI-C. Nature 377: 771–774, 1995.

[166958]

[166959]

[166960]16746.

[166961]Agalioti, T.; Lomvardas, S.; Parekh, B.; Yie, J.; Maniatis, T.;

[166962]Thanos, D.: Ordered recruitment of chromatin modifying and general

[166963]transcription factors to the IFN- β promoter. Cell 103: 667–678,

[166964]2000.

[166965]

[166966]

[166967]16747.

[166968]Munshi, N.; Agalioti, T.; Lomvardas, S.; Merika, M.; Chen, G.;

[166969]Thanos, D.: Coordination of a transcriptional switch by

HMGI(Y) acetylation. Science 293:

[166970]1133–1136, 2001.

[166971]

[166972]

[166973]16748.

[166974]Thanos, D.; Maniatis, T.: Virus induction of human IFN-
beta gene

[166975]expression requires the assembly of an enhanceosome.
Cell 83: 1091–1100,

[166976]1995.

[166977]

[166978]

[166979]16749.

[166980]Wood, L. J.; Mukherjee, M.; Dolde, C. E.; Xu, Y.; Maher, J.
F.;

[166981]Bunton, T. E.; Williams, J. B.; Resar, L. M. S.: HMG-I/Y, a
new c-Myc

[166982]target gene and potential oncogene. Molec. Cell Biol. 20:
5490–5502,

[166983]2000.

[166984]

[166985]

[166986]16750.

[166987]Burgess, D. L.; Kohrman, D. C.; Galt, J.; Plummer, N. W.;
Jones,

[166988]J. M.; Spear, B.; Meisler, M. H.: Mutation of a new sodium
channel

[166989]gene, Scn8a, in the mouse mutant 'motor endplate dis-
ease'. Nature

[166990]Genet. 10: 461–465, 1995.

[166991]

[166992]

[166993]16751.

[166994]DeRepentigny, Y.; Cote, P. D.; Pool, M.; Bernier, G.; Girard,
S.;

[166995]Vidal, S. M.; Kothary, R.: Pathological and genetic analysis
of the

[166996]degenerating muscle (dmu) mouse: a new allele of Scn8a.
Hum. Molec.

[166997]Genet. 10: 1819–1827, 2001.

[166998]

[166999]

[167000]16752.

[167001]Kohrman, D. C.; Harris, J. B.; Meisler, M. H.: Mutation detection

[167002]in the med and med(j) alleles of the sodium channel Scn8a: unusual

[167003]splicing due to a minor class AT-AC intron. J. Biol. Chem. 271:

[167004]17576-17581, 1996.

[167005]

[167006]

[167007]16753.

[167008]Kohrman, D. C.; Plummer, N. W.; Schuster, T.; Jones, J. M.; Jang,

[167009]W.; Burgess, D. L.; Galt, J.; Spear, B. T.; Meisler, M. H.: Insertional

[167010]mutation of the motor endplate disease (med) locus on mouse chromosome

[167011]15. Genomics 26: 171-177, 1995.

[167012]

[167013]

[167014]16754.

[167015]Kohrman, D. C.; Smith, M. R.; Goldin, A. L.; Harris, J.;
Meisler,

[167016]M. H.: A missense mutation in the sodium channel Scn8a
is responsible

[167017]for cerebellar ataxia in the mouse mutant jolting. J. Neu-
rosci. 16:

[167018]5993–5999, 1996.

[167019]

[167020]

[167021]16755.

[167022]Meisler, M. H.; Sprunger, L. K.; Plummer, N. W.; Escayg, A.;
Jones,

[167023]J. M.: Ion channel mutations in mouse models of inherited
neurological

[167024]disease. Ann. Med. 29: 569–574, 1997.

[167025]

[167026]

[167027]16756.

[167028]Plummer, N. W.; Galt, J.; Jones, J. M.; Burgess, D. L.;

Sprunger,

[167029]L. K.; Kohrman, D. C.; Meisler, M. H.: Exon organization,
coding

[167030]sequence, physical mapping, and polymorphic intragenic
markers for

[167031]the human neuronal sodium channel gene SCN8A. Ge-
nomics 54: 287-296,

[167032]1998.

[167033]

[167034]

[167035]16757.

[167036]Plummer, N. W.; McBurney, M. W.; Meisler, M. H.: Alterna-
tive splicing

[167037]of the sodium channel SCN8A predicts a truncated two-
domain protein

[167038]in fetal brain and non-neuronal cells. J. Biol. Chem. 272:
24008-24015,

[167039]1997.

[167040]

[167041]

[167042]16758.

[167043]Reid, E.; Escayg, A.; Dearlove, A. M.; Lee, D. D.; Meisler, M.

[167044]H.; Rubinsztein, D. C.: The spastic paraplegia SPG10 locus:
narrowing

[167045]of critical region and exclusion of sodium channel gene
SCN8A as a

[167046]candidate. J. Med. Genet. 38: 65–67, 2001.

[167047]

[167048]

[167049]16759.

[167050]Sprunger, L. K.; Escayg, A.; Tallaksen–Greene, S.; Albin, R.
L.;

[167051]Meisler, M. H.: Dystonia associated with mutation of the
neuronal

[167052]sodium channel Scn8a and identification of the modifier
locus Scnm1

[167053]on mouse chromosome 3. Hum. Molec. Genet. 8:
471–479, 1999.

[167054]

[167055]

[167056]16760.

[167057]Chang, D. Y.; Nelson, B.; Bilyeu, T.; Hsu, K.; Darlington, G. J.;

[167058]Maraia, R. J.: A human Alu RNA-binding protein whose expression is

[167059]associated with accumulation of small cytoplasmic Alu RNA. Molec.

[167060]Cell. Biol. 14: 3949–3959, 1994.

[167061]

[167062]

[167063]16761.

[167064]Chen, M.; She, H.; Davis, E. M.; Spicer, C. M.; Kim, L.; Ren, R.;

[167065]LeBeau, M. M.; Li, W.: Identification of Nck family genes, chromosomal

[167066]localization, expression, and signaling specificity. J. Biol. Chem. 273:

[167067]25171–25178, 1998.

[167068]

[167069]

[167070]16762.

[167071]Eden, S.; Rohatgi, R.; Podtelejnikov, A. V.; Mann, M.;
Kirschner,

[167072]M. W.: Mechanism of regulation of WAVE1-induced actin
nucleation

[167073]by Rac1 and Nck. Nature 418: 790–793, 2002.

[167074]

[167075]

[167076]16763.

[167077]Casey, J. L.; Di Jeso, B.; Rao, K.; Klausner, R. D.; Harford, J.

[167078]B.: Two genetic loci participate in the regulation by iron of
the

[167079]gene for the human transferrin receptor. Proc. Nat. Acad.
Sci. 85:

[167080]1787–1791, 1988.

[167081]

[167082]

[167083]16764.

[167084]Enns, C. A.; Suomalainen, H. A.; Gebhardt, J. E.; Schroder, J.;

[167085]Sussman, H. H.: Human transferrin receptor: expression of the receptor

[167086]is assigned to chromosome 3. Proc. Nat. Acad. Sci. 79: 3241–3245,

[167087]1982.

[167088]

[167089]

[167090]16765.

[167091]Gareau, R.; Gagnon, M. G.; Thellend, C.; Chenard, C.; Audran, M.;

[167092]Chanal, J.–L.; Ayotte, C.; Brisson, G. R.: Transferrin soluble receptor:

[167093]a possible probe for detection of erythropoietin abuse by athletes. Horm.

[167094]Metab. Res. 26: 311–312, 1994.

[167095]

[167096]

[167097]16766.

[167098]Goodfellow, P. N.; Banting, G.; Sutherland, R.; Greaves, M.;
Solomon,

[167099]E.; Povey, S.: Expression of human transferrin receptor is
controlled

[167100]by a gene on chromosome 3: assignment using species
specificity of

[167101]a monoclonal antibody. Somat. Cell Genet. 8: 197–206,
1982.

[167102]

[167103]

[167104]16767.

[167105]Larrick, J. W.; Hyman, E. S.: Acquired iron–deficiency ane–
mia

[167106]caused by an antibody against the transferrin receptor.
New Eng.

[167107]J. Med. 311: 214–218, 1984.

[167108]

[167109]

[167110]16768.

[167111]Levy, J. E.; Jin, O.; Fujiwara, Y.; Kuo, F.; Andrews, N. C.:
Transferrin

[167112]receptor is necessary for development of erythrocytes and
the nervous

[167113]system. Nature Genet. 21: 396–399, 1999.

[167114]

[167115]

[167116]16769.

[167117]Miller, Y. E.; Jones, C.; Scoggin, C.; Morse, H.; Seligman, P.

[167118]: Chromosome 3q (22–ter) encodes the human transferrin
receptor. Am.

[167119]J. Hum. Genet. 35: 573–583, 1983.

[167120]

[167121]

[167122]16770.

[167123]Newman, R.; Schneider, C.; Sutherland, R.; Vodinelich, L.;
Greaves,

[167124]M.: The transferrin receptor. Trends Biochem. Sci. 7:
397–400,

[167125]1982.

[167126]

[167127]

[167128]16771.

[167129]Nikinmaa, B.; Schroder, J.: Two antigens, the transferrin
receptor

[167130]and p90 assigned to human chromosome 3, are probably
the same protein. Hereditas 107:

[167131]55–58, 1987.

[167132]

[167133]

[167134]16772.

[167135]Omary, M. B.; Trowbridge, I. S.: Biosynthesis of the human transferrin

[167136]receptor in cultured cells. J. Biol. Chem. 256: 12888–12892, 1981.

[167137]

[167138]

[167139]16773.

[167140]Kashuba, V. I.; Gizatullin, R. Z.; Protopopov, A. I.; Allikmets,

[167141]R.; Korolev, S.; Li, J.; Boldog, F.; Tory, K.; Zabarovska, V.; Marcsek,

[167142]Z.; Sumegi, J.; Klein, G.; Zabarovsky, E. R.; Kisselev, L.: Notl

[167143]linking/jumping clones of human chromosome 3: mapping of the TFRC,

[167144]RAB7 and HAUSP genes to regions rearranged in leukemia and deleted

[167145]in solid tumors. FEBS Lett. 419: 181–185, 1997.

[167146]

[167147]

[167148]16774.

[167149]Rabin, M.; McClelland, A.; Kuhn, L.; Ruddle, F. H.: Regional
[167150]localization of the human transferrin receptor gene to
3q26.2-qter. Am.

[167151]J. Hum. Genet. 37: 1112–1116, 1985.

[167152]

[167153]

[167154]16775.

[167155]Schneider, C.; Kurkinen, M.; Greaves, M.: Isolation of cDNA
clones

[167156]for the human transferrin receptor. EMBO J. 2:
2259–2263, 1983.

[167157]

[167158]

[167159]16776.

[167160]Schneider, C.; Owen, M. J.; Banville, D.; Williams, J. G.: Pri-
mary

[167161]structure of human transferrin receptor deduced from the
mRNA sequence. Nature 311:

[167162]675–678, 1984.

[167163]

[167164]

[167165]16777.

[167166]Valenzuela, C. Y.; Avendano, A.; Harb, Z.: Association between

[167167]Rh and plasma iron binding (transferrin). Hum. Genet. 87: 438–440,

[167168]1991.

[167169]

[167170]

[167171]16778.

[167172]Vodinelich, L.; Sutherland, R.; Schneider, C.; Newman, R.; Greaves,

[167173]M.: Receptor for transferrin may be a 'target' structure for natural

[167174]killer cells. Proc. Nat. Acad. Sci. 80: 835–839, 1983.

[167175]

[167176]

[167177]16779.

[167178]Hamm, C. W.; Ravkilde, J.; Gerhardt, W.; Jorgensen, P.; Pe–

heim,

[167179]E.; Ljungdahl, L.; Goldmann, B.; Katus, H. A.: The prognostic value

[167180]of serum troponin T in unstable angina. New Eng. J. Med. 327: 146–150,

[167181]1992.

[167182]

[167183]

[167184]16780.

[167185]Carmellet, P.; Ferreira, V.; Breler, G.; Pollefeyt, S.; Kleckens,

[167186]L.; Gertsenstein, M.; Fahrig, M.; Vandenhoeck, A.; Harpal, K.; Eberhardt,

[167187]C.; Declercq, C.; Pawling, J.; Moons, L.; Collen, D.; Risau, W.; Nagy,

[167188]A.: Abnormal blood vessel development and lethality in embryos lacking

[167189]a single VEGF allele. Nature 380: 435–439, 1996.

[167190]

[167191]

[167192]16781.

[167193]Martin, L. H.; Calabi, F.; Lefebvre, F.-A.; Bilisland, C. A. G.;

[167194]Milstein, C.: Structure and expression of the human thymocyte antigens

[167195]CD1a, CD1b, and CD1c. Proc. Nat. Acad. Sci. 84: 9189–9193, 1987.

[167196]

[167197]

[167198]16782.

[167199]Martin, L. H.; Calabi, F.; Milstein, C.: Isolation of CD1 genes:

[167200]a family of major histocompatibility complex-related differentiation

[167201]antigens. Proc. Nat. Acad. Sci. 83: 9154–9158, 1986.

[167202]

[167203]

[167204]16783.

[167205]Moseley, W. S.; Watson, M. L.; Kingsmore, S. F.; Seldin, M. F.

[167206]: CD1 defines conserved linkage group border between human chromosomes

[167207]1 and mouse chromosomes 1 and 3. Immunogenetics 30:
378–382, 1989.

[167208]

[167209]

[167210]16784.

[167211]Samson, F.; de Jong, P. J.; Trask, B. J.; Koza–Taylor, P.;
Speer,

[167212]M. C.; Potter, T.; Roses, A. D.; Gilbert, J. R.: Assignment of
the

[167213]human slow skeletal troponin T gene to 19q13.4 using
somatic cell

[167214]hybrids and fluorescence in situ hybridization analysis.
Genomics 13:

[167215]1374–1375, 1992.

[167216]

[167217]

[167218]16785.

[167219]Samson, F.; Gilbert, J. R.; Lee, J. E.; Potter, T. G.; Koza–
Taylor,

[167220]P.; Speer, M. C.; Bachinski, L. L.; Siciliano, M. J.; Roses, A.
D.

[167221]: Isolation and localization of a human slow troponin T gene on chromosome

[167222]19q. (Abstract) Cytogenet. Cell Genet. 58: 2025, 1991.

[167223]

[167224]

[167225]16786.

[167226]Samson, F.; Lee, J. E.; Hung, W. Y.; Potter, T. G.; Herbstreith,

[167227]M.; Roses, A. D.; Gilbert, J. R.: Isolation and localisation of a

[167228]slow troponin (TNT) gene on chromosome 19 by subtraction of a cDNA

[167229]muscle library using myotonic muscle cDNA. J. Neurosci. Res. 27:

[167230]441-451, 1990.

[167231]

[167232]

[167233]16787.

[167234]Samson, F.; Mesnard, L.; Mihovilovic, M.; Potter, T. G.; Mercadier,

[167235]J.-J.; Roses, A. D.; Gilbert, J. R.: A new human slow skeletal troponin

[167236]T (TnTs) mRNA isoform derived from alternative splicing of a single

[167237]gene. Biochem. Biophys. Res. Commun. 199: 841–847, 1994.

[167238]

[167239]

[167240]16788.

[167241]Johnston, J. J.; Kelley, R. I.; Crawford, T. O.; Morton, D. H.;

[167242]Agarwala, R.; Koch, T.; Schaffer, A. A.; Francomano, C. A.; Biesecker,

[167243]L. G.: A novel nemaline myopathy in the Amish caused by a mutation

[167244]in troponin T1. Am. J. Hum. Genet. 67: 814–821, 2000.

[167245]

[167246]

[167247]16789.

[167248]Novelli, G.; Gennarelli, M.; Rocchi, M.; Dallapiccola, B.: Assignment

[167249]of the slow troponin T (TNNT1) gene to chromosome 19
using polymerase

[167250]chain reaction. Hum. Genet. 88: 697–698, 1992.

[167251]

[167252]

[167253]16790.

[167254]Novelli, G.; Gennarelli, M.; Zelano, G.; Sangiuolo, F.; Roc-
chi,

[167255]M.; Dallapiccola, B.: Isolation and mapping of the slow
skeletal

[167256]troponin T using the polymerase chain reaction. (Abstract)
Cytogenet.

[167257]Cell Genet. 58: 2023, 1991.

[167258]

[167259]

[167260]16791.

[167261]Crow, R. S.: Peripheral neuritis in myelomatosis. Brit. Med.

[167262]J. 2: 802–804, 1956.

[167263]

[167264]

[167265]16792.

[167266]Dantz, D.; Bewersdorf, J.; Fruehwald-Schultes, B.; Kern, W.; Jelkmann,

[167267]W.; Born, J.; Fehm, H. L.; Peters, A.: Vascular endothelial growth

[167268]factor: a novel endocrine defensive response to hypoglycemia. J.

[167269]Clin. Endocr. Metab. 87: 835–840, 2002.

[167270]

[167271]

[167272]16793.

[167273]Ferrara, N.; Carver-Moore, K.; Chen, H.; Dowd, M.; Lu, L.; O'Shea,

[167274]K. S.; Powell-Braxton, L.; Hillan, K. J.; Moore, M. W.: Heterozygous

[167275]embryonic lethality induced by targeted inactivation of the VEGF gene. Nature 380:

[167276]439–442, 1996.

[167277]

[167278]

[167279]16794.

[167280]Folkman, J.: Angiogenesis in cancer, vascular, rheumatoid
and

[167281]other disease. Nature Med. 1: 27–31, 1995.

[167282]

[167283]

[167284]16795.

[167285]Fukumura, D.; Xavier, R.; Sugiura, T.; Chen, Y.; Park, E.–C.;
Lu,

[167286]N.; Selig, M.; Nielsen, G.; Taksir, T.; Jain, R. K.; Seed, B.:
Tumor

[167287]induction of VEGF promoter activity in stromal cells. Cell
94: 715–725,

[167288]1998.

[167289]

[167290]

[167291]16796.

[167292]Gerber, H.–P.; Malik, A. K.; Solar, G. P.; Sherman, D.;

Liang,

[167293]X. H.; Meng, G.; Hong, K.; Marsters, J. C.; Ferrara, N.: VEGF
regulates

[167294]haematopoietic stem cell survival by an internal autocrine
loop mechanism. Nature 417:

[167295]954–958, 2002.

[167296]

[167297]

[167298]16797.

[167299]Gerber, H.–P.; Vu, T. H.; Ryan, A. M.; Kowalski, J.; Werb, Z.;

[167300]Ferrara, N.: VEGF couples hypertrophic cartilage remodel-
ing, ossification

[167301]and angiogenesis during endochondral bone formation.

Nature Med. 5:

[167302]623–628, 1999.

[167303]

[167304]

[167305]16798.

[167306]Giordano, F. J.; Gerber, H.–P.; Williams, S.–P.; VanBruggen,
N.;

[167307]Bunting, S.; Ruiz–Lozano, P.; Gu, Y.; Nath, A. K.; Huang, Y.; Hickey,

[167308]R.; Dalton, N.; Peterson, K. L.; Ross, J., Jr.; Chien, K. R.; Ferrara,

[167309]N.: A cardiac myocyte vascular endothelial growth factor paracrine

[167310]pathway is required to maintain cardiac function. Proc. Nat. Acad.

[167311]Sci. 98: 5780–5785, 2001.

[167312]

[167313]

[167314]16799.

[167315]Helmlinger, G.; Endo, M.; Ferrara, N.; Hlatky, L.; Jain, R. K.

[167316]: Formation of endothelial cell networks. Nature 405: 139–141, 2000.

[167317]

[167318]

[167319]16800.

[167320]Hofman, P.; van Blijswijk, B. C.; Gaillard, P. J.; Vrensen, G.

[167321]F. J. M.; Schlingemann, R. O.: Endothelial cell hypertrophy

induced

[167322]by vascular endothelial growth factor in the retina: new insights

[167323]into the pathogenesis of capillary nonperfusion. Arch. Ophthalmol. 119:

[167324]861–866, 2001.

[167325]

[167326]

[167327]16801.

[167328]Holash, J.; Maisonpierre, P. C.; Compton, D.; Boland, P.; Alexander,

[167329]C. R.; Zagzag, D.; Yancopoulos, G. D.; Wiegand, S. J.: Vessel cooption,

[167330]regression, and growth in tumors mediated by angiopoietins and VEGF. Science 284:

[167331]1994–1998, 1999.

[167332]

[167333]

[167334]16802.

[167335]Jin, K.; Zhu, Y.; Sun, Y.; Mao, X. O.; Xie, L.; Greenberg, D.

[167336]A.: Vascular endothelial growth factor (VEGF) stimulates neurogenesis

[167337]in vitro and in vivo. Proc. Nat. Acad. Sci. 99: 11946–11950, 2002.

[167338]

[167339]

[167340]16803.

[167341]Mattei, M.–G.; Borg, J.–P.; Rosnet, O.; Marme, D.; Birnbaum, D.

[167342]: Assignment of vascular endothelial growth factor (VEGF) and placenta

[167343]growth factor (PIGF) genes to human chromosome 6p12–p21 and 14q24–q31

[167344]regions, respectively. Genomics 32: 168–169, 1996.

[167345]

[167346]

[167347]16804.

[167348]Miralles, G. D.; O'Fallon, J. R.; Talley, N. J.: Plasma-cell

[167349]dyscrasia with polyneuropathy: the spectrum of POEMS syndrome. New

[167350]Eng. J. Med. 327: 1919–1923, 1992.

[167351]

[167352]

[167353]16805.

[167354]Mueller, M. D.; Vigne, J.–L.; Minchenko, A.; Lebovic, D. I.;
Leitman,

[167355]D. C.; Taylor, R. N.: Regulation of vascular endothelial
growth factor

[167356](VEGF) gene transcription by estrogen receptors alpha and
beta. Proc.

[167357]Nat. Acad. Sci. 97: 10972–10977, 2000.

[167358]

[167359]

[167360]16806.

[167361]Nakanishi, T.; Sobue, I.; Toyokura, Y.; Nishitani, H.;
Kuroiwa,

[167362]Y.; Satoyoshi, E.; Tsubaki, T.; Igata, A.; Ozaki, Y.: The
Crow–Fukase

[167363]syndrome: a study of 102 cases in Japan. Neurology 34:
712–720,

[167364]1984.

[167365]

[167366]

[167367]16807.

[167368]Niimi, H.; Arimura, K.; Jonosono, M.; Hashiguchi, T.;

Kawabata,

[167369]M.; Osame, M.; Kitajima, I.: VEGF is causative for pulmonary hypertension

[167370]in a patient with Crow–Fukase (POEMS) syndrome. Intern. Med. 39:

[167371]1101–1104, 2000.

[167372]

[167373]

[167374]16808.

[167375]Oosthuyse, B.; Moons, L.; Storkebaum, E.; Beck, H.;

Nuyens, D.;

[167376]Brusselmans, K.; Van Dorpe, J.; Hellings, P.; Gorselink, M.;

Heymans,

[167377]S.; Theilmeier, G.; Dewerchin, M.; and 20 others: Deletion of the

[167378]hypoxia-response element in the vascular endothelial growth factor

[167379]promoter causes motor neuron degeneration. Nature Genet. 28: 131–138,

[167380]2001.

[167381]

[167382]

[167383]16809.

[167384]Poltorak, Z.; Cohen, T.; Sivan, R.; Kandelis, Y.; Spira, G.; Vlodavsky,

[167385]I.; Keshet, E.; Neufeld, G.: VEGF145, a secreted vascular endothelial

[167386]growth factor isoform that binds to extracellular matrix. J. Biol.

[167387]Chem. 272: 7151–7158, 1997.

[167388]

[167389]

[167390]16810.

[167391]Shimpo, S.: Solitary myeloma causing polyneuritis and endocrine

[167392]disorders. Nippon Rinsho 26: 2444–2456, 1968.

[167393]

[167394]

[167395]16811.

[167396]Soker, S.; Takashima, S.; Miao, H. Q.; Neufeld, G.; Klags-
brun,

[167397]M.: Neuropilin–1 is expressed by endothelial and tumor
cells as an

[167398]isoform–specific receptor for vascular endothelial growth
factor. Cell 92:

[167399]735–745, 1998.

[167400]

[167401]

[167402]16812.

[167403]Sone, H.; Kawakami, Y.; Sakauchi, M.; Nakamura, Y.; Taka-
hashi,

[167404]A.; Shimano, H.; Okuda, Y.; Segawa, T.; Suzuki, H.; Ya-
mada, N.: Neutralization

[167405]of vascular endothelial growth factor prevents collagen-
induced arthritis

[167406]and ameliorates established disease in mice. Biochem.
Biophys. Res.

[167407]Commun. 281: 562–568, 2001.

[167408]

[167409]

[167410]16813.

[167411]Springer, M. L.; Chen, A. S.; Kraft, P. E.; Bednarski, M.;
Blau,

[167412]H. M.: VEGF gene delivery to muscle: potential role for
vasculogenesis

[167413]in adults. Molec. Cell 2: 549–558, 1998.

[167414]

[167415]

[167416]16814.

[167417]Thurston, G.; Suri, C.; Smith, K.; McClain, J.; Sato, T. N.;
Yancopoulos,

[167418]G. D.; McDonald, D. M.: Leakage-resistant blood vessels in
mice transgenically

[167419]overexpressing angiopoietin–1. Science 286: 2511–2514,
1999.

[167420]

[167421]

[167422]16815.

[167423]Tischer, E.; Mitchell, R.; Hartman, T.; Silva, M.; Gospo-
darowicz,

[167424]D.; Fiddes, J. C.; Abraham, J. A.: The human gene for vas-
cular endothelial

[167425]growth factor: multiple protein forms are encoded
through alternative

[167426]exon splicing. J. Biol. Chem. 266: 11947–11954, 1991.

[167427]

[167428]

[167429]16816.

[167430]Osborne, L. R.; Martindale, D.; Scherer, S. W.; Shi, X.–M.;
Huizenga,

[167431]J.; Heng, H. H. Q.; Costa, T.; Pober, B.; Lew, L.; Brinkman,
J.; Rommens,

[167432]J.; Koop, B.; Tsui, L.–C.: Identification of genes from a
500–kb

[167433]region at 7q11.23 that is commonly deleted in Williams
syndrome patients. Genomics 36:

[167434]328–336, 1996.

[167435]

[167436]

[167437]16817.

[167438]Rampoldi, L.; Dobson–Stone, C.; Rubio, J. P.; Danek, A.;
Chalmers,

[167439]R. M.; Wood, N. W.; Verellen, C.; Ferrer, X.; Malandrini, A.;
Fabrizi,

[167440]G. M.; Brown, R.; Vance, J.; Pericak–Vance, M.; Rudolf, G.;
Carre,

[167441]S.; Alonso, E.; Manfredi, M.; Nemeth, A. H.; Monaco, A. P.:
A conserved

[167442]sorting–associated protein is mutant in chorea–
acanthocytosis. Nature

[167443]Genet. 28: 119–120, 2001.

[167444]

[167445]

[167446]16818.

[167447]Weese–Mayer, D. E.; Bolk, S.; Silvestri, J. M.; Chakravarti,
A.

[167448]: Idiopathic congenital central hypoventilation syndrome:
evaluation

[167449]of brain-derived neurotrophic factor genomic DNA se-
quence variation. Am.

[167450]J. Med. Genet. 107: 306–310, 2002.

[167451]

[167452]

[167453]16819.

[167454]Belden, W. J.; Barlowe, C.: Role of Erv29p in collecting sol-
uble

[167455]secretory proteins into ER-derived transport vesicles. Sci-
ence 294:

[167456]1528–1531, 2001.

[167457]

[167458]

[167459]16820.

[167460]Duhig, T.; Ruhrberg, C.; Mor, O.; Fried, M.: The human
surfeit

[167461]locus. Genomics 52: 72–78, 1998.

[167462]

[167463]

[167464]16821.

[167465]Reeves, J. E.; Fried, M.: The surf-4 gene encodes a novel
30 kDa

[167466]integral membrane protein. Molec. Membr. Biol. 12:
201-208, 1995.

[167467]

[167468]

[167469]16822.

[167470]Archer, B. T., III; Ozcelik, T.; Jahn, R.; Francke, U.; Sudhof,
[167471]T. C.: Structures and chromosomal localizations of two
human genes

[167472]encoding synaptobrevins 1 and 2. J. Biol. Chem. 265:
17267-17273,

[167473]1990.

[167474]

[167475]

[167476]16823.

[167477]Hu, K.; Carroll, J.; Fedorovich, S.; Rickman, C.; Sukhodub,

A.;

[167478]Davietov, B.: Vesicular restriction of synaptobrevin suggests a role

[167479]for calcium in membrane fusion. Nature 415: 646–650, 2002.

[167480]

[167481]

[167482]16824.

[167483]Franchi, F.; Biguzzi, E.; Cetin, I.; Facchetti, F.; Radaelli, T.;

[167484]Bozzo, M.; Pardi, G.; Faioni, E. M.: Mutations in the thrombomodulin

[167485]and endothelial protein C receptor genes in women with late fetal

[167486]loss. Brit. J. Haemat. 114: 641–646, 2001.

[167487]

[167488]

[167489]16825.

[167490]Perez, F.; Diamantopoulos, G. S.; Stalder, R.; Kreis, T. E.: CLIP–170

[167491]highlights growing microtubule ends in vivo. Cell 96:

517–527, 1999.

[167492]

[167493]

[167494]16826.

[167495]Pierre, P.; Pepperkok, R.; Kreis, T. E.: Molecular characteri-
zation

[167496]of two functional domains of CLIP–170 in vivo. J. Cell Sci.
107:

[167497]1909–1920, 1994.

[167498]

[167499]

[167500]16827.

[167501]Pierre, P.; Scheel, J.; Rickard, J. E.; Kreis, T. E.: CLIP–170

[167502]links endocytic vesicles to microtubules. Cell 70:
887–900, 1992.

[167503]

[167504]

[167505]16828.

[167506]Acland, G. M.; Aguirre, G. D.; Ray, J.; Zhang, Q.; Aleman,

T. S.;

[167507]Cideciyan, A. V.; Pearce-Kelling, S. E.; Anand, V.; Zeng, Y.;

Maguire,

[167508]A. M.; Jacobson, S. G.; Hauswirth, W. W.; Bennett, J.: Gene

therapy

[167509]restores vision in a canine model of childhood blindness.

Nature 28:

[167510]92–95, 2001.

[167511]

[167512]

[167513]16829.

[167514]Aguirre, G. D.; Baldwin, V.; Pearce-Kelling, S.; Narfstrom,

K.;

[167515]Ray, K.; Acland, G. M.: Congenital stationary night blind-

ness in

[167516]the dog: common mutation in the RPE65 gene indicates

founder effect. Molec.

[167517]Vision 4: 23, 1998. Note: Electronic Article.

[167518]

[167519]

[167520]16830.

[167521]Bavik, C.-O.; Busch, C.; Eriksson, U.: Characterization of a
plasma

[167522]retinol-binding protein membrane receptor expressed in
the retinal

[167523]pigment epithelium. J. Biol. Chem. 267: 23035–23042,
1992.

[167524]

[167525]

[167526]16831.

[167527]Felius, J.; Thompson, D. A.; Khan, N. W.; Bingham, E. L.;
Jamison,

[167528]J. A.; Kemp, J. A.; Sieving P. A.: Clinical course and visual
function

[167529]in a family with mutations in the RPE65 gene. Arch. Oph-
thal. 120:

[167530]55–61, 2002.

[167531]

[167532]

[167533]16832.

[167534]Grimm, C.; Wenzel, A.; Hafezi, F.; Yu, S.; Redmond, T. M.;
Reme,

[167535]C. E.: Protection of Rpe65–deficient mice identifies
rhodopsin as
[167536]a mediator of light–induced retinal degeneration. Nature
Genet. 25:
[167537]63–66, 2000.

[167538]

[167539]

[167540]16833.

[167541]Gu, S.; Thompson, D. A.; Srikumari, C. R. S.; Lorenz, B.;
Finckh,
[167542]U.; Nicoletti, A.; Murthy, K. R.; Rathmann, M.; Kumara–
manickavel,
[167543]G.; Denton, M. J.; Gal, A.: Mutations in RPE65 cause auto–
somal recessive
[167544]childhood–onset severe retinal dystrophy. Nature Genet.
17: 194–197,
[167545]1997.

[167546]

[167547]

[167548]16834.

[167549]Hamel, C. P.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.;
[167550]Redmond, T. M.: The gene for the retinal pigment epithe-
lium-specific
[167551]protein RPE65 is localized to human 1p31 and mouse 3.
Genomics 20:
[167552]509–512, 1994.
[167553]
[167554]
[167555]16835.
[167556]Hamel, C. P.; Tsilou, E.; Pfeffer, B. A.; Hooks, J. J.; Detrick,
[167557]B.; Redmond, T. M.: Molecular cloning and expression of
RPE65, a
[167558]novel retinal pigment epithelium-specific microsomal
protein that
[167559]is post-transcriptionally regulated in vitro. J. Biol. Chem.
268:
[167560]15751–15757, 1993.
[167561]
[167562]
[167563]16836.

[167564]Marlhens, F.; Bareil, C.; Griffoin, J.-M.; Zrenner, E.; Amal-
ric,

[167565]P.; Eliaou, C.; Liu, S.-Y.; Harris, E.; Redmond, T. M.; Ar-
naud, B.;

[167566]Claustres, M.; Hamel, C. P.: Mutations in RPE65 cause
Leber's congenital

[167567]amaurosis. (Letter) Nature Genet. 17: 139-141, 1997.

[167568]

[167569]

[167570]16837.

[167571]Morimura, H.; Fishman, G. A.; Grover, S. A.; Fulton, A. B.;
Berson,

[167572]E. L.; Dryja, T. P.: Mutations in the RPE65 gene in patients
with

[167573]autosomal recessive retinitis pigmentosa or Leber con-
genital amaurosis. Proc.

[167574]Nat. Acad. Sci. 95: 3088-3093, 1998.

[167575]

[167576]

[167577]16838.

- [167578]Narfstrom, K.; Wrigstad, A.; Nilsson, S. E.: The Briard dog:
[167579]a new animal model of congenital stationary night blind-
ness. Brit.
[167580]J. Ophthalm. 73: 750–756, 1989.
- [167581]
- [167582]
- [167583]16839.
- [167584]Nicoletti, A.; Wong, D. J.; Kawase, K.; Gibson, L. H.; Yang-
Feng,
[167585]T. L.; Richards, J. E.; Thompson, D. A.: Molecular charac-
terization
[167586]of the human gene encoding an abundant 61 kDa protein
specific to
[167587]the retinal pigment epithelium. Hum. Molec. Genet. 4:
641–649, 1995.
- [167588]
- [167589]
- [167590]16840.
- [167591]Adeokun, A. M.; West, S. P.; Ellis, F. R.; Halsall, P. J.; Hop-
kins,

[167592]P. M.; Foroughmand, A. M.; Iles, D. E.; Robinson, R. L.;
Stewart,

[167593]A. D.; Curran, J. L.: The G1021A substitution in the RYR1
gene does

[167594]not cosegregate with malignant hyperthermia susceptibil-
ity in a British

[167595]pedigree. Am. J. Hum. Genet. 60: 833–341, 1997.

[167596]

[167597]

[167598]16841.

[167599]Alestrom, A.; Fagerlund, T. H.; Berg, K.: A simple method
to detect

[167600]the RYR1 mutation G1021A, a cause of malignant hyper-
thermia susceptibility. Clin.

[167601]Genet. 47: 274–275, 1995.

[167602]

[167603]

[167604]16842.

[167605]Avila, G.; O'Brien, J. J.; Dirksen, R. T.: Excitation–contrac-
tion

[167606]uncoupling by a human central core disease mutation in
the ryanodine

[167607]receptor. Proc. Nat. Acad. Sci. 98: 4215–4220, 2001.

[167608]

[167609]

[167610]16843.

[167611]Barone, V.; Bertocchini, F.; Bottinelli, R.; Protasi, F.; Allen,

[167612]P. D.; Armstrong, C. F.; Reggiani, C.; Sorrentino, V.: Con-
tractile

[167613]impairment and structural alterations of skeletal muscles
from knockout

[167614]mice lacking type 1 and type 3 ryanodine receptors. FEBS
Lett. 422:

[167615]160–164, 1998.

[167616]

[167617]

[167618]16844.

[167619]Brandt, A.; Schleithoff, L.; Jurkat–Rott, K.; Klingler, W.;
Baur,

[167620]C.; Lehmann–Horn, F.: Screening of ryanodine receptor

gene in 105

[167621]malignant hyperthermia families: novel mutations and
concordance with

[167622]the in vitro contracture test. Hum. Molec. Genet. 8:
2055–2062,

[167623]1999.

[167624]

[167625]

[167626]16845.

[167627]Brown, R. L.; Pollock, A. N.; Couchman, K. G.; Hodges, M.;
Hutchinson,

[167628]D. O.; Waaka, R.; Lynch, P.; McCarthy, T. V.; Stowell, K. M.:
A novel

[167629]ryanodine receptor mutation and genotype–phenotype
correlation in

[167630]a large malignant hyperthermia New Zealand Maori pedi-
gree. Hum. Molec.

[167631]Genet. 9: 1515–1524, 2000.

[167632]

[167633]

[167634]16846.

[167635]Cavanna, J. S.; Greenfield, A. J.; Johnson, K. J.; Marks, A.
R.;

[167636]Nadal-Ginard, B.; Brown, S. D. M.: Establishment of the
mouse chromosome

[167637]7 region with homology to the myotonic dystrophy region
of human chromosome

[167638]19q. Genomics 7: 12–18, 1990.

[167639]

[167640]

[167641]16847.

[167642]Cheng, H.; Lederer, W. J.; Cannell, M. B.: Calcium sparks:
elementary

[167643]events underlying excitation–contraction coupling in heart
muscle. Science 262:

[167644]740–744, 1993.

[167645]

[167646]

[167647]16848.

[167648]Deufel, T.; Sudbrak, R.; Feist, Y.; Rubsam, B.; Du Chesne,
I.;

[167649]Schafer, K.-L.; Roewer, N.; Grimm, T.; Lehmann-Horn, F.;
Hartung,
[167650]E. J.; Muller, C. R.: Discordance, in a malignant hyperther-
mia pedigree,
[167651]between in vitro contracture-test phenotypes and haplo-
types for the
[167652]MHS1 region on chromosome 19q12-13.2, comprising
the C1840T transition
[167653]in the RYR1 gene. Am. J. Hum. Genet. 56: 1334-1342,
1995.
[167654]
[167655]
[167656]16849.
[167657]Eu, J. P.; Sun, J.; Xu, L.; Stamler, J. S.; Meissner, G.: The
[167658]skeletal muscle calcium release channel: coupled O₂ sen-
sor and NO
[167659]signaling functions. Cell 102: 499-509, 2000.
[167660]
[167661]
[167662]16850.

[167663]Fagerlund, T.; Ording, H.; Bendixen, D.; Berg, K.: Search
for

[167664]three known mutations in the RYR gene in 48 Danish fam-
ilies with malignant

[167665]hyperthermia. Clin. Genet. 46: 401–404, 1994.

[167666]

[167667]

[167668]16851.

[167669]Fagerlund, T.; Ording, H.; Bendixen, D.; Islander, G.; Ran-
klev–Twetman,

[167670]E.; Berg, K.: RYR1 mutation G1021A (gly341–to–arg) is not
frequent

[167671]in Danish and Swedish families with malignant hyperther-
mia susceptibility. Clin.

[167672]Genet. 49: 186–188, 1996.

[167673]

[167674]

[167675]16852.

[167676]Fagerlund, T. H.; Islander, G.; Ranklev–Twetman, E.; Berg,
K.

[167677]: Recombination between the postulated CCD/MHE/MHS
locus and RYR1

[167678]gene markers. Clin. Genet. 50: 455–458, 1996.

[167679]

[167680]

[167681]16853.

[167682]Fagerlund, T. H.; Islander, G.; Twetman, E. R.; Berg, K.: A
search

[167683]for three known RYR1 gene mutations in 41 Swedish fam-
ilies with predisposition

[167684]to malignant hyperthermia. Clin. Genet. 48: 12–16, 1995.

[167685]

[167686]

[167687]16854.

[167688]Fagerlund, T. H.; Ording, H.; Bendixen, D.; Islander, G.;
Ranklev

[167689]Twetman, E.; Berg, K.: Discordance between malignant hy-
perthermia

[167690]susceptibility and RYR1 mutation C1840T in two Scandi-
navian MH families

[167691]exhibiting this mutation. Clin. Genet. 52: 416–421, 1997.

[167692]

[167693]

[167694]16855.

[167695]Ferreiro, A.; Monnier, N.; Romero, N. B.; Leroy, J.–P.; Bon–
nemann,

[167696]C.; Haenggeli, C.–A.; Straub, V.; Voss, W. D.; Nivoche, Y.;
Jungbluth,

[167697]H.; Lemainque, A.; Voit, T.; Lunardi, J.; Fardeau, M.;
Guicheney,

[167698]P.: A recessive form of central core disease, transiently
presenting

[167699]as multi–minicore disease, is associated with a homozy–
gous mutation

[167700]in the ryanodine receptor type 1 gene. Ann. Neurol. 51:
750–759,

[167701]2002.

[167702]

[167703]

[167704]16856.

[167705]Fujii, J.; Otsu, K.; Zorzato, F.; de Leon, S.; Khanna, V. K.;
[167706]Weiler, J. E.; O'Brien, P. J.; MacLennan, D. H.: Identification
of
[167707]a mutation in porcine ryanodine receptor associated with
malignant
[167708]hyperthermia. Science 253: 448–451, 1991.
[167709]
[167710]
[167711]16857.
[167712]Gillard, E. F.; Otsu, K.; Fujii, J.; Duff, C.; de Leon, S.;
Khanna,
[167713]V. K.; Britt, B. A.; Worton, R. G.; MacLennan, D. H.: Poly-
morphisms
[167714]and deduced amino acid substitutions in the coding se-
quence of the
[167715]ryanodine receptor (RYR1) gene in individuals with malig-
nant hyperthermia. Genomics 13:
[167716]1247–1254, 1992.
[167717]
[167718]

[167719]16858.

[167720]Gillard, E. F.; Otsu, K.; Fujii, J.; Khanna, V. K.; de Leon, S.;

[167721]Derdemezi, J.; Britt, B. A.; Duff, C. L.; Worton, R. G.;

MacLennan,

[167722]D. H.: A substitution of cysteine for arginine 614 in the
ryanodine

[167723]receptor is potentially causative of human malignant hy-
perthermia. Genomics 11:

[167724]751-755, 1991.

[167725]

[167726]

[167727]16859.

[167728]Hall-Curran, J. L.; Stewart, A. D.; Ball, S. P.; Halsall, J. P.;

[167729]Hopkins, P. M.; Ellis, F. R.: No C1840 to T mutation in
RYR1 in malignant

[167730]hyperthermia.(Letter) Hum. Mutat. 2: 330, 1993.

[167731]

[167732]

[167733]16860.

[167734]Quane, K. A.; Keating, K. E.; Manning, B. M.; Healy, J. M. S.;

[167735]Monsieurs, K.; Heffron, J. J. A.; Lehane, M.; Heytens, L.;

Krivosic-Horber,

[167736]R.; Adnet, P.; Ellis, F. R.; Monnier, N.; Lunardi, J.; Mc-

Carthy, T.

[167737]V.: Detection of a novel common mutation in the ryan-
odine receptor

[167738]gene in malignant hyperthermia: implications for diagno-
sis and heterogeneity

[167739]studies. Hum. Molec. Genet. 3: 471-476, 1994.

[167740]

[167741]

[167742]16861.

[167743]Vincenti, V.; Cassano, C.; Rocchi, M.; Persico, M. G.: As-
signment

[167744]of the vascular endothelial growth factor gene to human
chromosome

[167745]6p21.3. Circulation 93: 1493-1495, 1996.

[167746]

[167747]

[167748]16862.

[167749]Alonso, M.; Barton, D. E.; Francke, U.: MAL, a membrane protein

[167750]associated with human T-cell differentiation, is encoded on human

[167751]chromosome 2, region cen-q13. (Abstract) Cytogenet. Cell Genet. 46:

[167752]570 only, 1987.

[167753]

[167754]

[167755]16863.

[167756]Alonso, M. A.; Barton, D. E.; Francke, U.: Assignment of the T-cell

[167757]differentiation gene MAL to human chromosome 2, region cen-q13. Immunogenetics 27:

[167758]91-95, 1988.

[167759]

[167760]

[167761]16864.

[167762]Alonso, M. A.; Weissman, S. M.: cDNA cloning and se-

quence of MAL,

[167763]a hydrophobic protein associated with human T-cell differentiation. Proc.

[167764]Nat. Acad. Sci. 84: 1997–2001, 1987.

[167765]

[167766]

[167767]16865.

[167768]Millan, J.; Puertollano, R.; Fan, L.; Rancano, C.; Alonso, M. A.

[167769]: The MAL proteolipid is a component of the detergent-insoluble membrane

[167770]subdomains of human T-lymphocytes. Biochem. J. 321: 247–252, 1997.

[167771]

[167772]

[167773]16866.

[167774]Rancano, C.; Rubio, T.; Alonso, M. A.: Alternative splicing of

[167775]human T-cell-specific MAL mRNA and its correlation with the exon/intron

[167776]organization of the gene. Genomics 21: 447–450, 1994.

[167777]

[167778]

[167779]16867.

[167780]Wong–Staal, F.; Dalla–Favera, R.; Franchini, G.; Gelmann, E. P.;

[167781]Gallo, R. C.: Three distinct genes in human DNA related to the transforming

[167782]genes of mammalian sarcoma retroviruses. Science 213: 226–228, 1981.

[167783]

[167784]

[167785]16868.

[167786]Aurias, A.; Rimbaut, C.; Buffe, D.; Dubousset, J.; Mazabraud, A.

[167787]: Chromosomal translocations in Ewing's sarcoma. (Letter) New Eng.

[167788]J. Med. 309: 496–497, 1983.

[167789]

[167790]

[167791]16869.

[167792]Bartram, C. R.; de Klein, A.; Hagemeijer, A.; Grosveld, G.;
Heisterkamp,

[167793]N.; Groffen, J.: Localization of the human c-sis oncogene
in Ph-1-positive

[167794]and Ph-1-negative chronic myelocytic leukemia by in situ
hybridization. Blood 63:

[167795]223-225, 1984.

[167796]

[167797]

[167798]16870.

[167799]Bechet, J.-M.; Bornkamm, G.; Freese, U.-K.; Lenoir, G. M.:
The

[167800]c-sis oncogene is not activated in Ewing's sarcoma.

(Letter) New

[167801]Eng. J. Med. 310: 393 only, 1984.

[167802]

[167803]

[167804]16871.

[167805]Bishop, J. M.: Enemies within: the genesis of retrovirus
oncogenes. Cell 23:

[167806]5–6, 1981.

[167807]

[167808]

[167809]16872.

[167810]Bolger, G. B.; Stamberg, J.; Kirsch, I. R.; Hollis, G. F.;
Schwarz,

[167811]D. F.; Thomas, G. H.: Chromosomal translocation t(14;22)
and oncogene

[167812](c-sis) variant in a pedigree with familial meningioma.
New Eng.

[167813]J. Med. 312: 564–567, 1985.

[167814]

[167815]

[167816]16873.

[167817]Cohen, J. B.; Levinson, A. D.: A point mutation in the last
intron

[167818]responsible for increased expression and transforming
activity of

[167819]the c-Ha-ras oncogene. Nature 334: 119–124, 1988.

[167820]

[167821]

[167822]16874.

[167823]Collins, T.; Ginsburg, D.; Boss, J. M.; Orkin, S. H.; Pober, J.

[167824]S.: Cultured human endothelial cells express platelet-derived growth

[167825]factor B chain: cDNA cloning and structural analysis. Nature 316:

[167826]748–750, 1985.

[167827]

[167828]

[167829]16875.

[167830]Dalla-Favera, R.; Gallo, R. C.; Giallongo, A.; Croce, C.: Chromosomal

[167831]localization of the human homolog (c-sis) of the simian sarcoma virus

[167832]onc gene. Science 218: 686–688, 1982.

[167833]

[167834]

[167835]16876.

[167836]Dalla-Favera, R.; Gelmann, E. P.; Gallo, R. C.; Wong-Staal, F.

[167837]: A human onc gene homologous to the transforming gene (v-sis) of

[167838]simian sarcoma virus. Nature 292: 31–35, 1981.

[167839]

[167840]

[167841]16877.

[167842]Deuel, T. F.; Huang, J. S.; Huang, S. S.; Stroobant, P.; Waterfield,

[167843]M. D.: Expression of a platelet-derived growth factor-like protein

[167844]in simian sarcoma virus transformed cells. Science 221: 1348–1350,

[167845]1983.

[167846]

[167847]

[167848]16878.

[167849]Devare, S. G.; Reddy, E. P.; Law, J. D.; Robbins, K. C.; Aaronson,

[167850]S. A.: Nucleotide sequence of the simian sarcoma virus
genome: demonstration

[167851]that its acquired cellular sequences encode the transform-
ing gene

[167852]product p28-sis. Proc. Nat. Acad. Sci. 80: 731-735, 1983.

[167853]

[167854]

[167855]16879.

[167856]Charlet-B., N.; Logan, P.; Singh, G.; Cooper, T. A.: Dy-
namic antagonism

[167857]between ETR-3 and PTB regulates cell type-specific alter-
native splicing. Molec.

[167858]Cell 9: 649-658, 2002.

[167859]

[167860]

[167861]16880.

[167862]Borglum, A. D.; Flint, T.; Tommerup, N.; Fleckner, J.;
Justesen,

[167863]J.; Kruse, T. A.: Assignment of the human tryptophanyl-
tRNA synthetase

[167864]gene (WARS) to chromosome 14q32.2–q32.32. Cytogenet.

Cell Genet. 73:

[167865]99–103, 1996.

[167866]

[167867]

[167868]16881.

[167869]Denney, R. M.; Borgaonkar, D.; Ruddle, F. H.: Order of
genes for

[167870]NP and TRPRS on chromosome 14. Cytogenet. Cell Genet.
22: 493–497,

[167871]1978.

[167872]

[167873]

[167874]16882.

[167875]Francke, U.; Denney, R. M.; Ruddle, F. H.: Intrachromoso-
mal gene

[167876]mapping in man: the gene for tryptophanyl–tRNA syn-
thetase maps in

[167877]region q21–qter of chromosome 14. Somat. Cell Genet. 3:
381–389,

[167878]1977.

[167879]

[167880]

[167881]16883.

[167882]Benson, K. F.; Horwitz, M.; Wolff, J.; Friend, K.; Thompson, E.;

[167883]White, S.; Richards, R. I.; Raskind, W. H.; Bird, T. D.: CAG repeat

[167884]expansion in autosomal dominant familial spastic paraparesis: novel

[167885]expansion in a subset of patients. Hum. Molec. Genet. 7: 1779–1786,

[167886]1998.

[167887]

[167888]

[167889]16884.

[167890]Fonknechten, N.; Mavel, D.; Byrne, P.; Davoine, C.–S.; Cruaud,

[167891]C.; Boentsch, D.; Samson, D.; Coutinho, P.; Hutchinson, M.; McMonagle,

[167892]P.; Burgunder, J.-M.; Tartaglione, A.; and 10 others: Spectrum
trum

[167893]of SPG4 mutations in autosomal dominant spastic paraplegia. Hum.

[167894]Molec. Genet. 9: 637-644, 2000.

[167895]

[167896]

[167897]16885.

[167898]Nielsen, J. E.; Koefoed, P.; Abell, K.; Hasholt, L.; Eiberg, H.;

[167899]Fenger, K.; Niebuhr, E.; Sorensen, S. A.: CAG repeat expansion in
pansion in

[167900]autosomal dominant pure spastic paraplegia linked to
chromosome 2p21-p24. Hum.

[167901]Molec. Genet. 6: 1811-1816, 1997.

[167902]

[167903]

[167904]16886.

[167905]Svenson, I. K.; Ashley-Koch, A. E.; Gaskell, P. C.; Riney, T.

[167906]J.; Cumming, W. J. K.; Kingston, H. M.; Hogan, E. L.; Bous-

tany, R.-M.

[167907]N.; Vance, J. M.; Nance, M. A.; Pericak-Vance, M. A.;
Marchuk, D.

[167908]A.: Identification and expression analysis of spastin gene
mutations

[167909]in hereditary spastic paraplegia. Am. J. Hum. Genet. 68:
1077-1085,

[167910]2001.

[167911]

[167912]

[167913]16887.

[167914]Ala-Kapee, M.; Nevanlinna, H.; Mali, M.; Jalkanen, M.;
Schroder,

[167915]J.: Localization of gene for human syndecan, an integral
membrane

[167916]proteoglycan and a matrix receptor, to chromosome 2.
Somat. Cell

[167917]Molec. Genet. 16: 501-505, 1990.

[167918]

[167919]

[167920]16888.

[167921]Alexander, C. M.; Reichsman, F.; Hinkes, M. T.; Lincecum, J.; Becker,

[167922]K. A.; Cumberledge, S.; Bernfield, M.: Syndecan-1 is required for

[167923]Wnt-1-induced mammary tumorigenesis in mice. *Nature Genet.* 25: 329-332,

[167924]2000.

[167925]

[167926]

[167927]16889.

[167928]Mali, M.; Jaakkola, P.; Arvilommi, A.-M.; Jalkanen, M.: Sequence

[167929]of human syndecan indicates a novel gene family of integral membrane

[167930]proteoglycans. *J. Biol. Chem.* 265: 6884-6889, 1990.

[167931]

[167932]

[167933]16890.

[167934]Oettinger, H. F.; Streeter, H.; Lose, E.; Copeland, N. G.; Gilbert,

[167935]D. J.; Justice, M. J.; Jenkins, N. A.; Mohandas, T.; Bernfield, M.

[167936]: Chromosome mapping of the murine syndecan gene. Genomics 11: 334–338,

[167937]1991.

[167938]

[167939]

[167940]16891.

[167941]Reizes, O.; Lincecum, J.; Wang, Z.; Goldberger, O.; Huang, L.;

[167942]Kaksonen, M.; Ahima, R.; Hinkes, M. T.; Barsh, G. S.; Rauvala, H.;

[167943]Bernfield, M.: Transgenic expression of syndecan–1 uncovers a physiological

[167944]control of feeding behavior by syndecan–3. Cell 106: 105–116, 2001.

[167945]

[167946]

[167947]16892.

[167948]Sanderson, R. D.; Lalor, P.; Bernfield, M.: B lymphocytes express

[167949]and lose syndecan at specific stages of differentiation.

Cell Regulation 1:

[167950]27–35, 1989.

[167951]

[167952]

[167953]16893.

[167954]Westman, P.; Hsieh, C.–L.; Mali, M.; Jalkanen, M.; Francke, U.;

[167955]Schroder, J.: Assignment of the human syndecan (SDC) gene to short

[167956]arm of chromosome 2. (Abstract) Cytogenet. Cell Genet. 58: 1873–1874,

[167957]1991.

[167958]

[167959]

[167960]16894.

[167961]Burns, A. L.; Magendzo, K.; Shirvan, A.; Srivastava, M.; Rojas,

[167962]E.; Alijani, M. R.; Pollard, H. B.: Calcium channel activity of purified

[167963]human synexin and structure of the human synexin gene.

Proc. Nat.

[167964]Acad. Sci. 86: 3798–3802, 1989.

[167965]

[167966]

[167967]16895.

[167968]Caohuy, H.; Srivastava, M.; Pollard, H. B.: Membrane fusion
protein

[167969]synexin (annexin VII) as a Ca^{2+} /GTP sensor in exocytotic
secretion. Proc.

[167970]Nat. Acad. Sci. 93: 10797–10802, 1996.

[167971]

[167972]

[167973]16896.

[167974]Herr, C.; Smyth, N.; Ullrich, S.; Yun, F.; Sasse, P.; Hes-
cheler,

[167975]J.; Fleischmann, B.; Lasek, K.; Brixius, K.; Schwinger, R. H.
G.;

[167976]Fassler, R.; Schroder, R.; Noegel, A. A.: Loss of annexin A7
leads

[167977]to alterations in frequency-induced shortening of isolated murine

[167978]cardiomyocytes. Molec. Cell. Biol. 21: 4119–4128, 2001.

[167979]

[167980]

[167981]16897.

[167982]Magendzo, K.; Shirvan, A.; Cultraro, C.; Srivastava, M.; Pollard,

[167983]H. B.; Burns, A. L.: Alternative splicing of human synexin mRNA in

[167984]brain, cardiac, and skeletal muscle alters the unique N-terminal domain. J.

[167985]Biol. Chem. 266: 3228–3232, 1991.

[167986]

[167987]

[167988]16898.

[167989]Shirvan, A.; Srivastava, M.; Wang, M. G.; Cultraro, C.; Magendzo,

[167990]K.; McBride, O. W.; Pollard, H. B.; Burns, A. L.: Divergent structure

[167991]of the human synexin (annexin VII) gene and assignment
to chromosome

[167992]10. Biochemistry 33: 6888–6901, 1994.

[167993]

[167994]

[167995]16899.

[167996]Srivastava, M.; Atwater, I.; Glasman, M.; Leighton, X.; Gop-
ing,

[167997]G.; Caohuy, H.; Miller, G.; Pichel, J.; Westphal, H.; Mears,
D.; Rojas,

[167998]E.; Pollard, H. B.: Defects in inositol 1,4,5–trisphosphate
receptor

[167999]expression, Ca(2+) signaling, and insulin secretion in the
anx7(+/-)

[168000]knockout mouse. Proc. Nat. Acad. Sci. 96: 13783–13788,
1999.

[168001]

[168002]

[168003]16900.

[168004]Srivastava, M.; Bubendorf, L.; Srikantan, V.; Fossom, L.;
Nolan,

[168005]L.; Glasman, M.; Leighton, X.; Fehrle, W.; Pittaluga, S.; Raffeld,

[168006]M.; Koivisto, P.; Willi, N.; Gasser, T. C.; Kononen, J.; Sauter, G.;

[168007]Kallioniemi, O. P.; Srivastava, S.; Pollard, H. B.: ANX7, a candidate

[168008]tumor suppressor gene for prostate cancer. Proc. Nat. Acad. Sci. 98:

[168009]4575–4580, 2001.

[168010]

[168011]

[168012]16901.

[168013]Zhang–Keck, Z.–Y.; Srivastava, M.; Kozak, C. A.; Caohuy, H.; Shirvan,

[168014]A.; Burns, A. L.; Pollard, H. B.: Genomic organization and chromosomal

[168015]localization of the mouse synexin gene. Biochem. J. 301: 835–845,

[168016]1994.

[168017]

[168018]

[168019]16902.

[168020]Asch, A. S.; Barnwell, J.; Silverstein, R. L.; Nachman, R. L.:

[168021]Isolation of the thrombospondin membrane receptor. J.

Clin. Invest. 79:

[168022]1054–1061, 1987.

[168023]

[168024]

[168025]16903.

[168026]de Fraipont, F.; El Atifi, M.; Gicquel, C.; Bertagna, X.;

Chambaz,

[168027]E. M.; Feige, J. J.: Expression of the angiogenesis markers

vascular

[168028]endothelial growth factor–A, thrombospondin–1, and

platelet–derived

[168029]endothelial cell growth factor in human sporadic adreno–

cortical tumors:

[168030]correlation with genotypic alterations. J. Clin. Endocr.

Metab. 85:

[168031]4734–4741, 2000.

[168032]

[168033]

[168034]16904.

[168035]Dixit, V. M.; Hennessy, S. W.; Grant, G. A.; Rotwein, P.;

Frazier,

[168036]W. A.: Characterization of a cDNA encoding the heparin
and collagen

[168037]binding domains of human thrombospondin. Proc. Nat.
Acad. Sci. 83:

[168038]5449–5453, 1986.

[168039]

[168040]

[168041]16905.

[168042]Frazier, W. A.: Thrombospondin: a modular adhesive gly-
coprotein

[168043]of platelets and nucleated cells. J. Cell Biol. 105: 625–632,
1987.

[168044]

[168045]

[168046]16906.

[168047]Jaffe, E.; Bornstein, P.; Disteche, C. M.: Mapping of the

thrombospondin

[168048]gene to human chromosome 15 and mouse chromosome
2 by in situ hybridization. Genomics 7:

[168049]123–126, 1990.

[168050]

[168051]

[168052]16907.

[168053]Lawler, J.; Sunday, M.; Thibert, V.; Duquette, M.; George,
E. L.;

[168054]Rayburn, H.; Hynes, R. O.: Thrombospondin–1 is required
for normal

[168055]murine pulmonary homeostasis and its absence causes
pneumonia. J.

[168056]Clin. Invest. 101: 982–992, 1998.

[168057]

[168058]

[168059]16908.

[168060]Rodriguez–Manzaneque, J. C.; Lane, T. F.; Ortega, M. A.;
Hynes,

[168061]R. O.; Lawler, J.; Iruela–Arispe, M. L.: Thrombospondin–1

suppresses

[168062]spontaneous tumor growth and inhibits activation of matrix metalloproteinase-9

[168063]and mobilization of vascular endothelial growth factor.

Proc. Nat.

[168064]Acad. Sci. 98: 12485–12490, 2001.

[168065]

[168066]

[168067]16909.

[168068]Wolf, F. W.; Eddy, R. L.; Shows, T. B.; Dixit, V. M.: Structure

[168069]and chromosomal localization of the human thrombospondin gene. Genomics 6:

[168070]685–691, 1990.

[168071]

[168072]

[168073]16910.

[168074]Imai, K.; Sarker, A. H.; Akiyama, K.; Ikeda, S.; Yao, M.;

Tsutsui,

[168075]K.; Shohmori, T.; Seki, S.: Genomic structure and sequence of a human

[168076]homologue (NTHL1/NTH1) of Escherichia coli endonuclease III with those

[168077]of the adjacent parts of TSC2 and SLC9A3R2 genes. Gene 222: 287–295,

[168078]1998.

[168079]

[168080]

[168081]16911.

[168082]Mira, J.–P.; Cariou, A.; Grall, F.; Delclaux, C.; Losser, M.–R.;

[168083]Heshmati, F.; Cheval, C.; Monchi, M.; Teboul, J.–L.; Riche, F.; Leleu,

[168084]G.; Arbibe, L.; Mignon, A.; Delpech, M.; Dhainaut, J.–F.: Association

[168085]of TNF2, a TNF–alpha promoter polymorphism, with septic shock susceptibility

[168086]and mortality: a multicenter study. J.A.M.A. 282: 561–568, 1999.

[168087]

[168088]

[168089]16912.

[168090]Moffatt, M. F.; Cookson, W. O. C. M.: Tumour necrosis factor

[168091]haplotypes and asthma. Hum. Molec. Genet. 6: 551–554, 1997.

[168092]

[168093]

[168094]16913.

[168095]Moraes, M. O.; Duppre, N. C.; Suffys, P. N.; Santos, A. R.; Almeida,

[168096]A. S.; Nery, J. A. C.; Sampaio, E. P.; Sarno, E. N.: Tumor necrosis

[168097]factor–alpha promoter polymorphism TNF2 is associated with a stronger

[168098]delayed–type hypersensitivity reaction in the skin of borderline tuberculoid

[168099]leprosy patients. Immunogenetics 53: 45–47, 2001.

[168100]

[168101]

[168102]16914.

[168103]Mulcahy, B.; Waldron–Lynch, F.; McDermott, M. F.; Adams, C.; Amos,

[168104]C. I.; Zhu, D. K.; Ward, R. H.; Clegg, D. O.; Shanahan, F.;
Molloy,

[168105]M. G.; O'Gara, F.: Genetic variability in the tumor necrosis
factor–lymphotoxin

[168106]region influences susceptibility to rheumatoid arthritis.
Am. J.

[168107]Hum. Genet. 59: 676–683, 1996.

[168108]

[168109]

[168110]16915.

[168111]Muller, U.; Jongeneel, C. V.; Nedospasov, S. A.; Lindahl, K.
F.;

[168112]Steinmetz, M.: Tumour necrosis factor and lymphotoxin
genes map close

[168113]to H-2D in the mouse major histocompatibility complex.
Nature 325:

[168114]265–267, 1987.

[168115]

[168116]

[168117]16916.

[168118]Nadel, S.; Newport, M. J.; Booy, R.; Levin, M.: Variation in
[168119]the tumor necrosis factor- α gene promoter region
may be associated
[168120]with death from meningococcal disease. J. Infect. Dis.
174: 878-880,
[168121]1996.

[168122]

[168123]

[168124]16917.

[168125]Nedospasov, S. A.; Hirt, B.; Shakhov, A. N.; Dobrynin, V.
N.;

[168126]Kawashima, E.; Accolla, R. S.; Jongeneel, C. V.: The genes
for tumor

[168127]necrosis factor (TNF- α) and lymphotoxin (TNF- β)
are tandemly

[168128]arranged on chromosome 17 of the mouse. Nucleic Acids
Res. 14: 7713-7725,

[168129]1986.

[168130]

[168131]

[168132]16918.

[168133]Nedwin, G. E.; Naylor, S. L.; Sakaguchi, A. Y.; Smith, D.;
Jarrett–Nedwin,

[168134]J.; Pennica, D.; Goeddel, D. V.; Gray, P. W.: Human lym-
photoxin and

[168135]tumor necrosis factor genes: structure, homology and
chromosomal localization. Nucleic

[168136]Acids Res. 13: 6361–6373, 1985.

[168137]

[168138]

[168139]16919.

[168140]Norman, R. A.; Bogardus, C.; Ravussin, E.: Linkage be-
tween obesity

[168141]and a marker near the tumor necrosis factor–alpha locus
in Pima Indians. J.

[168142]Clin. Invest. 96: 158–162, 1995.

[168143]

[168144]

[168145]16920.

[168146]Obayashi, H.; Hasegawa, G.; Fukui, M.; Kamiuchi, K.; Kita–

mura,

[168147]A.; Ogata, M.; Kanaitzuka, T.; Shigeta, H.; Kitagawa, Y.;

Nakano,

[168148]K.; Nishimura, M.; Ohta, M.; Nakamura, N.: Tumor necro-

sis factor

[168149]microsatellite polymorphism influences the development

of insulin

[168150]dependency in adult-onset diabetes patients with the

DRB1*1502-DQB1*0601

[168151]allele and anti-glutamic acid decarboxylase antibodies. J.

Clin.

[168152]Endocr. Metab. 85: 3348-3351, 2000.

[168153]

[168154]

[168155]16921.

[168156]Old, L. J.: Tumor necrosis factor (TNF). Science 230:
630–632,

[168157]1985.

[168158]

[168159]

[168160]16922.

[168161]Pennica, D.; Nedwin, G. E.; Hayflick, J. S.; Seeburg, P. H.;
Derynck,

[168162]R.; Palladino, M. A.; Kohr, W. J.; Aggarwal, B. B.; Goeddel,
D. V.

[168163]: Human tumour necrosis factor: precursor structure, ex-
pression and

[168164]homology to lymphotoxin. Nature 312: 724–729, 1984.

[168165]

[168166]

[168167]16923.

[168168]Ragoussis, J.; Bloemer, K.; Weiss, E. H.; Ziegler, A.: Local–

ization

[168169]of the genes for tumor necrosis factor and lymphotoxin
between the

[168170]HLA class I and III regions by field inversion gel elec-
trophoresis. Immunogenetics 27:

[168171]66–69, 1988.

[168172]

[168173]

[168174]16924.

[168175]Rasmussen, S. K.; Urhammer, S. A.; Jensen, J. N.; Hansen,
T.;

[168176]Borch-Johnsen, K.; Pedersen, O.: The –238 and –308 G6A
polymorphisms

[168177]of the tumor necrosis factor alpha gene promoter are not
associated

[168178]with features of the insulin resistance syndrome or altered
birth

[168179]weight in Danish Caucasians. J. Clin. Endocr. Metab. 85:
1731–1734,

[168180]2000.

[168181]

[168182]

[168183]16925.

[168184]Rosmond, R.; Chagnon, M.; Bouchard, C.; Bjorntorp, P.: G-
308A

[168185]polymorphism of the tumor necrosis factor alpha gene
promoter and

[168186]salivary cortisol secretion. J. Clin. Endocr. Metab. 86:
2178-2180,

[168187]2001.

[168188]

[168189]

[168190]16926.

[168191]Roy, S.; McGuire, W.; Mascie-Taylor, C. G.; Saha, B.; Hazra,
S.

[168192]K.; Hill, A. V.; Kwiatkowski, D.: Tumor necrosis factor pro-
moter

[168193]polymorphism and susceptibility to lepromatous leprosy.
J. Infect.

[168194]Dis. 176: 530-532, 1997.

[168195]

[168196]

[168197]16927.

[168198]Ruuls, S. R.; Sedgwick, J. D.: Unlinking tumor necrosis factor

[168199]biology from the major histocompatibility complex: lessons from human

[168200]genetics and animal models. Am. J. Hum. Genet. 65: 294–301, 1999.

[168201]

[168202]

[168203]16928.

[168204]Spies, T.; Morton, C. C.; Nedospasov, S. A.; Fiers, W.; Pious,

[168205]D.; Strominger, J. L.: Genes for the tumor necrosis factors alpha

[168206]and beta are linked to the human major histocompatibility complex. Proc.

[168207]Nat. Acad. Sci. 83: 8699–8702, 1986.

[168208]

[168209]

[168210]16929.

[168211]van Hensbroek, M. B.; Palmer, A.; Onyiorah, E.; Schneider, G.;

[168212]Jaffar, S.; Dolan, G.; Memming, H.; Frenkel, J.; Enwere, G.; Bennett,

[168213]S.; Kwiatkowski, D.; Greenwood, B.: The effect of a monoclonal antibody

[168214]to tumor necrosis factor on survival from childhood cerebral malaria. J.

[168215]Infect. Dis. 174: 1091–1097, 1996.

[168216]

[168217]

[168218]16930.

[168219]Sashio, H.; Tamura, K.; Ito, R.; Yamamoto, Y.; Bamba, H.; Kosaka,

[168220]T.; Fukui, S.; Sawada, K.; Fukuda, Y.; Tamura, K.; Satomi, M.; Shimoyama,

[168221]T.; Furuyama, J.: Polymorphisms of the TNF gene and the TNF receptor

[168222]superfamily member 1B gene are associated with susceptibility to ulcerative

[168223]colitis and Crohn's disease, respectively. Immunogenetics
53: 1020–1027,
[168224]2002.

[168225]

[168226]

[168227]16931.

[168228]Van Ostade, X.; Vandenabeele, P.; Everaerd, B.; Loetscher,
H.;

[168229]Gentz, R.; Brockhaus, M.; Lesslauer, W.; Tavernier, J.;
Brouckaert,

[168230]P.; Fiers, W.: Human TNF mutants with selective activity on
the p55

[168231]receptor. Nature 361: 266–269, 1993.

[168232]

[168233]

[168234]16932.

[168235]Wang, A. M.; Creasey, A. A.; Ladner, M. B.; Lin, L. S.;
Strickler,

[168236]J.; Van Arsdel, J. N.; Yamamoto, R.; Mark, D. F.: Molecular
cloning

[168237]of the complementary DNA for human tumor necrosis factor. Science 228:

[168238]149–154, 1985.

[168239]

[168240]

[168241]16933.

[168242]Wilson, A. G.; Symons, J. A.; McDowell, T. L.; et al: Effects

[168243]of a polymorphism in the human tumor necrosis factor alpha promoter

[168244]on transcriptional activation. Proc. Nat. Acad. Sci. 94:
3195–3199,

[168245]1997.

[168246]

[168247]

[168248]16934.

[168249]Winchester, E. C.; Millwood, I. Y.; Rand, L.; Penny, M. A.;
Kessling,

[168250]A. M.: Association of the TNF-alpha-308 (G-A) polymorphism with self-reported

[168251]history of childhood asthma. Hum. Genet. 107: 591–596,

2000.

[168252]

[168253]

[168254]16935.

[168255]Witte, J. S.; Palmer, L. J.; O'Connor, R. D.; Hopkins, P. J.;

[168256]Hall, J. M.: Relation between tumour necrosis factor polymorphism

[168257]TNF-alpha-308 and risk of asthma. Europ. J. Hum. Genet. 10: 82-85,

[168258]2002.

[168259]

[168260]

[168261]16936.

[168262]Peter, D.; Finn, J. P.; Klisak, I.; Liu, Y.; Kojis, T.; Heinzmann,

[168263]C.; Roghani, A.; Sparkes, R. S.; Edwards, R. H.: Chromosomal localization

[168264]of the human vesicular amine transporter genes. Genomics 18: 720-723,

[168265]1993.

[168266]

[168267]

[168268]16937.

[168269]Roghani, A.; Welch, C.; Xia, Y.-R.; Liu, Y.; Peter, D.; Finn, J.

[168270]P.; Edwards, R. H.; Luskis, A. J.: Assignment of the mouse
vesicular

[168271]monoamine transporter genes, Slc18a1 and Slc18a2, to
chromosomes 8

[168272]and 19 by linkage analysis. Mammalian Genome 7:
393–394, 1996.

[168273]

[168274]

[168275]16938.

[168276]Lania, L.; Donti, E.; Pannuti, A.; Pascucci, A.; Pengue, G.;
Feliciello,

[168277]I.; La Mantia, G.; Lanfranccone, L.; Pelicci, P.-G.: cDNA iso-
lation,

[168278]expression analysis, and chromosomal localization of two
human zinc

[168279]finger genes. Genomics 6: 333–340, 1990.

[168280]

[168281]

[168282]16939.

[168283]Thiesen, H.-J.: Multiple genes encoding zinc finger domains are

[168284]expressed in human T cells. New Biologist 2: 363–374, 1990.

[168285]

[168286]

[168287]16940.

[168288]Seite, P.; Huebner, K.; Rousseau–Merck, M. F.; Berger, R.; Thiesen,

[168289]H. J.: Two human genes encoding zinc finger proteins, ZNF12 (KOX3)

[168290]and ZNF26 (KOX20), map to chromosomes 7p22–p21 and 12q24.33, respectively. Hum.

[168291]Genet. 86: 585–590, 1991.

[168292]

[168293]

[168294]16941.

[168295]Gaynor, R. B.; Muchardt, C.; Diep, A.; Mohandas, T. K.;
Sparkes,

[168296]R. S.; Lusi, A. J.: Localization of the zinc finger DNA-
binding

[168297]protein HIV-EP1/MBP-1/PRDII-BF1 to human chromosome
6p22.3-p24. Genomics 9:

[168298]758-761, 1991.

[168299]

[168300]

[168301]16942.

[168302]Tripathi, R. K.; Droetto, S.; Spritz, R. A.: Many patients
with

[168303]'tyrosinase-positive' oculocutaneous albinism have tyrosi-
nase gene

[168304]mutations. (Abstract) Am. J. Hum. Genet. 51 (suppl.):
A179, 1992.

[168305]

[168306]

[168307]16943.

[168308]Brilliant, M. H.: The mouse pink-eyed dilution locus: a
model

[168309]for aspects of Prader–Willi syndrome, Angelman syndrome, and a form

[168310]of hypomelanosis of Ito. Mammalian Genome 3: 187–191, 1992.

[168311]

[168312]

[168313]16944.

[168314]Castle, D.; Kromberg, J.; Kowalsky, R.; Moosa, R.; Gillman, N.;

[168315]Zwane, E.; Fritz, V.: Visual evoked potentials in Negro carriers

[168316]of the gene for tyrosinase positive oculocutaneous albinism. J. Med.

[168317]Genet. 25: 835–837, 1988.

[168318]

[168319]

[168320]16945.

[168321]Colman, M. A.; Stevens, G.; Ramsay, M.; Kwon, B.; Jenkins, T.:

[168322]Exclusion of two candidate pigment loci, c and b, part of chromosome

[168323]11p, and 33 random polymorphic markers as the locus for
tyrosinase-positive

[168324]oculocutaneous albinism. Hum. Genet. 90: 556–560,
1993.

[168325]

[168326]

[168327]16946.

[168328]Donlon, T. A.; Lalande, M.; Wyman, A.; Bruns, G.; Latt, S.
A.:

[168329]Isolation of molecular probes associated with the chromo-
some 15 instability

[168330]in the Prader–Willi syndrome. Proc. Nat. Acad. Sci. 83:
4408–4412,

[168331]1986.

[168332]

[168333]

[168334]16947.

[168335]Durham–Pierre, D.; Gardner, J. M.; Nakatsu, Y.; King, R. A.;
Francke,

[168336]U.; Ching, A.; Aquaron, R.; del Marmol, V.; Brilliant, M. H.:
African

[168337]origin of an intragenic deletion of the human P gene in tyrosinase

[168338]positive oculocutaneous albinism. Nature Genet. 7: 176–179, 1994.

[168339]

[168340]

[168341]16948.

[168342]Durham–Pierre, D.; King, R. A.; Naber, J. M.; Laken, S.; Brilliant,

[168343]M. H.: Estimation of carrier frequency of a 2.7 kb deletion allele

[168344]of the P gene associated with OCA2 in African–Americans. Hum. Mutat. 7:

[168345]370–373, 1996.

[168346]

[168347]

[168348]16949.

[168349]Gardner, J. M.; Nakatsu, Y.; Gondo, Y.; Lee, S.; Lyon, M. F.; King,

[168350]R. A.; Brilliant, M. H.: The mouse pink–eyed dilution gene: association

[168351]with human Prader–Willi and Angelman syndromes. Sci-
ence 257: 1121–1124,
[168352]1992.

[168353]

[168354]

[168355]16950.

[168356]Gondo, Y.; Gardner, J. M.; Nakatsu, Y.; Durham–Pierre, D.;
Deveau,

[168357]S. A.; Kuper, C.; Brilliant, M. H.: High–frequency genetic
reversion

[168358]mediated by a DNA duplication: the mouse pink–eyed un-
stable mutation. Proc.

[168359]Nat. Acad. Sci. 90: 297–301, 1993.

[168360]

[168361]

[168362]16951.

[168363]Hagiwara, N.; Klewer, S. E.; Samson, R. A.; Erickson, D. T.;
Lyon,

[168364]M. F.; Brilliant, M. H.: Sox6 is a candidate gene for p100H
myopathy,

[168365]heart block, and sudden neonatal death. Proc. Nat. Acad.

Sci. 97:

[168366]4180–4185, 2000.

[168367]

[168368]

[168369]16952.

[168370]Hamabe, J.; Fukushima, Y.; Harada, N.; Abe, K.; Matsuo,
N.; Nagai,

[168371]T.; Yoshioka, A.; Tonoki, H.; Tsukino, R.; Niikawa, N.:
Molecular

[168372]study of the Prader–Willi syndrome: deletion, RFLP, and
phenotype

[168373]analyses of 50 patients. Am. J. Med. Genet. 41: 54–63,
1991.

[168374]

[168375]

[168376]16953.

[168377]Heim, R. A.; Dunn, D. S.; Candy, S. E.; Zwane, E.;
Kromberg, J.

[168378]G. R.; Jenkins, T.: The tyrosine–positive oculocutaneous
albinism

[168379]locus is not linked to the beta-globin locus in man. Hum.

Genet. 79:

[168380]89, 1988.

[168381]

[168382]

[168383]16954.

[168384]Kagore, F.; Lund, P. M.: Oculocutaneous albinism among
schoolchildren

[168385]in Harare, Zimbabwe. J. Med. Genet. 32: 859–861, 1995.

[168386]

[168387]

[168388]16955.

[168389]Kedda, M. A.; Stevens, G.; Manga, P.; Viljoen, C.; Jenkins,
T.;

[168390]Ramsay, M.: The tyrosinase-positive oculocutaneous al-
binism gene

[168391]shows locus homogeneity on chromosome 15q11–q13
and evidence of multiple

[168392]mutations in Southern African negroids. Am. J. Hum.
Genet. 54: 1078–1084,

[168393]1994.

[168394]

[168395]

[168396]16956.

[168397]Kerr, R.; Stevens, G.; Manga, P.; Salm, S.; John, P.; Haw, T.;

[168398]Ramsay, M.: Identification of P gene mutations in individuals with

[168399]oculocutaneous albinism in sub-Saharan Africa. Hum. Mutat. 15: 166-172,

[168400]2000.

[168401]

[168402]

[168403]16957.

[168404]King, R. A.: Personal Communication. Minneapolis, Minn. 12/31/1992.

[168405]

[168406]

[168407]16958.

[168408]Ansari-Lari, M. A.; Muzny, D. M.; Lu, J.; Lu, F.; Lilley, C. E.;

[168409]Spanos, S.; Malley, T.; Gibbs, R. A.: A gene-rich cluster
between

[168410]the CD4 and triosephosphate isomerase genes at human
chromosome 12p13. Genome

[168411]Res. 6: 314–326, 1996.

[168412]

[168413]

[168414]16959.

[168415]Bach, M. A.; Phan-Dinh-Tuy, F.; Bach, J. F.; Wallach, D.;
Biddison,

[168416]W. E.; Sharrow, S. O.; Goldstein, G.; Kung, P. C.: Unusual
phenotypes

[168417]of human inducer T cells as measured by OKT4 and re-
lated monoclonal

[168418]antibodies. J. Immun. 127: 980–982, 1981.

[168419]

[168420]

[168421]16960.

[168422]Browning, J.; Horner, J. W.; Pettoello-Mantovani, M.; Raker,
C.;

[168423]Yurasov, S.; DePinho, R. A.; Goldstein, H.: Mice transgenic
for human

[168424]CD4 and CCR5 are susceptible to HIV infection. Proc. Nat.
Acad. Sci. 94:

[168425]14637–14641, 1997.

[168426]

[168427]

[168428]16961.

[168429]Buttini, M.; Westland, C. E.; Masliah, E.; Yafeh, A. M.;
Wyss–Coray,

[168430]T.; Mucke, L.: Novel role of human CD4 molecule identi-
fied in neurodegeneration. Nature

[168431]Med. 4: 441–446, 1998.

[168432]

[168433]

[168434]16962.

[168435]Committee on Human Leukocyte Differentiation Antigens,
IUIS WHO

[168436]Nomenclature Subcommittee: Proposed nomenclature for
human leukocyte

[168437]differentiation antigens. Bull. WHO 5: 809–811, 1984.

Note: Alternate:

[168438]Immunology Today 5: 280 only, 1984.

[168439]

[168440]

[168441]16963.

[168442]Fukuda, T.; Matsunaga, M.; Kurata, A.; Mine, M.; Ikari, N.;
Katamine,

[168443]S.; Kanazawa, H.; Eguchi, K.; Nagataki, S.: Hereditary defi-
ciency

[168444]of OKT4–positive cells: studies for mode of inheritance
and lymphocyte

[168445]functions. Immunology 53: 643–649, 1984.

[168446]

[168447]

[168448]16964.

[168449]Fuller, T. C.; Trevithick, J. E.; Fuller, A. A.; Colvin, R. B.;

[168450]Cosimi, A. B.; Kung, P. C.: Antigenic polymorphism of the
T4 differentiation

[168451]antigen expressed on human T helper/inducer lympho–

cytes. Hum. Immun. 9:

[168452]89–102, 1984.

[168453]

[168454]

[168455]16965.

[168456]Gill, J. C.; Maples, J.; Nikaein, A.; Kirchner, P.; Lockhart, D.;

[168457]Snyder, A. J.; Montgomery, R. R.; Casper, J. T.: Inherited
absence

[168458]of OKT4 lymphocyte antigen in a chronically transfused
patient with

[168459]homozygous sickle cell disease. J. Pediat. 107: 251–253,
1985.

[168460]

[168461]

[168462]16966.

[168463]Irvine, D. J.; Purbhoo, M. A.; Krogsgaard, M.; Davis, M. M.:

[168464]Direct observation of ligand recognition by T cells. Nature
419:

[168465]845–849, 2002.

[168466]

[168467]

[168468]16967.

[168469]Isobe, M.; Huebner, K.; Maddon, P. J.; Littman, D. R.; Axel, R.;

[168470]Croce, C. M.: The gene encoding the T-cell surface protein T4 is

[168471]located on human chromosome 12. Proc. Nat. Acad. Sci. 83: 4399-4402,

[168472]1986.

[168473]

[168474]

[168475]16968.

[168476]Karol, R. A.; Eng, J.; Dennison, D. K.; Faris, E.; Marcus, D.

[168477]M.: Hereditary abnormalities of the OKT4 human lymphocyte epitope

[168478]in two families. J. Clin. Immun. 4: 71-74, 1984.

[168479]

[168480]

[168481]16969.

[168482]Kozbor, D.; Finan, J.; Nowell, P. C.; Croce, C. M.: The gene

[168483]encoding the T4 antigen maps to human chromosome 12.

J. Immun. 136:

[168484]1141–1143, 1986.

[168485]

[168486]

[168487]16970.

[168488]Maddon, P. J.; Molineaux, S. M.; Maddon, D. E.; Zimmerman, K.

[168489]A.; Godfrey, M.; Alt, F. W.; Chess, L.; Axel, R.: Structure
and expression

[168490]of the human and mouse T4 genes. Proc. Nat. Acad. Sci.
84: 9155–9159,

[168491]1987.

[168492]

[168493]

[168494]16971.

[168495]Piguet, V.; Gu, F.; Foti, M.; Demaurex, N.; Gruenberg, J.;

Carpentier,

[168496]J.-L.; Trono, D.: Nef-induced CD4 degradation: a diacidic-based motif

[168497]in Nef functions as a lysosomal targeting signal through the binding

[168498]of beta-COP in endosomes. Cell 97: 63-73, 1999.

[168499]

[168500]

[168501]16972.

[168502]Sato, M.; Hayashi, Y.; Yoshida, H.; Yanagawa, T.; Yura, Y.:

A

[168503]family with hereditary lack of T4+ inducer/helper T cell subsets in

[168504]peripheral blood lymphocytes. J. Immun. 132: 1071-1073, 1984.

[168505]

[168506]

[168507]16973.

[168508]Sawada, S.; Scarborough, J. D.; Killeen, N.; Littman, D. R.:

[168509]A lineage-specific transcriptional silencer regulates CD4

gene expression

[168510]during T lymphocyte development. Cell 77: 917–929,
1994.

[168511]

[168512]

[168513]16974.

[168514]Zinman, B.; Hanley, A. J. G.; Harris, S. B.; Kwan, J.; Fantus,

[168515]I. G.: Circulating tumor necrosis factor–alpha concentra-
tions in

[168516]a Native Canadian population with high rates of type 2 di-
abetes mellitus. J.

[168517]Clin. Endocr. Metab. 84: 272–278, 1999.

[168518]

[168519]

[168520]16975.

[168521]Spurr, N. K.; Goodfellow, P. N.; Sheer, D.; Bodmer, W. F.;
Vennstrom,

[168522]B.: Mapping of cellular oncogenes: ERBA1 is on chromo-
some 17. (Abstract) Cytogenet.

[168523]Cell Genet. 37: 591 only, 1984.

[168524]

[168525]

[168526]16976.

[168527]Thompson, C. C.; Weinberger, C.; Lebo, R.; Evans, R. M.:

Identification

[168528]of a novel thyroid hormone receptor expressed in the
mammalian central

[168529]nervous system. Science 237: 1610–1614, 1987.

[168530]

[168531]

[168532]16977.

[168533]Weinberger, C.; Thompson, C. C.; Ong, E. S.; Lebo, R.;

Gruol,

[168534]D. J.; Evans, R. M.: The c-erb-A gene encodes a thyroid
hormone receptor. Nature 324:

[168535]641–646, 1986.

[168536]

[168537]

[168538]16978.

[168539]Zabel, B. U.; Fournier, R. E. K.; Lalley, P. A.; Naylor, S. L.;

[168540]Sakaguchi, A. Y.: Cellular homologs of the avian erythroblastosis

[168541]virus erb-A and erb-B genes are syntenic in mouse but asyntenic in

[168542]man. Proc. Nat. Acad. Sci. 81: 4874-4878, 1984.

[168543]

[168544]

[168545]16979.

[168546]Carraway, K. L., III; Sliwkowski, M. X.; Akita, R.; Platko, J.

[168547]V.; Guy, P. M.; Nuijens, A.; Diamonti, A. J.; Vandlen, R. L.; Cantley,

[168548]L. C.; Cerione, R. A.: The erbB3 gene product is a receptor for heregulin. J.

[168549]Biol. Chem. 269: 14303-14306, 1994.

[168550]

[168551]

[168552]16980.

[168553]Cho, H.-S.; Leahy, D. J.: Structure of the extracellular re-

gion

[168554]of HER3 reveals an interdomain tether. Science 297:
1330–1333, 2002.

[168555]

[168556]

[168557]16981.

[168558]Kraus, M. H.; Issing, W.; Miki, T.; Popescu, N. C.; Aaron-
son, S.

[168559]A.: Isolation and characterization of ERBB3, a third mem-
ber of the

[168560]ERBB/epidermal growth factor receptor family: evidence
for overexpression

[168561]in a subset of human mammary tumors. Proc. Nat. Acad.
Sci. 86: 9193–9197,

[168562]1989.

[168563]

[168564]

[168565]16982.

[168566]Plowman, G. D.; Whitney, G. S.; Neubauer, M. G.; Green, J.
M.;

[168567]McDonald, V. L.; Todaro, G. J.; Shoyab, M.: Molecular

cloning and

[168568]expression of an additional epidermal growth factor re-
ceptor-related

[168569]gene. Proc. Nat. Acad. Sci. 87: 4905–4909, 1990.

[168570]

[168571]

[168572]16983.

[168573]Elchebly, M.; Payette, P.; Michaliszyn, E.; Cromlish, W.;
Collins,

[168574]S.; Loy, A. L.; Normandin, D.; Cheng, A.; Himms-Hagen, J.;
Chan, C.–C.;

[168575]Ramachandran, C.; Gresser, M. J.; Tremblay, M. L.;
Kennedy, B. P.

[168576]: Increased insulin sensitivity and obesity resistance in
mice lacking

[168577]the protein tyrosine phosphatase-1B gene. Science 283:
1544–1548,

[168578]1999.

[168579]

[168580]

[168581]16984.

[168582]Forsell, P. K. A. L.; Boie, Y.; Montalibet, J.; Collins, S.;

Kennedy,

[168583]B. P.: Genomic characterization of the human and mouse
protein tyrosine

[168584]phosphatase-1B genes. Gene 260: 145-153, 2000.

[168585]

[168586]

[168587]16985.

[168588]Gu, H. F.; Almgren, P.; Lindholm, E.; Frittitta, L.; Pizzuti,
A.;

[168589]Trischitta, V.; Groop, L. C.: Association between the hu-
man glycoprotein

[168590]PC-1 gene and elevated glucose and insulin levels in
paired-sibling

[168591]analysis. Diabetes 49: 1601-1603, 2000.

[168592]

[168593]

[168594]16986.

[168595]Haj, F. G.; Verveer, P. J.; Squire, A.; Neel, B. G.; Bastiaens,

[168596]P. I. H.: Imaging sites of receptor dephosphorylation by

PTP1B on

[168597]the surface of the endoplasmic reticulum. Science 295:
1708–1711,

[168598]2002.

[168599]

[168600]

[168601]16987.

[168602]Jia, Z.; Barford, D.; Flint, A. J.; Tonks, N. K.: Structural

[168603]basis for phosphotyrosine peptide recognition by protein
tyrosine

[168604]phosphatase 1B. Science 268: 1754–1758, 1995.

[168605]

[168606]

[168607]16988.

[168608]Kennedy, B. P.; Ramachandran, C.: Protein tyrosine phos-
phatase–1B

[168609]in diabetes. Biochem. Pharm. 60: 877–883, 2000.

[168610]

[168611]

[168612]16989.

[168613]Mok, A.; Cao, H.; Zinman, B.; Hanley, A. J. G.; Harris, S. B.;

[168614]Kennedy, B. P.; Hegele, R. A.: A single nucleotide polymorphism in

[168615]protein tyrosine phosphatase PTP-1B is associated with protection

[168616]from diabetes or impaired glucose tolerance in Oji-Cree. J. Clin.

[168617]Endocr. Metab. 87: 724-727, 2002.

[168618]

[168619]

[168620]16990.

[168621]Tonks, N. K.; Diltz, C. D.; Fischer, E. H.: Purification of the

[168622]major protein-tyrosine-phosphatases of human placenta. J. Biol. Chem. 263:

[168623]6722-6730, 1988.

[168624]

[168625]

[168626]16991.

[168627]Barnea, G.; Silvennoinen, O.; Shaanan,B.; Honegger, A. M.;
Canoll,

[168628]P. D.; D'Eustachio, P.; Morse, B.; Levy, J. B.; Laforgia, S.;
Huebner,

[168629]K.; Musacchio, J. M.; Sap, J.; Schlessinger, J.: Identification
of

[168630]a carbonic anhydrase-like domain in the extracellular re-
gion of RPTP-gamma

[168631]defines a new subfamily of receptor tyrosine phos-
phatases. Molec.

[168632]Cell. Biol. 13: 1497-1506, 1993.

[168633]

[168634]

[168635]16992.

[168636]Kastury, K.; Ohta, M.; Lasota, J.; Moir, D.; Dorman, T.;
LaForgia,

[168637]S.; Druck, T.; Huebner, K.: Structure of the human recep-
tor tyrosine

[168638]phosphatase gamma gene (PTPRG) and relation to the fa-
milial RCC t(3;8)

[168639]chromosome translocation. Genomics 32: 225-235, 1996.

[168640]

[168641]

[168642]16993.

[168643]LaForgia, S.; Morse, B.; Levy, J.; Barnea, G.; Cannizzaro, L.
A.;

[168644]Li, F.; Nowell, P. C.; Boghosian-Sell, L.; Glick, J.; Weston,
A.;

[168645]Harris, C. C.; Drabkin, H.; Patterson, D.; Croce, C. M.; Sch-
lessinger,

[168646]J.; Huebner, K.: Receptor protein-tyrosine phosphatase
gamma is a

[168647]candidate tumor suppressor gene at human chromosome
region 3p21. Proc.

[168648]Nat. Acad. Sci. 88: 5036-5040, 1991.

[168649]

[168650]

[168651]16994.

[168652]Latif, F.; Tory, K.; Modi, W.; Geil, L.; LaForgia, S.; Huebner,

[168653]K.; Zbar, B.; Lerman, M. I.: A MspI polymorphism and link-
age mapping

[168654]of the human protein-tyrosine phosphatase G (PTPRG)
gene. Hum. Molec.

[168655]Genet. 2: 91, 1993.

[168656]

[168657]

[168658]16995.

[168659]Ariyama, T.; Hasegawa, K.; Inazawa, J.; Mizuno, K.; Ogi-
moto, M.;

[168660]Katagiri, T.; Yakura, H.: Assignment of the human protein
tyrosine

[168661]phosphatase, receptor-type, zeta (PTPRZ) gene to chro-
mosome band 7q31.3. Cytogenet.

[168662]Cell Genet. 70: 52-54, 1995.

[168663]

[168664]

[168665]16996.

[168666]Schols, L.; Szymanski, S.; Peters, S.; Przuntek, H.; Epplen,
J.

[168667]T.; Hardt, C.; Riess, O.: Genetic background of apparently
idiopathic

[168668]sporadic cerebellar ataxia. Hum. Genet. 107: 132–137,
2000.

[168669]

[168670]

[168671]16997.

[168672]Chan, J. Y.; Cheung, M.–C.; Moi, P.; Chan, K.; Kan, Y. W.:
Chromosomal

[168673]localization of the human NF–E2 family of bZIP transcrip-
tion factors

[168674]by fluorescence in situ hybridization. Hum. Genet. 95:
265–269,

[168675]1995.

[168676]

[168677]

[168678]16998.

[168679]Chan, J. Y.; Han, X.–L.; Kan, Y. W.: Cloning of Nrf1, an NF–
E2–related

[168680]transcription factor, by genetic selection in yeast. Proc.
Nat. Acad.

[168681]Sci. 90: 11371–11375, 1993.

[168682]

[168683]

[168684]16999.

[168685]Chan, J. Y.; Kwong, M.; Lu, R.; Chang, J.; Wang, B.; Yen, T. S.

[168686]B.; Kan, Y. W.: Targeted disruption of the ubiquitous CNC-bZIP transcription

[168687]factor, Nrf-1, results in anemia and embryonic lethality in mice. EMBO

[168688]J. 17: 1779-1787, 1998.

[168689]

[168690]

[168691]17000.

[168692]Luna, L.; Johnsen, O.; Skartlien, A.; Pedeutour, F.; Turc-Carel,

[168693]C.; Prydz, H.; Kolsto, A.-B.: Molecular cloning of a putative novel

[168694]human bZIP transcription factor on chromosome 17q22. Genomics 22:

[168695]553-562, 1994.

[168696]

[168697]

[168698]17001.

[168699]Luna, L.; Johnsen, O.; Skartlien, A. H.; Pedeutour, F.; Turc-Carel,

[168700]C.; Prydz, H.; Kolsto, A.-B.: Molecular cloning of a putative novel

[168701]human bZIP transcription factor on chromosome 17q22.

Genomics 22:

[168702]553–562, 1994.

[168703]

[168704]

[168705]17002.

[168706]McKie, J.; Johnstone, K.; Mattei, M.-G.; Scambler, P.:

Cloning

[168707]and mapping of murine Nfe2l1. Genomics 25: 716–719, 1995.

[168708]

[168709]

[168710]17003.

[168711]McKie, J.; Scambler, P. J.: The Nfe2l1 gene maps to distal mouse

[168712]chromosome 11. Mammalian Genome 7: 89–90, 1996.

[168713]

[168714]

[168715]17004.

[168716]Kas, K.; Wlodarska, I.; Meyen, E.; Van den Berghe, H.; Van de Ven,

[168717]W. J. M.: Assignment of the gene encoding human Kruppel-related zinc

[168718]finger protein 4 (GLI4) to 8q24.3 by fluorescent in situ hybridization. Cytogenet.

[168719]Cell Genet. 72: 297–298, 1996.

[168720]

[168721]

[168722]17005.

[168723]Goldblum, S. E.; Ding, X.; Funk, S. E.; Sage, E. H.: SPARC (secreted

[168724]protein acidic and rich in cysteine) regulates endothelial cell shape

[168725]and barrier function. Proc. Nat. Acad. Sci. 91: 3448–3452,
1994.

[168726]

[168727]

[168728]17006.

[168729]Lossie, A. C.; Gordon, D. F.; Camper, S. A.: Localization of
thyrotropin–releasing

[168730]hormone receptor and thyrotroph embryonic factor on
mouse chromosome

[168731]15. Mammalian Genome 4: 621–623, 1993.

[168732]

[168733]

[168734]17007.

[168735]Matre, V.; Karlsen, H. E.; Wright, M. S.; Lundell, I.; Fjeld–
heim,

[168736]A. K.; Gabrielsen, O. S.; Larhammar, D.; Gautvik, K. M.:
Molecular

[168737]cloning of a functional human thyrotropin–releasing hor–
mone receptor. Biochem.

[168738]Biophys. Res. Commun. 195: 179–185, 1993.

[168739]

[168740]

[168741]17008.

[168742]Morrison, N.; Duthie, S. M.; Boyd, E.; Eidne, K. A.; Connor, J.

[168743]M.: Assignment of the gene encoding the human thyrotropin-releasing

[168744]hormone receptor to 8q23 by fluorescence in situ hybridization. Hum.

[168745]Genet. 93: 716-718, 1994.

[168746]

[168747]

[168748]17009.

[168749]Straub, R. E.; Frech, G. C.; Joho, R. H.; Gershengorn, M. C.:

[168750]Expression cloning of a cDNA encoding the mouse pituitary thyrotropin-releasing

[168751]hormone receptor. Proc. Nat. Acad. Sci. 87: 9514-9518, 1990.

[168752]

[168753]

[168754]17010.

[168755]Yamada, M.; Monden, T.; Konaka, S.; Mori, M.: Assignment
of human

[168756]thyrotropin-releasing hormone (TRH) receptor gene to
chromosome 8. Somat.

[168757]Cell Molec. Genet. 19: 577-580, 1993.

[168758]

[168759]

[168760]17011.

[168761]Zhao, D.; Yang, J.; Jones, K. E.; Gerald, C.; Suzuki, Y.;
Hogan,

[168762]P. G.; Chin, W. W.; Tashjian, A. H., Jr.: Molecular cloning of
a

[168763]complementary deoxyribonucleic acid encoding the thy-
rotropin-releasing

[168764]hormone receptor and regulation of its messenger ri-
bonucleic acid

[168765]in rat GH cells. Endocrinology 130: 3529-3536, 1992.

[168766]

[168767]

[168768]17012.

[168769]Bongarzone, I.; Butti, M. G.; Coronelli, S.; Borrello, M. G.;
Santoro,

[168770]M.; Mondellini, P.; Pilotti, S.; Fusco, A.; Della Porta, G.;
Pierotti,

[168771]M. A.: Frequent activation of ret protooncogene by fusion
with a

[168772]new activating gene in papillary thyroid carcinomas. Can-
cer Res. 54:

[168773]2979–2985, 1994.

[168774]

[168775]

[168776]17013.

[168777]Bongarzone, I.; Butti, M. G.; Fugazzola, L.; Pacini, F.;
Pinchera,

[168778]A.; Vorontsova, T. V.; Demidchik, E. P.; Pierotti, M. A.:
Comparison

[168779]of the breakpoint regions of ELE1 and RET genes involved
in the generation

[168780]of RET/PTC3 oncogene in sporadic and in radiation-as-
sociated papillary

[168781]thyroid carcinomas. Genomics 42: 252–259, 1997.

[168782]

[168783]

[168784]17014.

[168785]Klein, M.; Vignaud, J.–M.; Hennequin, V.; Toussaint, B.;
Bresler,

[168786]L.; Plenat, F.; Leclere, J.; Duprez, A.; Weryha, G.: Increased
expression

[168787]of the vascular endothelial growth factor is a pejorative
prognosis

[168788]marker in papillary thyroid carcinoma. J. Clin. Endocr.
Metab. 86:

[168789]656–658, 2001.

[168790]

[168791]

[168792]17015.

[168793]Katyal, S. L.; Singh, G.; Locker, J.: Characterization of a
second

[168794]human pulmonary surfactant–associated protein SP–A
gene. Am. J. Resp.

[168795]Cell Molec. Biol. 6: 446–452, 1992.

[168796]

[168797]

[168798]17016.

[168799]Ramet, M.; Lofgren, J.; Albo, O.–P.; Hallman, M.: Surfactant
protein–A

[168800]gene locus associated with recurrent otitis media. J. Pe-
diat. 138:

[168801]266–268, 2001.

[168802]

[168803]

[168804]17017.

[168805]Wilson, C. L.; Ouellette, A. J.; Satchell, D. P.; Ayabe, T.;
Lopez–Boado,

[168806]Y. S.; Stratman, J. L.; Hultgren, S. J.; Matrisian, L. M.; Parks,

[168807]W. C.: Regulation of intestinal alpha–defensin activation
by the

[168808]metalloproteinase matrilysin in innate host defense. Sci-
ence 286:

[168809]113–117, 1999.

[168810]

[168811]

[168812]17018.

[168813]Robinson, P. R.; Cohen, G. B.; Zhukovsky, E. A.; Oprian, D.
D.

[168814]: Constitutively active mutants of rhodopsin. Neuron 9:
719–725,

[168815]1992.

[168816]

[168817]

[168818]17019.

[168819]Mason, I. J.; Murphy, D.; Munke, M.; Francke, U.; Elliott, R.
W.;

[168820]Hogan, B. L. M.: Developmental and transformation–sensi–
tive expression

[168821]of the SPARC gene on mouse chromosome 11. EMBO J. 5:
1831–1837,

[168822]1986.

[168823]

[168824]

[168825]17020.

[168826]Mason, I. J.; Taylor, A.; Williams, J. G.; Sage, H.; Hogan, B.

[168827]L. M.: Evidence from molecular cloning that SPARC, a major product

[168828]of mouse embryo parietal endoderm, is related to an endothelial cell

[168829]'culture shock' glycoprotein of Mr 43,000. EMBO J. 5: 1465–1472,

[168830]1986.

[168831]

[168832]

[168833]17021.

[168834]Naylor, S. L.; Helen–Davis, D.; Villarreal, X. C.; Long, G. L.

[168835]: The human osteonectin gene on chromosome 5 is polymorphic. (Abstract) Cytogenet.

[168836]Cell Genet. 51: 1051 only, 1989.

[168837]

[168838]

[168839]17022.

[168840]Schwartz, R. C.; Young, M. F.; Tsipouras, P.: Two RFLPs in
the
[168841]5-prime end of the human osteonectin (ON) gene. Nucleic
Acids Res. 16:
[168842]9076 only, 1988.

[168843]

[168844]

[168845]17023.

[168846]Stenner, D. D.; Tracy, R. P.; Riggs, B. L.; Mann, K. G.: Hu-
man
[168847]platelets contain and secrete osteonectin, a major protein
of mineralized
[168848]bone. Proc. Nat. Acad. Sci. 83: 6892-6896, 1986.

[168849]

[168850]

[168851]17024.

[168852]Swaroop, A.; Francke, U.: Molecular cloning, cDNA se-
quence, and
[168853]expression of human SPARC (osteonectin). (Abstract) Am.
J. Hum. Genet. 41:

[168854]A240 only, 1987.

[168855]

[168856]

[168857]17025.

[168858]Swaroop, A.; Hogan, B. L. M.; Francke, U.: Molecular analysis

[168859]of the cDNA for human SPARC/osteonectin/BM-40: sequence, expression,

[168860]and localization of the gene to chromosome 5q31-q33.

Genomics 2:

[168861]37-47, 1988.

[168862]

[168863]

[168864]17026.

[168865]Termine, J. D.; Kleinman, H. K.; Whitson, S. W.; Conn, K. M.;

[168866]McGarvey, M. L.; Martin, G. R.: Osteonectin, a bone-specific protein

[168867]linking mineral to collagen. Cell 26: 99-105, 1981.

[168868]

[168869]

[168870]17027.

[168871]Blau, N.; Thony, B.; Renneberg, A.; Arnold, L. A.; Hyland, K.:

[168872]Dihydropteridine reductase deficiency localized to the central nervous

[168873]system. J. Inherit. Metab. Dis. 21: 433–434, 1998.

[168874]

[168875]

[168876]17028.

[168877]Blau, N.; Thony, B.; Renneberg, A.; Penzien, J. M.; Hyland, K.;

[168878]Hoffmann, G.: Variant of dihydropteridine reductase deficiency without

[168879]hyperphenylalaninemia: effect of oral phenylalanine loading. J. Inherit.

[168880]Metab. Dis. 22: 216–220, 1999.

[168881]

[168882]

[168883]17029.

[168884]Bonafe, L.; Thony, B.; Penzien, J. M.; Czarnecki, B.; Blau, N.

[168885]: Mutations in the sepiapterin reductase gene cause a
novel tetrahydrobiopterin-dependent

[168886]monoamine-neurotransmitter deficiency without hyper-
phenylalaninemia. Am.

[168887]J. Hum. Genet. 69: 269–277, 2001.

[168888]

[168889]

[168890]17030.

[168891]Ichinose, H.; Katoh, S.; Sueoka, T.; Titani, K.; Fujita, K.;
Nagatsu,

[168892]T.: Cloning and sequencing of cDNA encoding human
sepiapterin reductase:

[168893]an enzyme involved in tetrahydrobiopterin biosynthesis.
Biochem.

[168894]Biophys. Res. Commun. 179: 183–189, 1991.

[168895]

[168896]

[168897]17031.

[168898]Murdoch, J. N.; Eddleston, J.; Stanier, P.; Copp, A. J.: Local–

ization

[168899]of the mouse gene encoding tyrosine kinase receptor type
10 on distal

[168900]chromosome 1. Mammalian Genome 8: 941–952, 1997.

[168901]

[168902]

[168903]17032.

[168904]Thony, B.; Heizmann, C. W.; Mattei, M.–G.: Human GTP–
cyclohydrolase

[168905]I gene and sepiapterin reductase gene map to region
14q21–q22 and

[168906]2p14–p12, respectively, by in situ hybridization. Ge–
nomics 26: 168–170,

[168907]1995.

[168908]

[168909]

[168910]17033.

[168911]Jin, H.; Oksenberg, D.; Ashkenazi, A.; Peroutka, S. J.; Dun–
can,

[168912]A. M. V.; Rozmahel, R.; Yang, Y.; Mengod, G.; Palacios, J.

M.; O'Dowd,

[168913]B. F.: Characterization of the human
5-hydroxytryptamine(1B) receptor. J.

[168914]Biol. Chem. 267: 5735–5738, 1992.

[168915]

[168916]

[168917]17034.

[168918]Levy, F. O.; Gudermann, T.; Birnbaumer, M.; Kaumann, A.
J.; Birnbaumer,

[168919]L.: Molecular cloning of a human gene (S31) encoding a
novel serotonin

[168920]receptor mediating inhibition of adenylyl cyclase. FEBS
Lett. 296:

[168921]201–206, 1992.

[168922]

[168923]

[168924]17035.

[168925]Levy, F. O.; Holtgreve-Grez, H.; Tasken, K.; Solberg, R.;
Ried,

[168926]T.; Gudermann, T.: Assignment of the gene encoding the

5-HT-1E serotonin

[168927]receptor (S31) (locus HTR1E) to human chromosome

6q14-q15. Genomics 22:

[168928]637-640, 1994.

[168929]

[168930]

[168931]17036.

[168932]McAllister, G.; Charlesworth, A.; Snodin, C.; Beer, M. S.;
Noble,

[168933]A. J.; Middlemiss, D. N.; Iversen, L. L.; Whiting, P.: Molecu-
lar

[168934]cloning of a serotonin receptor from human brain
(5HT1E): a fifth

[168935]5HT1-like subtype. Proc. Nat. Acad. Sci. 89: 5517-5521,
1992.

[168936]

[168937]

[168938]17037.

[168939]Weinshank, R. L.; Zgombick, J. M.; Macchi, M. J.; Branchek,
T.

[168940]A.; Hartig, P. R.: Human serotonin 1D receptor is encoded

by a subfamily

[168941]of two distinct genes: 5-HT(1D-alpha) and
5-HT(1D-beta). Proc. Nat.

[168942]Acad. Sci. 89: 3630-3634, 1992.

[168943]

[168944]

[168945]17038.

[168946]Evans, T.; Reitman, M.; Felsenfeld, G.: An erythrocyte-spe-
cific

[168947]DNA-binding factor recognizes a regulatory sequence
common to all

[168948]chicken globin genes. Proc. Nat. Acad. Sci. 85:
5976-5980, 1988.

[168949]

[168950]

[168951]17039.

[168952]Chen, Z.-Y.; Battinelli, E. M.; Fielder, A.; Bunday, S.; Sims,

[168953]K.; Breakefield, X. O.; Craig, I. W.: A mutation in the Norrie
disease

[168954]gene (NDP) associated with X-linked familial exudative

vitreoretinopathy. Nature

[168955]Genet. 5: 180–183, 1993.

[168956]

[168957]

[168958]17040.

[168959]Dudgeon, J.: Familial exudative vitreo–retinopathy. Trans.
Ophthal.

[168960]Soc. U.K. 99: 45–49, 1979.

[168961]

[168962]

[168963]17041.

[168964]Sinke, R. J.; de Leeuw, B.; Janssen, H. A. P.; Olde Weghuis,
D.;

[168965]Suijkerbuijk, R. F.; Meloni, A. M.; Gilgenkrantz, S.; Berger,
W.;

[168966]Ropers, H. H.; Sandberg, A. A.; Geurts van Kessel, A.: Lo-
calization

[168967]of X chromosome short arm markers relative to synovial
sarcoma– and

[168968]renal adenocarcinoma–associated translocation break–

points. Hum. Genet. 92:

[168969]305–308, 1993.

[168970]

[168971]

[168972]17042.

[168973]Smith, S.; Reeves, B. R.; Wong, L.; Fisher, C.: A consistent

[168974]chromosome translocation in synovial sarcoma. (Letter)

Cancer Genet.

[168975]Cytogenet. 26: 179–180, 1987.

[168976]

[168977]

[168978]17043.

[168979]Turc-Carel, C.; Dal Cin, P.; Limon, J.; Rao, U.; Li, F. P.; Cor-

son,

[168980]J. M.; Zimmerman, R.; Parry, D. M.; Cowan, J. M.; Sand-

berg, A. A.

[168981]: Involvement of chromosome X in primary cytogenetic
change in human

[168982]neoplasia: nonrandom translocation in synovial sarcoma.

Proc. Nat.

[168983]Acad. Sci. 84: 1981–1985, 1987.

[168984]

[168985]

[168986]17044.

[168987]Ueda, T.; Aozasa, K.; Tsujimoto, M.; Uchida, A.; Taniwaki, M.;

[168988]Abe, T.: Translocation X;18 and insertion 15;11 in a case of synovial

[168989]sarcoma. Cancer Genet. Cytogenet. 30: 183–185, 1988.

[168990]

[168991]

[168992]17045.

[168993]Wang–Wuu, S.; Soukup, S. W.; Lange, B. J.: Another synovial sarcoma

[168994]with t(X;18). Cancer Genet. Cytogenet. 29: 179–181, 1987.

[168995]

[168996]

[168997]17046.

[168998]Brennan, T. J.; Seeley, W. W.; Kilgard, M.; Schreiner, C. E.;
Tecott,

[168999]L. H.: Sound-induced seizures in serotonin 5-HT-2C re-
ceptor mutant

[169000]mice. Nature Genet. 16: 387-390, 1997.

[169001]

[169002]

[169003]17047.

[169004]Gurevich, I.; Tamir, H.; Arango, V.; Dwork, A. J.; Mann, J. J.;

[169005]Schmauss, C.: Altered editing of serotonin 2C receptor
pre-mRNA in

[169006]the prefrontal cortex of depressed suicide victims. Neuron
34: 349-356,

[169007]2002.

[169008]

[169009]

[169010]17048.

[169011]Hall, C. S.: Genetic differences in fatal audiogenic
seizures:

[169012]between two inbred strains of house mice. J. Hered. 38:

3-6, 1947.

[169013]

[169014]

[169015]17049.

[169016]Lappalainen, J.; Zhang, L.; Dean, M.; Oz, M.; Ozaki, N.; Yu, D.;

[169017]Virkkunen, M.; Weight, F.; Linnoila, M.; Goldman, D.: Identification,

[169018]expression, and pharmacology of a cys(23)-ser(23) substitution in

[169019]the human 5-HT(2C) receptor gene (HTR2C). Genomics 27: 274-279,

[169020]1995.

[169021]

[169022]

[169023]17050.

[169024]Milatovich, A.; Hsieh, C.-L.; Bonaminio, G.; Tecott, L.; Julius,

[169025]D.; Francke, U.: Serotonin receptor 1c gene assigned to X chromosome

[169026]in human (band q24) and mouse (bands D-F4). Hum.

Molec. Genet. 1:

[169027]681–684, 1992.

[169028]

[169029]

[169030]17051.

[169031]Tecott, L. H.; Sun, L. M.; Akana, S. F.; Strack, A. M.;
Lowenstein,

[169032]D. H.; Dallman, M. F.; Julius, D.: Eating disorder and
epilepsy in

[169033]mice lacking 5-HT_{2C} serotonin receptors. Nature 374:
542–546, 1995.

[169034]

[169035]

[169036]17052.

[169037]Blaschke, R. J.; Monaghan, A. P.; Schiller, S.; Schechinger,
B.;

[169038]Rao, E.; Padilla–Nash, H.; Ried, T.; Rappold, G. A.: SHOT, a
SHOX–related

[169039]homeobox gene, is implicated in craniofacial, brain, heart,
and limb

[169040]development. Proc. Nat. Acad. Sci. 95: 2406–2411, 1998.

[169041]

[169042]

[169043]17053.

[169044]Clement-Jones, M.; Schiller, S.; Rao, E.; Blaschke, R. J.; Zuniga,
niga,

[169045]A.; Zeller, R.; Robson, S. C.; Binder, G.; Glass, I.; Strachan,
T.;

[169046]Lindsay, S.; Rappold, G. A.: The short stature homeobox
gene SHOX

[169047]is involved in skeletal abnormalities in Turner syndrome.
Hum. Molec.

[169048]Genet. 9: 695-702, 2000.

[169049]

[169050]

[169051]17054.

[169052]Affara, N. A.; Chalmers, I. J.; Ferguson-Smith, M. A.: Anal-
ysis

[169053]of the SRY gene in 22 sex-reversed XY females identifies
four new

[169054]point mutations in the conserved DNA binding domain.
Hum. Molec.

[169055]Genet. 2: 785–789, 1993.

[169056]

[169057]

[169058]17055.

[169059]Behlke, M. A.; Bogan, J. S.; Beer–Romero, P.; Page, D. C.:
Evidence

[169060]that the SRY protein is encoded by a single exon on the
human Y chromosome. Genomics 17:

[169061]736–739, 1993.

[169062]

[169063]

[169064]17056.

[169065]Berkovitz, G. D.; Fechner, P. Y.; Zacur, H. W.; Rock, J. A.;
Snyder,

[169066]H. M., III; Migeon, C. J.; Perlman, E. J.: Clinical and patho-
logic

[169067]spectrum of 46,XY gonadal dysgenesis: its relevance to
the understanding

[169068]of sex differentiation. Medicine 70: 375–383, 1991.

[169069]

[169070]

[169071]17057.

[169072]Bowles, J.; Cooper, L.; Berkman, J.; Koopman, P.: Sry re-
quires

[169073]a CAG repeat domain for male sex determination in *Mus*
musculus. *Nature*

[169074]*Genet.* 22: 405–408, 1999.

[169075]

[169076]

[169077]17058.

[169078]Braun, A.; Kammerer, S.; Cleve, H.; Lohrs, U.; Schwarz,
H.–P.;

[169079]Kuhnle, U.: True hermaphroditism in a 46,XY individual,
caused by

[169080]a postzygotic somatic point mutation in the male gonadal
sex-determining

[169081]locus (SRY): molecular genetics and histological findings
in a sporadic

[169082]case. *Am. J. Hum. Genet.* 52: 578–585, 1993.

[169083]

[169084]

[169085]17059.

[169086]Brown, S.; Yu, C. C.; Lanzano, P.; Heller, D.; Thomas, L.;

Warburton,

[169087]D.; Kitajewski, J.; Stadtmauer, L.: A de novo mutation

(gln2stop)

[169088]at the 5-prime end of the SRY gene leads to sex reversal

with partial

[169089]ovarian function. (Letter) Am. J. Hum. Genet. 62: 189–192,

1998.

[169090]

[169091]

[169092]17060.

[169093]Canto, P.; de la Chesnaye, E.; Lopez, M.; Cervantes, A.;

Chavez,

[169094]B.; Vilchis, F.; Reyes, E.; Ulloa-Aguirre, A.; Kofman-Alfaro,

S.;

[169095]Mendez, J. P.: A mutation in the 5-prime non-high mobil-

ity group

[169096]box region of the SRY gene in patients with Turner syn-

drome and Y

[169097]mosaicism. J. Clin. Endocr. Metab. 85: 1908–1911, 2000.

[169098]

[169099]

[169100]17061.

[169101]Cherfas, J.: Sex and the single gene. Science 252: 782,
1991.

[169102]

[169103]

[169104]17062.

[169105]Clepet, C.; Schafer, A. J.; Sinclair, A. H.; Palmer, M. S.;
Lovell–Badge,

[169106]R.; Goodfellow, P. N.: The human SRY transcript. Hum.
Molec. Genet. 2:

[169107]2007–2012, 1993.

[169108]

[169109]

[169110]17063.

[169111]Davis, R. M.: Localisation of male determining factors in
man:

[169112]a thorough review of structural anomalies of the Y chromosome. J.

[169113]Med. Genet. 18: 161–195, 1981.

[169114]

[169115]

[169116]17064.

[169117]Dork, T.; Stuhrmann, M.; Miller, K.; Schmidtke, J.: Independent

[169118]observation of SRY mutation I90M in a patient with complete gonadal

[169119]dysgenesis. (Letter) Hum. Mutat. 11: 90–91, 1998.

[169120]

[169121]

[169122]17065.

[169123]Eicher, E. M.; Washburn, L. L.: Genetic control of primary sex

[169124]determination in mice. Annu. Rev. Genet. 20: 327–360, 1986.

[169125]

[169126]

[169127]17066.

[169128]Foster, J. W.; Brennan, F. E.; Hampikian, G. K.; Goodfellow, P.

[169129]N.; Sinclair, A. H.; Lovell-Badge, R.; Selwood, L.; Renfree, M. B.;

[169130]Cooper, D. W.; Graves, J. A. M.: Evolution of sex determination and

[169131]the Y chromosome: SRY-related sequences in marsupials. Nature 359:

[169132]531–533, 1992.

[169133]

[169134]

[169135]17067.

[169136]Fuqua, J. S.; McLaughlin, J.; Perlman, E. J.; Berkowitz, G. D.

[169137]: Analysis of the SRY gene in gonadal tissue of subjects with 46,XY

[169138]gonadal dysgenesis (Letter) J. Clin. Endocr. Metab. 82: 701–702,

[169139]1997.

[169140]

[169141]

[169142]17068.

[169143]Domenice, S.; Nishi, M. Y.; Billerbeck, A. E. C.; Latronico, A.

[169144]C.; Medeiros, M. A.; Russell, A. J.; Vass, K.; Carvalho, F. M.; Frade,

[169145]E. M. C.; Arnhold, I. J. P.; Mendonca, B. B.: A novel mis-sense mutation

[169146](S18N) in the 5-prime non-HMG box region of the SRY gene in a patient

[169147]with partial gonadal dysgenesis and his normal male relatives. Hum.

[169148]Genet. 102: 213-215, 1998.

[169149]

[169150]

[169151]17069.

[169152]Giese, K.; Pagel, J.; Grosschedl, R.: Distinct DNA-binding properties

[169153]of the high mobility group domain of murine and human SRY sex-determining

[169154]factors. Proc. Nat. Acad. Sci. 91: 3368-3372, 1994.

[169155]

[169156]

[169157]17070.

[169158]Goodfellow, P. N.; Lovell-Badge, R.: SRY and sex determination
nation

[169159]in mammals. Annu. Rev. Genet. 27: 71–92, 1993.

[169160]

[169161]

[169162]17071.

[169163]Gubbay, J.; Collignon, J.; Koopman, P.; Capel, B.;
Economou, A.;

[169164]Munsterberg, A.; Vivian, N.; Goodfellow, P. N.; Lovell-Badge, R.:

[169165]A gene mapping to the sex-determining region of the
mouse Y chromosome

[169166]is a member of a novel family of embryonically expressed
genes. Nature 346:

[169167]245–250, 1990.

[169168]

[169169]

[169170]17072.

[169171]Haqq, C. M.; King, C.-Y.; Ukiyama, E.; Falsafi, S.; Haqq, T.
N.;

[169172]Donahoe, P. K.; Weiss, M. A.: Molecular basis of mam-
malian sexual

[169173]determination: activation of Mullerian inhibiting substance
gene expression

[169174]by SRY. Science 266: 1494–1500, 1994.

[169175]

[169176]

[169177]17073.

[169178]Cannizzaro, L. A.; Dal Cin, P.; Mendez, M. J.; Sandberg, A.
A.

[169179]: Sublocalization of Xp breakpoint in the X;18 chromo-
some translocation

[169180]of synovial sarcoma. (Abstract) Cytogenet. Cell Genet. 51:
974,

[169181]1989.

[169182]

[169183]

[169184]17074.

[169185]Chand, A.; Clark, J.; Cooper, C. S.; Craig, I. W.: Long-range

[169186]organization of reiterated sequences, including the SSX1
cDNA, at

[169187]the OATL1 cluster in Xp11.23. Genomics 30: 545–552,

1995.

[169188]

[169189]

[169190]17075.

[169191]Clark, J.; Rocques, P. J.; Crew, A. J.; Gill, S.; Shipley, J.;

[169192]Chan, A. M.-L.; Gusterson, B. A.; Cooper, C. S.: Identifica-
tion of

[169193]novel genes, SYT and SSX, involved in the
t(X;18)(p11.2;q11.2) translocation

[169194]found in human synovial sarcoma. Nature Genet. 7:
502-508, 1994.

[169195]

[169196]

[169197]17076.

[169198]Enzinger, F. M.; Weiss, S. W.: Soft Tissue Tumors. St.
Louis:

[169199]C. V. Mosby (pub.) 1983. Pp. 519-549.

[169200]

[169201]

[169202]17077.

[169203]Gilgenkrantz, S.; Mujica, P.; Chery, M.; Gregoire, M.-J.;

Leotard,

[169204]B.; Mandel, J.-L.: Mapping the breakpoint at Xp11.2 in
synovial sarcoma.

[169205](Abstract) Cytogenet. Cell Genet. 51: 1004, 1989.

[169206]

[169207]

[169208]17078.

[169209]Griffin, C. A.; Emanuel, B. S.: Translocation (X;18) in a syn-
ovial

[169210]sarcoma. (Letter) Cancer Genet. Cytogenet. 26: 181-183,
1987.

[169211]

[169212]

[169213]17079.

[169214]Knight, J. C.; Martin, C. A.; Warren, W.; Smith, S.; West-
bury,

[169215]G.; Fisher, C.; Gusterson, B. A.; Cooper, C. S.; Reeves, B.
R.: Mapping

[169216]the translocation t(X;18)(p11.2;q11.2) in human synovial sarcoma.

[169217](Abstract) Cytogenet. Cell Genet. 51: 1023, 1989.

[169218]

[169219]

[169220]17080.

[169221]Knight, J. C.; Reeves, B. R.; Kearney, L.; Monaco, A. P.; Lehrach,

[169222]H.; Cooper, C. S.: Localization of the synovial sarcoma t(X;18)(p11.2;q11.2)

[169223]breakpoint by fluorescence in situ hybridization. Hum. Molec. Genet. 1:

[169224]633–637, 1992.

[169225]

[169226]

[169227]17081.

[169228]Miozzo, M.; Sozzi, G.; Cariani, C. T.; Di Palma, S.; Luksch, R.;

[169229]Azzarelli, A.; Pierotti, M. A.; Della Porta, G.: A synovial sarcoma

[169230]with t(X;18)(p11;q11) in a patient with Turner's syndrome.

Genes

[169231]Chromosomes Cancer 4: 192–193, 1992.

[169232]

[169233]

[169234]17082.

[169235]Yorifuji, T.; Wilson, R. W.; Beaudet, A. L.: Retroviral mediated

[169236]expression of CD18 in normal and deficient human bone marrow progenitor

[169237]cells. Hum. Molec. Genet. 2: 1443–1448, 1993.

[169238]

[169239]

[169240]17083.

[169241]Ramulu, P.; Kennedy, M.; Xiong, W.–H.; Williams, J.; Cowan, M.;

[169242]Blesh, D.; Yau, K.–W.; Hurley, J. B.; Nathans, J.: Normal light response,

[169243]photoreceptor integrity, and rhodopsin dephosphorylation in mice lacking

[169244]both protein phosphatases with EF hands (PPEF-1 and PPEF-2). Molec.

[169245]Cell. Biol. 21: 8605-8614, 2001.

[169246]

[169247]

[169248]17084.

[169249]van de Vosse, E.; Franco, B.; van der Bent, P.; Montini, E.; Orth,

[169250]U.; Hanauer, A.; Tijmes, N.; van Ommen, G.-J. B.; Ballabio, A.; den

[169251]Dunnen, J. T.; Bergen, A. A. B.: Exclusion of PPEF as the gene causing

[169252]X-linked juvenile retinoschisis. Hum. Genet. 101: 235-237, 1997.

[169253]

[169254]

[169255]17085.

[169256]Fisher, S. E.; Ciccodicola, A.; Tanaka, K.; Curci, A.; Desicato,

[169257]S.; D'Urso, M.; Craig, I. W.: Sequence-based exon prediction around

[169258]the synaptophysin locus reveals a gene-rich area containing novel

[169259]genes in human proximal Xp. Genomics 45: 340–347, 1997.

[169260]

[169261]

[169262]17086.

[169263]Oliva, M. M.; Wu, T. C.; Yang, V. W.: Isolation and characterization

[169264]of a differentiation-dependent gene in the human colonic cell line

[169265]HT29–18. Arch. Biochem. Biophys. 302: 183–192, 1993.

[169266]

[169267]

[169268]17087.

[169269]Prakash, S. K.; Paylor, R.; Jenna, S.; Lamarche–Vane, N.; Armstrong,

[169270]D. L.; Xu, B.; Mancini, M. A.; Zoghbi, H. Y.: Functional analysis

[169271]of ARHGAP6, a novel GTPase-activating protein for RhoA. Hum. Molec.

[169272]Genet. 9: 477–488, 2000.

[169273]

[169274]

[169275]17088.

[169276]Schaefer, L.; Prakash, S.; Zoghbi, H. Y.: Cloning and characterization

[169277]of a novel rho-type GTPase-activating protein gene (ARHGAP6) from

[169278]the critical region for microphthalmia with linear skin defects. Genomics 46:

[169279]268–277, 1997.

[169280]

[169281]

[169282]17089.

[169283]Aman, M. J.; Tayebi, N.; Obiri, N. I.; Puri, R. K.; Modi, W. S.;

[169284]Leonard, W. J.: cDNA cloning and characterization of the human interleukin

[169285]13 receptor alpha chain. J. Biol. Chem. 271: 29265–29270, 1996.

[169286]

[169287]

[169288]17090.

[169289]Guo, J.; Apiou , F.; Mellerin, M.-P.; Lebeau, B.; Jacques, Y.;

[169290]Minvielle, S.: Chromosome mapping and expression of the
human interleukin-13

[169291]receptor. Genomics 42: 141-5, 1997.

[169292]

[169293]

[169294]17091.

[169295]Hilton, D. J.; Zhang, J.-G.; Metcalf, D.; Alexander, W. S.;
Nicola,

[169296]N. A.; Willson, T. A.: Cloning and characterization of a
binding

[169297]subunit of the interleukin 13 receptor that is also a com-
ponent of

[169298]the interleukin 4 receptor. Proc. Nat. Acad. Sci. 93:
497-501, 1996.

[169299]

[169300]

[169301]17092.

[169302]Kaye, F. J.; Modi, S.; Ivanovska, I.; Koonin, E. V.; Thress, K.;

[169303]Kubo, A.; Kornbluth, S.; Rose, M. D.: A family of ubiquitin-like

[169304]proteins binds the ATPase domain of Hsp70-like Stch.
FEBS Lett. 467:

[169305]348–352, 2000.

[169306]

[169307]

[169308]17093.

[169309]Kaye, F. J.; Shows, T. B.: Assignment of ubiquilin 2
(UBQLN2)

[169310]to human chromosome xp11.23–p11.1 by GeneBridge radiation hybrids. Cytogenet.

[169311]Cell Genet. 89: 116–117, 2000.

[169312]

[169313]

[169314]17094.

[169315]Kleijnen, M. F.; Shih, A. H.; Zhou, P.; Kumar, S.; Soccio, R.
E.;

[169316]Kedersha, N. L.; Gill, G.; Howley, P. M.: The hPLIC proteins

may

[169317]provide a link between the ubiquitination machinery and the proteasome. Molec.

[169318]Cell 6: 409–419, 2000.

[169319]

[169320]

[169321]17095.

[169322]Marchand, J.-B.; Kaiser, D. A.; Pollard, T. D.; Higgs, H. N.:

[169323]Interaction of WASP/Scar proteins with actin and vertebrate Arp2/3

[169324]complex. Nature Cell Biol. 3: 76–82, 2001.

[169325]

[169326]

[169327]17096.

[169328]Lee, D. K.; Nguyen, T.; Lynch, K. R.; Cheng, R.; Vanti, W. B.;

[169329]Arkhitko, O.; Lewis, T.; Evans, J. F.; George, S. R.; O'Dowd, B. F.

[169330]: Discovery and mapping of ten novel G protein-coupled receptor genes. Gene 275:

[169331]83-91, 2001.

[169332]

[169333]

[169334]17097.

[169335]Furneau, H. M.; Rosenblum, M. K.; Dalmau, J.; Wong, E.;
Woodruff,

[169336]P.; Graus, F.; Posner, J. B.: Selective expression of Purk-
inje-cell

[169337]antigens in tumor tissue from patients with paraneoplastic
cerebellar

[169338]degeneration. New Eng. J. Med. 322: 1844-1851, 1990.

[169339]

[169340]

[169341]17098.

[169342]Furneau, H. M.; Wong, E.; Posner, J. B.: Isolation of cDNA
clones

[169343]encoding the major Yo paraneoplastic antigen. (Abstract)
Neurology 40

[169344](suppl. 1): 166 only, 1990.

[169345]

[169346]

[169347]17099.

[169348]Knight, J. C.; Renwick, P. J.; Downing, J. R.; Okuda, T.:

Physical

[169349]linkage of the cdc2-related gene (PCTK1) and the ubiquitin-activating

[169350]enzyme E1 gene (UBE1) on human Xp11.3. Cytogenet. Cell Genet. 71:

[169351]155-157, 1995.

[169352]

[169353]

[169354]17100.

[169355]Aarskog, D.: A familial syndrome of short stature associated with

[169356]facial dysplasia and genital anomalies. J. Pediat. 77: 856-861,

[169357]1970.

[169358]

[169359]

[169360]17101.

[169361]Baldellou, A.; Galve, L.; Bassecourt, M.: Risk of medullary damage

[169362]in Aarskog–Scott syndrome. (Abstract) Clin. Genet. 23: 225 only,

[169363]1983.

[169364]

[169365]

[169366]17102.

[169367]Bawle, E.; Tyrkus, M.; Lipman, S.; Bozimowski, D.: Aarskog syndrome:

[169368]full male and female expression associated with an X–autosome translocation. Am.

[169369]J. Med. Genet. 17: 595–602, 1984.

[169370]

[169371]

[169372]17103.

[169373]Berman, P. A.; Desjardins, C.; Fraser, F. C.: Inheritance of the

[169374]Aarskog syndrome. Birth Defects Orig. Art. Ser. X(7): 151–159, 1974.

[169375]

[169376]

[169377]17104.

[169378]Buechner, M.; Hall, D. H.; Bhatt, H.; Hedgecock, E. M.: Cys-
tic

[169379]canal mutants in *Caenorhabditis elegans* are defective in
the apical

[169380]membrane domain of the renal (excretory) cell. *Dev. Biol.*
214: 227–241,

[169381]1999.

[169382]

[169383]

[169384]17105.

[169385]Casteels, M.; Samain, H.; Penninckx, F.; Coremans, G.;
Beirinckx,

[169386]J.; Fryns, J. P.: Megadolichosigmoid in a young male with
Aarskog

[169387]syndrome. *Genet. Counsel.* 5: 81–83, 1994.

[169388]

[169389]

[169390]17106.

[169391]Estrada, L.; Caron, E.; Gorski, J. L.: Fgd1, the Cdc42 guanine
nine

[169392]nucleotide exchange factor responsible for faciogenital
dysplasia,

[169393]is localized to the subcortical actin cytoskeleton and Golgi
membrane. Hum.

[169394]Molec. Genet. 10: 485–495, 2001.

[169395]

[169396]

[169397]17107.

[169398]Fernandez, I.; Tsukahara, M.; Mito, H.; Yoshii, H.; Uchida,
M.;

[169399]Matsuo, K.; Kajii, T.: Congenital heart defects in Aarskog
syndrome. Am.

[169400]J. Med. Genet. 50: 318–322, 1994.

[169401]

[169402]

[169403]17108.

[169404]Fryns, J.-P.: Dolichomegasigmoid in Aarskog syndrome.

(Letter) Am.

[169405]J. Med. Genet. 45: 122 only, 1993.

[169406]

[169407]

[169408]17109.

[169409]Fryns, J. P.: Aarskog syndrome: the changing phenotype
with age. Am.

[169410]J. Med. Genet. 43: 420–427, 1992.

[169411]

[169412]

[169413]17110.

[169414]Fryns, J. P.; Macken, J.; Vinken, L.; Igodt–Ameye, L.; van
den

[169415]Berghe, H.: The Aarskog syndrome. Hum. Genet. 42:
129–135, 1978.

[169416]

[169417]

[169418]17111.

[169419]Funderburk, S. J.; Crandall, B. F.: The Aarskog syndrome in

three

[169420]brothers. Clin. Genet. 6: 119–124, 1974.

[169421]

[169422]

[169423]17112.

[169424]Furukawa, C. T.; Hall, B. D.; Smith, D. W.: The Aarskog
syndrome. J.

[169425]Pediat. 81: 1117–1122, 1972.

[169426]

[169427]

[169428]17113.

[169429]Gao, J.; Estrada, L.; Cho, S.; Ellis, R. E.; Gorski, J. L.: The

[169430]Caenorhabditis elegans homolog of FGD1, the human
Cdc42 GEF gene responsible

[169431]for faciogenital dysplasia, is critical for excretory cell mor-
phogenesis. Hum.

[169432]Molec. Genet. 10: 3049–3062, 2001.

[169433]

[169434]

[169435]17114.

[169436]Glover, T. W.; Verga, V.; Rafael, J.; Barcroft, C.; Gorski, J.

[169437]L.; Bawle, E. V.; Higgins, J. V.: Translocation breakpoint in
Aarskog

[169438]syndrome maps to Xp11.21 between ALAS2 and DXS323.

Hum. Molec. Genet. 2:

[169439]1717–1718, 1993.

[169440]

[169441]

[169442]17115.

[169443]Grier, R. E.; Farrington, F. H.; Kendig, R.; Mamunes, P.:

Autosomal

[169444]dominant inheritance of the Aarskog phenotype.

(Abstract) Am. J.

[169445]Hum. Genet. 33: 64A only, 1981.

[169446]

[169447]

[169448]17116.

[169449]Hoo, J. J.: The Aarskog (facio–digito–genital) syndrome.

Clin.

[169450]Genet. 16: 269–276, 1979.

[169451]

[169452]

[169453]17117.

[169454]Kodama, M.; Fujimoto, S.; Namikawa, T.; Matsuda, I.:

Aarskog

[169455]syndrome with isolated growth hormone deficiency. Europ. J. Pediat. 135:

[169456]273–276, 1981.

[169457]

[169458]

[169459]17118.

[169460]Lebel, R. R.; May, M.; Pouls, S.; Lubs, H. A.; Stevenson, R. E.;

[169461]Schwartz, C. E.: Non-syndromic X-linked mental retardation associated

[169462]with a missense mutation (P312L) in the FGD1 gene. Clin. Genet. 61:

[169463]139–145, 2002.

[169464]

[169465]

[169466]17119.

[169467]Logie, L. J.; Porteous, M. E. M.: Intelligence and development

[169468]in Aarskog syndrome. Arch. Dis. Child. 79: 359–360, 1998.

[169469]

[169470]

[169471]17120.

[169472]Mikelsaar, R. V.–A.; Lurie, I. W.: Atypical case of Aarskog syndrome. J.

[169473]Med. Genet. 29: 349–350, 1992.

[169474]

[169475]

[169476]17121.

[169477]Nielsen, K. B.: Aarskog syndrome in a Danish family: an illustration

[169478]of the need for dysmorphology in paediatrics. Clin. Genet.

33: 315–317,

[169479]1988.

[169480]

[169481]

[169482]17122.

[169483]Oberiter, V.; Lovrencic, M. K.; Schmutzer, L.; Kraus, O.:

The

[169484]Aarskog syndrome. *Acta Paediat. Scand.* 69: 567–570,
1980.

[169485]

[169486]

[169487]17123.

[169488]Orrico, A.; Galli, L.; Falciani, M.; Bracci, M.; Cavaliere, M.

[169489]L.; Rinaldi, M. M.; Musacchio, A.; Sorrentino, V.: A muta-
tion in

[169490]the pleckstrin homology (PH) domain of the FGD1 gene in
an Italian

[169491]family with faciogenital dysplasia (Aarskog–Scott syn-
drome). *FEBS*

[169492]*Lett.* 478: 216–220, 2000.

[169493]

[169494]

[169495]17124.

[169496]Pasteris, N. G.; Cadle, A.; Logie, L. J.; Porteous, M. E. M.;

[169497]Schwartz, C. E.; Stevenson, R. E.; Glover, T. W.; Wilroy, R.

S.; Gorski,

[169498]J. L.: Isolation and analysis of the faciogenital dysplasia

(Aarskog-Scott

[169499]syndrome) gene: a putative, rho/rac guanine nucleotide

exchange factor. Cell 79:

[169500]669-678, 1994.

[169501]

[169502]

[169503]17125.

[169504]Pasteris, N. G.; de Gouyon, B.; Cadle A. B.; Campbell, K.;

Herman,

[169505]G. E.; Gorski, J. L.: Cloning and regional localization of the

mouse

[169506]faciogenital dysplasia (Fgd1) gene. Mammalian Genome 6:

658-661,

[169507]1995.

[169508]

[169509]

[169510]17126.

[169511]Pedersen, J. C.; Fryns, J. P.; Bracke, P.; Geeraert, M.; Van
Den

[169512]Berghe, H.: The Aarskog syndrome. Ann. Genet. 23:
108–110, 1980.

[169513]

[169514]

[169515]17127.

[169516]Porteous, M. E. M.; Goudie, D. R.: Aarskog syndrome. J.
Med.

[169517]Genet. 28: 44–47, 1991.

[169518]

[169519]

[169520]17128.

[169521]Rafael, J.; Verga, V.; Hall, B.; Burrigh, E.; Gorski, J.; Bawle,

[169522]J.; Higgins, J. V.; Glover, T. W.: Assignment of the translo-

cation

[169523]breakpoint in a patient with Aarskog syndrome to
Xp11.21. (Abstract) Am.

[169524]J. Hum. Genet. 51 (suppl.): A116 only, 1992.

[169525]

[169526]

[169527]17129.

[169528]Schwartz, C. E.; Gillesen-Kaesbach, G.; May, M.; Cappa,
M.; Gorski,

[169529]J.; Steindl, K.; Neri, G.: Two novel mutations confirm FGD1
is responsible

[169530]for the Aarskog syndrome. Europ. J. Hum. Genet. 8:
869-874, 2000.

[169531]

[169532]

[169533]17130.

[169534]Scott, C. I., Jr.: Unusual facies, joint hypermobility, genital

[169535]anomaly and short stature: a new dysmorphic syndrome.

Birth Defects

[169536]Orig. Art. Ser. VII(6): 240-246, 1971.

[169537]

[169538]

[169539]17131.

[169540]Demarest, S. J.; Martinez-Yamout, M.; Chung, J.; Chen, H.;
Xu,

[169541]W.; Dyson, H. J.; Evans, R. M.; Wright, P. E.: Mutual syner-
gistic

[169542]folding in recruitment of CBP/p300 by p160 nuclear re-
ceptor coactivators. Nature 415:

[169543]549–553, 2002.

[169544]

[169545]

[169546]17132.

[169547]Eckner, R.; Ewen, M. E.; Newsome, D.; Gerdes, M.; De-
Caprio, J.

[169548]A.; Lawrence, J. B.; Livingston, D. M.: Molecular cloning
and functional

[169549]analysis of the adenovirus E1A-associated 300-kD protein
(p300) reveals

[169550]a protein with properties of a transcriptional adaptor.
Genes Dev. 15:

[169551]869–884, 1994.

[169552]

[169553]

[169554]17133.

[169555]Lin, C. H.; Hare, B. J.; Wagner, G.; Harrison, S. C.; Maniatis,

[169556]T.; Fraenkel, E.: A small domain of CBP/p300 binds diverse proteins:

[169557]solution structure and functional studies. Molec. Cell 8: 581–590,

[169558]2001.

[169559]

[169560]

[169561]17134.

[169562]Cohen–Haguenauer, O.; Picard, J. Y.; Mattei, M.–G.; Serero, S.;

[169563]Van Cong, N.; de Tand, M.–F.; Guerrier, D.; Hors–Cayla, M.–C.; Josso,

[169564]N.; Frezal, J.: Mapping of the gene for anti–mullerian hormone to

[169565]the short arm of human chromosome 19. Cytogenet. Cell

Genet. 44:

[169566]2–6, 1987.

[169567]

[169568]

[169569]17135.

[169570]Forest, M. G.: Serum Mullerian inhibiting substance assay
-- a

[169571]new diagnostic test for disorders of gonadal development.
(Editorial) New

[169572]Eng. J. Med. 336: 1519–1521, 1997.

[169573]

[169574]

[169575]17136.

[169576]Cate, R. L.; Mattaliano, R. J.; Hession, C.; Tizard, R.; Far-
ber,

[169577]N. M.; Cheung, A.; Ninfa, E. G.; Frey, A. Z.; Gash, D. J.;
Chow, E.

[169578]P.; Fisher, R. A.; Bertonis, J. M.; Torres, G.; Wallner, B. P.;
Ramachandran,

[169579]K. L.; Ragin, R. C.; Manganaro, T. F.; MacLaughlin, D. T.;

Donahoe,

[169580]P. K.: Isolation of the bovine and human genes for mullerian inhibiting

[169581]substance and expression of the human gene in animal cells. Cell 45:

[169582]685–698, 1986.

[169583]

[169584]

[169585]17137.

[169586]Guerrier, D.; Tran, D.; Vanderwinden, J. M.; Hideux, S.; Van Outryve,

[169587]L.; Legeai, L.; Bouchard, M.; Van Vliet, G.; De Laet, M. H.; Picard,

[169588]J. Y.; Kahn, A.; Josso, N.: The persistent mullerian duct syndrome:

[169589]a molecular approach. J. Clin. Endocr. Metab. 68: 46–52, 1989.

[169590]

[169591]

[169592]17138.

[169593]Harbison, M. D.; Magid, M. L. S.; Josso, N.; Mininberg, D.

T.;

[169594]New, M. I.: Anti-Mullerian hormone in three intersex conditions. Ann.

[169595]Genet. 34: 226-232, 1991.

[169596]

[169597]

[169598]17139.

[169599]King, T. R.; Lee, B. K.; Behringer, R. R.; Eicher, E. M.: Mapping
ping

[169600]anti-Mullerian hormone (Amh) and related sequences in
the mouse: identification

[169601]of a new region of homology between MMU10 and
HSA19p. Genomics 11:

[169602]273-283, 1991.

[169603]

[169604]

[169605]17140.

[169606]Knebelmann, B.; Boussin, L.; Guerrier, D.; Legeai, L.; Kahn,
A.;

[169607]Josso, N.; Picard, J.-Y.: Anti-Mullerian hormone Bruxelles:

a nonsense

[169608]mutation associated with the persistent Mullerian duct syndrome. Proc.

[169609]Nat. Acad. Sci. 88: 3767–3771, 1991.

[169610]

[169611]

[169612]17141.

[169613]Lee, M. M.; Donahoe, P. K.; Silverman, B. L.; Hasegawa, T.;
Hasegawa,

[169614]Y.; Gustafson, M. L.; Chang, Y.; MacLaughlin, D. T.: Measurements
surements

[169615]of serum Mullerian inhibiting substance in the evaluation
of children

[169616]with nonpalpable gonads. New Eng. J. Med. 336:
1480–1486, 1997.

[169617]

[169618]

[169619]17142.

[169620]Rey, R. A.; Belville, C.; Nihoul-Fekete, C.; Michel-Cale-
mard,

[169621]L.; Forest, M. G.; Lahlou, N.; Jaubert, F.; Mowszowicz, I.;

David,

[169622]M.; Saka, N.; Bouvattier, C.; Bertrand, A.-M.; and 16 others: Evaluation

[169623]of gonadal function in 107 intersex patients by means of serum antimullerian

[169624]hormone measurement. J. Clin. Endocr. Metab. 84: 627-631, 1999.

[169625]

[169626]

[169627]17143.

[169628]Rogers, D. S.; Gallagher, D. S.; Womack, J. E.: Somatic cell

[169629]mapping of the genes for anti-mullerian hormone and osteonectin in

[169630]cattle: identification of a new bovine syntenic group. Genomics 9:

[169631]298-300, 1991.

[169632]

[169633]

[169634]17144.

[169635]Riesewijk, A. M.; Blagitko, N.; Schinzel, A. A.; Hu, L.;

Schulz,

[169636]U.; Hamel, B. C. J.; Ropers, H.-H.; Kalscheuer, V. M.: Evidence against

[169637]a major role of PEG1/MEST Silver-Russell syndrome. Europ. J. Hum.

[169638]Genet. 6: 114-120, 1998.

[169639]

[169640]

[169641]17145.

[169642]Riesewijk, A. M.; Hu, L.; Schulz, U.; Tariverdian, G.; Hoglund,

[169643]P.; Kere, J.; Ropers, H.-H.; Kalscheuer, V. M.: Monoallelic expression

[169644]of human PEG1/MEST is paralleled by parent-specific methylation in

[169645]fetuses. Genomics 42: 236-244, 1997.

[169646]

[169647]

[169648]17146.

[169649]Sado, T.; Nakajima, N.; Tada, M.; Takagi, N.: A novel

mesoderm-specific

[169650]cDNA isolated from a mouse embryonal carcinoma cell
line. Dev. Growth

[169651]Differ. 35: 551-560, 1993.

[169652]

[169653]

[169654]17147.

[169655]Jayakumar, A.; Chirala, S. S.; Chinault, A. C.; Baldini, A.;
Abu-Elheiga,

[169656]L.; Wakil, S. J.: Isolation and chromosomal mapping of ge-
nomic clones

[169657]encoding the human fatty acid synthase gene. Genomics
23: 420-424,

[169658]1994.

[169659]

[169660]

[169661]17148.

[169662]Jayakumar, A.; Tai, M.-H.; Huang, W.-Y.; Al-Feel, W.; Hsu,
M.;

[169663]Abu-Elheiga, L.; Chirala, S. S.; Wakil, S. J.: Human fatty

acid synthase:

[169664]properties and molecular cloning. Proc. Nat. Acad. Sci. 92:
8695–8699,

[169665]1995.

[169666]

[169667]

[169668]17149.

[169669]Loftus, T. M.; Jaworsky, D. E.; Frehywot, G. L.; Townsend,
C. A.;

[169670]Ronnett, G. V.; Lane, M. D.; Kuhajda, F. P.: Reduced food
intake

[169671]and body weight in mice treated with fatty acid synthase
inhibitors. Science 288:

[169672]2379–2381, 2000.

[169673]

[169674]

[169675]17150.

[169676]Wakil, S. J.: Fatty acid synthase, a proficient multifunc-
tional

[169677]enzyme. Biochemistry 28: 4523–4530, 1989.

[169678]

[169679]

[169680]17151.

[169681]Ye, Q.; Chung, L. W. K.; Li, S.; Zhau, H. E.: Identification of

[169682]a novel FAS/ER- α fusion transcript expressed in human cancer cells. *Biochim.*

[169683]*Biophys. Acta* 1493: 373–377, 2000.

[169684]

[169685]

[169686]17152.

[169687]Hofmann, M. A.; Drury, S.; Fu, C.; Qu, W.; Taguchi, A.; Lu, Y.;

[169688]Avila, C.; Kambham, N.; Bierhaus, A.; Nawroth, P.; Neurath, M. F.;

[169689]Slattery, T.; Beach, D.; McClary, J.; Nagashima, M.; Morser, J.; Stern,

[169690]D.; Schmidt, A. M.: RAGE mediates a novel proinflammatory axis: a

[169691]central cell surface receptor for S100/calgranulin polypeptides. *Cell* 97:

[169692]889–901, 1999.

[169693]

[169694]

[169695]17153.

[169696]Yan, S. D.; Chen, X.; Fu, J.; Chen, M.; Zhu, H.; Roher, A.;
Slattery,

[169697]T.; Zhao, L.; Nagashima, M.; Morser, J.; Migheli, A.;
Nawroth, P.;

[169698]Stern, D.; Schmidt, A. M.: RAGE and amyloid–beta peptide
neurotoxicity

[169699]in Alzheimer's disease. Nature 382: 685–691, 1996.

[169700]

[169701]

[169702]17154.

[169703]Argeson, A. C.; Druck, T.; Veronese, M. L.; Knopf, J. L.;
Buchberg,

[169704]A. M.; Huebner, K.; Siracusa, L. D.: Phospholipase C
gamma–2 (Plcg2)

[169705]and phospholipase C gamma–1 (Plcg1) map to distinct re-
gions in the

[169706]human and mouse genomes. Genomics 25: 29–35, 1995.

[169707]

[169708]

[169709]17155.

[169710]Hernandez, D.; Egan, S. E.; Yulug, I. G.; Fisher, E. M. C.:

Mapping

[169711]the gene that encodes phosphatidylinositol-specific phospholipase

[169712]C-gamma-2 in the human and the mouse. Genomics 23: 504–507, 1994.

[169713]

[169714]

[169715]17156.

[169716]Kang, J. S.; Kohlhuber, F.; Hug, H.; Marme, D.; Eick, D.;

Ueffing,

[169717]M.: Cloning and functional analysis of the hematopoietic cell-specific

[169718]phospholipase C-gamma-2 promoter. FEBS Lett. 399: 14–20, 1996.

[169719]

[169720]

[169721]17157.

[169722]Dumont, D. J.; Anderson, L.; Breitman, M. L.; Duncan, A.
M. V.

[169723]: Assignment of the endothelial-specific protein receptor
tyrosine

[169724]kinase gene (TEK) to human chromosome 9p21. Genomics
23: 512-513,

[169725]1994.

[169726]

[169727]

[169728]17158.

[169729]Delot, E.; King, L. M.; Briggs, M. D.; Wilcox, W. R.; Cohn, D.

[169730]H.: Trinucleotide expansion mutations in the cartilage
oligomeric

[169731]matrix protein (COMP) gene. Hum. Molec. Genet. 8:
123-128, 1999.

[169732]

[169733]

[169734]17159.

[169735]Back, A. L.; Kerkerling, M.; Baker, D.; Bauer, T. R.; Embree, L.

[169736]J.; Hickstein, D. D.: A point mutation associated with leukocyte

[169737]adhesion deficiency type 1 of moderate severity. Biochem. Biophys.

[169738]Res. Commun. 193: 912–918, 1993.

[169739]

[169740]

[169741]17160.

[169742]Back, A. L.; Kwok, W. W.; Hickstein, D. D.: Identification of

[169743]two molecular defects in a child with leukocyte adherence deficiency. J.

[169744]Biol. Chem. 267: 5482–5487, 1992.

[169745]

[169746]

[169747]17161.

[169748]Barclay, A. N.; Birkeland, M. L.; Brown, M. H.; Beyers, A. D.;

[169749]Davis, S. J.; Somoza, C.; Williams, A. F.: The Leukocyte

Antigen

[169750]Facts Book. New York: Academic Press (pub.) 1993. Pp.
124–127

[169751]and 140–141.

[169752]

[169753]

[169754]17162.

[169755]Beatty, P. G.; Ochs, H. D.; Harlan, J. M.; Price, T. H.; Rosen,
[169756]H.; Taylor, R. F.; Hansen, J. A.; Klebanoff, S. J.: Absence of
monoclonal–antibody–defined

[169757]protein complex in a boy with abnormal leucocyte func-
tion. Lancet I:

[169758]535–537, 1984.

[169759]

[169760]

[169761]17163.

[169762]Bianchi, E.; Denti, S.; Granata, A.; Bossi, G.; Geginat, J.;
Villa,

[169763]A.; Rogge, L.; Pardi, R.: Integrin LFA–1 interacts with the
transcriptional

[169764]co-activator JAB1 to modulate AP-1 activity. Nature 404:
617-621,
[169765]2000.

[169766]

[169767]

[169768]17164.

[169769]Bissenden, J. G.; Haeney, M. R.; Tarlow, M. J.; Thompson,
R. A.

[169770]: Delayed separation of the umbilical cord, severe
widespread infections,

[169771]and immunodeficiency. Arch. Dis. Child. 56: 397-399,
1981.

[169772]

[169773]

[169774]17165.

[169775]Boucheix, C.: Personal Communication. Villejuif, France
1/31/1987.

[169776]

[169777]

[169778]17166.

- [169779]Bowen, T. J.; Ochs, H. D.; Altman, L. C.; Price, T. H.; Van Epps,
[169780]D. E.; Brautigan, D. L.; Rosin, R. E.; Perkins, W. D.; Babior,
B.
[169781]M.; Klebanoff, S. J.; Wedgwood, R. J.: Severe recurrent bac-
terial
[169782]infections associated with defective adherence and
chemotaxis in two
[169783]patients with neutrophils deficient in a cell-associated
glycoprotein. J.
[169784]Pediat. 101: 932–940, 1982.
- [169785]
- [169786]
- [169787]17167.
- [169788]Bullard, D. C.; Scharffetter–Kochanek, K.; McArthur, M. J.;
Chosay,
[169789]J. G.; McBride, M. E.; Montgomery, C. A.; Beaudet, A. L.: A
polygenic
[169790]mouse model of psoriasiform skin disease in
CD18-deficient mice. Proc.
[169791]Nat. Acad. Sci. 93: 2116–2121, 1996.

[169792]

[169793]

[169794]17168.

[169795]Crowley, C. A.; Curnutte, J. T.; Rosin, R. E.; Andre-
Schwartz,

[169796]J.; Gallin, J. I.; Klempner, M.; Snyderman, R.; Southwick, F.
S.;

[169797]Stossel, T. P.; Babior, B. M.: An inherited abnormality of
neutrophil

[169798]adhesion: its genetic transmission and its association with
a missing

[169799]protein. New Eng. J. Med. 302: 1163–1168, 1980.

[169800]

[169801]

[169802]17169.

[169803]Dana, N.; Clayton, L. K.; Tennen, D. G.; Pierce, M. W.;
Lachmann,

[169804]P. J.; Law, S. A.; Arnaout, M. A.: Leukocytes from four pa-
tients

[169805]with complete or partial Leu–CAM deficiency contain the
common beta–subunit

[169806]precursor and beta-subunit messenger RNA. J. Clin. Invest. 79: 1010-1015,
[169807]1987.

[169808]

[169809]

[169810]17170.

[169811]Dana, N.; Todd, R. F., III; Pitt, J.; Springer, T. A.; Arnaout,

[169812]M. A.: Deficiency of a surface membrane glycoprotein
(Mo1) in man. J.

[169813]Clin. Invest. 73: 153-159, 1984.

[169814]

[169815]

[169816]17171.

[169817]Fischer, A.; Griscelli, C.; Friedrich, W.; Kubanek, B.; Levin-
sky,

[169818]R.; Morgan, G.; Vossen, J.; Wagemaker, G.; Landais, P.:
Bone-marrow

[169819]transplantation for immunodeficiencies and osteopetrosis:
European

[169820]survey, 1968-1985. Lancet II: 1080-1084, 1986.

[169821]

[169822]

[169823]17172.

[169824]Fujita, K.; Kobayashi, K.; Kajii, T.: Impaired neutrophil adhesion:

[169825]a new patient in a previously reported family. Acta Paediat. Jpn. 27:

[169826]527–534, 1985.

[169827]

[169828]

[169829]17173.

[169830]Fujita, K.; Kobayashi, K.; Okino, F.: Juvenile rheumatoid arthritis

[169831]in two siblings with congenital leucocyte adhesion deficiency. Europ.

[169832]J. Pediat. 148: 118–119, 1988.

[169833]

[169834]

[169835]17174.

[169836]Harvath, L.; Andersen, B. R.: Defective initiation of oxidative

[169837]metabolism in polymorphonuclear leukocytes. New Eng. J. Med. 300:

[169838]1130–1135, 1979.

[169839]

[169840]

[169841]17175.

[169842]Hayward, A. R.; Leonard, J.; Harvey, B. A. M.; Greenwood, M. C.;

[169843]Wood, C. B. S.; Soothill, J. F.: Delayed separation of the umbilical

[169844]cord, widespread infections, and defective neutrophil mobility. Lancet I:

[169845]1099–1101, 1979.

[169846]

[169847]

[169848]17176.

[169849]Hibbs, M. L.; Wardlaw, A. J.; Stacker, S. A.; Anderson, D. C.;

[169850]Lee, A.; Roberts, T. M.; Springer, T. A.: Transfection of cells from

[169851]patients with leukocyte adhesion deficiency with an integrin beta

[169852]subunit (CD18) restores lymphocyte function-associated antigen-1 expression

[169853]and function. J. Clin. Invest. 85: 674-681, 1990.

[169854]

[169855]

[169856]17177.

[169857]Hogg, N.; Stewart, M. P.; Scarth, S. L.; Newton, R.; Shaw, J.

[169858]M.; Law, S. K. A.; Klein, N.: A novel leukocyte adhesion deficiency

[169859]caused by expressed but nonfunctional beta-2 integrins Mac-1 and LFA-1. J.

[169860]Clin. Invest. 103: 97-106, 1999.

[169861]

[169862]

[169863]17178.

[169864]Hynes, R. O.: Integrins: versatility, modulation and signal-

ing

[169865]in cell adhesion. Cell 69: 11–25, 1992.

[169866]

[169867]

[169868]17179.

[169869]Kehrli, M. E., Jr.; Ackermann, M. R.; Shuster, D. E.; van der

[169870]Maaten, M. J.; Schmalstieg, F. C.; Anderson, D. C.; Hughes,
B. J.

[169871]: Bovine leukocyte adhesion deficiency: beta(2) integrin
deficiency

[169872]in young Holstein cattle. Am. J. Path. 140: 1489–1492,
1992.

[169873]

[169874]

[169875]17180.

[169876]Kishimoto, T. K.; Hollander, N.; Roberts, T. M.; Anderson,
D.

[169877]C.; Springer, T. A.: Heterogeneous mutations in the beta
subunit

[169878]common to the LFA–1, Mac–1, and p150,95 glycoproteins

cause leukocyte

[169879]adhesion deficiency. Cell 50: 193–202, 1987.

[169880]

[169881]

[169882]17181.

[169883]Kishimoto, T. K.; O'Connor, K.; Lee, A.; Roberts, T. M.;
Springer,

[169884]T. A.: Cloning of the beta subunit of the leukocyte adhe-
sion proteins:

[169885]homology to an extracellular matrix receptor defines a
novel supergene

[169886]family. Cell 48: 681–690, 1987.

[169887]

[169888]

[169889]17182.

[169890]Kobayashi, K.; Fujita, K.; Okino, F.; Kajii, T.: An abnormal-
ity

[169891]of neutrophil adhesion: autosomal recessive inheritance
associated

[169892]with missing neutrophil glycoproteins. Pediatrics 73:

606–610, 1984.

[169893]

[169894]

[169895]17183.

[169896]Krauss, J. C.; Mayo–Bond, L. A.; Rogers, C. E.; Weber, K. L.;

[169897]Todd, R. F., III; Wilson, J. M.: An in vivo animal model of
gene

[169898]therapy for leukocyte adhesion deficiency. J. Clin. Invest.
88:

[169899]1412–1417, 1991.

[169900]

[169901]

[169902]17184.

[169903]Matsuura, S.; Kishi, F.; Tsukahara, M.; Nunoi, H.; Matsuda,
I.;

[169904]Kobayashi, K.; Kajii, T.: Leukocyte adhesion deficiency:
identification

[169905]of novel mutations in two Japanese patients with a severe
form. Biochem.

[169906]Biophys. Res. Commun. 184: 1460–1467, 1992.

[169907]

[169908]

[169909]17185.

[169910]Nelson, C.; Rabb, H.; Arnaout, M. A.: Genetic cause of
leukocyte

[169911]adhesion molecule deficiency: abnormal splicing and a
missense mutation

[169912]in a conserved region of CD18 impair cell surface expres-
sion of beta-2

[169913]integrins. J. Biol. Chem. 267: 3351-3357, 1992.

[169914]

[169915]

[169916]17186.

[169917]Tini, M.; Benecke, A.; Um, S.-J.; Torchia, J.; Evans, R. M.;
Chambon,

[169918]P.: Association of CBP/p300 acetylase and thymine DNA
glycosylase

[169919]links DNA repair and transcription. Molec. Cell 9:
265-277, 2002.

[169920]

[169921]

[169922]17187.

[169923]Weaver, B. K.; Kumar, K. P.; Reich, N. C.: Interferon regulatory

[169924]factor 3 and CREB-binding protein/p300 are subunits of double-stranded

[169925]RNA-activated transcription factor DRAF1. Molec. Cell. Biol. 18:

[169926]1359–1368, 1998.

[169927]

[169928]

[169929]17188.

[169930]Hansen, J. J.; Durr, A.; Cournu-Rebeix, I.; Georgopoulos, C.; Ang,

[169931]D.; Nielsen, M. N.; Davoine, C.-S.; Brice, A.; Fontaine, B.; Gregersen,

[169932]N.; Bross, P.: Hereditary spastic paraplegia SPG13 is associated

[169933]with a mutation in the gene encoding the mitochondrial chaperonin

[169934]Hsp60. Am. J. Hum. Genet. 70: 1328–1332, 2002.

[169935]

[169936]

[169937]17189.

[169938]Todd, M. J.; Viitanen, P. V.; Lorimer, G. H.: Dynamics of the

[169939]chaperonin ATPase cycle: implications for facilitated protein folding. Science 265:

[169940]659–666, 1994.

[169941]

[169942]

[169943]17190.

[169944]Hellevuo, K.; Berry, R.; Sikela, J. M.; Tabakoff, B.: Localization

[169945]of the gene for a novel human adenylyl cyclase (ADCY7) to chromosome

[169946]16. Hum. Genet. 95: 197–200, 1995.

[169947]

[169948]

[169949]17191.

[169950]Hellevuo, K.; Yoshimura, M.; Kao, M.; Hoffman, P. L.;

Cooper, D.

[169951]M. F.; Tabakoff, B.: A novel adenylyl cyclase sequence
cloned from

[169952]the human erythroleukemia cell line. Biochem. Biophys.
Res. Commun. 192:

[169953]311–318, 1993.

[169954]

[169955]

[169956]17192.

[169957]Moerman, P.; Fryns, J.–P.; Sastrowijoto, S. H.; Vanden-
berghe, K.;

[169958]Lauweryns, J. M.: Hereditary renal adysplasia: new obser-
vations and

[169959]hypotheses. Pediat. Path. 14: 405–410, 1994.

[169960]

[169961]

[169962]17193.

[169963]Beri, R. K.; Marley, A. E.; See, C. G.; Sopwith, W. F.; Aguan,

[169964]K.; Carling, D.; Scott, J.; Carey, F.: Molecular cloning, ex-
pression

[169965]and chromosomal localisation of human AMP-activated
protein kinase. FEBS

[169966]Lett. 356: 117-121, 1994.

[169967]

[169968]

[169969]17194.

[169970]Hardie, D. G.: Regulation of fatty acid and cholesterol
metabolism

[169971]by the AMP-activated protein kinase. Biochim. Biophys.
Acta 1123:

[169972]231-238, 1992.

[169973]

[169974]

[169975]17195.

[169976]Hardie, D. G.; MacKintosh, R. W.: AMP-activated protein
kinase:

[169977]an archetypal protein kinase cascade? Bioessays 14:
699-704, 1992.

[169978]

[169979]

[169980]17196.

[169981]Mu, J.; Brozinick, J. T., Jr.; Valladares, O.; Bucan, M.; Birnbaum,

[169982]M. J.: A role for AMP-activated protein kinase in contraction- and

[169983]hypoxia-regulated glucose transport in skeletal muscle.

Molec. Cell 7:

[169984]1085-1094, 2001.

[169985]

[169986]

[169987]17197.

[169988]Stapleton, D.; Woollatt, E.; Mitchelhill, K. I.; Nicholl, J. K.;

[169989]Fernandez, C. S.; Michell, B. J.; Witters, L. A.; Power, D. A.; Sutherland,

[169990]G. R.; Kemp, B. E.: AMP-activated protein kinase isoenzyme family:

[169991]subunit structure and chromosomal location. FEBS Lett. 409: 452-456,

[169992]1997.

[169993]

[169994]

[169995]17198.

[169996]Budarf, M. L.; Perier, F.; Barnoski, B. L.; Bell, C. J.; Vandenberg,

[169997]C. A.: Assignment of the human hippocampal inward rectifier potassium

[169998]channel (HIR) gene to 22q13.1. Genomics 26: 625–629, 1995.

[169999]

[170000]

[170001]17199.

[170002]Makhina, E. N.; Kelly, A. J.; Lopatin, A. N.; Mercer, R. W.; Nichols,

[170003]C. G.: Cloning and expression of a novel human brain inward rectifier

[170004]potassium channel. J. Biol. Chem. 269: 20468–20474, 1994.

[170005]

[170006]

[170007]17200.

[170008]Perier, F.; Radeke, C. M.; Vandenberg, C. A.: Primary structure

[170009]and characterization of a small-conductance inwardly rectifying potassium

[170010]channel from human hippocampus. Proc. Nat. Acad. Sci. 91: 6240–6244,

[170011]1994.

[170012]

[170013]

[170014]17201.

[170015]Gal, A.; Wieringa, B.; Smeets, D. F. C. M.; Bleeker-Wagemakers,

[170016]L.; Ropers, H. H.: Submicroscopic interstitial deletion of the X

[170017]chromosome explains a complex genetic syndrome dominated by Norrie

[170018]disease. Cytogenet. Cell Genet. 42: 219–224, 1986.

[170019]

[170020]

[170021]17202.

[170022]Simpson, E.: The H-Y antigen and sex reversal. Cell 44:
813–814,

[170023]1986.

[170024]

[170025]

[170026]17203.

[170027]Di Paola, R.; Frittitta, L.; Miscio, G.; Bozzali, M.; Baratta,

[170028]R.; Centra, M.; Spampinato, D.; Santagati, M. G.; Ercolino,
T.; Cisternino,

[170029]C.; Soccio, T. Mastroianno, S.; Tassi, V.; Almgren, P.; Piz-
zuti, A.;

[170030]Vigneri, R.; Trischitta, V.: A variation in 3-prime UTR of
hPTP1B

[170031]increases specific gene expression and associates with in-
sulin resistance. Am.

[170032]J. Hum. Genet. 70: 806–812, 2002.

[170033]

[170034]

[170035]17204.

[170036]Eidne, K.; Taylor, P.; Connor, M.; Duthie, S.: Isolation,

characterization

[170037]and chromosomal localization of the human thyrotropin releasing hormone

[170038]receptor. (Abstract) 75th Annual Meeting of the Endocr. Soc. 437

[170039]only, 1993.

[170040]

[170041]

[170042]17205.

[170043]Castedo, M.; Ferri, K. F.; Blanco, J.; Roumier, T.; Laro-chette,

[170044]N.; Barretina, J.; Amendola, A.; Nardacci, R.; Metivier, D.; Este,

[170045]J. A.; Piacentini, M.; Kroemer, G.: Human immunodeficiency virus

[170046]1 envelope glycoprotein complex-induced apoptosis involves mammalian

[170047]target of rapamycin/FKBP12-rapamycin-associated protein-mediated p53

[170048]phosphorylation. J. Exp. Med. 194: 1097-1110, 2001.

[170049]

[170050]

[170051]17206.

[170052]El-Deiry, W. S.; Tokino, T.; Velculescu, V. E.; Levy, D. B.;

Parsons,

[170053]R.; Trent, J. M.; Lin, D.; Mercer, E.; Kinzler, K. W.; Vogel-

stein,

[170054]B.: WAF1, a potential mediator of p53 tumor suppression.

Cell 75:

[170055]817-825, 1993.

[170056]

[170057]

[170058]17207.

[170059]Moore, K. J.; D'Amore-Bruno, M. A.; Korfhagen, T. R.;

Glasser,

[170060]S. W.; Whitsett, J. A.; Jenkins, N. A.; Copeland, N. G.: Chro-

mosomal

[170061]localization of three pulmonary surfactant protein genes

in the mouse. Genomics 12:

[170062]388-393, 1992.

[170063]

[170064]

[170065]17208.

[170066]Nogee, L. M.; Dunbar, A. E., III; Wert, S. E.; Askin, F.; Hamvas,

[170067]A.; Whitsett, J. A.: A mutation in the surfactant protein C gene

[170068]associated with familial interstitial lung disease. New Eng. J. Med. 344:

[170069]573–579, 2001.

[170070]

[170071]

[170072]17209.

[170073]Warr, R. G.; Hawgood, S.; Buckley, D. I.; Crisp, T. M.; Schilling,

[170074]J.; Benson, B. J.; Ballard, P. L.; Clements, J. A.; White, R. T.:

[170075]Low molecular weight human pulmonary surfactant protein (SP5): isolation,

[170076]characterization, and cDNA and amino acid sequences. Proc. Nat. Acad.

[170077]Sci. 84: 7915–7919, 1987.

[170078]

[170079]

[170080]17210.

[170081]Wood, S.; Yaremko, M. L.; Schertzer, M.; Kelemen, P. R.;
Minna,

[170082]J.; Westbrook, C. A.: Mapping of the pulmonary surfactant
SP5 (SFTP2)

[170083]locus to 8p21 and characterization of a microsatellite re-
peat marker

[170084]that shows frequent loss of heterozygosity in human car-
cinomas. Genomics 24:

[170085]597–600, 1994.

[170086]

[170087]

[170088]17211.

[170089]Kolble, K.; Lu, J.; Mole, S. E.; Kaluz, S.; Reid, K. B. M.: As-
signment

[170090]of the human pulmonary surfactant protein D gene
(SFTP4) to 10q22–q23

[170091]close to the surfactant protein A gene cluster. Genomics
17: 294–298,

[170092]1993.

[170093]

[170094]

[170095]17212.

[170096]Botas, C.; Poulain, F.; Akiyama, J.; Brown, C.; Allen, L.; Go-
erke,

[170097]J.; Clements, J.; Carlson, E.; Gillespie, A. M.; Epstein, C.;
Hawgood,

[170098]S.: Altered surfactant homeostasis and alveolar type II cell
morphology

[170099]in mice lacking surfactant protein D. Proc. Nat. Acad. Sci.
95:

[170100]11869–11874, 1998.

[170101]

[170102]

[170103]17213.

[170104]Crouch, E.; Rust, K.; Veile, R.; Donis–Keller, H.; Grosso, L.:

[170105]Genomic organization of human surfactant protein D
(SP–D): SP–D is

[170106]encoded on chromosome 10q22.2–23.1. J. Biol. Chem.

268: 2976–2983,

[170107]1993.

[170108]

[170109]

[170110]17214.

[170111]Holmskov, U.; Lawson, P.; Teisner, B.; Tornoe, I.; Willis, A.
C.;

[170112]Morgan, C.; Koch, C.; Reid, K. B. M.: Isolation and charac-
terization

[170113]of a new member of the scavenger receptor superfamily,
glycoprotein–340

[170114](gp–340), as a lung surfactant protein–D binding
molecule. J. Biol.

[170115]Chem. 272: 13743–13749, 1997.

[170116]

[170117]

[170118]17215.

[170119]Holmskov, U.; Mollenhauer, J.; Madsen, J.; Vitved, L.;
Gronlund,

[170120]J.; Tornoe, I.; Kliem, A.; Reid, K. B. M.; Poustka, A.; Skjodt,

K.

[170121]: Cloning of gp-340, a putative opsonin receptor for lung surfactant

[170122]protein D. Proc. Nat. Acad. Sci. 96: 10794-10799, 1999.

[170123]

[170124]

[170125]17216.

[170126]LeVine, A. M.; Whitsett, J. A.; Hartshorn, K. L.; Crouch, E. C.;

[170127]Korfhagen, T. R.: Surfactant protein D enhances clearance of influenza

[170128]A virus from the lung in vivo. J. Immun. 167: 5868-5873, 2001.

[170129]

[170130]

[170131]17217.

[170132]Lu, J.; Willis, A. C.; Reid, K. B. M.: Purification, characterization

[170133]and cDNA cloning of human lung surfactant protein D. Biochem. J. 284:

[170134]795-802, 1992.

[170135]

[170136]

[170137]17218.

[170138]Rust, K.; Grosso, L.; Zhang, V.; Chang, D.; Persson, A.;
Longmore,

[170139]W.; Cai, G.-Z.; Crouch, E.: Human surfactant protein D:
SP-D contains

[170140]a C-type lectin carbohydrate recognition domain. Arch.
Biochem. Biophys. 290:

[170141]116–126, 1991.

[170142]

[170143]

[170144]17219.

[170145]Wert, S. E.; Yoshida, M.; LeVine, A. M.; Ikegami, M.; Jones,
T.;

[170146]Ross, G. F.; Fisher, J. H.; Korfhagen, T. R.; Whitsett, J. A.:
Increased

[170147]metalloproteinase activity, oxidant production, and em-
physema in surfactant

[170148]protein D gene-inactivated mice. Proc. Nat. Acad. Sci. 97:
5972–5977,

[170149]2000.

[170150]

[170151]

[170152]17220.

[170153]Gedde-Dahl, T., Jr.; Olaisen, B.; Teisberg, P.; Wilhelmy, M.
C.;

[170154]Mevag, B.; Helland, R.: The locus for apolipoprotein E
(apoE) is

[170155]close to the Lutheran (Lu) blood group locus on chromo-
some 19. Hum.

[170156]Genet. 67: 178-182, 1984.

[170157]

[170158]

[170159]17221.

[170160]Grubb, R.: Zur Genetik des Lewis-Systems. Naturwis-
senschaften 21:

[170161]560-561, 1953.

[170162]

[170163]

[170164]17222.

[170165]Kolanus, W.; Nagel, W.; Schiller, B.; Zeitlmann, L.; Godar, S.;

[170166]Stockinger, H.; Seed, B.: Alpha-L-beta-2 integrin/LFA-1 binding to

[170167]ICAM-1 induced by cytohesin-1, a cytoplasmic regulatory molecule. Cell 86:

[170168]233-242, 1996.

[170169]

[170170]

[170171]17223.

[170172]Ogasawara, M.; Kim, S.-C.; Adamik, R.; Togawa, A.; Ferrans, V.

[170173]J.; Takeda, K.; Kirby, M.; Moss, J.; Vaughan, M.: Similarities in

[170174]function and gene structure of cytohesin-4 and cytohesin-1, guanine

[170175]nucleotide-exchange proteins for ADP-ribosylation factors. J. Biol.

[170176]Chem. 275: 3221-3230, 2000.

[170177]

[170178]

[170179]17224.

[170180]Findlay, D. M.; Fisher, L. W.; McQuillan, C. I.; Termine, J. D.;

[170181]Young, M. F.: Isolation of the osteonectin gene: evidence that a

[170182]variable region of the osteonectin molecule is encoded within one

[170183]exon. *Biochemistry* 27: 1483–1489, 1988.

[170184]

[170185]

[170186]17225.

[170187]Gilmour, D. T.; Lyon, G. J.; Carlton, M. B. L.; Sanes, J. R.; Cunningham,

[170188]J. M.; Anderson, J. R.; Hogan, B. L. M.; Evans, M. J.; Colledge, W.

[170189]H.: Mice deficient for the secreted glycoprotein SPARC/osteonectin/BM40

[170190]develop normally but show severe age-onset cataract formation and

[170191]disruption of the lens. *EMBO J.* 17: 1860–1870, 1998.

[170192]

[170193]

[170194]17226.

[170195]Sanyanusin, P.; McNoe, L. A.; Sullivan, M. J.; Weaver, R. G.;

[170196]Eccles, M. R.: Mutation of PAX2 in two siblings with renal-
coloboma

[170197]syndrome. Hum. Molec. Genet. 4: 2183–2184, 1995.

[170198]

[170199]

[170200]17227.

[170201]Sanyanusin, P.; Norrish, J. H.; Ward, T. A.; Nebel, A.; Mc-
Noe,

[170202]L. A.; Eccles, M. R.: Genomic structure of the human PAX2
gene. Genomics 35:

[170203]258–261, 1996.

[170204]

[170205]

[170206]17228.

[170207]Sanyanusin, P.; Schimmenti, L. A.; McNoe, L. A.; Ward, T.

A.;

[170208]Pierpont, M. E. M.; Sullivan, M. J.; Dobyns, W. B.; Eccles, M.
R.

[170209]: Mutation of the PAX2 gene in a family with optic nerve
colobomas,

[170210]renal anomalies and vesicoureteral reflux. Nature Genet.
9: 358–364,

[170211]1995.

[170212]

[170213]

[170214]17229.

[170215]Schimmenti, L. A.; Cunliffe, H. E.; McNoe, L. A.; Ward, T.
A.;

[170216]French, M. C.; Shim, H. H.; Zhang, Y.–H.; Proesmans, W.;
Leys, A.;

[170217]Byerly, K. A.; Braddock, S. R.; Masuno, M.; Imaizumi, K.;
Devriendt,

[170218]K.; Eccles, M. R.: Further delineation of renal–coloboma
syndrome

[170219]in patients with extreme variability of phenotype and
identical PAX2

[170220]mutations. Am. J. Hum. Genet. 60: 869–878, 1997.

[170221]

[170222]

[170223]17230.

[170224]Schimmenti, L. A.; Pierpont, M. E.; Carpenter, B. L. M.;
Kashtan,

[170225]C. E.; Johnson, M. R.; Dobyns, W. B.: Autosomal dominant
optic nerve

[170226]colobomas, vesicoureteral reflux, and renal anomalies.
Am. J. Med.

[170227]Genet. 59: 204–208, 1995.

[170228]

[170229]

[170230]17231.

[170231]Schimmenti, L. A.; Shim, H. H.; Wirtschafter, J. D.; Pan-
zarino,

[170232]V. A.; Kashtan, C. E.; Kirkpatrick, S. J.; Wargowski, D. S.;
France,

[170233]T. D.; Michel, E.; Dobyns, W. B.: Homonucleotide expan-
sion and contraction

[170234]mutations of PAX2 and inclusion of Chiari 1 malformation
as part of

[170235]renal-coloboma syndrome. Hum. Mutat. 14: 369–376,
1999.

[170236]

[170237]

[170238]17232.

[170239]Stapleton, P.; Weith, A.; Urbanek, P.; Kozmik, Z.; Bus-
slinger,

[170240]M.: Chromosomal localization of seven PAX genes and
cloning of a

[170241]novel family member, PAX-9. Nature Genet. 3: 292–298,
1993.

[170242]

[170243]

[170244]17233.

[170245]Tellier, A.-L.; Amiel, J.; Delezoide, A.-L.; Audollent, S.;
Auge,

[170246]J.; Esnault, D.; Encha-Razavi, F.; Munnich, A.; Lyonnet, S.;
Vekemans,

[170247]M.; Attie-Bitach, T.: Expression of the PAX2 gene in hu-
man embryos

[170248]and exclusion in the CHARGE syndrome. Am. J. Med.

Genet. 93: 85–88,
[170249]2000.

[170250]

[170251]

[170252]17234.

[170253]Tellier, A.–L.; Amiel, J.; Salomon, R.; Jolly, D.; Delezoide,
[170254]A.–L.; Auge, J.; Gubler, M.–C.; Munnich, A.; Lyonnet, S.;
Antignac,
[170255]C.; Vekemans, M.; Broyer, M.; Attie–Bitach, T.: PAX2 ex–
pression during
[170256]early human development and its mutations in renal hy–
poplasia with
[170257]or without coloboma. (Abstract) Am. J. Hum. Genet. 63
(suppl.):
[170258]A7 only, 1998.

[170259]

[170260]

[170261]17235.

[170262]Ward, T. A.; Nebel, A.; Reeve, A. E.; Eccles, M. R.: Alterna–
tive

[170263]messenger RNA forms and open reading frames within an additional conserved

[170264]region of the human PAX-2 gene. Cell Growth Differ. 5: 1015-1021,

[170265]1994.

[170266]

[170267]

[170268]17236.

[170269]Weaver, R. G.; Cashwell, L. F.; Lorentz, W.; Whiteman, D.; Geisinger,

[170270]K. R.; Ball, M.: Optic nerve coloboma associated with renal disease. Am.

[170271]J. Med. Genet. 29: 597-605, 1988.

[170272]

[170273]

[170274]17237.

[170275]Barr, F. G.; Nauta, L. E.; Davis, R. J.; Schafer, B. W.; Nycum,

[170276]L. M.; Biegel, J. A.: In vivo amplification of the PAX3-FKHR and

[170277]PAX7-FKHR fusion genes in alveolar rhabdomyosarcoma.

Hum. Molec.

[170278]Genet. 5: 15–21, 1996.

[170279]

[170280]

[170281]17238.

[170282]Burri, M.; Tromvoukis, Y.; Bopp, D.; Frigerio, G.; Noll, M.:

Conservation

[170283]of the paired domain in metazoans and its structure in
three isolated

[170284]human genes. EMBO J. 8: 1183–1190, 1989.

[170285]

[170286]

[170287]17239.

[170288]Gruss, P.; Walther, C.: Pax in development. Cell 69:

719–722,

[170289]1992.

[170290]

[170291]

[170292]17240.

[170293]Schafer, B. W.; Mattei, M. G.: The human paired domain gene PAX7

[170294](Hup1) maps to chromosome 1p35–1p36.2. Genomics 17: 249–251, 1993.

[170295]

[170296]

[170297]17241.

[170298]Seale, P.; Sabourin, L. A.; Girgis–Gabardo, A.; Mansouri, A.; Gruss,

[170299]P.; Rudnicki, M. A.: Pax7 is required for the specification of myogenic

[170300]satellite cells. Cell 102: 777–786, 2000.

[170301]

[170302]

[170303]17242.

[170304]Shapiro, D. N.; Sublett, J. E.; Li, B.; Valentine, M. B.; Morris,

[170305]S. W.; Noll, M.: The gene for PAX7, a member of the paired–box–containing

[170306]genes, is localized on human chromosome arm 1p36. Genomics 17: 767–769,

[170307]1993.

[170308]

[170309]

[170310]17243.

[170311]Helwig, U.; Imai, K.; Schmahl, W.; Thomas, B. E.; Varnum, D. S.;

[170312]Nadeau, J. H.; Balling, R.: Interaction between undulated and Patch

[170313]leads to an extreme form of spina bifida in double-mutant mice. Nature

[170314]Genet. 11: 60-63, 1995.

[170315]

[170316]

[170317]17244.

[170318]Sun, J.; Rose, J. B.; Bird, P.: Gene structure, chromosomal localization,

[170319]and expression of the murine homologue of human proteinase inhibitor

[170320]6 (PI-6) suggests divergence of PI-6 from the ovalbumin serpins. J.

[170321]Biol. Chem. 270: 16089–16096, 1995.

[170322]

[170323]

[170324]17245.

[170325]Aberle, H.; Bierkamp, C.; Torchard, D.; Serova, O.; Wagner,
T.;

[170326]Natt, E.; Wirsching, J.; Heidkamper, C.; Montagna, M.;
Lynch, H. T.;

[170327]Lenoir, G. M.; Scherer, G.; Feunteun, J.; Kemler, R.: The
human plakoglobin

[170328]gene localizes on chromosome 17q21 and is subjected to
loss of heterozygosity

[170329]in breast and ovarian cancers. Proc. Nat. Acad. Sci. 92:
6384–6388,

[170330]1995.

[170331]

[170332]

[170333]17246.

[170334]Bierkamp, C.; Mclaughlin, K. J.; Schwarz, H.; Huber, O.;
Kemler,

[170335]R.: Embryonic heart and skin defects in mice lacking plakoglobin. Dev.

[170336]Biol. 180: 780–785, 1996.

[170337]

[170338]

[170339]17247.

[170340]Cowley, C. M. E.; Simrak, D.; Spurr, N. K.; Arnemann, J.; Buxton,

[170341]R. S.: The plakophilin 1 (PKP1) and plakoglobin (JUP) genes map to

[170342]human chromosomes 1q and 17, respectively. Hum. Genet. 100: 486–488,

[170343]1997.

[170344]

[170345]

[170346]17248.

[170347]Franke, W. W.; Goldschmidt, M. D.; Zimbelmann, R.; Mueller, H.

[170348]M.; Schiller, D. L.; Cowin, P.: Molecular cloning and amino acid

[170349]sequence of human plakoglobin, the common junctional plaque protein. Proc.

[170350]Nat. Acad. Sci. 86: 4027–4031, 1989.

[170351]

[170352]

[170353]17249.

[170354]Guenet, J.–L.; Simon–Chazottes, D.; Ringwald, M.; Kemler, R.:

[170355]The genes coding for alpha and beta catenin (Catna1 and Catnb) and

[170356]plakoglobin (Jup) map to mouse chromosomes 18, 9, and 11, respectively. Mammalian

[170357]Genome 6: 363–366, 1995.

[170358]

[170359]

[170360]17250.

[170361]Knudsen, K. A.; Wheelock, M. J.: Plakoglobin, or an 83–kD homologue

[170362]distinct from beta–catenin, interacts with E–cadherin and N–cadherin. J.

[170363]Cell Biol. 118: 671–679, 1992.

[170364]

[170365]

[170366]17251.

[170367]Kolligs, F. T.; Kolligs, B.; Hajra, K. M.; Hu, G.; Tani, M.;
Cho,

[170368]K. R.; Fearon, E. R.: Gamma–catenin is regulated by the
APC tumor

[170369]suppressor and its oncogenic activity is distinct from that
of beta–catenin. Genes

[170370]Dev. 14: 1319–1331, 2000.

[170371]

[170372]

[170373]17252.

[170374]Mathur, M.; Goodwin, L.; Cowin, P.: Interactions of the cy-
toplasmic

[170375]domain of the desmosomal cadherin Dsg1 with
plakoglobin. J. Biol.

[170376]Chem. 269: 14075–14080, 1994.

[170377]

[170378]

[170379]17253.

[170380]McKoy, G.; Protonotarios, N.; Crosby, A.; Tsatsopoulou, A.;
Anastasakis,

[170381]A.; Coonar, A.; Norman, M.; Baboonian, C.; Jeffery, S.;
McKenna, W.

[170382]J.: Identification of a deletion in plakoglobin in arrhythmo-
genic

[170383]right ventricular cardiomyopathy with palmoplantar kera-
toderma and

[170384]woolly hair (Naxos disease). Lancet 355: 2119–2124,
2000.

[170385]

[170386]

[170387]17254.

[170388]Ruiz, P.; Brinkmann, V.; Ledermann, B.; Behrend, M.;
Grund, C.;

[170389]Thalhammer, C.; Vogel, F.; Birchmeier, C.; Gunthert, U.;
Franke, W.

[170390]W.; Birchmeier, W.: Targeted mutation of plakoglobin in
mice reveals

[170391]essential functions of desmosomes in the embryonic heart. J. Cell

[170392]Biol. 135: 215–225, 1996.

[170393]

[170394]

[170395]17255.

[170396]Buckley, M. F.; Goding, J. W.: Plasma cell membrane glycoprotein

[170397]gene Pca-1 (alkaline phosphodiesterase I) is linked to the proto-oncogene

[170398]Myb on mouse chromosome 10. Immunogenetics 36: 199–201, 1992.

[170399]

[170400]

[170401]17256.

[170402]Buckley, M. F.; Loveland, K. A.; McKinstry, W. J.; Garson, O. M.;

[170403]Goding, J. W.: Plasma cell membrane glycoprotein PC-1: cDNA cloning

[170404]of the human molecule, amino acid sequence, and chromosomal location. J.

[170405]Biol. Chem. 265: 17506–17511, 1990.

[170406]

[170407]

[170408]17257.

[170409]Frittitta, L.; Baratta, R.; Spampinato, D.; Di Paola, R.; Pizzuti,

[170410]A.; Vigneri, R.; Trischitta, V.: The Q121 PC-1 variant and obesity

[170411]have additive and independent effects in causing insulin resistance. J.

[170412]Clin. Endocr. Metab. 86: 5888–5891, 2001.

[170413]

[170414]

[170415]17258.

[170416]Harahap, A. R.; Goding, J. W.: Distribution of the murine plasma

[170417]cell antigen PC-1 in non-lymphoid tissues. J. Immun. 141: 2317–2320,

[170418]1988.

[170419]

[170420]

[170421]17259.

[170422]Huang, R.; Rosenbach, M.; Vaughn, R.; Provvedini, D.;
Rebbe, N.;

[170423]Hickman, S.; Goding, J.; Terkeltaub, R.: Expression of the
murine

[170424]plasma cell nucleotide pyrophosphohydrolase PC-1 is
shared by human

[170425]liver, bone, and cartilage cells: regulation of PC-1 expres-
sion in

[170426]osteosarcoma cells by transforming growth factor-beta. J.
Clin. Invest. 94:

[170427]560-567, 1994.

[170428]

[170429]

[170430]17260.

[170431]Kahn, C. R.: Causes of insulin resistance. Nature 373:
384-385,

[170432]1995.

[170433]

[170434]

[170435]17261.

[170436]Maddux, B. A.; Goldfine, I. D.: Membrane glycoprotein PC-
1 inhibition

[170437]of insulin receptor function occurs via direct interaction
with the

[170438]receptor alpha-subunit. Diabetes 49: 13-19, 2000.

[170439]

[170440]

[170441]17262.

[170442]Maddux, B. A.; Sbraccia, P.; Kumakura, S.; Sasson, S.;
Youngren,

[170443]J.; Fisher, A.; Spencer, S.; Grupe, A.; Henzel, W.; Stewart, T.
A.;

[170444]Reaven, G. M.; Goldfine, I. D.: Membrane glycoprotein PC-
1 and insulin

[170445]resistance in non-insulin-dependent diabetes mellitus.
Nature 373:

[170446]448-451, 1995.

[170447]

[170448]

[170449]17263.

[170450]Nakamura, I.; Ikegawa, S.; Okawa, A.; Okuda, S.;

Koshizuka, Y.;

[170451]Kawaguchi, H.; Nakamura, K.; Koyama, T.; Goto, S.;

Toguchida, J.;

[170452]Matsushita, M.; Ochi, T.; Takaoka, K.; Nakamura, Y.: Association
of the human NPPS gene with ossification of the posterior

[170453]longitudinal
ligament of the spine (OPLL). Hum. Genet. 104: 492–497,

[170454]1999.

[170455]

[170456]

[170457]17264.

[170458]Okawa, A.; Nakamura, I.; Goto, S.; Moriya, H.; Nakamura,
Y.; Ikegawa,

[170459]S.: Mutation in Npps in a mouse model of ossification of
the posterior

[170460]longitudinal ligament of the spine. Nature Genet. 19:
271–273, 1998.

[170461]

[170462]

[170463]17265.

[170464]Pizzuti, A.; Frittitta, L.; Argiolas, A.; Baratta, R.; Goldfine,

[170465]I. D.; Bozzali, M.; Ercolino, T.; Scarlato, G.; Iacoviello, L.;

Vigneri,

[170466]R.; Tassi, V.; Trischitta, V.: A polymorphism (K121Q) of
the human

[170467]glycoprotein PC-1 gene coding region is strongly associ-
ated with insulin

[170468]resistance. Diabetes 48: 1881-1884, 1999.

[170469]

[170470]

[170471]17266.

[170472]Rebbe, N. F.; Tong, B. D.; Finley, E. M.; Hickman, S.: Iden-
tification

[170473]of nucleotide pyrophosphatase/alkaline phosphodi-
esterase I activity

[170474]associated with the mouse plasma cell differentiation
antigen PC-1. Proc.

[170475]Nat. Acad. Sci. 88: 5192–5196, 1991.

[170476]

[170477]

[170478]17267.

[170479]Sakaguchi, A. Y.; Lalley, P. A.; Zabel, B. U.; Ellis, R. W.;
Scolnick,

[170480]E. M.; Naylor, S. L.: Chromosome assignments of four
mouse cellular

[170481]homologs of sarcoma and leukemia virus oncogenes.
Proc. Nat. Acad.

[170482]Sci. 81: 525–529, 1984.

[170483]

[170484]

[170485]17268.

[170486]Takahashi, T.; Old, L. J.; Boyse, E. A.: Surface alloantigens

[170487]of plasma cells. J. Exp. Med. 131: 1325–1341, 1970.

[170488]

[170489]

[170490]17269.

[170491]van Driel, I. R.; Goding, J. W.: Plasma cell membrane glycoprotein

[170492]PC-1: primary structure deduced from cDNA clones. J. Biol. Chem. 262:

[170493]4882-4887, 1987.

[170494]

[170495]

[170496]17270.

[170497]Frank, S. L.; Klisak, I.; Sparkes, R. S.; Lusk, A. J.: A gene

[170498]homologous to plasminogen located on human chromosome 2q11-p11. Genomics 4:

[170499]449-451, 1989.

[170500]

[170501]

[170502]17271.

[170503]Westergaard, J. G.; Chemnitz, J.; Teisner, B.; Poulsen, H. K.;

[170504]Ipsen, L.; Beck, B.; Grudzinskas, J. G.: Pregnancy-associated plasma

[170505]protein A: a possible marker in the classification and pre-

natal diagnosis

[170506]of Cornelia de Lange syndrome. Prenatal Diag. 3:
225–232, 1983.

[170507]

[170508]

[170509]17272.

[170510]Barnett, T.; Pickle, W., II; Rae, P. M. M.; Hart, J.; Kamarck,

[170511]M.; Elting, J.: Pregnancy-specific beta-1-glycoproteins are
related

[170512]to carcinoembryonic antigens and map to chromosome
19. (Abstract) Cytogenet.

[170513]Cell Genet. 51: 958, 1989.

[170514]

[170515]

[170516]17273.

[170517]Barnett, T. R.; Pickle, W., II; Rae, P. M. M.; Hart, J.; Ka-
marck,

[170518]M.; Elting, J.: Human pregnancy-specific
beta(1)-glycoproteins are

[170519]coded within chromosome 19. Am. J. Hum. Genet. 44:

890–893, 1989.

[170520]

[170521]

[170522]17274.

[170523]Bartels, I.; Lindemann, A.: Maternal levels of pregnancy-specific
cific

[170524]beta-1-glycoprotein (SP-1) are elevated in pregnancies
affected by

[170525]Down's syndrome. Hum. Genet. 80: 46–48, 1988.

[170526]

[170527]

[170528]17275.

[170529]Brandriff, B. F.; Gordon, L. A.; Tynan, K. T.; Olsen, A. S.;
Mohrenweiser,

[170530]H. W.; Fertitta, A.; Carrano, A. V.; Trask, B. J.: Order and
genomic

[170531]distances among members of the carcinoembryonic anti-
gen (CEA) gene

[170532]family determined by fluorescence in situ hybridization.

Genomics 12:

[170533]773–779, 1992.

[170534]

[170535]

[170536]17276.

[170537]Chan, W.-Y.; Qiu, W.-R.: Human pregnancy-specific beta-1 glycoprotein

[170538]is encoded by multiple genes localized on two chromosomes. Am. J.

[170539]Hum. Genet. 43: 152-159, 1988.

[170540]

[170541]

[170542]17277.

[170543]Khan, W. N.; Teglund, S.; Bremer, K.; Hammarstrom, S.: The pregnancy-specific

[170544]glycoprotein family of the immunoglobulin superfamily: identification

[170545]of new members and estimation of family size. Genomics 12: 780-787,

[170546]1992.

[170547]

[170548]

[170549]17278.

[170550]Niemann, S. C.; Flake, A.; Bohn, H.; Bartels, I.: Pregnancy-specific

[170551]beta-1-glycoprotein: cDNA cloning, tissue expression, and species

[170552]specificity of one member of the PSBG family. Hum. Genet. 82: 239-243,

[170553]1989.

[170554]

[170555]

[170556]17279.

[170557]Niemann, S. C.; Schonk, D.; van Dijk, P.; Wieringa, B.; Grzeschik,

[170558]K.-H.; Bartels, I.: Regional localization of the gene encoding pregnancy

[170559]specific beta-1-glycoprotein 1 (PSBG1) to human chromosome 19q13.1. Cytogenet.

[170560]Cell Genet. 52: 95-97, 1989.

[170561]

[170562]

[170563]17280.

[170564]Niemann, S. C.; Schonk, D.; van Dijk, P. E.; Grzeschik, K.-H.;

[170565]Bartels, I.: Chromosomal assignment of a cDNA clone encoding pregnancy-specific

[170566]beta-1-glycoprotein to chromosome 19. (Abstract) Cytogenet. Cell

[170567]Genet. 51: 1053, 1989.

[170568]

[170569]

[170570]17281.

[170571]Olsen, A.; Teglund, S.; Nelson, D.; Gordon, L.; Copeland, A.;

[170572]Georgescu, A.; Carrano, A.; Hammarstrom, S.: Gene organization of

[170573]the pregnancy-specific glycoprotein region on human chromosome 19:

[170574]assembly and analysis of a 700-kb cosmid contig spanning the region. Genomics 23:

[170575]659-668, 1994.

[170576]

[170577]

[170578]17282.

[170579]Streydio, C.; Swillens, S.; Georges, M.; Szpirer, C.; Vassart,

[170580]G.: Structure, evolution and chromosomal localization of
the human

[170581]pregnancy-specific beta-1 glycoprotein gene family. Ge-
nomics 6:

[170582]579-592, 1990. Note: Erratum: Genomics 7: 661-662,
1990.

[170583]

[170584]

[170585]17283.

[170586]Teglund, S.; Olsen, A.; Khan, W. N.; Frangsmyr, L.; Ham-
marstrom,

[170587]S.: The pregnancy-specific glycoprotein (PSG) gene cluster
on human

[170588]chromosome 19: fine structure of the 11 PSG genes and
identification

[170589]of 6 new genes forming a third subgroup within the carci-
noembryonic

[170590]antigen (CEA) family. Genomics 23: 669-684, 1994.

[170591]

[170592]

[170593]17284.

[170594]Thompson, J.; Koumari, R.; Wagner, K.; Barnert, S.;

Schleussner,

[170595]C.; Schrewe, H.; Zimmermann, W.; Muller, G.; Schempp,

W.; Zaninetta,

[170596]D.; Ammaturo, D.; Hardman, N.: The human pregnancy-specific glycoprotein

[170597]genes are tightly linked on the long arm of chromosome 19 and are

[170598]coordinately expressed. Biochem. Biophys. Res. Commun. 167: 848-859,

[170599]1990.

[170600]

[170601]

[170602]17285.

[170603]Watanabe, S.; Chou, J. Y.: Isolation and characterization of

[170604]complementary DNAs encoding human pregnancy-specific beta-1-glycoprotein. J.

[170605]Biol. Chem. 263: 2049–2054, 1988.

[170606]

[170607]

[170608]17286.

[170609]Leslie, K. K.; Watanabe, S.; Lei, K.-J.; Chou, D. Y.; Plouzek,

[170610]C. A.; Deng, H.-C.; Torres, J.; Chou, J. Y.: Linkage of two
human

[170611]pregnancy-specific beta-1-glycoprotein genes: one is as-
sociated with

[170612]hydatidiform mole. Proc. Nat. Acad. Sci. 87: 5822–5826,
1990.

[170613]

[170614]

[170615]17287.

[170616]Olsen, A.; Teglund, S.; Nelson, D.; Gordon, L.; Copeland,
A.; Georgescu,

[170617]A.; Carrano, A.; Hammarstrom, S.: Gene organization of
the pregnancy-specific

[170618]glycoprotein region on human chromosome 19: assembly
and analysis

[170619]of a 700-kb cosmid contig spanning the region. Genomics
23: 659–668,
[170620]1994.

[170621]

[170622]

[170623]17288.

[170624]Thompson, J.; Zimmermann, W.; Osthus–Bugat, P.;
Schleussner, C.;

[170625]Eades–Perner, A.–M.; Barnert, S.; von Kleist, S.; Willcocks,
T.; Craig,

[170626]I.; Tynan, K.; Olsen, A.; Mohrenweiser, H.: Long–range
chromosomal

[170627]mapping of the carcinoembryonic antigen (CEA) gene
family cluster. Genomics 12:

[170628]761–772, 1992.

[170629]

[170630]

[170631]17289.

[170632]Tynan, K.; Olsen, A.; Trask, B.; de Jong, P.; Thompson, J.;
Zimmermann,

[170633]W.; Carrano, A.; Mohrenweiser, H.: Assembly and analysis
of cosmid

[170634]contigs in the CEA-gene family region of human chromo-
some 19. Nucleic

[170635]Acids Res. 20: 1629-1636, 1992.

[170636]

[170637]

[170638]17290.

[170639]Sheer, D.; Sheppard, D. M.; Le Beau, M.; Rowley, J. D.; San
Roman,

[170640]C.; Solomon, E.: Localization of the oncogene c-erbA1 im-
mediately

[170641]proximal to the acute promyelocytic leukaemia breakpoint
on chromosome

[170642]17. Ann. Hum. Genet. 49: 167-171, 1985.

[170643]

[170644]

[170645]17291.

[170646]Nobukuni, Y.; Watanabe, A.; Takeda, K.; Skarka, H.;
Tachibana,

[170647]M.: Analyses of loss-of-function mutations of the MITF gene suggest

[170648]that haploinsufficiency is a cause of Waardenburg syndrome type 2A. Am.

[170649]J. Hum. Genet. 59: 76–83, 1996.

[170650]

[170651]

[170652]17292.

[170653]Steingrimsson, E.; Moore, K. J.; Lamoreux, M. L.; Ferre-D'Amare,

[170654]A. R.; Burley, S. K.; Sanders Zimring, D. C.; Skow, L. C.; Hodgkinson,

[170655]C. A.; Arnheiter, H.; Copeland, N. G.; Jenkins, N. A.: Molecular

[170656]basis of mouse microphthalmia (mi) mutations helps explain their developmental

[170657]and phenotypic consequences. Nature Genet. 8: 256–263, 1994.

[170658]

[170659]

[170660]17293.

[170661]Steingrimsen, E.; Tessarollo, L.; Pathak, B.; Hou, L.; Arnheiter,

[170662]H.; Copeland, N. G.; Jenkins, N. A.: Mitf and Tfe3, two members of

[170663]the Mitf-Tfe family of bHLH-Zip transcription factors, have important

[170664]but functionally redundant roles in osteoclast development. Proc.

[170665]Nat. Acad. Sci. 99: 4477-4482, 2002.

[170666]

[170667]

[170668]17294.

[170669]Tachibana, M.; Perez-Jurado, L. A.; Nakayama, A.; Hodgkinson,

[170670]C. A.; Li, X.; Schneider, M.; Miki, T.; Fex, J.; Francke, U.; Arnheiter,

[170671]H.: Cloning of MITF, the human homolog of the mouse microphthalmia

[170672]gene and assignment to chromosome 3p14.1-p12.3. Hum. Molec. Genet. 3:

[170673]553-557, 1994.

[170674]

[170675]

[170676]17295.

[170677]Tachibana, M.; Takeda, K.; Nobukuni, Y.; Urabe, K.; Long, J. E.;

[170678]Meyers, K. A.; Aaronson, S. A.; Miki, T.: Ectopic expression of MITF,

[170679]a gene for Waardenburg syndrome type 2, converts fibroblasts to cells

[170680]with melanocytes characteristics. Nature Genet. 14: 50–54, 1996.

[170681]

[170682]

[170683]17296.

[170684]Takeda, K.; Takemoto, C.; Kobayashi, I.; Watanabe, A.; Nobukuni,

[170685]Y.; Fisher, D. E.; Tachibana, M.: Ser298 of MITF, a mutation site

[170686]in Waardenburg syndrome type 2, is a phosphorylation site with functional

[170687]significance. Hum. Molec. Genet. 9: 125–132, 2000.

[170688]

[170689]

[170690]17297.

[170691]Tassabehji, M.; Newton, V. E.; Read, A. P.: Waardenburg
syndrome

[170692]type 2 caused by mutations in the human microphthalmia
(MITF) gene. Nature

[170693]Genet. 8: 251–255, 1994.

[170694]

[170695]

[170696]17298.

[170697]Watanabe, A.; Takeda, K.; Ploplis, B.; Tachibana, M.:
Epistatic

[170698]relationship between Waardenburg syndrome genes MITF
and PAX3. Nature

[170699]Genet. 18: 283–286, 1998.

[170700]

[170701]

[170702]17299.

[170703]Yoshitake, H.; Rittling, S. R.; Denhardt, D. T.; Noda, M.:

Osteopontin-deficient

[170704]mice are resistant to ovariectomy-induced bone resorption. Proc.

[170705]Nat. Acad. Sci. 96: 8156–8160, 1999.

[170706]

[170707]

[170708]17300.

[170709]Edelmann, W.; Zervas, M.; Costello, P.; Roback. L.; Fischer, I.;

[170710]Hammarback, J. A.; Cowan, N.; Davies, P.; Wainer, B.; Kucherlapati,

[170711]R.: Neuronal abnormalities in microtubule-associated protein 1B mutant

[170712]mice. Proc. Nat. Acad. Sci. 93: 1270–1275, 1996.

[170713]

[170714]

[170715]17301.

[170716]Hammarback, J. A.; Obar, R. A.; Hughes, S. M.; Vallee, R. B.:

[170717]MAP1B is encoded as a polyprotein that is processed to form a complex

[170718]N-terminal microtubule-binding domain. Neuron 7: 129-139, 1991.

[170719]

[170720]

[170721]17302.

[170722]Lien, L. L.; Boyce, F. M.; Kleyn, P.; Brzustowicz, L. M.; Men-
ninger,

[170723]J.; Ward, D. C.; Gilliam, T. C.; Kunkel, L. M.: Mapping of
human

[170724]microtubule-associated protein 1B in proximity to the
spinal muscular

[170725]atrophy locus at 5q13. Proc. Nat. Acad. Sci. 88:
7873-7876, 1991.

[170726]

[170727]

[170728]17303.

[170729]Lien, L. L.; Feener, C. A.; Fischbach, N.; Kunkel, L. M.:
Cloning

[170730]of human microtubule-associated protein 1B and the

identification

[170731]of a related gene on chromosome 15. Genomics 22:
273–280, 1994.

[170732]

[170733]

[170734]17304.

[170735]Wirth, B.; Voosen, B.; Rohrig, D.; Knapp, M.; Piechaczek,
B.; Rudnik–Schoneborn,

[170736]S.; Zerres, K.: Fine mapping and narrowing of the genetic
interval

[170737]of the spinal muscular atrophy region by linkage studies.
Genomics 15:

[170738]113–118, 1993.

[170739]

[170740]

[170741]17305.

[170742]Zhang, Y. Q.; Bailey, A. M.; Matthies, H. J. G.; Renden, R.
B.;

[170743]Smith, M. A.; Speese, S. D.; Rubin, G. M.; Broadie, K.:
Drosophila

[170744]fragile X–related gene regulates the MAP1B homolog

Futsch to control

[170745]synaptic structure and function. Cell 107: 591–603, 2001.

[170746]

[170747]

[170748]17306.

[170749]Garner, C. C.; Tucker, R. P.; Matus, A.: Selective localization
tion

[170750]of messenger RNA for cytoskeletal protein MAP2 in dendrites. Nature 336:

[170751]674–677, 1988.

[170752]

[170753]

[170754]17307.

[170755]Kalcheva, N.; Albala, J.; O'Guin, K.; Rubino, H.; Garner, C.;
Shafit-Zagardo,

[170756]B.: Genomic structure of human microtubule-associated
protein 2 (MAP-2)

[170757]and characterization of additional MAP-2 isoforms. Proc.
Nat. Acad.

[170758]Sci. 92: 10894–10898, 1995.

[170759]

[170760]

[170761]17308.

[170762]Kindler, S.; Garner, C. C.: Four repeat MAP2 isoforms in
human

[170763]and rat brain. Molec. Brain Res. 26: 218–224, 1994.

[170764]

[170765]

[170766]17309.

[170767]Marsden, K. M.; Doll, T.; Ferralli, J.; Botteri, F.; Matus, A.

[170768]: Transgenic expression of embryonic MAP2 in adult
mouse brain: implications

[170769]for neuronal polarization. J. Neurosci. 16: 3265–3273,
1996.

[170770]

[170771]

[170772]17310.

[170773]Neve, R. L.; Harris, P.; Kosik, K. S.; Kurnit, D. M.; Donlon,
T.

[170774]A.: Identification of cDNA clones for the human micro-
tubule-associated

[170775]protein tau and chromosomal localization of the genes for
tau and

[170776]microtubule-associated protein 2. Molec. Brain Res. 1:
271-280,

[170777]1986.

[170778]

[170779]

[170780]17311.

[170781]Chapin, S. J.; Bulinski, J. C.: Non-neuronal 210 x 10(3) M(r)

[170782]microtubule-associated protein (MAP4) contains a domain
homologous

[170783]to the microtubule-binding domains of neuronal MAP2
and tau. J. Cell

[170784]Sci. 98: 27-36, 1991.

[170785]

[170786]

[170787]17312.

[170788]Adriaansen, H. J.; Geurts Van Kessel, A. H. M.; Wijdenes-

De Bresser,

[170789]J. H. F. M.; Van Drunen–Schoenmaker, E.; Van Dongen, J. J.

M.: Expression

[170790]of the myeloid differentiation antigen CD33 depends on
the presence

[170791]of human chromosome 19 in human–mouse hybrids. Ann.

Hum. Genet. 54:

[170792]115–119, 1990.

[170793]

[170794]

[170795]17313.

[170796]Bonthron, D. T.; Dunlop, N.; Barr, D. G. D.; El Sanousi, A.

A.;

[170797]Al–Gazali, L. I.: Organisation of the human PAX4 gene and
its exclusion

[170798]as a candidate for the Wolcott–Rallison syndrome. J. Med.

Genet. 35:

[170799]288–292, 1998.

[170800]

[170801]

[170802]17314.

[170803]Inoue, H.; Nomiya, J.; Nakai, K.; Matsutani, A.;

Tanizawa, Y.;

[170804]Oka, Y.: Isolation of full-length cDNA of mouse PAX4 gene
and identification

[170805]of its human homologue. Biochem. Biophys. Res. Com-
mun. 243: 628-633,

[170806]1998.

[170807]

[170808]

[170809]17315.

[170810]Mansouri, A.; St-Onge, L.; Gruss, P.: Role of Pax genes in
endoderm-derived

[170811]organs. Trends Endocr. Metab. 10: 164-167, 1999.

[170812]

[170813]

[170814]17316.

[170815]Matsushita, T.; Yamaoka, T.; Otsuka, S.; Moritani, M.; Mat-
sumoto,

[170816]T.; Itakura, M.: Molecular cloning of mouse paired-
box-containing

[170817]gene (Pax)–4 from an islet beta cell line and deduced sequence of

[170818]human Pax–4. Biochem. Biophys. Res. Commun. 242: 176–180, 1998.

[170819]

[170820]

[170821]17317.

[170822]Sosa–Pineda, B.; Chowdhury, K.; Torres, M.; Oliver, G.; Gruss,

[170823]P.: The Pax4 gene is essential for differentiation of insulin–producing

[170824]beta cells in the mammalian pancreas. Nature 386: 399–402, 1997.

[170825]

[170826]

[170827]17318.

[170828]St–Onge, L.; Sosa–Pineda, B.; Chowdhury, K.; Mansouri, A.; Gruss,

[170829]P.: Pax6 is required for differentiation of glucagon–producing alpha–cells

[170830]in mouse pancreas. Nature 387: 406–409, 1997.

[170831]

[170832]

[170833]17319.

[170834]Tamura, T.; Izumikawa, Y.; Kishino, T.; Soejima, H.; Jinno, Y.;

[170835]Niikawa, N.: Assignment of the human PAX4 gene to chromosome band

[170836]7q32 by fluorescence in situ hybridization. Cytogenet. Cell Genet. 66:

[170837]132–134, 1994.

[170838]

[170839]

[170840]17320.

[170841]Adams, B.; Dorfler, P.; Aguzzi, A.; Kozmik, Z.; Urbanek, P.; Maurer–Fogy,

[170842]I.; Busslinger, M.: Pax–5 encodes the transcription factor BSAP and

[170843]is expressed in B lymphocytes, the developing CNS, and adult testis. Genes

[170844]Dev. 6: 1589–1607, 1992.

[170845]

[170846]

[170847]17321.

[170848]Busslinger, M.; Klix, N.; Pfeffer, P.; Graninger, P. G.;

Kozmik,

[170849]Z.: Deregulation of PAX-5 by translocation of the E- μ
enhancer of

[170850]the IgH locus adjacent to two alternative PAX-5 promoters
in a diffuse

[170851]large-cell lymphoma. Proc. Nat. Acad. Sci. 93:

6129-6134, 1996.

[170852]

[170853]

[170854]17322.

[170855]Mikkola, I.; Heavey, B.; Horcher, M.; Busslinger, M.: Rever-
sion

[170856]of B cell commitment upon loss of Pax5 expression. Sci-
ence 297:

[170857]110-113, 2002.

[170858]

[170859]

[170860]17323.

[170861]Nutt, S. L.; Heavey, B.; Rolink, A. G.; Busslinger, M.: Com-
mitment

[170862]to the B-lymphoid lineage depends on the transcription
factor Pax5. Nature 401:

[170863]556-562, 1999.

[170864]

[170865]

[170866]17324.

[170867]Nutt, S. L.; Vambrie, S.; Steinlein, P.; Kozmik, Z.; Rolink,
A.;

[170868]Weith, A.; Busslinger, M.: Independent regulation of the
two Pax5

[170869]alleles during B-cell development. Nature Genet. 21:
390-395, 1999.

[170870]

[170871]

[170872]17325.

[170873]Ohno, H.; Furukawa, T.; Fukuhara, S.; Zong, S. Q.; Kame-

saki, H.;

[170874]Shows, T. B.; Le Beau, M. M.; McKeithan, T. W.; Kawakami, T.; Honjo,

[170875]T.: Molecular analysis of a chromosomal translocation, t(9;14)(p13;q32),

[170876]in a diffuse large-cell lymphoma cell line expressing the Ki-1 antigen. Proc.

[170877]Nat. Acad. Sci. 87: 628-632, 1990.

[170878]

[170879]

[170880]17326.

[170881]Rolink, A. G.; Nutt, S. L.; Melchers, F.; Busslinger, M.: Long-term

[170882]in vivo reconstitution of T-cell development by Pax5-deficient B-cell

[170883]progenitors. Nature 401: 603-606, 1999.

[170884]

[170885]

[170886]17327.

[170887]Vorechovsky, I.; Koskinen, S.; Paganelli, R.; Smith, C. I. E.;

[170888]Busslinger, M.; Hammarstrom, L.: The PAX5 gene: a linkage and mutation

[170889]analysis in candidate human primary immunodeficiencies.

Immunogenetics 42:

[170890]149–152, 1995.

[170891]

[170892]

[170893]17328.

[170894]Walther, C.; Guenet, J.–L.; Simon, D.; Deutsch, U.; Jostes, B.;

[170895]Goulding, M. D.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a murine

[170896]multigene family of paired box containing genes. Genomics 11: 424–434,

[170897]1991.

[170898]

[170899]

[170900]17329.

[170901]Koseki, H.; Zachgo, J.; Mizutani, Y.; Simon–Chazottes, D.; Guenet,

[170902]J.-L.; Balling, R.; Gossler, A.: Fine genetic mapping of the proximal

[170903]part of mouse chromosome 2 excludes Pax-8 as a candidate gene for

[170904]Danforth's short tail (Sd). Mammalian Genome 4: 324-327, 1993.

[170905]

[170906]

[170907]17330.

[170908]Kroll, T. G.; Sarraf, P.; Pecciarini, L.; Chen, C.-J.; Mueller,

[170909]E.; Splegeman, B. M.; Fletcher, J. A.:

PAX8-PPAR-gamma-1 fusion

[170910]in oncogene human thyroid carcinoma. Science 289: 1357-1360, 2000.

[170911]

[170912]

[170913]17331.

[170914]Macchia, P. E.; Lapi, P.; Krude, H.; Pirro, M. T.; Missero, C.;

[170915]Chiovato, L.; Souabni, A.; Baserga, M.; Tassi, V.; Pinchera, A.; Fenzi,

[170916]G.; Gruters, A.; Busslinger, M.; Di Lauro, R.: PAX8 mutations associated

[170917]with congenital hypothyroidism caused by thyroid dysgenesis. Nature

[170918]Genet. 19: 83–86, 1998.

[170919]

[170920]

[170921]17332.

[170922]Mansouri, A.; Chowdhury, K.; Gruss, P.: Follicular cells of the

[170923]thyroid gland require Pax8 gene function. Nature Genet. 19: 87–90,

[170924]1998.

[170925]

[170926]

[170927]17333.

[170928]Pasca di Magliano, M.; Di Lauro, R.; Zannini, M.: Pax8 has a key

[170929]role in thyroid cell differentiation. Proc. Nat. Acad. Sci. 97:

[170930]13144–13149, 2000.

[170931]

[170932]

[170933]17334.

[170934]Plachov, D.; Chowdhury, K.; Walther, C.; Simon, D.;
Guenet, J.-L.;

[170935]Gruss, P.: Pax-8, a murine paired box gene expressed in
the developing

[170936]excretory system and thyroid gland. Development 110:
643-651, 1990.

[170937]

[170938]

[170939]17335.

[170940]Tell, G.; Pellizzari, L.; Esposito, G.; Pucillo, C.; Macchia,

[170941]P. E.; Di Lauro, R.; Damante, G.: Structural defects of a
Pax8 mutant

[170942]that give rise to congenital hypothyroidism. Biochem. J.
341: 89-93,

[170943]1999.

[170944]

[170945]

[170946]17336.

[170947]Bongarzone, I.; Vigano, E.; Alberti, L.; Borrello, M. G.;

Pasini,

[170948]B.; Greco, A.; Mondellini, P.; Smith, D. P.; Ponder, B. A. J.;

Romeo,

[170949]G.; Pierotti, M. A.: Full activation of MEN2B mutant RET by

an additional

[170950]MEN2A mutation or by ligand GDNF stimulation. Onco-

gene 16: 2295-2301,

[170951]1998.

[170952]

[170953]

[170954]17337.

[170955]Carlson, K. M.; Bracamontes, J.; Jackson, C. E.; Clark, R.;

Lacroix,

[170956]A.; Wells, S. A., Jr.; Goodfellow, P. J.: Parent-of-origin ef-

fects

[170957]in multiple endocrine neoplasia type 2B. Am. J. Hum.

Genet. 55:

[170958]1076-1082, 1994.

[170959]

[170960]

[170961]17338.

[170962]Carlson, K. M.; Dou, S.; Chi, D.; Scavarda, N.; Toshima, K.;
Jackson,

[170963]C. E.; Wells, S. A., Jr.; Goodfellow, P. J.; Donis-Keller, H.:
Single

[170964]missense mutation in the tyrosine kinase catalytic domain
of the RET

[170965]protooncogene is associated with multiple endocrine neo-
plasia type

[170966]2B. Proc. Nat. Acad. Sci. 91: 1579-1583, 1994.

[170967]

[170968]

[170969]17339.

[170970]Mulligan, L. M.; Eng, C.; Healey, C. S.; Clayton, D.; Kwok, J.

[170971]B. J.; Gardner, E.; Ponder, M. A.; Frilling, A.; Jackson, C. E.;
Lehnert,

[170972]H.; Neumann, H. P. H.; Thibodeau, S. N.; Ponder, B. A. J.:
Specific

[170973]mutations of the RET proto-oncogene are related to dis-
ease phenotype

[170974]in MEN 2A and FMTC. Nature Genet. 6: 70–74, 1994.

[170975]

[170976]

[170977]17340.

[170978]Bruss, M.; Kunz, J.; Lingen, B.; Bonisch, H.: Chromosomal mapping

[170979]of the human gene for the tricyclic antidepressant-sensitive noradrenaline

[170980]transporter. Hum. Genet. 91: 278–280, 1993.

[170981]

[170982]

[170983]17341.

[170984]Esler, M.; Jennings, G.; Lambert, G.; Meredith, I.; Horne, M.;

[170985]Eisenhofer, G.: Overflow of catecholamine neurotransmitters to the

[170986]circulation: source, fate, and functions. Physiol. Rev. 70: 963–985,

[170987]1990.

[170988]

[170989]

[170990]17342.

[170991]Fritz, J. D.; Jayanthi, L. D.; Thoreson, M. A.; Blakely, R. D.

[170992]: Cloning and chromosomal mapping of the murine norepinephrine transporter. J.

[170993]Neurochem. 70: 2241-2251, 1998.

[170994]

[170995]

[170996]17343.

[170997]Gelernter, J.; Kruger, S.; Kidd, K. K.; Amara, S.: TaqI RFLP
at

[170998]norepinephrine transporter protein (NET) locus. Hum.
Molec. Genet. 2:

[170999]820 only, 1993.

[171000]

[171001]

[171002]17344.

[171003]Gelernter, J.; Kruger, S.; Pakstis, A. J.; Pacholczyk, T.;
Sparkes,

[171004]R. S.; Kidd, K. K.; Amara, S.: Assignment of the norepinephrine transporter

[171005]protein (NET1) locus to chromosome 16. Genomics 18: 690–692, 1993.

[171006]

[171007]

[171008]17345.

[171009]Kaye, W. H.; Jimerson, D. C.; Lake, C. R.; Ebert, M. H.: Altered

[171010]norepinephrine metabolism following long-term weight recovery in patients

[171011]with anorexia nervosa. Psychiat. Res. 14: 333–342, 1985.

[171012]

[171013]

[171014]17346.

[171015]Pacholczyk, T.; Blakely, R. D.; Amara, S. G.: Expression cloning

[171016]of a cocaine- and antidepressant-sensitive human norepinephrine transporter. Nature 350:

[171017]350–354, 1991.

[171018]

[171019]

[171020]17347.

[171021]Paczkowski, F. A.; Bonisch, H.; Bryan-Lluka, L. J.: Pharma-
cological

[171022]properties of the naturally occurring ala457pro variant of
the human

[171023]norepinephrine transporter. Pharmacogenetics 12:
165–173, 2002.

[171024]

[171025]

[171026]17348.

[171027]Pirke, K. M.; Kellner, M.; Philipp, E.; Laessle, R.; Krieg, J.

[171028]C.; Fichter, M. M.: Plasma norepinephrine after a stan-
dardized test

[171029]meal in acute and remitted patients with anorexia nervosa
and in healthy

[171030]controls. Biol. Psychiat. 31: 1074–1077, 1992.

[171031]

[171032]

[171033]17349.

[171034]Porzgen, P.; Bonisch, H.; Bruss, M.: Molecular cloning and organization

[171035]of the coding region of the human norepinephrine transporter gene. Biochem.

[171036]Biophys. Res. Commun. 215: 1145–1150, 1995. Notes: Erratum: Biochem.

[171037]Biophys. Res. Commun. 227: 642–643, 1996.

[171038]

[171039]

[171040]17350.

[171041]Shannon, J. R.; Flattem, N. L.; Jordan, J.; Jacob, G.; Black,

[171042]B. K.; Biaggioni, I.; Blakely, R. D.; Robertson, D.: Orthostatic

[171043]intolerance and tachycardia associated with norepinephrine-transporter

[171044]deficiency. New Eng. J. Med. 342: 541–549, 2000.

[171045]

[171046]

[171047]17351.

[171048]Urwin, R. E.; Bennetts, B.; Wilcken, B.; Lampropoulos, B.;
Beumont,

[171049]P.; Clarke, S.; Russell, J.; Tanner, S.; Nunn, K. P.: Anorexia
nervosa

[171050](restrictive subtype) is associated with a polymorphism in
the novel

[171051]norepinephrine transporter gene promoter polymorphic
region. Molec.

[171052]Psychiat. 7: 652–657, 2002.

[171053]

[171054]

[171055]17352.

[171056]Tomilin, A.; Remenyi, A.; Lins, K.; Bak, H.; Leidel, S.;
Vriend,

[171057]G.; Wilmanns, M.; Scholer, H. R.: Synergism with the coac-
tinator

[171058]OBF–1 (OCA–B, BOB–1) is mediated by a specific POU
dimer configuration. Cell 103:

[171059]853–864, 2000.

[171060]

[171061]

[171062]17353.

[171063]Ko, H.-S.; Fast, P.; McBride, W.; Staudt, L. M.: A human protein

[171064]specific for the immunoglobulin octamer DNA motif contains a functional

[171065]homeobox domain. Cell 55: 135-144, 1988.

[171066]

[171067]

[171068]17354.

[171069]Schubart, K.; Massa, S.; Schubart, D.; Corcoran, L. M.; Rolink,

[171070]A. G.; Matthias, P.: B cell development and immunoglobulin gene transcription

[171071]in the absence of Oct-2 and OBF-1. Nature Immun. 2: 69-74, 2001.

[171072]

[171073]

[171074]17355.

[171075]Staudt, L. M.; Clerc, R. G.; Singh, H.; LeBowitz, J. H.; Sharp,

[171076]P. A.; Baltimore, D.: Cloning of a lymphoid-specific cDNA

encoding

[171077]a protein binding the regulatory octamer DNA motif. Science 241:

[171078]577–580, 1988.

[171079]

[171080]

[171081]17356.

[171082]Crouau–Roy, B.; Amadou, C.; Bouissou, C.; Clayton, J.; Vernet,

[171083]C.; Ribouchon, M.–T.; Pontarotti, P.: Localization of the OTF3 gene

[171084]within the human MHC class I region by physical and meiotic mapping. Genomics 21:

[171085]241–243, 1994.

[171086]

[171087]

[171088]17357.

[171089]Guillaudeux, T.; Mattei, M. G.; Depetris, D.; Le Bouteiller, P.;

[171090]Pontarotti, P.: In situ hybridization localizes the human

OTF3 to

[171091]chromosome 6p21.3–p22 and OTF3L to 12p13. Cyto-
genet. Cell Genet. 63:

[171092]212–214, 1993.

[171093]

[171094]

[171095]17358.

[171096]Nichols, J.; Zevnik, B.; Anastassiadis, K.; Niwa, H.; Klewe-
Nebenius,

[171097]D.; Chambers, I.; Scholer, H.; Smith, A.: Formation of
pluripotent

[171098]stem cells in the mammalian embryo depends on the POU
transcription

[171099]factor Oct4. Cell 95: 379–391, 1998.

[171100]

[171101]

[171102]17359.

[171103]Niwa, H.; Miyazaki, J.; Smith, A. G.: Quantitative expres-
sion

[171104]of Oct–3/4 defines differentiation, dedifferentiation or

self-renewal

[171105]of ES cells. Nature Genet. 24: 372–376, 2000.

[171106]

[171107]

[171108]17360.

[171109]Rosner, M. H.; De Santo, R. J.; Arnheiter, H.; Staudt, L. M.:

[171110]Oct-3 is a maternal factor required for the first mouse
embryonic

[171111]division. Cell 64: 1103–1110, 1991. Note: Retraction: Cell
69: 724

[171112]only, 1992.

[171113]

[171114]

[171115]17361.

[171116]Takeda, J.; Seino, S.; Bell, G. I.: Human Oct3 gene family:
cDNA

[171117]sequences, alternative splicing, gene organization, chro-
mosomal location,

[171118]and expression at low levels in adult tissues. Nucleic Acids
Res. 20:

[171119]4613–4620, 1992.

[171120]

[171121]

[171122]17362.

[171123]Parker, R. C.; Mardon, G.; Lebo, R. V.; Varmus, H. E.;
Bishop,

[171124]J. M.: Isolation of duplicated human c-src genes located
on chromosomes

[171125]1 and 20. Molec. Cell. Biol. 5: 831–838, 1985.

[171126]

[171127]

[171128]17363.

[171129]Tronick, S. R.; Popescu, N. C.; Cheah, M. S. C.; Swan, D. C.;
Amsbaugh,

[171130]S. C.; Lengel, C. R.; DiPaolo, J. A.; Robbins, K. C.: Isolation
and

[171131]chromosomal localization of the human fgr protoonco-
gene, a distinct

[171132]member of the tyrosine kinase gene family. Proc. Nat.
Acad. Sci. 82:

[171133]6595–6599, 1985.

[171134]

[171135]

[171136]17364.

[171137]Krebs, L. T.; Xue, Y.; Norton, C. R.; Shutter, J. R.; Maguire,

[171138]M.; Sundberg, J. P.; Gallahan, D.; Closson, V.; Kitajewski,
J.; Callahan,

[171139]R.; Smith, G. H.; Stark, K. L.; Gridley, T.: Notch signaling is
essential

[171140]for vascular morphogenesis in mice. *Genes Dev.* 14:
1343–1352, 2000.

[171141]

[171142]

[171143]17365.

[171144]Li, L.; Huang, G. M.; Banta, A. B.; Deng, Y.; Smith, T.;
Dong,

[171145]P.; Friedman, C.; Chen, L.; Trask, B. J.; Spies, T.; Rowen, L.;
Hood,

[171146]L.: Cloning, characterization, and the complete
56.8–kilobase DNA

[171147]sequence of the human NOTCH4 gene. Genomics 51:
45–58, 1998.

[171148]

[171149]

[171150]17366.

[171151]McGinnis, R. E.; Fox, H.; Yates, P.; Cameron, L.–A.; Barnes,
M.

[171152]R.; Gray, I. C.; Spurr, N. K.; Hurko, O.; St. Clair, D.: Failure

[171153]to confirm NOTCH4 association with schizophrenia in a
large population–based

[171154]sample from Scotland. Nature Genet. 28: 128–129, 2001.

[171155]

[171156]

[171157]17367.

[171158]Sklar, P.; Schwab, S. G.; Williams, N. M.; Daly, M.;
Schaffner,

[171159]S.; Maier, W.; Albus, M.; Trixler, M.; Eichhammer, P.; Lerer,
B.;

[171160]Hallmayer, J.; Norton, N.; and 11 others: Association anal-
ysis

[171161]of NOTCH4 loci in schizophrenia using family and population-based

[171162]controls. Nature Genet. 28: 126–128, 2001.

[171163]

[171164]

[171165]17368.

[171166]Sugaya, K.; Sasanuma, S.; Nohata, J.; Kimura, T.; Fukagawa, T.;

[171167]Nakamura, Y.; Ando, A.; Inoko, H.; Ikemura, T.; Mita, K.:
Gene organization

[171168]of human NOTCH4 and (CTG)_n polymorphism in this human counterpart

[171169]gene of mouse proto-oncogene Int3. Gene 189: 235–244, 1997.

[171170]

[171171]

[171172]17369.

[171173]Uyttendaele, H.; Marazzi, G.; Wu, G.; Yan, Q.; Sassoon, D.; Kitajewski,

[171174]J.: Notch4/int-3, a mammary proto-oncogene, is an endothelial cell-specific

[171175]mammalian Notch gene. Development 122: 2251–2259,
1996.

[171176]

[171177]

[171178]17370.

[171179]Wei, J.; Hemmings, G. P.: The NOTCH4 locus is associated
with

[171180]susceptibility to schizophrenia. (Letter) Nature Genet. 25:
376–377,

[171181]2000.

[171182]

[171183]

[171184]17371.

[171185]Bork, P.: The modular architecture of a new family of
growth regulators

[171186]related to connective tissue growth factor. FEBS Lett. 327:
125–130,

[171187]1993.

[171188]

[171189]

[171190]17372.

[171191]Sugaya, K.; Fukagawa, T.; Matsumoto, K.; Mita, K.; Takahashi, E.;

[171192]Ando, A.; Inoko, H.; Ikemura, T.: Three genes in the human MHC class

[171193]III region near the junction with the class II: gene for receptor

[171194]of advanced glycosylation end products, PBX2 homeobox gene and a Notch

[171195]homolog, human counterpart of mouse mammary tumor gene int-3. Genomics 23:

[171196]408-419, 1994.

[171197]

[171198]

[171199]17373.

[171200]Joliot, V.; Martinerie, C.; Dambrine, G.; Plassiart, G.; Brisac,

[171201]M.; Crochet, J.; Perbal, B.: Proviral rearrangements and overexpression

[171202]of a new cellular gene (nov) in myeloblastosis-associated virus type

[171203]1-induced nephroblastomas. Molec. Cell. Biol. 12: 10-21,

1992.

[171204]

[171205]

[171206]17374.

[171207]Kim, H.-S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T.,

[171208]Jr.; Rosenfeld, R. G.: Identification of a family of low-
affinity

[171209]insulin-like growth factor binding proteins (IFGBPs): char-
acterization

[171210]of connective tissue growth factor as a member of the
IGFBP superfamily. Proc.

[171211]Nat. Acad. Sci. 94: 12981–12986, 1997.

[171212]

[171213]

[171214]17375.

[171215]Martinerie, C.; Perbal, B.: Expression of a gene encoding a novel

[171216]potential IGF binding protein in human tissues. C. R. Acad. Sci.

[171217](Paris) 313 (ser. 3): 345–351, 1991.

[171218]

[171219]

[171220]17376.

[171221]Snaith, M. R.; Natarajan, D.; Taylor, L. B.; Choi, C.–P.; Martinerie,

[171222]C.; Perbal, B.; Schofield, P. N.; Boulter, C. A.: Genomic structure

[171223]and chromosomal mapping of the mouse nov gene. Genomics 38: 425–428,

[171224]1996.

[171225]

[171226]

[171227]17377.

[171228]Soret, J.; Dambrine, G.; Perbal, B.: Induction of nephroblastoma

[171229]by myeloblastosis-associated virus type 1: state of proviral DNAs

[171230]in tumor cells. J. Virol. 63: 1803–1807, 1989.

[171231]

[171232]

[171233]17378.

[171234]Amson, R.; Sigaux, F.; Przedborski, S.; Flandrin, G.; Givol, D.;

[171235]Telerman, A.: The human protooncogene product p33pim is expressed

[171236]during fetal hematopoiesis and in diverse leukemias. Proc. Nat. Acad.

[171237]Sci. 86: 8857–8861, 1989.

[171238]

[171239]

[171240]17379.

[171241]Ark, B.; Gummere, G.; Bennett, D.; Artzt, K.: Mapping of
the Pim-1

[171242]oncogene in mouse t-haplotypes and its use to define the
relative

[171243]map positions of the tcl loci t-0(t-6) and t-w12 and the
marker tf

[171244](tufted). Genomics 10: 385-389, 1991.

[171245]

[171246]

[171247]17380.

[171248]Cuypers, H. T.; Selten, G.; Berns, A.; Geurts van Kessel, A.
H.

[171249]M.: Assignment of the human homologue of Pim-1, a
mouse gene implicated

[171250]in leukemogenesis, to the pter-q12 region of chromo-
some 6. Hum. Genet. 72:

[171251]262-265, 1986.

[171252]

[171253]

[171254]17381.

[171255]Domen, J.; von Lindern, M.; Hermans, A.; Breuer, M.;
Grosveld,

[171256]G.; Berns, A.: Comparison of the human and mouse PIM-1
cDNAs: nucleotide

[171257]sequence and immunological identification of the in vitro
synthesized

[171258]PIM-1 protein. Oncogene Res. 1: 103-112, 1987.

[171259]

[171260]

[171261]17382.

[171262]Ben Othmane, K.; Loeb, D.; Hayworth-Hodgte, R.; Hentati,
F.; Rao,

[171263]N.; Roses, A. D.; Ben Hamida, M.; Pericak-Vance, M. A.;
Vance, J.

[171264]M.: Physical and genetic mapping of the CMT4A locus and
exclusion

[171265]of PMP-2 as the defect in CMT4A. Genomics 28: 286-290,
1995.

[171266]

[171267]

[171268]17383.

- [171269]Hayasaka, K.; Himoro, M.; Takada, G.; Takahashi, E.; Minoshima,
[171270]S.; Shimizu, N.: Structure and localization of the gene encoding
[171271]human peripheral myelin protein 2 (PMP2). Genomics 18:
244–248,
[171272]1993.
- [171273]
- [171274]
- [171275]17384.
- [171276]Hayasaka, K.; Nanao, K.; Tahara, M.; Sato, W.; Takada, G.;
Miura,
[171277]M.; Uyemura, K.: Isolation and sequence determination of
cDNA encoding
[171278]P2 protein of human peripheral myelin. Biochem. Biophys.
Res. Commun. 181:
[171279]204–207, 1991.
- [171280]
- [171281]
- [171282]17385.

[171283]Narayanan, V.; Ripepi, B.; Jabs, E. W.; Hawkins, A.; Griffin, C.;

[171284]Tennekoon, G.: Partial structure and mapping of the human myelin

[171285]P2 protein gene. J. Neurochem. 63: 2010–2013, 1994.

[171286]

[171287]

[171288]17386.

[171289]Manser, E.; Loo, T.-H.; Koh, C.-G.; Zhao, Z.-S.; Chen, X.-Q.; Tan,

[171290]L.; Tan, I.; Leung, T.; Lim, L.: PAK kinases are directly coupled

[171291]to the PIX family of nucleotide exchange factors. Molec. Cell 1:

[171292]183–192, 1998.

[171293]

[171294]

[171295]17387.

[171296]Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Kawarabayashi,

[171297]Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.:

Prediction

[171298]of the coding sequences of unidentified human genes. I.

The coding

[171299]sequences of 40 new genes (KIAA0001–KIAA0040) deduced by analysis

[171300]of randomly sampled cDNA clones from human immature myeloid cell line,

[171301]KG-1. DNA Res. 1: 27–35, 1994.

[171302]

[171303]

[171304]17388.

[171305]Buggy, J. J.; Sideris, M. L.; Mak, P.; Lorimer, D. D.; McIntosh,

[171306]B.; Clark, J. M.: Cloning and characterization of a novel human histone

[171307]deacetylase, HDAC8. Biochem. J. 350: 199–205, 2000.

[171308]

[171309]

[171310]17389.

[171311]Hu, E.; Chen, Z.; Fredrickson, T.; Zhu, Y.; Kirkpatrick, R.;
Zhang,

[171312]G.-F.; Johanson, K.; Sung, C.-M.; Liu, R.; Winkler, J.:
Cloning and

[171313]characterization of a novel human class I histone deacetylase that

[171314]functions as a transcription repressor. J. Biol. Chem. 275:
15254–15264,

[171315]2000.

[171316]

[171317]

[171318]17390.

[171319]Van den Wyngaert, I.; de Vries, W.; Kremer, A.; Neefs,
J.-M.; Verhasselt,

[171320]P.; Luyten, W. H. M. L.; Kass, S. U.: Cloning and characteri-
zation

[171321]of human histone deacetylase 8. FEBS Lett. 478: 77–83,
2000.

[171322]

[171323]

[171324]17391.

[171325]Grozinger, C. M.; Hassig, C. A.; Schreiber, S. L.: Three proteins

[171326]define a class of human histone deacetylases related to yeast Hda1p. Proc.

[171327]Nat. Acad. Sci. 96: 4868–4873, 1999.

[171328]

[171329]

[171330]17392.

[171331]Hubbert, C.; Guardiola, A.; Shao, R.; Kawaguchi, Y.; Ito, A.; Nixon,

[171332]A.; Yoshida, M.; Wang, X.-F.; Yao, T.-P.: HDAC6 is a microtubule-associated

[171333]deacetylase. Nature 417: 455–458, 2002.

[171334]

[171335]

[171336]17393.

[171337]Mahlknecht, U.; Schnittger, S.; Landgraf, F.; Schoch, C.; Ottmann,

[171338]O. G.; Hiddemann, W.; Hoelzer, D.: Assignment of the human histone

[171339]deacetylase 6 gene (HDAC6) to X chromosome p11.23 by
in situ hybridization. Cytogenet.

[171340]Cell Genet. 93: 135–136, 2001.

[171341]

[171342]

[171343]17394.

[171344]Nagase, T.; Isikawa, K.; Suyama, M.; Kikuno, R.; Hirose, M.;
M.;

[171345]Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara,
O.: Prediction

[171346]of the coding sequences of unidentified human genes. XII.
The complete

[171347]sequences of 100 new cDNA clones from brain which code
for large proteins

[171348]in vitro. DNA Res. 5: 355–364, 1998.

[171349]

[171350]

[171351]17395.

[171352]Fortenberry, Y.; Hwang, J.-R.; Apletalina, E. V.; Lindberg, I.

[171353]: Functional characterization of proSAAS: similarities and

differences

[171354]with 7B2. J. Biol. Chem. 277: 5175–5186, 2002.

[171355]

[171356]

[171357]17396.

[171358]Fricker, L. D.; McKinzie, A. A.; Sun, J.; Curran, E.; Qian, Y.;

[171359]Yan, L.; Patterson, S. D.; Courchesne, P. L.; Richards, B.;

Levin,

[171360]N.; Mzhavia, N.; Devi, L. A.; Douglass, J.: Identification and
characterization

[171361]of proSAAS, a granin-like neuroendocrine peptide precursor that inhibits

[171362]prohormone processing. J. Neurosci. 20: 639–648, 2000.

[171363]

[171364]

[171365]17397.

[171366]Disteche, C. M.; Brannan, C. I.; Larsen, A.; Adler, D. A.;

Schorderet,

[171367]D. F.; Gearing, D.; Copeland, N. G.; Jenkins, N. A.; Park, L.

S.:

[171368]The human pseudoautosomal GM-CSF receptor alpha subunit gene is autosomal

[171369]in mouse. Nature Genet. 1: 333–336, 1992.

[171370]

[171371]

[171372]17398.

[171373]Sugarman, G. I.; Rimoin, D. L.; Lachman, R. S.: The facial–digital–genital

[171374](Aarskog) syndrome. Am. J. Dis. Child. 126: 248–252, 1973.

[171375]

[171376]

[171377]17399.

[171378]Tsukahara, M.; Fernandez, G. I.: Umbilical findings in Aarskog

[171379]syndrome. Clin. Genet. 45: 260–265, 1994.

[171380]

[171381]

[171382]17400.

[171383]Tyrkus, M.; Bawle, E.; Lipman, S.; Bozimowski, D.; Woolley, P.

[171384]V., Jr.: Aarskog–Scott syndrome inherited as an X–linked dominant

[171385]with full male–female expression. (Abstract) Am. J. Hum. Genet. 32:

[171386]134A only, 1980.

[171387]

[171388]

[171389]17401.

[171390]van den Bergh, P.; Fryns, J. P.; Wilms, G.; Piot, R.; Dralands,

[171391]G.; van den Bergh, R.: Anomalous cerebral venous drainage in Aarskog

[171392]syndrome. Clin. Genet. 25: 288–294, 1984.

[171393]

[171394]

[171395]17402.

[171396]Zheng, Y.; Fischer, D. J.; Santos, M. F.; Tigyi, G.; Pasteris,

[171397]N. G.; Gorski, J. L.; Xu, Y.: The faciogenital dysplasia gene

product

[171398]FGD1 functions as a Cdc42Hs-specific guanine-nucleotide exchange factor. J.

[171399]Biol. Chem. 271: 33169–33172, 1996.

[171400]

[171401]

[171402]17403.

[171403]Inaba, H.; Fujimaki, M.; Kazazian, H. H., Jr.; Antonarakis, S.

[171404]E.: MspI polymorphic site in intron 22 of the factor VIII gene in

[171405]the Japanese population. Hum. Genet. 84: 214–215, 1990.

[171406]

[171407]

[171408]17404.

[171409]Levinson, B.; Bermingham, J. R., Jr.; Metzenberg, A.; Ken-
wick,

[171410]S.; Chapman, V.; Gitschier, J.: Sequence of the human fac-
tor VIII-associated

[171411]gene is conserved in mouse. Genomics 13: 862–865,
1992.

[171412]

[171413]

[171414]17405.

[171415]Levinson, B.; Kenwrick, S.; Lakich, D.; Hammonds, G.;
Gitschier,

[171416]J.: A transcribed gene in an intron of the human factor VIII
gene. Genomics 7:

[171417]1–11, 1990.

[171418]

[171419]

[171420]17406.

[171421]Peters, M. F.; Ross, C. A.: Isolation of a 40–kDa hunt-
ingtin-associated

[171422]protein. J. Biol. Chem. 276: 3188–3194, 2001.

[171423]

[171424]

[171425]17407.

[171426]Berra, B.; Gornati, R.; Rapelli, S.; Gatti, R.; Mancini, G. M.

[171427]S.; Ciana, G.; Bembi, B.: Infantile sialic acid storage dis-

ease:

[171428]biochemical studies. Am. J. Med. Genet. 58: 24–31, 1995.

[171429]

[171430]

[171431]17408.

[171432]Cameron, P. D.; Dubowitz, V.; Besley, G. T. N.; Fensom, A.
H.:

[171433]Sialic acid storage disease. Arch. Dis. Child. 65: 314–315,
1990.

[171434]

[171435]

[171436]17409.

[171437]Haataja, L.; Schleutker, J.; Laine, A.–P.; Renlund, M.;
Savontaus,

[171438]M.–L.; Dib, C.; Weissenbach, J.; Peltonen, L.; Aula, P.: The
genetic

[171439]locus for free sialic acid storage disease maps to the long
arm of

[171440]chromosome 6. Am. J. Hum. Genet. 54: 1042–1049, 1994.

[171441]

[171442]

[171443]17410.

[171444]Fojo, A.; Lebo, R.; Shimizu, N.; Chin, J. E.; Roninson, I. B.;

[171445]Merlino, G. T.; Gottesman, M. M.; Pastan, I.: Localization
of multidrug

[171446]resistance-associated DNA sequences to human chromo-
some 7. Somat.

[171447]Cell Molec. Genet. 12: 415-420, 1986.

[171448]

[171449]

[171450]17411.

[171451]Fojo, A. T.; Ueda, K.; Slamon, D. J.; Poplack, D. G.; Gottes-
man,

[171452]M. M.; Pastan, I.: Expression of a multidrug-resistance
gene in human

[171453]tumors and tissues. Proc. Nat. Acad. Sci. 84: 265-269,
1987.

[171454]

[171455]

[171456]17412.

[171457]Fujii, J.; Zarain-Herzberg, A.; Willard, H. F.; Tada, M.;
MacLennan,

[171458]D. H.: Structure of the rabbit phospholamban gene,
cloning of the

[171459]human cDNA, and assignment of the gene to human
chromosome 6. J.

[171460]Biol. Chem. 266: 11669–11675, 1991.

[171461]

[171462]

[171463]17413.

[171464]McTiernan, C. F.; Frye, C. S.; Lemster, B. H.; Kinder, E. A.;
Ogletree-Hughes,

[171465]M. L.; Moravec, C. S.; Feldman, A. M.: The human phos-
pholamban gene:

[171466]structure and expression. J. Molec. Cell Cardiol. 31:
679–692, 1999.

[171467]

[171468]

[171469]17414.

[171470]Otsu, K.; Fujii, J.; Periasamy, M.; Difilippantonio, M.; Up-
pender,

[171471]M.; Ward, D. C.; MacLennan, D. H.: Chromosome mapping
of five human

[171472]cardiac and skeletal muscle sarcoplasmic reticulum pro-
tein genes. Genomics 17:

[171473]507–509, 1993.

[171474]

[171475]

[171476]17415.

[171477]Barhanin, J.; Lesage, F.; Guillemare, E.; Fink, M.; Lazdun-
ski,

[171478]M.; Romey, G.: K(v)LQT1 and IsK (mink) proteins associate
to form

[171479]the I(Ks) cardiac potassium current. Nature 384: 78–80,
1996.

[171480]

[171481]

[171482]17416.

[171483]Marx, S. O.; Kurokawa, J.; Reiken, S.; Motoike, H.;
D'Armiento,

[171484]J.; Marks, A. R.; Kass, R. S.: Requirement of a macromolec-
ular signaling

[171485]complex for beta adrenergic receptor modulation of the
KCNQ1–KCNE1

[171486]potassium channel. Science 295: 496–499, 2002.

[171487]

[171488]

[171489]17417.

[171490]Sanguinetti, M. C.; Curran, M. E.; Zou, A.; Shen, J.; Spector,

[171491]P. S.; Atkinson, D. L.; Keating, M. T.: Coassembly of
K(v)LQT1 and

[171492]minK (IsK) proteins to form cardiac I(Ks) potassium chan-
nel. Nature 384:

[171493]80–83, 1996.

[171494]

[171495]

[171496]17418.

[171497]Coleman, R. A.; Smith, W. L.; Narumiya, S.: VIII. Interna-
tional

[171498]union of pharmacology classification of prostanoid recep-
tors: properties,

[171499]distribution, and structure of the receptors and their sub-

types. Pharm.

[171500]Rev. 46: 205–229, 1994.

[171501]

[171502]

[171503]17419.

[171504]Duncan, A. M. V.; Anderson, L. L.; Funk, C. D.;
Abramovitz, M.;

[171505]Adam, M.: Chromosomal localization of the human
prostanoid receptor

[171506]gene family. Genomics 25: 740–742, 1995.

[171507]

[171508]

[171509]17420.

[171510]Funk, C. D.; Furci, L.; FitzGerald, G. A.; Grygorczyk, R.;
Rochette,

[171511]C.; Bayne, M. A.; Abramovitz, M.; Adam, M.; Metters, K.
M.: Cloning

[171512]and expression of a cDNA for the human prostaglandin E
receptor EP–1

[171513]subtype. J. Biol. Chem. 268: 26767–26772, 1993.

[171514]

[171515]

[171516]17421.

[171517]Nagata, A.; Suzuki, Y.; Igarashi, M.; Eguchi, N.; Toh, H.;
Urade,

[171518]Y.; Hayaishi, O.: Human brain prostaglandin D synthase
has been evolutionarily

[171519]differentiated from lipophilic–ligand carrier proteins. Proc.
Nat.

[171520]Acad. Sci. 88: 4020–4024, 1991.

[171521]

[171522]

[171523]17422.

[171524]Onoe, H.; Ueno, R.; Fujita, I.; Nishino, H.; Oomura, Y.;
Hayashi, O.

[171525]:Proc. Nat. Acad. Sci. 85: 4082–4086, 1988.

[171526]

[171527]

[171528]17423.

[171529]Pinzar, E.; Kanaoka, Y.; Inui, T.; Eguchi, N.; Urade, Y.;

Hayaishi,

[171530]O.: Prostaglandin D synthase gene is involved in the regulation of

[171531]non-rapid eye movement sleep. Proc. Nat. Acad. Sci. 97: 4903-4907,

[171532]2000.

[171533]

[171534]

[171535]17424.

[171536]Tanaka, T.; Urade, Y.; Kimura, H.; Eguchi, N.; Nishikawa, A.; Hayaishi,

[171537]O.: Lipocalin-type prostaglandin D synthase (beta-trace) is a newly

[171538]recognized type of retinoid transporter. J. Biol. Chem. 272: 15789-15795,

[171539]1997.

[171540]

[171541]

[171542]17425.

[171543]White, D. M.; Mikol, D. D.; Espinosa, R.; Weimer, B.; Le
Beau,

[171544]M. M.; Stefansson, K.: Structure and chromosomal local-
ization of

[171545]the human gene for a brain form of prostaglandin D-2
synthase. J.

[171546]Biol. Chem. 267: 23202-23208, 1992.

[171547]

[171548]

[171549]17426.

[171550]Sales, K. J.; Katz, A. A.; Davis, M.; Hinz, S.; Soeters, R. P.;

[171551]Hofmeyr, M. D.; Millar, R. P.; Jabbour, H. N.: Cyclooxyge-
nase-2 expression

[171552]and prostaglandin E2 synthesis are up-regulated in carci-
nomas of the

[171553]cervix: a possible autocrine/paracrine regulation of neo-
plastic cell

[171554]function via EP2/EP4 receptors. J. Clin. Endocr. Metab. 86:
2243-2249,

[171555]2001.

[171556]

[171557]

[171558]17427.

[171559]Smock, S. L.; Pan, L. C.; Castleberry, T. A.; Lu, B.; Mather, R.

[171560]J.; Owen, T. A.: Cloning, structural characterization, and chromosomal

[171561]localization of the gene encoding the human prostaglandin E2 receptor

[171562]EP2 subtype. *Gene* 237: 393–402, 1999.

[171563]

[171564]

[171565]17428.

[171566]Taketo, M.; Rochelle, J. M.; Sugimoto, Y.; Namba, T.; Honda, A.;

[171567]Negishi, M.; Ichikawa, A.; Narumiya, S.; Seldin, M. F.: Mapping of

[171568]the genes encoding mouse thromboxane A2 receptor and prostaglandin

[171569]E receptor subtypes EP2 and EP3. *Genomics* 19: 585–588, 1994.

[171570]

[171571]

[171572]17429.

[171573]Tilley, S. L.; Audoly, L. P.; Hicks, E. H.; Kim, H.-S.; Flannery,

[171574]P. J.; Coffman, T. M.; Koller, B. H.: Reproductive failure and reduced

[171575]blood pressure in mice lacking the EP2 prostaglandin E-2 receptor. J.

[171576]Clin. Invest. 103: 1539–1545, 1999.

[171577]

[171578]

[171579]17430.

[171580]Catella-Lawson, F.; Reilly, M. P.; Kapoor, S. C.; Cucchiara, A.

[171581]J.; DeMarco, S.; Tournier, B.; Vyas, S. N.; FitzGerald, G. A.: Cyclooxygenase

[171582]inhibitors and the antiplatelet effects of aspirin. New Eng. J. Med. 345:

[171583]1809–1817, 2001.

[171584]

[171585]

[171586]17431.

[171587]Dube, J.-N.; Drouin, J.; Aminian, M.; Plant, M. H.; La-neuville,

[171588]O.: Characterization of a partial prostaglandin endoperox-
ide H synthase-1

[171589]deficiency in a patient with a bleeding disorder. Brit. J.
Haemat. 113:

[171590]878-885, 2001.

[171591]

[171592]

[171593]17432.

[171594]Funk, C. D.; Funk, L. B.; Kennedy, M. E.; Pong, A. S.;
Fitzgerald,

[171595]G. A.: Human platelet/erythroleukemia cell prostaglandin
G/H synthase:

[171596]cDNA cloning, expression, and gene chromosomal as-
signment. FASEB

[171597]J. 5: 2304-2312, 1991.

[171598]

[171599]

[171600]17433.

[171601]Gavett, S. H.; Madison, S. L.; Chulada, P. C.; Scarborough, P.

[171602]E.; Qu, W.; Boyle, J. E.; Tiano, H. F.; Lee, C. A.; Langenbach, R.;

[171603]Roggli, V. L.; Zeldin, D. C.: Allergic lung responses are increased

[171604]in prostaglandin H synthase-deficient mice. J. Clin. Invest. 104:

[171605]721-732, 1999.

[171606]

[171607]

[171608]17434.

[171609]Kirschenbaum, A.; Liotta, D. R.; Yao, S.; Liu, X.-H.; Klausner,

[171610]A. P.; Unger, P.; Shapiro, E.; Leav, I.; Levine, A. C.: Immunohistochemical

[171611]localization of cyclooxygenase-1 and cyclooxygenase-2 in the human

[171612]fetal and adult male reproductive tracts. J. Clin. Endocr.

Metab. 85:

[171613]3436–3441, 2000.

[171614]

[171615]

[171616]17435.

[171617]Langenbach, R.; Morham, S. G.; Tiano, H. F.; Loftin, C. D.;
Ghanayem,

[171618]B. I.; Chulada, P. C.; Mahler, J. F.; Lee, C. A.; Goulding, E.
H.;

[171619]Kluckman, K. D.; Kim, H. S.; Smithies, O.: Prostaglandin
synthase

[171620]1 gene disruption in mice reduces arachidonic acid–
induced inflammation

[171621]and indomethacin–induced gastric ulceration. Cell 83:
483–492, 1995.

[171622]

[171623]

[171624]17436.

[171625]Malkowski, M. G.; Ginell, S. L.; Smith, W. L.; Garavito, R. M.

[171626]: The productive conformation of arachidonic acid bound

to prostaglandin

[171627]synthase. Science 289: 1933–1937, 2000.

[171628]

[171629]

[171630]17437.

[171631]Pareti, F. I.; Manucci, P. M.; D'Angelo, A.; Smith, J. B.;
Sautebin,

[171632]L.; Galli, G.: Congenital deficiency of thromboxane and
prostacyclin. Lancet I:

[171633]898–900, 1980.

[171634]

[171635]

[171636]17438.

[171637]Picot, D.; Loll, P. J.; Garavito, R. M.: The x-ray crystal
structure

[171638]of the membrane protein prostaglandin H-2 synthase-1.
Nature 367:

[171639]243–249, 1994.

[171640]

[171641]

[171642]17439.

[171643]Niswender, K. D.; Morton, G. J.; Stearns, W. H.; Rhodes, C. J.;

[171644]Myers, M. G., Jr.; Schwartz, M. W.: Key enzyme in leptin-induced

[171645]anorexia. Nature 413: 794–795, 2001.

[171646]

[171647]

[171648]17440.

[171649]Shimomura, I.; Matsuda, M.; Hammer, R. E.; Bashmakov, Y.; Brown,

[171650]M. S.; Goldstein, J. L.: Decreased IRS–2 and increased SREBP–1c lead

[171651]to mixed insulin resistance and sensitivity in livers of lipodystrophic

[171652]and ob/ob mice. Molec. Cell 6: 77–86, 2000.

[171653]

[171654]

[171655]17441.

[171656]Siracusa, L. D.; Rosner, M. H.; Vigano, M. A.; Gilbert, D. J.;

[171657]Staudt, L. M.; Copeland, N. G.; Jenkins, N. A.: Chromosomal location

[171658]of the octamer transcription factors, Otf-1, Otf-2, and Otf-3, defines

[171659]multiple Otf-3-related sequences dispersed in the mouse genome. Genomics 10:

[171660]313-326, 1991.

[171661]

[171662]

[171663]17442.

[171664]Sturm, R. A.; Eyre, H. J.; Baker, E.; Sutherland, G. R.: The human

[171665]OTF1 locus which overlaps the CD3Z gene is located at 1q22-q23. Cytogenet.

[171666]Cell Genet. 68: 231-232, 1995.

[171667]

[171668]

[171669]17443.

[171670]Marshall, J. B.; Diaz-Arias, A. A.; Bochna, G. S.; Vogele, K.

[171671]A.: Achalasia due to diffuse esophageal leiomyomatosis
and inherited

[171672]as an autosomal dominant disorder. Gastroenterology 98:
1358–1365,

[171673]1990.

[171674]

[171675]

[171676]17444.

[171677]Mattei, M. G.; d'Auriol, L.; Andre, C.; Passage, E.; Mattei, J.

[171678]F.; Galibert, F.: Assignment of the human c-kit proto-
oncogene to

[171679]the q11–q12 region of chromosome 4, using in situ hy-
bridization. (Abstract) Cytogenet.

[171680]Cell Genet. 46: 657, 1987.

[171681]

[171682]

[171683]17445.

[171684]Nagata, H.; Worobec, A. S.; Oh, C. K.; Chowdhury, B. A.;
Tannenbaum,

[171685]S.; Suzuki, Y.; Metcalfe, D. D.: Identification of a point mu-

tation

[171686]in the catalytic domain of the protooncogene c-kit in peripheral blood

[171687]mononuclear cells of patients who have mastocytosis with an associated

[171688]hematologic disorder. Proc. Nat. Acad. Sci. 92: 10560–10564, 1995.

[171689]

[171690]

[171691]17446.

[171692]Nishida, T.; Hirota, S.; Taniguchi, M.; Hashimoto, K.; Isozaki,

[171693]K.; Nakamura, H.; Kanakura, Y.; Tanaka, T.; Takabayashi, A.; Matsuda,

[171694]H.; Kitamura, Y.: Familial gastrointestinal stromal tumours with

[171695]germline mutation of the KIT gene. (Letter) Nature Genet. 19: 323–324,

[171696]1998.

[171697]

[171698]

[171699]17447.

[171700]Nocka, K.; Tan, J. C.; Chiu, E.; Chu, T. Y.; Ray, P.; Traktman,

[171701]P.; Besmer, P.: Molecular bases of dominant negative and loss of

[171702]function mutations at the murine c-kit/white spotting locus: W-37,

[171703]W-v, W-41 and W. EMBO J. 9: 1805-1813, 1990.

[171704]

[171705]

[171706]17448.

[171707]Nomura, K.; Hatayama, I.; Narita, T.; Kaneko, T.; Shiraishi, M.

[171708]: A novel KIT gene missense mutation in a Japanese family with piebaldism.

[171709](Letter) J. Invest. Derm. 111: 337-338, 1998.

[171710]

[171711]

[171712]17449.

[171713]Pignon, J.-M.; Giraudier, S.; Duquesnoy, P.; Jouault, H.;

Imbert,

[171714]M.; Vainchenker, W.; Vernant, J.-P.; Tulliez, M.: A new c-kit mutation

[171715]in a case of aggressive mast cell disease. Brit. J. Haemat. 96:

[171716]374–376, 1997.

[171717]

[171718]

[171719]17450.

[171720]Reinsch, N.; Thomsen, H.; Xu, N.; Brink, M.; Looft, C.; Kalm,

[171721]E.; Brockmann, G. A.; Grupe, S.; Kuhn, C.; Schwerin, M.; Leyhe, B.;

[171722]Hiendleder, S.; Erhardt, G.; Medjugorac, I.; Russ, I.; Forster, M.;

[171723]Reents, R.; Averdunk, G.: A QTL for the degree of spotting in cattle

[171724]shows synteny with the KIT locus on chromosome 6. J. Hered. 90:

[171725]629–634, 1999.

[171726]

[171727]

[171728]17451.

[171729]Selmanowitz, V. J.; Rabinowitz, A. D.; Orentreich, N.;
Wenk, E.

[171730]: Pigmentary correction of piebaldism by autografts. I.
Procedures

[171731]and clinical findings. J. Derm. Surg. Oncol. 3: 615–622,
1977.

[171732]

[171733]

[171734]17452.

[171735]Spritz, R. A.; Beighton, P.: Piebaldism with deafness:
molecular

[171736]evidence for an expanded syndrome. Am. J. Med. Genet.
75: 101–103,

[171737]1998.

[171738]

[171739]

[171740]17453.

[171741]Spritz, R. A.; Droetto, S.; Fukushima, Y.: Deletion of the

KIT

[171742]and PDGFRA genes in a patient with piebaldism. Am. J.

Med. Genet. 44:

[171743]492–495, 1992.

[171744]

[171745]

[171746]17454.

[171747]Spritz, R. A.; Giebel, L. B.: Mutation of the c-kit

(mast/stem

[171748]cell growth factor receptor) proto-oncogene in human

piebaldism. (Abstract) Am.

[171749]J. Hum. Genet. 49 (suppl.): 38, 1991.

[171750]

[171751]

[171752]17455.

[171753]Spritz, R. A.; Giebel, L. B.; Holmes, S. A.: Dominant nega-

tive

[171754]and loss of function mutations of the c-kit (mast/stem

cell growth

[171755]factor receptor) proto-oncogene in human piebaldism.

Am. J. Hum.

[171756]Genet. 50: 261–269, 1992.

[171757]

[171758]

[171759]17456.

[171760]Spritz, R. A.; Holmes, S. A.; Ramesar, R.; Greenberg, J.;
Curtis,

[171761]D.; Beighton, P.: Mutations of the KIT (mast/stem cell
growth factor

[171762]receptor) proto–oncogene account for a continuous range
of phenotypes

[171763]in human piebaldism. Am. J. Hum. Genet. 51: 1058–1065,
1992.

[171764]

[171765]

[171766]17457.

[171767]Syrris, P.; Malik, N. M.; Murday, V. A.; Patton, M. A.; Carter,

[171768]N. D.; Hughes, H. E.; Metcalfe, K.: Three novel mutations
of the

[171769]proto–oncogene KIT cause human piebaldism. (Letter) Am.

J. Med. Genet. 95:

[171770]79–81, 2000.

[171771]

[171772]

[171773]17458.

[171774]Tan, J. C.; Nocka, K.; Ray, P.; Traktman, P.; Besmer, P.: The

[171775]dominant W42 spotting phenotype results from a mis-
sense mutation in

[171776]the c-kit receptor kinase. Science 247: 209–212, 1990.

[171777]

[171778]

[171779]17459.

[171780]Taylor, M. L.; Dastych, J.; Sehgal, D.; Sundstrom, M.; Nils-
son,

[171781]G.; Akin, C.; Mage, R. G.; Metcalfe, D. D.: The Kit-
activating mutation

[171782]D816V enhances stem cell factor-dependent chemotaxis.

Blood 98:

[171783]1195–1199, 2001.

[171784]

[171785]

[171786]17460.

[171787]Thomsen, L.; Robinson, T. L.; Lee, J. C. F.; Farraway, L. A.;

[171788]Hughes, M. J. G.; Andrews, D. W.; Huizinga, J. D.: Interstitial cells
of Cajal generate a rhythmic pacemaker current. Nature

Med. 4: 848–851,
[171790]1998.

[171791]

[171792]

[171793]17461.

[171794]Tian, Q.; Frierson, H. F., Jr.; Krystal, G. W.; Moskaluk, C. A.

[171795]: Activating c-kit gene mutations in human germ cell tumors. Am.

[171796]J. Path. 154: 1643–1647, 1999.

[171797]

[171798]

[171799]17462.

[171800]Tsujimura, T.; Furitsu, T.; Morimoto, M.; Isozaki, K.; No-
mura,

[171801]S.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Ligand-
independent

[171802]activation of c-kit receptor tyrosine kinase in a murine
mastocytoma

[171803]cell line P-815 generated by a point mutation. Blood 83:
2619-2626,

[171804]1994.

[171805]

[171806]

[171807]17463.

[171808]Vandenbark, G. R.; deCastro, C. M.; Taylor, H.; Dew-
Knight, S.;

[171809]Kaufman, R. E.: Cloning and structural analysis of the hu-
man c-kit

[171810]gene. Oncogene 7: 1259-1266, 1992.

[171811]

[171812]

[171813]17464.

[171814]Worobec, A. S.; Semere, T.; Nagata, H.; Metcalfe, D. D.:

Clinical

[171815]correlates of the presence of the asp816val c-kit mutation
in the

[171816]peripheral blood mononuclear cells of patients with mas-
tocyctosis. Cancer 83:

[171817]2120–2129, 1998.

[171818]

[171819]

[171820]17465.

[171821]Yarden, Y.; Kuang, W.-J.; Yang-Feng, T.; Coussens, L.;
Munemitsu,

[171822]S.; Dull, T. J.; Chen, E.; Schlessinger, J.; Francke, U.; Ull-
rich,

[171823]A.: Human proto-oncogene c-kit: a new cell surface re-
ceptor tyrosine

[171824]kinase for an unidentified ligand. EMBO J. 6: 3341–3351,
1987.

[171825]

[171826]

[171827]17466.

[171828]Cheah, M. S. C.; Ley, T. J.; Tronick, S. R.; Robbins, K. C.:

Fgr

[171829]proto-oncogene mRNA induced in B lymphocytes by Epstein-Barr virus

[171830]infection. Nature 319: 238-240, 1986.

[171831]

[171832]

[171833]17467.

[171834]Dracopoli, N. C.; Stanger, B. Z.; Lager, M.; Housman, D. E.:

Localization

[171835]of the FGR protooncogene on the genetic linkage map of human chromosome

[171836]1p. Genomics 3: 124-128, 1988.

[171837]

[171838]

[171839]17468.

[171840]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J.

D.:

[171841]Evidence for two distinct c-src loci on human chromosomes 1 and 20. Nature 312:

[171842]70–71, 1984.

[171843]

[171844]

[171845]17469.

[171846]Lebo, R. V.; Cheung, M.–C.; Bruce, B. D.: Rapid gene mapping by

[171847]dual laser chromosome sorting and spot blot DNA analysis. (Abstract) Am.

[171848]J. Hum. Genet. 36: 101S, 1984.

[171849]

[171850]

[171851]17470.

[171852]Nishizawa, M.; Semba, K.; Yoshida, M. C.; Yamamoto, T.; Sasaki,

[171853]M.; Toyoshima, K.: Structure, expression, and chromosomal location

[171854]of the human c-fgr gene. Molec. Cell. Biol. 6: 511–517, 1986.

[171855]

[171856]

[171857]17471.

[171858]Muyan, M.; Furuhashi, M.; Sugahara, T.; Boime, I.: The carboxy-terminal

[171859]region of the beta-subunits of luteinizing hormone and chorionic gonadotropin

[171860]differentially influence secretion and assembly of the heterodimers. Molec.

[171861]Endocr. 10: 1678–1687, 1996.

[171862]

[171863]

[171864]17472.

[171865]Ye, K.; Aghdasi, B.; Luo, H. R.; Moriarity, J. L.; Wu, F. Y.; Hong,

[171866]J. J.; Hurt, K. J.; Bae, S. S.; Suh, P.-G.; Snyder, S. H.: Phospholipase

[171867]C-gamma-1 is a physiological guanine nucleotide exchange factor for

[171868]the nuclear GTPase PIKE. Nature 415: 541–544, 2002.

[171869]

[171870]

[171871]17473.

[171872]Tsuji, M.; Kawano, S.; Tsuji, S.; Sawaoka, H.; Hori, M.;

DuBois,

[171873]R. N.: Cyclooxygenase regulates angiogenesis induced by
colon cancer

[171874]cells. Cell 93: 705–716, 1998.

[171875]

[171876]

[171877]17474.

[171878]Vane, J. R.; Mitchell, J. A.; Appleton, I.; Tomlinson, A.;

Bishop–Bailey,

[171879]D.; Croxtall, J.; Willoughby, D. A.: Inducible isoforms of
cyclooxygenase

[171880]and nitric–oxide synthase in inflammation. Proc. Nat.

Acad. Sci. 91:

[171881]2046–2050, 1994.

[171882]

[171883]

[171884]17475.

[171885]Yokoyama, C.; Tanabe, T.: Cloning of human gene encod–

ing prostaglandin

[171886]endoperoxide synthase and primary structure of the enzyme. Biochem.

[171887]Biophys. Res. Commun. 165: 888–894, 1989.

[171888]

[171889]

[171890]17476.

[171891]Adam, M.; Boie, Y.; Rushmore, T. H.; Muller, G.; Bastien, L.; McKee,

[171892]K. T.; Metters, K. M.; Abramovitz, M.: Cloning and expression of

[171893]three isoforms of the human EP(3) prostanoid receptor. FEBS Lett. 338:

[171894]170–174, 1994.

[171895]

[171896]

[171897]17477.

[171898]Kotani, M.; Tanaka, I.; Ogawa, Y.; Suganami, T.; Matsumoto, T.;

[171899]Muro, S.; Yamamoto, Y.; Sugawara, A.; Yoshimasa, Y.;

Sagawa, N.; Narumiya,

[171900]S.; Nakao, K.: Multiple signal transduction pathways
through two

[171901]prostaglandin E receptor EP3 subtype isoforms expressed
in human uterus. J.

[171902]Clin. Endocr. Metab. 85: 4315–4322, 2000.

[171903]

[171904]

[171905]17478.

[171906]Kotani, M.; Tanaka, I.; Ogawa, Y.; Usui, T.; Mori, K.;
Ichikawa,

[171907]A.; Narumiya, S.; Yoshimi, T.; Nakao, K.: Molecular cloning
and expression

[171908]of multiple isoforms of human prostaglandin E receptor
EP(3) subtype

[171909]generated by alternative messenger RNA splicing: multiple
second messenger

[171910]systems and tissue-specific distributions. Molec. Pharm.
48: 869–879,

[171911]1995.

[171912]

[171913]

[171914]17479.

[171915]Kotani, M.; Tanaka, I.; Ogawa, Y.; Usui, T.; Tamura, N.;
Mori,

[171916]K.; Narumiya, S.; Yoshimi, T.; Nakao, K.: Structural organi-
zation

[171917]of the human prostaglandin EP(3) receptor subtype gene
(PTGER3). Genomics 40:

[171918]425–434, 1997.

[171919]

[171920]

[171921]17480.

[171922]Schmid, A.; Thierauch, K.–H.; Schleuning, W.–D.; Dinter,
H.: Splice

[171923]variants of the human EP(3) receptor for prostaglandin
E(2). Europ.

[171924]J. Biochem. 228: 23–30, 1995.

[171925]

[171926]

[171927]17481.

[171928]Ushikubi, F.; Segi, E.; Sugimoto, Y.; Murata, T.; Matsuoka, T.;

[171929]Kobayashi, T.; Hizaki, H.; Tuboi, K.; Katsuyama, M.; Ichikawa, A.;

[171930]Tanaka, T.; Yoshida, N.; Narumiya, S.: Impaired febrile response

[171931]in mice lacking the prostaglandin E receptor subtype EP(3). Nature 395:

[171932]281–284, 1998.

[171933]

[171934]

[171935]17482.

[171936]Wurz, H.; Geiger, W.; Kunzig, H. J.; Jabs–Lehmann, A.; Bohn, H.;

[171937]Luben, G.: Radioimmunoassay of SP1 (pregnancy–specific beta–1–glycoprotein)

[171938]in maternal blood and in amniotic fluid in normal and pathologic pregnancies. J.

[171939]Perinat. Med. 2: 67–78, 1981.

[171940]

[171941]

[171942]17483.

[171943]Kaplan, R.; Morse, B.; Huebner, K.; Croce, C.; Howk, R.;
Ravera,

[171944]M.; Ricca, G.; Jaye, M.; Schlessinger, J.: Cloning of three
human

[171945]tyrosine phosphatases reveals a multigene family of re-
ceptor-linked

[171946]protein-tyrosine-phosphatases expressed in brain. Proc.
Nat. Acad.

[171947]Sci. 87: 7000-7004, 1990.

[171948]

[171949]

[171950]17484.

[171951]Banville, D.; Stocco, R.; Shen, S.-H.: Human protein tyro-
sine

[171952]phosphatase 1C (PTPN6) gene structure: alternate pro-
moter usage and

[171953]exon skipping generate multiple transcripts. Genomics
27: 165-173,

[171954]1995.

[171955]

[171956]

[171957]17485.

[171958]Beghini, A.; Ripamonti, C. B.; Peterlongo, P.; Roversi, G.;
Cairolì,

[171959]R.; Morra, E.; Larizza, L.: RNA hyperediting and alternative
splicing

[171960]of hematopoietic cell phosphatase (PTPN6) gene in acute
myeloid leukemia. Hum.

[171961]Molec. Genet. 9: 2297–2304, 2000.

[171962]

[171963]

[171964]17486.

[171965]Matthews, R. J.; Bowne, D. B.; Flores, E.; Thomas, M. L.:
Characterization

[171966]of hematopoietic intracellular protein tyrosine phos-
phatases: description

[171967]of a phosphatase containing an SH2 domain and another
enriched in

[171968]proline-, glutamic acid-, serine-, and threonine-rich se-
quences. Molec.

[171969]Cell. Biol. 12: 2396–2405, 1992.

[171970]

[171971]

[171972]17487.

[171973]Plutzky, J.; Neel, B. G.; Rosenberg, R. D.; Eddy, R. L.; Byers,

[171974]M. G.; Jani-Sait, S.; Shows, T. B.: Chromosomal localization
of an

[171975]SH2-containing tyrosine phosphatase (PTPN6). Genomics
13: 869–872,

[171976]1992.

[171977]

[171978]

[171979]17488.

[171980]Shen, S.-H.; Bastien, L.; Posner, B. I.; Chretien, P.: A pro-
tein-tyrosine

[171981]phosphatase with sequence similarity to the SH2 domain
of the protein-tyrosine

[171982]kinases. Nature 352: 736–739, 1991.

[171983]

[171984]

[171985]17489.

[171986]Shultz, L. D.; Schweitzer, P. A.; Rajan, T. V.; Yi, T.; Ihle, J.

[171987]N.; Matthews, R. J.; Thomas, M. L.; Beier, D. R.: Mutations
at the

[171988]murine motheaten locus are within the hematopoietic cell
protein-tyrosine

[171989]phosphatase (Hcph) gene. Cell 73: 1445-1454, 1993.

[171990]

[171991]

[171992]17490.

[171993]Tsui, H. W.; Siminovitch, K. A.; de Souza, L.; Tsui, F. W. L.:

[171994]Motheaten and viable motheaten mice have mutations in
the haematopoietic

[171995]cell phosphatase gene. Nature Genet. 4: 124-129, 1993.

[171996]

[171997]

[171998]17491.

[171999]Yi, T.; Cleveland, J. L.; Ihle, J. N.: Identification of novel

[172000]protein tyrosine phosphatases of hematopoietic cells by

PCR amplification. Blood 78:

[172001]2222–2228, 1991.

[172002]

[172003]

[172004]17492.

[172005]Yi, T.; Cleveland, J. L.; Ihle, J. N.: Protein tyrosine phosphatase

[172006]containing SH2 domains: characterization, preferential expression

[172007]in hematopoietic cells, and localization to human chromosome 12p12–p13. Molec.

[172008]Cell. Biol. 12: 836–846, 1992.

[172009]

[172010]

[172011]17493.

[172012]Yi, T.; Gilbert, D. J.; Jenkins, N. A.; Copeland, N. G.; Ihle,

[172013]J. N.: Assignment of a novel protein tyrosine phosphatase gene (Hcph)

[172014]to mouse chromosome 6. Genomics 14: 793–795, 1992.

[172015]

[172016]

[172017]17494.

[172018]Jirik, F. R.; Anderson, L. L.; Duncan, A. M. V.: The human
protein-tyrosine

[172019]phosphatase PTP-alpha/LRP gene (PTPA) is assigned to
chromosome 20p13. Cytogenet.

[172020]Cell Genet. 60: 117-118, 1992.

[172021]

[172022]

[172023]17495.

[172024]Jirik, F. R.; Janzen, N. M.; Melhado, I. G.; Harder, K. W.:
Cloning

[172025]and chromosomal assignment of a widely expressed hu-
man receptor-like

[172026]protein-tyrosine phosphatase. FEBS Lett. 273: 239-242,
1990.

[172027]

[172028]

[172029]17496.

[172030]Matthews, R. J.; Cahir, E. D.; Thomas, M. L.: Identification
of

[172031]an additional member of the protein-tyro-
sine-phosphatase family: evidence

[172032]for alternative splicing in the tyrosine phosphatase do-
main. Proc.

[172033]Nat. Acad. Sci. 87: 4444-4448, 1990.

[172034]

[172035]

[172036]17497.

[172037]Rao, V. V. N. G.; Loffler, C.; Sap, J.; Schlessinger, J.; Hans-
mann,

[172038]I.: The gene for receptor-linked protein-tyro-
sine-phosphatase (PTPA)

[172039]is assigned to human chromosome 20p12-pter by in situ
hybridization

[172040](ISH and FISH). Genomics 13: 906-907, 1992.

[172041]

[172042]

[172043]17498.

[172044]Schnittger, S.; Rao, V. V. N. G.; Deutsch, U.; Gruss, P.;
Balling,

[172045]R.; Hansmann, I.: PAX1, a member of the paired box-
containing class

[172046]of developmental control genes, is mapped to human
chromosome 20p11.2

[172047]by in situ hybridization (ISH and FISH). Genomics 14:
740–744, 1992.

[172048]

[172049]

[172050]17499.

[172051]Ahmad, F.; Azevedo, J. L., Jr.; Cortright, R.; Dohm, G. L.;
Goldstein,

[172052]B. J.: Alternations in skeletal muscle protein-tyrosine
phosphatase

[172053]activity and expression in insulin-resistant human obesity
and diabetes. J.

[172054]Clin. Invest. 100: 449–458, 1997.

[172055]

[172056]

[172057]17500.

[172058]Brown-Shimer, S.; Johnson, K. A.; Lawrence, J. B.; Johnson, C.;

[172059]Bruskin, A.; Green, N. R.; Hill, D. E.: Molecular cloning and chromosome

[172060]mapping of the human gene encoding protein phosphotyrosyl phosphatase

[172061]1B. Proc. Nat. Acad. Sci. 87: 5148–5152, 1990.

[172062]

[172063]

[172064]17501.

[172065]Charbonneau, H.; Tonks, N. K.; Kumar, S.; Diltz, C. D.; Harrylock,

[172066]M.; Cool, D. E.; Krebs, E. G.; Fischer, E. H.; Walsh, K. A.: Human

[172067]placenta protein-tyrosine-phosphatase: amino acid sequence and relationship

[172068]to a family of receptor-like proteins. Proc. Nat. Acad. Sci. 86:

[172069]5252–5256, 1989.

[172070]

[172071]

[172072]17502.

[172073]Chernoff, J.; Schievella, A. R.; Jost, C. A.; Erikson, R. L.;
Neel,

[172074]B. G.: Cloning of a cDNA for a major human protein-tyro-
sine-phosphatase. Proc.

[172075]Nat. Acad. Sci. 87: 2735–2739, 1990.

[172076]

[172077]

[172078]17503.

[172079]Matilla, A.; Gorbea, C.; Einum, D. D.; Townsend, J.; Micha-
lik,

[172080]A.; van Broeckhoven, C.; Jensen, C. C.; Murphy, K. J.;
Ptacek, L.

[172081]J.; Fu, Y.-H.: Association of ataxin-7 with the proteasome
subunit

[172082]S4 of the 19S regulatory complex. Hum. Molec. Genet. 10:
2821–2831,

[172083]2001.

[172084]

[172085]

[172086]17504.

[172087]Cazzaniga, G.; Tosi, S.; Aloisi, A.; Giudici, G.; Daniotti, M.;

[172088]Pioltelli, P.; Kearney, L.; Biondi, A.: The tyrosine kinase
Abl-related

[172089]gene ARG is fused to ETV6 in an AML-M4Eo patient with a
t(1;12)(q25;p13):

[172090]molecular cloning of both reciprocal transcripts. Blood 94:
4370-4373,

[172091]1999.

[172092]

[172093]

[172094]17505.

[172095]Iijima, Y.; Ito, T.; Oikawa, T.; Eguchi, M.; Eguchi-Ishimae,
M.;

[172096]Kamada, N.; Kishi, K.; Asano, S.; Sakaki, Y.; Sato, Y.: A new
ETV6/TEL

[172097]partner gene, ARG (ABL-related gene or ABL2), identified
in an AML-M3

[172098]cell line with a t(1;12)(q25;p13) translocation. Blood 95:
2126-2131,

[172099]2000.

[172100]

[172101]

[172102]17506.

[172103]Joazeiro, C. A. P.; Wing, S. S.; Huang, H.; Levenson, J. D.;
Hunter,

[172104]T.; Liu, Y.-C.: The tyrosine kinase negative regulator c-Cbl
as a

[172105]RING-type, E2-dependent ubiquitin-protein ligase. Sci-
ence 286: 309-312,

[172106]1999.

[172107]

[172108]

[172109]17507.

[172110]Langdon, W. Y.; Hartley, J. W.; Klinken, S. P.; Ruscetti, S.
K.;

[172111]Morse, H. C., III: v-cbl, an oncogene from a dual-
recombinant murine

[172112]retrovirus that induces early B-lineage lymphomas. Proc.
Nat. Acad.

[172113]Sci. 86: 1168-1172, 1989.

[172114]

[172115]

[172116]17508.

[172117]Mastick, C. C.; Brady, M. J.; Saltiel, A. R.: Insulin stimulates

[172118]the tyrosine phosphorylation of caveolin. J. Cell Biol. 129:

1523–1531,

[172119]1995.

[172120]

[172121]

[172122]17509.

[172123]Petrelli, A.; Gilestro, G. F.; Lanzardo, S.; Comoglio, P. M.;

Migone,

[172124]N.; Giordano, S.: The endophilin–CIN85–Cbl complex me-

diates ligand-dependent

[172125]downregulation of c-Met. Nature 416: 187–190, 2002.

[172126]

[172127]

[172128]17510.

[172129]Savage, P. D.; Shapiro, M.; Langdon, W. Y.; Geurts van

Kessel,

[172130]A. D.; Seuanez, H. N.; Akao, Y.; Croce, C.; Morse, H. C., III;

Kersey,

[172131]J. H.: Relationship of the human protooncogene CBL2 on

11q23 to the

[172132]t(4;11), t(11;22), and t(11;14) breakpoints. Cytogenet. Cell

Genet. 56:

[172133]112–115, 1991.

[172134]

[172135]

[172136]17511.

[172137]Soubeyran, P.; Kowanetz, K.; Szymkiewicz, I.; Langdon, W.

Y.; Dikic,

[172138]I.: Cbl–CIN85–endophilin complex mediates ligand–in–

duced downregulation

[172139]of EGF receptors. Nature 416: 183–187, 2002.

[172140]

[172141]

[172142]17512.

[172143]Thien, C. B. F.; Walker, F.; Langdon, W. Y.: RING finger

mutations

[172144]that abolish c-Cbl-directed polyubiquitination and down-regulation

[172145]of the EGF receptor are insufficient for cell transformation.
Molec.

[172146]Cell 7: 355-365, 2001.

[172147]

[172148]

[172149]17513.

[172150]Wei, S.; Rocchi, M.; Archidiacono, N.; Sacchi, N.; Romeo, G.;

[172151]Gatti, R. A.: Physical mapping of the human chromosome 11q23 region

[172152]containing the ataxia-telangiectasia locus. Cancer Genet. Cytogenet. 46:

[172153]1-8, 1990.

[172154]

[172155]

[172156]17514.

[172157]Yoon, C. H.; Lee, J.; Jongeward, G. D.; Sternberg, P. W.:

Similarity

[172158]of sli-1, a regulator of vulval development in *C. elegans*,
to the

[172159]mammalian proto-oncogene c-cbl. *Science* 269:
1102-1105, 1995.

[172160]

[172161]

[172162]17515.

[172163]Cannizzaro, L. A.; Madaule, P.; Hecht, F.; Axel, R.; Croce,
C.

[172164]M.; Huebner, K.: Chromosome localization of human ARH
genes, a ras-related

[172165]gene family. *Genomics* 6: 197-203, 1990.

[172166]

[172167]

[172168]17516.

[172169]Chardin, P.; Madaule, P.; Tavitian, A.: Coding sequence of
human

[172170]rho cDNAs clone 6 and clone 9. *Nucleic Acid Res.* 16:
2717 only,

[172171]1988.

[172172]

[172173]

[172174]17517.

[172175]Madaule, P.; Axel, R.: A novel ras-related gene family. Cell
41:

[172176]31–40, 1985.

[172177]

[172178]

[172179]17518.

[172180]Maekawa, M.; Ishizaki, T.; Boku, S.; Watanabe, N.; Fujita,
A.;

[172181]Iwamatsu, A.; Obinata, T.; Ohashi, K.; Mizuno, K.; Naru-
miya, S.:

[172182]Signaling from Rho to the actin cytoskeleton through pro-
tein kinases

[172183]ROCK and LIM-kinase. Science 285: 895–898, 1999.

[172184]

[172185]

[172186]17519.

[172187]Ridley, A. J.; Hall, A.: The small GTP-binding protein rho regulates

[172188]the assembly of focal adhesions and actin stress fibers in response

[172189]to growth factors. Cell 70: 389–399, 1992.

[172190]

[172191]

[172192]17520.

[172193]Fagan, K. P.; Oliveira, L.; Pittler, S. J.: Sequence of rho small

[172194]GTP-binding protein cDNAs from human retina and identification of

[172195]novel 5-prime end cloning artifacts. Exp. Eye Res. 59: 235–237,

[172196]1994.

[172197]

[172198]

[172199]17521.

[172200]Morris, S. W.; Valentine, M. B.; Kirstein, M. N.; Huebner, K.:

[172201]Reassignment of the human ARH9 RAS-related gene to

chromosome 1p13–p21. Genomics 15:

[172202]677–679, 1993.

[172203]

[172204]

[172205]17522.

[172206]Maesaki, R.; Ihara, K.; Shimizu, T.; Kuroda, S.; Kaibuchi, K.;

[172207]Hakoshima, T.: The structural basis of Rho effector recognition revealed

[172208]by the crystal structure of human RhoA complexed with the effector

[172209]domain of PKN/PRK1. Molec. Cell 4: 793–803, 1999.

[172210]

[172211]

[172212]17523.

[172213]Nakamura, M.; Nagano, T.; Chikama, T.; Nishida, T.: Role of the

[172214]small GTP-binding protein Rho in epithelial cell migration in the

[172215]rabbit cornea. Invest. Ophthalm. Vis. Sci. 42: 941–947, 2001.

[172216]

[172217]

[172218]17524.

[172219]Rao, P. V.; Deng, P.-F.; Kumar, J.; Epstein, D. L.: Modulation
tion

[172220]of aqueous humor outflow facility by the Rho kinase-specific inhibitor

[172221]Y-27632. Invest. Ophthal. Vis. Sci. 42: 1029-1037, 2001.

[172222]

[172223]

[172224]17525.

[172225]Gharib, B.; Fox, M. F.; Bartoli, C.; Giorgi, D.; Sansonetti, A.;

[172226]Swallow, D. M.; Dagorn, J. C.; Berge-Lefranc, J. L.: Human
regeneration

[172227]protein/lithostathine genes map to chromosome 2p12.

Ann. Hum. Genet. 57:

[172228]9-16, 1993.

[172229]

[172230]

[172231]17526.

[172232]Miyashita, H.; Nakagawara, K.; Mori, M.; Narushima, Y.;
Noguchi,

[172233]N.; Moriizumi, S.; Takasawa, S.; Yonekura, H.; Takeuchi,
T.; Okamoto,

[172234]H.: Human REG family genes are tandemly ordered in a
95-kilobase

[172235]region of chromosome 2p12. FEBS Lett. 377: 429–433,
1995.

[172236]

[172237]

[172238]17527.

[172239]Moriizumi, S.; Watanabe, T.; Unno, M.; Nakagawara, K.;
Suzuki,

[172240]Y.; Miyashita, H.; Yonekura, H.; Okamoto, H.: Isolation,
structural

[172241]determination and expression of a novel reg gene, human
regI-beta. Biochim.

[172242]Biophys. Acta 1217: 199–202, 1994.

[172243]

[172244]

[172245]17528.

[172246]Chen, K.; Yang, W.; Grimsby, J.; Shih, J. C.: The human
5-HT₂

[172247]receptor is encoded by a multiple intron-exon gene. Brain
Res. Molec.

[172248]Brain Res. 14: 20-26, 1992.

[172249]

[172250]

[172251]17529.

[172252]Holmes, C.; Arranz, M. J.; Powell, J. F.; Collier, D. A.; Love-
stone,

[172253]S.: 5-HT-2A and 5-HT-2C receptor polymorphisms and
psychopathology

[172254]in late onset Alzheimer's disease. Hum. Molec. Genet. 7:
1507-1509,

[172255]1998.

[172256]

[172257]

[172258]17530.

[172259]Hsieh, C.-L.; Bowcock, A. M.; Farrer, L. A.; Hebert, J. M.;
Huang,

[172260]K. N.; Cavalli-Sforza, L. L.; Julius, D.; Francke, U.: The
serotonin

[172261]receptor subtype 2 locus HTR2 is on human chromosome
13 near genes

[172262]for esterase D and retinoblastoma-1 and on mouse chro-
mosome 14. Somat.

[172263]Cell Molec. Genet. 16: 567-574, 1990.

[172264]

[172265]

[172266]17531.

[172267]Kato, M. V.; Shimizu, T.; Nagayoshi, M.; Kaneko, A.;
Sasaki, M.

[172268]S.; Ikawa, Y.: Genomic imprinting of the human serotonin-
receptor

[172269](HTR2) gene involved in development of retinoblastoma.
Am. J. Hum.

[172270]Genet. 59: 1084-1090, 1996.

[172271]

[172272]

[172273]17532.

[172274]Liu, J.; Chen, Y.; Kozak, C. A.; Yu, L.: The 5-HT(2) serotonin

[172275]receptor gene Htr-2 is tightly linked to Es-10 on mouse chromosome

[172276]14. Genomics 11: 231-234, 1991.

[172277]

[172278]

[172279]17533.

[172280]Peroutka, S. J.: 5-Hydroxytryptamine receptor subtypes. Ann.

[172281]Rev. Neurosci. 11: 45-60, 1988.

[172282]

[172283]

[172284]17534.

[172285]Sparkes, R. S.; Lan, N.; Klisak, I.; Mohandas, T.; Diep, A.; Kojis,

[172286]T.; Heinzmann, C.; Shih, J. C.: Assignment of a serotonin 5HT-2 receptor

[172287]gene (HTR2) to human chromosome 13q14-q21 and

mouse chromosome 14. Genomics 9:

[172288]461–465, 1991.

[172289]

[172290]

[172291]17535.

[172292]Williams, J.; McGuffin, P.; Nothen, M.; Owen, M. J.; EMASS
Collaborative

[172293]Group: Meta-analysis of association between the 5-HT
(2A) receptor

[172294]T102C polymorphism and schizophrenia. (Letter) Lancet
349: 1221

[172295]only, 1997.

[172296]

[172297]

[172298]17536.

[172299]Williams, J.; Spurlock, G.; McGuffin, P.; Mallet, J.; Nothen,
M.

[172300]M.; Gill, M.; Aschauer, H.; Nylander, P. O.; Macciardi, F.;
Owen,

[172301]M. J.: Association between schizophrenia and T102C poly-

morphism of

[172302]the 5-hydroxytryptophan type 2a-receptor gene. European Multicentre

[172303]Association Study of Schizophrenia (EMASS) Group. Lancet 347: 1294-1296,

[172304]1996.

[172305]

[172306]

[172307]17537.

[172308]Philipson, L. H.; Hice, R. E.; Schaefer, K.; LaMendola, J.; Bell,

[172309]G. I.; Neldon, D. J.; Steiner, D. F.: Sequence and functional expression

[172310]in *Xenopus* oocytes of a human insulinoma and islet potassium channel. Proc.

[172311]Nat. Acad. Sci. 88: 53-57, 1991.

[172312]

[172313]

[172314]17538.

[172315]Ghanshani, S.; Pak, M.; McPherson, J. D.; Strong, M.; Deth-

lefs,

[172316]B.; Wasmuth, J. J.; Salkoff, L.; Gutman, G. A.; Chandy, K.

G.: Genomic

[172317]organization, nucleotide sequence, and cellular distribution of a

[172318]Shaw-related potassium channel gene, Kv3.3, and mapping of Kv3.3 and

[172319]Kv3.4 to human chromosomes 19 and 1. Genomics 12: 190–196, 1992.

[172320]

[172321]

[172322]17539.

[172323]McPherson, J. D.; Wasmuth, J. J.; Chandy, K. G.; Swanson, R.; Dethlefs,

[172324]B.; Chandy, G.; Wymore, R.; Ghanshani, S.: Chromosomal localization

[172325]of 7 potassium channel genes. (Abstract) Cytogenet. Cell Genet. 58:

[172326]1979 only, 1991.

[172327]

[172328]

[172329]17540.

[172330]Rudy, B.; Sen, K.; Vega-Saenz de Miera, E.; Lau, D.; Ried, T.;

[172331]Ward, D. C.: Cloning of a human cDNA expressing a high voltage-activating,

[172332]TEA-sensitive, type-A K⁺ channel which maps to chromosome 1 band p21. J.

[172333]Neurosci. Res. 29: 401-412, 1991.

[172334]

[172335]

[172336]17541.

[172337]Gessler, M.; Grupe, A.; Grzeschik, K.-H.; Pongs, O.: The potassium

[172338]channel gene HK1 maps to human chromosome 11p14.1, close to the FSHB

[172339]gene. Hum. Genet. 90: 319-321, 1992.

[172340]

[172341]

[172342]17542.

[172343]Grandy, D.; Mathew, M. K.; Ramaswami, M.; Tanouye, M.;

Sheffield,

[172344]V.; Jones, C. A.; Al-Dhalimi, M.; Zhang, Y.; Saez, C.; Litt, M.:

[172345]A human voltage-gated potassium channel gene, HuKII, maps to chromosome

[172346]11p14-p13. (Abstract) Am. J. Hum. Genet. 51 (suppl.):

A396 only,

[172347]1992.

[172348]

[172349]

[172350]17543.

[172351]Philipson, L. H.; Eddy, R. L.; Shows, T. B.; Bell, G. I.: Assignment
segment

[172352]of human potassium channel gene KCNA4 (Kv1.4, PCN2) to chromosome

[172353]11q13.4-q14.1. Genomics 15: 463-464, 1993.

[172354]

[172355]

[172356]17544.

[172357]Philipson, L. H.; Schaefer, K.; LaMendola, J.; Bell, G. I.;

Steiner,

[172358]D. F.: Sequence of a human fetal skeletal muscle potassium channel

[172359]cDNA related to RCK4. Nucleic Acids Res. 18: 7160 only, 1990.

[172360]

[172361]

[172362]17545.

[172363]Stuehmer, W.; Ruppersberg, J. P.; Schroter, K. H.; Sakmann, B.;

[172364]Stocker, M.; Giese, K. P.; Perschke, A.; Baumann, A.; Pongs, O.:

[172365]Molecular basis of functional diversity of voltage-gated potassium

[172366]channels in mammalian brain. EMBO J. 8: 3235–3244, 1989.

[172367]

[172368]

[172369]17546.

[172370]Tamkun, M. M.; Knoth, K. M.; Walbridge, J. A.; Kroemer, H.; Roden,

[172371]D. M.; Glover, D. M.: Molecular cloning and characterization of two

[172372]voltage-gated K⁺ channel cDNAs from human ventricle.

FASEB J. 5:

[172373]331–337, 1991.

[172374]

[172375]

[172376]17547.

[172377]Phromchotikul, T.; Browne, D. L.; Curran, M. E.; Keating, M. T.;

[172378]Litt, M.: Dinucleotide repeat polymorphism at the KCNA5 locus. Hum.

[172379]Molec. Genet. 2: 1512 only, 1993.

[172380]

[172381]

[172382]17548.

[172383]Bardien-Kruger, S.; Wulff, H.; Arieff, Z.; Brink, P.; Chandy, K.

[172384]G.; Corfield, V.: Characterisation of the human voltage-gated potassium

[172385]channel gene, KCNA7, a candidate gene for inherited cardiac disorders,

[172386]and its exclusion as cause of progressive familial heart block I (PFHBI). Europ.

[172387]J. Hum. Genet. 10: 36–43, 2002.

[172388]

[172389]

[172390]17549.

[172391]Day, J. R.; Albers, J. J.; Lofton–Day, C. E.; Gilbert, T. L.; Ching,

[172392]A. F. T.; Grant, F. J.; O'Hara, P. J.; Marcovina, S. M.; Adolphson,

[172393]J. L.: Complete cDNA encoding human phospholipid transfer protein

[172394]from human endothelial cells. J. Biol. Chem. 269: 9388–9391, 1994.

[172395]

[172396]

[172397]17550.

[172398]Jiang, X.; Bruce, C.; Mar, J.; Lin, M.; Ji, Y.; Francone, O. L.;

[172399]Tall, A. R.: Targeted mutation of plasma phospholipid transfer protein

[172400]gene markedly reduces high-density lipoprotein levels. J. Clin. Invest. 103:

[172401]907-914, 1999.

[172402]

[172403]

[172404]17551.

[172405]LeBoeuf, R. C.; Caldwell, M.; Tu, A.-Y.; Albers, J. J.: Phospholipid

[172406]transfer protein maps to distal mouse chromosome 2. Genomics 34:

[172407]259-260, 1996.

[172408]

[172409]

[172410]17552.

[172411]Tu, A.-Y.; Wolfbauer, G.; Albers, J. J.: Functional characterization

[172412]of the promoter region of the human phospholipid transfer protein

[172413]gene. Biochem. Biophys. Res. Commun. 217: 705–711,
1995.

[172414]

[172415]

[172416]17553.

[172417]Whitmore, T. E.; Day, J. R.; Albers, J. J.: Localization of the

[172418]human phospholipid transfer protein gene to chromo-
some 20q12–q13.1. Genomics 28:

[172419]599–600, 1995.

[172420]

[172421]

[172422]17554.

[172423]Comings, D. E.: Evidence for ancient tetraploidy and con-
servation

[172424]of linkage groups in mammalian chromosomes. Nature
238: 455–457,

[172425]1972.

[172426]

[172427]

[172428]17555.

[172429]D'Ancona, G. G.; Chern, C. J.; Benn, P.; Croce, C. M.: Assignment
segment

[172430]of the human gene for enolase 1 to region pter-p36 of
chromosome 1. Cytogenet.

[172431]Cell Genet. 18: 327-332, 1977.

[172432]

[172433]

[172434]17556.

[172435]D'Ancona, G. G.; Croce, C. M.: Assignment of the gene for
enolase

[172436]to mouse chromosome 4 using somatic cell hybrids. Cyto-
genet. Cell

[172437]Genet. 19: 1-6, 1977.

[172438]

[172439]

[172440]17557.

[172441]Feo, S.; Oliva, D.; Arico, B.; Barba, G.; Cali, L.; Giallongo,

[172442]A.: The human genome contains a single processed pseu-

dogene for alpha

[172443]enolase located on chromosome 1. DNA Sequence 1:
79–83, 1990.

[172444]

[172445]

[172446]17558.

[172447]Giallongo, A.; Feo, S.; Moore, R.; Croce, C. M.; Showe, L.
C.:

[172448]Molecular cloning and nucleotide sequence of a full–
length cDNA for

[172449]human alpha enolase. Proc. Nat. Acad. Sci. 83:
6741–6745, 1986.

[172450]

[172451]

[172452]17559.

[172453]Giblett, E. R.; Chen, S.–H.; Anderson, J. E.; Lewis, M.: A
family

[172454]study suggesting genetic linkage of phosphopyruvate hy–
dratase (enolase)

[172455]to the Rh blood group system. Cytogenet. Cell Genet. 13:
91–92,

[172456]1974.

[172457]

[172458]

[172459]17560.

[172460]Lachant, N. A.; Jennings, M. A.; Tanaka, K. R.: Partial erythrocyte

[172461]enolase deficiency: a hereditary disorder with variable clinical expression.

[172462](Abstract) Blood 68: 55a only, 1986.

[172463]

[172464]

[172465]17561.

[172466]Lachant, N. A.; Tanaka, K. R.: Enolase kinetic properties in

[172467]partial erythrocyte enolase deficiency. (Abstract) Clin. Res. 35:

[172468]426A only, 1987.

[172469]

[172470]

[172471]17562.

[172472]Lalley, P. A.; Francke, U.; Minna, J. D.: Homologous genes
for

[172473]enolase, phosphogluconate dehydrogenase, phosphoglu-
comutase, and adenylate

[172474]kinase are syntenic on mouse chromosome 4 and human
chromosome 1p. Proc.

[172475]Nat. Acad. Sci. 75: 2382–2386, 1978.

[172476]

[172477]

[172478]17563.

[172479]Ribaud, M. R.; Di Leonardo, A.; Rubino, P.; Giallongo, A.;
Feo,

[172480]S.: Assignment of enolase processed pseudogene (ENO1P)
to human chromosome

[172481]1 bands 1q41–q42. Cytogenet. Cell Genet. 74: 201–202,
1996.

[172482]

[172483]

[172484]17564.

[172485]Van Cong, N.; Weil, D.; Rebourcet, R.; Frezal, J.: Localisa-
tion

[172486]des enolases 1 et 2 respectivement sur les chromosomes
1 et 12 par

[172487]l'analyse des hybrids homme-souris. Ann. Genet. 20:
153-157, 1977.

[172488]

[172489]

[172490]17565.

[172491]Wistow, G. J.; Lietman, T.; Williams, L. A.; Stapel, S. O.; de

[172492]Jong, W. W.; Horwitz, J.; Piatigorsky, J.: Tau-crys-
tallin/alpha-enolase:

[172493]one gene encodes both an enzyme and a lens structural
protein. J.

[172494]Cell Biol. 107: 2729-2736, 1988.

[172495]

[172496]

[172497]17566.

[172498]Barton, J. W.; Hart, I. M.; Patterson, D.: Mapping of a locus

[172499]correcting lack of phosphoribosylaminoimidazole car-
boxylase activity

[172500]in Chinese hamster ovary cell Ade(-)D mutants to human

chromosome

[172501]4. Genomics 9: 314–321, 1991.

[172502]

[172503]

[172504]17567.

[172505]Brayton, K. A.; Chen, Z.; Zhou, G.; Nagy, P. L.; Gavalas, A.;
Trent,

[172506]J. M.; Deaven, L. L.; Dixon, J. E.; Zalkin, H.: Two genes for
de

[172507]novo purine nucleotide synthesis on human chromosome
4 are closely

[172508]linked and divergently transcribed. J. Biol. Chem. 269:
5313–5321,

[172509]1994.

[172510]

[172511]

[172512]17568.

[172513]Torban, E.; Pelletier, J.; Goodyer, P.: F329L polymorphism
in

[172514]the human PAX8 gene. Am. J. Med. Genet. 72: 186–187,

1997.

[172515]

[172516]

[172517]17569.

[172518]Gartner, J.; Kearns, W.; Pearson, P.; Valle, D.: Characteri-
zation

[172519]and localization of the human 70-kD peroxisomal mem-
brane protein (PMP70)

[172520]gene. (Abstract) Am. J. Hum. Genet. 51 (suppl.): A168,
1992.

[172521]

[172522]

[172523]17570.

[172524]Gartner, J.; Kearns, W.; Rosenberg, C.; Pearson, P.;
Copeland,

[172525]N. G.; Gilbert, D. J.; Jenkins, N. A.; Valle, D.: Localization of

[172526]the 70-kDa peroxisomal membrane protein to human
1p21-p22 and mouse

[172527]

[172528]17571.

[172529]Genomics 15: 412–414, 1993.

[172530]

[172531]3. Gartner, J.; Moser, H.; Valle, D.: Mutations in the 70K
peroxisomal

[172532]membrane protein gene in Zellweger syndrome. Nature
Genet. 1: 16–23,

[172533]1992.

[172534]

[172535]

[172536]17572.

[172537]Vilain, C.; Rydlewski, C.; Duprez, L.; Heinrichs, C.;
Abramowicz,

[172538]M.; Malvaux, P.; Renneboog, B.; Parma, J.; Costagliola, S.;
Vassart,

[172539]G.: Autosomal dominant transmission of congenital thy-
roid hypoplasia

[172540]due to loss-of-function mutation of PAX8. J. Clin. Endocr.
Metab. 86:

[172541]234–238, 2001.

[172542]

[172543]

[172544]17573.

[172545]Walther, C.; Guenet, J.-L.; Simon, D.; Deutsch, U.; Jostes, B.;

[172546]Goulding, M.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a murine

[172547]multigene family of paired box-containing genes. Genomics 11: 424-434,

[172548]1991.

[172549]

[172550]

[172551]17574.

[172552]Das, P.; Stockton, D. W.; Bauer, C.; Shaffer, L. G.; D'Souza, R.

[172553]N.; Wright, J. T.; Patel, P. I.: Haploinsufficiency of PAX9 is associated

[172554]with autosomal dominant hypodontia. Hum. Genet. 110: 371-376, 2002.

[172555]

[172556]

[172557]17575.

[172558]Nieminen, P.; Arte, S.; Tanner, D.; Paulin, L.; Alaluusua, S.;

[172559]Thesleff, I.; Pirinen, S.: Identification of a nonsense mutation
tion

[172560]in the PAX9 gene in molar oligodontia. Europ. J. Hum.

Genet. 9:

[172561]743–746, 2001.

[172562]

[172563]

[172564]17576.

[172565]Peters, H.; Neubuser, A.; Kratochwil, K.; Balling, R.:

Pax9-deficient

[172566]mice lack pharyngeal pouch derivatives and teeth and exhibit craniofacial

[172567]and limb abnormalities. Genes Dev. 12: 2735–2747,
1998.

[172568]

[172569]

[172570]17577.

[172571]Santagati, F.; Gerber, J.-K.; Blusch, J. H.; Kokubu, C.; Pe-

ters,

[172572]H.; Adamski, J.; Werner, T.; Balling, R.; Imai, K.: Compara-
tive analysis

[172573]of the genomic organization of Pax9 and its conserved
physical association

[172574]with Nkx2-9 in the human, mouse, and pufferfish
genomes. Mammalian

[172575]Genome 12: 232-237, 2001.

[172576]

[172577]

[172578]17578.

[172579]Stockton, D. W.; Das, P.; Goldenberg, M.; D'Souza, R. N.;
Patel,

[172580]P. I.: Mutation of PAX9 is associated with oligodontia.
(Letter) Nature

[172581]Genet. 24: 18-19, 2000.

[172582]

[172583]

[172584]17579.

[172585]Wallin, J.; Mizutani, Y.; Imai, K.; Miyashita, N.; Moriwaki,

K.;

[172586]Taniguchi, M.; Koseki, H.; Balling, R.: A new Pax gene,
Pax-9, maps

[172587]to mouse chromosome 12. Mammalian Genome 4:
354-358, 1993.

[172588]

[172589]

[172590]17580.

[172591]Grueneberg, D. A.; Natesan, S.; Alexandre, C.; Gilman, M.
Z.:

[172592]Human and Drosophila homeodomain proteins that en-
hance the DNA-binding

[172593]activity of serum response factor. Science 257:
1089-1095, 1992.

[172594]

[172595]

[172596]17581.

[172597]Kern, M. J.; Argao, E. A.; Birkenmeier, E. H.; Rowe, L. B.;
Potter,

[172598]S. S.: Genomic organization and chromosome localization
of the murine

[172599]homeobox gene Pmx. Genomics 19: 334–340, 1994.

[172600]

[172601]

[172602]17582.

[172603]Nakamura, T.; Yamazaki, Y.; Hatano, Y.; Miura, I.: NUP98
is fused

[172604]to PMX1 homeobox gene in human acute myelogenous
leukemia with chromosome

[172605]translocation t(1;11)(q23;p15). Blood 94: 741–747, 1999.

[172606]

[172607]

[172608]17583.

[172609]Norris, R. A.; Scott, K. K.; Moore, C. S.; Stetten, G.; Brown,

[172610]C. R.; Jabs, E. W.; Wulfsberg, E. A.; Yu, J.; Kern, M. J.: Hu-
man

[172611]PRRX1 and PRRX2 genes: cloning, expression, genomic lo-
calization,

[172612]and exclusion as disease genes for Nager syndrome.

Mammalian Genome 11:

[172613]1000–1005, 2000.

[172614]

[172615]

[172616]17584.

[172617]Riazuddin, S.; Castelein, C. M.; Ahmed, Z. M.; Lalwani, A. K.;

[172618]Mastroianni, M. A.; Naz, S.; Smith, T. N.; Liburd, N. A.; Friedman,

[172619]T. B.; Griffith, A. J.; Riazuddin, S.; Wilcox, E. R.: Dominant modifier

[172620]DFNM1 suppresses recessive deafness DFNB26. Nature Genet. 26: 431–434,

[172621]2000.

[172622]

[172623]

[172624]17585.

[172625]Hebrok, M.; Kim, S. K.; St Jacques, B.; McMahon, A. P.; Melton,

[172626]D. A.: Regulation of pancreas development by hedgehog signaling. Development 127:

[172627]4905–4913, 2000.

[172628]

[172629]

[172630]17586.

[172631]Reidenberg, M. M.; Levy, M.; Drayer, D. E.; Zylber-Katz, E.;
Robbins,

[172632]W. C.: Acetylator phenotype in idiopathic systemic lupus
erythematosus. Arthritis

[172633]Rheum. 23: 569-573, 1980.

[172634]

[172635]

[172636]17587.

[172637]Stohl, W.; Crow, M. K.; Kunkel, H. G.: Systemic lupus ery-
thematosus

[172638]with deficiency of the T4 epitope on T helper/inducer
cells. New

[172639]Eng. J. Med. 312: 1671-1678, 1985.

[172640]

[172641]

[172642]17588.

[172643]Theofilopoulos, A. N.; Dixon, F. J.: Murine models of systemic

[172644]lupus erythematosus. Adv. Immun. 37: 269–390, 1985.

[172645]

[172646]

[172647]17589.

[172648]Friedl, W.; Uhlhaas, S.; Schulmann, K.; Stolte, M.; Loff, S.;
Back,

[172649]W.; Mangold, E.; Stern, M.; Knaebel, H. P.; Sutter, C.; We-
ber, R.

[172650]G.; Pistorius, S.; Burger, B.; Propping, P.: Juvenile polyposis:

[172651]massive gastric polyposis is more common in MADH4 mutation carriers

[172652]than in BMPR1A mutation carriers. Hum. Genet. 111:
108–111, 2002.

[172653]

[172654]

[172655]17590.

[172656]Houlston, R.; Bevan, S.; Williams, A.; Young, J.; Dunlop, M.;
Rozen,

[172657]P.; Eng, C.; Markie, D.; Woodford-Richens, K.; Rodriguez-Bigas, M.

[172658]A.; Leggett, B.; Neale, K.; Phillips, R.; Sheridan, E.; Hodgson, S.;

[172659]Iwama, T.; Eccles, D.; Bodmer, W.; Tomlinson, I.: Mutations in DPC4

[172660](SMAD4) cause juvenile polyposis syndrome, but only account for a

[172661]minority of cases. Hum. Molec. Genet. 7: 1907-1912, 1998.

[172662]

[172663]

[172664]17591.

[172665]Howe, J. R.; Ringold, J. C.; Summers, R. W.; Mitros, F. A.; Nishimura,

[172666]D. Y.; Stone, E. M.: A gene for familial juvenile polyposis maps

[172667]to chromosome 18q21.1. Am. J. Hum. Genet. 62: 1129-1136, 1998.

[172668]

[172669]

[172670]17592.

[172671]Howe, J. R.; Roth, S.; Ringold, J. C.; Summers, R. W.; Jarvinen,

[172672]H. J.; Sistonen, P.; Tomlinson, I. P. M.; Houlston, R. S.; Bevan,

[172673]S.; Mitros, F. A.; Stone, E. M.; Aaltonen, L. A.: Mutations in the

[172674]SMAD4/DPC4 gene in juvenile polyposis. Science 280: 1086–1088, 1998.

[172675]

[172676]

[172677]17593.

[172678]Leslie, K. K.; Watanabe, S.; Lei, K.-J.; Chou, D. Y.; Plouzek,

[172679]C. A.; Deng, H.-C.; Torres, J.; Chou, J. Y.: Linkage of two human

[172680]pregnancy-specific beta 1-glycoprotein genes: one is associated with

[172681]hydatidiform mole. Proc. Nat. Acad. Sci. 87: 5822–5826, 1990.

[172682]

[172683]

[172684]17594.

[172685]Tatarinov, Y. S.; Falaleeva, D. M.; Kalashnikov, V. V.: Human
man

[172686]pregnancy-specific beta-1-globulin and its relation to
chorioepithelioma. Int.

[172687]J. Cancer 17: 626-632, 1976.

[172688]

[172689]

[172690]17595.

[172691]Berry, C.; Cree, J.; Mann, T.: Aarskog's syndrome. Arch.
Dis.

[172692]Child. 55: 706-710, 1980.

[172693]

[172694]

[172695]17596.

[172696]Escobar, V.; Weaver, D. D.: Aarskog syndrome: new find-
ings and

[172697]genetic analysis. J.A.M.A. 240: 2638-2641, 1978.

[172698]

[172699]

[172700]17597.

[172701]Hanley, W. B.; McKusick, V. A.; Barranco, F. T.: Osteochondritis

[172702]dissecans and associated malformations in brothers: a review of familial

[172703]aspects. J. Bone Joint Surg. 49A: 925–937, 1967.

[172704]

[172705]

[172706]17598.

[172707]Coleman, M. P.; Ambrose, H. J.; Carrel, L.; Nemeth, A. H.; Willard,

[172708]H. F.; Davies, K. E.: A novel gene, DXS8237E, lies within 20 kb upstream

[172709]of UBE1 in Xp11.23 and has a different X inactivation status. Genomics 31:

[172710]135–138, 1996.

[172711]

[172712]

[172713]17599.

[172714]Inoue, A.; Takahashi, K. P.; Kimura, M.; Watanabe, T.;
Morisawa,

[172715]S.: Molecular cloning of a RNA binding protein, S1-1. Nu-
cleic Acids

[172716]Res. 24: 2990-2997, 1996.

[172717]

[172718]

[172719]17600.

[172720]Nagase, T.; Seki, N.; Tanaka, A.; Ishikawa, K.; Nomura, N.:
Prediction

[172721]of the coding sequences of unidentified human genes. IV.
The coding

[172722]sequences of 40 new genes (KIAA0121-KIAA0160) de-
duced by analysis

[172723]of cDNA clones from human cell line KG-1. DNA Res. 2:
167-174, 1995.

[172724]

[172725]

[172726]17601.

[172727]Parrish, J. E.; Ciccodicola, A.; Wehnert, M.; Cox, G. F.;
Chen,

[172728]E.; Nelson, D. L.: A muscle-specific DNase I-like gene in human Xq28. Hum.

[172729]Molec. Genet. 4: 1557–1564, 1995.

[172730]

[172731]

[172732]17602.

[172733]Pergolizzi, R.; Appierto, V.; Bosetti, A.; DeBellis, G. L.; Rovida,

[172734]E.; Biunno, I.: Cloning of a gene encoding a DNase I-like endonuclease

[172735]in the human Xq28 region. Gene 168: 267–270, 1996.

[172736]

[172737]

[172738]17603.

[172739]Klink, A.; Schiebel, K.; Winkelmann, M.; Rao, E.; Hors-themke, B.;

[172740]Ludecke, H.-J.; Claussen, U.; Scherer, G.; Rappold, G.: The human

[172741]protein kinase gene PKX1 on Xp22.3 displays Xp/Yp homology and is

[172742]a site of chromosomal instability. Hum. Molec. Genet. 4:
869–878,
[172743]1995.

[172744]

[172745]

[172746]17604.

[172747]Schiebel, K.; Mertz, A.; Winkelmann, B.; Glaser, B.;
Schempp, W.;

[172748]Rappold, G.: FISH localization of the human Y-homolog of
protein

[172749]kinase PRKX (PRKY) to Yp11.2 and two pseudogenes to
15q26 and Xq12–q13. Cytogenet.

[172750]Cell Genet. 76: 49–52, 1997.

[172751]

[172752]

[172753]17605.

[172754]Brown, C. J.; Carrel, L.; Willard, H. F.: Expression of genes

[172755]from the human active and inactive X chromosomes. Am.
J. Hum. Genet. 60:

[172756]1333–1343, 1997.

[172757]

[172758]

[172759]17606.

[172760]Dong, B.; Horowitz, D. S.; Kobayashi, R.; Krainer, A. R.:

Purification

[172761]and cDNA cloning of HeLa cell p54(nrb), a nuclear protein
with two

[172762]RNA recognition motifs and extensive homology to human
splicing factor

[172763]PSF and Drosophila NONA/BJ6. Nucleic Acids Res. 21:
4085–4092, 1993.

[172764]

[172765]

[172766]17607.

[172767]Nelson, D. L.; Ballabio, A.; Cremers, F.; Monaco, A. P.;

Schlessinger,

[172768]D.: Report of the sixth international workshop on X chro-
mosome mapping

[172769]1995. Cytogenet. Cell Genet. 71: 308–342, 1995.

[172770]

[172771]

[172772]17608.

[172773]Janssens, R.; Boeynaems, J.-M.; Godart, M.; Communi, D.:

Cloning

[172774]of a human heptahelical receptor closely related to the

P2Y(5) receptor. Biochem.

[172775]Biophys. Res. Commun. 236: 106–112, 1997.

[172776]

[172777]

[172778]17609.

[172779]O'Dowd, B. F.; Nguyen, T.; Jung, B. P.; Marchese, A.;

Cheng, R.;

[172780]Heng, H. H. Q.; Kolakowski, L. F., Jr.; Lynch, K. R.; George,

S. R.

[172781]: Cloning and chromosomal mapping of four putative

novel human G-protein-coupled

[172782]receptor genes. Gene 187: 75–81, 1997.

[172783]

[172784]

[172785]17610.

[172786]Knight, S. W.; Heiss, N. S.; Vulliamy, T. J.; Aalfs, C. M.;

McMahon,

[172787]C.; Richmond, P.; Jones, A.; Hennekam, R. C. M.; Poustka,

A.; Mason,

[172788]P. J.; Dokal, I.: Unexplained aplastic anaemia, immunodeficiency,

[172789]and cerebellar hypoplasia (Hoyeraal–Hreidarsson syndrome) due to mutations

[172790]in the dyskeratosis congenita gene, DKC1. Brit. J. Haemat. 107:

[172791]335–339, 1999.

[172792]

[172793]

[172794]17611.

[172795]Knight, S. W.; Heiss, N. S.; Vulliamy, T. J.; Greschner, S.;

Stavrides,

[172796]G.; Pai, G. S.; Lestringant, G.; Varma, N.; Mason, P. J.;

Dokal, I.;

[172797]Poustka, A.: X-linked dyskeratosis congenita is predominantly caused

[172798]by missense mutations in the DKC1 gene. Am. J. Hum.

Genet. 65: 50–58,

[172799]1999.

[172800]

[172801]

[172802]17612.

[172803]Knight, S. W.; Vulliamy, T. J.; Morgan, B.; Devriendt, K.;
Mason,

[172804]P. J.; Dokal, I.: Identification of novel DKC1 mutations in
patients

[172805]with dyskeratosis congenita: implications for pathophysiology and

[172806]diagnosis. Hum. Genet. 108: 299–303, 2001.

[172807]

[172808]

[172809]17613.

[172810]Luzzatto, L.; Karadimitris, A.: Dyskeratosis and ribosomal
rebellion. Nature

[172811]Genet. 19: 6–7, 1998.

[172812]

[172813]

[172814]17614.

[172815]McGrath, J. A.: Dyskeratosis congenita: new clinical and
molecular

[172816]insights into ribosome function. Lancet 353: 1204–1205,
1999.

[172817]

[172818]

[172819]17615.

[172820]Mitchell, J. R.; Wood, E.; Collins, K.: A telomerase compo-
nent

[172821]is defective in the human disease dyskeratosis congenita.
Nature 402:

[172822]551–555, 1999.

[172823]

[172824]

[172825]17616.

[172826]Salowsky, R.; Heiss, N. S.; Benner, A.; Wittig, R.; Poustka,
A.

[172827]: Basal transcription activity of the dyskeratosis congenita
gene

[172828]is mediated by Sp1 and Sp3 and a patient mutation in a
Sp1 binding

[172829]site is associated with decreased promoter activity. Gene
293: 9–19,

[172830]2002.

[172831]

[172832]

[172833]17617.

[172834]Vulliamy, T. J.; Knight, S. W.; Heiss, N. S.; Smith, O. P.;
Poustka,

[172835]A.; Dokal, I.; Mason, P. J.: Dyskeratosis congenita caused
by a 3–prime

[172836]deletion: germline and somatic mosaicism in a female
carrier. Blood 94:

[172837]1254–1260, 1999.

[172838]

[172839]

[172840]17618.

[172841]Yaghmai, R.; Kimyai–Asadi, A.; Rostamiani, K.; Heiss, N. S.;
Poustka,

[172842]A.; Eyaid, W.; Bodurtha, J.; Nousari, H. C.; Hamosh, A.;
Metzenberg,

[172843]A.: Overlap of dyskeratosis congenita with the Hoyeraal–
Hreidarsson

[172844]syndrome. J. Pediat. 136: 390–393, 2000.

[172845]

[172846]

[172847]17619.

[172848]Greenfield, A.; Carrel, L.; Pennisi, D.; Philippe, C.; Quaderi,

[172849]N.; Siggers, P.; Steiner, K.; Tam, P. P. L.; Monaco, A. P.;
Willard,

[172850]H. F.; Koopman, P.: The UTX gene escapes X inactivation
in mice and

[172851]humans. Hum. Molec. Genet. 7: 737–742, 1998.

[172852]

[172853]

[172854]17620.

[172855]Greenfield, A. J.; Scott, D.; Pennisi, D.; Ehrmann, I.; Ellis,

[172856]P.; Cooper, L.; Simpson, E.; Koopman, P.: An H-YDb epi–

tope is encoded

[172857]by a novel mouse Y chromosome gene. Nature Genet. 14:
474–478, 1996.

[172858]

[172859]

[172860]17621.

[172861]Caput, D.; Laurent, P.; Kaghad, M.; Lelias, J.–M.; Lefort, S.;

[172862]Vita, N.; Ferrara, P.: Cloning and characterization of a spe-
cific

[172863]interleukin (IL)–13 binding protein structurally related to
the IL–5

[172864]receptor alpha chain. J. Biol. Chem. 271: 16921–16926,
1996.

[172865]

[172866]

[172867]17622.

[172868]Guo, J.; Apiou, F.; Mellerin, M.–P.; Lebeau, B.; Jacques, Y.;
Minvielle,

[172869]S.: Chromosome mapping and expression of the human
interleukin–13

[172870]receptor. Genomics 42: 141–145, 1997.

[172871]

[172872]

[172873]17623.

[172874]Lin, C.-S.; Aebersold, R. H.; Leavitt, J.: Correction of the
N-terminal

[172875]sequences of the human plastin isoforms by using an-
chored polymerase

[172876]chain reaction: identification of a potential calcium-bind-
ing domain. Molec.

[172877]Cell. Biol. 10: 1818-1821, 1990.

[172878]

[172879]

[172880]17624.

[172881]Fukuda, M. N.; Sato, T.; Nakayama, J.; Klier, G.; Mikami,
M.; Aoki,

[172882]D.; Nozawa, S.: Trophinin and tastin, a novel cell adhesion
molecule

[172883]complex with potential involvement in embryo implanta-
tion. Genes

[172884]Dev. 9: 1199-1210, 1995.

[172885]

[172886]

[172887]17625.

[172888]Pack, S. D.; Tanigami, A.; Ledbetter, D. H.; Sato, T.;
Fukuda,

[172889]M. N.: Assignment of trophoblast/endometrial epithelium
cell adhesion

[172890]molecule trophinin gene TRO to human chromosome
bands Xp11.22–p11.21

[172891]by in situ hybridization. Cytogenet. Cell Genet. 79:
123–124, 1997.

[172892]

[172893]

[172894]17626.

[172895]Suzuki, N.; Nakayama, J.; Shih, I. M.; Aoki, D.; Nozawa, S.;
Fukuda,

[172896]M. N.: Expression of trophinin, tastin, and bystin by tro-
phoblast

[172897]and endometrial cells in human placenta. Biol. Reprod. 60:
621–627,

[172898]1999.

[172899]

[172900]

[172901]17627.

[172902]Sana, T. R.; Debets, R.; Timans, J. C.; Bazan, J. F.;
Kastelein,

[172903]R. A.: Computational identification, cloning, and charac-
terization

[172904]of IL1R9, a novel interleukin-1 receptor-like gene en-
coded over an

[172905]unusually large interval of human chromosome
Xq22.2-q22.3. Genomics 69:

[172906]252-262, 2000.

[172907]

[172908]

[172909]17628.

[172910]Bech-Hansen, N. T.; Naylor, M. J.; Maybaum, T. A.;
Sparkes, R.

[172911]L.; Koop, B.; Birch, D. G.; Bergen, A. A. B.; Prinsen, C. F.
M.; Polomeno,

[172912]R. C.; Gal, A.; Drack, A. V.; Musarella, M. A.; Jacobson, S.
G.; Young,

[172913]R. S. L.; Weleber, R. G.: Mutations in NYX, encoding the leucine-rich

[172914]proteoglycan nyctalopin, cause X-linked complete congenital stationary

[172915]night blindness. Nature Genet. 26: 319–323, 2000.

[172916]

[172917]

[172918]17629.

[172919]Pusch, C. M.; Zeitze, C.; Brandau, O.; Pesch, K.; Achatz, H.; Feil,

[172920]S.; Scharfe, C.; Maurer, J.; Jacobi, F. K.; Pinckes, A.; Andreasson,

[172921]S.; Hardcastle, A.; Wissinger, B.; Berger, W.; Meindl, A.: The complete

[172922]form of X-linked congenital stationary night blindness is caused by

[172923]mutations in a gene encoding a leucine-rich repeat protein. Nature

[172924]Genet. 26: 324–327, 2000.

[172925]

[172926]

[172927]17630.

[172928]Isbrandt, D.; Leicher, T.; Waldschutz, R.; Zhu, X.; Luhmann, U.;

[172929]Michel, U.; Sauter, K.; Pongs, O.: Gene structures and expression

[172930]profiles of three human KCND (Kv4) potassium channels mediating A-type

[172931]currents I(to) and I(sa). Genomics 64: 144–154, 2000.

[172932]

[172933]

[172934]17631.

[172935]Davies, J. P.; Cotter, P. D.; Ioannou, Y. A.: Cloning and mapping

[172936]of human Rab7 and Rab9 cDNA sequences and identification of a Rab9

[172937]pseudogene. Genomics 41: 131–134, 1997.

[172938]

[172939]

[172940]17632.

[172941]Brinkmann, U.; Vasmatzis, G.; Lee, B.; Pastan, I.: Novel

genes

[172942]in the PAGE and GAGE family of tumor antigens found by
homology walking

[172943]in the dbEST database. Cancer Res. 59: 1445–1448, 1999.

[172944]

[172945]

[172946]17633.

[172947]Liu, X. F.; Helman, L. J.; Yeung, C.; Bera, T. K.; Lee, B.; Pas-
tan,

[172948]L.: XAGE-1, a new gene that is frequently expressed in Ew-
ing's sarcoma. Cancer

[172949]Res. 60: 4752–4755, 2000.

[172950]

[172951]

[172952]17634.

[172953]Lahn, B. T.; Page, D. C.: Functional coherence of the hu-
man Y

[172954]chromosome. Science 278: 675–680, 1997.

[172955]

[172956]

[172957]17635.

[172958]Chung, J.; Lee, S.-G.; Song, K.: Identification of a human
homolog

[172959]of a putative RNA helicase gene (mDEAD3) expressed in
mouse erythroid

[172960]cells. Korean J. Biochem. 27: 193–197, 1995.

[172961]

[172962]

[172963]17636.

[172964]Park, S. H.; Lee, S.-G.; Kim, Y.; Song, K.: Assignment of a
human

[172965]putative RNA helicase gene, DDX3, to human X chromo-
some bands p11.3–p11.23. Cytogenet.

[172966]Cell Genet. 81: 178–179, 1998.

[172967]

[172968]

[172969]17637.

[172970]Gasper, N. J.; Kinzy, T. G.; Scherer, B. J.; Humbelin, M.;

Hershey,

[172971]J. W. B.; Merrick, W. C.: Translation initiation factor eIF-2:
cloning

[172972]and expression of the human cDNA encoding the gamma-
subunit. J. Biol.

[172973]Chem. 269: 3415-3422, 1994.

[172974]

[172975]

[172976]17638.

[172977]Ehrmann, I. E.; Ellis, P. S.; Mazeyrat, S.; Duthie, S.; Brock-
dorff,

[172978]N.; Mattei, M. G.; Gavin, M. A.; Affara, N. A.; Brown, G. M.;
Simpson,

[172979]E.; Mitchell, M. J.; Scott, D. M.: Characterization of genes
encoding

[172980]translation initiation factor eIF-2-gamma in mouse and
human: sex

[172981]chromosome localization, escape from X-inactivation and
evolution. Hum.

[172982]Molec. Genet. 7: 1725-1737, 1998.

[172983]

[172984]

[172985]17639.

[172986]Lee, S. M. Y.; Tsui, S. K. W.; Chan, K. K.; Garcia-Barcelo, M.;

[172987]Waye, M. M. Y.; Fung, K. P.; Liew, C. C.; Lee, C. Y.: Chromosomal

[172988]mapping, tissue distribution and cDNA sequence of four-and-a-half

[172989]LIM domain protein 1 (FHL1). Gene 216: 163–170, 1998.

[172990]

[172991]

[172992]17640.

[172993]Morgan, M. J.; Madgwick, A. J.; Charleston, B.; Pell, J. M.; Loughna,

[172994]P. T.: The developmental regulation of a novel muscle LIM-protein. Biochem.

[172995]Biophys. Res. Commun. 212: 840–846, 1995.

[172996]

[172997]

[172998]17641.

[172999]Morgan, M. J.; Madgwick, A. J. A.: Slim defines a novel family

[173000]of LIM–proteins expressed in skeletal muscle. Biochem. Biophys. Res.

[173001]Commun. 225: 632–638, 1996.

[173002]

[173003]

[173004]17642.

[173005]Bard, J. A.; Zgombick, J.; Adham, N.; Vaysse, P.; Branchek, T.

[173006]A.; Weinshank, R. L.: Cloning of a novel human serotonin receptor

[173007](5–HT–7) positively linked to adenylate cyclase. J. Biol. Chem. 268:

[173008]23422–23426, 1993.

[173009]

[173010]

[173011]17643.

[173012]Gelernter, J.; Rao, P. A.; Pauls, D. L.; Hamblin, M. W.; Sibley,

[173013]D. R.; Kidd, K. K.: Assignment of the 5HT7 receptor gene
(HTR7) to

[173014]chromosome 10q and exclusion of genetic linkage with
Tourette syndrome. Genomics 26:

[173015]207–209, 1995.

[173016]

[173017]

[173018]17644.

[173019]Kenakin, T. P.; Bond, R. A.; Bonner, T. I.: II. Definition of

[173020]pharmacological receptors. Pharm. Rev. 44: 351–362,
1992.

[173021]

[173022]

[173023]17645.

[173024]Lassig, J. P.; Vachirasomtoon, K.; Hartzell, K.; Leventhal,
M.;

[173025]Courchesne, E.; Courchesne, R.; Lord, C.; Leventhal, B. L.;
Cook,

[173026]E. H., Jr.: Physical mapping of the serotonin 5-HT(7) re-
ceptor gene

[173027](HTR7) to chromosome 10 and pseudogene (HTR7P) to chromosome 12, and

[173028]testing of linkage disequilibrium between HTR7 and autistic disorder. Am.

[173029]J. Med. Genet. 88: 472–475, 1999.

[173030]

[173031]

[173032]17646.

[173033]Ruat, M.; Traiffort, E.; Leurs, R.; Tardivel–Lacombe, J.; Diaz,

[173034]J.; Arrang, J.–M.; Schwartz, J.–C.: Molecular cloning, characterization,

[173035]and localization of a high–affinity serotonin receptor (5–HT–7) activating

[173036]cAMP formation. Proc. Nat. Acad. Sci. 90: 8547–8551, 1993.

[173037]

[173038]

[173039]17647.

[173040]Tsou, A.; Kosaka, A.; Bach, C.; Zuppan, P.; Yee, C.; Tom, L.; Alvarez,

[173041]R.; Ramsey, S.; Bonhaus, D. W.; Stefanich, E.; Jakeman, L.;
Eglen,

[173042]R. M.; Chan, H. W.: Cloning and expression of a
5-hydroxytryptamine-7

[173043]receptor positively coupled to adenylyl cyclase. J. Neu-
rochem. 63:

[173044]456-464, 1994.

[173045]

[173046]

[173047]17648.

[173048]Gelernter, J.; Kruger, S.; Pakstis, A. J.; Amara, S.; Kidd, K.

[173049]K.: Linkage mapping of norepinephrine transporter pro-
tein and serotonin

[173050]transporter protein genes. (Abstract) Am. J. Hum. Genet.
53 (suppl.):

[173051]A1003, 1993.

[173052]

[173053]

[173054]17649.

[173055]Gelernter, J.; Pakstis, A. J.; Kidd, K. K.: Linkage mapping of

[173056]serotonin transporter protein gene SLC6A4 on chromosome 17. Hum.

[173057]Genet. 95: 677–680, 1995.

[173058]

[173059]

[173060]17650.

[173061]Gregor, P.; Patel, A.; Shimada, S.; Lin, C.–L.; Rochelle, J. M.;

[173062]Kitayama, S.; Seldin, M. F.; Uhl, G. R.: Murine serotonin transporter:

[173063]sequence and localization to chromosome 11. Mammalian Genome 4:

[173064]283–284, 1993.

[173065]

[173066]

[173067]17651.

[173068]Hariri, A. R.; Mattay, V. S.; Tessitore, A.; Kolachana, B.; Fera,

[173069]F.; Goldman, D.; Egan, M. F.; Weinberger, D. R.: Serotonin transporter

[173070]genetic variation and the response of the human amygdala. Science 297:

[173071]400–403, 2002.

[173072]

[173073]

[173074]17652.

[173075]Heils, A.; Teufel, A.; Petri, S.; Seemann, M.; Bengel, D.;
Balling,

[173076]U.; Riederer, P.; Lesch, K. P.: Functional promoter and
polyadenylation

[173077]site mapping of the human serotonin (5-HT) transporter
gene. J. Neural

[173078]Transm. 102: 247–254, 1995.

[173079]

[173080]

[173081]17653.

[173082]Heils, A.; Teufel, A.; Petri, S.; Stober, G.; Riederer, P.; Bengel,

[173083]D.; Lesch, K. P.: Allelic variation of human serotonin
transporter

[173084]gene expression. J. Neurochem. 66: 2621–2624, 1996.

[173085]

[173086]

[173087]17654.

[173088]Kim, S.–J.; Cox, N.; Courchesne, R.; Lord, C.; Corsello, C.;
Akshoomoff,

[173089]N.; Guter, S.; Leventhal, B. L.; Courchesne, E.; Cook, E. H.,
Jr.

[173090]: Transmission disequilibrium mapping at the serotonin
transporter

[173091]gene (SLC6A4) region in autistic disorder. Molec. Psychiat.
7: 278–288,

[173092]2002.

[173093]

[173094]

[173095]17655.

[173096]Lesch, K. P.; Balling, U.; Gross, J.; Strauss, K.; Wolozin, B.

[173097]L.; Murphy, D. L.: Organisation of the human serotonin
transporter

[173098]gene. J. Neural Trans. 95: 157–162, 1994.

[173099]

[173100]

[173101]17656.

[173102]Chern, C. J.; Beutler, E.: Biochemical and electrophoretic studies

[173103]of erythrocyte pyridoxine kinase in white and black Americans. Am.

[173104]J. Hum. Genet. 28: 9–17, 1976.

[173105]

[173106]

[173107]17657.

[173108]Hanna, M. C.; Turner, A. J.; Kirkness, E. F.: Human pyridoxal

[173109]kinase: cDNA cloning, expression, and modulation by ligands of the

[173110]benzodiazepine receptor. J. Biol. Chem. 272: 10756–10760, 1997.

[173111]

[173112]

[173113]17658.

[173114]Dougherty, K. M.; Brandriss, M. C.; Valle, D.: Cloning human pyrroline-5-carboxylate

[173115]reductase cDNA by complementation in *Saccharomyces cerevisiae*. J.

[173116]Biol. Chem. 267: 871-875, 1992.

[173117]

[173118]

[173119]17659.

[173120]Merrill, M. J.; Yeh, G. C.; Phang, J. M.: Purified human erythrocyte

[173121]pyrroline-5-carboxylate reductase: preferential oxidation of NADPH. J.

[173122]Biol. Chem. 264: 9352-9358, 1989.

[173123]

[173124]

[173125]17660.

[173126]Brissenden, J. E.; Caras, I.; Thelander, L.; Francke, U.: The

[173127]structural gene for the M1 subunit of human ribonucleotide reductase

[173128]maps to chromosome 11, band p15, in human and to

chromosome 7 in mouse. Exp.

[173129]Cell Res. 174: 302–308, 1988.

[173130]

[173131]

[173132]17661.

[173133]Byrne, J.; Smith, P.: Human polymorphic probe pE1.8 de-
tects SacI

[173134]polymorphism in the ribonucleotide reductase M1 subunit
gene. Hum.

[173135]Genet. 87: 376 only, 1991.

[173136]

[173137]

[173138]17662.

[173139]Barletta, C.; Druck, T.; LaForgia, S.; Calabretta, B.;
Drabkin,

[173140]H.; Patterson, D.; Croce, C. M.; Huebner, K.: Chromosome
locations

[173141]of the MYB related genes, AMYB and BMYB. Cancer Res.
51: 3821–3824,

[173142]1991.

[173143]

[173144]

[173145]17663.

[173146]Nomura, N.; Takahashi, M.; Matsui, M.; Ishii, S.; Date, T.;
Sasamoto,

[173147]S.; Ishizaki, R.: Isolation of human cDNA clones of MYB-
related genes,

[173148]A-MYB and B-MYB. Nucleic Acids Res. 16: 11075-11089,
1988.

[173149]

[173150]

[173151]17664.

[173152]Takahashi, T.; Nakagoshi, H.; Sarai, A.; Nomura, N.; Ya-
mamoto,

[173153]T.; Ishii, S.: Human A-myb gene encodes a transcriptional
activator

[173154]containing the negative regulatory domains. FEBS Lett.
358: 89-96,

[173155]1995.

[173156]

[173157]

[173158]17665.

[173159]Boylan, K. B.; Takahashi, N.; Diamond, M.; Hood, L. E.;
Prusiner,

[173160]S. B.: DNA length polymorphism located 5-prime to the
human myelin

[173161]basic protein gene. Am. J. Hum. Genet. 40: 387-400,
1987.

[173162]

[173163]

[173164]17666.

[173165]Dayhoff, M. O.: Atlas of Protein Sequence and Structure.
Myelin

[173166]membrane encephalitogenic protein. Washington: National
Biomedical

[173167]Research Foundation (pub.) 5: 1972. Pp. D324 only.

[173168]

[173169]

[173170]17667.

[173171]Eylar, E. H.; Brostoff, S.; Hashim, G.; Westall, F. C.: Basic

[173172]A1 protein of the myelin membrane: the complete amino acid sequence. J.

[173173]Biol. Chem. 246: 5770–5784, 1971.

[173174]

[173175]

[173176]17668.

[173177]Gomez, C. M.; Muggleton–Harris, A. L.; Whittingham, D. G.; Hood,

[173178]L. E.; Readhead, C.: Rapid preimplantation detection of mutant (shiverer)

[173179]and normal alleles of the mouse myelin basic protein gene allowing

[173180]selective implantation and birth of live young. Proc. Nat. Acad.

[173181]Sci. 87: 4481–4484, 1990.

[173182]

[173183]

[173184]17669.

[173185]Kamholz, J.; de Ferra, F.; Puckett, C.; Lazzarini, R.: Identification

[173186]of three forms of human myelin basic protein by cDNA cloning. Proc.

[173187]Nat. Acad. Sci. 83: 4962–4966, 1986.

[173188]

[173189]

[173190]17670.

[173191]Kamholz, J.; Spielman, R.; Gogolin, K.; Modi, W.; O'Brien, S.;

[173192]Lazzarini, R.: The human myelin–basic–protein gene: chromosomal localization

[173193]and RFLP analysis. Am. J. Hum. Genet. 40: 365–373, 1987.

[173194]

[173195]

[173196]17671.

[173197]Lalley, P. A.; McKusick, V. A.: Report of the committee on comparative

[173198]mapping (HGM8). Cytogenet. Cell Genet. 40: 536–566, 1985.

[173199]

[173200]

[173201]17672.

[173202]Martensen, R. E.: Myelin basic protein speciation. Prog.
Clin.

[173203]Biol. Res. 146: 511–521, 1984.

[173204]

[173205]

[173206]17673.

[173207]Marty, M. C.; Alliot, F.; Rutin, J.; Fritz, R.; Trisler, D.; Pes-
sac,

[173208]B.: The myelin basic protein gene is expressed in differen-
tiated

[173209]blood cell lineages and in hemopoietic progenitors. Proc.
Nat. Acad.

[173210]Sci. 99: 8856–8861, 2002.

[173211]

[173212]

[173213]17674.

[173214]Molineaux, S. M.; Engh, H.; de Ferra, F.; Hudson, L.; Laz-
zarini,

[173215]R. A.: Recombination within the myelin basic protein gene

created

[173216]the dysmyelinating shiverer mouse mutation. Proc. Nat. Acad. Sci. 83:

[173217]7542–7546, 1986.

[173218]

[173219]

[173220]17675.

[173221]Popko, B.; Puckett, C.; Lai, E.; Shine, H. D.; Readhead, C.;
Takahashi,

[173222]N.; Hunt, S. W., III; Sidman, R. L.; Hood, L.: Myelin deficient
mice:

[173223]expression of myelin basic protein and generation of mice
with varying

[173224]levels of myelin. Cell 48: 713–721, 1987.

[173225]

[173226]

[173227]17676.

[173228]Readhead, C.; Popko, B.; Takahashi, N.; Shine, H. D.;
Saavedra,

[173229]R. A.; Sidman, R. L.; Hood, L.: Expression of a myelin basic

protein

[173230]gene in transgenic shiverer mice: correction of the dys-
myelinating

[173231]phenotype. Cell 48: 703–712, 1987.

[173232]

[173233]

[173234]17677.

[173235]Roach, A.; Boylan, K.; Horvath, S.; Prusiner, S. B.; Hood, L.

[173236]E.: Characterization of cloned cDNA representing rat
myelin basic

[173237]protein: absence of expression in shiverer mutant mice.
Cell 34:

[173238]799–806, 1983.

[173239]

[173240]

[173241]17678.

[173242]Roach, A.; Takahashi, N.; Pravtcheva, D.; Ruddle, F.; Hood,
L.

[173243]: Chromosomal mapping of mouse myelin basic protein
gene and structure

[173244]and transcription of the partially deleted gene in shiverer
mutant

[173245]mice. Cell 42: 149–155, 1985.

[173246]

[173247]

[173248]17679.

[173249]Saxe, D. F.; Takahashi, N.; Hood, L.; Simon, M. I.: Local-
ization

[173250]of the human myelin basic protein gene (MBP) to region
18q22-qter

[173251]by in situ hybridization. Cytogenet. Cell Genet. 39:
246-249, 1985.

[173252]

[173253]

[173254]17680.

[173255]Sheremata, W.; Cosgrove, J. B. R.; Hylar, E. H.: Cellular hy-
persensitivity

[173256]to basic myelin (A1) protein and clinical multiple sclerosis.
New

[173257]Eng. J. Med. 291: 14-17, 1974.

[173258]

[173259]

[173260]17681.

[173261]Sidman, R.: Personal Communication. Boston, Mass. 1983.

[173262]

[173263]

[173264]17682.

[173265]Sidman, R. L.; Conover, C. S.; Carson, J. H.: Shiverer gene
maps

[173266]near the distal end of chromosome 18 in the house
mouse. Cytogenet.

[173267]Cell Genet. 39: 241–245, 1985.

[173268]

[173269]

[173270]17683.

[173271]Sparkes, R. S.; Mohandas, T.; Heinzmann, C.; Roth, H. J.;
Klisak,

[173272]I.; Campagnoni, A. T.: Assignment of the myelin basic
protein gene

[173273]to human chromosome 18q22–qter. Hum. Genet. 75:
147–150, 1987.

[173274]

[173275]

[173276]17684.

[173277]Takahashi, N.; Roach, A.; Teplow, D. B.; Prusiner, S. B.;
Hood,

[173278]L.: Cloning and characterization of the myelin basic protein gene

[173279]from mouse: one gene can encode both 14 kd and 18.5 kd MBPs by alternate

[173280]use of exons. Cell 42: 139–148, 1985.

[173281]

[173282]

[173283]17685.

[173284]Barouch, L. A.; Harrison, R. W.; Skaf, M. W.; Rosas, G. O.; Cappola,

[173285]T. P.; Kobeissi, Z. A.; Hobai, I. A.; Lemmon, C. A.; Burnett, A. L.;

[173286]O'Rourke, B.; Rodriguez, E. R.; Huang, P. L.; Lima, J. A. C.; Berkowitz,

[173287]D. E.; Hare, J. M.: Nitric oxide regulates the heart by spatial confinement

[173288]of nitric oxide synthase isoforms. Nature 416: 337–340, 2002.

[173289]

[173290]

[173291]17686.

[173292]Dimmeler, S.; Fleming, I.; Fisslthaler, B.; Hermann, C.;
Busse,

[173293]R.; Zeiher, A. M.: Activation of nitric oxide synthase in en-
dothelial

[173294]cells by Akt-dependent phosphorylation. Nature 399:
601–605, 1999.

[173295]

[173296]

[173297]17687.

[173298]Fulton, D.; Gratton, J.–P.; McCabe, T. J.; Fontana, J.; Fujio,

[173299]Y.; Walsh, K.; Franke, T. F.; Papapetropoulos, A.; Sessa, W.
C.:

[173300]Regulation of endothelium-derived nitric oxide production
by the protein

[173301]kinase Akt. Nature 399: 597–601, 1999.

[173302]

[173303]

[173304]17688.

[173305]Kruh, G. D.; King, C. R.; Kraus, M. H.; Popescu, N. C.;
Amsbaugh,

[173306]S. C.; McBride, W. O.; Aaronson, S. A.: A novel human gene
closely

[173307]related to the abl proto-oncogene. Science 234:
1545–1548, 1986.

[173308]

[173309]

[173310]17689.

[173311]Arnaud, J.; Vavrusa, B.; Sevin, J.; Constans, J.: Human red-
cell

[173312]acid phosphatase (ACP1): a new mutant (ACP1*KUK) de-
tected by isoelectric

[173313]focusing, kinetics of thermostability and substrate activity.
Hum.

[173314]Hered. 39: 288–293, 1989.

[173315]

[173316]

[173317]17690.

[173318]Arnaud, J.; Vavrusa, B.; Wiederanders, G.; Constans, J.:
Human

[173319]red-cell acid phosphatase (ACP1): kinetic and thermody-
namic characterization

[173320]of the KUK variant. Hum. Hered. 42: 140–142, 1992.

[173321]

[173322]

[173323]17691.

[173324]Beemer, F. A.; van der Heiden, C.; Van Hemel, J. O.;
Jansen, M.

[173325]: Letter to the editors. (Letter) Clin. Genet. 24: 151, 1983.

[173326]

[173327]

[173328]17692.

[173329]Berg, K.: Close linkage between APOB and ACP1 excluded.
(Abstract) Cytogenet.

[173330]Cell Genet. 46: 580, 1987.

[173331]

[173332]

[173333]17693.

[173334]Bottini, E.; Carapella, E.; Orzalesi, M.; Lucarelli, P.; Pas-
cone,

[173335]R.; Gloria-Bottini, F.; Coccia, M.: Is there a role of erythro-

cyte

[173336]acid phosphatase polymorphism in intrauterine development? (Letter) Am.

[173337]J. Hum. Genet. 32: 764–767, 1980.

[173338]

[173339]

[173340]17694.

[173341]Seldin, M. F.; Kruh, G. D.: Mapping of Abll within a conserved

[173342]linkage group on distal mouse chromosome 1 syntenic with human chromosome

[173343]1 using an interspecific cross. Genomics 4: 221–223, 1989.

[173344]

[173345]

[173346]17695.

[173347]Bello, M. J.; Salagnon, N.; Rey, J. A.; Guichaoua, M. R.; Berge-Lefranc,

[173348]J. L.; Jordan, B. R.; Luciani, J. M.: Precise in situ localization

[173349]of NCAM, ETS1, and D11S29 on human meiotic chromo-

somes. Cytogenet.

[173350]Cell Genet. 52: 7–10, 1989.

[173351]

[173352]

[173353]17696.

[173354]Gold, D. P.; van Dongen, J. J. M.; Morton, C. C.; Bruns, G.
A.

[173355]P.; van den Elsen, P.; Geurts van Kessel, A. H. M.; Terhorst,
C.:

[173356]The gene encoding the epsilon subunit of the T3/T-cell
receptor complex

[173357]maps to chromosome 11 in humans and to chromosome
9 in mice. Proc.

[173358]Nat. Acad. Sci. 84: 1664–1668, 1987.

[173359]

[173360]

[173361]17697.

[173362]Ohtani, N.; Zebedee, Z.; Huot, T. J. G.; Stinson, J. A.; Sugi-
moto,

[173363]M.; Ohashi, Y.; Sharrocks, A. D.; Peters, G.; Hara, E.: Op–

posing

[173364]effects of Ets and Id proteins on p16(INK4A) expression
during cellular

[173365]senescence. Nature 409: 1067–1070, 2001.

[173366]

[173367]

[173368]17698.

[173369]Sacchi, N.; Watson, D. K.; Geurts van Kessel, A. H. M.;
Hagemeijer,

[173370]A.; Kersey, J.; Drabkin, H. D.; Patterson, D.; Papas, T. S.:
Hu-ets-1

[173371]and Hu-ets-2 genes are transposed in acute leukemias
with (4;11) and

[173372](8;21) translocations. Science 231: 379–382, 1986.

[173373]

[173374]

[173375]17699.

[173376]Watson, D. K.; McWilliams-Smith, M. J.; Kozak, C.; Reeves,
R.;

[173377]Gearhart, J.; Nunn, M. F.; Nash, W.; Fowle, J. R., III; Dues–

berg,

[173378]P.; Papas, T. S.; O'Brien, S. J.: Conserved chromosomal positions

[173379]of dual domains of the ets protooncogene in cats, mice, and humans. Proc.

[173380]Nat. Acad. Sci. 83: 1792–1796, 1986.

[173381]

[173382]

[173383]17700.

[173384]Watson, D. K.; McWilliams–Smith, M. J.; Nunn, M. F.; Duesberg,

[173385]P. H.; O'Brien, S. J.; Papas, T. S.: The ets sequence from the transforming

[173386]gene of avian erythroblastosis virus, E26, has unique domains on human

[173387]chromosomes 11 and 21: both loci are transcriptionally active. Proc.

[173388]Nat. Acad. Sci. 82: 7294–7298, 1985.

[173389]

[173390]

[173391]17701.

[173392]Bellacosa, A.; Franke, T. F.; Gonzalez-Portal, M. E.; Datta, K.;

[173393]Taguchi, T.; Gardner, J.; Cheng, J. Q.; Testa, J. R.; Tsichlis, P.

[173394]N.: Structure, expression and chromosomal mapping of c-akt: relationship

[173395]to v-akt and its implications. Oncogene 8: 745-754, 1993.

[173396]

[173397]

[173398]17702.

[173399]Brunet, A.; Bonni, A.; Zigmond, M. J.; Lin, M. Z.; Juo, P.; Hu,

[173400]L. S.; Anderson, M. J.; Arden, K. C.; Blenis, J.; Greenberg, M. E.

[173401]: Akt promotes cell survival by phosphorylating and inhibiting a Forkhead

[173402]transcription factor. Cell 96: 857-868, 1999.

[173403]

[173404]

[173405]17703.

[173406]Chen, W. S.; Xu, P.-Z.; Gottlob, K.; Chen, M.-L.; Sokol, K.;
Shiyanova,

[173407]T.; Roninson, I.; Weng, W.; Suzuki, R.; Tobe, K.; Kadowaki,
T.; Hay,

[173408]N.: Growth retardation and increased apoptosis in mice
with homozygous

[173409]disruption of the akt1 gene. Genes Dev. 15: 2203–2208,
2001.

[173410]

[173411]

[173412]17704.

[173413]Condorelli, G.; Drusco, A.; Stassi, G.; Bellacosa, A.; Ron-
carati,

[173414]R.; Iaccarino, G.; Russo, M. A.; Gu, Y.; Dalton, N.; Chung,
C.; Latronico,

[173415]M. V. G.; Napoli, C.; Sadoshima, J.; Croce, C. M.; Ross, J.,
Jr.:

[173416]Akt induces enhanced myocardial contractility and cell
size in vivo

[173417]in transgenic mice. Proc. Nat. Acad. Sci. 99:
12333–12338, 2002.

[173418]

[173419]

[173420]17705.

[173421]Ellis, N. A.: Ecce Ohno. Nature Genet. 10: 373–375, 1995.

[173422]

[173423]

[173424]17706.

[173425]Haldane, J. B. S.: Sex ratio and unisexual sterility in hybrid

[173426]animals. J. Genet. 12: 101–109, 1922.

[173427]

[173428]

[173429]17707.

[173430]Milatovich, A.; Kitamura, T.; Miyajima, A.; Francke, U.:

Gene

[173431]for the alpha-subunit of the human interleukin-3 receptor

(IL3RA)

[173432]localized to the X-Y pseudoautosomal region. Am. J. Hum.

Genet. 53:

[173433]1146–1153, 1993.

[173434]

[173435]

[173436]17708.

[173437]Ohno, S.: Sex Chromosomes and Sex-linked Genes. Berlin
and New

[173438]York: Springer (pub.) 1967.

[173439]

[173440]

[173441]17709.

[173442]Palmer, S.; Perry, J.; Ashworth, A.: A contravention of
Ohno's

[173443]law in mice. Nature Genet. 10: 472–476, 1995.

[173444]

[173445]

[173446]17710.

[173447]Rugarli, E. I.; Adler, D. A.; Borsani, G.; Tsuchiya, K.;
Franco,

[173448]B.; Hauge, X.; Disteche, C.; Chapman, V.; Ballabio, A.: Dif-
ferent

[173449]chromosomal localization of the Clcn4 gene in Mus spre-

tus and C57BL/6J

[173450]mice. Nature Genet. 10: 466–471, 1995.

[173451]

[173452]

[173453]17711.

[173454]Schnur, R. E.; Wick, P. A.: Intragenic Taql restriction fragment
ment

[173455]length polymorphism (RFLP) in CLCN4, between the loci
for X-linked

[173456]ocular albinism (OA1) and microphthalmia with linear skin
defects

[173457]syndrome (MLS). Hum. Genet. 95: 594–595, 1995.

[173458]

[173459]

[173460]17712.

[173461]van Slegtenhorst, M. A.; Bassi, M. T.; Borsani, G.; Wape-
naar, M.

[173462]C.; Ferrero, G. B.; de Conciliis, L.; Rugarli, E. I.; Grillo, A.;

[173463]Franco, B.; Zoghbi, H. Y.; Ballabio, A.: A gene from the
Xp22.3 region

[173464]shares homology with voltage-gated chloride channels.

Hum. Molec.

[173465]Genet. 3: 547-552, 1994.

[173466]

[173467]

[173468]17713.

[173469]Curry, C. J. R.; Magenis, R. E.; Brown, M.; Lanman, J. T., Jr.;

[173470]Tsai, J.; O'Lague, P.; Goodfellow, P.; Mohandas, T.;

Bergner, E. A.;

[173471]Shapiro, L. J.: Inherited chondrodysplasia punctata due to
a deletion

[173472]of the terminal short arm of an X chromosome. New Eng.
J. Med. 311:

[173473]1010-1015, 1984.

[173474]

[173475]

[173476]17714.

[173477]Maslen, G. L.; Boyd, Y.: Comparative mapping of the Grpr
locus

[173478]on the X chromosomes of man and mouse. Genomics 17:

106–109, 1993.

[173479]

[173480]

[173481]17715.

[173482]Schantz, L. J.; Naylor, S. L.; Giladi, E.; Spindel, E. R.: Assignment
segment

[173483]of the GRP receptor gene to the human X chromosome.
(Abstract) Cytogenet.

[173484]Cell Genet. 58: 2085–2086, 1991.

[173485]

[173486]

[173487]17716.

[173488]Shiraishi, M.; Alitalo, T.; Sekiya, T.: The chromosomal organization
ganization

[173489]of the human endogenous retrovirus–like sequence HERV–
H: clustering

[173490]of the HERV–H sequences in a 300–kb region close to the
GRPR locus

[173491]on the X chromosome. DNA Res. 3: 425–429, 1996.

[173492]

[173493]

[173494]17717.

[173495]Shriver, S. P.; Bourdeau, H. A.; Gubish, C. T.; Tirpak, D. L.;

[173496]Davis, A. L. G.; Luketich, J. D.; Siegfried, J. M.: Sex-specific

[173497]expression of gastrin-releasing peptide receptor: relationship to

[173498]smoking history and risk of lung cancer. J. Nat. Cancer Inst. 92:

[173499]24-33, 2000.

[173500]

[173501]

[173502]17718.

[173503]Spindel, E. R.; Giladi, E.; Brehm, P.; Goodman, R. H.;

Segerson,

[173504]T. P.: Cloning and functional characterization of a complementary

[173505]DNA encoding the murine fibroblast bombesin/gastrin-releasing peptide

[173506]receptor. Molec. Endocr. 4: 1956-1963, 1990.

[173507]

[173508]

[173509]17719.

[173510]Ma, K.; Sharkey, A.; Kirsch, S.; Vogt, P.; Keil, R.; Hargreave,

[173511]T. B.; McBeath, S.; Chandley, A. C.: Towards the molecular
localisation

[173512]of the AZF locus: mapping of microdeletions in azoosper-
mic men within

[173513]14 subintervals of interval 6 of the human Y chromosome.
Hum. Molec.

[173514]Genet. 1: 29-33, 1992.

[173515]

[173516]

[173517]17720.

[173518]Adam, A.; Tippett, P.; Gavin, J.; Noades, J.; Sanger, R.;
Race,

[173519]R. R.: The linkage relation of Xg to G6PD in Israelis: the
evidence

[173520]of a second series of families. Am. J. Hum. Genet. 30:
211-218,

[173521]1966.

[173522]

[173523]

[173524]17721.

[173525]Ahluwalia, A.; Corcoran, C. M.; Vulliamy, T. J.; Ishwad, C. S.;

[173526]Naidu, J. M.; Argusti, A.; Stevens, D. J.; Mason, P. J.; Luz-zatto,

[173527]L.: G6PD Kalyan and G6PD Kerala; two deficient variants in India

[173528]caused by the same 317 glu-to-lys mutation. Hum. Molec. Genet. 3:

[173529]209-210, 1992.

[173530]

[173531]

[173532]17722.

[173533]Aksoy, K.; Yuregir, G. T.; Dikmen, N.; Unlukurt, I.: Three new

[173534]G6PD variants, G6PD Adana, G6PD Samandag, and G6PD Balcali in Cukurova,

[173535]Turkey. Hum. Genet. 76: 199-201, 1987.

[173536]

[173537]

[173538]17723.

[173539]Alfinito, F.; Cimmino, A.; Ferraro, F.; Cubellis, M. V.;
Vitagliano,

[173540]L.; Francese, M.; Zagari, A.; Rotoli, B.; Filosa, S.; Martini, G.

[173541]: Molecular characterization of G6PD deficiency in South-
ern Italy:

[173542]heterogeneity, correlation genotype-phenotype and de-
scription of a

[173543]new variant (G6PD Neapolis). Brit. J. Haemat. 98: 41-46,
1997.

[173544]

[173545]

[173546]17724.

[173547]Alperin, J. B.; Mills, G. C.: New variants of glucose-
6-phosphate

[173548]dehydrogenase (G6PD). Clin. Res. 20: 76, 1972.

[173549]

[173550]

[173551]17725.

[173552]Azevedo, E.; Kirkman, H. N.; Morrow, A. C.; Motulsky, A.
G.: Variants

[173553]of red cell glucose-6-phosphate dehydrogenase among
Asiatic Indians. Ann.

[173554]Hum. Genet. 31: 373-379, 1968.

[173555]

[173556]

[173557]17726.

[173558]Azevedo, E. S.; Yoshida, A.: Brazilian variant of glucose-
6-phosphate

[173559]dehydrogenase (GD Minas Gerais). Nature 222: 380-382,
1969.

[173560]

[173561]

[173562]17727.

[173563]Babalola, A. O. G.; Beetlestone, J. G.; Luzzatto, L.: Genetic

[173564]variants of human erythrocyte glucose-6-phosphate de-
hydrogenase: kinetic

[173565]and thermodynamic parameters of variants A, B, and A- in

relation

[173566]to quaternary structure. J. Biol. Chem. 251: 2993–3002,
1976.

[173567]

[173568]

[173569]17728.

[173570]Balinsky, D.; Cayanis, E.; Carter, G.; Jenkins, T.; Bersohn, I.

[173571]: A new variant of human erythrocyte glucose–
6–phosphate dehydrogenase:

[173572]G6PD Port Elizabeth. Int. J. Biochem. 4: 235–244, 1973.

[173573]

[173574]

[173575]17729.

[173576]Balinsky, D.; Gomperts, E.; Cayanis, E.; Jenkins, T.; Bryer,
D.;

[173577]Bersohn, I.; Metz, J.: Glucose 6–phosphate dehydrogenase
Johannesburg:

[173578]a new variant with reduced activity in a patient with con–
genital non–spherocytic

[173579]haemolytic anaemia. Brit. J. Haemat. 25: 385–392, 1973.

[173580]

[173581]

[173582]17730.

[173583]Balinsky, D.; Rootman, A. J.; Nurse, G. T.; Cayanis, E.;
Lane,

[173584]A.; Jenkins, T.; Bersohn, I.: G6PD Kuanyama: a new variant
of human

[173585]erythrocyte glucose 6-phosphate dehydrogenase showing
slower than

[173586]normal electrophoretic mobility. S. Afr. J. Med. Sci. 39:
5-13,

[173587]1974.

[173588]

[173589]

[173590]17731.

[173591]Barretto, O. C.; Nonoyama, K.: Gd(+) Cuiaba, a new rare
glucose-6-phosphate

[173592]dehydrogenase variant presenting normal activity. Hum.
Genet. 77:

[173593]201-202, 1987.

[173594]

[173595]

[173596]17732.

[173597]Kornak, U.; Kasper, D.; Bosl, M. R.; Kaiser, E.; Schweizer, M.;

[173598]Schulz, A.; Friedrich, W.; Delling, G.; Jentsch, T. J.: Loss of the

[173599]ClC-7 chloride channel leads to osteopetrosis in mice and man. Cell 104:

[173600]205–215, 2001.

[173601]

[173602]

[173603]17733.

[173604]Kornak, U.; Schulz, A.; Friedrich, W.; Uhlhaas, S.; Kremens, B.;

[173605]Voit, T.; Hasan, C.; Bode, U.; Jentsch, T. J.; Kubisch, C.: Mutations

[173606]in the $\alpha 3$ subunit of the vacuolar H(+)-ATPase cause infantile malignant

[173607]osteopetrosis. Hum. Molec. Genet. 9: 2059–2063, 2000.

[173608]

[173609]

[173610]17734.

[173611]Rosenberg, C. L.; Wong, E.; Petty, E. M.; Bale, A. E.; Tsujimoto,
moto,

[173612]Y.; Harris, N. L.; Arnold, A.: PRAD1, a candidate BCL1
oncogene:

[173613]mapping and expression in centrocytic lymphoma. Proc.
Nat. Acad.

[173614]Sci. 88: 9638–9642, 1991.

[173615]

[173616]

[173617]17735.

[173618]Tsujimoto, Y.; Jaffe, E.; Cossman, J.; Gorham, J.; Nowell, P.

[173619]C.; Croce, C. M.: Clustering of breakpoints on chromo-
some 11 in human

[173620]B-cell neoplasms with the t(11;14) chromosome translo-
cation. Nature 315:

[173621]340–343, 1985.

[173622]

[173623]

[173624]17736.

[173625]Tsujiimoto, Y.; Yunis, J.; Onorato-Showe, L.; Erikson, J.;
Nowell,

[173626]P. C.; Croce, C. M.: Molecular cloning of the chromosomal
breakpoint

[173627]of B-cell lymphomas and leukemias with the t(11;14)
chromosome translocation. Science 224:

[173628]1403–1406, 1984.

[173629]

[173630]

[173631]17737.

[173632]Barbany, G.; Hoglund, M.; Simonsson, B.: Complete
molecular remission

[173633]in chronic myelogenous leukemia after imatinib therapy.
(Letter) New

[173634]Eng. J. Med. 347: 539–540, 2002.

[173635]

[173636]

[173637]17738.

[173638]Budarf, M.; Canaani, E.; Emanuel, B. S.: Linear order of the
four

[173639]BCR-related loci in 22q11. Genomics 3: 168–171, 1988.

[173640]

[173641]

[173642]17739.

[173643]Castellanos, A.; Pintado, B.; Weruaga, E.; Arevalo, R.;
Lopez,

[173644]A.; Orfao, A.; Sanchez-Garcia, I.: A BCR-ABL(p190) fusion
gene made

[173645]by homologous recombination causes B-cell acute lym-
phoblastic leukemias

[173646]in chimeric mice with independence of the endogenous
bcr product. Blood 90:

[173647]2168–2174, 1997. 1. Brunning, R. D.: Philadelphia chro-
mosome positive

[173648]leukemia. Hum. Path. 11: 307–309, 1980.

[173649]

[173650]

[173651]17740.

[173652]Chissoe, S. L.; Bodenteich, A.; Wang, Y.-F.; Wang, Y.-P.;
Burian,

[173653]D.; Clifton, S. W.; Crabtree, J.; Freeman, A.; Iyer, K.; Jian,
L.;

[173654]Ma, Y.; McLaury, H.-J.; Pan, H.-Q.; Sarhan, O. H.; Toth, S.;
Wang,

[173655]Z.; Zhang, G.; Heisterkamp, N.; Groffen, J.; Roe, B. A.: Se-
quence

[173656]and analysis of the human ABL gene, the BCR gene, and
regions involved

[173657]in the Philadelphia chromosomal translocation. Genomics
27: 67-82,

[173658]1995.

[173659]

[173660]

[173661]17741.

[173662]Court Brown, W. M.; Doll, R.: Mortality from cancer and
other

[173663]causes after radiotherapy for ankylosing spondylitis. Brit.
Med.

[173664]J. 2: 1327-1332, 1965.

[173665]

[173666]

[173667]17742.

[173668]Croce, C. M.; Huebner, K.; Isobe, M.; Fainstain, E.; Lifshitz,

[173669]B.; Shtivelman, E.; Canaani, E.: Mapping of four distinct
BCR-related

[173670]loci to chromosome region 22q11: order of BCR loci rela-
tive to chronic

[173671]myelogenous leukemia and acute lymphoblastic leukemia
breakpoints. Proc.

[173672]Nat. Acad. Sci. 84: 7174-7178, 1987.

[173673]

[173674]

[173675]17743.

[173676]Daley, G. Q.; Van Etten, R. A.; Baltimore, D.: Induction of
chronic

[173677]myelogenous leukemia in mice by the P210(bcr/abl) gene
of the Philadelphia

[173678]chromosome. Science 247: 824-830, 1990.

[173679]

[173680]

[173681]17744.

[173682]de Klein, A.; Geurts van Kessel, A.; Grosveld, G.; Bartram, C.

[173683]R.; Hagemeijer, A.; Bootsma, D.; Spurr, N. K.; Heisterkamp, N.; Groffen,

[173684]J.; Stephenson, J. R.: A cellular oncogene is translocated to the

[173685]Philadelphia chromosome in chronic myelocytic leukaemia. Nature 300:

[173686]765–767, 1982.

[173687]

[173688]

[173689]17745.

[173690]Diekmann, D.; Brill, S.; Garrett, M. D.; Totty, N.; Hsuan, J.;

[173691]Monfries, C.; Hall, C.; Lim, L.; Hall, A.: Bcr encodes a GT-Pase-activating

[173692]protein for p21(rac). Nature 351: 400–402, 1991.

[173693]

[173694]

[173695]17746.

[173696]Druker, B. J.; Sawyers, C. L.; Kantarjian, H.; Resta, D. J.;
Reese,

[173697]S. F.; Ford, J. M.; Capdeville, R.; Talpaz, M.: Activity of a
specific

[173698]inhibitor of the BCR–ABL tyrosine kinase in the blast crisis
of chronic

[173699]myeloid leukemia and acute lymphoblastic leukemia with
the Philadelphia

[173700]chromosome. New Eng. J. Med. 344: 1038–1042, 2001.

[173701]

[173702]

[173703]17747.

[173704]Druker, B. J.; Talpaz, M.; Resta, D. J.; Peng, B.; Buch-
dunger,

[173705]E.; Ford, J. M.; Lydon, N. B.; Kantarjian, H.; Capdeville, R.;
Ohno–Jones,

[173706]S.; Sawyers, C. L.: Efficacy and safety of a specific inhibitor
of

[173707]the BCR–ABL tyrosine kinase in chronic myeloid leukemia.
New Eng.

[173708]J. Med. 344: 1031–1037, 2001.

[173709]

[173710]

[173711]17748.

[173712]Emanuel, B. S.; Selden, J. R.; Wang, E.; Nowell, P. C.; Croce,

[173713]C. M.: In situ hybridization and translocation breakpoint mapping.

[173714]I. Non-identical 22q11 breakpoints for the t(9;22) of CML and the

[173715]t(8;22) of Burkitt lymphoma. Cytogenet. Cell Genet. 38: 127–131,

[173716]1984.

[173717]

[173718]

[173719]17749.

[173720]Grandori, C.; Mac, J.; Siebelt, F.; Ayer, D. E.; Eisenman, R. N.

[173721]: Myc–Max heterodimers activate a DEAD box gene and interact with

[173722]multiple E box-related sites in vivo. EMBO J. 15:

4344–4357, 1996.

[173723]

[173724]

[173725]17750.

[173726]Gallagher, A. R.; Cedzich, A.; Gretz, N.; Somlo, S.; Witzgall,

[173727]R.: The polycystic kidney disease protein PKD2 interacts
with Hax-1,

[173728]a protein associated with the actin cytoskeleton. Proc. Nat.
Acad.

[173729]Sci. 97: 4017–4022, 2000.

[173730]

[173731]

[173732]17751.

[173733]Cleutjens, K. B. J. M.; van Eekelen, C. C. E. M.; van der Ko-
rput,

[173734]H. A. G. M.; Brinkmann, A. O.; Trapman, J.: Two androgen
response

[173735]regions cooperate in steroid hormone regulated activity of
the prostate-specific

[173736]antigen promoter. J. Biol. Chem. 271: 6379–6388, 1996.

[173737]

[173738]

[173739]17752.

[173740]Evans, B. A.; Drinkwater, C. C.; Richards, R. I.: Mouse
glandular

[173741]kallikrein genes: structure and partial sequence analysis of
the kallikrein

[173742]gene locus. J. Biol. Chem. 262: 8027–8034, 1987.

[173743]

[173744]

[173745]17753.

[173746]Lundwall, A.; Lilja, H.: Molecular cloning of human
prostate specific

[173747]antigen cDNA. FEBS Lett. 214: 317–322, 1987.

[173748]

[173749]

[173750]17754.

[173751]Melegos, D. N.; Yu, H.; Ashok, M.; Wang, C.; Stanczyk, F.;
Diamandis,

[173752]E. P.: Prostate-specific antigen in female serum, a potential new

[173753]marker of androgen excess. J. Clin. Endocr. Metab. 82: 777-780,

[173754]1997.

[173755]

[173756]

[173757]17755.

[173758]Prendergast, G. C.; Lawe, D.; Ziff, E. B.: Association of Myn,

[173759]the murine homolog of Max, with c-Myc stimulates methylation-sensitive

[173760]DNA binding and Ras cotransformation. Cell 65: 395-408, 1991.

[173761]

[173762]

[173763]17756.

[173764]Wagner, A. J.; Le Beau, M. M.; Diaz, M. O.; Hay, N.: Expression,

[173765]regulation, and chromosomal localization of the Max gene. Proc. Nat.

[173766]Acad. Sci. 89: 3111–3115, 1992.

[173767]

[173768]

[173769]17757.

[173770]Zervos, A. S.; Faccio, L.; Gatto, J. P.; Kyriakis, J. M.; Brent,

[173771]R.: Mxi2, a mitogen-activated protein kinase that recognizes and

[173772]phosphorylates Max protein. Proc. Nat. Acad. Sci. 92: 10531–10534,

[173773]1995.

[173774]

[173775]

[173776]17758.

[173777]Emi, M.; Katagiri, T.; Harada, Y.; Saito, H.; Inazawa, J.; Ito,

[173778]I.; Kasumi, F.; Nakamura, Y.: A novel metalloprotease/disintegrin-like

[173779]gene at 17q21.3 is somatically rearranged in two primary breast cancers. Nature

[173780]Genet. 5: 151–157, 1993.

[173781]

[173782]

[173783]17759.

[173784]Katagiri, T.; Harada, Y.; Emi, M.; Nakamura, Y.: Human metalloprotease/disintegrin-like

[173785](MDC) gene: exon-intron organization and alternative splicing. Cytogenet.

[173786]Cell Genet. 68: 39-44, 1995.

[173787]

[173788]

[173789]17760.

[173790]Begley, C. G.; Lipkowitz, S.; Gobel, V.; Mahon, K. A.; Bertness,

[173791]V.; Green, A. R.; Gough, N. M.; Kirsch, I. R.: Molecular characterization

[173792]of NSCL, a gene encoding a helix-loop-helix protein expressed in the

[173793]developing nervous system. Proc. Nat. Acad. Sci. 89: 38-42, 1992.

[173794]

[173795]

[173796]17761.

[173797]Brown, L.; Espinosa, R., III; Le Beau, M. M.; Siciliano, M. J.;

[173798]Baer, R.: HEN1 and HEN2: a subgroup of basic helix–
loop–helix genes

[173799]that are coexpressed in a human neuroblastoma. Proc.
Nat. Acad. Sci. 89:

[173800]8492–8496, 1992.

[173801]

[173802]

[173803]17762.

[173804]Cogliati, T.; Good, D. J.; Haigney, M.; Delgado–Romero, P.;
Eckhaus,

[173805]M. A.; Koch, W. J.; Kirsch, I. R.: Predisposition to arrhyth–
mia and

[173806]autonomic dysfunction in Nhlh1–deficient mice. Molec.
Cell. Biol. 22:

[173807]4977–4983, 2002.

[173808]

[173809]

[173810]17763.

[173811]Lipkowitz, S.; Gobel, V.; Varterasian, M. L.; Nakahara, K.;
Tchorz,

[173812]K.; Kirsch, I. R.: A comparative structural characterization
of the

[173813]human NSCL-1 and NSCL-2 genes: two basic helix-
loop-helix genes expressed

[173814]in the developing nervous system. J. Biol. Chem. 267:
21065-21071,

[173815]1992.

[173816]

[173817]

[173818]17764.

[173819]Mullick, A.; Groulx, N.; Trasler, D.; Gros, P.: Nhlh1, a basic

[173820]helix-loop-helix transcription factor, is very tightly linked
to the

[173821]mouse looptail (Lp) mutation. Mammalian Genome 6:
700-704, 1995.

[173822]

[173823]

[173824]17765.

[173825]Meeker, T. C.; Nagarajan, L.; ar-Rushdi, A.; Rovera, G.;

Huebner,

[173826]K.; Croce, C. M.: Characterization of the human PIM-1

gene: a putative

[173827]proto-oncogene coding for a tissue specific member of

the protein

[173828]kinase family. Oncogene Res. 1: 87-101, 1987.

[173829]

[173830]

[173831]17766.

[173832]Pasqualucci, L.; Neumeister, P.; Goossens, T.; Nanjangud,

G.; Chaganti,

[173833]R. S. K.; Kuppers, R.; Dalla-Favera, R.: Hypermutation of

multiple

[173834]proto-oncogenes in B-cell diffuse large-cell lymphomas.

Nature 412:

[173835]341-346, 2001.

[173836]

[173837]

[173838]17767.

[173839]Ragoussis, J.; Senger, G.; Mockridge, I.; Sanseau, P.;
Ruddy, S.;

[173840]Dudley, K.; Sheer, D.; Trowsdale, J.: A testis-expressed Zn
finger

[173841]gene (ZNF76) in human 6p21.3 centromeric to the MHC is
closely linked

[173842]to the human homolog of the t-complex gene tcp-11.

Genomics 14:

[173843]673-679, 1992.

[173844]

[173845]

[173846]17768.

[173847]Saris, C. J. M.; Domen, J.; Berns, A.: The pim-1 oncogene
encodes

[173848]two related protein-serine/threonine kinases by alterna-
tive initiation

[173849]at AUG and CUG. EMBO J. 10: 655-664, 1991.

[173850]

[173851]

[173852]17769.

[173853]Selten, G.; Cuypers, H. T.; Boelens, W.; Robanus-Maandag, E.;

[173854]Verbeek, J.; Domen, J.; van Beveren, C.; Berns, A.: The primary structure

[173855]of the putative oncogene pim-1 shows extensive homology with protein

[173856]kinases. Cell 46: 603-611, 1986.

[173857]

[173858]

[173859]17770.

[173860]Zoghbi, H. Y.; Ballantyne, C. M.; O'Brien, W. E.; McCall, A. E.;

[173861]Kwiatkowski, T. J., Jr.; Ledbetter, S. A.; Beaudet, A. L.: Deletion

[173862]and linkage mapping of eight markers from the proximal short arm of

[173863]chromosome 6. Genomics 6: 352-357, 1990.

[173864]

[173865]

[173866]17771.

[173867]Chapin, S. J.; Lue, C.-M.; Yu, M. T.; Bulinski, J. C.: Differential
ential

[173868]expression of alternatively spliced forms of MAP4: a
repertoire of

[173869]structurally different microtubule-binding domains. Bio-
chemistry 34:

[173870]2289-2301, 1995.

[173871]

[173872]

[173873]17772.

[173874]Laird, P. W.; van der Lugt, N. M. T.; Clarke, A.; Domen, J.;
Linders,

[173875]K.; McWhir, J.; Berns, A.; Hooper, M.: In vivo analysis of
Pim-1

[173876]deficiency. Nucleic Acids Res. 21: 4750-4755, 1993.

[173877]

[173878]

[173879]17773.

[173880]Gartner, J.; Obie, C.; Watkins, P.; Valle, D.: Restoration of

[173881]peroxisome biogenesis in a peroxisome-deficient mammalian cell line

[173882]by expression of either the 35 kDa or the 70 kDa peroxisomal membrane

[173883]proteins. J. Inherit. Metab. Dis. 17: 327-329, 1994.

[173884]

[173885]

[173886]17774.

[173887]Maric, S. C.; Crozat, A.; Janne, O. A.: Structure and organization

[173888]of the human S-adenosylmethionine decarboxylase gene. J. Biol. Chem. 267:

[173889]18915-18923, 1992.

[173890]

[173891]

[173892]17775.

[173893]Maric, S. C.; Crozat, A.; Louhimo, J.; Knuutila, S.; Janne, O.

[173894]A.: The human S-adenosylmethionine decarboxylase gene: nucleotide

[173895]sequence of a pseudogene and chromosomal localization

of the active

[173896]gene (AMD1) and the pseudogene (AMD2). Cytogenet. Cell
Genet. 70:

[173897]195–199, 1995.

[173898]

[173899]

[173900]17776.

[173901]Mangan, M. E.; Olmsted, J. B.: The gene for microtubule-as-
sociated

[173902]protein 4 (Mtap4) maps to the distal region of mouse
chromosome 9. Mammalian

[173903]Genome 7: 918–925, 1996.

[173904]

[173905]

[173906]17777.

[173907]West, R. R.; Tenbarger, K. M.; Olmsted, J. B.: A model for
microtubule-associated

[173908]protein 4 structure: domains defined by comparisons of
human, mouse,

[173909]and bovine sequences. J. Biol. Chem. 266: 21886–21896,

1991.

[173910]

[173911]

[173912]17778.

[173913]Abel, K. J.; Boehnke, M.; Prahalad, M.; Ho, P.; Flejter, W. L.;

[173914]Watkins, M.; VanderStoep, J.; Chandrasekharappa, S. C.;
Collins, F.

[173915]S.; Glover, T. W.; Weber, B. L.: A radiation hybrid map of
the BRCA1

[173916]region of chromosome 17q12–q21. Genomics 17:
632–641, 1993.

[173917]

[173918]

[173919]17779.

[173920]Alonso, A. D. C.; Grundke-Iqbal, I.; Iqbal, K.: Alzheimer's
disease

[173921]hyperphosphorylated tau sequesters normal tau into tan-
gles of filaments

[173922]and disassembles microtubules. Nature Med. 2: 783–787,
1996.

[173923]

[173924]

[173925]17780.

[173926]Arima, K.; Kowalska, A.; Hasegawa, M.; Mukoyama, M.;
Watanabe,

[173927]R.; Kawai, M.; Takahashi, K.; Iwatsubo, T.; Tabira, T.;
Sunohara,

[173928]N.: Two brothers with frontotemporal dementia and
parkinsonism with

[173929]an N279K mutation of the tau gene. Neurology 54:
1787–1795, 2000.

[173930]

[173931]

[173932]17781.

[173933]Baker, M.; Litvan, I.; Houlden, H.; Adamson, J.; Dickson,
D.; Perez-Tur,

[173934]J.; Hardy, J.; Lynch, T.; Bigio, E.; Hutton, M.: Association of
an

[173935]extended haplotype in the tau gene with progressive
supranuclear palsy. Hum.

[173936]Molec. Genet. 8: 711–715, 1999.

[173937]

[173938]

[173939]17782.

[173940]Brown, J.; Lantos, P. L.; Roques, P.; Fidani, L.; Rossor, M.
N.

[173941]: Familial dementia with swollen achromatic neurons and
corticobasal

[173942]inclusion bodies: a clinical and pathological study. J. Neu-
rol. Sci. 135:

[173943]21–30, 1996.

[173944]

[173945]

[173946]17783.

[173947]Clark, L. N.; Poorkaj, P.; Wszolek, Z.; Geschwind, D. H.;
Nasreddine,

[173948]Z. S.; Miller, B.; Li, D.; Payami, H.; Awert, F.; Markopoulou,
K.;

[173949]Andreadis, A.; D'Souza, I.; Lee, V. M.-Y.; Reed, L.; Tro-
janowski,

[173950]J. Q.; Zhukareva, V.; Bird, T.; Schellenberg, G.; Wilhelmsen,
K. C.

[173951]: Pathogenic implications of mutations in the tau gene in

pallido-ponto-nigral

[173952]degeneration and related neurodegenerative disorders
linked to chromosome

[173953]17. Proc. Nat. Acad. Sci. 95: 13103–13107, 1998.

[173954]

[173955]

[173956]17784.

[173957]Connell, J. W.; Gibb, G. M.; Betts, J. C.; Blackstock, W. P.;
Gallo,

[173958]J.–M.; Lovestone, S.; Hutton, M.; Anderton, B. H.: Effects of
FTDP–17

[173959]mutations on the in vitro phosphorylation of tau by glyco-
gen synthase

[173960]kinase 3–beta identified by mass spectrometry demon-
strate certain

[173961]mutations exert long–range conformational changes. FEBS
Lett. 493:

[173962]40–44, 2001.

[173963]

[173964]

[173965]17785.

[173966]Conrad, C.; Andreadis, A.; Trojanowski, J. Q.; Dickson, D.
W.; Kang,

[173967]D.; Chen, X.; Weiderholt, W.; Hansen, L.; Masliah, E.; Thal,
L. J.;

[173968]Katzman, R.; Xia, Y.; Saitoh, T.: Genetic evidence for the
involvement

[173969]of tau in progressive supranuclear palsy. Ann. Neurol. 41:
277–281,

[173970]1997.

[173971]

[173972]

[173973]17786.

[173974]Conrad, C.; Vianna, C.; Freeman, M.; Davies, P.: A poly-
morphic

[173975]gene nested within an intron of the tau gene: implications
for Alzheimer's

[173976]disease. Proc. Nat. Acad. Sci. 99: 7751–7756, 2002.

[173977]

[173978]

[173979]17787.

[173980]Delisle, M. B.; Murrell, J. R.; Richardson, R.; Trofatter, J.

[173981]A.; Rascol, O.; Soulages, X.; Mohr, M.; Calvas, P.; Ghetti,
B.: A

[173982]mutation at codon 279 (N279K) in exon 10 of the tau
gene causes a

[173983]tauopathy with dementia and supranuclear palsy. Acta
Neuropath. 98:

[173984]62–77, 1999.

[173985]

[173986]

[173987]17788.

[173988]Donlon, T. A.; Harris, P.; Neve, R. L.: Localization of mi-
crotubule-associated

[173989]protein tau (MTBT1) to chromosome 17q21. (Abstract) Cy-
togenet. Cell

[173990]Genet. 46: 607, 1987.

[173991]

[173992]

[173993]17789.

[173994]Goedert, M.; Crowther, R. A.; Spillantini, M. G.: Tau muta-

tions

[173995]cause frontotemporal dementias. Neuron 21: 955–958, 1998.

[173996]

[173997]

[173998]17790.

[173999]Goedert, M.; Spillantini, M. G.; Potier, M. C.; Ulrich, J.; Crowther,

[174000]R. A.: Cloning and sequencing of the cDNA encoding an isoform of

[174001]microtubule-associated protein tau containing four tandem repeats:

[174002]differential expression of tau protein mRNAs in human brain. EMBO

[174003]J. 8: 393–399, 1989.

[174004]

[174005]

[174006]17791.

[174007]Goedert, M.; Wischik, C. M.; Crowther, R. A.; Walker, J. E.; Klug,

[174008]A.: Cloning and sequencing of the cDNA encoding a core

protein of

[174009]the paired helical filament of Alzheimer disease: identification as

[174010]the microtubule-associated protein tau. Proc. Nat. Acad. Sci. 85:

[174011]4051–4055, 1988.

[174012]

[174013]

[174014]17792.

[174015]Goode, B. L.; Chau, M.; Denis, P. E.; Feinstein, S. C.: Structural

[174016]and functional differences between 3-repeat and 4-repeat tau isoforms:

[174017]implications for normal tau function and the onset of neurodegenerative

[174018]disease. J. Biol. Chem. 275: 38182–38189, 2000.

[174019]

[174020]

[174021]17793.

[174022]Heutink, P.: Untangling tau-related dementia. Hum.

Molec. Genet. 9:

[174023]979–986, 2000.

[174024]

[174025]

[174026]17794.

[174027]Heutink, P.; Stevens, M.; Rizzu, P.; Bakker, E.; Kros, J. M.;

[174028]Tibben, A.; Niermeijer, M. F.; van Duijn, C. M.; Oostra, B.

A.; van

[174029]Swieten, J. C.: Hereditary frontotemporal dementia is
linked to chromosome

[174030]17q21–q22: a genetic and clinicopathological study of
three Dutch

[174031]families. Ann. Neurol. 41: 150–159, 1997.

[174032]

[174033]

[174034]17795.

[174035]Hiesberger, T.; Trommsdorff, M.; Howell, B. W.; Goffinet,
A.;

[174036]Mumby, M. C.; Cooper, J. A.; Herz, J.: Direct binding of
reelin to

[174037]VLDL receptor and apoE receptor 2 induces tyrosine phosphorylation

[174038]of disabled-1 and modulates tau phosphorylation. Neuron
24: 481-489,

[174039]1999.

[174040]

[174041]

[174042]17796.

[174043]Hong, M.; Zhukareva, V.; Vogelsberg-Ragaglia, V.; Ws-
zolek, Z.;

[174044]Reed, L.; Miller, B. I.; Geschwind, D. H.; Bird, T. D.; McKeel,
D.;

[174045]Goate, A.; Morris, J. C.; Wilhelmsen, K. C.; Schellenberg, G.
D.;

[174046]Trojanowski, J. Q.; Lee, V. M.-Y.: Mutation-specific func-
tional impairments

[174047]in distinct tau isoforms of hereditary FTDP-17. Science
282: 1914-1917,

[174048]1998.

[174049]

[174050]

[174051]17797.

[174052]Hutton, M.: Missense and splice site mutations in tau associated

[174053]with FTDP-17: multiple pathogenic mechanisms. Neurology 56 (suppl.

[174054]4): S21-S25, 2001.

[174055]

[174056]

[174057]17798.

[174058]Hutton, M.; Lendon, C. L.; Rizzu, P.; Baker, M.; Froelich, S.;

[174059]Houlden, H.; Pickering-Brown, S.; Chakraverty, S.; Isaacs, A.; Grover,

[174060]A.; Hackett, J.; Adamson, J.; and 39 others: Association of missense

[174061]and 5-prime-splice-site mutations in tau with the inherited dementia

[174062]FTDP-17. Nature 393: 702-705, 1998.

[174063]

[174064]

[174065]17799.

[174066]Iijima, M.; Tabira, T.; Poorkaj, P.; Schellenberg, G. D.; Trojanowski,

[174067]J. Q.; Lee, V. M.; Schmidt, M. L.; Takahashi, K.; Nabika, T.; Matsumoto,

[174068]T.; Yamashita, Y.; Yoshioka, S.; Ishino, H.: A distinct familial

[174069]presenile dementia with a novel missense mutation in the tau gene. Neuroreport 10:

[174070]497–501, 1999.

[174071]

[174072]

[174073]17800.

[174074]Ishihara, T.; Hong, M.; Zhang, B.; Nakagawa, Y.; Lee, M. K.; Trojanowski,

[174075]J. Q.; Lee, V. M.-Y.: Age-dependent emergence and progression of

[174076]a tauopathy in transgenic mice overexpressing the shortest human tau

[174077]isoform. Neuron 24: 751–762, 1999.

[174078]

[174079]

[174080]17801.

[174081]Janssen, J. C.; Warrington, E. K.; Morris, H. R.; Lantos, P.;

[174082]Brown, J.; Revesz, T.; Wood, N.; Khan, M. N.; Cipolotti, L.;

Fox,

[174083]N. C.; Rossor, M. N.: Clinical features of frontotemporal
dementia

[174084]due to the intronic tau 10 +16 mutation. *Neurology* 58:
1161–1168,

[174085]2002.

[174086]

[174087]

[174088]17802.

[174089]Bergstrom, D. A.; Penn, B. H.; Strand, A.; Perry, R. L. S.;
Rudnicki,

[174090]M. A.; Tapscott, S. J.: Promoter-specific regulation of
MyoD binding

[174091]and signal transduction cooperate to pattern gene ex-
pression. *Molec.*

[174092]*Cell* 9: 587–600, 2002.

[174093]

[174094]

[174095]17803.

[174096]Braun, T.; Grzeschik, K.-H.; Bober, E.; Arnold, H.-H.: The
MYF

[174097]genes, a group of human muscle determining factors, are
localized

[174098]on different human chromosomes. (Abstract) Cytogenet.
Cell Genet. 51:

[174099]969 only, 1989.

[174100]

[174101]

[174102]17804.

[174103]Davis, R. L.; Weintraub, H.; Lassar, A. B.: Expression of a
single

[174104]transfected cDNA converts fibroblasts to myoblasts. Cell
51: 987-1000,

[174105]1987.

[174106]

[174107]

[174108]17805.

[174109]de la Serna, I. L.; Carlson, K. A.; Imbalzano, A. N.: Mam-
malian

[174110]SWI/SNF complexes promote MyoD-mediated muscle dif-
ferentiation. Nature

[174111]Genet. 27: 187-190, 2001.

[174112]

[174113]

[174114]17806.

[174115]Gessler, M.; Hameister, H.; Henry, I.; Junien, C.; Braun, T.;
Arnold,

[174116]H. H.: The human MyoD1 (MYF3) gene maps on the short
arm of chromosome

[174117]11 but is not associated with the WAGR locus or the re-
gion for the

[174118]Beckwith-Wiedemann syndrome. Hum. Genet. 86:
135-138, 1990.

[174119]

[174120]

[174121]17807.

[174122]Guttridge, D. C.; Mayo, M. W.; Madrid, L. V.; Wang, C.-Y.;
Baldwin,

[174123]A. S, Jr.: NF-kappa-B-induced loss of MyoD messenger
RNA: possible

[174124]role in muscle decay and cachexia. Science 289:
2363-2366, 2000.

[174125]

[174126]

[174127]17808.

[174128]Henry, I.; Puech, A.; Antignac, C.; Couillin, P.; Jeanpierre,
M.;

[174129]Ahnine, L.; Barichard, F.; Boehm, T.; Augereau, P.; Scrable,
H.; Rabbitts,

[174130]T. H.; Rochefort, H.; Cavenee, W.; Junien, C.: Subregional
mapping

[174131]of BWS, CTSD, MYOD1, and a T-ALL breakpoint in 11p15.
(Abstract) Cytogenet.

[174132]Cell Genet. 51: 1013 only, 1989.

[174133]

[174134]

[174135]17809.

[174136]Kim, Y.-J.; Noguchi, S.; Hayashi, Y. K.; Tsukahara, T.;
Shimizu,

[174137]T.; Arahata, K.: The product of an oculopharyngeal muscular dystrophy

[174138]gene, poly(A)-binding protein 2, interacts with SKIP and stimulates

[174139]muscle-specific gene expression. Hum. Molec. Genet. 10: 1129-1139,

[174140]2001.

[174141]

[174142]

[174143]17810.

[174144]Olson, E. N.: MyoD family: a paradigm for development. Genes

[174145]Dev. 4: 1454-1461, 1990.

[174146]

[174147]

[174148]17811.

[174149]Sartorelli, V.; Puri, P. L.; Hamamori, Y.; Ogryzko, V.; Chung,

[174150]G.; Nakatani, Y.; Wang, J. Y. J.; Kedes, L.: Acetylation of MyoD

[174151]directed by PCAF is necessary for the execution of the muscle program. Molec.

[174152]Cell 4: 725–734, 1999.

[174153]

[174154]

[174155]17812.

[174156]Tapscott, S. J.; Davis, R. L.; Thayer, M. J.; Cheng, P. F.; Weintraub,

[174157]H.; Lassar, A. B.: MyoD1: a nuclear phosphoprotein requiring a Myc

[174158]homology region to convert fibroblasts to myoblasts. Science 242:

[174159]405–411, 1988.

[174160]

[174161]

[174162]17813.

[174163]Bonner, T. I.; Young, A. C.; de Miguel, C.; Detera-Wadleigh, S.;

[174164]Modi, W. S.; O'Brien, S. J.: The chromosomal location of the two

[174165]human tackykinin (sic) genes: human substance P and neurokinin K.

[174166](Abstract) Cytogenet. Cell Genet. 46: 584 only, 1987.

[174167]

[174168]

[174169]17814.

[174170]Cao, Y. Q.; Mantyh, P. W.; Carlson, E. J.; Gillespie, A.-M.; Epstein,

[174171]C. J.; Basbaum, A. I.: Primary afferent tachykinins are required

[174172]to experience moderate to intense pain. Nature 392: 390–393, 1998.

[174173]

[174174]

[174175]17815.

[174176]Krause, J. E.; Chirgwin, J. M.; Carter, M. S.; Xu, Z. S.; Hershey,

[174177]A. D.: Three rat preprotachykinin mRNAs encode the neuropeptides

[174178]substance P and neurokinin A. Proc. Nat. Acad. Sci. 84: 881–885,

[174179]1987.

[174180]

[174181]

[174182]17816.

[174183]Liu, H.; Cao, Y.; Basbaum, A. I.; Mazarati, A. M; Sankar, R.;
Wasterlain,

[174184]C. G.: Resistance to excitotoxin-induced seizures and
neuronal death

[174185]in mice lacking the preprotachykinin A gene. Proc. Nat.
Acad. Sci. 96:

[174186]12096–12101, 1999.

[174187]

[174188]

[174189]17817.

[174190]Troger, J.; Neyer, S.; Heufler, C.; Huemer, H.; Schmid, E.;
Griesser,

[174191]U.; Kralinger, M.; Kremser, B.; Baldissera, I.; Kieselbach,
G.: Substance

[174192]P and vasoactive intestinal polypeptide in the streptozotocin-induced

[174193]diabetic rat retina. Invest. Ophthal. Vis. Sci. 42:
1045–1050, 2001.

[174194]

[174195]

[174196]17818.

[174197]Zimmer, A.; Zimmer, A. M.; Baffi, J.; Usdin, T.; Reynolds,
K.;

[174198]Konig, M.; Palkovits, M.; Mezey, E.: Hypoalgesia in mice
with a targeted

[174199]deletion of the tachykinin 1 gene. Proc. Nat. Acad. Sci. 95:
2630–2635,

[174200]1998.

[174201]

[174202]

[174203]17819.

[174204]Gregory, C. A.; Schwartz, J. S.: The cDNA of the human
neuromedin

[174205]B gene (NMB) mapped to 15q11–qter recognizes an XbaI
RFLP. Nucleic

[174206]Acids Res. 19: 1167 only, 1991.

[174207]

[174208]

[174209]17820.

[174210]Krane, I. M.; Naylor, S. L.; Helin-Davis, D.; Chin, W. W.;
Spindel,

[174211]E. R.: Molecular cloning of cDNAs encoding the human
bombesin-like

[174212]peptide neuromedin B: chromosomal localization and
comparison to cDNAs

[174213]encoding its amphibian homolog ranatensin. J. Biol. Chem.
263:

[174214]13317-13323, 1988. Note: Erratum: J. Biol. Chem. 265:
7091 only, 1990.

[174215]

[174216]

[174217]17821.

[174218]Mattei, M.-G.; Riviere, M.; Krust, A.; Ingvarsson, S.;
Vennstrom,

[174219]B.; Islam, M. Q.; Levan, G.; Kautner, P.; Zelent, A.; Cham-
bon, P.;

[174220]Szpirer, J.; Szpirer, C.: Chromosomal assignment of

retinoic acid

[174221]receptor (RAR) genes in the human, mouse, and rat
genomes. Genomics 10:

[174222]1061–1069, 1991.

[174223]

[174224]

[174225]17822.

[174226]Abramson, D. H.; Ellsworth, R. M.; Zimmerman, L. E.:
Monocular

[174227]cancer in retinoblastoma survivors. Trans. Am. Acad.
Ophthalmol. Otolaryng. 81:

[174228]454–457, 1976.

[174229]

[174230]

[174231]17823.

[174232]Aherne, G. E. S.; Roberts, D. F.: Retinoblastoma—a clinical
survey

[174233]and its genetic implications. Clin. Genet. 8: 275–290,
1975.

[174234]

[174235]

[174236]17824.

[174237]Alonso, J.; Garcia-Miguel, P.; Abelairas, J.; Mendiola, M.;
Sarret,

[174238]E.; Vendrell, M. T.; Navajas, A.; Pestana, A.: Spectrum of
germline

[174239]RB1 gene mutations in Spanish retinoblastoma patients:
phenotypic

[174240]and molecular epidemiological implications. Hum. Mutat.
17: 412-422,

[174241]2001.

[174242]

[174243]

[174244]17825.

[174245]Amoaku, W. M. K.; Willshaw, H. E.; Parkes, S. E.; Shah, K. J.;

[174246]Mann, J. R.: Trilateral retinoblastoma: a report of five pa-
tients. Cancer 78:

[174247]858-863, 1996.

[174248]

[174249]

[174250]17826.

[174251]Bader, J. L.; Meadows, A. T.; Zimmerman, L. E.; Rorke, L.
B.; Voute,

[174252]P. A.; Champion, L. A. A.; Miller, R. W.: Bilateral
retinoblastoma

[174253]with ectopic intracranial retinoblastoma: trilateral
retinoblastoma. Cancer

[174254]Genet. Cytogenet. 5: 203–213, 1982.

[174255]

[174256]

[174257]17827.

[174258]Balaban–Malenbaum, G.; Gilbert, F.; Nichols, W. W.; Hill,
R.; Shields,

[174259]J.; Meadows, A. T.: A deleted chromosome no. 13 in hu-
man retinoblastoma

[174260]cells: relevance to tumorigenesis. Cancer Genet. Cyto-
genet. 3: 243–250,

[174261]1981.

[174262]

[174263]

[174264]17828.

[174265]Bandara, L. R.; Adamczewski, J. P.; Hunt, T.; La Thangue, N. B.

[174266]: Cyclin A and the retinoblastoma gene product complex with a common

[174267]transcription factor. Nature 352: 249–251, 1991.

[174268]

[174269]

[174270]17829.

[174271]Benedict, W. F.; Murphree, A. L.; Banerjee, A.; Spina, C. A.;
Sparkes,

[174272]M. C.; Sparkes, R. S.: Patient with 13 chromosome deletion: evidence

[174273]that the retinoblastoma gene is a recessive cancer gene.
Science 219:

[174274]973–975, 1983.

[174275]

[174276]

[174277]17830.

[174278]Benedict, W. F.; Xu, H.-J.; Hu, S.-X.; Takahashi, R.: Role of

[174279]the retinoblastoma gene in the initiation and progression
of human

[174280]cancer. J. Clin. Invest. 85: 988–993, 1990.

[174281]

[174282]

[174283]17831.

[174284]Bia, B.; Cowell, J. K.: Independent constitutional germline mutations

[174285]occurring in the RB1 gene in cousins with bilateral retinoblastoma. Oncogene 11:

[174286]977–979, 1995.

[174287]

[174288]

[174289]17832.

[174290]Blanquet, V.; Turleau, C.; de Grouchy, J.; Creau–Goldberg, N.

[174291]: Physical map around the retinoblastoma gene: possible genomic imprinting

[174292]suggested by Nrul digestion. Genomics 10: 350–355, 1991.

[174293]

[174294]

[174295]17833.

[174296]Blanquet, V.; Turleau, C.; Gross–Morand, M. S.; Sena–maud–Beaufort,

[174297]C.; Doz, F.; Besmond, C.: Spectrum of germline mutations
in the RB1

[174298]gene: a study of 232 patients with hereditary and non
hereditary retinoblastoma. Hum.

[174299]Molec. Genet. 4: 383–388, 1995.

[174300]

[174301]

[174302]17834.

[174303]Jiang, Z.; Cote, J.; Kwon, J. M.; Goate, A. M.; Wu, J. Y.:
Aberrant

[174304]splicing of tau pre-mRNA caused by intronic mutations
associated with

[174305]the inherited dementia frontotemporal dementia with
Parkinsonism linked

[174306]to chromosome 17. Molec. Cell. Biol. 20: 4036–4048,
2000.

[174307]

[174308]

[174309]17835.

[174310]Tapscott, S. J.; Weintraub, H.: MyoD and the regulation of
myogenesis

[174311]by helix-loop-helix proteins. J. Clin. Invest. 87:
1133-1138, 1991.

[174312]

[174313]

[174314]17836.

[174315]Futterweit, W.; Ritch, R.; Teekhasaene, C.; Nelson, E. S.:
Coexistence

[174316]of Prader-Willi syndrome, congenital ectropion uveae with
glaucoma,

[174317]and factor XI deficiency. J.A.M.A. 255: 3280-3282, 1986.

[174318]

[174319]

[174320]17837.

[174321]Trapani, J. A.; Browne, K. A.; Dawson, M. J.; Ramsay, R. G.;
Eddy,

[174322]R. L.; Shows, T. B.; White, P. C.; Dupont, B.: A novel gene
constitutively

[174323]expressed in human lymphoid cells is inducible with in-
terferon-gamma

[174324]in myeloid cells. Immunogenetics 36: 369-376, 1992.

[174325]

[174326]

[174327]17838.

[174328]Bowcock, A. M.; Kidd, J. R.; Lathrop, G. M.; Daneshvar, L.;
May,

[174329]L. T.; Ray, A.; Sehgal, P. B.; Kidd, K. K.; Cavalli-Sforza, L. L.

[174330]: The human 'interferon-beta-2/hepatocyte stimulating
factor/interleukin-6'

[174331]gene: DNA polymorphism studies and localization to
chromosome 7p21. Genomics 3:

[174332]8-16, 1988.

[174333]

[174334]

[174335]17839.

[174336]Chen, Y.; Ferguson-Smith, A. C.; Newman, M. S.; May, L.
T.; Sehgal,

[174337]P. B.; Ruddle, F. H.: Regional localization of the human
beta 2-interferon

[174338]gene. (Abstract) Am. J. Hum. Genet. 41: A161, 1987.

[174339]

[174340]

[174341]17840.

[174342]Chow, D.; He, X.; Snow, A. L.; Rose-John, S.; Garcia, K. C.:

Structure

[174343]of an extracellular gp130 cytokine receptor signaling

complex. Science 291:

[174344]2150–2155, 2001.

[174345]

[174346]

[174347]17841.

[174348]Chung, U.; Tanaka, Y.; Fujita, T.: Association of inter-

leukin-6

[174349]and hypoaldosteronism in patients with cancer. (Letter)

New Eng.

[174350]J. Med. 334: 473, 1996.

[174351]

[174352]

[174353]17842.

[174354]Cressman, D. E.; Greenbaum, L. E.; DeAngelis, R. A.; Cilib-

erto,

[174355]G.; Furth, E. E.; Poli, V.; Taub, R.: Liver failure and defective

[174356]hepatocyte regeneration in interleukin-6-deficient mice.

Science 274:

[174357]1379-1382, 1996.

[174358]

[174359]

[174360]17843.

[174361]De Benedetti, F.; Alonzi, T.; Moretta, A.; Lazzaro, D.;

Costa,

[174362]P.; Poli, V.; Martini, A.; Ciliberto, G.; Fattori, E.: Interleukin

[174363]6 causes growth impairment in transgenic mice through a decrease in

[174364]insulin-like growth factor-I. J. Clin. Invest. 99: 643-650, 1997.

[174365]

[174366]

[174367]17844.

[174368]Ferguson-Smith, A. C.; Chen, Y.-F.; Newman, M. S.; May,

L. T.;

[174369]Sehgal, P. B.; Ruddle, F. H.: Regional localization of the interferon

[174370]beta-2/B-cell stimulatory factor 2/hepatocyte stimulating factor gene

[174371]to human chromosome 7p15-p21. Genomics 2: 203-208, 1988.

[174372]

[174373]

[174374]17845.

[174375]Fernandez-Real, J.-M.; Broch, M.; Vendrell, J.; Richart, C.; Ricart,

[174376]W.: Interleukin-6 gene polymorphism and lipid abnormalities in healthy

[174377]subjects. J. Clin. Endocr. Metab. 85: 1334-1339, 2000.

[174378]

[174379]

[174380]17846.

[174381]Fishman, D.; Faulds, G.; Jeffery, R.; Mohamed-Ali, V.; Yudkin,

[174382]J. S.; Humphries, S.; Woo, P.: The effect of novel polymor-

phisms

[174383]in the interleukin-6 (IL-6) gene on IL-6 transcription and
plasma

[174384]IL-6 levels, and an association with systemic-onset juvenile chronic

[174385]arthritis. J. Clin. Invest. 102: 1369-1376, 1998.

[174386]

[174387]

[174388]17847.

[174389]Foster, C. B.; Lehrnbecher, T.; Samuels, S.; Stein, S.; Mol,
F.;

[174390]Metcalf, J. A.; Wyvill, K.; Steinberg, S. M.; Kovacs, J.; Blau-
velt,

[174391]A.; Yarchoan, R.; Chanock, S. J.: An IL6 promoter polymor-
phism is

[174392]associated with a lifetime risk of development of Kaposi
sarcoma in

[174393]men infected with human immunodeficiency virus. Blood
96: 2562-2567,

[174394]2000.

[174395]

[174396]

[174397]17848.

[174398]Funatsu, H.; Yamashita, H.; Noma, H.; Mimura, T.; Yamashita, T.;

[174399]Hori, S.: Increased levels of vascular endothelial growth factor

[174400]and interleukin-6 in the aqueous humor of diabetics with macular edema. Am.

[174401]J. Ophthal. 133: 70-77, 2002.

[174402]

[174403]

[174404]17849.

[174405]Hirano, T.; Yasukawa, K.; Harada, H.; Taga, T.; Watanabe, Y.;

[174406]Matsuda, T.; Kashiwamura, S.; Nakajima, K.; Koyama, K.; Iwamatsu,

[174407]A.; Tsunasawa, S.; Sakiyama, F.; Matsui, H.; Takahara, Y.; Taniguchi,

[174408]T.; Kishimoto, T.: Complementary DNA for a novel human interleukin

[174409](BSF-2) that induces B lymphocytes to produce im-

munoglobulin. Nature 324:

[174410]73–76, 1986.

[174411]

[174412]

[174413]17850.

[174414]Kawano, M.; Hirano, T.; Matsuda, T.; Taga, T.; Horii, Y.;
Iwato,

[174415]K.; Asaoku, H.; Tang, B.; Tanabe, O.; Tanaka, H.; Ku-
ramoto, A.; Kishimoto,

[174416]T.: Autocrine generation and requirement of BSF-2/IL-6
for human

[174417]multiple myelomas. Nature 332: 83–85, 1988.

[174418]

[174419]

[174420]17851.

[174421]Kovalchuk, A. L.; Kim, J. S.; Park, S. S.; Coleman, A. E.;
Ward,

[174422]J. M.; Morse, H. C, III; Kishimoto, T.; Potter, M.; Janz, S.: IL-
6

[174423]transgenic mouse model for extraosseous plasmacytoma.

Proc. Nat.

[174424] Acad. Sci. 99: 1509–1514, 2002.

[174425]

[174426]

[174427] 17852.

[174428] May, L. T.; Ghrayeb, J.; Santhanam, U.; Tatter, S. B.;

Sthoeger,

[174429] Z.; Helfgott, D. C.; Chiorazzi, N.; Grieninger, G.; Sehgal, P.

B.

[174430]: Synthesis and secretion of multiple forms of beta–

2–interferon/B–cell

[174431] differentiation factor 2/hepatocyte–stimulating factor by

human fibroblasts

[174432] and monocytes. J. Biol. Chem. 263: 7760–7766, 1988.

[174433]

[174434]

[174435] 17853.

[174436] Ota, N.; Hunt, S. C.; Nakajima, T.; Suzuki, T.; Hosoi, T.;

Orimo,

[174437] H.; Shirai, Y.; Emi, M.: Linkage of interleukin 6 locus to

human

[174438]osteopenia by sibling pair analysis. Hum. Genet. 105:
253–257, 1999.

[174439]

[174440]

[174441]17854.

[174442]Ota, N.; Nakajima, T.; Nakazawa, I.; Suzuki, T.; Hosoi, T.;
Orimo,

[174443]H.; Inoue, S.; Shirai, Y.; Emi, M.: A nucleotide variant in the
promoter

[174444]region of the interleukin–6 gene associated with de-
creased bone mineral

[174445]density. J. Hum. Genet. 46: 267–272, 2001.

[174446]

[174447]

[174448]17855.

[174449]Redwine, L.; Hauger, R. L.; Gillin, J. C.; Irwin, M.: Effects

[174450]of sleep and sleep deprivation on interleukin–6, growth
hormone, cortisol,

[174451]and melatonin levels in humans. J. Clin. Endocr. Metab.

85: 3597–3603,
[174452]2000.

[174453]

[174454]

[174455]17856.

[174456]Roodman, G. D.; Kurihara, N.; Ohsaki, Y.; Kukita, A.;
Hosking,

[174457]D.; Demulder, A.; Smith, J. F.; Singer, F. R.: Interleukin 6: a
potential

[174458]autocrine/paracrine factor in Paget's disease of bone. J.
Clin. Invest. 89:

[174459]46–52, 1992.

[174460]

[174461]

[174462]17857.

[174463]Rooney, M.; David, J.; Symons, J.; Di Giovine, F.; Varsani,
H.;

[174464]Woo, P.: Inflammatory cytokine responses in juvenile
chronic arthritis. Brit.

[174465]J. Rheum. 34: 454–460, 1995.

[174466]

[174467]

[174468]17858.

[174469]Santhanam, U.; Ray, A.; Sehgal, P. B.: Repression of the interleukin

[174470]6 gene promoter by p53 and the retinoblastoma susceptibility gene

[174471]product. Proc. Nat. Acad. Sci. 88: 7605–7609, 1991.

[174472]

[174473]

[174474]17859.

[174475]Scheidt–Nave, C.; Bismar, H.; Leidig–Bruckner, G.; Woitge, H.;

[174476]Seibel, M. J.; Ziegler, R.; Pfeilschifter, J.: Serum interleukin

[174477]6 is a major predictor of bone loss in women specific to the first

[174478]decade past menopause. J. Clin. Endocr. Metab. 86: 2032–2042, 2001.

[174479]

[174480]

[174481]17860.

[174482]Sehgal, P. B.; May, L. T.; Tamm, I.; Vilcek, J.: Human beta-
2

[174483]interferon and B-cell differentiation factor BSF-2 are identical. Science 235:

[174484]731-732, 1987.

[174485]

[174486]

[174487]17861.

[174488]Sehgal, P. B.; Walther, Z.; Tamm, I.: Rapid enhancement of
beta(2)-interferon/B-cell

[174489]differentiation factor BSF-2 gene expression in human fibroblasts

[174490]by diacylglycerols and the calcium ionophore A23187.
Proc. Nat. Acad.

[174491]Sci. 84: 3663-3667, 1987.

[174492]

[174493]

[174494]17862.

[174495]Sehgal, P. B.; Zilberstein, A.; Ruggieri, R.-M.; May, L. T.;

Ferguson-Smith,

[174496]A.; Slate, D. L.; Revel, M.; Ruddle, F. H.: Human chromosome 7 carries

[174497]the beta-2 interferon gene. Proc. Nat. Acad. Sci. 83: 5219-5222,

[174498]1986.

[174499]

[174500]

[174501]17863.

[174502]Lewis, J.; McGowan, E.; Rockwood, J.; Melrose, H.; Nacharaju,

[174503]P.; Van Slegtenhorst, M.; Gwinn-Hardy, K.; Murphy, M. P.; Baker, M.;

[174504]Yu, X.; Duff, K.; Hardy, J.; Corral, A.; Lin, W.-L.; Yen, S.-H.; Dickson,

[174505]D. W.; Davies, P.; Hutton, M.: Neurofibrillary tangles, amyotrophy

[174506]and progressive motor disturbance in mice expressing mutant (P301L)

[174507]tau protein. Nature Genet. 25: 402-405, 2000. Note: Erratum: Nature

[174508]Genet. 26: 127 only, 2000.

[174509]

[174510]

[174511]17864.

[174512]Lippa, C. F.; Zhukareva, V.; Kawarai, T.; Uryu, K.; Shafiq, M.;

[174513]Nee, L. E.; Grafman, J.; Liang, Y.; St George-Hyslop, P. H.; Trojanowski,

[174514]J. Q.; Lee, V. M.-Y.: Frontotemporal dementia with novel tau pathology

[174515]and a glu342val tau mutation. Ann. Neurol. 48: 850–858, 2000.

[174516]

[174517]

[174518]17865.

[174519]Litvan, I.; Baker, M.; Hutton, M.: Tau genotype: no effect on

[174520]onset, symptom severity, or survival in progressive supranuclear palsy. Neurology 57:

[174521]138–140, 2001.

[174522]

[174523]

[174524]17866.

[174525]Martin, E. R.; Scott, W. K.; Nance, M. A.; Watts, R. L.; Hub-
ble,

[174526]J. P.; Koller, W. C.; Lyons, K.; Pahwa, R.; Stern, M. B.;
Colcher,

[174527]A.; Hiner, B. C.; Jankovic, J.; and 20 others: Association of
single-nucleotide

[174528]polymorphisms of the Tau gene with late-onset Parkinson
disease. J.A.M.A. 286:

[174529]2245-2250, 2001.

[174530]

[174531]

[174532]17867.

[174533]Moser, A. B.; Rasmussen, M.; Naidu, S.; Watkins, P. A.;
McGuinness,

[174534]M.; Hajra, A. K.; Chen, G.; Raymond, G.; Liu, A.; Gordon,
D.; Garnaas,

[174535]K.; Walton, D. S.; Skjedal, O. H.; Guggenheim, M. A.; Jack-
son, L.

[174536]G.; Elias, E. R.; Moser, H. W.: Phenotype of patients with

peroxisomal

[174537]disorders subdivided into sixteen complementation
groups. J. Pediat. 127:

[174538]13–22, 1995.

[174539]

[174540]

[174541]17868.

[174542]Paton, B. C.; Heron, S. E.; Nelson, P. V.; Morris, C. P.; Pou-
los,

[174543]A.: Absence of mutations raises doubts about the role of
the 70-kD

[174544]peroxisomal membrane protein in Zellweger syndrome.
(Letter) Am.

[174545]J. Hum. Genet. 60: 1535–1539, 1997.

[174546]

[174547]

[174548]17869.

[174549]Aljada, A.; Ghanim, H.; Friedman, J.; Garg, R.; Mohanty, P.;
Dandona,

[174550]P.: Troglitazone reduces the expression of PPAR–gamma

while stimulating

[174551]that of PPAR-alpha in mononuclear cells in obese subjects. J. Clin.

[174552]Endocr. Metab. 86: 3130-3133, 2001.

[174553]

[174554]

[174555]17870.

[174556]Costet, P.; Legendre, C.; More, J.; Edgar, A.; Galtier, P.; Pineau,

[174557]T.: Peroxisome proliferator-activated receptor alpha-isoform deficiency

[174558]leads to progressive dyslipidemia with sexually dimorphic obesity

[174559]and steatosis. J. Biol. Chem. 273: 29577-29585, 1998.

[174560]

[174561]

[174562]17871.

[174563]Djouadi, F.; Weinheimer, C. J.; Saffitz, J. E.; Pitchford, C.;

[174564]Bastin, J.; Gonzalez, F. J.; Kelly, D. P.: A gender-related defect

[174565]in lipid metabolism and glucose homeostasis in peroxi-
some proliferator-activated

[174566]receptor alpha-deficient mice. J. Clin. Invest. 102:
1083-1091,

[174567]1998.

[174568]

[174569]

[174570]17872.

[174571]Kersten, S.; Desvergne, B.; Wahli, W.: Roles of PPARs in
health

[174572]and disease. Nature 405: 421-424, 2000.

[174573]

[174574]

[174575]17873.

[174576]Kersten, S.; Seydoux, J.; Peters, J. M.; Gonzalez, F. J.;
Desvergne,

[174577]B.; Wahli, W.: Peroxisome proliferator-activated receptor
alpha mediates

[174578]the adaptive response to fasting. J. Clin. Invest. 103:
1489-1498,

[174579]1999.

[174580]

[174581]

[174582]17874.

[174583]Lee, S. S.-T.; Pineau, T.; Drago, J.; Lee, E. J.; Owens, J. W.;

[174584]Kroetz, D. L.; Fernandez-Salguero, P. M.; Westphal, H.;

Gonzalez,

[174585]F. J.: Targeted disruption of the alpha isoform of the per-
oxisome

[174586]proliferator-activated receptor gene in mice results in
abolishment

[174587]of the pleiotropic effects of peroxisome proliferators.

Molec. Cell.

[174588]Biol. 15: 3012-3022, 1995.

[174589]

[174590]

[174591]17875.

[174592]Lee, Y.; Yu, X.; Gonzales, F.; Mangelsdorf, D. J.; Wang,

M.-Y.;

[174593]Richardson, C.; Witters, L. A.; Unger, R. H.: PPAR-alpha is

necessary

[174594]for the lipopenic action of hyperleptinemia on white adipose and liver

[174595]tissue. Proc. Nat. Acad. Sci. 99: 11848–11853, 2002.

[174596]

[174597]

[174598]17876.

[174599]Leone, T. C.; Weinheimer, C. J.; Kelly, D. P.: A critical role

[174600]for the peroxisome proliferator-activated receptor alpha (PPAR-alpha)

[174601]in the cellular fasting response: the PPAR-alpha-null mouse as a model

[174602]of fatty acid oxidation disorders. Proc. Nat. Acad. Sci. 96: 7473–7478,

[174603]1999.

[174604]

[174605]

[174606]17877.

[174607]Michalik, L.; Desvergne, B.; Tan, N. S.; Basu-Modak, S.; Escher,

[174608]P.; Rieusset, J.; Peters, J. M.; Kaya, G.; Gonzalez, F. J.; Zaky,

[174609]J.; Metzger, D.; Chambon, P.; Duboule, D.; Wahli, W.: Impaired skin

[174610]wound healing in peroxisome proliferator-activated receptor (PPAR)-alpha

[174611]and PPAR-beta mutant mice. J. Cell Biol. 154: 799-814, 2001.

[174612]

[174613]

[174614]17878.

[174615]Sher, T.; Yi, H.-F.; McBride, O. W.; Gonzalez, F. J.: cDNA cloning,

[174616]chromosomal mapping, and functional characterization of the human

[174617]peroxisome proliferator activated receptor. Biochemistry 32: 5598-5604,

[174618]1993.

[174619]

[174620]

[174621]17879.

[174622]Xu, H. E.; Stanley, T. B.; Montana, V. G.; Lambert, M. H.;
Shearer,

[174623]B. G.; Cobb, J. E.; McKee, D. D.; Galardi, C. M.; Plunket, K.
D.;

[174624]Nolte, R. T.; Parks, D. J.; Moore, J. T.; Kliwer, S. A.; Will-
son,

[174625]T. M.; Stimmel, J. B.: Structural basis for antagonist-medi-
ated recruitment

[174626]of nuclear co-repressors by PPAR-alpha. Nature 415:
813-817, 2002.

[174627]

[174628]

[174629]17880.

[174630]Ames, G. F.-L.: The basis of multidrug resistance in mam-
malian

[174631]cells: homology with bacterial transport. Cell 47:
323-324, 1986.

[174632]

[174633]

[174634]17881.

[174635]Baldini, N.; Scotlandi, K.; Barbanti-Brodano, G.; Manara, M.

C.;

[174636]Maurici, D.; Bacci, G.; Bertoni, F.; Picci, P.; Sottili, S.; Campanacci,

[174637]M.; Serra, M.: Expression of P-glycoprotein in high-grade osteosarcomas

[174638]in relation to clinical outcome. New Eng. J. Med. 333: 1380-1385,

[174639]1995.

[174640]

[174641]

[174642]17882.

[174643]Bell, D. R.; Trent, J. M.; Willard, H. F.; Riordan, J. R.; Ling,

[174644]V.: Chromosomal location of human P-glycoprotein gene sequences. Cancer

[174645]Genet. Cytogenet. 25: 141-148, 1987.

[174646]

[174647]

[174648]17883.

[174649]Callen, D. F.; Baker, E.; Simmers, R. N.; Seshadri, R.; Roninson,

[174650]I. B.: Localization of the human multiple drug resistance gene, MDR1,

[174651]to 7q21.1. Hum. Genet. 77: 142–144, 1987.

[174652]

[174653]

[174654]17884.

[174655]Chan, H. S. L.; Haddad, G.; Thorner, P. S.; DeBoer, G.; Lin, Y.

[174656]P.; Ondrusek, N.; Yeager, H.; Ling, V.: P-glycoprotein expression

[174657]as a predictor of the outcome of therapy for neuroblastoma. New Eng.

[174658]J. Med. 325: 1608–1614, 1991.

[174659]

[174660]

[174661]17885.

[174662]Chang, G.; Roth, C. B.: Structure of MsbA from E. coli: a homolog

[174663]of the multidrug resistance ATP binding cassette (ABC) transporters. Science 293:

[174664]1793–1800, 2001.

[174665]

[174666]

[174667]17886.

[174668]Chen, C.; Chin, J. E.; Ueda, K.; Clark, D. P.; Pastan, I.;

Gottesman,

[174669]M. M.; Roninson, I. B.: Internal duplication and homology
with bacterial

[174670]transport proteins in the *mdr1* (P-glycoprotein) gene from
multidrug-resistant

[174671]human cells. *Cell* 47: 381–389, 1986.

[174672]

[174673]

[174674]17887.

[174675]Choi, K. H.; Chen, C.-J.; Kriegler, M.; Roninson, I. B.: An
altered

[174676]pattern of cross-resistance in multidrug-resistant human
cells results

[174677]from spontaneous mutations in the *mdr1* (P-glycoprotein)
gene. *Cell* 53:

[174678]519–529, 1988.

[174679]

[174680]

[174681]17888.

[174682]Croop, J. M.; Gros, P.; Housman, D. E.: Genetics of mul-
tidrug

[174683]resistance. J. Clin. Invest. 81: 1303–1309, 1988.

[174684]

[174685]

[174686]17889.

[174687]de Lannoy, I. A. M.; Silverman, M.: The MDR1 gene prod-
uct, P-glycoprotein,

[174688]mediates the transport of the cardiac glycoside, digoxin.
Biochem.

[174689]Biophys. Res. Commun. 189: 551–557, 1992.

[174690]

[174691]

[174692]17890.

[174693]Cameron, P.; Limjuco, G.; Rodkey, J.; Bennett, C.; Schmidt,

J.

[174694]A.: Amino acid sequence analysis of human interleukin 1
(IL-1): evidence

[174695]for biochemically distinct forms of IL-1. J. Exp. Med. 162:
790-801,

[174696]1985.

[174697]

[174698]

[174699]17891.

[174700]D'Eustachio, P.; Jadidi, S.; Fuhlbrigge, R. C.; Gray, P. W.;
Chaplin,

[174701]D. D.: Interleukin-1 alpha and beta genes: linkage on
chromosome

[174702]2 in the mouse. Immunogenetics 26: 339-343, 1987.

[174703]

[174704]

[174705]17892.

[174706]Dinarello, C. A.: An update on human interleukin-1: from
molecular

[174707]biology to clinical relevance. J. Clin. Immun. 5: 287-297,

1985.

[174708]

[174709]

[174710]17893.

[174711]El-Omar, E. M.; Carrington, M.; Chow, W.-H.; McColl, K. E.
L.;

[174712]Bream, J. H.; Young, H. A.; Herrera, J.; Lissowska, J.; Yuan,
C.-C.;

[174713]Rothman, N.; Lanyon, G.; Martin, M.; Fraumeni, J. F., Jr.;
Rabkin,

[174714]C. S.: Interleukin-1 polymorphisms associated with in-
creased risk

[174715]of gastric cancer. Nature 404: 398-402, 2000. Note: Erra-
tum: Nature

[174716]412: 99 only, 2001.

[174717]

[174718]

[174719]17894.

[174720]Furutani, Y.; Notake, M.; Yamayoshi, M.; Yamagishi, J.;
Nomura,

[174721]H.; Ohue, M.; Furuta, R.; Fukui, T.; Yamada, M.; Nakamura,

S.: Cloning

[174722]and characterization of the cDNAs for human and rabbit
interleukin-1

[174723]precursor. Nucleic Acids Res. 13: 5869-5882, 1985.

[174724]

[174725]

[174726]17895.

[174727]Hamajima, N.; Matsuo, K.; Saito, T.; Tajima, K.; Okuma, K.;
Yamao,

[174728]K.; Tominaga, S.: Interleukin 1 polymorphisms, lifestyle
factors,

[174729]and helicobacter pylori infection. Jpn. J. Cancer Res. 92:
383-389,

[174730]2001.

[174731]

[174732]

[174733]17896.

[174734]Le Beau, M. M.; Rowley, J. D.: Personal Communication.
Chicago,

[174735]III. 6/18/1986.

[174736]

[174737]

[174738]17897.

[174739]March, C. J.; Mosley, B.; Larsen, A.; Cerretti, D. P.; Braedt,

[174740]G.; Price, V.; Gillis, S.; Henney, C. S.; Kronheim, S. R.;

Grabstein,

[174741]K.; Conlon, P. J.; Hopp, T. P.; Cosman, D.: Cloning, se-

quence and

[174742]expression of two distinct human interleukin-1 comple-

mentary DNAs. Nature 315:

[174743]641-647, 1985.

[174744]

[174745]

[174746]17898.

[174747]Nicklin, M. J. H.; Weith, A.; Duff, G. W.: A physical map of

[174748]the region encompassing the human interleukin-1-alpha,

interleukin-1-beta,

[174749]and interleukin-1 receptor antagonist genes. Genomics

19: 382-384,

[174750]1994.

[174751]

[174752]

[174753]17899.

[174754]Samad, T. A.; Moore, K. A.; Sapirstein, A.; Billet, S.; All-
chorne,

[174755]A.; Poole, S.; Bonventre, J. V.; Woolf, C. J.: Interleukin-
1-beta-mediated

[174756]induction of Cox-2 in the CNS contributes to inflamma-
tory pain hypersensitivity. Nature 410:

[174757]471-475, 2001.

[174758]

[174759]

[174760]17900.

[174761]Vidal-Vanaclocha, F.; Fantuzzi, G.; Mendoza, L.; Fuentes,
A. M.;

[174762]Anasagasti, M. J.; Martin, J.; Carrascal, T.; Walsh, P.;
Reznikov,

[174763]L. L.; Kim, S.-H.; Novick, D.; Rubinstein, M.; Dinarello, C.
A.:

[174764]IL-18 regulates IL-1-beta-dependent hepatic melanoma
metastasis via

[174765]vascular cell adhesion molecule-1. Proc. Nat. Acad. Sci.
97: 734-739,
[174766]2000.

[174767]

[174768]

[174769]17901.

[174770]Webb, A.; Collins, K.; Auron, P.; Eddy, R.; Nakai, H.; Byers,

[174771]M.; Shows, T. B.: Genetics of acute phase response: a gene
for interleukin-1

[174772]is on chromosome 2. (Abstract) Am. J. Hum. Genet. 37:
A142 only,

[174773]1985.

[174774]

[174775]

[174776]17902.

[174777]Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai,

[174778]H.; Byers, M.; Shows, T. B.: The gene for interleukin-1 (IL1)
is

[174779]on human chromosome 2. (Abstract) Cytogenet. Cell

Genet. 40: 774

[174780]only, 1985.

[174781]

[174782]

[174783]17903.

[174784]Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai,

[174785]H.; Byers, M. G.; Haley, L. L.; Henry, W. M.; Shows, T. B.:

Interleukin-1

[174786]gene (IL1) assigned to long arm of human chromosome 2.

Lymphokine

[174787]Res. 5: 77-85, 1986.

[174788]

[174789]

[174790]17904.

[174791]Ansano, M.; Toda, M.; Sakaguchi, N.; Sakaguchi, S.: Au-

toimmune

[174792]disease as a consequence of developmental abnormality

of a T cell

[174793]subpopulation. J. Exp. Med. 184: 387-396, 1996.

[174794]

[174795]

[174796]17905.

[174797]Cosman, D.; Cerretti, D. P.; Larsen, A.; Park, L.; March, C.;
Dower,

[174798]S.; Gillis, S.; Urdal, D.: Cloning, sequence and expression
of human

[174799]interleukin-2 receptor. Nature 312: 768-771, 1984.

[174800]

[174801]

[174802]17906.

[174803]Ferrari, S.; Cannizzaro, L. A.; Battini, R.; Huebner, K.;
Baserga,

[174804]R.: The gene encoding human vimentin is located on the
short arm

[174805]of chromosome 10. Am. J. Hum. Genet. 41: 616-626,
1987.

[174806]

[174807]

[174808]17907.

[174809]Greene, W. C.; Leonard, W. J.; Depper, J. M.; Nelson, D. L.; Waldmann,

[174810]T. A.: The human interleukin-2 receptor: normal and abnormal expression

[174811]in T cells and in leukemias induced by the human T-lymphotropic retroviruses. Ann.

[174812]Intern. Med. 105: 560-572, 1986.

[174813]

[174814]

[174815]17908.

[174816]Hatakeyama, M.; Minamoto, S.; Taniguchi, T.: Intracytoplasmic

[174817]phosphorylation sites of Tac antigen (p55) are not essential for the

[174818]conformation, function, and regulation of the human interleukin 2

[174819]receptor. Proc. Nat. Acad. Sci. 83: 9650-9654, 1986.

[174820]

[174821]

[174822]17909.

[174823]Hatakeyama, M.; Minamoto, S.; Uchiyama, T.; Hardy, R. R.; Yamada,

[174824]G.; Taniguchi, T.: Reconstitution of functional receptor for human

[174825]interleukin-2 in mouse cells. Nature 318: 467-470, 1985.

[174826]

[174827]

[174828]17910.

[174829]Ihle, J. N.; Kerr, I. M.: Jaks and Stats in signaling by the cytokine

[174830]receptor superfamily. Trends Genet. 11: 69-74, 1995.

[174831]

[174832]

[174833]17911.

[174834]Ishida, N.; Kanamori, H.; Noma, T.; Nikaido, T.; Sabe, H.; Suzuki,

[174835]N.; Shimizu, A.; Honjo, T.: Molecular cloning and structure of the

[174836]human interleukin 2 receptor gene. Nucleic Acids Res. 13: 7579-7589,

[174837]1985.

[174838]

[174839]

[174840]17912.

[174841]Kondo, S.; Shimizu, A.; Maeda, M.; Tagaya, Y.; Yodoi, J.;

Honjo,

[174842]T.: Expression of functional human interleukin-2 receptor
in mouse

[174843]T cells by cDNA transfection. Nature 320: 75-77, 1986.

[174844]

[174845]

[174846]17913.

[174847]Leonard, W. J.; Depper, J. M.; Crabtree, G. R.; Rudikoff, S.;

[174848]Pumphrey, J.; Robb, R. J.; Kronke, M.; Svetlik, P. B.; Pfeffer,
N.

[174849]J.; Waldmann, T. A.; Greene, W. C.: Molecular cloning and
expression

[174850]of cDNAs for the human interleukin-2 receptor. Nature
311: 626-631,

[174851]1984.

[174852]

[174853]

[174854]17914.

[174855]Dahia, P. L. M.; Aguiar, R. C. T.; Alberta, J.; Kum, J. B.;
Caron,

[174856]S.; Sill, H.; Marsh, D. J.; Ritz, J.; Freedman, A.; Stiles, C.;
Eng,

[174857]C.: PTEN is inversely correlated with the cell survival factor
Akt/PKB

[174858]and is inactivated via multiple mechanisms in haemato-
logical malignancies. Hum.

[174859]Molec. Genet. 8: 185–193, 1999.

[174860]

[174861]

[174862]17915.

[174863]Bryson, G. L. M.; Massa, H.; Trask, B. J.; Van Etten, R. L.:
Gene

[174864]structure, sequence, and chromosomal localization of the
human red

[174865]cell-type low molecular weight acid phosphotyrosyl phos-
phatase gene,

[174866]ACP1. Genomics 30: 133–140, 1995.

[174867]

[174868]

[174869]17916.

[174870]Scrabble, H. J.; Johnson, D. K.; Rinchik, E. M.; Cavenee, W. K.

[174871]: Rhabdomyosarcoma-associated locus and MYOD1 are syntenic but separate

[174872]loci on the short arm of human chromosome 11. Proc. Nat. Acad. Sci. 87:

[174873]2182–2186, 1990.

[174874]

[174875]

[174876]17917.

[174877]Chen, T.–R.; McMorris, F. A.; Creagan, R.; Ricciuti, F. C.; Tischfield,

[174878]J.; Ruddle, F. H.: Assignment of the genes for malate oxidoreductase

[174879]decarboxylating to chromosome 6 and peptidase B and lactate dehydrogenase

[174880]B to chromosome 12 in man. Am. J. Hum. Genet. 25:
200–207, 1973.

[174881]

[174882]

[174883]17918.

[174884]Hamerton, J. L.; Mohandas, T.; McAlpine, P. J.; Douglas, G.
R.

[174885]: Localization of human gene loci using spontaneous
chromosome rearrangements

[174886]in human–Chinese hamster somatic cell hybrids. Am. J.
Hum. Genet. 27:

[174887]595–608, 1975.

[174888]

[174889]

[174890]17919.

[174891]Herbschleb–Voogt, E.; Meera Khan, P.: Defining the locus
of origin

[174892]of a genetically determined electrophoretic variant of a
multilocus

[174893]enzyme system; the Calcutta–1 of human LDH system is a
B–locus variant. Hum.

[174894]Genet. 57: 290–295, 1981.

[174895]

[174896]

[174897]17920.

[174898]Houki, N.; Matsushima, Y.; Kitamura, M.; Tukada, T.;
Nishina, T.;

[174899]Nakayama, T.: A case of deficiency of lactate dehydroge-
nase H-subunit. Jpn.

[174900]J. Clin. Chem. 15: 85–90, 1986.

[174901]

[174902]

[174903]17921.

[174904]Kitamura, M.; Iijima, N.; Hashimoto, F.; Hiratsuka, A.:
Hereditary

[174905]deficiency of subunit H of lactate dehydrogenase. Clin.
Chim. Acta 34:

[174906]419–423, 1971.

[174907]

[174908]

[174909]17922.

[174910]Maekawa, M.; Sudo, K.; Kitajima, M.; Matsuura, Y.; Li, S.
S.-L.;

[174911]Kanno, T.: Analysis of a genetic mutation in an elec-
trophoretic variant

[174912]of the human lactate dehydrogenase-B(H) subunit. Hum.
Genet. 91:

[174913]423-426, 1993.

[174914]

[174915]

[174916]17923.

[174917]Maekawa, M.; Sudo, K.; Nagura, K.; Li, S. S.-L.; Kanno, T.:
Population

[174918]screening of lactate dehydrogenase deficiencies in
Fukuoka Prefecture

[174919]in Japan and molecular characterization of three indepen-
dent mutations

[174920]in the lactate dehydrogenase-B(H) gene. Hum. Genet. 93:
74-76, 1994.

[174921]

[174922]

[174923]17924.

[174924]Malpuech, G.; Kaplan, J. C.; Rethore, M. O.; Junien, C.;
Geneix,

[174925]A.: Une observation de deletion partielle du bras court du
chromosome

[174926]12: localisation du gene de la lacticodeshydrogenase B.
Lyon Med. 233:

[174927]275–279, 1975.

[174928]

[174929]

[174930]17925.

[174931]Mayeda, K.; Weiss, L.; Lindahl, R.; Dully, M.: Localization of
[174932]the human lactate dehydrogenase B gene on the short arm
of chromosome

[174933]12. Am. J. Hum. Genet. 26: 59–64, 1974.

[174934]

[174935]

[174936]17926.

[174937]Miwa, S.; Nishima, T.; Kanehashi, Y.; Kitamura, M.; Hirat–

suka,

[174938]A.; Shizume, K.: Studies on erythrocyte metabolism in a case with

[174939]hereditary deficiency of H-subunit of lactate dehydrogenase. Acta

[174940]Haemat. Jpn. 34: 228–232, 1971.

[174941]

[174942]

[174943]17927.

[174944]Bell, D. W.; Varley, J. M.; Szydlo, T. E.; Kang, D. H.; Wahrer,

[174945]D. C. R.; Shannon, K. E.; Lubratovich, M.; Verselis, S. J.; Isselbacher,

[174946]K. J.; Fraumeni, J. F.; Birch, J. M.; Li, F. P.; Garber, J. E.; Haber,

[174947]D. A.: Heterozygous germ line hCHK2 mutations in Li-Fraumeni syndrome. Science 286:

[174948]2528–2531, 1999.

[174949]

[174950]

[174951]17928.

[174952]Hamalainen, E.-R.; Jones, T. A.; Sheer, D.; Taskinen, K.;
Pihlajaniemi,

[174953]T.; Kivirikko, K. I.: Molecular cloning of human lysyl oxi-
dase and

[174954]assignment of the gene to chromosome 5q23.3-31.2. Ge-
nomics 11: 508-516,

[174955]1991.

[174956]

[174957]

[174958]17929.

[174959]Hamalainen, E.-R.; Kempainen, R.; Pihlajaniemi, T.;
Kivirikko,

[174960]K. I.: Structure of the human lysyl oxidase gene. Genomics
17:

[174961]544-548, 1993.

[174962]

[174963]

[174964]17930.

[174965]Kenyon, K.; Contente, S.; Trackman, P. C.; Tang, J.; Kagan,
H.

[174966]M.; Friedman, R. M.: Lysyl oxidase and rrg messenger
RNA. Science 253:

[174967]802 only, 1991.

[174968]

[174969]

[174970]17931.

[174971]Khakoo, A.; Thomas, R.; Trompeter, R.; Duffy, P.; Price, R.;
Pope,

[174972]F. M.: Congenital cutis laxa and lysyl oxidase deficiency.
Clin.

[174973]Genet. 51: 109–114, 1997.

[174974]

[174975]

[174976]17932.

[174977]Kuivaniemi, H.; Savolainen, E.–R.; Kivirikko, K. I.: Human
placental

[174978]lysyl oxidase: purification, partial characterization, and
preparation

[174979]of two specific antisera to the enzyme. J. Biol. Chem. 259:
6996–7002,

[174980]1984.

[174981]

[174982]

[174983]17933.

[174984]Li, W.; Nellaiappan, K.; Strassmaier, T.; Graham, L.;
Thomas, K.

[174985]M.; Kagan, H. M.: Localization and activity of lysyl oxidase
within

[174986]nuclei of fibrogenic cells. Proc. Nat. Acad. Sci. 94:
12817–12822,

[174987]1997.

[174988]

[174989]

[174990]17934.

[174991]Lossie, A. C.; Buckwalter, M. S.; Camper, S. A.: Lysyl oxi-
dase

[174992](Lox) maps between Grl-1 and Adrb-2 on mouse chromo-
some 18. Mammalian

[174993]Genome 4: 177–178, 1993.

[174994]

[174995]

[174996]17935.

[174997]Mariani, T. J.; Trackman, P. C.; Kagan, H. M.; Eddy, R. L.;
Shows,

[174998]T. B.; Boyd, C. D.; Deak, S. B.: The complete derived amino
acid

[174999]sequence of human lysyl oxidase and assignment of the
gene to chromosome

[175000]5 (extensive sequence homology with the murine RAS re-
cision gene). Matrix 12:

[175001]242–248, 1992.

[175002]

[175003]

[175004]17936.

[175005]Mock, B. A.; Contente, S.; Kenyon, K.; Friedman, R. M.;
Kozak,

[175006]C. A.: The gene for lysyl oxidase maps to mouse chromo-
some 18. Genomics 14:

[175007]822–823, 1992.

[175008]

[175009]

[175010]17937.

[175011]Svinarich, D. M.; Twomey, T. A.; Macauley, S. P.; Krebs, C. J.;

[175012]Yang, T. P.; Krawetz, S. A.: Characterization of the human lysyl

[175013]oxidase gene locus. J. Biol. Chem. 267: 14382–14387, 1992.

[175014]

[175015]

[175016]17938.

[175017]Goy, A.; Gilles, F.; Remache, Y.; Zelenetz, A. D.: Physical linkage

[175018]of the lysyl oxidase–like (LOXL1) gene to the PML gene on human chromosome

[175019]15q22. Cytogenet. Cell Genet. 88: 22–24, 2000.

[175020]

[175021]

[175022]17939.

[175023]Kenyon, K.; Modi, W. S.; Contente, S.; Friedman, R. M.: A

novel

[175024]human cDNA with a predicted protein similar to lysyl oxidase maps

[175025]to chromosome 15q24–q25. J. Biol. Chem. 268: 18435–18437, 1993.

[175026]

[175027]

[175028]17940.

[175029]Kim, Y.; Boyd, C. D.; Csiszar, K.: A new gene with sequence and

[175030]structural similarity to the gene encoding human lysyl oxidase. J.

[175031]Biol. Chem. 270: 7176–7182, 1995.

[175032]

[175033]

[175034]17941.

[175035]Wydner, K. S.; Kim, Y.; Csiszar, K.; Boyd, C. D.; Passmore, H.

[175036]C.: An intron capture strategy used to identify and map a lysyl oxidase-like

[175037]gene on chromosome 9 in the mouse. Genomics 40:

342–345, 1997.

[175038]

[175039]

[175040]17942.

[175041]Dudek, H.; Datta, S. R.; Franke, T. F.; Birnbaum, M. J.; Yao, R.;

[175042]Cooper, G. M.; Segal, R. A.; Kaplan, D. R.; Greenberg, M. E.: Regulation

[175043]of neuronal survival by the serine–threonine protein kinase Akt. Science 275:

[175044]661–663, 1997.

[175045]

[175046]

[175047]17943.

[175048]Franke, T. F.; Kaplan, D. R.; Cantley, L. C.; Toker, A.: Direct

[175049]regulation of the Akt proto–oncogene product by phosphatidylinositol–3,4–bisphosphate. Science 275:

[175050]665–667, 1997.

[175051]

[175052]

[175053]17944.

[175054]Franke, T. F.; Yang, S.-I.; Chan, T. O.; Datta, K.; Kazi-
auskas,

[175055]A.; Morrison, D. K.; Kaplan, D. R.; Tschlis, P. N.: The pro-
tein

[175056]kinase encoded by the Akt proto-oncogene is a target of
the PDGF-activated

[175057]phosphatidylinositol 3-kinase. Cell 81: 727-736, 1995.

[175058]

[175059]

[175060]17945.

[175061]Furnari, F. B.; Huang, H. J.; Cavenee, W. K.: The phospho-
inositol

[175062]phosphatase activity of PTEN mediates a serum-sensitive
G1 growth

[175063]arrest in glioma cells. Cancer Res. 58: 5002-5008, 1998.

[175064]

[175065]

[175066]17946.

[175067]Hajdуч, E.; Litherland, G. J.; Hundal, H. S.: Protein kinase

[175068]B (PKB/Akt)--a key regulator of glucose transport? FEBS

Lett. 492:

[175069]199–203, 2001.

[175070]

[175071]

[175072]17947.

[175073]Hemmings, B. A.: Akt signaling: linking membrane events
to life

[175074]and death decisions. Science 275: 628–630, 1997.

[175075]

[175076]

[175077]17948.

[175078]Jones, P. F.; Jakubowicz, T.; Pitossi, F. J.; Maurer, F.; Hem-
mings,

[175079]B. A.: Molecular cloning and identification of a serine/
threonine

[175080]protein kinase of the second-messenger subfamily. Proc.
Nat. Acad.

[175081]Sci. 88: 4171–4175, 1991.

[175082]

[175083]

[175084]17949.

[175085]Lee, M.-J.; Thangada, S.; Paik, J.-H.; Sapkota, G. P.; An-
cellin,

[175086]N.; Chae, S.-S.; Wu, M.; Morales-Ruiz, M.; Sessa, W. C.;
Alessi, D.

[175087]R.; Hla, T.: Akt-mediated phosphorylation of the G pro-
tein-coupled

[175088]receptor EDG-1 is required for endothelial cell chemo-
taxis. Molec.

[175089]Cell 8: 693-704, 2001.

[175090]

[175091]

[175092]17950.

[175093]Maira, S.-M.; Galetic, I.; Brazil, D. P.; Kaech, S.; Ingley, E.;

[175094]Thelen, M.; Hemmings, B. A.: Carboxyl-terminal modula-
tor protein

[175095](CTMP), a negative regulator of PKB/Akt and v-Akt at the
plasma membrane. Scien

[175096]ce 294: 374–380, 2001.

[175097]

[175098]

[175099]17951.

[175100]Powell, D. W.; Rane, M. J.; Chen, Q.; Singh, S.; McLeish, K.
R.

[175101]: Identification of 14–3–3–zeta as a protein kinase B/Akt
substrate. J.

[175102]Biol. Chem. 277: 21639–21642, 2002.

[175103]

[175104]

[175105]17952.

[175106]Staal, S. P.: Molecular cloning of the akt oncogene and its
human

[175107]homologues AKT1 and AKT2: amplification of AKT1 in a
primary human

[175108]gastric adenocarcinoma. Proc. Nat. Acad. Sci. 84:
5034–5037, 1987.

[175109]

[175110]

[175111]17953.

[175112]Staal, S. P.; Huebner, K.; Croce, C. M.; Parsa, N. Z.; Testa,

[175113]J. R.: The AKT1 proto-oncogene maps to human chromosome 14, band

[175114]q32. Genomics 2: 96–98, 1988.

[175115]

[175116]

[175117]17954.

[175118]Vanhaesebroeck, B.; Alessi, D. R.: The PI3K–PDK1 connection:

[175119]more than just a road to PKB. Biochem. J. 346: 561–576, 2000.

[175120]

[175121]

[175122]17955.

[175123]Weng, L.–P.; Brown, J. L.; Eng, C.: PTEN induces apoptosis and

[175124]cell cycle arrest through phosphoinositol–3–kinase/Akt–dependent and

[175125]–independent pathways. Hum. Molec. Genet. 10:

237–242, 2001.

[175126]

[175127]

[175128]17956.

[175129]Yang, J.; Cron, P.; Thompson, V.; Good, V. M.; Hess, D.;
Hemmings,

[175130]B. A.; Barford, D.: Molecular mechanism for the regulation
of protein

[175131]kinase B/Akt by hydrophobic motif phosphorylation.
Molec. Cell 9:

[175132]1227–1240, 2002.

[175133]

[175134]

[175135]17957.

[175136]Le Beau, M. M.; Rowley, J. D.; Sacchi, N.; Watson, D. K.; Pa-
pas,

[175137]T. S.; Diaz, M. O.: Hu-ets-2 is translocated to chromo-
some 8 in the

[175138]t(8;21) in acute myelogenous leukemia. Cancer Genet. Cy-
togenet. 23:

[175139]269–274, 1986.

[175140]

[175141]

[175142]17958.

[175143]Mavrothalassitis, G. J.; Watson, D. K.; Papas, T. S.: Molecular
lar

[175144]and functional characterization of the promoter of ETS2,
the human

[175145]c-ets-2 gene. Proc. Nat. Acad. Sci. 87: 1047-1051, 1990.

[175146]

[175147]

[175148]17959.

[175149]Sacchi, N.; Cheng, S. V.; Tanzi, R. E.; Gusella, J. F.;
Drabkin,

[175150]H. A.; Patterson, D.; Haines, J. H.; Papas, T. S.: The ETS
genes

[175151]on chromosome 21 are distal to the breakpoint of the
acute myelogenous

[175152]leukemia translocation (8;21). Genomics 3: 110-116,
1988.

[175153]

[175154]

[175155]17960.

[175156]Sumarsono, S. H.; Wilson, T. J.; Tymms, M. J.; Venter, D. J.;
Corrick,

[175157]C. M.; Kola, R.; Lahoud, M. H.; Papas, T. S.; Seth, A.; Kola,
I.:

[175158]Down's syndrome-like skeletal abnormalities in Ets2
transgenic mice. Nature 379:

[175159]534–540, 1996.

[175160]

[175161]

[175162]17961.

[175163]Chu, E. H. Y.; Chang, C. C.; Sun, N. C.: Synteny of the hu-
man

[175164]genes for gal-1-PT, ACP-1, MDH-1, and gal-
plus-activator and assignment

[175165]to chromosome 2. Birth Defects Orig. Art. Ser. XI(3):
103–106, 1975.

[175166]Note: Alternate: Cytogenet. Cell Genet. 14: 273–276,
1975...

[175167]

[175168]

[175169]17962.

[175170]Rasmussen, S. K.; Lautier, C.; Hansen, L.; Echwald, S. M.;

Hansen,

[175171]T.; Ekstrom, C. T.; Urhammer, S. A.; Borch-Johnsen, K.;

Grigorescu,

[175172]F.; Smith, R. J.; Pedersen, O.: Studies of the variability of
the

[175173]genes encoding the insulin-like growth factor I receptor
and its ligand

[175174]in relation to type 2 diabetes mellitus. J. Clin. Endocr.

Metab. 85:

[175175]1606–1610, 2000.

[175176]

[175177]

[175178]17963.

[175179]Dissing, J.; Johnsen, A. H.: Human red cell acid phos-
phatase (ACP1):

[175180]the primary structure of the two pairs of isozymes en-
coded by the

[175181]ACP1*A and ACP1*C alleles. Biochim. Biophys. Acta 1121:

261-268,

[175182]1992.

[175183]

[175184]

[175185]17964.

[175186]Dissing, J.; Johnsen, A. H.; Sensabaugh, G. F.: Human red
cell

[175187]acid phosphatase (ACP1): the amino acid sequence of the
two isozymes

[175188]Bf and Bs encoded by the ACP1*B allele. J. Biol. Chem.
266: 20619-20625,

[175189]1991.

[175190]

[175191]

[175192]17965.

[175193]Emanuel, B. S.; Zackai, E. H.; Van Dyke, D. C.; Swallow, D.
M.;

[175194]Allen, F. H.; Mellman, W. J.: Deletion mapping: further evi-
dence

[175195]for the location of acid phosphatase (ACP-1) within 2p23.

Am. J.

[175196]Med. Genet. 4: 167–172, 1979.

[175197]

[175198]

[175199]17966.

[175200]Ferguson-Smith, M. A.; Newman, B. F.; Ellis, P. M.; Thomson, D.

[175201]M. G.; Riley, I. D.: Assignment by deletion of human red cell acid

[175202]phosphatase gene locus to the short arm of chromosome 2. Nature 243:

[175203]271–273, 1973.

[175204]

[175205]

[175206]17967.

[175207]Fisher, R. A.; Harris, H.: Studies on the separate isoenzymes

[175208]of red cell acid phosphatase phenotypes A and B: chromatographic separation

[175209]of the isoenzymes. Ann. Hum. Genet. 34: 431–438, 1971.

[175210]

[175211]

[175212]17968.

[175213]Fuhrmann, W.; Lichte, K. H.: Human red cell acid phosphatase

[175214]polymorphism: a study on gene frequency and forensic use of the system

[175215]in cases of disputed paternity. Humangenetik 3: 121–126, 1966.

[175216]

[175217]

[175218]17969.

[175219]Giblett, E. R.; Scott, N. M.: Red cell acid phosphatase: racial

[175220]distribution and report of a new phenotype. Am. J. Hum. Genet. 17:

[175221]425–432, 1965.

[175222]

[175223]

[175224]17970.

[175225]Herbich, J.; Fisher, R. A.; Hopkinson, D. A.: Atypical segregation

[175226]of human red cell acid phosphatase phenotypes: evidence for a rare

[175227]'silent' allele P(O). Ann. Hum. Genet. 34: 145–152, 1970.

[175228]

[175229]

[175230]17971.

[175231]Herbich, J.; Meinhart, K.: The rare 'silent' allele P(O) or P(V)

[175232](P Vienna) of human red cell acid phosphatase, typed in a second family. Humangenetik 15:

[175233]345–348, 1972.

[175234]

[175235]

[175236]17972.

[175237]Hopkinson, D. A.; Spencer, N.; Harris, H.: Red cell acid phosphatase

[175238]variants: a new human polymorphism. Nature 199: 969–971, 1963.

[175239]

[175240]

[175241]17973.

[175242]Hulten, M.; Lindsten, J.; Pen-Ming, L. M.; Fraccaro, M.;
Mannini,

[175243]A.; Tiepolo, L.; Robson, E. B.; Heiken, A.; Tillinger, K. G.:
Possible

[175244]localization of the genes for the Kidd blood group on an
autosome

[175245]involved in a reciprocal translocation. Nature 211:
1067-1068, 1966.

[175246]

[175247]

[175248]17974.

[175249]Junien, C.; Kaplan, J.-C.; Bernheim, A.; Berger, R.: Regional

[175250]assignment of red cell acid phosphatase locus to band
2p25. Hum.

[175251]Genet. 48: 17-21, 1979.

[175252]

[175253]

[175254]17975.

[175255]Karp, G. W., Jr.; Sutton, H. E.: Some new phenotypes of human

[175256]red cell acid phosphatase. Am. J. Hum. Genet. 19: 54–62, 1967.

[175257]

[175258]

[175259]17976.

[175260]Lothe, R. A.; Gedde–Dahl, T.; Olaisen, B.; Bakker, E.; Pearson, son,

[175261]P.: Very close linkage between D2S1 and ACP1 on chromosome 2p. Ann.

[175262]Hum. Genet. 50: 361–367, 1986.

[175263]

[175264]

[175265]17977.

[175266]Mace, M. A.; Cook, P. J. L.; Robson, E. B.: Linkage data on red

[175267]cell acid phosphatase from family studies. Ann. Hum. Genet. 38:

[175268]471–477, 1975.

[175269]

[175270]

[175271]17978.

[175272]Mace, M. A.; Robson, E. B.: Linkage data on ACP-1 and
MNSs. Cytogenet.

[175273]Cell Genet. 13: 123-125, 1974.

[175274]

[175275]

[175276]17979.

[175277]Magenis, R. E.; Koler, R. D.; Lovrien, E. W.; Bigley, R. H.;
Duval,

[175278]M. C.; Overton, K. M.: Gene dosage: evidence for assign-
ment of erythrocyte

[175279]acid phosphatase locus to chromosome 2. Proc. Nat.
Acad. Sci. 72:

[175280]4526-4530, 1975.

[175281]

[175282]

[175283]17980.

[175284]Mayr, W. R.: No close linkage between MNSs and red cell
acid

[175285]phosphatase. Hum. Hered. 26: 1–3, 1976.

[175286]

[175287]

[175288]17981.

[175289]Miller, S. A.; Nelson, M. S.; Dykes, D. D.; Polesky, H. F.:

Comparison

[175290]of acid phosphatase ACP1 variants by isoelectric focusing
and conventional

[175291]electrophoresis: identification of three new alleles,

ACP1*N, ACP1*P

[175292]and ACP1*S. Hum. Hered. 37: 371-375, 1987.

[175293]

[175294]

[175295]17982.

[175296]Mohrenweiser, H. W.; Novotny, J. E.: ACP-1-GUA-1: a low-
activity

[175297]variant of human erythrocyte acid phosphatase: associa-
tion with increased

[175298]glutathione reductase activity. Am. J. Hum. Genet. 34:
425-433,

[175299]1982.

[175300]

[175301]

[175302]17983.

[175303]Nezbeda, P.: Occurrence of the ACP-1 null allele in
Czechoslovakia. Hum.

[175304]Genet. 46: 227-229, 1979.

[175305]

[175306]

[175307]17984.

[175308]Palmarino, R.; Agostino, R.; Gloria, F.; Lucarelli, P.; Bus-
inco,

[175309]L.; Antognoni, G.; Maggioni, G.; Workman, P. L.; Bottini, E.:
Red

[175310]cell acid phosphatase: another polymorphism correlated
with malaria? Am.

[175311]J. Phys. Anthrop. 43: 177-186, 1975.

[175312]

[175313]

[175314]17985.

[175315]Povey, S.; Swallow, D. M.; Bobrow, M.; Craig, I.; Van

Heyningen,

[175316]V.: Probable assignment of the locus determining human red cell acid

[175317]phosphatase ACP(1) to chromosome 2 using somatic cell hybrids. Ann.

[175318]Hum. Genet. 38: 1–5, 1974.

[175319]

[175320]

[175321]17986.

[175322]Renwick, J. H.: Assignment and map–positioning of human loci

[175323]using chromosomal variation. Ann. Hum. Genet. 35: 79–97, 1971.

[175324]

[175325]

[175326]17987.

[175327]Sensabaugh, G. F.; Golden, V. L.: Phenotype dependence in the

[175328]inhibition of red cell acid phosphatase (ACP) by folates. Am. J.

[175329]Hum. Genet. 30: 553–560, 1978.

[175330]

[175331]

[175332]17988.

[175333]Swallow, D. M.; Povey, S.; Harris, H.: Activity of the 'red
cell'

[175334]acid phosphatase locus in other tissues. Ann. Hum. Genet.
37: 31–38,

[175335]1973.

[175336]

[175337]

[175338]17989.

[175339]Wu, R. M.; Cheng, C. W.; Chen, K. H.; Lu, S. L.; Shan, D. E.;

[175340]Ho, Y. F.; Chern, H. D.: The COMT L allele modifies the as-
sociation

[175341]between MAOB polymorphism and PD in Taiwanese. Neu-
rology 56: 375–382,

[175342]2001.

[175343]

[175344]

[175345]17990.

[175346]Levy, J. B.; Canoll, P. D.; Silvennoinen, O.; Barnea, G.;

Morse,

[175347]B.; Honegger, A. M.; Huang, J.-T.; Cannizzaro, L. A.; Park,

S.-H.;

[175348]Druck, T.; Huebner, K.; Sap, J.; Ehrlich, M.; Musacchio, J.

M.; Schlessinger,

[175349]J.: The cloning of a receptor-type protein tyrosine phosphatase expressed

[175350]in the central nervous system. J. Biol. Chem. 268:

10573-10581,

[175351]1993.

[175352]

[175353]

[175354]17991.

[175355]Berube, D.; Simard, J.; Sandberg, M.; Grzeschik, K.-H.;

Gagne,

[175356]R.; Hansson, V.; Jahnsen, T.: Assignment of the gene encoding the

[175357]catalytic subunit C(beta) of cAMP-dependent protein kinase to the

[175358]p36 band on chromosome 1. (Abstract) Cytogenet. Cell
Genet. 58:

[175359]1850 only, 1991.

[175360]

[175361]

[175362]17992.

[175363]Leonard, W. J.; Depper, J. M.; Kanehisa, M.; Kronke, M.;
Peffer,

[175364]N. J.; Svetlik, P. B.; Sullivan, M.; Greene, W. C.: Structure of

[175365]the human interleukin-2 receptor gene. Science 230:
633-639, 1985.

[175366]

[175367]

[175368]17993.

[175369]Leonard, W. J.; Depper, J. M.; Robb, R. J.; Waldmann, T. A.;
Greene,

[175370]W. C.: Characterization of the human receptor for T-cell
growth factor. Proc.

[175371]Nat. Acad. Sci. 80: 6957-6961, 1983.

[175372]

[175373]

[175374]17994.

[175375]Marx, J. L.: The interleukin-2 receptor gene is cloned. Science 226:

[175376]1064-1065, 1985.

[175377]

[175378]

[175379]17995.

[175380]Nikaido, T.; Shimizu, A.; Ishida, N.; Sabe, H.; Teshigawara, K.;

[175381]Maeda, M.; Uchiyama, T.; Yodoi, J.; Honjo, T.: Molecular cloning

[175382]of a cDNA encoding human interleukin-2 receptor. Nature 311: 631-635,

[175383]1984.

[175384]

[175385]

[175386]17996.

[175387]Robb, R. J.; Rusk, C. M.; Neeper, M. P.: Structure-function

relationships

[175388]for the interleukin 2 receptor: location of ligand and anti-body binding

[175389]sites on the Tac receptor chain by mutational analysis.

Proc. Nat.

[175390]Acad. Sci. 85: 5654–5658, 1988.

[175391]

[175392]

[175393]17997.

[175394]Sharfe, N.; Dadi, H. K.; Shahar, M.; Roifman, C. M.: Human immune

[175395]disorder arising from mutation of the alpha chain of the interleukin–2

[175396]receptor. Proc. Nat. Acad. Sci. 94: 3168–3171, 1997.

[175397]

[175398]

[175399]17998.

[175400]Shevach, E. M.: Certified professionals: CD4(+)CD25(+) suppressor

[175401]T cells. J. Exp. Med. 193: F41–F45, 2001.

[175402]

[175403]

[175404]17999.

[175405]Tsudo, M.; Kozak, R. W.; Goldman, C. K.; Waldmann, T. A.:

Demonstration

[175406]of a non-Tac peptide that binds interleukin 2: a potential
participant

[175407]in a multichain interleukin 2 receptor complex. Proc. Nat.
Acad.

[175408]Sci. 83: 9694–9698, 1986.

[175409]

[175410]

[175411]18000.

[175412]Urdal, D. L.; March, C. J.; Gillis, S.; Larsen, A.; Dower, S.

[175413]K.: Purification and chemical characterization of the re-
ceptor for

[175414]interleukin 2 from activated human T lymphocytes and
from a human

[175415]T-cell lymphoma cell line. Proc. Nat. Acad. Sci. 81:
6481–6485,

[175416]1984.

[175417]

[175418]

[175419]18001.

[175420]Celestin, J.; Rotschke, O.; Falk, K.; Ramesh, N.; Jabara, H.;

Strominger,

[175421]J.; Geha, R. S.: IL-3 induces B7.2 (CD86) expression and
costimulatory

[175422]activity in human eosinophils. J. Immun. 167: 6097-6104,
2001.

[175423]

[175424]

[175425]18002.

[175426]Mohrenweiser, H. W.; Neel, J. V.: Frequency of thermosta-
bility

[175427]variants: estimation of total 'rare' variant frequency in hu-
man populations. Proc.

[175428]Nat. Acad. Sci. 78: 5729-5733, 1981.

[175429]

[175430]

[175431]18003.

[175432]Gervais, F. G.; Singaraja, R.; Xanthoudakis, S.; Gutekunst, C.-A.;

[175433]Leavitt, B. R.; Metzler, M.; Hackam, A. S.; Tam, J.; Vaillancourt,

[175434]J. P.; Houtzager, V.; Rasper, D. M.; Roy, S.; Hayden, M. R.; Nicholson,

[175435]D. W.: Recruitment and activation of caspase-8 by the huntingtin-interacting

[175436]protein Hip-1 and a novel partner Hippi. *Nature Cell Biol.* 4: 95-105,

[175437]2002.

[175438]

[175439]

[175440]18004.

[175441]Abbott, A. M.; Bueno, R.; Pedrini, M. T.; Murray, J. M.; Smith,

[175442]R. J.: Insulin-like growth factor I receptor gene structure. J.

[175443]*Biol. Chem.* 267: 10759-10763, 1992.

[175444]

[175445]

[175446]18005.

[175447]All-Ericsson, C.; Girnita, L.; Seregard, S.; Bartolazzi, A.;
Jager,

[175448]M. J.; Larsson, O.: Insulin-like growth factor-1 receptor in
uveal

[175449]melanoma: a predictor for metastatic disease and a po-
tential therapeutic

[175450]target. Invest. Ophthalm. Vis. Sci. 43: 1-8, 2002.

[175451]

[175452]

[175453]18006.

[175454]Cooke, D. W.; Bankert, L. A.; Roberts, C. T., Jr.; LeRoith,
D.;

[175455]Casella, S. J.: Analysis of the human type I insulin-like
growth

[175456]factor receptor promoter region. Biochem. Biophys. Res.
Commun. 177:

[175457]1113-1120, 1991.

[175458]

[175459]

[175460]18007.

[175461]Dey, B. R.; Furlanetto, R. W.; Nissley, S. P.: Cloning of human
man

[175462]p55-gamma, a regulatory subunit of phosphatidylinositol
3-kinase,

[175463]by a yeast two-hybrid library screen with the insulin-like
growth

[175464]factor-I receptor. Gene 209: 175-183, 1998.

[175465]

[175466]

[175467]18008.

[175468]Fernandez, A. M.; Kim, J. K.; Yakar, S.; Dupont, J.; Hernan-
dez-Sanchez,

[175469]C.; Castle, A. L.; Filmore, J.; Shulman, G. I.; Le Roith, D.:
Functional

[175470]inactivation of the IGF-I and insulin receptors in skeletal
muscle

[175471]causes type 2 diabetes. Genes Dev. 15: 1926-1934, 2001.

[175472]

[175473]

[175474]18009.

[175475]Flier, J. S.; Usher, P.; Moses, A. C.: Monoclonal antibody to

[175476]the type I insulin-like growth factor (IGF-I) receptor blocks
IGF-I

[175477]receptor-mediated DNA synthesis: clarification of the mi-
togenic mechanisms

[175478]of IGF-I and insulin in human skin fibroblasts. Proc. Nat.
Acad.

[175479]Sci. 83: 664-668, 1986.

[175480]

[175481]

[175482]18010.

[175483]Francke, U.; Yang-Feng, T. L.; Brissenden, J. E.; Ullrich, A.:

[175484]Chromosomal mapping of genes involved in growth con-
trol. Cold Spring

[175485]Harbor Symp. Quant. Biol. 51: 855-866, 1986.

[175486]

[175487]

[175488]18011.

[175489]Grant, E. S.; Ross, M. B.; Ballard, S.; Naylor, A.; Habib, F. K.

[175490]: The insulin-like growth factor type I receptor stimulates growth

[175491]and suppresses apoptosis in prostatic stromal cells. J.

Clin. Endocr.

[175492]Metab. 83: 3252–3257, 1998.

[175493]

[175494]

[175495]18012.

[175496]Herskowitz, I.: Functional inactivation of genes by dominant negative

[175497]mutations. Nature 329: 219–222, 1987.

[175498]

[175499]

[175500]18013.

[175501]Howard, T. K.; Algar, E. M.; Glatz, J. A.; Reeve, A. E.; Smith,

[175502]P. J.: The insulin-like growth factor 1 receptor gene is normally

[175503]biallelically expressed in human juvenile tissue and tumours. Hum.

[175504]Molec. Genet. 2: 2089–2092, 1993.

[175505]

[175506]

[175507]18014.

[175508]Kulkarni, R. N.; Holzenberger, M.; Shih, D. Q.; Ozcan, U.; Stoffel,

[175509]M.; Magnuson, M. A.; Kahn, C. R.: Beta-cell-specific deletion of

[175510]the Igf1 receptor leads to hyperinsulinemia and glucose intolerance

[175511]but does not alter beta-cell mass. Nature Genet. 31: 111–115, 2002.

[175512]

[175513]

[175514]18015.

[175515]Maor, S. B.; Abramovitch, S.; Erdos, M. R.; Brody, L. C.; Werner,

[175516]H.: BRCA1 suppresses insulin-like growth factor-I receptor promoter

[175517]activity: potential interaction between BRCA1 and Sp1.

Molec. Genet.

[175518]Metab. 69: 130–136, 2000.

[175519]

[175520]

[175521]18016.

[175522]Poduslo, S. E.; Dean, M.; Kolch, U.; O'Brien, S. J.: Detecting

[175523]high-resolution polymorphisms in human coding loci by
combining PCR

[175524]and single-strand conformation polymorphism (SSCP)
analysis. Am.

[175525]J. Hum. Genet. 49: 106–111, 1991.

[175526]

[175527]

[175528]18017.

[175529]Prager, D.; Yamasaki, H.; Weber, M. M.; Gebremedhin, S.;
Melmed,

[175530]S.: Human insulin-like growth factor I receptor function in
pituitary

[175531]cells is suppressed by a dominant negative mutant. J. Clin.

Invest. 90:

[175532]2117-2122, 1992.

[175533]

[175534]

[175535]18018.

[175536]Roback, E. W.; Barakat, A. J.; Dev, V. G.; Mbikay, M.; Chretien,

[175537]M.; Butler, M. G.: An infant with deletion of the distal long arm

[175538]of chromosome 15 (q26.1-qter) and loss of insulin-like growth factor

[175539]1 receptor gene. Am. J. Med. Genet. 38: 74-79, 1991.

[175540]

[175541]

[175542]18019.

[175543]Okumura, N.; Terasawa, F.; Ueno, I.; Oki, K.; Yamauchi, K.; Hidaka,

[175544]H.; Tozuka, M.; Okura, M.; Katsuyama, T.: Genetic analyses in homozygous

[175545]and heterozygous variants of lactate dehydrogenase-B (H)

subunit:

[175546]LD-B Matsumoto I and II (LD-B W323R). Clin. Chim. Acta
287: 163–171,

[175547]1999.

[175548]

[175549]

[175550]18020.

[175551]Rethore, M.–O.; Junien, C.; Malpuech, G.; Baccichetti, C.;
Tenconi,

[175552]R.; Kaplan, J. C.; de Romeuf, J.; Lejeune, J.: Localisation du
gene

[175553]de la glyceraldehyde 3–phosphate dehydrogenase (G3PD)
sur le segment

[175554]distal du bras court du chromosome 12. Ann. Genet. 19:
140–142,

[175555]1976.

[175556]

[175557]

[175558]18021.

[175559]Rethore, M.–O.; Kaplan, J.–C.; Junien, C.; Cruveiller, J.;

Dutrillaux,

[175560]B.; Aurias, A.; Carpentier, S.; Lafourcade, J.; Lejeune, J.:

Augmentation

[175561]de l'activite de la LDH-B chez un garcon trisomique 12p

par malsegregation

[175562]d'une translocation maternelle t(12;14)(q12;p11). Ann.

Genet. 18:

[175563]81-87, 1975.

[175564]

[175565]

[175566]18022.

[175567]Sakai, I.; Sharief, F. S.; Pan, Y.-C. E.; Li, S. S.-L.: The cDNA

[175568]and protein sequences of human lactate dehydrogenase B.

Biochem.

[175569]J. 248: 933-936, 1987.

[175570]

[175571]

[175572]18023.

[175573]Santachiara, A. S.; Nabholz, M.; Miggiano, V.; Darlington,

A.

[175574]J.; Bodmer, W. F.: Linkage between human lactate dehydrogenase B

[175575]and peptidase B genes. Nature 227: 248–251, 1970.

[175576]

[175577]

[175578]18024.

[175579]Steinbach, P.; Rehder, H.: Tetrasomy for the short arm of chromosome

[175580]12 with accessory isochromosome (+i[12p]) and a marked LDH-B gene

[175581]dosage effect. Clin. Genet. 32: 1–4, 1987.

[175582]

[175583]

[175584]18025.

[175585]Sudo, K.: Personal Communication. Komae City, Japan
7/12/1993.

[175586]

[175587]

[175588]18026.

[175589]Sudo, K.; Maekawa, M.; Ikawa, S.; Machida, K.; Kitamura, M.; Li,

[175590]S. S.-L.: A missense mutation found in human lactate dehydrogenase-B

[175591](H) variant gene. Biochem. Biophys. Res. Commun. 168: 672-676, 1990.

[175592]

[175593]

[175594]18027.

[175595]Sudo, K.; Maekawa, M.; Luedemann, M. M.; Deaven, L. L.; Li, S.

[175596]S.-L.: Human lactate dehydrogenase-B processed pseudo-gene: nucleotide

[175597]sequence analysis and assignment to the X-chromosome. Biochem. Biophys.

[175598]Res. Commun. 171: 67-74, 1990.

[175599]

[175600]

[175601]18028.

[175602]Sudo, K.; Maekawa, M.; Tomonaga, A.; Tsukada, T.; Nakayama, T.;

[175603]Kitamura, M.; Li, S. S.-L.; Kanno, T.; Toriumi, J.: Molecular
characterization

[175604]of genetic mutations in human lactate dehydrogenase
(LDH)B (H) variant. Hum.

[175605]Genet. 89: 158-162, 1992.

[175606]

[175607]

[175608]18029.

[175609]Van Someren, H.; Meera Khan, P.; Westerveld, A.; Bootsma,
D.:

[175610]Human genetics--two new linkage groups carrying differ-
ent loci for

[175611]LDH and glutamic-pyruvic transaminase found. Nature
240: 221-222,

[175612]1972.

[175613]

[175614]

[175615]18030.

[175616]Weiss, L.; Mayeda, K.; Lindahl, R.; Dully, M.: Localization of

[175617]human LDH-B gene of the short arm of chromosome 12.

(Abstract) Am.

[175618]J. Hum. Genet. 25: 85A only, 1973.

[175619]

[175620]

[175621]18031.

[175622]Davies, H.; Bignell, G. R.; Cox, C.; Stephens, P.; Edkins, S.;

[175623]Clegg, S.; Teague, J.; Woffendin, H.; Garnett, M. J.; Bot-
tomley, W.;

[175624]Davis, N.; Dicks, E.; and 40 others: Mutations of the BRAF
gene

[175625]in human cancer. Nature 417: 949–954, 2002.

[175626]

[175627]

[175628]18032.

[175629]Eychene, A.; Barnier, J. V.; Apiou, F.; Dutrillaux, B.;
Calothy,

[175630]G.: Chromosomal assignment of two human B-raf(Rmil)
proto-oncogene

[175631]loci: B-raf-1 encoding the p94(Braf/Rmil) and B-raf-2, a
processed

[175632]pseudogene. Oncogene 7: 1657–1660, 1992.

[175633]

[175634]

[175635]18033.

[175636]Rajagopalan, H.; Bardelli, A.; Lengauer, C.; Kinzler, K. W.;
Vogelstein,

[175637]B.; Velculescu, V. E.: RAF/RAS oncogenes and mismatch–re–
pair status.

[175638](Letter) Nature 418: 934 only, 2002.

[175639]

[175640]

[175641]18034.

[175642]Sithanandam, G.; Druck, T.; Cannizzaro, L. A.; Leuzzi, G.;
Huebner,

[175643]K.; Rapp, U. R.: B–raf and a B–raf pseudogene are located
on 7q in

[175644]man. Oncogene 7: 795–799, 1992.

[175645]

[175646]

[175647]18035.

[175648]Wojnowski, L.; Zimmer, A. M.; Beck, T. W.; Hahn, H.;
Bernal, R.;

[175649]Rapp, U. R.; Zimmer, A.: Endothelial apoptosis in Braf-
deficient

[175650]mice. Nature Genet. 16: 293–297, 1997.

[175651]

[175652]

[175653]18036.

[175654]Zhong, S.; Delva, L.; Rachez, C.; Cenciarelli, C.; Gandini,
D.;

[175655]Zhang, H.; Kalantry, S.; Freedman, L. P.; Pandolfi, P. P.: A
RA-dependent,

[175656]tumour-growth suppressive transcription complex is the
target of the

[175657]PML-RAR-alpha and T18 oncoproteins. Nature Genet. 23:
287–295, 1999.

[175658]

[175659]

[175660]18037.

[175661]Sakamoto, O.; Iwama, A.; Amitani, R.; Takehara, T.; Yamaguchi,

[175662]N.; Yamamoto, T.; Masuyama, K.; Yamanaka, T.; Ando, M.; Suda, T.:

[175663]Role of macrophage-stimulating protein and its receptor, RON tyrosine

[175664]kinase, in ciliary motility. J. Clin. Invest. 99: 701-709, 1997.

[175665]

[175666]

[175667]18038.

[175668]Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura, N.; Shimizu,

[175669]N.: Assignment of hepatocyte growth factor (HGF) gene to chromosome

[175670]7q21.1. (Abstract) Cytogenet. Cell Genet. 58: 1921 only, 1991.

[175671]

[175672]

[175673]18039.

[175674]Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura, N.;

Shimizu,

[175675]N.: Regional localization of the hepatocyte growth factor
(HGF) gene

[175676]to human chromosome 7 band q21.1. Genomics 11:
410-415, 1991.

[175677]

[175678]

[175679]18040.

[175680]Gherardi, E.; Stoker, M.: Hepatocytes and scatter factor.
Nature 346:

[175681]228 only, 1990.

[175682]

[175683]

[175684]18041.

[175685]Gohda, E.; Tsubouchi, H.; Nakayama, H.; Hirono, S.;
Sakiyama, O.;

[175686]Takahashi, K.; Miyazaki, H.; Hashimoto, S.; Daikuhara, Y.:
Purification

[175687]and partial characterization of hepatocyte growth factor
from plasma

[175688]of a patient with fulminant hepatic failure. J. Clin. Invest.

81:

[175689]414–419, 1988.

[175690]

[175691]

[175692]18042.

[175693]Kilby, M. D.; Afford, S.; Li, X. F.; Strain, A. J.; Ahmed, A.;

[175694]Whittle, M. J.: Localisation of hepatocyte growth factor and
its

[175695]receptor (c-met) protein and mRNA in human term pla-
centa. Growth

[175696]Factors 13: 133–139, 1996.

[175697]

[175698]

[175699]18043.

[175700]Lai, L.; Goldschneider, I.: Cutting edge: identification of a

[175701]hybrid cytokine consisting of IL-7 and the beta-chain of
the hepatocyte

[175702]growth factor/scatter factor. J. Immun. 167: 3550–3554,
2001.

[175703]

[175704]

[175705]18044.

[175706]Maina, F.; Casagrande, F.; Audero, E.; Simeone, A.; Comoglio, P.

[175707]M.; Klein, R.; Ponzetto, C.: Uncoupling of Grb2 from the Met receptor

[175708]in vivo reveals complex roles in muscle development. Cell 87: 531–542,

[175709]1996.

[175710]

[175711]

[175712]18045.

[175713]Miyazawa, K.; Tsubouchi, H.; Naka, D.; Takahashi, K.; Oikawa, K.

[175714]M.; Arakaki, N.; Nakayama, H.; Hirono, S.; Sakiyama, O.; Takahashi,

[175715]K.; Gohda, E.; Daikuhara, Y.; Kitamura, N.: Molecular cloning and

[175716]sequence analysis of cDNA for human hepatocyte growth factor. Biochem.

[175717]Biophys. Res. Commun. 163: 967–973, 1989.

[175718]

[175719]

[175720]18046.

[175721]Nakamura, T.; Nishizawa, T.; Hagiya, M.; Seki, T.; Shimon-
ishi,

[175722]M.; Sugimura, A.; Tashiro, K.; Shimizu, S.: Molecular
cloning and

[175723]expression of human hepatocyte growth factor. Nature
342: 440–443,

[175724]1989.

[175725]

[175726]

[175727]18047.

[175728]Noonan, F. P.; Recio, J. A.; Takayama, H.; Duray, P.; Anver,
M.

[175729]R.; Rush, W. L.; De Fabo, E. C.; Merlino, G.: Neonatal sun-
burn and

[175730]melanoma in mice: severe sunburn in newborn, but not
adult, mice is

[175731]linked with melanoma in later life. Nature 413: 271–272, 2001.

[175732]

[175733]

[175734]18048.

[175735]Powell, E. M.; Mars, W. M.; Levitt, P.: Hepatocyte growth factor/scatter

[175736]factor is a motogen for interneurons migrating from the ventral to

[175737]dorsal telencephalon. Neuron 30: 79–89, 2001.

[175738]

[175739]

[175740]18049.

[175741]Rubin, J. S.; Chan, A. M.–L.; Bottaro, D. P.; Burgess, W. H.;

[175742]Taylor, W. G.; Cech, A. C.; Hirschfield, D. W.; Wong, J.; Miki, T.;

[175743]Finch, P. W.; Aaronson, S. A.: A broad–spectrum human lung fibroblast–derived

[175744]mitogen is a variant of hepatocyte growth factor. Proc. Nat. Acad.

[175745]Sci. 88: 415–419, 1991.

[175746]

[175747]

[175748]18050.

[175749]Saccone, S.; Narsimhan, R. P.; Gaudino, G.; Dalpra, L.; Comoglio,

[175750]P. M.; Della Valle, G.: Regional mapping of the human hepatocyte

[175751]growth factor (HGF)–scatter factor gene to chromosome 7q21.1. Genomics 13:

[175752]912–914, 1992.

[175753]

[175754]

[175755]18051.

[175756]Schmidt, C.; Bladt, F.; Goedecke, S.; Brinkmann, V.; Zschiesche,

[175757]W.; Sharpe, M.; Gherardi, E.; Birchmeier, C.: Scatter factor/hepatocyte

[175758]growth factor is essential for liver development. Nature 373: 699–702,

[175759]1995.

[175760]

[175761]

[175762]18052.

[175763]Uehara, Y.; Minowa, O.; Mori, C.; Shiota, K.; Kuno, J.;

Noda,

[175764]T.; Kitamura, N.: Placental defect and embryonic lethality
in mice

[175765]lacking hepatocyte growth factor/scatter factor. Nature
373: 702–705,

[175766]1995.

[175767]

[175768]

[175769]18053.

[175770]Weidner, K. M.; Arakaki, N.; Hartmann, G.; Vandekerck-
hove, J.;

[175771]Weingart, S.; Rieder, H.; Fonatsch, C.; Tsubouchi, H.;
Hishida, T.;

[175772]Daikuhara, Y.; Birchmeier, W.: Evidence for the identity of
human

[175773]scatter factor and human hepatocyte growth factor. Proc.
Nat. Acad.

[175774]Sci. 88: 7001–7005, 1991.

[175775]

[175776]

[175777]18054.

[175778]Zarnegar, R.; Petersen, B.; DeFrances, M. C.; Michalopoulos, G.

[175779]: Localization of hepatocyte growth factor (HGF) gene on
human chromosome

[175780]7. Genomics 12: 147–150, 1992.

[175781]

[175782]

[175783]18055.

[175784]Calandro, L. M.; Baer, D. M.; Sensabaugh, G. F.: Character-
ization

[175785]of a recombinant that locates the hereditary hemochro-
matosis gene

[175786]telomeric to HLA-F. Hum. Genet. 96: 339–342, 1995.

[175787]

[175788]

[175789]18056.

[175790]Geraghty, D. E.; Wei, X. H.; Orr, H. T.; Koller, B. H.: Human

[175791]leukocyte antigen F (HLA-F): an expressed HLA gene composed of a class

[175792]I coding sequence linked to a novel transcribed repetitive element. J.

[175793]Exp. Med. 171: 1-18, 1990.

[175794]

[175795]

[175796]18057.

[175797]Shiina, T.; Tamiya, G.; Oka, A.; Takishima, N.; Yamagata, T.; Kikkawa,

[175798]E.; Iwata, K.; Tomizawa, M.; Okuaki, N.; Kuwano, Y.; Watanabe, K.;

[175799]Fukuzumi, Y.; and 11 others: Molecular dynamics of MHC genesis

[175800]unraveled by sequence analysis of the 1,796,938-bp HLA class I region. Proc.

[175801]Nat. Acad. Sci. 96: 13282-13287, 1999.

[175802]

[175803]

[175804]18058.

[175805]Gasparini, P.; Borgato, L.; Piperno, A.; Girelli, D.; Olivieri,

[175806]O.; Gottardi, E.; Roetto, A.; Dianzani, I.; Fargion, S.; Schi-
naia,

[175807]G.; Cappellini, M. D.; Gandini, G.; Pignatti, P. F.; Fiorelli,
G.;

[175808]De Sandre, G.; Camaschella, C.: Linkage analysis of 6p21
polymorphic

[175809]markers and the hereditary hemochromatosis: localization
of the gene

[175810]centromeric to HLA-F. Hum. Molec. Genet. 2: 571-576,
1993.

[175811]

[175812]

[175813]18059.

[175814]van den Boogaard, M.-J. H.; Dorland, M.; Beemer, F. A.;
Ploos

[175815]van Amstel, H. K.: MSX1 mutation is associated with oro-
facial clefting

[175816]and tooth agenesis in humans. (Letter) Nature Genet. 24:
342–343,

[175817]2000. Note: Erratum: Nature Genet. 25: 125 only, 2000.

[175818]

[175819]

[175820]18060.

[175821]Del Campo, M.; Jones, M. C.; Veraksa, A. N.; Curry, C. J.;
Jones,

[175822]K. L.; Mascarello, J. T.; Ali-Kahn-Catts, Z.; Drumheller, T.;
McGinnis,

[175823]W.: Monodactylous limbs and abnormal genitalia are asso-
ciated with

[175824]hemizygosy for the human 2q31 region that includes the
HOXD cluster. Am.

[175825]J. Hum. Genet. 65: 104–110, 1999.

[175826]

[175827]

[175828]18061.

[175829]Zakany, J.; Kmita, M.; Alarcon, P.; de la Pompa, J.–L.;
Duboule,

[175830]D.: Localized and transient transcription of Hox genes
suggests a

[175831]link between patterning and the segmentation clock. Cell
106: 207–217,

[175832]2001.

[175833]

[175834]

[175835]18062.

[175836]Perez, O. D.; Kinoshita, S.; Hitoshi, Y.; Payan, D. G.; Kita-
mura,

[175837]T.; Nolan, G. P.; Lorens, J. B.: Activation of the PKB/AKT
pathway

[175838]by ICAM–2. Immunity 16: 51–65, 2002.

[175839]

[175840]

[175841]18063.

[175842]Sansom, D.; Borrow, J.; Solomon, E.; Trowsdale, J.: The hu-
man

[175843]ICAM2 gene maps to 17q23–25. Genomics 11: 462–464,
1991.

[175844]

[175845]

[175846]18064.

[175847]Staunton, D. E.; Dustin, M. L.; Springer, T. A.: Functional
cloning

[175848]of ICAM-2, a cell adhesion ligand for LFA-1 homologous
to ICAM-1. Nature 339:

[175849]61-64, 1989.

[175850]

[175851]

[175852]18065.

[175853]Rotem-Yehudar, R.; Galperin, E.; Horowitz, M.: Association
of

[175854]insulin-like growth factor 1 receptor with EHD1 and
SNAP29. J. Biol.

[175855]Chem. 276: 33054-33060, 2001.

[175856]

[175857]

[175858]18066.

[175859]Scott, B. A.; Avidan, M. S.; Crowder, C. M.: Regulation of hypoxic

[175860]death in *C. elegans* by the insulin/IGF receptor homolog DAF-2. *Science* 296:

[175861]2388–2391, 2002.

[175862]

[175863]

[175864]18067.

[175865]Tatar, M.; Kopelman, A.; Epstein, D.; Tu, M.-P.; Yin, C.-M.; Garofalo,

[175866]R. S.: A mutant *Drosophila* insulin receptor homolog that extends

[175867]life-span and impairs neuroendocrine function. *Science* 292: 107–110,

[175868]2001.

[175869]

[175870]

[175871]18068.

[175872]Ullrich, A.; Gray, A.; Tam, A. W.; Yang-Feng, T.; Tsubokawa, M.;

[175873]Collins, C.; Henzel, W.; Le Bon, T.; Kathuria, S.; Chen, E.;
Jacobs,

[175874]S.; Francke, U.; Ramachandran, J.; Fujita–Yamaguchi, Y.:
Insulin–like

[175875]growth factor I receptor primary structure: comparison
with insulin

[175876]receptor suggests structural determinants that define
functional specificity. EMBO

[175877]J. 5: 2503–2512, 1986.

[175878]

[175879]

[175880]18069.

[175881]Werner, H.; Karnieli, E.; Rauscher, F. J., III; LeRoth, D.:
Wild–type

[175882]and mutant p53 differentially regulate transcription of the
insulin–like

[175883]growth factor I receptor gene. Proc. Nat. Acad. Sci. 93:
8318–8323,

[175884]1996.

[175885]

[175886]

[175887]18070.

[175888]Barton, D. E.; Yang-Feng, T. L.; Mason, A. J.; Seeburg, P.
H.;

[175889]Francke, U.: INHA, INHBA and INHBB, the loci for the three
subunits

[175890]of inhibin, mapped in mouse and man. (Abstract) Cyto-
genet. Cell Genet. 46:

[175891]578 only, 1987.

[175892]

[175893]

[175894]18071.

[175895]Barton, D. E.; Yang-Feng, T. L.; Mason, A. J.; Seeburg, P.
H.;

[175896]Francke, U.: Mapping of genes for inhibin subunits alpha,
beta(A)

[175897]and beta(B) on human and mouse chromosomes and
studies of jsd mice. Genomics 5:

[175898]91-99, 1989.

[175899]

[175900]

[175901]18072.

[175902]Bremner, W. J.: Inhibin: from hypothesis to clinical application.

[175903](Editorial) New Eng. J. Med. 321: 826–827, 1989.

[175904]

[175905]

[175906]18073.

[175907]Lappohn, R. E.; Burger, H. G.; Bouma, J.; Bangah, M.; Krans, M.;

[175908]de Bruijn, H. W. A.: Inhibin as a marker for granulosa-cell tumors. New

[175909]Eng. J. Med. 321: 790–793, 1989.

[175910]

[175911]

[175912]18074.

[175913]Mason, A. J.; Hayflick, J. S.; Ling, N.; Esch, F.; Ueno, N.; Ying,

[175914]S.-Y.; Guillemin, R.; Niall, H.; Seeburg, P. H.: Complementary DNA

[175915]sequences of ovarian follicular fluid inhibin show precur-

sor structure

[175916]and homology with transforming growth factor-beta. Nature 318: 659-663,

[175917]1985.

[175918]

[175919]

[175920]18075.

[175921]Mason, A. J.; Niall, H. D.; Seeburg, P. H.: Structure of two human

[175922]ovarian inhibins. Biochem. Biophys. Res. Commun. 135: 957-964, 1986.

[175923]

[175924]

[175925]18076.

[175926]Matzuk, M. M.; Finegold, M. J.; Mather, J. P.; Krummen, L.; Lu,

[175927]H.; Bradley, A.: Development of cancer cachexia-like syndrome and

[175928]adrenal tumors in inhibin-deficient mice. Proc. Nat. Acad. Sci. 91:

[175929]8817-8821, 1994.

[175930]

[175931]

[175932]18077.

[175933]Matzuk, M. M.; Finegold, M. J.; Su, J.-G. J.; Hsueh, A. J. W.;

[175934]Bradley, A.: Alpha-inhibin is a tumour-suppressor gene
with gonadal

[175935]specificity in mice. Nature 360: 313-319, 1992.

[175936]

[175937]

[175938]18078.

[175939]Mellor, S. L.; Richards, M. G.; Pedersen, J. S.; Robertson, D.

[175940]M.; Risbridger, G. P.: Loss of the expression and localization of

[175941]inhibin alpha-subunit in high grade prostate cancer. J.
Clin. Endocr.

[175942]Metab. 83: 969-975, 1998.

[175943]

[175944]

[175945]18079.

[175946]Meunier, H.; Rivier, C.; Evans, R. M.; Vale, W.: Gonadal and
[175947]extragonadal expression of inhibin alpha, beta-A and
beta-B subunits
[175948]in various tissues predicts diverse functions. Proc. Nat.
Acad. Sci. 85:
[175949]247-251, 1988.

[175950]

[175951]

[175952]18080.

[175953]Ramasharma, K.; Li, C. H.: Characteristics of binding of
human
[175954]seminal alpha-inhibin-92 to human pituitary membranes.
Proc. Nat.
[175955]Acad. Sci. 84: 3595-3598, 1987.

[175956]

[175957]

[175958]18081.

[175959]Schmitt, J. F.; Millar, D. S.; Pedersen, J. S.; Clark, S. L.;

[175960]Venter, D. J.; Frydenberg, M.; Molloy, P. L.; Risbridger, G.

P.:

[175961]Hypermethylation of the inhibin alpha-subunit gene in prostate carcinoma. Molec.

[175962]Endocr. 16: 213–220, 2002.

[175963]

[175964]

[175965]18082.

[175966]Thirunavukarasu, P.; Stephenson, T.; Forray, J.; Stanton, P. G.;

[175967]Groome, N.; Wallace, E.; Robertson, D. M.: Changes in molecular weight

[175968]forms of inhibin A and pro-alpha-C in maternal serum during human

[175969]pregnancy. J. Clin. Endocr. Metab. 86: 5794–5804, 2001.

[175970]

[175971]

[175972]18083.

[175973]Lafuse, W. P.; Zwillling, B. S.: Localization of the inhibin beta-B

[175974]gene on mouse chromosome 1. Mammalian Genome 4:

399–400, 1993.

[175975]

[175976]

[175977]18084.

[175978]Salmenkivi, K.; Arola, J.; Voutilainen, R.; Ilvesmaki, V.;
Haglund,

[175979]C.; Kahri, A. I.; Heikkila, P.; Liu, J.: Inhibin/activin beta-
B-subunit

[175980]expression in pheochromocytomas favors benign diagnosis. J. Clin.

[175981]Endocr. Metab. 86: 2231–2235, 2001.

[175982]

[175983]

[175984]18085.

[175985]Schrewe, H.; Gendron-Maguire, M.; Harbison, M. L.; Grid-
ley, T.

[175986]: Mice homozygous for a null mutation of activin beta B
are viable

[175987]and fertile. Mech. Dev. 47: 43–51, 1994.

[175988]

[175989]

[175990]18086.

[175991]Nicoll, J. A. R.; Mrak, R. E.; Graham, D. I.; Stewart, J.;
Wilcock,

[175992]G.; MacGowan, S.; Esiri, M. M.; Murray, L. S.; Dewar, D.;
Love, S.;

[175993]Moss, T.; Griffin, W. S. T.: Association of interleukin-1
gene polymorphisms

[175994]with Alzheimer's disease. Ann. Neurol. 47: 365-368,
2000.

[175995]

[175996]

[175997]18087.

[175998]Rogers, J.: An IL-1-alpha susceptibility polymorphism in
Alzheimer's

[175999]disease: new fuel for the inflammation hypothesis.

(Editorial) Neurology 55:

[176000]464-465, 2000.

[176001]

[176002]

[176003]18088.

[176004]Sabatino, M.; Boyce, B.; Aufdemorte, T.; Bonewald, L.;
Mundy,

[176005]G. R.: Infusions of recombinant human interleukins 1 al-
pha and 1

[176006]beta cause hypercalcemia in normal mice. Proc. Nat. Acad.
Sci. 85:

[176007]5235–5239, 1988.

[176008]

[176009]

[176010]18089.

[176011]Silver, A. R. J.; Masson, W. K.; George, A. M.; Adam, J.;
Cox,

[176012]R.: The Il–1 alpha and beta genes are closely linked (less
than 70

[176013]kb) on mouse chromosome 2. Somat. Cell Molec. Genet.
16: 549–556,

[176014]1990.

[176015]

[176016]

[176017]18090.

[176018]Arai, N.; Nomura, D.; Villaret, D.; DeWaal Malefijt, R.; Seiki,

[176019]M.; Yoshida, M.; Minoshima, S.; Fukuyama, R.; Maekawa,
M.; Kudoh,

[176020]J.; Shimizu, N.; Yokota, K.; Abe, E.; Yokota, T.; Takebe, Y.;
Arai,

[176021]K.: Complete nucleotide sequence of the chromosomal
gene for human

[176022]IL-4 and its expression. J. Immun. 142: 274-282, 1989.

[176023]

[176024]

[176025]18091.

[176026]Carvalho, L. H.; Sano, G.; Hafalla, J. C. R.; Morrot, A.;
Curotto

[176027]de Lafaille, M. A.; Zavala, F.: IL-4-secreting CD4+ T cells
are crucial

[176028]to the development of CD8+ T-cell responses against
malaria liver

[176029]stages. Nature Med. 8: 166-170, 2002.

[176030]

[176031]

[176032]18092.

[176033]Dickensheets, H. L.; Venkataraman, C.; Schindler, U.; Donnelly,

[176034]R. P.: Interferons inhibit activation of STAT6 by interleukin 4 in

[176035]human monocytes by inducing SOCS-1 gene expression. Proc. Nat. Acad.

[176036]Sci. 96: 10800-10805, 1999.

[176037]

[176038]

[176039]18093.

[176040]Eder, A.; Krafft-Czepa, H.; Krammer, P. H.: The 5-prime region

[176041]of the human interleukin 4 gene: structure and potential regulatory

[176042]elements. Nucleic Acids Res. 16: 772 only, 1988.

[176043]

[176044]

[176045]18094.

- [176046]Hunt, P. J.; Marshall, S. E.; Weetman, A. P.; Bell, J. I.; Wass,
[176047]J. A. H.; Welsh, K. I.: Cytokine gene polymorphisms in au-
toimmune
[176048]thyroid disease. J. Clin. Endocr. Metab. 85: 1984–1988,
2000.
- [176049]
- [176050]
- [176051]18095.
- [176052]Inoue, T.; Kira, R.; Nakao, F.; Ihara, K.; Bassuny, W. M.;
Kusuhara,
[176053]K.; Nihei, K.; Takeshita, K.; Hara, T.: Contribution of the
interleukin
[176054]4 gene to susceptibility to subacute sclerosing panen-
cephalitis. Arch.
[176055]Neurol. 59: 822–827, 2002.
- [176056]
- [176057]
- [176058]18096.
- [176059]Kotanides, H.; Reich, N. C.: Interleukin–4–induced STAT6
recognizes

[176060]and activates a target site in the promoter of the inter-
leukin-4 receptor

[176061]gene. J. Biol. Chem. 271: 25555-25561, 1996.

[176062]

[176063]

[176064]18097.

[176065]Le Beau, M. M.; Lemons, R. S.; Espinosa, R., III; Larson, R.
A.;

[176066]Arai, N.; Rowley, J. D.: Interleukin-4 and interleukin-5
map to human

[176067]chromosome 5 in a region encoding growth factors and
receptors and

[176068]are deleted in myeloid leukemias with a del(5q). Blood 73:
647-650,

[176069]1989.

[176070]

[176071]

[176072]18098.

[176073]Lewis, D. B.; Liggitt, H. D.; Effmann, E. L.; Motley, S. T.;
Teitelbaum,

[176074]S. L.; Jepsen, K. J.; Goldstein, S. A.; Bonadio, J.; Carpenter, J.;

[176075]Perlmutter, R. M.: Osteoporosis induced in mice by over-production

[176076]of interleukin 4. Proc. Nat. Acad. Sci. 90: 11618–11622, 1993.

[176077]

[176078]

[176079]18099.

[176080]Saltman, D. L.; Dolganov, G. M.; Warrington, J. A.; Wasmuth, J.

[176081]J.; Lovett, M.: A physical map of 15 loci on human chromosome 5q23–q33

[176082]by two-color fluorescence in situ hybridization. Genomics 16: 726–732,

[176083]1993.

[176084]

[176085]

[176086]18100.

[176087]Takahashi, M.; Yoshida, M. C.; Satoh, H.; Hilgers, J.; Yaoita,

[176088]Y.; Honjo, T.: Chromosomal mapping of the mouse IL-4
and human IL-5

[176089]genes. Genomics 4: 47-52, 1989.

[176090]

[176091]

[176092]18101.

[176093]Yokota, T.; Otsuka, T.; Mosmann, T.; Banchereau, J.; De-
France,

[176094]T.; Blanchard, D.; De Vries, J. E.; Lee, F.; Arai, K.: Isolation

[176095]and characterization of a human interleukin cDNA clone,
homologous

[176096]to mouse B-cell stimulatory factor 1, that expresses B-
cell- and T-cell-stimulating

[176097]activities. Proc. Nat. Acad. Sci. 83: 5894-5898, 1986.

[176098]

[176099]

[176100]18102.

[176101]Caggana, M.; Walker, K.; Reilly, A. A.; Conroy, J. M.; Duva,
S.;

[176102]Walsh, A. C.: Population-based studies reveal differences

in the

[176103]allelic frequencies of two functionally significant human
interleukin-4

[176104]receptor polymorphisms in several ethnic groups. Genet.
Med. 1:

[176105]267-271, 1999.

[176106]

[176107]

[176108]18103.

[176109]Deichmann, K. A.; Heinzmann, A.; Forster, J.; Dischinger,
S.; Mehl,

[176110]C.; Brueggenolte, E.; Hildebrandt, F.; Moseler, M.; Kuehr,
J.: Linkage

[176111]and allelic association of atopy and markers flanking the
IL4-receptor

[176112]gene. Clin. Exp. Allergy 28: 151-155, 1998.

[176113]

[176114]

[176115]18104.

[176116]MacLeod, J. N.; Lee, A. K.; Liebhaber, S. A.; Cooke, N. E.:

Developmental

[176117]control and alternative splicing of the placentally expressed transcripts

[176118]from the human growth hormone gene cluster. J. Biol. Chem. 267:

[176119]14219–14226, 1992.

[176120]

[176121]

[176122]18105.

[176123]Wurzel, J. M.; Parks, J. S.; Herd, J. E.; Nielsen, P. V.: A gene

[176124]deletion is responsible for absence of human chorionic somatomammotropin. DNA 1:

[176125]251–257, 1982.

[176126]

[176127]

[176128]18106.

[176129]Gallin, J. I.: Personal Communication. Bethesda, Md.
10/12/1990.

[176130]

[176131]

[176132]18107.

[176133]McCombs, J. L.; Teng, C. T.; Pentecost, B. T.; Magnuson, V. L.;

[176134]Moore, C. M.; McGill, J. R.: Chromosomal localization of human lactotransferrin

[176135]gene (LTF) by in situ hybridization. Cytogenet. Cell Genet. 47:

[176136]16–17, 1988.

[176137]

[176138]

[176139]18108.

[176140]Moriuchi, M.; Moriuchi, H.: A milk protein lactoferrin enhances

[176141]human T cell leukemia virus type I and suppresses HIV–1 infection. J.

[176142]Immun. 166: 4231–4236, 2001.

[176143]

[176144]

[176145]18109.

- [176146]Naylor, S. L.; Marshall, A.; Solomon, A.; McGill, J. R.; Mc-Combs,
[176147]J.; Magnuson, V. L.; Moore, C. M.; Lalley, P. A.; Pentecost,
B. T.;
- [176148]Teng, C.: Lactoferrin maps to human chromosome
3(q21-q23) and mouse
[176149]chromosome 9. (Abstract) Cytogenet. Cell Genet. 46: 669
only, 1987.
- [176150]
- [176151]
- [176152]18110.
- [176153]Powell, M. J.; Ogden, J. E.: Nucleotide sequence of human
lactoferrin
[176154]cDNA. Nucleic Acids Res. 18: 4013 only, 1990.
- [176155]
- [176156]
- [176157]18111.
- [176158]Qiu, J.; Hendrixson, D. R.; Baker, E. N.; Murphy, T. F.; St.
Geme,
[176159]J. W., III; Plaut, A. G.: Human milk lactoferrin inactivates
two

[176160]putative colonization factors expressed by *Haemophilus influenzae*. Proc.

[176161]Nat. Acad. Sci. 95: 12641–12646, 1998.

[176162]

[176163]

[176164]18112.

[176165]Rado, T. A.; Wei, X.; Benz, E. J., Jr.: Isolation of lactoferrin

[176166]cDNA from a human myeloid library and expression of mRNA during normal

[176167]and leukemic myelopoiesis. Blood 70: 989–993, 1987.

[176168]

[176169]

[176170]18113.

[176171]Singh, P. K.; Parsek, M. R.; Greenberg, E. P.; Welsh, M. J.: A

[176172]component of innate immunity prevents bacterial biofilm development. Nature 417:

[176173]552–555, 2002.

[176174]

[176175]

[176176]18114.

[176177]Teng, C. T.; Pentecost, B. T.; Marshall, A.; Solomon, A.;
Bowman,

[176178]B. H.; Lalley, P. A.; Naylor, S. L.: Assignment of the lacto-
transferrin

[176179]gene to human chromosome 3 and to mouse chromosome
9. Somat. Cell

[176180]Molec. Genet. 13: 689–693, 1987.

[176181]

[176182]

[176183]18115.

[176184]Yang, F.; Lum, J.; Baldwin, W. D.; Brune, J. L.; van Bragt, P.;

[176185]Bowman, B. H.: Genetic analysis of human iron binding
glycoproteins.

[176186](Abstract) Am. J. Hum. Genet. 35: 184A only, 1983.

[176187]

[176188]

[176189]18116.

[176190]Nonaka, S.; Shiratori, H.; Saijoh, Y.; Hamada, H.: Determi-

nation

[176191]of left–right patterning of the mouse embryo by artificial
nodal flow. Nature 418:

[176192]96–99, 2002.

[176193]

[176194]

[176195]18117.

[176196]Tanaka, A. R.; Ikeda, Y.; Abe–Dohmae, S.; Arakawa, R.;
Sadanami,

[176197]K.; Kidera, A.; Nakagawa, S.; Nagase, T.; Aoki, R.; Kioka,
N.; Amachi,

[176198]T.; Yokoyama, S.; Ueda, K.: Human ABCA1 contains a large
amino terminal

[176199]extracellular domain homologous to an epitope of Sjo–
gren's syndrome. Biochem.

[176200]Biophys. Res. Commun. 283: 1019–1025, 2001.

[176201]

[176202]

[176203]18118.

[176204]Xu, W.; Gelber, S.; Orr–Urtreger, A.; Armstrong, D.; Lewis,

R.

[176205]A.; Ou, C.-N.; Patrick, J.; Role, L.; De Biasi, M.; Beaudet, A.

L.

[176206]: Megacystis, mydriasis, and ion channel defect in mice
lacking the

[176207]alpha-3 neuronal nicotinic acetylcholine receptor. Proc.
Nat. Acad.

[176208]Sci. 96: 5746–5751, 1999.

[176209]

[176210]

[176211]18119.

[176212]Gallardo, M. E.; Lopez-Rios, J.; Fernaud-Espinosa, I.;
Granadino,

[176213]B.; Sanz, R.; Ramos, C.; Ayuso, C.; Seller, M. J.; Brunner, H.
G.;

[176214]Bovolenta, P.; Rodriguez de Cordoba, S.: Genomic cloning
and characterization

[176215]of the human homeobox gene SIX6 reveals a cluster of SIX
genes in

[176216]chromosome 14 and associates SIX6 hemizyosity with
bilateral anophthalmia

[176217]and pituitary anomalies. Genomics 61: 82–91, 1999.

[176218]

[176219]

[176220]18120.

[176221]Sun, X.-H.: Constitutive expression of the Id1 gene impairs mouse

[176222]B cell development. Cell 79: 893-900, 1994.

[176223]

[176224]

[176225]18121.

[176226]Nowell, P. C.; Hungerford, D. A.: Chromosome studies on normal

[176227]and leukemic human leukocytes. J. Nat. Cancer Inst. 25: 85-109,

[176228]1960.

[176229]

[176230]

[176231]18122.

[176232]Dowbenko, D. J.; Diep, A.; Taylor, B. A.; Lysis, A. J.; Lasky,

[176233]L. A.: Characterization of the murine homing receptor

gene reveals

[176234]correspondence between protein domains and coding ex-
ons. Genomics 9:

[176235]270–277, 1991.

[176236]

[176237]

[176238]18123.

[176239]Couturier, J.; Sastre–Garau, X.; Schneider–Maunoury, S.;
Labib,

[176240]A.; Orth, G.: Integration of papillomavirus DNA near myc
genes in

[176241]genital carcinomas and its consequences for proto-
oncogene expression. J.

[176242]Virol. 65: 4534–4538, 1991.

[176243]

[176244]

[176245]18124.

[176246]Cooper, P. K.; Nospikel, T.; Clarkson, S. G.; Leadon, S. A.:

[176247]Defective transcription–coupled repair of oxidative base
damage in

[176248]Cockayne syndrome patients from XP group G. Science
275: 990–993,

[176249]1997.

[176250]

[176251]

[176252]18125.

[176253]Gersen, S.; Warburton, D.; Jackson, C. L.; Housman, D.:
Regional

[176254]localization of the excision repair gene ERCC5 on chro-
mosome 13. (Abstract) Cytogenet.

[176255]Cell Genet. 51: 1003 only, 1989.

[176256]

[176257]

[176258]18126.

[176259]Habraken, Y.; Sung, P.; Prakash, L.; Prakash, S.: Human
xeroderma

[176260]pigmentosum group G gene encodes a DNA endonucle-
ase. Nucleic Acids

[176261]Res. 22: 3312–3316, 1994.

[176262]

[176263]

[176264]18127.

[176265]Habraken, Y.; Sung, P.; Prakash, S.; Prakash, L.: Transcription

[176266]factor TFIIH and DNA endonuclease Rad2 constitute yeast nucleotide

[176267]excision repair factor 3: implications for nucleotide excision repair

[176268]and Cockayne syndrome. Proc. Nat. Acad. Sci. 93: 10718–10722, 1996.

[176269]

[176270]

[176271]18128.

[176272]Hamel, B. C. J.; Raams, A.; Schuitema–Dijkstra, A. R.; Simons,

[176273]P.; van der Burgt, I.; Jaspers, N. G. J.; Kleijer, W. J.: Xeroderma

[176274]pigmentosum–Cockayne syndrome complex: a further case. J. Med. Genet. 33:

[176275]607–610, 1996.

[176276]

[176277]

[176278]18129.

[176279]Harada, Y.-N.; Matsuda, Y.; Shiomi, N.; Shiomi, T.: Complementary

[176280]DNA sequence and chromosomal localization of xpg, the mouse counterpart

[176281]of human repair gene XPG/ERCC5. Genomics 28: 59–65, 1995.

[176282]

[176283]

[176284]18130.

[176285]Hori, T.; Shiomi, T.; Sato, K.: Human chromosome 13 compensates

[176286]a DNA repair defect in UV-sensitive mouse cells by mouse-human cell

[176287]hybridization. Proc. Nat. Acad. Sci. 80: 5655–5659, 1983.

[176288]

[176289]

[176290]18131.

[176291]Jaeken, J.; Klocker, H.; Schwaiger, H.; Bellmann, R.; Hirsch-

Kauffmann,

[176292]M.; Schweiger, M.: Clinical and biochemical studies in
three patients

[176293]with severe early infantile Cockayne syndrome. Hum.
Genet. 83: 339–346,

[176294]1989.

[176295]

[176296]

[176297]18132.

[176298]Lee, S.–K.; Yu, S.–L.; Prakash, L.; Prakash, S.: Requirement
[176299]of yeast RAD2, a homolog of human XPG gene, for effi-
cient RNA polymerase

[176300]II transcription: implications for Cockayne syndrome. Cell
109:

[176301]823–834, 2002.

[176302]

[176303]

[176304]18133.

[176305]Lehmann, A. R.; Bootsma, D.; Clarkson, S. G.; Cleaver, J.
E.;

[176306]McAlpine, P. J.; Tanaka, K.; Thompson, L. H.; Wood, R. D.:
Nomenclature

[176307]of human DNA repair genes. Mutat. Res. 315: 41–42,
1994.

[176308]

[176309]

[176310]18134.

[176311]MacInnes, M. A.; Dickson, J. A.; Hernandez, R. R.; Lear-
month,

[176312]D.; Lin, G. Y.; Mudgett, J. S.; Park, M. S.; Schauer, S.;
Reynolds,

[176313]R. J.; Strniste, G. F.; Yu, J. Y.: Human ERCC5 cDNA–cosmid
complementation

[176314]for excision repair and bipartite amino acid domains con-
served with

[176315]RAD proteins of *Saccharomyces cerevisiae* and *Schizosac-*
charomyces pombe. Molec.

[176316]Cell. Biol. 13: 6393–6402, 1993.

[176317]

[176318]

[176319]18135.

- [176320]Mudgett, J. S.; MacInnes, M. A.: Isolation of the functional
- [176321]human excision repair gene ERCC5 by intercosmid recombination. Genomics 8:
- [176322]623–633, 1990.
- [176323]
- [176324]
- [176325]18136.
- [176326]Norris, P. G.; Hawk, J. L. M.; Avery, J. A.; Giannelli, F.: Xeroderma
- [176327]pigmentosum complementation group G--report of two cases. Brit. J.
- [176328]Derm. 116: 861–866, 1987.
- [176329]
- [176330]
- [176331]18137.
- [176332]Noussipikel, T.; Clarkson, S. G.: Mutations that disable the
- DNA
- [176333]repair gene XPG in a xeroderma pigmentosum group G patient. Hum.
- [176334]Molec. Genet. 3: 963–967, 1994.

[176335]

[176336]

[176337]18138.

[176338]Noussipiel, T.; Lalle, P.; Leadon, S. A.; Cooper, P. K.; Clarkson,
son,

[176339]S. G.: A common mutational pattern in Cockayne syndrome patients
drome patients

[176340]from xeroderma pigmentosum group G: implications for a
second XPG

[176341]function. Proc. Nat. Acad. Sci. 94: 3116–3121, 1997.

[176342]

[176343]

[176344]18139.

[176345]O'Donovan, A.; Davies, A. A.; Moggs, J. G.; West, S. C.;
Wood,

[176346]R. D.: XPG endonuclease makes the 3–prime incision in
human DNA nucleotide

[176347]excision repair. Nature 371: 432–435, 1994.

[176348]

[176349]

[176350]18140.

[176351]O'Donovan, A.; Scherly, D.; Clarkson, S. G.; Wood, R. D.:

Isolation

[176352]of active recombinant XPG protein, a human DNA repair
endonuclease. J.

[176353]Biol. Chem. 269: 15965–15968, 1994.

[176354]

[176355]

[176356]18141.

[176357]O'Donovan, A.; Wood, R. D.: Identical defects in DNA re-
pair in

[176358]xeroderma pigmentosum group G and rodent ERCC group
5. Nature 363:

[176359]185–188, 1993.

[176360]

[176361]

[176362]18142.

[176363]Samec, S.; Jones, T. A.; Corlet, J.; Scherly, D.; Sheer, D.;
Wood,

[176364]R. D.; Clarkson, S. G.: The human gene for xeroderma

pigmentosum

[176365]complementation group G (XPG) maps to 13q33 by fluorescence in situ

[176366]hybridization. Genomics 21: 283–285, 1994.

[176367]

[176368]

[176369]18143.

[176370]Scherly, D.; Nospikel, T.; Corlet, J.; Ucla, C.; Bairoch, A.;

[176371]Clarkson, S. G.: Complementation of the DNA repair defect in xeroderma

[176372]pigmentosum group G cells by a human cDNA related to yeast RAD2. Nature 363:

[176373]182–185, 1993.

[176374]

[176375]

[176376]18144.

[176377]Abruzzo, L. V.; Jaffe, E. S.; Cotelingam, J. D.; Whang-Peng, J.;

[176378]Del Duca, V., Jr.; Medeiros, L. J.: T-cell lymphoblastic lymphoma

[176379]with eosinophilia associated with subsequent myeloid malignancy. Am.

[176380]J. Surg. Path. 16: 236–245, 1992.

[176381]

[176382]

[176383]18145.

[176384]Chesi, M.; Nardini, E.; Brents, L. A.; Schrock, E.; Ried, T.; Kuehl,

[176385]W. M.; Bergsagel, P. L.: Frequent translocation
t(4;14)(p16.3;q32.3)

[176386]in multiple myeloma is associated with increased expression and activating

[176387]mutations of fibroblast growth factor receptor 3. Nature Genet. 16:

[176388]260–264, 1997.

[176389]

[176390]

[176391]18146.

[176392]Demiroglu, A.; Steer, E. J.; Heath, C.; Taylor, K.; Bentley, M.;

- [176393] Allen, S. L.; Koduru, P.; Brody, J. P.; Hawson, G.; Rodwell, R.; Doody, M.-L.; Carnicero, F.; Reiter, A.; Goldman, J. M.; Melo, J. V.; Cross, N. C. P.: The t(8;22) in chronic myeloid leukemia fuses BCR to FGFR1: transforming activity and specific inhibition of FGFR1 fusion proteins. Blood 98: 3778–3783, 2001.
- [176397]
- [176398]
- [176399]
- [176400] 18147.
- [176401] Guasch, G.; Mack, G. J.; Popovici, C.; Dastugue, N.; Birnbaum, D.; Rattner, J. B.; Pebusque, M.-J.: FGFR1 is fused to the centrosome-associated protein CEP110 in the 8p12 stem cell myeloproliferative disorder with t(8;9)(p12;q33). Blood 95: 1788–1796, 2000.
- [176404]
- [176405]
- [176406]

[176407]18148.

[176408]Baroni, M. G.; Oelbaum, R. S.; Pozzilli, P.; Stocks, J.; Li, S.-R.;

[176409]Fiore, V.; Galton, D. J.: Polymorphisms at the GLUT1 (HepG2) and

[176410]GLUT4 (muscle/adipocyte) glucose transporter genes and non-insulin-dependent

[176411]diabetes mellitus (NIDDM). Hum. Genet. 88: 557-561, 1992.

[176412]

[176413]

[176414]18149.

[176415]Kelley, M. J.; Pech, M.; Seuanez, H. N.; Rubin, J. S.; O'Brien,

[176416]S. J.; Aaronson, S. A.: Emergence of the keratinocyte growth factor

[176417]multigene family during the great ape radiation. Proc. Nat. Acad. Acad.

[176418]Sci. 89: 9287-9291, 1992.

[176419]

[176420]

[176421]18150.

[176422]Mattei, M.-G.; deLapeyriere, O.; Bresnick, J.; Dickson, C.;
Birnbaum,

[176423]D.; Mason, I.: Mouse Fgf7 (fibroblast growth factor 7) and
Fgf8 (fibroblast

[176424]growth factor 8) genes map to chromosomes 2 and 19 re-
spectively. Mammalian

[176425]Genome 6: 196-197, 1995.

[176426]

[176427]

[176428]18151.

[176429]Rubin, J. S.; Osada, H.; Finch, P. W.; Taylor, W. G.;
Rudikoff,

[176430]S.; Aaronson, S. A.: Purification and characterization of a
newly

[176431]identified growth factor specific for epithelial cells. Proc.
Nat.

[176432]Acad. Sci. 86: 802-806, 1989.

[176433]

[176434]

[176435]18152.

[176436]Werner, S.; Smola, H.; Liao, X.; Longaker, M. T.; Krieg, T.;
Hofschneider,

[176437]P. H.; Williams, L. T.: The function of KGF in morphogene-
sis of epithelium

[176438]and reepithelialization of wounds. Science 266: 819–822,
1994.

[176439]

[176440]

[176441]18153.

[176442]Zimonjic, D. B.; Kelley, M. J.; Rubin, J. S.; Aaronson, S. A.;

[176443]Popescu, N. C.: Fluorescence in situ hybridization analysis
of keratinocyte

[176444]growth factor gene amplification and dispersion in evolu-
tion of great

[176445]apes and humans. Proc. Nat. Acad. Sci. 94: 11461–11465,
1997.

[176446]

[176447]

[176448]18154.

[176449]Edwards, Y. H.; Parkar, M.; Povey, S.; West, L. F.; Parrington,
ton,

[176450]J. M.; Solomon, E.: Human myosin heavy chain genes assigned to chromosome

[176451]17 using a human cDNA clone as probe. Ann. Hum. Genet.
49: 101–109,

[176452]1985.

[176453]

[176454]

[176455]18155.

[176456]Nadal–Ginard, B.; Mahdavi, V.: Molecular basis of cardiac
performance:

[176457]plasticity of the myocardium generated through protein
isoform switches. J.

[176458]Clin. Invest. 84: 1693–1700, 1989.

[176459]

[176460]

[176461]18156.

[176462]Rappold, G. A.; Vosberg, H.–P.: Chromosomal localization
of a

[176463]human myosin heavy-chain gene by in situ hybridization.

Hum. Genet. 65:

[176464]195–197, 1983.

[176465]

[176466]

[176467]18157.

[176468]Sparkes, R. S.; Zollman, S.; Klisak, I.; Kirchgessner, T. G.;

[176469]Komaromy, M. C.; Mohandas, T.; Schotz, M. C.; Lysis, A. J.:

Human

[176470]genes involved in lipolysis of plasma lipoproteins: mapping of loci

[176471]for lipoprotein lipase to 8p22 and hepatic lipase to 15q21.

Genomics 1:

[176472]138–144, 1987.

[176473]

[176474]

[176475]18158.

[176476]Baumgartner, M.; Holzfeind, P.; Redl, B.: Assignment of the gene

[176477]for human tear prealbumin (LCN1), a member of the

lipocalin superfamily,

[176478]to chromosome 8q24. Cytogenet. Cell Genet. 65:
101–103, 1994.

[176479]

[176480]

[176481]18159.

[176482]Blaker, M.; Kock, K.; Ahlers, C.; Buck, F.; Schmale, H.:
Molecular

[176483]cloning of human von Ebner's gland protein, a member of
the lipocalin

[176484]superfamily highly expressed in lingual salivary glands.
Biochim.

[176485]Biophys. Acta 1172: 131–137, 1993.

[176486]

[176487]

[176488]18160.

[176489]Liu, Y.; Wu, Y.–P.; Wada, R.; Neufeld, E. B.; Mullin, K. A.;
Howard,

[176490]A. C.; Pentchev, P. G.; Vanier, M. T.; Suzuki, K.; Proia, R. L.:

[176491]Alleviation of neuronal ganglioside storage does not im–

prove the clinical

[176492]course of the Niemann–Pick C disease mouse. Hum.

Molec. Genet. 9:

[176493]1087–1092, 2000.

[176494]

[176495]

[176496]18161.

[176497]Ito, M.; Yuan, C.–X.; Malik, S.; Gu, W.; Fondell, J. D.; Yama-
mura,

[176498]S.; Fu, Z.–Y.; Zhang, X.; Qin, J.; Roeder, R. G.: Identity be-
tween

[176499]TRAP and SMCC complexes indicates novel pathways for
the function

[176500]of nuclear receptors and diverse mammalian activators.

Molec. Cell 3:

[176501]361–370, 1999.

[176502]

[176503]

[176504]18162.

[176505]Nagase, T.; Seki, N.; Ishikawa, K.; Tanaka, A.; Nomura, N.:

Prediction

[176506]of the coding sequences of unidentified human genes. V.

The coding

[176507]sequences of 40 new genes (KIAA0161–KIAA0200) deduced by analysis

[176508]of cDNA clones from human cell line KG–1. DNA Res. 3:

17–24, 1996.

[176509]

[176510]

[176511]18163.

[176512]Makino, K.; Kuwahara, H.; Masuko, N.; Nishiyama, Y.; Morisaki,

[176513]T.; Sasaki, J.; Nakao, M.; Kuwano, A.; Nakata, M.; Ushio, Y.; Saya,

[176514]H.: Cloning and characterization of NE–dlg: a novel human homolog

[176515]of the Drosophila discs large (dlg) tumor suppressor protein interacts

[176516]with the APC protein. Oncogene 14: 2425–2433, 1997.

[176517]

[176518]

[176519]18164.

[176520]Stathakis, D. G.; Lee, D.; Bryant, P. J.: DLG3, the gene encoding

[176521]human neuroendocrine Dlg (NE-Dlg), is located within the 1.8-Mb dystonia-parkinsonism

[176522]region at Xq13.1. Genomics 49: 310-313, 1998.

[176523]

[176524]

[176525]18165.

[176526]Maeda, K.; Matsushashi, S.; Hori, K.; Xin, Z.; Mukai, T.; Tabuchi,

[176527]K.; Egashira, M.; Niikawa, N.: Cloning and characterization of a

[176528]novel human gene, TM4SF6, encoding a protein belonging to the transmembrane

[176529]4 superfamily, and mapped to Xq22. Genomics 52: 240-242, 1998.

[176530]

[176531]

[176532]18166.

[176533]Todd, S. C.; Doctor, V. S.; Levy, S.: Sequences and expres-
sion

[176534]of six new members of the tetraspanin/TM4SF family.
Biochim. Biophys.

[176535]Acta 1399: 101–104, 1998.

[176536]

[176537]

[176538]18167.

[176539]Crew, A. J.; Clark, J.; Fisher, C.; Gill, S.; Grimer, R.; Chand,

[176540]A.; Shipley, J.; Gusterson, B. A.; Cooper, C. S.: Fusion of
SYT to

[176541]two genes, SSX1 and SSX2, encoding proteins with homol-
ogy to the Kruppel-associated

[176542]box in human synovial sarcoma. EMBO J. 14: 2333–2340,
1995.

[176543]

[176544]

[176545]18168.

[176546]Vaccari, T.; Beltrame, M.; Ferrari, S.; Bianchi, M. E.: Hmg4,

[176547]a new member of the Hmg1/2 gene family. Genomics 49:

247–252, 1998.

[176548]

[176549]

[176550]18169.

[176551]Wilke, K.; Wiemann, S.; Gaul, R.; Gong, W.; Poustka, A.:

Isolation

[176552]of human and mouse HMG2a cDNAs: evidence for an

HMG2a-specific 3-prime

[176553]untranslated region. Gene 198: 269–274, 1997.

[176554]

[176555]

[176556]18170.

[176557]Anson, D. S.; Blake, D. J.; Winship, P. R.; Birnbaum, D.;

Brownlee,

[176558]G. G.: Nullisomic deletion of the mcf.2 transforming gene

in two

[176559]haemophilia B patients. EMBO J. 7: 2795–2799, 1988.

[176560]

[176561]

[176562]18171.

[176563]Imbeaud, S.; Belville, C.; Messika-Zeitoun, L.; Rey, R.; di
Clemente,

[176564]N.; Josso, N.; Picard, J.-Y.: A 27 base-pair deletion of the
anti-Mullerian

[176565]type II receptor gene is the most common cause of the
persistent Mullerian

[176566]duct syndrome. Hum. Molec. Genet. 5: 1269-1277, 1996.

[176567]

[176568]

[176569]18172.

[176570]Imbeaud, S.; Carre-Eusebe, D.; Rey, R.; Belville, C.; Josso,
N.;

[176571]Picard, J.-Y.: Molecular genetics of the persistent Mullerian
duct

[176572]syndrome: a study of 19 families. Hum. Molec. Genet. 3:
125-131,

[176573]1994.

[176574]

[176575]

[176576]18173.

[176577]Imbeaud, S.; Faure, E.; Lamarre, I.; Mattei, M.-G.; di
Clemente,

[176578]N.; Tizard, R.; Carre-Eusebe, D.; Belville, C.; Tragethon, L.;
Tonkin,

[176579]C.; Nelson, J.; McAuliffe, M.; Bidart, J.-M.; Lababidi, A.;
Josso,

[176580]N.; Cate, R. L.; Picard, J.-V.: Insensitivity to anti-Mullerian
hormone

[176581]due to a mutation in the human anti-Mullerian hormone
receptor. Nature

[176582]Genet. 11: 382-388, 1995.

[176583]

[176584]

[176585]18174.

[176586]Lang-Muritano, M.; Biason-Lauber, A.; Gitzelmann, C.;
Belville,

[176587]C.; Picard, Y.; Schoenle, E. J.: A novel mutation in the anti-
mullerian

[176588]hormone gene as cause of persistent mullerian duct syn-
drome. Europ.

[176589]J. Pediat. 160: 652–654, 2001.

[176590]

[176591]

[176592]18175.

[176593]Picard, J.–Y.; Benarous, R.; Guerrier, D.; Josso, N.; Kahn, A.

[176594]: Cloning and expression of cDNA for anti–mullerian hormone. Proc.

[176595]Nat. Acad. Sci. 83: 5464–5468, 1986.

[176596]

[176597]

[176598]18176.

[176599]Tezcan, I.; Xu, W.; Gurgey, A.; Tuncer, M.; Cetin, M.; Oner, C.;

[176600]Yetgin, S.; Ersoy, F.; Aizencang, G.; Astrin, K. H.; Desnick, R. J.

[176601]: Congenital erythropoietic porphyria successfully treated by allogeneic

[176602]bone marrow transplantation. Blood 92: 4053–4058, 1998.

[176603]

[176604]

[176605]18177.

[176606]Xu, W.; Astrin, K. H.; Desnick, R. J.: Molecular basis of
congenital

[176607]erythropoietic porphyria: mutations in the human uropor-
phyrinogen

[176608]III synthase gene. Hum. Mutat. 7: 187–192, 1996.

[176609]

[176610]

[176611]18178.

[176612]Pennarun, G.; Escudier, E.; Chapelin, C.; Bridoux, A.–M.;
Cacheux,

[176613]V.; Roger, G.; Clement, A.; Goossens, M.; Amselem, S.;
Duriez, B.

[176614]: Loss-of-function mutations in a human gene related to
Chlamydomonas

[176615]reinhardtii dynein IC78 result in primary ciliary dyskinesia.
Am.

[176616]J. Hum. Genet. 65: 1508–1519, 1999.

[176617]

[176618]

[176619]18179.

[176620]Weiss, P.; Tietze, F.; Gahl, W. A.; Seppala, R.; Ashwell, G.:

[176621]Identification of the metabolic defect in sialuria. J. Biol.
Chem. 264:

[176622]17635–17636, 1989.

[176623]

[176624]

[176625]18180.

[176626]Bassi, M. T.; Ramesar, R. S.; Caciotti, B.; Winship, I. M.; De

[176627]Grandi, A.; Riboni, M.; Townes, P. L.; Beighton, P.; Ballabio,
A.;

[176628]Borsani, G.: X-linked late-onset sensorineural deafness
caused by

[176629]a deletion involving OA1 and a novel gene containing WD-
40 repeats. Am.

[176630]J. Hum. Genet. 64: 1604–1616, 1999.

[176631]

[176632]

[176633]18181.

[176634]Converse, P. J.: Personal Communication. Baltimore, Md.
8/24/2001.

[176635]

[176636]

[176637]18182.

[176638]Disteche, C. M.; Dinulos, M. B.; Bassi, M. T.; Elliott, R. W.;

[176639]Rugarli, E. I.: Mapping of the murine Tbl1 gene reveals a
new rearrangement

[176640]between mouse and human X chromosomes. Mammalian
Genome 9: 1062–1064,

[176641]1998.

[176642]

[176643]

[176644]18183.

[176645]Dong, X.; Tsuda, L.; Zavitz, K. H.; Lin, M.; Li, S.; Carthew,
R.

[176646]W.; Zipursky, S. L.: ebi regulates epidermal growth factor
receptor

[176647]signaling pathways in Drosophila. Genes Dev. 13:

954–965, 1999.

[176648]

[176649]

[176650]18184.

[176651]Matsuzawa, S.; Reed, J. C.: Siah–1, SIP, and Ebi collaborate
in

[176652]a novel pathway for beta–catenin degradation linked to
p53 responses. Molec.

[176653]Cell 7: 915–926, 2001.

[176654]

[176655]

[176656]18185.

[176657]Zhang, J.; Kalkum, M.; Chait, B. T.; Roeder, R. G.: The N–
CoR–HDAC3

[176658]nuclear receptor corepressor complex inhibits the JNK
pathway through

[176659]the integral subunit GPS2. Molec. Cell 9: 611–623, 2002.

[176660]

[176661]

[176662]18186.

[176663]Mu, J.; Roach, P. J.: Characterization of human glycogenin-2,

[176664]a self-glucosylating initiator of liver glycogen metabolism. J. Biol.

[176665]Chem. 273: 34850–34856, 1998.

[176666]

[176667]

[176668]18187.

[176669]Mu, J.; Skurat, A. V.; Roach, P. J.: Glycogenin-2, a novel self-glucosylating

[176670]protein involved in liver glycogen biosynthesis. J. Biol. Chem. 272:

[176671]27589–27597, 1997.

[176672]

[176673]

[176674]18188.

[176675]Delbridge, M. L.; Lingenfelter, P. A.; Disteché, C. M.; Marshall

[176676]Graves, J. A.: The candidate spermatogenesis gene RBMY

has a homologue

[176677]on the human X chromosome. (Letter) Nature Genet. 22:
223–224, 1999.

[176678]

[176679]

[176680]18189.

[176681]Le Coniat, M.; Soulard, M.; Della Valle, V.; Larsen, C.–J.;
Berger,

[176682]R.: Localization of the human gene encoding heteroge-
neous nuclear

[176683]RNA ribonucleoprotein G (hnRNP–G) to chromosome
6p12. Hum. Genet. 88:

[176684]593–595, 1992.

[176685]

[176686]

[176687]18190.

[176688]Mazeyrat, S.; Saut, N.; Mattei, M.–G.; Mitchell, M. J.: RBMY
evolved

[176689]on the Y chromosome from a ubiquitously transcribed X–Y
identical

[176690]gene. (Letter) Nature Genet. 22: 224–226, 1999.

[176691]

[176692]

[176693]18191.

[176694]Venables, J. P.; Elliott, D. J.; Makarova, O. V.; Makarov, E. M.;

[176695]Cooke, H. J.; Eperon, I. C.: RBMY, a probable human spermatogenesis

[176696]factor, and other hnRNP G proteins interact with Tra2-beta and affect

[176697]splicing. Hum. Molec. Genet. 9: 685-694, 2000.

[176698]

[176699]

[176700]18192.

[176701]Vogt, P. H.; Affara, N.; Davey, P.; Hammer, M.; Jobling, M. A.;

[176702]Lau, Y. F.; Mitchell, M.; Schempp, W.; Tyler-Smith, C.; Williams,

[176703]G.; Yen, P.; Rappold, G. A: Report of the Third International Workshop

[176704]on Y Chromosome Mapping 1997. Heidelberg, Germany, April 13-16, 1997. Cytogenet.

[176705]Cell Genet. 79: 1–20, 1997.

[176706]

[176707]

[176708]18193.

[176709]Wilcken, B.; Don, N.; Greenaway, R.; Hammond, J.; Sosula,
L.:

[176710]Sialuria: a second case. J. Inherit. Metab. Dis. 10: 97–102,
1987.

[176711]

[176712]

[176713]18194.

[176714]Sutherland, G. R.; Baker, E.; Callen, D. F.; Hyland, V. J.;
Wong,

[176715]G.; Clark, S.; Jones, S. S.; Eglinton, L. K.; Shannon, M. F.;
Lopez,

[176716]A. F.; Vadas, M. A.: Interleukin 4 is at 5q31 and interleukin
6 is

[176717]at 7p15. Hum. Genet. 79: 335–337, 1988.

[176718]

[176719]

[176720]18195.

[176721]Symmons, D. P.; Jones, M.; Osborne, J.; Sills, J.; South-
wood,

[176722]T. R.; Woo, P.: Pediatric rheumatology in the United King-
dom: data

[176723]from the British Pediatric Rheumatology Group National
Diagnostic

[176724]Register. J. Rheum. 23: 1975–1980, 1996.

[176725]

[176726]

[176727]18196.

[176728]Tosato, G.; Seamon, K. B.; Goldman, N. D.; Sehgal, P. B.;
May,

[176729]L. T.; Washington, G. C.; Jones, K. D.; Pike, S. E.: Mono-
cyte-derived

[176730]human B-cell growth factor identified as interferon-
beta-2 (BSF-2,

[176731]IL-6). Science 239: 502–504, 1988.

[176732]

[176733]

[176734]18197.

[176735]Villuendas, G.; San Millan, J. L.; Sancho, J. and Escobar-Morreale,

[176736]H. F.: The -597 G-A and -174 G-C polymorphisms in the promoter of

[176737]the IL-6 gene are associated with hyperandrogenism. J. Clin. Endocr.

[176738]Metab. 87: 1134-1141, 2002.

[176739]

[176740]

[176741]18198.

[176742]Zilberstein, A.; Ruggieri, R.; Korn, J. H.; Revel, M.: Structure

[176743]and expression of cDNA and genes for human interferon-beta-2, a distinct

[176744]species inducible by growth-stimulatory cytokines. EMBO J. 5: 2529-2537,

[176745]1986.

[176746]

[176747]

[176748]18199.

[176749]Gaedigk, R.; Duncan, A. M. V.; Miyazaki, I.; Robinson, B.
H.; Dosch,

[176750]H.-M.: ICA1 encoding p69, a protein linked to the devel-
opment of

[176751]type 1 diabetes, maps to human chromosome 7p22. Cy-
togenet. Cell Genet. 66:

[176752]274-276, 1994.

[176753]

[176754]

[176755]18200.

[176756]Gaedigk, R.; Karges, W.; Hui, M. F.; Scherer, S. W.; Dosch,
H.-M.

[176757]: Genomic organization and transcript analysis of ICAp69,
a target

[176758]antigen in diabetic autoimmunity. Genomics 38: 382-391,
1996.

[176759]

[176760]

[176761]18201.

[176762]Pietropaolo, M.; Castano, L.; Babu, S.; Buelow, R.; Kuo, Y.-L.

[176763]S.; Martin, S.; Martin, A.; Powers, A. C.; Prochazka, M.; Naggert,

[176764]J.; Leiter, E. H.; Eisenbarth, G. S.: Islet cell autoantigen 69 kD

[176765](ICA69): molecular cloning and characterization of a novel diabetes-associated

[176766]autoantigen. J. Clin. Invest. 92: 359-371, 1993.

[176767]

[176768]

[176769]18202.

[176770]Cavalieri, R. L.; Havell, E. A.; Vilcek, J.; Pestka, S.: Synthesis
[176771]of human interferon by *Xenopus laevis* oocytes: two structural genes

[176772]for interferons in human cells. Proc. Nat. Acad. Sci. 74: 3287-3291,

[176773]1977.

[176774]

[176775]

[176776]18203.

[176777]Chany, C.; Finaz, C.; Weil, D.; Vignal, M.; Van Cong, N.; de Grouchy,

[176778]J.: Investigations on the chromosomal localizations of the human

[176779]and chimpanzee interferon genes: possible role of chromosomes 9 and

[176780]13. Ann. Genet. 23: 201–207, 1980.

[176781]

[176782]

[176783]18204.

[176784]Derynck, R.; Content, J.; DeClercq, E.; Volckaert, G.; Tavernier,

[176785]J.; Devos, R.; Fiers, W.: Isolation and structure of a human fibroblast

[176786]interferon gene. Nature 285: 542–547, 1980.

[176787]

[176788]

[176789]18205.

[176790]Diaz, M. O.; Le Beau, M. M.; Pitha, P.; Rowley, J. D.: Inter–

feron

[176791]and c-ets-1 genes in the translocation (9;11)(p22;q23) in human acute

[176792]monocytic leukemia. Science 231: 265-267, 1986.

[176793]

[176794]

[176795]18206.

[176796]Diaz, M. O.; Ziemin, S.; Le Beau, M. M.; Pitha, P.; Smith, S. D.;

[176797]Chilcote, R. R.; Rowley, J. D.: Homozygous deletion of the alpha-

[176798]and beta(1)-interferon genes in human leukemia and derived cell lines. Proc.

[176799]Nat. Acad. Sci. 85: 5259-5263, 1988.

[176800]

[176801]

[176802]18207.

[176803]Erickson, B. W.; May, L. T.; Sehgal, P. B.: Internal duplication
tion

[176804]in human alpha-1 and beta-1 interferons. Proc. Nat. Acad.

Sci. 81:

[176805]7171–7175, 1984.

[176806]

[176807]

[176808]18208.

[176809]Henry, L.; Sizun, J.; Turleau, C.; Boue, J.; Azoulay, M.; Ju-
nien,

[176810]C.: The gene for human fibroblast interferon (IFB) maps to
9p21. Hum.

[176811]Genet. 68: 67–69, 1984.

[176812]

[176813]

[176814]18209.

[176815]Houghton, M.; Eaton, M. A. W.; Stewart, A. G.; Smith, J. C.;
Doel,

[176816]S. M.; Catlin, G. H.; Lewis, H. M.; Patel, T. P.; Emtage, J. S.;
Carey,

[176817]N. H.; Porter, A. G.: The complete amino acid sequence of
human fibroblast

[176818]interferon as deduced using synthetic oligodeoxyribonu–

cleotide primers

[176819]of reverse transcriptase. Nucleic Acids Res. 8: 2885–2894,
1980.

[176820]

[176821]

[176822]18210.

[176823]Houghton, M.; Jackson, I. J.; Porter, A. G.; Doel, S. M.;
Catlin,

[176824]G. H.; Barber, C.; Carey, N. H.: The absence of introns
within a

[176825]human fibroblast interferon gene. Nucleic Acids Res. 9:
247–266,

[176826]1981.

[176827]

[176828]

[176829]18211.

[176830]Knight, E., Jr.: Human fibroblast interferon: amino acid
analysis

[176831]and amino terminal amino acid sequence. Science 207:
525–527, 1980.

[176832]

[176833]

[176834]18212.

[176835]May, L. T.; Landsberger, F. R.; Inouye, M.; Sehgal, P. B.:

Significance

[176836]of similarities in patterns: an application to beta interferon-related

[176837]DNA on human chromosome 2. Proc. Nat. Acad. Sci. 82: 4090-4094,

[176838]1985.

[176839]

[176840]

[176841]18213.

[176842]Meager, A.; Graves, H.; Burke, D. C.; Swallow, D. M.: Involvement

[176843]of a gene on chromosome 9 in human fibroblast interferon production. Nature 280:

[176844]493-495, 1979.

[176845]

[176846]

[176847]18214.

[176848]Meager, A.; Graves, H. E.; Walker, J. R.; Burke, D. C.; Swallow,

[176849]D. M.; Westerveld, A.: Tentative assignment of the gene for human

[176850]fibroblast interferon to chromosome 9. (Abstract) Cytogenet. Cell

[176851]Genet. 25: 183–184, 1979.

[176852]

[176853]

[176854]18215.

[176855]Ohlsson, M.; Feder, J.; Cavalli-Sforza, L. L.; von Gabain, A.

[176856]: Close linkage of alpha and beta interferons and infrequent duplication

[176857]of beta interferon in humans. Proc. Nat. Acad. Sci. 82: 4473–4476,

[176858]1985.

[176859]

[176860]

[176861]18216.

[176862]Ohno, S.; Taniguchi, T.: Structure of a chromosomal gene

for

[176863]human interferon beta. Proc. Nat. Acad. Sci. 78:
5305–5309, 1981.

[176864]

[176865]

[176866]18217.

[176867]Owerbach, D.; Rutter, W. J.; Shows, T. B.; Gray, P.; Goed-
del,

[176868]D. V.; Lawn, R. M.: Leukocyte and fibroblast interferon
genes are

[176869]located on human chromosome 9. Proc. Nat. Acad. Sci.
78: 3123–3127,

[176870]1981.

[176871]

[176872]

[176873]18218.

[176874]Pitha, P. M.; Slate, D. L.; Raj, N. B. K.; Ruddle, F. H.: Human

[176875]beta interferon gene localization and expression in so-
matic cell hybrids. Molec.

[176876]Cell. Biol. 2: 564–570, 1982.

[176877]

[176878]

[176879]18219.

[176880]Sagar, A. D.; Sehgal, P. B.; May, L. T.; Inouye, M.; Slate, D.

[176881]L.; Shulman, L.; Ruddle, F. H.: Interferon-beta-related

DNA is dispersed

[176882]in the human genome. Science 223: 1312-1315, 1984.

[176883]

[176884]

[176885]18220.

[176886]Sehgal, P. B.; May, L. T.; Sagar, A. D.; LaForge, K. S.; In-

ouye,

[176887]M.: Isolation of novel human genomic DNA clones related

to human

[176888]interferon-beta(1) cDNA. Proc. Nat. Acad. Sci. 80:

3632-3636, 1983.

[176889]

[176890]

[176891]18221.

[176892]Shepard, H. M.; Leung, D.; Stebbing, N.; Goeddel, D. V.: A
single

[176893]amino acid change in IFN-beta(1) abolishes its antiviral
activity. Nature 294:

[176894]563-567, 1981.

[176895]

[176896]

[176897]18222.

[176898]Siegal, F. P.; Kadowaki, N.; Shodell, M.; Fitzgerald-Bo-
carsly,

[176899]P. A.; Shah, K.; Ho, S.; Antonenko, S.; Liu, Y.-J.: The nature
of

[176900]the principal type 1 interferon-producing cells in human
blood. Science 284:

[176901]1835-1837, 1999.

[176902]

[176903]

[176904]18223.

[176905]Heath, P.; Elvin, P.; Jenner, D.; Gammack, A.; Morten, J.;
Markham,

[176906]A.: Localisation of a cDNA clone for human cytokeratin 18
to chromosome

[176907]17p11–p12 by in situ hybridisation. Hum. Genet. 85:
669–670, 1990.

[176908]

[176909]

[176910]18224.

[176911]Ku, N.–O.; Michie, S.; Oshima, R. G.; Omary, M. B.: Chronic
hepatitis,

[176912]hepatocyte fragility, and increased soluble phosphogly-
cokeratins in

[176913]transgenic mice expressing a keratin 18 conserved argi-
nine mutant. J.

[176914]Cell Biol. 131: 1303–1314, 1995.

[176915]

[176916]

[176917]18225.

[176918]Ku, N.–O.; Michie, S. A.; Soetikno, R. M.; Resurreccion, E.
Z.;

[176919]Broome, R. L.; Oshima, R. G.; Omary, M. B.: Susceptibility
to hepatotoxicity

[176920]in transgenic mice that express a dominant-negative human keratin

[176921]18 mutant. J. Clin. Invest. 98: 1034–1046, 1996.

[176922]

[176923]

[176924]18226.

[176925]Kulesh, D. A.; Oshima, R. G.: Complete structure of the gene for

[176926]human keratin 18. Genomics 4: 339–347, 1989.

[176927]

[176928]

[176929]18227.

[176930]Lendahl, U.; Zimmerman, L. B.; McKay, R. D. G.: CNS stem cells

[176931]express a new class of intermediate filament protein. Cell 60: 585–595,

[176932]1990.

[176933]

[176934]

[176935]18228.

[176936]Steinert, P. M.; Roop, D. R.: Molecular and cellular biology
of
[176937]intermediate filaments. Ann. Rev. Biochem. 57: 593–625,
1988.

[176938]

[176939]

[176940]18229.

[176941]Waseem, A.; Gough, A. C.; Spurr, N. K.; Lane, E. B.: Local-
ization
[176942]of the gene for human simple epithelial keratin 18 to
chromosome 12

[176943]using polymerase chain reaction. Genomics 7: 188–194,
1990.

[176944]

[176945]

[176946]18230.

[176947]Erikson, J.; Griffin, C.; ar-Rushdi, A.; Valtieri, M.; Hoxie,
[176948]J.; Finan, J.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.;
Croce, C.

[176949]M.: Heterogeneity of chromosome 22 breakpoint in
Philadelphia-positive

[176950](Ph+) acute lymphocytic leukemia. Proc. Nat. Acad. Sci.
83: 1807-1811,

[176951]1986.

[176952]

[176953]

[176954]18231.

[176955]Fioretos, T.; Heisterkamp, N.; Groffen, J.: No evidence for
genomic

[176956]imprinting of the human BCR gene. Blood 83: 3441-3444,
1994.

[176957]

[176958]

[176959]18232.

[176960]Fitzgerald, P. H.: Evidence that chromosome band 22q12
is concerned

[176961]with cell proliferation in chronic myeloid leukemia. Hum.
Genet. 33:

[176962]269-274, 1976.

[176963]

[176964]

[176965]18233.

[176966]Ganesan, T. S.; Rassool, F.; Guo, A.-P.; Th'ng, K. H.;
Dowding,

[176967]C.; Hibbin, J. A.; Young, B. D.; White, H.; Kumaran, T. O.;
Galton,

[176968]D. A. G.; Goldman, J. M.: Rearrangement of the bcr gene in
Philadelphia

[176969]chromosome-negative chronic myeloid leukemia. Blood
68: 957-960,

[176970]1986.

[176971]

[176972]

[176973]18234.

[176974]Goldman, J. M.; Melo, J. V.: Targeting the BCR-ABL tyro-
sine kinase

[176975]in chronic myeloid leukemia. (Editorial) New Eng. J. Med.
344: 1084-1086,

[176976]2001.

[176977]

[176978]

[176979]18235.

[176980]Gorre, M. E.; Mohammed, M.; Ellwood, K.; Hsu, N.; Paquette, R.;

[176981]Rao, P. N.; Sawyers, C. L.: Clinical resistance to STI-571 cancer

[176982]therapy caused by BCR-ABL gene mutation or amplification. Science 293:

[176983]876-883, 2001.

[176984]

[176985]

[176986]18236.

[176987]Groffen, J.; Stephenson, J. R.; Heisterkamp, N.; de Klein, A.;

[176988]Bartram, C. R.; Grosveld, G.: Philadelphia chromosomal breakpoints

[176989]are clustered within a limited region, bcr, on chromosome 22. Cell 36:

[176990]93-99, 1984.

[176991]

[176992]

[176993]18237.

[176994]Grossman, A.; Silver, R. T.; Arlin, Z.; Coleman, M.; Cam-
posano,

[176995]E.; Gascon, P.; Benn, P. A.: Fine mapping of chromosome
22 breakpoints

[176996]within the breakpoint cluster region (bcr) implies a role for
bcr

[176997]exon 3 in determining disease duration in chronic myeloid
leukemia. Am.

[176998]J. Hum. Genet. 45: 729–738, 1989.

[176999]

[177000]

[177001]18238.

[177002]Grosveld, G.; Verwoerd, T.; van Agthoven, T.; de Klein, A.;
Ramachandran,

[177003]K. L.; Heisterkamp, N.; Stam, K.; Groffen, J.: The chronic
myelocytic

[177004]cell line K562 contains a breakpoint in bcr and produces a
chimeric

[177005]bcr/c-abl transcript. Molec. Cell. Biol. 6: 607–616, 1986.

[177006]

[177007]

[177008]18239.

[177009]Haas, O. A.: Are ABL and BCR imprinted? No definitive answers,

[177010]but more questions. Leukemia 9: 740–745, 1995.

[177011]

[177012]

[177013]18240.

[177014]Haas, O. A.; Argyriou–Tirita, A.; Lion, T.: Parental origin of

[177015]chromosomes involved in the translocation t(9;22). Nature
359: 414–416,

[177016]1992.

[177017]

[177018]

[177019]18241.

[177020]Hariharan, I. K.; Adams, J. M.: cDNA sequence for human

bcr,

[177021]the gene that translocates to the abl oncogene in chronic myeloid

[177022]leukaemia. EMBO J. 6: 115–119, 1987.

[177023]

[177024]

[177025]18242.

[177026]Heisterkamp, N.; Jenster, G.; ten Hoeve, J.; Zovich, D.; Pat-
tengale,

[177027]P. K.; Groffen, J.: Acute leukaemia in bcr/abl transgenic
mice. Nature 344:

[177028]251–253, 1990.

[177029]

[177030]

[177031]18243.

[177032]Heisterkamp, N.; Stam, K.; Groffen, J.; de Klein, A.;
Grosveld,

[177033]G.: Structural organization of the bcr gene and its role in
the Ph-1

[177034]translocation. Nature 315: 758–761, 1985.

[177035]

[177036]

[177037]18244.

[177038]Hermans, A.; Heisterkamp, N.; von Lindern, M.; van Baal,
S.; Meijer,

[177039]D.; van der Plas, D.; Wiedemann, L. M.; Groffen, J.;
Bootsma, D.;

[177040]Grosveld, G.: Unique fusion of bcr and c-abl genes in
Philadelphia

[177041]chromosome positive acute lymphoblastic leukemia. Cell
51: 33–40,

[177042]1987.

[177043]

[177044]

[177045]18245.

[177046]Huettner, C. S.; Zhang, P.; Van Etten, R. A.; Tenen, D. G.:
Reversibility

[177047]of acute B-cell leukaemia induced by BCR–ABL1. Nature
Genet. 24:

[177048]57–60, 2000.

[177049]

[177050]

[177051]18246.

[177052]Jacobs, A.: Benzene and leukemia. Brit. J. Haemat. 72:
119-121,

[177053]1989.

[177054]

[177055]

[177056]18247.

[177057]Klein, G.: Specific chromosomal translocations and the
genesis

[177058]of the B-cell-derived tumors in mice and men. Cell 32:
311-315,

[177059]1983.

[177060]

[177061]

[177062]18248.

[177063]Koeffler, H. P.; Golde, D. W.: Chronic myelogenous
leukemia--new

[177064]concepts. New Eng. J. Med. 304: 1201–1209 and
1269–1274, 1981.

[177065]

[177066]

[177067]18249.

[177068]Kohno, S.-I.; Sandberg, A. A.: Chromosomes and causa-
tion of human

[177069]cancer and leukemia. XXXIX. Usual and unusual findings in
Ph(1)–positive

[177070]CML. Cancer 46: 2227–2237, 1980.

[177071]

[177072]

[177073]18250.

[177074]Laurent, E.; Talpaz, M.; Kantarjian, H.; Kurzrock, R.: The
BCR

[177075]gene and Philadelphia chromosome–positive leukemogen-
esis. Cancer

[177076]Res. 61: 2343–2355, 2001.

[177077]

[177078]

[177079]18251.

[177080]Lillicrap, D. A.; Sterndale, H.: Familial chronic myeloid
leukaemia.

[177081](Letter) Lancet II: 699, 1984.

[177082]

[177083]

[177084]18252.

[177085]Lim, Y.-M.; Wong, S.; Lau, G.; Witte, O. N.; Colicelli, J.:
BCR/ABL

[177086]inhibition by an escort/phosphatase fusion protein. Proc.
Nat. Acad.

[177087]Sci. 97: 12233–12238, 2000.

[177088]

[177089]

[177090]18253.

[177091]Litz, C. E.; Copenhaver, C. M.: Paternal origin of the rear-
ranged

[177092]major breakpoint cluster region in chronic myeloid
leukemia. Blood 83:

[177093]3445–3448, 1994.

[177094]

[177095]

[177096]18254.

[177097]Maru, Y.; Witte, O. N.: The BCR gene encodes a novel ser-
ine/threonine

[177098]kinase activity within a single exon. Cell 67: 459–468,
1991.

[177099]

[177100]

[177101]18255.

[177102]Melo, J. V.; Yan, X.–H.; Diamond, J.; Goldman, J. M.: Lack
of

[177103]imprinting of the ABL gene. (Letter) Nature Genet. 8:
318–319, 1994.

[177104]

[177105]

[177106]18256.

[177107]Melo, J. V.; Yan, X.–H.; Diamond, J.; Goldman, J. M.: Bal-
anced

[177108]parental contribution to the ABL component of the BCR–

ABL gene in

[177109]chronic myeloid leukemia. Leukemia 9: 734–745, 1995.

[177110]

[177111]

[177112]18257.

[177113]Mes–Masson, A.–M.; McLaughlin, J.; Daley, G. Q.; Paskind, M.;

[177114]Witte, O. N.: Overlapping cDNA clones define the complete coding

[177115]region for the P210(c-abl) gene product associated with chronic myelogeneous

[177116]leukemia cells containing the Philadelphia chromosome. Proc. Nat.

[177117]Acad. Sci. 83: 9768–9772, 1986.

[177118]

[177119]

[177120]18258.

[177121]Mills, K. I.; MacKenzie, E. D.; Birnie, G. D.: The site of the

[177122]breakpoint within the bcr is a prognostic factor in Philadelphia–positive

[177123]CML patients. Blood 72: 1237–1241, 1988.

[177124]

[177125]

[177126]18259.

[177127]Mittelman, F.; Levan, G.: Clustering of aberrations to specific
cific

[177128]chromosomes in human neoplasms. III. Incidence and geographic distribution

[177129]of chromosome aberrations in 856 cases. Hereditas 89:
207–232, 1978.

[177130]

[177131]

[177132]18260.

[177133]Nowell, P. C.; Hungerford, D. A.: A minute chromosome in
human

[177134]chronic granulocytic leukemia. Science 132: 1497, 1960.

[177135]

[177136]

[177137]18261.

[177138]Kusari, J.; Takata, Y.; Hatada, E.; Freidenberg, G.; Kolter-
man,

[177139]O.; Olefsky, J. M.: Insulin resistance and diabetes due to
different

[177140]mutations in the tyrosine kinase domain of both insulin
receptor gene

[177141]alleles. J. Biol. Chem. 266: 5260–5267, 1991.

[177142]

[177143]

[177144]18262.

[177145]Lander, E. S.; Botstein, D.: Homozygosity mapping: a way
to map

[177146]human recessive traits with the DNA of inbred children.
Science 236:

[177147]1567–1570, 1987.

[177148]

[177149]

[177150]18263.

[177151]Kramer, F.; White, K.; Pauleikhoff, D.; Gehrig, A.; Pass-
more,

[177152]L.; Rivera, A.; Rudolph, G.; Kellner, U.; Andrassi, M.;
Lorenz, B.;

[177153]Rohrschneider, K.; Blankenagel, A.; Jurklies, B.; Schilling,
H.; Schutt,

[177154]F.; Holz, F. G.; Weber, B. H. F.: Mutations in the VMD2
gene are

[177155]associated with juvenile-onset vitelliform macular dystro-
phy (Best

[177156]disease) and adult vitelliform macular dystrophy but not
age-related

[177157]macular degeneration. Europ. J. Hum. Genet. 8: 286–292,
2000.

[177158]

[177159]

[177160]18264.

[177161]Sagane, K.; Ohya, Y.; Hasegawa, Y.; Tanaka, I.: Metallo-
proteinase-like,

[177162]disintegrin-like, cysteine-rich proteins MDC2 and MDC3:
novel human

[177163]cellular disintegrins highly expressed in the brain.
Biochem. J. 334:

[177164]93–98, 1998.

[177165]

[177166]

[177167]18265.

[177168]Alsobrook, J. P., II; Zohar, A. H.; Leboyer, M.; Chabane, N.;
Ebstein,

[177169]R. P.; Pauls, D. L.: Association between the COMT locus
and obsessive-compulsive

[177170]disorder in females but not males. Am. J. Med. Genet.
(Neuropsychiat.

[177171]Genet.) 114: 116–120, 2002.

[177172]

[177173]

[177174]18266.

[177175]Karayiorgou, M.; Sobin, C.; Blundell, M. L.; Galke, B. L.;
Malinova,

[177176]L.; Goldberg, P.; Ott, J.; Gogos, J. A.: Family-based associ-
ation

[177177]studies support a sexually dimorphic effect of COMT and
MAOA on genetic

[177178]susceptibility to obsessive-compulsive disorder. Biol. Psy-
chiat. 45:

[177179]1178–1189, 1999.

[177180]

[177181]

[177182]18267.

[177183]Taira, M.; Yoshida, T.; Miyagawa, K.; Sakamoto, H.; Terada, M.;

[177184]Sugimura, T.: cDNA sequence of human transforming gene hst and identification

[177185]of the coding sequence required for transforming activity. Proc.

[177186]Nat. Acad. Sci. 84: 2980–2984, 1987.

[177187]

[177188]

[177189]18268.

[177190]Krill, A. E.; Morse, P. A.; Potts, A. M.; Klien, B. A.: Hereditary

[177191]vitelliruptive macular degeneration. Am. J. Ophthal. 61: 1405–1415,

[177192]1966.

[177193]

[177194]

[177195]18269.

[177196]Maloney, W. F.; Robertson, D. M.; Miller, S. A.: Hereditary
vitelliform

[177197]macular degeneration--variable fundus findings within a
single pedigree. Arch.

[177198]Ophthal. 95: 979-983, 1977.

[177199]

[177200]

[177201]18270.

[177202]Mansergh, F. C.; Kenna, P. F.; Rudolph, G.; Meitinger, T.;
Farrar,

[177203]G. J.; Kumar-Singh, R.; Scorer, J.; Hally, A. M.; Mynett-
Johnson,

[177204]L.; Humphries, M. M.; Kiang, S.; Humphries, P.: Evidence
for genetic

[177205]heterogeneity in Best's vitelliform macular dystrophy. J.
Med. Genet. 32:

[177206]855-858, 1995.

[177207]

[177208]

[177209]18271.

[177210]Marmorstein, A. D.; Marmorstein, L. Y.; Rayborn, M.;
Wang, X.;

[177211]Hollyfield, J. G.; Petrukhin, K.: Bestrophin, the product of
the

[177212]Best vitelliform macular dystrophy gene (VMD2), localizes
to the basolateral

[177213]plasma membrane of the retinal pigment epithelium. Proc.
Nat. Acad.

[177214]Sci. 97: 12758–12763, 2000.

[177215]

[177216]

[177217]18272.

[177218]Marquardt, A.; Stohr, H.; Passmore, L. A.; Kramer, F.;
Rivera,

[177219]A.; Weber, B. H. F.: Mutations in a novel gene, VMD2, en-
coding a

[177220]protein of unknown properties cause juvenile-onset vitel-
liform macular

[177221]dystrophy (Best's disease). Hum. Molec. Genet. 7:

1517–1525, 1998.

[177222]

[177223]

[177224]18273.

[177225]Nichols, B. E.; Bascom, R.; Litt, M.; McInnes, R.; Sheffield,

[177226]V. C.; Stone, E. M.: Refining the locus for Best vitelliform
macular

[177227]dystrophy and mutation analysis of the candidate gene
ROM1. Am. J.

[177228]Hum. Genet. 54: 95–103, 1994.

[177229]

[177230]

[177231]18274.

[177232]Nordstrom, S.: Epidemiological studies of hereditary mac-
ular

[177233]degeneration (Best's disease) in Swedish and Swedish-
American populations.In:

[177234]Eriksson, A. W.; Forsius, H. R.; Nevanlinna, H. R.; Work-
man, P. L.;

[177235]Norio, R. K.: Population Structure and Genetic Disorders.

New York:

[177236]Academic Press (pub.) 1980. Pp. 431–443.

[177237]

[177238]

[177239]18275.

[177240]Nordstrom, S.: Personal Communication. Umea, Sweden
1978.

[177241]

[177242]

[177243]18276.

[177244]Nordstrom, S.; Thorburn, W.: Dominantly inherited macu-
lar degeneration

[177245](Best's disease) in a homozygous father with 11 children.

Clin. Genet. 18:

[177246]211–216, 1980.

[177247]

[177248]

[177249]18277.

[177250]O'Gorman, S.; Flaherty, W. A.; Fishman, G. A.; Berson, E.

L.:

[177251]Histopathologic findings in Best's vitelliform macular dystrophy. Arch.

[177252]Ophthalmol. 106: 1261–1268, 1988.

[177253]

[177254]

[177255]18278.

[177256]Petrukhin, K.; Koisti, M. J.; Bakall, B.; Li, W.; Xie, G.; Marknell,

[177257]T.; Sandgren, O.; Forsman, K.; Holmgren, G.; Andreasson, S.; Vujic,

[177258]M.; Bergen, A. A. B.; McGarty–Dugan, V.; Figueroa, D.; Austin, C.

[177259]P.; Metzker, M. L.; Caskey, C. T.; Wadelius, C.: Identification of

[177260]the gene responsible for Best macular dystrophy. Nature Genet. 19:

[177261]241–247, 1998.

[177262]

[177263]

[177264]18279.

[177265]Rivas, F.; Ruiz, C.; Rivera, H.; Moller, M.; Serrano-Lucas, J.

[177266]I.; Cantu, J. M.: De novo del(6)(q25) associated with macular degeneration. Ann.

[177267]Genet. 29: 42-44, 1986.

[177268]

[177269]

[177270]18280.

[177271]Rosas, F. E.: Maculopatía hereditaria viteliforme de Best.
Ann.

[177272]Soc. Mex. Oft. 50: 157-171, 1976.

[177273]

[177274]

[177275]18281.

[177276]Sorsby, A.; Savory, M.; Davey, J. B.; Fraser, R. J. L.: Macular

[177277]cysts: a dominantly inherited affection with a progressive
course. Brit.

[177278]J. Ophthalm. 40: 144-158, 1956.

[177279]

[177280]

[177281]18282.

[177282]Stohr, H.; Marquardt, A.; Rivera, A.; Cooper, P. R.; Nowak, N.

[177283]J.; Shows, T. B.; Gerhard, D. S.; Weber, B. H. F.: A gene map of

[177284]the Best's vitelliform macular dystrophy region in chromosome 11q12-q13.1. Genome

[177285]Res. 8: 48-56, 1998.

[177286]

[177287]

[177288]18283.

[177289]Stone, E. M.; Nichols, B. E.; Streb, L. M.; Kimura, A. E.; Sheffield,

[177290]V. C.: Genetic linkage of vitelliform macular degeneration (Best's

[177291]disease) to chromosome 11q13. Nature Genet. 1: 246-250, 1992.

[177292]

[177293]

[177294]18284.

[177295]Sun, H.; Tsunenari, T.; Yau, K.-W.; Nathans, J.: The vitelli-
form

[177296]macular dystrophy protein defines a new family of chlo-
ride channels. Proc.

[177297]Nat. Acad. Sci. 99: 4008–4013, 2002.

[177298]

[177299]

[177300]18285.

[177301]Vail, D.; Shoch, D.: Hereditary degeneration of the macula.
II.

[177302]Follow-up report and histopathologic study. Trans. Am.
Ophthal. Soc. 63:

[177303]51–63, 1965.

[177304]

[177305]

[177306]18286.

[177307]Vossius, A.: Ueber die Bestsche familiaere Maculadegener-
ation. Arch.

[177308]Ophthal. 105: 1050–1059, 1921.

[177309]

[177310]

[177311]18287.

[177312]Weber, B. H. F.; Walker, D.; Muller, B.: Molecular evidence
for

[177313]non-penetrance in Best's disease. J. Med. Genet. 31:
388-392, 1994.

[177314]

[177315]

[177316]18288.

[177317]Weber, B. H. F.; Walker, D.; Muller, B.; Mar, L.: Best's vitel-
liform

[177318]dystrophy (VMD2) maps between D11S903 and PYGM: no
evidence for locus

[177319]heterogeneity. Genomics 20: 267-274, 1994.

[177320]

[177321]

[177322]18289.

[177323]White, K.; Marquardt, A.; Weber, B. H. F.: VMD2 mutations
in

[177324]vitelliform macular dystrophy (Best disease) and other

maculopathies. Hum.

[177325]Mutat. 15: 301–308, 2000.

[177326]

[177327]

[177328]18290.

[177329]Yoder, F. E.; Cross, H. E.; Chase, G. A.; Fine, S. L.; Freid-
hoff,

[177330]L.; Machan, C. H.; Bias, W. B.: Linkage studies of Best's
macular

[177331]dystrophy. Clin. Genet. 34: 26–30, 1988.

[177332]

[177333]

[177334]18291.

[177335]Farndon, J. R.; Leight, G. S.; Dilley, W. G.; Baylin, S. B.;
Smallridge,

[177336]R. C.; Harrison, T. S.; Wells, S. A., Jr.: Familial medullary
thyroid

[177337]carcinoma without associated endocrinopathies: a distinct
clinical

[177338]entity. Brit. J. Surg. 73: 278–281, 1986.

[177339]

[177340]

[177341]18292.

[177342]Wallis, D. E.; Roessler, E.; Hehr, U.; Nanni, L.; Wiltshire, T.;

[177343]Richieri-Costa, A.; Gillessen-Kaesbach, G.; Zackai, E. H.;
Rommens,

[177344]J.; Muenke, M.: Mutations in the homeodomain of the hu-
man SIX3 gene

[177345]cause holoprosencephaly. Nature Genet. 22: 196-198,
1999.

[177346]

[177347]

[177348]18293.

[177349]McMichael, A.; Makgoba, W.: Complexity in human histo-
compatibility

[177350]loci. Nature 293: 701-702, 1981.

[177351]

[177352]

[177353]18294.

[177354]Moen, T.; Albrechtsen, D.; Flatmark, A.; Jakobsen, A.;

Jervell,

[177355]J.; Halvorsen, S.; Solheim, B. G.; Thorsby, E.: Importance of
HLA-DR

[177356]matching in cadaveric renal transplantation: a prospective
one-center

[177357]study of 170 transplants. New Eng. J. Med. 303: 850-854,
1980.

[177358]

[177359]

[177360]18295.

[177361]Olerup, O.; Troye-Blomberg, M.; Schreuder, G. M. T.; Ri-
ley, E.

[177362]M.: HLA-DR and -DQ gene polymorphism in West Africans
is twice as

[177363]extensive as in North European Caucasians: evolutionary
implications. Proc.

[177364]Nat. Acad. Sci. 88: 8480-8484, 1991.

[177365]

[177366]

[177367]18296.

[177368]Park, M. S.; Terasaki, P. I.; Bernoco, D.: Relationship be-

tween

[177369]MT and DR antigens.In: Terasaki, P. I.: Histocompatibility
Testing

[177370]1980. Los Angeles: UCLA Press (pub.) 1980.

[177371]

[177372]

[177373]18297.

[177374]Park, M. S.; Terasaki, P. I.; Bernoco, D.; Iwaki, Y.: Evidence
[177375]for a second B-cell locus separate from the DR locus.

Transplant.

[177376]Proc. 10: 823–828, 1978.

[177377]

[177378]

[177379]18298.

[177380]Pisella, P.–J.; Brignole, F.; Debbasch, C.; Lozato, P.–A.;
Creuzot–Garcher,

[177381]C.; Bara, J.; Saiag, P.; Warnet, J.–M.; Baudouin, C.: Flow cy-
tometric

[177382]analysis of conjunctival epithelium in ocular rosacea and
keratoconjunctivitis

[177383]sicca. Ophthalmology 107: 1841–1849, 2000.

[177384]

[177385]

[177386]18299.

[177387]Rollini, P.; Mach, B.; Gorski, J.: Linkage map of three HLA-
DR

[177388]beta-chain genes: evidence for a recent duplication event.
Proc.

[177389]Nat. Acad. Sci. 82: 7197–7201, 1985.

[177390]

[177391]

[177392]18300.

[177393]Sachs, J. A.; Jaraquemada, D.; Festenstein, H.: Intra HLA-D
region

[177394]recombinant maps HLA-DR between HLA-B and HLA-D.
Tissue Antigens 17:

[177395]43–56, 1981.

[177396]

[177397]

[177398]18301.

[177399]Schwartz, B. D.: Personal Communication. St. Louis, Mo.
2/22/1983.

[177400]

[177401]

[177402]18302.

[177403]Shackelford, D. A.; Mann, D. L.; van Rood, J. J.; Ferrara, G.

[177404]B.; Strominger, J. L.: Human B-cell alloantigens DC1, MT1,
and LB12

[177405]are identical to each other but distinct from the HLA-DR
antigen. Proc.

[177406]Nat. Acad. Sci. 78: 4566-4570, 1981.

[177407]

[177408]

[177409]18303.

[177410]Strominger, J. L.: Biology of the human histocompatibility
leukocyte

[177411]antigen (HLA) system and a hypothesis regarding the gen-
eration of

[177412]autoimmune diseases. J. Clin. Invest. 77: 1411-1415,

1986.

[177413]

[177414]

[177415]18304.

[177416]Suciu-Foca, N.; Weiner, J.; Rohowsky, C.; McKiernan, P.;

Susinno,

[177417]E.; Rubinstein, P.: Indications that Dw determinants are
controlled

[177418]by distinct (but closely linked) genes. Transplant. Proc. 10:
799-804,

[177419]1978.

[177420]

[177421]

[177422]18305.

[177423]Thursz, M. R.; Thomas, H. C.; Greenwood, B. M.; Hill, A. V.

S.

[177424]: Heterozygote advantage for HLA class-II type in hepatitis
B virus

[177425]infection. (Letter) Nature Genet. 17: 11-12, 1997.

[177426]

[177427]

[177428]18306.

[177429]Tiercy, J.-M.; Gorski, J.; Jeannet, M.; Mach, B.: Identifica-
tion

[177430]and distribution of three serologically undetected alleles
of HLA-DR

[177431]by oligonucleotide-DNA typing analysis. Proc. Nat. Acad.
Sci. 85:

[177432]198-202, 1988.

[177433]

[177434]

[177435]18307.

[177436]Tiwari, J. L.; Terasaki, P. I.: HLA and Disease Associations.

[177437]New York: Springer-Verlag (pub.) 1985.

[177438]

[177439]

[177440]18308.

[177441]Trowsdale, J.: Genetics and polymorphism: class II anti-
gens. Brit.

[177442]Med. Bull. 43: 15–36, 1987.

[177443]

[177444]

[177445]18309.

[177446]Tsubota, K.; Fukagawa, K.; Fujihara, T.; Shimmura, S.;
Saito,

[177447]I.; Saito, K.; Takeuchi, T.: Regulation of human leukocyte
antigen

[177448]expression in human conjunctival epithelium. Invest.
Ophthalm. Vis.

[177449]Sci. 40: 28–34, 1999.

[177450]

[177451]

[177452]18310.

[177453]Walker, L. E.; Hewick, R.; Hunkapiller, M. W.; Hood, L. E.;
Dreyer,

[177454]W. J.; Reisfeld, R. A.: N-terminal amino acid sequences of
the alpha

[177455]and beta chains of HLA-DR1 and HLA-DR2 antigens. Bio-
chemistry 23:

[177456]185–188, 1983.

[177457]

[177458]

[177459]18311.

[177460]Ellis, S. A.; Palmer, M. S.; McMichael, A. J.: Human trophoblast

[177461]and the choriocarcinoma cell line BeWo express a truncated HLA class

[177462]I molecule. J. Immun. 144: 731–735, 1990.

[177463]

[177464]

[177465]18312.

[177466]Geraghty, D. E.; Koller, B. H.; Orr, H. T.: A human major histocompatibility

[177467]complex class I gene that encodes a protein with a shortened cytoplasmic

[177468]segment. Proc. Nat. Acad. Sci. 84: 9145–9149, 1987.

[177469]

[177470]

[177471]18313.

[177472]Geraghty, D. E.; Pei, J.; Lipsky, B.; Hansen, J. A.; Taillon-Miller,

[177473]P.; Bronson, S. K.; Chaplin, D. D.: Cloning and physical mapping

[177474]of the HLA class I region spanning the HLA-E-to-HLA-F interval by

[177475]using yeast artificial chromosomes. Proc. Nat. Acad. Sci. 89: 2669-2673,

[177476]1992.

[177477]

[177478]

[177479]18314.

[177480]Hurks, H. M. H.; Valter, M. M.; Wilson, L.; Hilgert, I.; van den

[177481]Elsen, P. J.; Jager, M. J.: Uveal melanoma: no expression of HLA-G. Invest.

[177482]Ophthal. Vis. Sci. 42: 3081-3084, 2001.

[177483]

[177484]

[177485]18315.

[177486]Kirszenbaum, M.; Moreau, P.; Gluckman, E.; Dausset, J.;

Carosella,

[177487]E.: An alternatively spliced form of HLA-G mRNA in human trophoblasts

[177488]and evidence for the presence of HLA-G transcript in adult lymphocytes. Proc.

[177489]Nat. Acad. Sci. 91: 4209–4213, 1994.

[177490]

[177491]

[177492]18316.

[177493]Lila, N.; Carpentier, A.; Amrein, C.; Khalil-Daher, I.; Daus-

set,

[177494]J.; Carosella, E. D.: Implication of HLA-G molecule in heart-graft

[177495]acceptance. Lancet 355: 2138 only, 2000.

[177496]

[177497]

[177498]18317.

[177499]McAlpine, P. J.: Personal Communication. Winnipeg, Mani-

toba,

[177500]Canada 6/22/1988.

[177501]

[177502]

[177503]18318.

[177504]Morales, P.; Corell, A.; Martinez-Laso, J.; Martin-Villa, J.
M.;

[177505]Varela, P.; Paz-Artal, E.; Allende, L.-M.; Arnaiz-Villena, A.:
Three

[177506]new HLA-G alleles and their linkage disequilibria with
HLA-A. Immunogenetics 38:

[177507]323-331, 1993.

[177508]

[177509]

[177510]18319.

[177511]Onno, M.; Guillaudeux, T.; Amiot, L.; Renard, I.; Drenou,
B.;

[177512]Hirel, B.; Girr, M.; Semana, G.; Le Bouteiller, P.; Fauchet,
R.:

[177513]The HLA-G gene is expressed at a low mRNA level in dif-

ferent human

[177514]cells and tissues. Human Immun. 41: 79–86, 1994.

[177515]

[177516]

[177517]18320.

[177518]Paul, P.; Rouas–Freiss, N.; Khalil–Daher, I.; Moreau, P.;
Riteau,

[177519]B.; Le Gal, F. A.; Avril, M. F.; Dausset, J.; Guillet, J. G.;
Carosella,

[177520]E. D.: HLA–G expression in melanoma: a way for tumor
cells to escape

[177521]from immunosurveillance. Proc. Nat. Acad. Sci. 95:
4510–4515, 1998.

[177522]

[177523]

[177524]18321.

[177525]Rouas–Freiss, N.; Marchal, R. E.; Kirszenbaum, M.; Daus–
set, J.;

[177526]Carosella, E. D.: The alpha1 domain of HLA–G1 and HLA–
G2 inhibits

[177527]cytotoxicity induced by natural killer cells: is HLA–G the

public

[177528]ligand for natural killer cell inhibitory receptors? Proc. Nat. Acad.
Acad.

[177529]Sci. 94: 5249–5254, 1997.

[177530]

[177531]

[177532]18322.

[177533]Schmidt, C. M.; Orr, H. T.: A physical linkage map of HLA–
A,

[177534]–G, –7.5p, and –F. Hum. Immun. 31: 180–185, 1991.

[177535]

[177536]

[177537]18323.

[177538]Stewart, W. E., II: The Interferon System. Berlin: Springer

[177539](pub.) 1979.

[177540]

[177541]

[177542]18324.

[177543]Farfel, Z.; Iiri, T.; Shapira, H.; Roitman, A.; Mouallem, M.;

[177544]Bourne, H. R.: Pseudohypoparathyroidism, a novel mutation in the

[177545]beta/gamma- contact region of Gs-alpha impairs receptor stimulation. J.

[177546]Biol. Chem. 271: 19653-19655, 1996.

[177547]

[177548]

[177549]18325.

[177550]Heisler, L. K.; Cowley, M. A.; Tecott, L. H.; Fan, W.; Low, M.

[177551]J.; Smart, J. L.; Rubinstein, M.; Tatro, J. B.; Marcus, J. N.; Holstege,

[177552]H.; Lee, C. E.; Cone, R. D.; Elmquist, J. K.: Activation of central

[177553]melanocortin pathways by fenfluramine. Science 297: 609-611, 2002.

[177554]

[177555]

[177556]18326.

[177557]Folster-Holst, R.; Moises, H. W.; Yang, L.; Fritsch, W.;

Weissenbach,

[177558]J.; Christophers, E.: Linkage between atopy and the IgE
high-affinity

[177559]receptor gene at 11q13 in atopic dermatitis families.

Hum. Genet. 102:

[177560]236–239, 1998.

[177561]

[177562]

[177563]18327.

[177564]Hill, M. R.; Cookson, W. O. C. M.: A new variant of the beta
subunit

[177565]of the high-affinity receptor for immunoglobulin E
(Fc-epsilon-RI-beta

[177566]E237G): associations with measures of atopy and
bronchial hyper-responsiveness. Hum.

[177567]Molec. Genet. 5: 959–962, 1996.

[177568]

[177569]

[177570]18328.

[177571]Hupp, K.; Siwarski, D.; Mock, B. A.; Kinet, J.-P.: Gene map–

ping

[177572]of the three subunits of the high affinity FcR for IgE to
mouse chromosomes

[177573]1 and 19. J. Immun. 143: 3787–3791, 1989.

[177574]

[177575]

[177576]18329.

[177577]Kuster, H.; Zhang, L.; Brini, A. T.; MacGlashan, D. W. J.;
Kinet,

[177578]J.–P.: The gene and cDNA for the human high affinity im-
munoglobulin

[177579]E receptor beta chain and expression of the complete hu-
man receptor. J.

[177580]Biol. Chem. 267: 12782–12787, 1992.

[177581]

[177582]

[177583]18330.

[177584]Nagata, H.; Mutoh, H.; Kumahara, K.; Arimoto, Y.;
Tomemori, T.;

[177585]Sakurai, D.; Arase, K.; Ohno, K.; Yamakoshi, T.; Nakano,

K.; Okawa,

[177586]T.; Numata, T.; Konno, A.: Association between nasal allergy and

[177587]a coding variant of the Fc-epsilon-RI-beta gene Glu237Gly in a Japanese

[177588]population. Hum. Genet. 109: 262–266, 2001.

[177589]

[177590]

[177591]18331.

[177592]Shirakawa, T.; Li, A.; Dubowitz, M.; Dekker, J. W.; Shaw, A. E.;

[177593]Faux, J. A.; Ra, C.; Cookson, W. O. C. M.; Hopkin, J. M.: Association
sociation

[177594]between atopy and variants of the beta subunit of the
high-affinity

[177595]immunoglobulin E receptor. Nature Genet. 7: 125–130,
1994.

[177596]

[177597]

[177598]18332.

[177599]Shirakawa, T.; Mao, X.-Q.; Sasaki, S.; Enomoto, T.; Kawai,

M.;

[177600]Morimoto, K.; Hopkin, J.: Association between atopic asthma and a

[177601]coding variant of Fc-epsilon-RI-beta in a Japanese population. Hum.

[177602]Molec. Genet. 5: 1129-1130, 1996.

[177603]

[177604]

[177605]18333.

[177606]Szepietowski, P.; Gaudray, P.: FCER1B, a candidate gene for atopy,

[177607]is located in 11q13 between CD20 and TCN1. Genomics 19: 399-400,

[177608]1994.

[177609]

[177610]

[177611]18334.

[177612]Kuster, H.; Thompson, H.; Kinet, J.-P.: Characterization and expression

[177613]of the gene for the human Fc receptor gamma subunit:

definition of

[177614]a new gene family. J. Biol. Chem. 265: 6448–6452, 1990.

[177615]

[177616]

[177617]18335.

[177618]Takayanagi, H.; Kim, S.; Matsuo, K.; Suzuki, H.; Suzuki, T.;
Sato,

[177619]K.; Yokochi, T.; Oda, H.; Nakamura, K.; Ida, N.; Wagner, E.
F.; Taniguchi,

[177620]T.: RANKL maintains bone homeostasis through c-
Fos-dependent induction

[177621]of interferon- β . Nature 416: 744–749, 2002.

[177622]

[177623]

[177624]18336.

[177625]Taniguchi, T.; Fujii-Kuriyama, Y.; Muramatsu, M.: Molecu-
lar cloning

[177626]of human interferon cDNA. Proc. Nat. Acad. Sci. 77:
4003–4006, 1980.

[177627]

[177628]

[177629]18337.

[177630]Taniguchi, T.; Mantei, N.; Schwarzstein, M.; Nagata, S.;

Muramatsu,

[177631]M.; Weissmann, C.: Human leukocyte and fibroblast inter-

ferons are

[177632]structurally related. Nature 285: 547-549, 1980.

[177633]

[177634]

[177635]18338.

[177636]Taniguchi, T.; Ohno, S.; Fujii-Kuriyama, Y.; Muramatsu,

M.: The

[177637]nucleotide sequence of human fibroblast interferon cDNA.

Gene 10:

[177638]11-15, 1980.

[177639]

[177640]

[177641]18339.

[177642]Tavernier, J.; Derynck, R.; Fiers, W.: Evidence for a unique

[177643]human fibroblast interferon (IFN-beta1) chromosomal
gene devoid of

[177644]intervening sequences. Nucleic Acids Res. 9: 461-471,
1981.

[177645]

[177646]

[177647]18340.

[177648]Tavernier, J.; Gheysen, D.; Duerinck, F.; Van der Heyden,
J.;

[177649]Fiers, W.: Deletion mapping of the inducible promoter of
human IFN-beta

[177650]gene. Nature 301: 634-636, 1983.

[177651]

[177652]

[177653]18341.

[177654]Trent, J. M.; Olson, S.; Lawn, R.: Chromosomal localization
of

[177655]human leukocyte, fibroblast, and immune interferon
genes by means

[177656]of in situ hybridization. Proc. Nat. Acad. Sci. 79:
7809-7813, 1982.

[177657]

[177658]

[177659]18342.

[177660]Weissenbach, J.; Chernajovsky, Y.; Zeevi, M.; Shulman, L.;
Soreq,

[177661]H.; Nir, U.; Wallach, D.; Perricaudet, M.; Tiollais, P.; Revel,
M.

[177662]: Two interferon mRNAs in human fibroblasts: in vitro
translation

[177663]and Escherichia coli cloning studies. Proc. Nat. Acad. Sci.
77:

[177664]7152–7156, 1980.

[177665]

[177666]

[177667]18343.

[177668]Zinn, K.; DiMaio, D.; Maniatis, T.: Identification of two dis-
tinct

[177669]regulatory regions adjacent to the human beta-interferon
gene. Cell 34:

[177670]865–879, 1983.

[177671]

[177672]

[177673]18344.

[177674]Bruns, G. A. P.; Eisenman, R. E.; Gerald, P. S.: Human mitochondrial

[177675]NADP-dependent isocitrate dehydrogenase in man-mouse somatic cell

[177676]hybrids. Cytogenet. Cell Genet. 17: 200-211, 1976.

[177677]

[177678]

[177679]18345.

[177680]Champion, M. J.; Brown, J. A.; Shows, T. B.: Assignment of cytoplasmic

[177681]alpha-mannosidase (MAN-A) and confirmation of the mitochondrial isocitrate

[177682]dehydrogenase (IDH-M) genes to the q11-qter region of chromosome 15

[177683]in man. Cytogenet. Cell Genet. 22: 498-502, 1978.

[177684]

[177685]

[177686]18346.

[177687]Grzeschik, K.-H.: Assignment of a gene for human mitochondrial

[177688]isocitrate dehydrogenase (ICD-M, EC 1.1.1.41.) to chromosome 15. Hum.

[177689]Genet. 34: 23-28, 1976.

[177690]

[177691]

[177692]18347.

[177693]Huh, T.-L.; Kim, Y.-O.; Oh, I.-U.; Song, B. J.; Inazawa, J.: Assignment

[177694]of the human mitochondrial NAD(+)-specific isocitrate dehydrogenase

[177695]alpha subunit (IDH3A) gene to 15q25.1-q25.2 by in situ hybridization. Genomics 32:

[177696]295-296, 1996.

[177697]

[177698]

[177699]18348.

[177700]Oh, I.-U.; Inazawa, J.; Kim, Y.-O.; Song, B. J.; Huh, T.-L.:

Assignment

[177701]of the human mitochondrial NADP(+)-specific isocitrate dehydrogenase

[177702](IDH2) gene to 15q26.1 by in situ hybridization. Genomics 38: 104-106,

[177703]1996.

[177704]

[177705]

[177706]18349.

[177707]Shimizu, N.; Giles, R. E.; Kucherlapati, R. S.; Shimizu, Y.; Ruddle,

[177708]F. H.: Somatic cell genetic assignment of the human gene for mitochondrial

[177709]NADP-linked isocitrate dehydrogenase to the long arm of chromosome

[177710]15. Somat. Cell Genet. 3: 47-60, 1977.

[177711]

[177712]

[177713]18350.

[177714]Appel, S.; Filter, M.; Reis, A.; Hennies, H. C.; Bergheim, A.;

[177715]Ogilvie, E.; Arndt, S.; Simmons, A.; Lovett, M.; Hide, W.;
Ramsay,

[177716]M.; Reichwald, K.; Zimmermann, W.; Rosenthal, A.: Physi-
cal and transcriptional

[177717]map of the critical region for keratolytic winter erythema
(KWE) on

[177718]chromosome 8p22–p23 between D8S550 and D8S1759.
Europ. J. Hum. Genet. 10:

[177719]17–25, 2002.

[177720]

[177721]

[177722]18351.

[177723]Olavarria, E.; Craddock, C.; Dazzi, F.; Marin, D.; Markt, S.;

[177724]Apperley, J. F.; Goldman, J. M.: Imatinib mesylate (STI571)
in the

[177725]treatment of relapse of chronic myeloid leukemia after al-
logeneic

[177726]stem cell transplantation. Blood 99: 3861–3862, 2002.

[177727]

[177728]

[177729]18352.

[177730]Pegoraro, L.; Matera, L.; Ritz, J.; Levis, A.; Palumbo, A.; Biagini,

[177731]G.: Establishment of a Ph(1)-positive human cell line (BV173). J.

[177732]Nat. Cancer Inst. 70: 447-451, 1983.

[177733]

[177734]

[177735]18353.

[177736]Perrotti, D.; Cesi, V.; Trotta, R.; Guerzoni, C.; Santilli, G.;

[177737]Campbell, K.; Iervolino, A.; Condorelli, F.; Gambacorti-Passerini,

[177738]C.; Caligiuri, M. A.; Calabretta, B.: BCR-ABL suppresses C/EBP-alpha

[177739]expression through inhibitory action of hnRNP E2. Nature Genet. 30:

[177740]48-58, 2002.

[177741]

[177742]

[177743]18354.

[177744]Prakash, O.; Yunis, J. J.: High resolution chromosomes of
the

[177745]t(9;22) positive leukemias. Cancer Genet. Cytogenet. 11:
361–367,

[177746]1984.

[177747]

[177748]

[177749]18355.

[177750]Priest, J. R.; Robison, L. L.; McKenna, R. W.; Lindquist, L.
L.;

[177751]Warkentin, P. I.; LeBien, T. W.; Woods, W. G.; Kersey, J. H.;
Coccia,

[177752]P. F.; Nesbit, M. E., Jr.: Philadelphia chromosome positive
childhood

[177753]acute lymphoblastic leukemia. Blood 56: 15–22, 1980.

[177754]

[177755]

[177756]18356.

[177757]Rowley, J. D.: A new consistent chromosomal abnormality
in chronic

[177758]myelogenous leukemia identified by quinacrine fluorescence and Giemsa

[177759]staining. Nature 243: 290–293, 1973.

[177760]

[177761]

[177762]18357.

[177763]Rubin, C. M.; Carrino, J. J.; Dickler, M. N.; Leibowitz, D.;
Smith,

[177764]S. D.; Westbrook, C. A.: Heterogeneity of genomic fusion
of BCR and

[177765]ABL in Philadelphia chromosome–positive acute lymphoblastic leukemia. Proc.

[177766]Nat. Acad. Sci. 85: 2795–2799, 1988.

[177767]

[177768]

[177769]18358.

[177770]Saglio, G.; Storlazzi, C. T.; Giugliano, E.; Surace, C.; Anelli,

[177771]L.; Rege–Cambrin, G.; Zagaria, A.; Velasco, A. J.; Heiniger,
A.; Scaravaglio,

[177772]P.; Gomez, A. T.; Gomez, J. R.; Archidiacono, N.; Banfi, S.;

Rocchi,

[177773]M.: A 76-kb duplicon maps close to the BCR gene on chromosome 22

[177774]and the ABL gene on chromosome 9: possible involvement in the genesis

[177775]of the Philadelphia chromosome translocation. Proc. Nat. Acad. Sci. 99:

[177776]9882–9887, 2002.

[177777]

[177778]

[177779]18359.

[177780]Savage, D. G.; Antman, K. H.: Imatinib mesylate--a new oral targeted

[177781]therapy. New Eng. J. Med. 346: 683–693, 2002.

[177782]

[177783]

[177784]18360.

[177785]Sawyers, C. L.: Chronic myeloid leukemia. New Eng. J. Med. 340:

[177786]1330–1340, 1999.

[177787]

[177788]

[177789]18361.

[177790]Schaefer-Rego, K.; Dudek, H.; Popenoe, D.; Arlin, Z.;
Mears, J.

[177791]G.; Bank, A.; Leibowitz, D.: CML patients in blast crisis
have breakpoints

[177792]localized to a specific region of the BCR. Blood 70:
448–455, 1987.

[177793]

[177794]

[177795]18362.

[177796]Shtivelman, E.; Gale, R. P.; Dreazen, O.; Berrebi, A.; Zaizov,

[177797]R.; Kubonishi, I.; Miyoshi, I.; Canaani, E.: bcr-abl RNA in
patients

[177798]with chronic myelogenous leukemia. Blood 69: 971–973,
1987.

[177799]

[177800]

[177801]18363.

[177802]Shtivelman, E.; Lifshitz, B.; Gale, R. P.; Canaani, E.: Fused
[177803]transcript of abl and bcr genes in chronic myelogenous
leukaemia. Nature 315:
[177804]550–554, 1985.
[177805]
[177806]
[177807]18364.
[177808]Skorski, T.; Nieborowska-Skorska, M.; Nicolaides, N. C.;
Szczylik,
[177809]C.; Iversen, P.; Iozzo, R. V.; Zon, G.; Calabretta, B.: Sup-
pression
[177810]of Philadelphia-1 leukemia cell growth in mice by BCR-
ABL antisense
[177811]oligodeoxynucleotide. Proc. Nat. Acad. Sci. 91:
4504–4508, 1994.
[177812]
[177813]
[177814]18365.
[177815]Stam, K.; Heisterkamp, N.; Grosveld, G.; de Klein, A.;
Verma,

[177816]R. S.; Coleman, M.; Dosik, H.; Groffen, J.: Evidence of a new chimeric

[177817]bcr/c-abl mRNA in patients with chronic myelocytic leukemia and the

[177818]Philadelphia chromosome. New Eng. J. Med. 313: 1429–1433, 1985.

[177819]

[177820]

[177821]18366.

[177822]Stam, K.; Heisterkamp, N.; Reynolds, F. H., Jr.; Groffen, J.:

[177823]Evidence that the phl gene encodes a 160,000–dalton phosphoprotein

[177824]with associated kinase activity. Molec. Cell. Biol. 7: 1955–1960,

[177825]1987.

[177826]

[177827]

[177828]18367.

[177829]Swan, D. C.; McBride, O. W.; Robbins, K. C.; Keithley, D. A.;

[177830]Reddy, E. P.; Aaronson, S. A.: Chromosomal mapping of

the simian

[177831]sarcoma virus onc gene analogue in human cells. Proc.
Nat. Acad.

[177832]Sci. 79: 4691–4695, 1982.

[177833]

[177834]

[177835]18368.

[177836]Tanabe, T.; Kuwabara, T.; Warashina, M.; Tani, K.; Taira,
K.;

[177837]Asano, S.: Oncogene inactivation in a mouse model: tissue
invasion

[177838]by leukaemic cells is stalled by loading them with a de-
signer ribozyme. Nature 406:

[177839]473–474, 2000.

[177840]

[177841]

[177842]18369.

[177843]Teyssier, J. R.; Bartram, C. R.; Deville, J.; Potron, G.; Pi-
geon,

[177844]F.: C-abl oncogene and chromosome 22 'bcr' juxtaposi-

tion in chronic

[177845]myelogenous leukemia. New Eng. J. Med. 312:
1393–1394, 1985.

[177846]

[177847]

[177848]18370.

[177849]Tkachuk, D. C.; Westbrook, C. A.; Andreeff, M.; Donlon, T.
A.;

[177850]Cleary, M. L.; Suryanarayan, K.; Homge, M.; Redner, A.;
Gray, J.;

[177851]Pinkel, D.: Detection of bcr–abl fusion in chronic myeloge-
neous leukemia

[177852]by in situ hybridization. Science 250: 559–562, 1990.

[177853]

[177854]

[177855]18371.

[177856]Verhest, A.; Monsieur, R.: Philadelphia chromosome–posi-
tive thrombocythemia

[177857]with leukemic transformation. (Letter) New Eng. J. Med.
308: 1603,

[177858]1983.

[177859]

[177860]

[177861]18372.

[177862]Verma, R. S.; Dosik, H.: Heteromorphisms of the Philadelphia
phia

[177863](Ph-1) chromosome in patients with chronic myelogenous
leukaemia (CML).

[177864]I. Classification and clinical significance. Brit. J. Haemat.
45:

[177865]215-222, 1980.

[177866]

[177867]

[177868]18373.

[177869]Cockerham, G. C.; Hidayat, A. A.; Bijwaard, K. E.; Sheng,
Z.-M.

[177870]: Re-evaluation of 'reactive lymphoid hyperplasia of the
uvea': an

[177871]immunohistochemical and molecular analysis of 10 cases.
Ophthalmology 107:

[177872]151-158, 2000.

[177873]

[177874]

[177875]18374.

[177876]Guru, S. C.; Agarwal, S. K.; Manickam, P.; Olufemi, S.-E.;
Crabtree,

[177877]J. S.; Weisemann, J. M.; Kester, M. B.; Kim, Y. S.; Wang, Y.;
Emmert-Buck,

[177878]M. R.; Liotta, L. A.; Spiegel, A. M.; Boguski, M. S.; Roe, B.
A.;

[177879]Collins, F. S.; Marx, S. J.; Burns, L.; Chandrasekharappa, S.
C.:

[177880]A transcript map for the 2.8-Mb region containing the
multiple endocrine

[177881]neoplasia type 1 locus. *Genome Res.* 7: 725-735, 1997.

[177882]

[177883]

[177884]18375.

[177885]Van Leeuwen, A.: Di-allelic allo-antigenic systems on hu-
man T-lymphocyte

[177886]subsets. Thesis: London Hospital Medical College (pub.)
1982.

[177887]

[177888]

[177889]18376.

[177890]Walker, I. D.; Sandrin, M. S.; Hogarth, P. M.; Sutton, V. R.;

[177891]McKenzie, I. F. C.: Expression of Qa alloantigens on peripheral T

[177892]cells: the relationship of the Qa-m2, 7, 8, 9 specificities. Immunogenetics 24:

[177893]90-94, 1986.

[177894]

[177895]

[177896]18377.

[177897]Yokoyama, W. M.: The mother-child union: the case of missing-self

[177898]and protection of the fetus. Proc. Nat. Acad. Sci. 94: 5998-6000,

[177899]1997.

[177900]

[177901]

[177902]18378.

[177903]Frossard, P. M.; Lestringant, G. G.: Association between a
dimorphic

[177904]site on chromosome 12 and clinical diagnosis of hyper-
tension in three

[177905]independent populations. Clin. Genet. 48: 284–287, 1995.

[177906]

[177907]

[177908]18379.

[177909]Glenn, C. L.; Wang, W. Y. S.; Benjafield, A. V.; Morris, B. J.

[177910]: Linkage and association of tumor necrosis factor recep-
tor 2 locus

[177911]with hypertension, hypercholesterolemia and plasma shed
receptor. Hum.

[177912]Molec. Genet. 9: 1943–1949, 2000.

[177913]

[177914]

[177915]18380.

[177916]Lasky, L. A.; Singer, M. S.; Dowbenko, D.; Imai, Y.; Henzel,

W.

[177917]J.; Grimley, C.; Fennie, C.; Gillett, N.; Watson, S. R.; Rosen, S.

[177918]D.: An endothelial ligand for L-selectin is a novel mucin-like molecule. Cell 69:

[177919]927–938, 1992.

[177920]

[177921]

[177922]18381.

[177923]Lasky, L. A.; Singer, M. S.; Yednock, T. A.; Dowbenko, D.; Fennie,

[177924]C.; Rodriguez, H.; Nguyen, T.; Stachel, S.; Rosen, S. D.: Cloning

[177925]of a lymphocyte homing receptor reveals a lectin domain. Cell 56:

[177926]1045–1055, 1989.

[177927]

[177928]

[177929]18382.

[177930]Ord, D. C.; Ernst, T. J.; Zhou, L.-J.; Rambaldi, A.; Spertini,

[177931]O.; Griffin, J.; Tedder, T. F.: Structure of the gene encoding the

[177932]human leukocyte adhesion molecule-1 (TQ1, Leu-8) of lymphocytes and

[177933]neutrophils. J. Biol. Chem. 265: 7760-7767, 1990.

[177934]

[177935]

[177936]18383.

[177937]Siegelman, M. H.; Weissman, I. L.: Human homologue of mouse lymph

[177938]node homing receptor: evolutionary conservation at tandem cell interaction

[177939]domains. Proc. Nat. Acad. Sci. 86: 5562-5566, 1989.

[177940]

[177941]

[177942]18384.

[177943]Sitrin, R. G.; Pan, P. M.; Blackwood, R. A.; Huang, J.; Petty,

[177944]H. R.: Cutting edge: evidence for a signaling partnership between

[177945]urokinase receptors (CD87) and L-selectin (CD62L) in hu-

man polymorphonuclear

[177946]neutrophils. J. Immun. 166: 4822–4825, 2001.

[177947]

[177948]

[177949]18385.

[177950]Tedder, T. F.; Isaacs, C. M.; Ernst, T. J.; Demetri, G. D.;
Adler,

[177951]D. A.; Disteché, C. M.: Isolation and chromosomal localization of

[177952]cDNAs encoding a novel human lymphocyte cell surface
molecule, LAM-1:

[177953]homology with the mouse lymphocyte homing receptor
and other human

[177954]adhesion proteins. J. Exp. Med. 170: 123–133, 1989.

[177955]

[177956]

[177957]18386.

[177958]Aizawa, S.; Nakano, H.; Ishida, T.; Horie, R.; Nagai, M.; Ito,

[177959]K.; Yagita, H.; Okumura, K.; Inoue, J.; Watanabe, T.: Tumor
necrosis

[177960]factor receptor-associated factor (TRAF) 5 and TRAF2 are involved

[177961]in CD30-mediated NF-kappa-B activation. J. Biol. Chem. 272: 2042-2045,

[177962]1997.

[177963]

[177964]

[177965]18387.

[177966]Durkop, H.; Latza, U.; Hummel, M.; Eitelbach, F.; Seed, B.; Stein,

[177967]H.: Molecular cloning and expression of a new member of the nerve

[177968]growth factor receptor family that is characteristic for Hodgkin's

[177969]disease. Cell 68: 421-427, 1992.

[177970]

[177971]

[177972]18388.

[177973]Kurts, C.; Carbone, F. R.; Krummel, M. F.; Koch, K. M.; Miller,

[177974]J. F. A. P.; Heath, W. R.: Signalling through CD30 protects
against

[177975]autoimmune diabetes mediated by CD8 T cells. Nature
398: 341–344,

[177976]1999.

[177977]

[177978]

[177979]18389.

[177980]McClive, P. J.; Morahan, G.: Assignment of the mouse ho-
mologues

[177981]of 6 loci from HSA1p to chromosomes 3 and 4. Genomics
23: 243–246,

[177982]1994.

[177983]

[177984]

[177985]18390.

[177986]Stein, H.; Gerdes, J.; Schwab, U.; Lemke, H.; Mason, D. Y.;
Ziegler,

[177987]A.; Schienle, W.; Diehl, V.: Identification of Hodgkin and
Sternberg–Reed

[177988]cells as a unique cell type derived from a newly detected
small cell

[177989]population. *Int. J. Cancer* 30: 445–449, 1982.

[177990]

[177991]

[177992]18391.

[177993]de Lau, W.; Clevers, H.: LEF1 turns over a new leaf. *Nature*
Genet. 28:

[177994]3–5, 2001.

[177995]

[177996]

[177997]18392.

[177998]Hovanes, K.; Li, T. W. H.; Munguia, J. E.; Truong, T.; Milo-
vanovic,

[177999]T.; Marsh, J. L.; Holcombe, R. F.; Waterman, M. L.: Beta-
catenin-sensitive

[178000]isoforms of lymphoid enhancer factor-1 are selectively
expressed in

[178001]colon cancer. *Nature Genet.* 28: 53–57, 2001.

[178002]

[178003]

[178004]18393.

[178005]Love, J. J.; Li, X.; Case, D. A.; Giese, K.; Grosschedl, R.;
Wright,

[178006]P. E.: Structural basis for DNA bending by the architectural
transcription

[178007]factor LEF-1. Nature 376: 791-795, 1995.

[178008]

[178009]

[178010]18394.

[178011]Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.:
LEF1,

[178012]the gene for lymphoid enhancer-binding factor 1, mapped
to human chromosome

[178013]4 (q23-q25) and distal mouse chromosome 3 by Southern
blot analysis

[178014]and fluorescence in situ hybridization. (Abstract) Cyto-
genet. Cell

[178015]Genet. 58: 1888 only, 1991.

[178016]

[178017]

[178018]18395.

[178019]Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.: Gene

[178020]for lymphoid enhancer-binding factor 1 (LEF1) mapped to
human chromosome

[178021]4 (q23-q25) and mouse chromosome 3 near Egf. Ge-
nomics 11: 1040-1048,

[178022]1991.

[178023]

[178024]

[178025]18396.

[178026]van Genderen, C.; Okamura, R. M.; Farinas, I.; Quo, R. G.;
Parslow,

[178027]T. G.; Bruhn, L.; Grosschedl, R.: Development of several
organs that

[178028]require inductive epithelial-mesenchymal interactions is
impaired

[178029]in LEF-1-deficient mice. Genes Dev. 8: 2691-2703, 1994.

[178030]

[178031]

[178032]18397.

[178033]Waterman, M. L.; Fischer, W. H.; Jones, K. A.: A thymus-specific

[178034]member of the HMG protein family regulates the human T cell receptor

[178035]C alpha enhancer. Genes Dev. 5: 656-669, 1991.

[178036]

[178037]

[178038]18398.

[178039]Zhou, P.; Byrne, C.; Jacobs, J.; Fuchs, E.: Lymphoid enhancer

[178040]factor 1 directs hair follicle patterning and epithelial cell fate. Genes

[178041]Dev. 9: 700-713, 1995.

[178042]

[178043]

[178044]18399.

[178045]White, P. S.; Jensen, S. J.; Rajalingam, V.; Stairs, D.; Sulman,

[178046]E. P.; Maris, J. M.; Biegel, J. A.; Wooster, R.; Brodeur, G. M.:

[178047]Physical mapping of the CA6, ENO1, and SLC2A5 (GLUT5) genes and reassignment

[178048]of SLC2A5 to 1p36.2. Cytogenet. Cell Genet. 81: 60–64, 1998.

[178049]

[178050]

[178051]18400.

[178052]Contractor, A.; Swanson, G.; Heinemann, S. F.: Kainate receptors

[178053]are involved in short- and long-term plasticity at mossy fiber synapses

[178054]in the hippocampus. Neuron 29: 209–216, 2001.

[178055]

[178056]

[178057]18401.

[178058]Paschen, W.; Hedreen, J. C.; Ross, C. A.: RNA editing of the glutamate

[178059]receptor subunits GluR2 and GluR6 in human brain tissue. J. Neurochem. 63:

[178060]1596–1602, 1994.

[178061]

[178062]

[178063]18402.

[178064]Barbon, A.; Barlati, S.: Genomic organization, proposed alternative

[178065]splicing mechanisms, and RNA editing structure of GRIK1. Cytogenet.

[178066]Cell Genet. 88: 236–239, 2000.

[178067]

[178068]

[178069]18403.

[178070]Eubanks, J. H.; Puranam, R. S.; Kleckner, N. W.; Bettler, B.; Heinemann,

[178071]S. F.; McNamara, J. O.: The gene encoding the glutamate receptor

[178072]subunit GluR5 is located on human chromosome 21q21.1–22.1 in the vicinity

[178073]of the gene for familial amyotrophic lateral sclerosis. Proc. Nat.

[178074]Acad. Sci. 90: 178–182, 1993.

[178075]

[178076]

[178077]18404.

[178078]Gregor, P.; Gaston, S. M.; Yang, X.; O'Regan, J. P.; Rosen,
D.

[178079]R.; Tanzi, R. E.; Patterson, D.; Haines, J. L.; Horvitz, H. R.;
Uhl,

[178080]G. R.; Brown, R. H., Jr.: Genetic and physical mapping of
the GLUR5

[178081]glutamate receptor gene on human chromosome 21.
Hum. Genet. 94:

[178082]565–570, 1994.

[178083]

[178084]

[178085]18405.

[178086]Gregor, P.; Reeves, R. H.; Jabs, E. W.; Yang, X.; Dackowski,
W.;

[178087]Rochelle, J. M.; Brown, R. H., Jr.; Haines, J. L.; O'Hara, B. F.;

[178088]Uhl, G. R.; Seldin, M. F.: Chromosomal localization of glu-
tamate

[178089]receptor genes: relationship to familial amyotrophic lateral sclerosis

[178090]and other neurological disorders of mice and humans.

Proc. Nat. Acad.

[178091]Sci. 90: 3053–3057, 1993.

[178092]

[178093]

[178094]18406.

[178095]Gregor, P.; Seldin, M. F.; Reeve, R.; Jabs, E.; Yang, X.; Rochelle,

[178096]J. M.; O'Hara, B. F.; Uhl, G. R.: Chromosomal localization of glutamate

[178097]receptor genes: candidates for familial amyotrophic lateral sclerosis

[178098]and other neurological disorders of mice and man.

(Abstract) Neurosci.

[178099]Abstracts 18: 395 only, 1992.

[178100]

[178101]

[178102]18407.

[178103]Paschen, W.; Djuricic, B.: Extent of RNA editing of glutamate

[178104]receptor subunit GluR5 in different brain regions of the rat. Cell.

[178105]Molec. Neurobiol. 14: 259–270, 1994.

[178106]

[178107]

[178108]18408.

[178109]Potier, M.-C.; Dutriaux, A.; Lambolez, B.; Bochet, P.; Rossier,

[178110]J.: Assignment of the human glutamate receptor gene GLUR5 to 21q22

[178111]by screening a chromosome 21 YAC library. Genomics 15: 696–697,

[178112]1993.

[178113]

[178114]

[178115]18409.

[178116]Sander, T.; Hildmann, T.; Kretz, R.; Furst, R.; Sailer, U.; Bauer,

[178117]G.; Schmitz, B.; Beck–Mannagetta, G.; Wienker, T. F.; Janz, D.: Allelic

[178118]association of juvenile absence epilepsy with a gluR5 kainate receptor

[178119]gene (GRIK1) polymorphism. Am. J. Med. Genet. 74: 416–421, 1997.

[178120]

[178121]

[178122]18410.

[178123]Smolders, I.; Bortolotto, Z. A.; Clarke, V. R. J.; Warre, R.;

[178124]Khan, G. M.; O'Neill, M. J.; Ornstein, P. L.; Bleakman, D.; Ogden,

[178125]A.; Weiss, B.; Stables, J. P.; Ho, K. H.; Ebinger, G.; Collingridge,

[178126]G. L.; Lodge, D.; Michotte, Y.: Antagonists of GLU–K5–containing

[178127]kainate receptors prevent pilocarpine–induced limbic seizures. Nature

[178128]Neurosci. 5: 796–804, 2002.

[178129]

[178130]

[178131]18411.

[178132]McNamara, J. O.; Eubanks, J. H.; McPherson, J. D.; Wasmuth, J.

[178133]J.; Evans, G. A.; Heinemann, S. F.: Chromosomal localization of human

[178134]glutamate receptor genes. J. Neurosci. 12: 2555–2562, 1992.

[178135]

[178136]

[178137]18412.

[178138]Armstrong, N.; Gouaux, E.: Mechanisms for activation and antagonism

[178139]of an AMPA-sensitive glutamate receptor: crystal structures of the

[178140]GluR2 ligand binding core. Neuron 28: 165–181, 2000.

[178141]

[178142]

[178143]18413.

[178144]Brusa, R.; Zimmermann, F.; Koh, D.-S.; Feldmeyer, D.; Gass, P.;

[178145]Seeburg, P. H.; Sprengel, R.: Early-onset epilepsy and postnatal

[178146]lethality associated with an editing-deficient GluR-B allele in mice. Science 270:

[178147]1677–1680, 1995.

[178148]

[178149]

[178150]18414.

[178151]Hollmann, M.; Hartley, M.; Heinemann, S.: Ca(2+) permeability

[178152]of KA-AMPA-gated glutamate receptor channels depends on subunit composition. Science 252:

[178153]851–853, 1991.

[178154]

[178155]

[178156]18415.

[178157]Kohler, M.; Kornau, H. C.; Seeburg, P. H.: The organization of

[178158]the gene for the functionally dominant alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic

[178159]acid receptor subunit Glu4-B. J. Biol. Chem. 269:
17367-17370, 1994.

[178160]

[178161]

[178162]18416.

[178163]Sun, W.; Ferrer-Montiel, A. V.; Schinder, A. F.; McPherson,
J.

[178164]P.; Evans, G. A.; Montal, M.: Molecular cloning, chromoso-
mal mapping,

[178165]and functional expression of human brain glutamate re-
ceptors. Proc.

[178166]Nat. Acad. Sci. 89: 1443-1447, 1992.

[178167]

[178168]

[178169]18417.

[178170]Sun, Y.; Olson, R.; Horning, M.; Armstrong, N.; Mayer, M.;
Gouaux,

[178171]E.: Mechanism of glutamate receptor desensitization. Na-
ture 417:

[178172]245-253, 2002.

[178173]

[178174]

[178175]18418.

[178176]Chui, D.; Oh-Eda, M.; Liao, Y.-F.; Panneerselvam, K.; Lai, A.;

[178177]Marek, K. W.; Freeze, H. H.; Moremen, K. W.; Fukuda, M. N.; Marth,

[178178]J. D.: Alpha-mannosidase-II deficiency results in dyserythropoiesis

[178179]and unveils an alternate pathway in oligosaccharide biosynthesis. Cell 90:

[178180]157-167, 1997.

[178181]

[178182]

[178183]18419.

[178184]Chui, D.; Sellakumar, G.; Green, R. S.; Sutton-Smith, M.; McQuistan,

[178185]T.; Marek, K. W.; Morris, H. R.; Dell, A.; Marth, J. D.: Genetic

[178186]remodeling of protein glycosylation in vivo induces autoimmune disease. Proc.

[178187]Nat. Acad. Sci. 98: 1142–1147, 2001.

[178188]

[178189]

[178190]18420.

[178191]Gasparini, P.; del Giudice, E. M.; Delaunay, J.; Totaro, A.;
Granatiero,

[178192]M.; Melchionda, S.; Zelante, L.; Iolascon, A.: Localization of
the

[178193]congenital dyserythropoietic anemia II locus to chromo-
some 20q11.2

[178194]by genomewide search. Am. J. Hum. Genet. 61:
1112–1116, 1997.

[178195]

[178196]

[178197]18421.

[178198]Misago, M.; Liao, Y.-F.; Kudo, S.; Eto, S.; Mattei, M.-G.;
Moremen,

[178199]K. W.; Fukuda, M. N.: Molecular cloning and expression of
cDNAs encoding

[178200]human alpha-mannosidase II and a previously unrecog-
nized alpha-mannosidase

[178201]II(X) isozyme. Proc. Nat. Acad. Sci. 92: 11766–11770,
1995.

[178202]

[178203]

[178204]18422.

[178205]Moremen, K. W.; Robbins, P. W.: Isolation, characteriza-
tion, and

[178206]expression of cDNAs encoding murine alpha-
mannosidase II, a Golgi

[178207]enzyme that controls conversion of high mannose to
complex N-glycans. J.

[178208]Cell Biol. 115: 1521–1534, 1991.

[178209]

[178210]

[178211]18423.

[178212]Wada, A.; Sakamoto, H.; Katoh, O.; Yoshida, T.; Yokota, J.;
Little,

[178213]P. F. R.; Sugimura, T.; Terada, M.: Two homologous onco-
genes, HST1

[178214]and INT2, are closely located in human genome. Biochem.
Biophys.

[178215]Res. Commun. 157: 828–835, 1988.

[178216]

[178217]

[178218]18424.

[178219]Yoshida, T.; Tsutsumi, M.; Sakamoto, H.; Miyagawa, K.;
Teshima,

[178220]S.; Sugimura, T.; Terada, M.: Expression of the HST1
oncogene in

[178221]human germ cell tumors. Biochem. Biophys. Res. Com-
mun. 155: 1324–1329,

[178222]1988.

[178223]

[178224]

[178225]18425.

[178226]Nimmo, E.; Padua, R.–A.; Hughes, D.; Brook, J. D.;
Williamson,

[178227]R.; Johnson, K. J.: Confirmation and refinement of the lo-
calisation

[178228]of the c-MEL locus on chromosome 19 by physical and
genetic mapping. Hum.

[178229]Genet. 81: 382–384, 1989.

[178230]

[178231]

[178232]18426.

[178233]Nimmo, E.; Williamson, R.; Johnson, K.: Localization of the
c-MEL

[178234]gene to 19(cen-p13.2). (Abstract) Cytogenet. Cell Genet.
51: 1053

[178235]only, 1989.

[178236]

[178237]

[178238]18427.

[178239]Nimmo, E. R.; Sanders, P. G.; Padua, R. A.; Hughes, D.;
Williamson,

[178240]R.; Johnson, K. J.: The MEL gene: a new member of the
RAB/YPT class

[178241]of RAS-related genes. Oncogene 6: 1347–1351, 1991.

[178242]

[178243]

[178244]18428.

[178245]Padua, R. A.; Barrass, N.; Currie, G. A.: A novel transform-
ing

[178246]gene in a human malignant melanoma cell line. Nature
311: 671-673,

[178247]1984.

[178248]

[178249]

[178250]18429.

[178251]Spurr, N. K.; Hughes, D.; Goodfellow, P. N.; Brook, J. D.;
Padua,

[178252]R. A.: Chromosomal assignment of c-MEL, a human trans-
forming oncogene,

[178253]to chromosome 19(p13.2-q13.2). Somat. Cell Molec.
Genet. 12: 637-640,

[178254]1986.

[178255]

[178256]

[178257]18430.

[178258]Dabiri, G. A.; Young, C. L.; Rosenbloom, J.; Southwick, F.

S.:

[178259]Molecular cloning of human macrophage capping protein
cDNA: a unique

[178260]member of the gelsolin/villin family expressed primarily in
macrophages. J.

[178261]Biol. Chem. 267: 16545–16552, 1992.

[178262]

[178263]

[178264]18431.

[178265]Mishra, V. S.; Henske, E. P.; Kwiatkowski, D. J.; Southwick,
F.

[178266]S.: The human actin–regulatory protein cap G: gene struc-
ture and

[178267]chromosome location. Genomics 23: 560–565, 1994.

[178268]

[178269]

[178270]18432.

[178271]Southwick, F. S.; DiNubile, M. J.: Rabbit alveolar
macrophages

[178272]contain a Ca^{2+} –sensitive, 41,000–Dalton protein which

reversibly

[178273]blocks the 'barbed' ends of actin filaments but does not sever them. J.

[178274]Biol. Chem. 261: 14191–14195, 1986.

[178275]

[178276]

[178277]18433.

[178278]Eichbaum, Q.; Clerc, P.; Bruns, G.; McKeon, F.; Ezekowitz, R. A.

[178279]B.: Assignment of the human macrophage mannose receptor gene (MRC1)

[178280]to 10p13 by in situ hybridization and PCR-based somatic cell hybrid

[178281]mapping. Genomics 22: 656–658, 1994.

[178282]

[178283]

[178284]18434.

[178285]Harris, N.; Peters, L. L.; Eicher, E. M.; Rits, M.; Raspberry,

[178286]D.; Eichbaum, Q. G.; Super, M.; Ezekowitz, R. A. B.: The exon–intron

[178287]structure and chromosomal localization of the mouse
macrophage mannose

[178288]receptor gene, Mcr1: identification of a ricin-like domain
at the

[178289]N-terminus of the receptor. Biochem. Biophys. Res. Com-
mun. 198:

[178290]682-692, 1994.

[178291]

[178292]

[178293]18435.

[178294]Kim, S. J.; Ruiz, N.; Bezouska, K.; Drickamer, K.: Organiza-
tion

[178295]of the gene encoding the human macrophage mannose
receptor (MRC1). Genomics 14:

[178296]721-727, 1992.

[178297]

[178298]

[178299]18436.

[178300]Lee, S. J.; Evers, S.; Roeder, D.; Parlow, A. F.; Risteli, J.;

[178301]Risteli, L.; Lee, Y. C.; Feizi, T.; Langen, H.; Nussenzweig,

M. C.

[178302]: Mannose receptor-mediated regulation of serum glycoprotein homeostasis. Science 295:

[178303]1898–1901, 2002.

[178304]

[178305]

[178306]18437.

[178307]Bernhagen, J.; Calandra, T.; Mitchell, R. A.; Martin, S. B.; Tracey,

[178308]K. J.; Voelter, W.; Manogue, K. R.; Cerami, A.; Bucala, R.: MIF is

[178309]a pituitary-derived cytokine that potentiates lethal endotoxaemia. Nature 365:

[178310]756–759, 1993.

[178311]

[178312]

[178313]18438.

[178314]Bloom, B. R.; Bennett, B.: Mechanism of a reaction in vitro associated

[178315]with delayed-type hypersensitivity. Science 153: 80–82,

1966.

[178316]

[178317]

[178318]18439.

[178319]Bozza, M.; Kolakowski, L. F., Jr.; Jenkins, N. A.; Gilbert, D.

[178320]J.; Copeland, N. G.; David, J. R.; Gerard, C.: Structural
characterization

[178321]and chromosomal location of the mouse macrophage mi-
gration inhibitory

[178322]factor gene and pseudogenes. Genomics 27: 412–419,
1995.

[178323]

[178324]

[178325]18440.

[178326]Bucala, R.: MIF rediscovered: cytokine, pituitary hormone,
and

[178327]glucocorticoid-induced regulator of the immune re-
sponse. FASEB J. 10:

[178328]1607–1613, 1996.

[178329]

[178330]

[178331]18441.

[178332]Budarf, M.; McDonald, T.; Sellinger, B.; Kozak, C.; Graham, C.;

[178333]Wistow, G.: Localization of the human gene for
macrophage migration

[178334]inhibitory factor (MIF) to chromosome 22q11.2. Genomics
39: 235–236,

[178335]1997.

[178336]

[178337]

[178338]18442.

[178339]Cato, A. C. B.; Sillar, G. M.; Kioussis, J.; Burdon, R. H.:
Molecular

[178340]cloning of cDNA sequences coding for the major (beta–,
gamma–, delta–,

[178341]and epsilon) heat–shock polypeptides of HeLa cells. Gene
16: 27–34,

[178342]1981.

[178343]

[178344]

[178345]18443.

[178346]Cummings, C. J.; Sun, Y.; Opal, P.; Antalffy, B.; Mestril, R.;

[178347]Orr, H. T.; Dillmann, W. H.; Zoghbi, H. Y.: Over-expression
of inducible

[178348]HSP70 chaperone suppresses neuropathology and im-
proves motor function

[178349]in SCA1 mice. Hum. Molec. Genet. 10: 1511–1518, 2001.

[178350]

[178351]

[178352]18444.

[178353]Gaskins, H. R.; Prochazka, M.; Nadeau, J. H.; Henson, V.
W.; Leiter,

[178354]E. H.: Localization of a mouse heat shock Hsp70 gene
within the H-2

[178355]complex. Immunogenetics 32: 286–289, 1990.

[178356]

[178357]

[178358]18445.

[178359]Goate, A. M.; Cooper, D. N.; Hall, C.; Leung, T. K. C.;
Solomon,

[178360]E.; Lim, L.: Localization of a human heat-shock HSP70
gene sequence

[178361]to chromosome 6 and detection of two other loci by so-
matic-cell hybrid

[178362]and restriction fragment length polymorphism analysis.
Hum. Genet. 75:

[178363]123-128, 1987.

[178364]

[178365]

[178366]18446.

[178367]Fukai, K.; Oh, J.; Frenk, E.; Almodovar, C.; Spritz, R. A.:
Linkage

[178368]disequilibrium mapping of the gene for Hermansky-Pud-
lak syndrome to

[178369]chromosome 10q23.1-q23.3. Hum. Molec. Genet. 4:
1665-1669, 1995.

[178370]

[178371]

[178372]18447.

[178373]Gwynn, B.; Eicher, E. M.; Peters, L. L.: Genetic localization
[178374]of Cd63, a member of the transmembrane 4 superfamily,
reveals two
[178375]distinct loci in the mouse genome. Genomics 35:
389–391, 1996.

[178376]

[178377]

[178378]18448.

[178379]Hotta, H.; Ross, A. H.; Huebner, K.; Isobe, M.; Wendeborn,
S.;

[178380]Chao, M. V.; Ricciardi, R. P.; Tsujimoto, Y.; Croce, C. M.;
Koprowski,

[178381]H.: Molecular cloning and characterization of an antigen
associated

[178382]with early stages of melanoma tumor progression. Cancer
Res. 48:

[178383]2955–2962, 1988.

[178384]

[178385]

[178386]18449.

[178387]Metzelaar, M. J.; Wijngaard, P. L. J.; Peters, P. J.; Sixma, J.
[178388]J.; Nieuwenhuis, H. K.; Clevers, H. C.: CD63 antigen: a
novel lysosomal
[178389]membrane glycoprotein, cloned by a screening procedure
for intracellular
[178390]antigens in eukaryotic cells. J. Biol. Chem. 266:
3239–3245, 1991.
[178391]
[178392]
[178393]18450.
[178394]Nishibori, M.; Cham, B.; McNicol, A.; Shalev, A.; Jain, N.;
Gerrard,
[178395]J. M.: The protein CD63 is in platelet dense granules, is
deficient
[178396]in a patient with Hermansky–Pudlak syndrome, and ap-
pears identical
[178397]to granulophysin. J. Clin. Invest. 91: 1775–1782, 1993.
[178398]
[178399]
[178400]18451.

- [178401]Azizi, E.; Friedman, J.; Pavlovsky, F.; Iscovich, J.; Bornstein,
[178402]A.; Shafir, R.; Trau, H.; Brenner, H.; Nass, D.: Familial cu-
taneous
[178403]malignant melanoma and tumors of the nervous system.
Cancer 76:
[178404]1571–1578, 1995.
- [178405]
- [178406]
- [178407]18452.
- [178408]Doege, K.; Rhodes, C.; Sasaki, M.; Hassell, J. R.; Yamada,
Y.:
[178409]Molecular biology of cartilage proteoglycan (aggrecan)
and link protein.In:
[178410]Sandel, L. J.; Boyd, C. D.: Extracellular Matrix Genes. New
York:
[178411]Academic Press (pub.) 1990. Pp. 137–152.
- [178412]
- [178413]
- [178414]18453.
- [178415]Finkelstein, J. E.; Doege, K.; Yamada, Y.; Pyeritz, R. E.;

Graham,

[178416]J. M., Jr.; Moeschler, J. B.; Pauli, R. M.; Hecht, J. T.; Franco-
mano,

[178417]C. A.: Analysis of the chondroitin sulfate proteoglycan
core protein

[178418](CSPGCP) gene in achondroplasia and pseudoachon-
droplasia. Am. J.

[178419]Hum. Genet. 48: 97-102, 1991.

[178420]

[178421]

[178422]18454.

[178423]Just, W.; Klett, C.; Vetter, U.; Vogel, W.: Assignment of the

[178424]human aggrecan gene AGC1 to 15q25-q26.2 by in situ
hybridization. Hum.

[178425]Genet. 92: 516-518, 1993.

[178426]

[178427]

[178428]18455.

[178429]Kimata, K.; Barrach, H. J.; Brown, K. S.; Pennypacker, J. P.:

[178430]Absence of proteoglycan core protein in cartilage from the
cmd/cmd

[178431](cartilage deficiency) mouse. J. Biol. Chem. 256:
6961–6968, 1981.

[178432]

[178433]

[178434]18456.

[178435]Korenberg, J. R.; Chen, X. N.; Doege, K.; Grover, J.; Rough-
ley,

[178436]P. J.: Assignment of the human aggrecan gene (AGC1) to
15q26 using

[178437]fluorescence in situ hybridization analysis. Genomics 16:
546–548,

[178438]1993.

[178439]

[178440]

[178441]18457.

[178442]Lane, P. W.: New mutants and linkages: small with kinky
tail. Mouse

[178443]Newsletter 80: 165, 1988.

[178444]

[178445]

[178446]18458.

[178447]Watanabe, H.; Kimata, K.; Line, S.; Strong, D.; Gao, L.;
Kozak,

[178448]C. A.; Yamada, Y.: Mouse cartilage matrix deficiency (cmd)
caused

[178449]by a 7 bp deletion in the aggrecan gene. Nature Genet. 7:
154–157,

[178450]1994.

[178451]

[178452]

[178453]18459.

[178454]Watanabe, H.; Nakata, K.; Kimata, K.; Nakanishi, I.; Ya-
mada, Y.

[178455]: Dwarfism and age-associated spinal degeneration of
heterozygote

[178456]cmd mice defective in aggrecan. Proc. Nat. Acad. Sci. 94:
6943–6947,

[178457]1997.

[178458]

[178459]

[178460]18460.

[178461]Fong, L. Y. Y.; Fidanza, V.; Zanesi, N.; Lock, L. F.; Siracusa,

[178462]L. D.; Mancini, R.; Siprashvili, Z.; Ottey, M.; Martin, S. E.;

Druck,

[178463]T.; McCue, P. A.; Croce, C. M.; Huebner, K.: Muir–

Torre–like syndrome

[178464]in Fhit–deficient mice. Proc. Nat. Acad. Sci. 97:

4742–4747, 2000.

[178465]

[178466]

[178467]18461.

[178468]Chemelli, R. M.; Willie, J. T.; Sinton, C. M.; Elmquist, J. K.;

[178469]Scammell, T.; Lee, C.; Richardson, J. A.; Williams, S. C.;

Xiong,

[178470]Y.; Kisanuki, Y.; Fitch, T. E.; Nakazato, M.; Hammer, R. E.;

Saper,

[178471]C. B.; Yanagisawa, M.: Narcolepsy in orexin knockout

mice: molecular

[178472]genetics of sleep regulation. Cell 437–451, 1999.

[178473]

[178474]

[178475]18462.

[178476]Azen, E. A.; Denniston, C.: Polymorphism of Ps (parotid size variant)

[178477]and detection of a protein (PmS) related to the Pm (parotid middle

[178478]band) system with genetic linkage of Ps and Pm to G1, Db, and Pr genetic

[178479]determinants. Biochem. Genet. 18: 483–501, 1980.

[178480]

[178481]

[178482]18463.

[178483]Adkins, S.; Gan, K. N.; Mody, M.; La Du, B. N.: Molecular basis

[178484]for the polymorphic forms of human serum paraoxonase/arylesterase:

[178485]glutamine or arginine at position 191, for the respective A and B

[178486]allozymes. Am. J. Hum. Genet. 52: 598–608, 1993.

[178487]

[178488]

[178489]18464.

[178490]Antikainen, M.; Murtomaki, S.; Syvanne, M.; Pahlman, R.;
Tahvanainen,

[178491]E.; Jauhiainen, M.; Frick, M. H.; Ehnholm, C.: The gln-
arg191 polymorphism

[178492]of the human paraoxonase gene (HUMPONA) is not asso-
ciated with the

[178493]risk of coronary artery disease in Finns. J. Clin. Invest. 98:
883–885,

[178494]1996.

[178495]

[178496]

[178497]18465.

[178498]Augustinsson, K.-B.; Henricson, B.: A genetically con-
trolled esterase

[178499]in rat plasma. Biochim. Biophys. Acta 124: 323–331,
1966.

[178500]

[178501]

[178502]18466.

[178503]Barbieri, M.; Bonafe, M.; Marfella, R.; Ragno, E.; Giugliano, D.;

[178504]Franceschi, C.; Paolisso, G.: LL–paraoxonase genotype is associated

[178505]with a more severe degree of homeostasis model assessment IR in healthy

[178506]subjects. J. Clin. Endocr. Metab. 87: 222–225, 2002.

[178507]

[178508]

[178509]18467.

[178510]Brophy, V. H.; Jampsa, R. L.; Clendenning, J. B.; McKinstry, L.

[178511]A.; Jarvik, G. P.; Furlong, C. E.: Effects of 5–prime regulatory–region

[178512]polymorphisms on paraoxonase–gene (PON1) expression. Am. J. Hum.

[178513]Genet. 68: 1428–1436, 2001.

[178514]

[178515]

[178516]18468.

[178517]Cherry, N.; Mackness, M.; Durrington, P.; Povey, A.; Dipp-
nall,

[178518]M.; Smith, T.; Mackness, B.: Paraoxonase (PON1) polymor-
phisms in

[178519]farmers attributing ill health to sheep dip. Lancet 359:
763–764,

[178520]2002.

[178521]

[178522]

[178523]18469.

[178524]Clendenning, J. B.; Humbert, R.; Green, E. D.; Wood, C.;
Traver,

[178525]D.; Furlong, C. E.: Structural organization of the human
PON1 gene. Genomics 35:

[178526]586–589, 1996.

[178527]

[178528]

[178529]18470.

[178530]Coates, P. M.; Mestriner, M. A.; Hopkinson, D. A.: A preliminary

[178531]interpretation of the esterase isozymes of human tissues.
Ann. Hum.

[178532]Genet. 39: 1-20, 1975.

[178533]

[178534]

[178535]18471.

[178536]Davies, H. G.; Richter, R. J.; Keifer, M.; Broomfield, C. A.;
Sowalla,

[178537]J.; Furlong, C. E.: The effect of the human serum paraox-
onase polymorphism

[178538]is reversed with diazoxon, soman and sarin. Nature Genet.
14: 334-336,

[178539]1996.

[178540]

[178541]

[178542]18472.

[178543]Deakin, S.; Leviev, I.; Nicaud, V.; Meynet, M.-C. B.; Tiret,

L.;

[178544]James, R. W.: Paraoxonase-1 L55M polymorphism is associated with

[178545]an abnormal oral glucose tolerance test and differentiates high risk

[178546]coronary disease families. J. Clin. Endocr. Metab. 87: 1268-1273,

[178547]2002.

[178548]

[178549]

[178550]18473.

[178551]Eckerson, H. W.; Romson, J.; Wyte, C.; La Du, B. N.: The human

[178552]serum paraoxonase polymorphism: identification of phenotypes by their

[178553]response to salts. Am. J. Hum. Genet. 35: 214-227, 1983.

[178554]

[178555]

[178556]18474.

[178557]Eckerson, H. W.; Wyte, C. M.; La Du, B. N.: The human

serum paraoxonase/arylesterase

[178558]polymorphism. Am. J. Hum. Genet. 35: 1126–1136, 1983.

[178559]

[178560]

[178561]18475.

[178562]Eiberg, H.; Mohr, J.: Linkage relations of the paraoxonase
polymorphism

[178563]with 43 marker systems. (Abstract) Cytogenet. Cell Genet.
25: 150,

[178564]1979.

[178565]

[178566]

[178567]18476.

[178568]Rey–Campos, J.; Chouard, T.; Yaniv, M.; Cereghini, S.:
vHNF1

[178569]is a homeoprotein that activates transcription and forms
heterodimers

[178570]with HNF1. EMBO J. 10: 1445–1457, 1991.

[178571]

[178572]

[178573]18477.

[178574]Shih, D. Q.; Bussen, M.; Sehayek, E.; Ananthanarayanan,
M.; Shneider,

[178575]B. L.; Suchy, F. J.; Shefer, S.; Bollileni, J. S.; Gonzalez, F. J.;

[178576]Breslow, J. L.; Stoffel, M.: Hepatocyte nuclear factor-
1-alpha is

[178577]an essential regulator of bile acid and plasma cholesterol
metabolism. Nature

[178578]Genet. 27: 375-382, 2001.

[178579]

[178580]

[178581]18478.

[178582]Urhammer, S. A.; Hansen, T.; Ekstrom, C. T.; Eiberg, H.;
Pederson,

[178583]O.: The Ala/Val98 polymorphism of the hepatocyte nu-
clear factor-1-alpha

[178584]gene contributes to the interindividual variation in serum
C-peptide

[178585]response during an oral glucose tolerance test: evidence
from studies

[178586]of 231 glucose-tolerant first degree relatives of type 2 diabetic

[178587]probands. J. Clin. Endocr. Metab. 83: 4506–4509, 1998.

[178588]

[178589]

[178590]18479.

[178591]Urhammer, S. A.; Moller, A. M.; Nyholm, B.; Ekstrom, C. T.; Eiberg,

[178592]H.; Clausen, J. O.; Hansen, T.; Pedersen, O.; Schmitz, O.: The effect

[178593]of two frequent amino acid variants of the hepatocyte nuclear factor-1-alpha

[178594]gene on estimates of the pancreatic beta-cell function in Caucasian

[178595]glucose-tolerant first-degree relatives of type 2 diabetic patients. J.

[178596]Clin. Endocr. Metab. 83: 3992–3995, 1998.

[178597]

[178598]

[178599]18480.

- [178600]Urhammer, S. A.; Rasmussen, S. K.; Kaisaki, P. J.; Oda, N.; Yamagata, K.; Moller, A. M.; Fridberg, M.; Hansen, L.; Hansen, T.; Bell, G.
- [178602]J.; Pedersen, O.: Genetic variation in the hepatocyte nuclear factor-1-alpha gene in Danish Caucasians with late-onset NIDDM. Diabetesologia 40: 473-475, 1997.
- [178605]
- [178606]
- [178607]18481.
- [178608]Vaxillaire, M.; Rouard, M.; Yamagata, K.; Oda, N.; Kaisaki, P.
- [178609]J.; Boriraj, V. V.; Chevre, J.-C.; Boccio, V.; Cox, R. D.; Lathrop, G. M.; Dussoix, P.; Philippe, J.; Timsit, J.; Charpentier, G.; Velho, G.; Bell, G. I.; Froguel, P.: Identification of nine novel mutations in the hepatocyte nuclear factor 1 alpha gene associated with maturity-onset

[178613]diabetes of the young (MODY3). Hum. Molec. Genet. 6:
583–586, 1997.

[178614]

[178615]

[178616]18482.

[178617]Yamada, S.; Nishigori, H.; Onda, H.; Utsugi, T.; Yanagawa,
T.;

[178618]Maruyama, T.; Onigata, K.; Nagashima, K.; Nagai, R.;
Morikawa, A.;

[178619]Takeuchi, T.; Takeda, J.: Identification of mutations in the
hepatocyte

[178620]nuclear factor (HNF)–1–alpha gene in Japanese subjects
with IDDM. Diabetes 46:

[178621]1643–1647, 1997.

[178622]

[178623]

[178624]18483.

[178625]Yamagata, K.; Oda, N.; Kalsaki, P. J.; Menzel, S.; Furuta, H.;

[178626]Vaxillaire, M.; Southam, L.; Cox, R. D.; Lathrop, G. M.;
Borhaj, V.

[178627]V.; Chen, X.; Cox, N. J.; Oda, Y.; Yano, H.; Le Beau, M. M.; Yamada,

[178628]S.; Nishigori, H.; Takeda, J.; Fajans, S. S.; Hattersley, A. T.; Iwasaki,

[178629]N.; Hansen, T.; Pedersen, O.; Polonsky, K. S.; Turner, R. C.; Velho,

[178630]G.; Chevre, J.-C.; Froguel, P.; Bell, G. I.: Mutations in the hepatocyte

[178631]nuclear factor-1-alpha gene in maturity-onset diabetes of the young

[178632](MODY3). Nature 384: 455-457, 1996.

[178633]

[178634]

[178635]18484.

[178636]Yoshiuchi, I.; Yamagata, K.; Yoshimoto, M.; Zhu, Q.; Yang, Q.;

[178637]Nammo, T.; Uenaka, R.; Kinoshita, E.; Hanafusa, T.; Miyagawa, J.;

[178638]Matsuzawa, Y.: Analysis of a non-functional HNF-1-alpha (TCF1) mutation

[178639]in Japanese subjects with familial type 1 diabetes. Hum. Mutat. 18:

[178640]345–351, 2001.

[178641]

[178642]

[178643]18485.

[178644]Dhanasekaran, S. M.; Barrette, T. R.; Ghosh, D.; Shah, R.;
Varambally,

[178645]S.; Kurachi, K.; Pienta, K. J.; Rubin, M. A.; Chinnaiyan, A.
M.:

[178646]Delineation of prognostic biomarkers in prostate cancer.
Nature 412:

[178647]822–826, 2001.

[178648]

[178649]

[178650]18486.

[178651]Leytus, S. P.; Loeb, K. R.; Hagen, F. S.; Kurachi, K.; Davie,
E.

[178652]W.: A novel trypsin–like serine protease (hepsin) with a
putative

[178653]transmembrane domain expressed by human liver and
hepatoma cells. Biochemistry 27:

[178654]1067–1074, 1988.

[178655]

[178656]

[178657]18487.

[178658]Tsuji, A.; Torres–Rosado, A.; Arai, T.; Le Beau, M. M.;

Lemons,

[178659]R. S.; Chou, S.–H.; Kurachi, K.: Hepsin, a cell membrane–
associated

[178660]protease: characterization, tissue distribution, and gene
localization. J.

[178661]Biol. Chem. 266: 16948–16953, 1991.

[178662]

[178663]

[178664]18488.

[178665]Wu, Q.; Yu, D.; Post, J.; Halks–Miller, M.; Sadler, J. E.;

Morser,

[178666]J.: Generation and characterization of mice deficient in
hepsin,

[178667]a hepatic transmembrane serine protease. J. Clin. Invest.
101: 321–326,

[178668]1998.

[178669]

[178670]

[178671]18489.

[178672]Cannella, B.; Hoban, C. J.; Gao, Y.-L.; Garcia-Arenas, R.;
Lawson,

[178673]D.; Marchionni, M.; Gwynne, D.; Raine, C. S.: The neureg-
ulin, glial

[178674]growth factor 2, diminishes autoimmune demyelination
and enhances

[178675]remyelination in a chronic relapsing model for multiple
sclerosis. Proc.

[178676]Nat. Acad. Sci. 95: 10100–10105, 1998.

[178677]

[178678]

[178679]18490.

[178680]Falls, D. L.; Rosen, K. M.; Corfas, G.; Lane, W. S.; Fis-
chbach,

[178681]G. D.: ARIA, a protein that stimulates acetylcholine recep-
tor synthesis,

[178682]is a member of the Neu ligand family. Cell 72: 801–815,
1993.

[178683]

[178684]

[178685]18491.

[178686]Fernandez, P.–A.; Tang, D. G.; Cheng, L.; Prochiantz, A.;
Mudge,

[178687]A. W.; Raff, M. C.: Evidence that axon–derived neuregulin
promotes

[178688]oligodendrocyte survival in the developing rat optic nerve.
Neuron 28:

[178689]81–90, 2000.

[178690]

[178691]

[178692]18492.

[178693]Holmes, W. E.; Sliwkowski, M. X.; Akita, R. W.; Henzel, W.
J.;

[178694]Lee, J.; Park, J. W.; Yansura, D.; Abadi, N.; Raab, H.; Lewis,
G.

[178695]D.; Shepard, H. M.; Kuang, W.–J.; Wood, W. I.; Goeddel, D.
V.; Vandlen,

[178696]R. L.: Identification of heregulin, a specific activator of
p185(erbB2). Science 256:

[178697]1205–1210, 1992.

[178698]

[178699]

[178700]18493.

[178701]Huang, Y. Z.; Won, S.; Ali, D. W.; Wang, Q.; Tanowitz, M.;
Du,

[178702]Q. S.; Pelkey, K. A.; Yang, D. J.; Xiong, W. C.; Salter, M. W.;
Mei,

[178703]L.: Regulation of neuregulin signaling by PSD–95 interact-
ing with

[178704]ErbB4 at CNS synapses. Neuron 26: 443–455, 2000.

[178705]

[178706]

[178707]18494.

[178708]Lee, J.; Wood, W. I.: Assignment of heregulin (HGL) to hu-
man chromosome

[178709]8p22–p11 by PCR analysis of somatic cell hybrid DNA.
Genomics 16:

[178710]790–791, 1993.

[178711]

[178712]

[178713]18495.

[178714]Martinou, J.–C.; Falls, D. L.; Fischbach, G. D.; Merlie, J. P.

[178715]: Acetylcholine receptor–inducing activity stimulates expression of

[178716]the epsilon–subunit gene of the muscle acetylcholine receptor. Proc.

[178717]Nat. Acad. Sci. 88: 7669–7673, 1991.

[178718]

[178719]

[178720]18496.

[178721]Ni, C.–Y.; Murphy, M. P.; Golde, T. E.; Carpenter, G.:

Gamma–secretase

[178722]cleavage and nuclear localization of ErbB–4 receptor tyrosine kinase. Science 294:

[178723]2179–2181, 2001.

[178724]

[178725]

[178726]18497.

[178727]Orr-Urtreger, A.; Trakhtenbrot, L.; Ben-Levy, R.; Wen, D.;
Rechavi,

[178728]G.; Lonai, P.; Yarden, Y.: Neural expression and chromo-
somal mapping

[178729]of NEU differentiation factor to 8p12-p21. Proc. Nat.
Acad. Sci. 90:

[178730]1867-1871, 1993.

[178731]

[178732]

[178733]18498.

[178734]Boyd, D.; Jain, S. K.; Crampton, J.; Barrett, K. J.; Drysdale,

[178735]J.: Isolation and characterization of a cDNA clone for hu-
man ferritin

[178736]heavy chain. Proc. Nat. Acad. Sci. 81: 4751-4755, 1984.

[178737]

[178738]

[178739]18499.

[178740]Caskey, J. H.; Jones, C.; Miller, Y. E.; Seligman, P. A.: Human
man

[178741]ferritin gene is assigned to chromosome 19. Proc. Nat.
Acad. Sci. 80:

[178742]482–486, 1983.

[178743]

[178744]

[178745]18500.

[178746]Costanzo, F.; Colombo, M.; Staempfli, S.; Santoro, C.;
Marone,

[178747]M.; Frank, R.; Delius, H.; Cortese, R.: Structure of gene and
pseudogenes

[178748]of human apoferritin H. Nucleic Acids Res. 14: 721–736,
1986.

[178749]

[178750]

[178751]18501.

[178752]Cragg, S. J.; Drysdale, J.; Worwood, M.: Genes for the 'H'
subunit

[178753]of human ferritin are present on a number of human
chromosomes. Hum.

[178754]Genet. 71: 108–112, 1985.

[178755]

[178756]

[178757]18502.

[178758]Eisenstein, R. S.: Iron regulatory proteins and the molecular

[178759]control of mammalian iron metabolism. Annu. Rev. Nutr.
20: 627–662,

[178760]2000.

[178761]

[178762]

[178763]18503.

[178764]Ferreira, C.; Bucchini, D.; Martin, M.–E.; Levi, S.; Arosio, P.;

[178765]Grandchamp, B.; Beaumont, C.: Early embryonic lethality
of H ferritin

[178766]gene deletion in mice. J. Biol. Chem. 275: 3021–3024,
2000.

[178767]

[178768]

[178769]18504.

[178770]Ferreira, C.; Santambrogio, P.; Martin, M.-E.; Andrieu, V.;
Feldmann,

[178771]G.; Henin, D.; Beaumont, C.: H ferritin knockout mice: a
model of

[178772]hyperferritinemia in the absence of iron overload. Blood
98: 525-532,

[178773]2001.

[178774]

[178775]

[178776]18505.

[178777]Gailani, M. R.; Petty, E. M.; Horsthemke, B.; Arnold, A.;
Marx,

[178778]S. J.; Bale, A. E.: Physical mapping of chromosome
11q12-13 by pulsed

[178779]field gel electrophoresis (PFGE). (Abstract) Cytogenet. Cell
Genet. 58:

[178780]1959 only, 1991.

[178781]

[178782]

[178783]18506.

[178784]Gatti, R. A.; Shaked, R.; Mohandas, T. K.; Salser, W.: Human
man

[178785]ferritin genes: chromosomal assignments and polymor-
phisms. Am. J.

[178786]Hum. Genet. 41: 654–667, 1987.

[178787]

[178788]

[178789]18507.

[178790]Harrison, P. M.; Arosio, P.: The ferritins: molecular proper-
ties,

[178791]iron storage function and cellular regulation. Biochim.
Biophys.

[178792]Acta 1275: 161–203, 1996.

[178793]

[178794]

[178795]18508.

[178796]Hentze, M. W.; Keim, S.; Papadopoulos, P.; O'Brien, S.;
Modi,

[178797]W.; Drysdale, J.; Leonard, W. J.; Harford, J. B.; Klausner, R.

D.

[178798]: Cloning, characterization, expression, and chromosomal
localization

[178799]of a human ferritin heavy-chain gene. Proc. Nat. Acad. Sci.
83:

[178800]7226–7230, 1986.

[178801]

[178802]

[178803]18509.

[178804]Kato, J.; Fujikawa, K.; Kanda, M.; Fukuda, N.; Sasaki, K.;
Takayama,

[178805]T.; Kobune, M.; Takada, K.; Takimoto, R.; Hamada, H.;
Ikeda, T.; Niitsu,

[178806]Y.: A mutation, in the iron-responsive element of H fer-
ritin mRNA,

[178807]causing autosomal dominant iron overload. Am. J. Hum.
Genet. 69:

[178808]191–197, 2001.

[178809]

[178810]

[178811]18510.

[178812]Leibold, E. A.; Munro, H. N.: Cytoplasmic protein binds in vitro

[178813]to a highly conserved sequence in the 5-prime untranslated region

[178814]of ferritin heavy- and light-subunit mRNAs. Proc. Nat. Acad. Sci. 85:

[178815]2171-2175, 1988.

[178816]

[178817]

[178818]18511.

[178819]Anderson, L. A.; Friedman, L.; Osborne-Lawrence, S.; Lynch, E.;

[178820]Weissenbach, J.; Bowcock, A.; King, M.-C.: High-density genetic map

[178821]of the BRCA1 region of chromosome 17q12-q21. Genomics 17: 618-623,

[178822]1993.

[178823]

[178824]

[178825]18512.

[178826]Fryns, J. P.; Kleczkowska, A.; Moerman, P.; Vandenberghe, K.:

[178827]Hereditary hydronephrosis and the short arm of chromosome 6. (Letter) Hum.

[178828]Genet. 91: 514–515, 1993.

[178829]

[178830]

[178831]18513.

[178832]Groenen, P. M. A.; Garcia, E.; Debeer, P.; Devriendt, K.; Fryns,

[178833]J. P.; Van de Ven, W. J. M.: Structure, sequence, and chromosome

[178834]19 localization of human USF2 and its rearrangement in a patient with

[178835]multicystic renal dysplasia. Genomics 38: 141–148, 1996.

[178836]

[178837]

[178838]18514.

[178839]Groenen, P. M. A.; Vanderlinden, G.; Devriendt, K.; Fryns, J.-P;

[178840]Van de Ven, W. J. M.: Rearrangement of the human CDC5L gene by a

[178841]t(6;19)(p21;q13.1) in a patient with multicystic renal dysplasia. Genomics 49:

[178842]218–229, 1998.

[178843]

[178844]

[178845]18515.

[178846]Bost, F.; Bourguignon, J.; Martin, J.–P.; Sesboue, R.; Thiberville,

[178847]L.; Diarra–Mehrpour, M.: Isolation and characterization of the human

[178848]inter–alpha–trypsin inhibitor heavy–chain H1 gene. Europ. J. Biochem. 218:

[178849]283–291, 1993.

[178850]

[178851]

[178852]18516.

[178853]Galizzi, J.–P.; Zuber, C. E.; Harada, N.; Gorman, D. M.; Kastelein,

[178854]R.; Banchereau, J.; Howard, M.; Miyajima, A.: Molecular cloning of

[178855]a cDNA encoding the human interleukin 4 receptor. Int. Immun. 2:

[178856]669–675, 1990.

[178857]

[178858]

[178859]18517.

[178860]Grimbacher, B.; Holland, S. M.; Puck, J. M.: The interleukin-4

[178861]receptor variant Q576R in hyper-IgE syndrome. (Letter) New Eng. J.

[178862]Med. 338: 1073–1074, 1998.

[178863]

[178864]

[178865]18518.

[178866]Idzerda, R. L.; March, C. J.; Mosley, B.; Lyman, S. D.; VandenBos,

[178867]T.; Gimpel, S. D.; Din, W. S.; Grabstein, K. H.; Widmer, M. B.; Park,

[178868]L. S.; Cosman, D.; Beckmann, M. P.: Human interleukin-4 receptor

[178869]confers biological responsiveness and defines a novel receptor superfamily. J.

[178870]Exp. Med. 171: 861-873, 1990.

[178871]

[178872]

[178873]18519.

[178874]Khurana Hershey, G. K.; Friedrich, M. F.; Esswein, L. A.; Thomas,

[178875]M. L.; Chatila, T. A.: The association of atopy with a gain-of-function

[178876]mutation in the alpha subunit of the interleukin-4 receptor. New

[178877]Eng. J. Med. 337: 1720-1725, 1997.

[178878]

[178879]

[178880]18520.

[178881]Kruse, S.; Japha, T.; Tedner, M.; Sparholt, S. H.; Forster, J.;

[178882]Kuehr, J.; Deichmann, K. A.; Abello, A.: The polymor-

phisms S503P

[178883]and Q576R in the interleukin-4 receptor alpha gene are associated

[178884]with atopy and influence the signal transduction. Immunology 96:

[178885]365-371, 1999.

[178886]

[178887]

[178888]18521.

[178889]Mitsuyasu, H.; Izuhara, K.; Mao, X.-Q.; Gao, P.-S.; Arinobu, Y.;

[178890]Enomoto, T.; Kawai, M.; Sasaki, S.; Dake, Y.; Hamasaki, N.; Shirakawa,

[178891]T.; Hopkin, J. M.: Ile50val variant of IL4R-alpha upregulates IgE

[178892]synthesis and associates with atopic asthma. (Letter) Nature Genet. 19:

[178893]119-120, 1998.

[178894]

[178895]

[178896]18522.

[178897]Olson, M.; Hood, L.; Cantor, C.; Botstein, D.: A common language

[178898]for physical mapping of the human genome. Science 245: 1434–1435,

[178899]1989.

[178900]

[178901]

[178902]18523.

[178903]Sutherland, G. R.; Baker, E.; Fernandez, K. E. W.; Callen, D. F.;

[178904]Goodwin, R. G.; Lupton, S.; Namen, A. E.; Shannon, M. F.; Vadas, M.

[178905]A.: The gene for human interleukin 7 (IL7) is at 8q12–13. Hum. Genet. 82:

[178906]371–372, 1989.

[178907]

[178908]

[178909]18524.

[178910]Watanabe, M.; Ueno, Y.; Yajima, T.; Iwao, Y.; Tsuchiya, M.; Ishikawa,

[178911]H.; Aiso, S.; Hibi, T.; Ishii, H.: Interleukin 7 is produced by human

[178912]intestinal epithelial cells and regulates the proliferation of intestinal

[178913]mucosal lymphocytes. J. Clin. Invest. 95: 2945–2953, 1995.

[178914]

[178915]

[178916]18525.

[178917]Corcoran, A. E.; Riddell, A.; Krooshoop, D.; Venkitaraman, A. R.

[178918]: Impaired immunoglobulin gene rearrangement in mice lacking the IL-7

[178919]receptor. Nature 391: 904–907, 1998.

[178920]

[178921]

[178922]18526.

[178923]Ober, C.; Leavitt, S. A.; Tsalenko, A.; Howard, T. D.; Hoki, D.

[178924]M.; Daniel, R.; Newman, D. L.; Wu, X.; Parry, R.; Lester, L. A.; Solway,

- [178925]J.; Blumenthal, M.; King, R. A.; Xu, J.; Meyers, D. A.;
Bleecker,
[178926]E. R.; Cox, N. J.: Variation in the interleukin 4–receptor al-
pha
[178927]gene confers susceptibility to asthma and atopy in ethni-
cally diverse
[178928]populations. Am. J. Hum. Genet. 66: 517–526, 2000.
- [178929]
- [178930]
- [178931]18527.
- [178932]Ober, C.; Tsalenko, A.; Willadsen, S.; Newman, D.; Daniel,
R.;
- [178933]Wu, X.; Andal, J.; Hoki, D.; Schneider, D.; True, K.; Schou,
C.; Parry,
[178934]R.; Cox, N.: Genome–wide screen for atopy susceptibility
alleles
[178935]in the Hutterites. Clin. Exp. Allergy 29 (suppl. 4): 11–15,
1999.
- [178936]
- [178937]
- [178938]18528.

[178939]Patuzzo, C.; Trabetti, E.; Malerba, G.; Martinati, L. C.;

Boner,

[178940]A. L.; Pescollderungg, L.; Zanoni, G.; Pignatti, P. F.: No
linkage

[178941]or association of the IL-4R-alpha gene Q576R mutation
with atopic

[178942]asthma in Italian families. (Letter) J. Med. Genet. 37:
382-384,

[178943]2000.

[178944]

[178945]

[178946]18529.

[178947]Pritchard, M. A.; Baker, E.; Whitmore, S. A.; Sutherland, G.
R.;

[178948]Idzerda, R. L.; Park, L. S.; Cosman, D.; Jenkins, N. A.;
Gilbert,

[178949]D. J.; Copeland, N. G.; Beckmann, M. P.: The interleukin-4
receptor

[178950]gene (IL4R) maps to 16p11.2-16p12.1 in human and to
the distal region

[178951]of mouse chromosome 7. Genomics 10: 801-806, 1991.

[178952]

[178953]

[178954]18530.

[178955]Suzuki, H.; Chung, F.; Palmer, E.; Sasaki, T.; Ohara, N.;
Taylor,

[178956]B. A.; Ohara, J.-I.: Gene mapping of mouse IL-4 receptor:
the loci

[178957]of interleukin 4 (IL-4) receptor gene and lymphocyte func-
tion associated

[178958]antigen 1 (LFA-1) gene are closely linked on chromosome
7. Immunogenetics 34:

[178959]252-256, 1991.

[178960]

[178961]

[178962]18531.

[178963]Zurawski, S. M.; Vega, F., Jr.; Huyghe, B.; Zurawski, G.: Re-
ceptors

[178964]for interleukin-13 and interleukin-4 are complex and
share a novel

[178965]component that functions in signal transduction. EMBO J.
12: 2663-2670,

[178966]1993.

[178967]

[178968]

[178969]18532.

[178970]Koshland, M. E.: The coming of age of the immunoglobulin J chain. Annu.

[178971]Rev. Immun. 3: 425–453, 1985.

[178972]

[178973]

[178974]18533.

[178975]Max, E. E.; McBride, O. W.; Morton, C. C.; Robinson, M. A.:
Human

[178976]J chain gene: chromosomal localization and associated restriction

[178977]fragment length polymorphisms. Proc. Nat. Acad. Sci. 83:
5592–5596,

[178978]1986.

[178979]

[178980]

[178981]18534.

[178982]Jones, C.; Penny, L.; Mattina, T.; Yu, S.; Baker, E.; Voullaire,

[178983]L.; Langdon, W. Y.; Sutherland, G. R.; Richards, R. I.; Tun-
nacliffe,

[178984]A.: Association of a chromosome deletion syndrome with
a fragile

[178985]site within the proto-oncogene CBL2. Nature 376:
145-149, 1995.

[178986]

[178987]

[178988]18535.

[178989]Diarra-Mehrpour, M.; Bourguignon, J.; Sesboue, R.; Mattei,
M.-G.;

[178990]Passage, E.; Salier, J.-P.; Martin, J.-P.: Human plasma in-
ter-alpha-trypsin

[178991]inhibitor is encoded by four genes on three chromosomes.
Europ. J.

[178992]Biochem. 179: 147-154, 1989.

[178993]

[178994]

[178995]18536.

[178996]Enghild, J. J.; Thogersen, I. B.; Pizzo, S. V.; Salvesen, G.:

[178997]Analysis of inter-alpha-trypsin inhibitor and a novel
trypsin inhibitor,

[178998]pre-alpha-trypsin inhibitor, from human plasma:
polypeptide chain

[178999]stoichiometry and assembly by glycan. J. Biol. Chem. 264:
15975-15981,

[179000]1989.

[179001]

[179002]

[179003]18537.

[179004]Salier, J. P.; Diarra-Mehrpour, M.; Sesboue, R.; Bour-
guignon, J.;

[179005]Benarous, R.; Ohkubo, I.; Kurachi, S.; Kurachi, K.; Martin, J.
P.

[179006]: Isolation and characterization of cDNAs encoding the
heavy chain

[179007]of human inter-alpha-trypsin inhibitor (I-alpha-TI): un-
ambiguous evidence

[179008]for multipolypeptide chain structure of I-alpha-TI. Proc.

Nat. Acad.

[179009]Sci. 84: 8272–8276, 1987.

[179010]

[179011]

[179012]18538.

[179013]Salier, J. P.; Simon, D.; Rouet, P.; Raguenez, G.; Muscatelli,

[179014]F.; Gebhard, W.; Guenet, J. L.; Mattei, M. G.: Homologous
chromosomal

[179015]locations of the four genes for inter-alpha-inhibitor and
pre-alpha-inhibitor

[179016]family in human and mouse: assignment of the ancestral
gene for the

[179017]lipocalin superfamily. Genomics 14: 83–88, 1992.

[179018]

[179019]

[179020]18539.

[179021]Bourguignon, J.; Diarra-Mehrpour, M.; Thiberville, L.; Bost,
F.;

[179022]Sesboue, R.; Martin, J.-P.: Human pre-alpha-trypsin in-
hibitor-precursor

[179023]heavy chain cDNA and deduced amino-acid sequence. Europ. J. Biochem. 212:

[179024]771–776, 1993.

[179025]

[179026]

[179027]18540.

[179028]Diarra-Mehrpour, M.; Sarafan, N.; Bourguignon, J.; Bonnet, F.;

[179029]Bost, F.; Martin, J.-P.: Human inter-alpha-trypsin inhibitor heavy

[179030]chain H3 gene: genomic organization, promoter analysis, and gene linkage. J.

[179031]Biol. Chem. 273: 26809–26819, 1998.

[179032]

[179033]

[179034]18541.

[179035]Brunton, L. L.; Lupton, S. D.: An STS in the human IL7 gene located

[179036]at 18q12–13. Nucleic Acids Res. 18: 1315 only, 1990.

[179037]

[179038]

[179039]18542.

[179040]Carvalho, T. L.; Mota-Santos, T.; Cumano, A.; Demengeot, J.; Vieira,

[179041]P.: Arrested B lymphopoiesis and persistence of activated B cells

[179042]in adult interleukin 7-/- mice. J. Exp. Med. 194: 1141-1150, 2001.

[179043]

[179044]

[179045]18543.

[179046]Goodwin, R. G.; Lupton, S.; Schmierer, A.; Hjerrild, K. J.; Jerzy,

[179047]R.; Clevenger, W.; Gillis, S.; Cosman, D.; Namen, A. E.: Human interleukin

[179048]7: molecular cloning and growth factor activity on human and murine

[179049]B-lineage cells. Proc. Nat. Acad. Sci. 86: 302-306, 1989.

[179050]

[179051]

[179052]18544.

[179053]Namen, A. E.; Lupton, S.; Hjerrild, K.; Wignall, J.;

Mochizuki,

[179054]D. Y.; Schmierer, A.; Mosley, B.; March, C. J.; Urdal, D.;

Gillis,

[179055]S.; Cosman, D.; Goodwin, R. G.: Stimulation of B-cell pro-
genitors

[179056]by cloned murine interleukin-7. Nature 333: 571-573,
1988.

[179057]

[179058]

[179059]18545.

[179060]Namen, A. E.; Schmierer, A. E.; March, C. J.; Overell, R. W.;

Park,

[179061]L. S.; Urdal, D. L.; Mochizuki, D. Y.: B-cell precursor
growth-promoting

[179062]activity: purification and characterization of a growth fac-
tor active

[179063]on lymphocyte precursors. J. Exp. Med. 167: 988-1002,
1988.

[179064]

[179065]

[179066]18546.

[179067]Goodwin, R. G.; Friend, D.; Ziegler, S. F.; Jerzy, R.; Falk, B.

[179068]A.; Gimpel, S.; Cosman, D.; Dower, S. K.; March, C. J.; Namen, A.

[179069]E.; Park, L. S.: Cloning of the human and murine interleukin-7 receptors:

[179070]demonstration of a soluble form and homology to a new receptor superfamily. Cell 60:

[179071]941-951, 1990.

[179072]

[179073]

[179074]18547.

[179075]Kondo, M.; Takeshita, T.; Higuchi, M.; Nakamura, M.; Sudo, T.;

[179076]Nishikawa, S.-I.; Sugamura, K.: Functional participation of the IL-2

[179077]receptor gamma chain in IL-7 receptor complexes. Science 263: 1453-1454,

[179078]1994.

[179079]

[179080]

[179081]18548.

[179082]Lai, S. Y.; Molden, J.; Goldsmith, M. A.: Shared gamma(c)
subunit

[179083]within the human interleukin-7 receptor complex: a
molecular basis

[179084]for the pathogenesis of X-linked severe combined im-
munodeficiency. J.

[179085]Clin. Invest. 99: 169-177, 1997.

[179086]

[179087]

[179088]18549.

[179089]Lynch, M.; Baker, E.; Park, L. S.; Sutherland, G. R.; Good-
win,

[179090]R. G.: The interleukin-7 receptor gene is at 5p13. Hum.
Genet. 89:

[179091]566-568, 1992.

[179092]

[179093]

[179094]18550.

[179095]Maki, K.; Sunaga, S.; Komagata, Y.; Kodaira, Y.; Mabuchi, A.; Karasuyama,

[179096]H.; Yokomuro, K.; Miyazaki, J.; Ikuta, K.: Interleukin 7 receptor-deficient

[179097]mice lack gamma/delta T cells. Proc. Nat. Acad. Sci. 93: 7172-7177,

[179098]1996.

[179099]

[179100]

[179101]18551.

[179102]Noguchi, M.; Nakamura, Y.; Russell, S. M.; Ziegler, S. F.; Tsang,

[179103]M.; Cao, X.; Leonard, W. J.: Interleukin-2 receptor gamma chain:

[179104]a functional component of the interleukin-7 receptor. Science 262:

[179105]1877-1880, 1993.

[179106]

[179107]

[179108]18552.

[179109]Peschon, J. J.; Morrissey, P. J.; Grabstein, K. H. et al.: Early

[179110]lymphocyte expansion is severely impaired in interleukin
7 receptor-deficient

[179111]mice. J. Exp. Med. 180: 1955–1960, 1994.

[179112]

[179113]

[179114]18553.

[179115]Puel, A.; Ziegler, S. F.; Buckley, R. H.; Leonard, W. J.: De-
fective

[179116]IL7R expression in T-B+NK+ severe combined immunod-
efficiency. Nature

[179117]Genet. 20: 394–397, 1998.

[179118]

[179119]

[179120]18554.

[179121]Mezei, M. M.; Mankodi, A.; Brais, B.; Marineau, C.; Thorn-
ton, C.

[179122]A.; Rouleau, G. A.; Karpati, G.: Minimal expansion of the

GCG repeat

[179123]in the PABP2 gene does not predispose to sporadic inclusion body myositis. Neurology 52:

[179124]669–670, 1999.

[179125]

[179126]

[179127]18555.

[179128]Aleman, A.; Verhaar, H. J. J.; De Haan, E. H. F.; De Vries, W.

[179129]R.; Samson, M. M.; Drent, M. L.; Van Der Veen, E. A.;

Koppeschaar,

[179130]H. P. F.: Insulin-like growth factor-I and cognitive function in

[179131]healthy older men. J. Clin. Endocr. Metab. 84: 471–475, 1999.

[179132]

[179133]

[179134]18556.

[179135]Baker, J.; Liu, J.-P.; Robertson, E. J.; Efstratiadis, A.: Role

[179136]of insulin-like growth factors in embryonic and postnatal growth. Cell 75:

[179137]73–82, 1993.

[179138]

[179139]

[179140]18557.

[179141]Bowcock, A.; Sartorelli, V.: Polymorphism and mapping of
the IGF1

[179142]gene, and absence of association with stature among
African Pygmies. Hum.

[179143]Genet. 85: 349–354, 1990.

[179144]

[179145]

[179146]18558.

[179147]Camacho–Hubner, C.; Woods, K. A.; Miraki–Moud, F.;
Hindmarsh, P.

[179148]C.; Clark, A. J.; Hansson, Y.; Johnston, A.; Baxter, R. C.;
Savage,

[179149]M. O.: Effects of recombinant human insulin–like growth
factor I

[179150](IGF–I) therapy on the growth hormone–IGF system of a
patient with

[179151]a partial IGF-I gene deletion. J. Clin. Endocr. Metab. 84:
1611–1616,
[179152]1999.

[179153]

[179154]

[179155]18559.

[179156]Goddard, A. D.; Covello, R.; Luoh, S.–M.; Clackson, T.; At-
tie,
[179157]K. M.; Gesundheit, N.; Rundle, A. C.; Wells, J. A.; Carlsson,
L. M.

[179158]S.: Mutations of the growth hormone receptor in children
with idiopathic

[179159]short stature. New Eng. J. Med. 333: 1093–1098, 1995.

[179160]

[179161]

[179162]18560.

[179163]Guler, H.–P.; Binz, K.; Eigenmann, E.; Jaggi, S.; Zimmer-
mann, D.;

[179164]Zapf, J.; Froesch, E. R.: Small stature and insulin-like
growth factors:

[179165]prolonged treatment of mini-poodles with recombinant human insulin-like

[179166]growth factor I. Acta Endocr. 121: 456–464, 1989.

[179167]

[179168]

[179169]18561.

[179170]Hirai, Y.; Takebe, K.; Takashina, M.; Kobayashi, S.; Take-
ichi,

[179171]M.: Epimorphin: a mesenchymal protein essential for ep-
ithelial morphogenesis. Cell 69:

[179172]471–481, 1992.

[179173]

[179174]

[179175]18562.

[179176]Zha, H.; Remmers, E. F.; Szpirer, C.; Szpirer, J.; Zhang, H.;
Kozak,

[179177]C. A.; Wilder, R. L.: The epimorphin gene is highly con-
served among

[179178]humans, mice, and rats and maps to human chromosome
7, mouse chromosome

[179179]5, and rat chromosome 12. Genomics 37: 386–389, 1996.

[179180]

[179181]

[179182]18563.

[179183]Fajans, S. S.: Maturity-onset diabetes of the young
(MODY). Diabetes

[179184]Metab. Rev. 5: 579–606, 1989.

[179185]

[179186]

[179187]18564.

[179188]Grosz, M. D.; Womack, J. E.; Skow, L. C.: Syntenic conser-
vation

[179189]of HSP70 genes in cattle and humans. Genomics 14:
863–868, 1992.

[179190]

[179191]

[179192]18565.

[179193]Harrison, G. S.; Drabkin, H. A.; Kao, F.-T.; Hartz, J.; Hart, I.

[179194]M.; Chu, E. H. Y.; Wu, B. J.; Morimoto, R. I.: Chromosomal

location

[179195]of human genes encoding major heat-shock protein
HSP70. Somat. Cell

[179196]Molec. Genet. 13: 119-130, 1987.

[179197]

[179198]

[179199]18566.

[179200]Harrison, G. S.; Morimoto, R.; Kao, F.-T.; Chu, E. H. Y.; Wu,
B.

[179201]J.; Drabkin, H.: Chromosomal location of human genes
encoding the

[179202]major heat shock protein HSP70. (Abstract) Am. J. Hum.
Genet. 39:

[179203]A157 only, 1986.

[179204]

[179205]

[179206]18567.

[179207]Hickey, E.; Brandon, S. E.; Sadis, S.; Smale, G.; Weber, L. A.

[179208]: Molecular cloning of sequences encoding the human
heat-shock proteins

[179209]and their expression during hyperthermia. Gene 43:
147–154, 1986.

[179210]

[179211]

[179212]18568.

[179213]Imai, Y.; Soda, M.; Hatakeyama, S.; Akagi, T.; Hashikawa,
T.; Nakayama,

[179214]K.; Takahashi, R.: CHIP is associated with Parkin, a gene
responsible

[179215]for familial Parkinson's disease, and enhances its ubiquitin
ligase

[179216]activity. Molec. Cell 10: 55–67, 2002.

[179217]

[179218]

[179219]18569.

[179220]Laroia, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.: Control

[179221]of mRNA decay by heat shock–ubiquitin–proteasome
pathway. Science 284:

[179222]499–502, 1999.

[179223]

[179224]

[179225]18570.

[179226]Leung, T. K. C.; Hall, C.; Rajendran, M.; Spurr, N. K.; Lim, L.

[179227]: The human heat-shock genes HSPA6 and HSPA7 are both expressed and

[179228]localize to chromosome 1. Genomics 12: 74–79, 1992.

[179229]

[179230]

[179231]18571.

[179232]Milner, C. M.; Campbell, R. D.: Polymorphic analysis of the three

[179233]MHC-linked HSP70 genes. Immunogenetics 36: 357–362, 1992.

[179234]

[179235]

[179236]18572.

[179237]Milner, C. M.; Campbell, R. D.: Structure and expression of the

[179238]three MHC-linked HSP70 genes. Immunogenetics 32:

242–251, 1990.

[179239]

[179240]

[179241]18573.

[179242]Pelham, H. R. B.: Speculations on the functions of the major

[179243]heat shock and glucose-regulated proteins. Cell 46:
959–961, 1986.

[179244]

[179245]

[179246]18574.

[179247]Sargent, C. A.; Dunham, I.; Trowsdale, J.; Campbell, R. D.:
Human

[179248]major histocompatibility complex contains genes for the
major heat

[179249]shock protein HSP70. Proc. Nat. Acad. Sci. 86:
1968–1972, 1989.

[179250]

[179251]

[179252]18575.

[179253]Slater, A.; Cato, A. C. B.; Sillar, G. M.; Kioussis, J.; Burdon,

[179254]R. H.: The pattern of protein synthesis induced by heat shock of

[179255]HeLa cells. *Europ. J. Biochem.* 117: 341–346, 1981.

[179256]

[179257]

[179258]18576.

[179259]Leung, T. K. C.; Rajendran, M. Y.; Monfries, C.; Hall, C.; Lim,

[179260]L.: The human heat-shock protein family: expression of a novel heat-inducible

[179261]HSP70 (HSP70B-prime) and isolation of its cDNA and genomic DNA. *Biochem.*

[179262]*J.* 267: 125–132, 1990.

[179263]

[179264]

[179265]18577.

[179266]Schiller, P.; Amin, J.; Ananthan, J.; Brown, M. E.; Scott, W. A.;

[179267]Voellmy, R.: Cis-acting elements involved in the regulated

expression

[179268]of a human HSP70 gene. J. Molec. Biol. 203: 97–105,
1988.

[179269]

[179270]

[179271]18578.

[179272]Voellmy, R.; Ahmed, A.; Schiller, P.; Bromley, P.; Rungger,
D.

[179273]: Isolation and functional analysis of a human
70,000–dalton heat

[179274]shock protein gene segment. Proc. Nat. Acad. Sci. 82:
4949–4953,

[179275]1985.

[179276]

[179277]

[179278]18579.

[179279]Rebbe, N. F.; Hickman, W. S.; Ley, T. J.; Stafford, D. W.;
Hickman,

[179280]S.: Nucleotide sequence and regulation of a human
90–kDa heat shock

[179281]protein gene. J. Biol. Chem. 264: 15006–15011, 1989.

[179282]

[179283]

[179284]18580.

[179285]Wolpowitz, D.; Mason, T. B. A.; Dietrich, P.; Mendelsohn, M.;

[179286]Talmage, D. A.; Role, L. W.: Cysteine-rich domain isoforms of the

[179287]neuregulin-1 gene are required for maintenance of peripheral synapses. Neuron 25:

[179288]79-91, 2000.

[179289]

[179290]

[179291]18581.

[179292]Bolton, P.; Powell, J.; Rutter, M.; Buckle, V.; Yates, J. R. W.;

[179293]Ishikawa-Brush, Y.; Monaco, A. P.: Autism, mental retardation, multiple

[179294]exostoses and short stature in a female with 46,X,t(X;8)(p22.13;q22.1). Psychiat.

[179295]Genet. 5: 51-55, 1995.

[179296]

[179297]

[179298]18582.

[179299]David, G.; van der Schueren, B.; Marynen, P.; Cassiman, J.-J.;

[179300]van den Berghe, H.: Molecular cloning of amphiglycan, a novel integral

[179301]membrane heparan sulfate proteoglycan expressed by epithelial and

[179302]fibroblastic cells. J. Cell Biol. 118: 961–969, 1992.

[179303]

[179304]

[179305]18583.

[179306]Ishikawa–Brush, Y.; Powell, J. F.; Bolton, P.; Miller, A. P.; Francis,

[179307]F.; Willard, H. F.; Lehrach, H.; Monaco, A. P.: Autism and multiple

[179308]exostoses associated with an X;8 translocation occurring within the

[179309]GRPR gene and 3–prime to the SDC2 gene. Hum. Molec. Genet. 6: 1241–1250,

[179310]1997.

[179311]

[179312]

[179313]18584.

[179314]Spring, J.; Goldberger, O. A.; Jenkins, N. A.; Gilbert, D. J.;

[179315]Copeland, N. G.; Bernfield, M.: Mapping of the syndecan
genes in

[179316]the mouse: linkage with members of the Myc gene family.

Genomics 21:

[179317]597–601, 1994.

[179318]

[179319]

[179320]18585.

[179321]Grenningloh, G.; Pribilla, I.; Prior, P.; Multhaup, G.;

Beyreuther,

[179322]K.; Taleb, O.; Betz, H.: Cloning and expression of the 58
kd beta

[179323]subunit of the inhibitory glycine receptor. Neuron 4:
963–970, 1990.

[179324]

[179325]

[179326]18586.

[179327]Handford, C. A.; Lynch, J. W.; Baker, E.; Webb, G. C.; Ford, J.

[179328]H.; Sutherland, G. R.; Schofield, P. R.: The human glycine receptor

[179329]beta subunit: primary structure, functional characterisation and chromosomal

[179330]localisation of the human and murine genes. Molec. Brain Res. 35:

[179331]211–219, 1996.

[179332]

[179333]

[179334]18587.

[179335]Milani, N.; Mulhardt, C.; Weber, R. G.; Lichter, P.; Kioschis,

[179336]P.; Poustka, A.; Becker, C.–M.: The human glycine receptor beta subunit

[179337]gene (GLRB): structure, refined chromosomal localization, and population

[179338]polymorphism. Genomics 50: 341–345, 1998.

[179339]

[179340]

[179341]18588.

[179342]Mulhardt, C.; Fischer, M.; Gass, P.; Simon-Chazottes, D.;

Guenet,

[179343]J.-L.; Kuhse, J.; Betz, H.; Becker, C.-M.: The spastic mouse:

aberrant

[179344]splicing of glycine receptor beta subunit mRNA caused by

intronic

[179345]insertion of L1 element. Neuron 13: 1003–1015, 1994.

[179346]

[179347]

[179348]18589.

[179349]Browner, M. F.; Nakano, K.; Bang, A. G.; Fletterick, R. J.:

Human

[179350]muscle glycogen synthase cDNA sequence: a negatively

charged protein

[179351]with an asymmetric charge distribution. Proc. Nat. Acad.

Sci. 86:

[179352]1443–1447, 1989.

[179353]

[179354]

[179355]18590.

[179356]Groop, L. C.; Kankuri, M.; Schalin-Jantti, C.; Ekstrand, A.;

Nikula-Ijas,

[179357]P.; Widen, E.; Kuusmanen, E.; Eriksson, J.; Franssila-

Kallunki, A.;

[179358]Saloranta, C.; Koskimies, S.: Association between polymorphism of

[179359]the glycogen synthase gene and non-insulin-dependent diabetes mellitus. New

[179360]Eng. J. Med. 328: 10-14, 1993.

[179361]

[179362]

[179363]18591.

[179364]Huang, X.; Vaag, A.; Hansson, M.; Weng, J.; Laurila, E.;
Groop,

[179365]L.: Impaired insulin-stimulated expression of the glycogen
synthase

[179366]gene in skeletal muscle of type 2 diabetic patients is ac-
quired rather

[179367]than inherited. J. Clin. Endocr. Metab. 85: 1584-1590,
2000.

[179368]

[179369]

[179370]18592.

[179371]Kadowaki, T.; Kadowaki, H.; Yazaki, Y.: Polymorphism of
the glycogen

[179372]synthase gene and non-insulin-dependent diabetes melli-
tus. (Letter) New

[179373]Eng. J. Med. 328: 1569, 1993.

[179374]

[179375]

[179376]18593.

[179377]Lehto, M.; Stoffel, M.; Groop, L.; Espinosa, R., III; Le Beau,

[179378]M. M.; Bell, G. I.: Assignment of the gene encoding glyco-
gen synthase

[179379](GYS) to human chromosome 19, band q13.3. Genomics
15: 460–461,

[179380]1993.

[179381]

[179382]

[179383]18594.

[179384]Seldin, M. F.; Mott, D.; Bhat, D.; Petro, A.; Kuhn, C. M.;
Kingsmore,

[179385]S. F.; Bogardus, C.; Opara, E.; Feinglos, M. N.; Surwit, R. S.:
Glycogen

[179386]synthase: a putative locus for diet–induced hyperglycemia.
J. Clin.

[179387]Invest. 94: 269–276, 1994.

[179388]

[179389]

[179390]18595.

[179391]Zouali, H.; Velho, G.; Froguel, P.: Polymorphism of the

glycogen

[179392]synthase gene and non-insulin-dependent diabetes mellitus. (Letter) New

[179393]Eng. J. Med. 328: 1568, 1993.

[179394]

[179395]

[179396]18596.

[179397]David, J. R.: Delayed hypersensitivity in vitro: its mediation

[179398]by cell-free substances formed by lymphoid cell-antigen interaction. Proc.

[179399]Nat. Acad. Sci. 56: 72-77, 1966.

[179400]

[179401]

[179402]18597.

[179403]Reche, P. A.; Soumelis, V.; Gorman, D. M.; Clifford, T.; Liu,

[179404]M.; Travis, M.; Zurawski, S. M.; Johnston, J.; Liu, Y.-J.;

Spits,

[179405]H.; de Waal Malefyt, R.; Kastelein, R. A.; Bazan, J. F.: Human thymic

[179406]stromal lymphopoietin preferentially stimulates myeloid cells. J.

[179407]Immun. 167: 336–343, 2001.

[179408]

[179409]

[179410]18598.

[179411]Kalmar, G. B.; Kay, R. J.; LaChance, A. C.; Cornell, R. B.:
Primary

[179412]structure and expression of a human CTP:phosphocholine
cytidylyltransferase. Biochim.

[179413]Biophys. Acta 1219: 328–334, 1994.

[179414]

[179415]

[179416]18599.

[179417]Rutherford, M. S.; Rock, C. O.; Jenkins, N. A.; Gilbert, D. J.;

[179418]Tessner, T. G.; Copeland, N. G.; Jackowski, S.: The gene
for murine

[179419]CTP:phosphocholine cytidylyltransferase (Ctpct) is located
on mouse

[179420]chromosome 16. Genomics 18: 698–701, 1993.

[179421]

[179422]

[179423]18600.

[179424]Tabas, J. A.; Zasloff, M.; Wasmuth, J. J.; Emanuel, B. S.; Al-
therr,

[179425]M. R.; McPherson, J. D.; Wozney, J. M.; Kaplan, F. S.: Bone
morphogenetic

[179426]protein: chromosomal localization of human genes for
BMP1, BMP2A,

[179427]and BMP3. Genomics 9: 283–289, 1991.

[179428]

[179429]

[179430]18601.

[179431]Gopal Rao, V. V. N.; Loffler, C.; Wozney, J. M.; Hansmann,
I.:

[179432]The gene for bone morphogenetic protein 2A (BMP2A) is
localized to

[179433]human chromosome 20p12 by radioactive and nonra-
dioactive in situ hybridization. Hum.

[179434]Genet. 90: 299–302, 1992.

[179435]

[179436]

[179437]18602.

[179438]Wang, E. A.; Rosen, V.; D'Alessandro, J. S.; Bauduy, M.;
Cordes,

[179439]P.; Harada, T.; Israel, D. I.; Hewick, R. M.; Kerns, K. M.; La-
Pan,

[179440]P.; Luxenberg, D. P.; McQuaid, D.; Moutsatsos, I. K.; Nove,
J.; Wozney,

[179441]J. M.: Recombinant human bone morphogenetic protein
induces bone

[179442]formation. Proc. Nat. Acad. Sci. 87: 2220–2224, 1990.

[179443]

[179444]

[179445]18603.

[179446]White, P. M.; Morrison, S. J.; Orimoto, K.; Kubu, C. J.;
Verdi,

[179447]J. M.; Anderson, D. J.: Neural crest stem cells undergo
cell–intrinsic

[179448]developmental changes in sensitivity to instructive differ-
entiation

[179449]signals. Neuron 29: 57–71, 2001.

[179450]

[179451]

[179452]18604.

[179453]Wozney, J. M.; Rosen, V.; Celeste, A. J.; Mitsock, L. M.;
Whitters,

[179454]M. J.; Kriz, R. W.; Hewick, R. M.; Wang, E. A.: Novel regula-
tors

[179455]of bone formation: molecular clones and activities. Sci-
ence 242:

[179456]1528–1534, 1988.

[179457]

[179458]

[179459]18605.

[179460]Connor, J. M.: Fibrodysplasia ossificans progressiva:
lessons

[179461]from rare maladies. (Editorial) New Eng. J. Med. 335:
591–593, 1996.

[179462]

[179463]

[179464]18606.

[179465]Dooley, C. A.; Attia, G. R.; Rainey, W. E.; Moore, D. R.;
Carr,

[179466]B. R.: Bone morphogenetic protein inhibits ovarian andro-
gen production. J.

[179467]Clin. Endocr. Metab. 85: 3331–3337, 2000.

[179468]

[179469]

[179470]18607.

[179471]McAlpine, P. J.: Personal Communication. Winnipeg, Mani-
toba,

[179472]Canada 7/15/1992.

[179473]

[179474]

[179475]18608.

[179476]Monsoro–Burq, A.–H.; le Douarin, N. M.: BMP4 plays a key
role

[179477]in left–right patterning in chick embryos by maintaining
Sonic Hedgehog

[179478]asymmetry. Molec. Cell 7: 789–799, 2001.

[179479]

[179480]

[179481]18609.

[179482]Shafritz, A. B.; Shore, E. M.; Gannon, F. H.; Zasloff, M. A.;
Taub,

[179483]R.; Muenke, M.; Kaplan, F. S.: Overexpression of an os-
teogenic morphogen

[179484]in fibrodysplasia ossificans progressiva. New Eng. J. Med.
335:

[179485]555–561, 1996.

[179486]

[179487]

[179488]18610.

[179489]Tabas, J. A.; Hahn, G. V.; Cohen, R. B.; Seaunez, H. N.;
Modi,

[179490]W. S.; Wozney, J. M.; Zasloff, M.; Kaplan, F. S.: Chromoso-
mal assignment

[179491]of the human gene for bone morphogenetic protein 4.
Clin. Orthop.

[179492]Rel. Res. 293: 310–316, 1993.

[179493]

[179494]

[179495]18611.

[179496]Tucker, A. S.; Matthews, K. L.; Sharpe, P. T.: Transforma-
tion

[179497]of tooth type induced by inhibition of BMP signaling. Sci-
ence 282:

[179498]1136–1138, 1998.

[179499]

[179500]

[179501]18612.

[179502]van den Wijngaard, A.; Olde Weghuis, D.; Boersma, C. J.
C.; van

[179503]Zoelen, E. J. J.; Geurts van Kessel, A.; Olijve, W.: Fine map-
ping

[179504]of the human bone morphogenetic protein–4 gene (BMP4)
to chromosome

[179505]14q22–q23 by in situ hybridization. Genomics 27:
559–560, 1995.

[179506]

[179507]

[179508]18613.

[179509]van den Wijngaard, A.; van Kraay, M.; van Zoelen, E. J. J.;

Olijve,

[179510]W.; Boersma, C. J. C.: Genomic organization of the human

bone morphogenetic

[179511]protein-4 gene: molecular basis for multiple transcripts.

Biochem.

[179512]Biophys. Res. Commun. 219: 789–794, 1996.

[179513]

[179514]

[179515]18614.

[179516]Amano, S.; Scott, I. C.; Takahara, K.; Koch, M.; Champli-

aud, M.-F.;

[179517]Gerecke, D. R.; Keene, D. R.; Hudson, D. L.; Nishiyama, T.;

Lee, S.;

[179518]Greenspan, D. S.; Burgeson, R. E.: Bone morphogenetic

protein 1 is

[179519]an extracellular processing enzyme of the laminin 5

gamma-2 chain. J.

[179520]Biol. Chem. 275: 22728–22735, 2000.

[179521]

[179522]

[179523]18615.

[179524]Bond, J. S.; Beynon, R. J.: The astacin family of metalloen-
dopeptidases. Protein

[179525]Sci. 4: 1247–1261, 1995.

[179526]

[179527]

[179528]18616.

[179529]Ceci, J. D.; Kingsley, D. M.; Silan, C. M.; Copeland, N. G.;
Jenkins,

[179530]N. A.: An interspecific backcross linkage map of the prox-
imal half

[179531]of mouse chromosome 14. Genomics 6: 673–678, 1990.

[179532]

[179533]

[179534]18617.

[179535]Kessler, E.; Takahara, K.; Biniaminov, L.; Brusel, M.;
Greenspan,

[179536]D.: Bone morphogenic protein-1: the type I procollagen
C-proteinase. Science 271:
[179537]360-362, 1996.

[179538]

[179539]

[179540]18618.

[179541]Chen, K.-C.; Vannais, D. B.; Jones, C.; Patterson, D.;
Davidson,

[179542]J. N.: Chromosomal localization of the human CAD gene
to 2p21-22.

[179543](Abstract) Am. J. Hum. Genet. 41: A161 only, 1987.

[179544]

[179545]

[179546]18619.

[179547]Chen, K.-C.; Vannais, D. B.; Jones, C.; Patterson, D.;
Davidson,

[179548]J. N.: Mapping of the gene encoding the multifunctional
protein carrying

[179549]out the first three steps of pyrimidine biosynthesis to hu-
man chromosome

[179550]2. Hum. Genet. 82: 40–44, 1989.

[179551]

[179552]

[179553]18620.

[179554]Chen, S.; Bigner, S. H.; Modrich, P.: High rate of CAD gene
amplification

[179555]in human cells deficient in MLH1 or MSH6. Proc. Nat.
Acad. Sci. 98:

[179556]13802–13807, 2001.

[179557]

[179558]

[179559]18621.

[179560]Graves, L. M.; Guy, H. I.; Kozlowski, P.; Huang, M.;
Lazarowski,

[179561]E.; Pope, R. M.; Collins, M. A.; Dahlstrand, E. N.; Earp, H.
S., III;

[179562]Evans, D. R.: Regulation of carbamoyl phosphate syn-
thetase by MAP

[179563]kinase. Nature 403: 328–332, 2000.

[179564]

[179565]

[179566]18622.

[179567]Muenke, M.; Sosnoski, D. M.; Wilson, W. G.; Wassman, E.
R.; Davidson,

[179568]J. N.; Patterson, D.; Nussbaum, R. L.: Exclusion of the CAD
locus

[179569]from 2p2101–p23.3 using 2p interstitial deletions from
patients with

[179570]holoprosencephaly. (Abstract) Am. J. Hum. Genet. 45
(suppl.): A84

[179571]only, 1989.

[179572]

[179573]

[179574]18623.

[179575]Simmer, J. P.; Kelly, R. E.; Rinker, A. G., Jr.; Zimmermann,
B.

[179576]H.; Scully, J. L.; Kim, H.; Evans, D. R.: Mammalian dihy-
droorotase:

[179577]nucleotide sequence, peptide sequences, and evolution of
the dihydroorotase

[179578]domain of the multifunctional protein CAD. Proc. Nat.

Acad. Sci. 87:

[179579]174–178, 1990.

[179580]

[179581]

[179582]18624.

[179583]Donalies, M.; Cramer, M.; Ringwald, M.; Starzinski-Powitz,
A.:

[179584]Expression of M-cadherin, a member of the cadherin
multigene family,

[179585]correlates with differentiation of skeletal muscle cells.

Proc. Nat.

[179586]Acad. Sci. 88: 8024–8028, 1991.

[179587]

[179588]

[179589]18625.

[179590]Kaupmann, K.; Becker-Follmann, J.; Scherer, G.; Jockusch,
H.; Starzinski-Powitz,

[179591]A.: The gene for the cell adhesion molecule M-cadherin
maps to mouse

[179592]chromosome 8 and human chromosome 16q24.1–qter

and is near the E-cadherin

[179593](uvomorulin) locus in both species. Genomics 14:
488–490, 1992.

[179594]

[179595]

[179596]18626.

[179597]Kremmidiotis, G.; Baker, E.; Crawford, J.; Eyre, H. J.; Nah-
mias,

[179598]J.; Callen, D. F.: Localization of human cadherin genes to
chromosome

[179599]regions exhibiting cancer-related loss of heterozygosity.
Genomics 49:

[179600]467–471, 1998.

[179601]

[179602]

[179603]18627.

[179604]Garcia-Castro, M. I.; Vielmetter, E.; Bronner-Fraser, M.: N-
cadherin,

[179605]a cell adhesion molecule involved in establishment of em-
bryonic left-right

[179606]asymmetry. Science 288: 1047–1051, 2000.

[179607]

[179608]

[179609]18628.

[179610]Hermiston, M. L.; Gordon, J. I.: Inflammatory bowel disease and

[179611]adenomas in mice expressing a dominant negative N-cadherin. Science 270:

[179612]1203–1206, 1995.

[179613]

[179614]

[179615]18629.

[179616]Miyatani, S.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.;

[179617]Takeichi, M.: Genomic structure and chromosomal mapping of the mouse

[179618]N-cadherin gene. Proc. Nat. Acad. Sci. 89: 8443–8447, 1992.

[179619]

[179620]

[179621]18630.

[179622]Takeichi, M.: Cadherins: a molecular family essential for selective

[179623]cell-cell adhesion and animal morphogenesis. Trends Genet. 3: 213-217,

[179624]1987.

[179625]

[179626]

[179627]18631.

[179628]Tanaka, H.; Shan, W.; Phillips, G. R.; Arndt, K.; Bozdagi, O.;

[179629]Shapiro, L.; Huntley, G. W.; Benson, D. L.; Colman, D. R.: Molecular

[179630]modification of N-cadherin in response to synaptic activity. Neuron 25:

[179631]93-107, 2000.

[179632]

[179633]

[179634]18632.

[179635]Wallis, J.; Fox, M. F.; Walsh, F. S.: Structure of the human N-cadherin

[179636]gene: YAC analysis and fine chromosomal mapping to

18q11.2. Genomics 22:

[179637]172–179, 1994.

[179638]

[179639]

[179640]18633.

[179641]Walsh, F. S.; Barton, C. H.; Putt, W.; Moore, S. E.; Kelsell, D.;

[179642]Spurr, N.; Goodfellow, P. N.: N-cadherin gene maps to human chromosome

[179643]18 and is not linked to the E-cadherin gene. J. Neurochem. 55: 805–812,

[179644]1990.

[179645]

[179646]

[179647]18634.

[179648]Claverie, J.–M.; Hardelin, J.–P.; Legouis, R.; Levilliers, J.;

[179649]Bougueleret, L.; Mattei, M.–G.; Petit, C.: Characterization and chromosomal

[179650]assignment of a human cDNA encoding a protein related to the murine

[179651]102-kDa cadherin-associated protein (alpha-catenin).

Genomics 15:

[179652]13-20, 1993.

[179653]

[179654]

[179655]18635.

[179656]Cook, S. A.; Bronson, R. T.; Donahue, L. R.; Ben-Arie, N.;
Davisson,

[179657]M. T.: Cerebellar deficient folia (cdf): a new mutation on
mouse

[179658]chromosome 6. Mammalian Genome 8: 108-112, 1997.

[179659]

[179660]

[179661]18636.

[179662]Herrenknecht, K.; Ozawa, M.; Eckerskorn, C.; Lottspeich,
F.; Lenter,

[179663]M.; Kemler, R.: The uvomorulin-anchorage protein alpha-
catenin is

[179664]a vinculin homologue. Proc. Nat. Acad. Sci. 88:
9156-9160, 1991.

[179665]

[179666]

[179667]18637.

[179668]Park, C.; Falls, W.; Finger, J. H.; Longo-Guess, C. M.; Ackerman,

[179669]S. L.: Deletion in Catna2, encoding alpha-N-catenin, causes cerebellar

[179670]and hippocampal lamination defects and impaired startle modulation. Nature

[179671]Genet. 31: 279–284, 2002.

[179672]

[179673]

[179674]18638.

[179675]Magnuson, V. L.; McCombs, J. L.; Lee, C.-C.; Yang, F.; Bowman,

[179676]B. H.; McGill, J. R.: Human alpha-2-HS-glycoprotein localized to

[179677]3q27–q29 by in situ hybridization. Cytogenet. Cell Genet. 47: 72–74,

[179678]1988.

[179679]

[179680]

[179681]18639.

[179682]Seroussi, E.; Kedra, D.; Kost-Alimova, M.; Sandberg-Nordqvist,

[179683]A.-C.; Fransson, I.; Jacobs, J. F. M.; Fu, Y.; Pan, H.-Q.; Roe, B.

[179684]A.; Imreh, S.; Dumanski, J. P.: TOM1 genes map to human chromosome

[179685]22q13.1 and mouse chromosome 8C1 and encode proteins similar to the

[179686]endosomal proteins HGS and STAM. Genomics 57: 380-388, 1999.

[179687]

[179688]

[179689]18640.

[179690]Hanson, I.; Churchill, A.; Love, J.; Axton, R.; Moore, T.; Clarke,

[179691]M.; Meire, F.; van Heyningen, V.: Missense mutations in the most

[179692]ancient residues of the PAX6 paired domain underlie a

spectrum of

[179693]human congenital eye malformations. Hum. Molec. Genet.

8: 165–172,

[179694]1999.

[179695]

[179696]

[179697]18641.

[179698]O'Farrell, P. H.: High resolution two-dimensional electrophoresis

[179699]of proteins. J. Biol. Chem. 250: 4007–4021, 1975.

[179700]

[179701]

[179702]18642.

[179703]Osawa, M.; Umetsu, K.; Ohki, T.; Nagasawa, T.; Suzuki, T.; Takeichi,

[179704]S.: Molecular evidence for human alpha2-HS glycoprotein (AHSG) polymorphism. Hum.

[179705]Genet. 99: 18–21, 1997.

[179706]

[179707]

[179708]18643.

[179709]Pedersen, K. O.: Fetuin, a new globulin isolated from
serum.

[179710](Letter) Nature 154: 575, 1944.

[179711]

[179712]

[179713]18644.

[179714]Rizzu, P.; Baldini, A.: Three members of the human cys-
tatin gene

[179715]superfamily, AHSG, HRG, and KNG, map within one
megabase of genomic

[179716]DNA at 3q27. Cytogenet. Cell Genet. 70: 26–28, 1995.

[179717]

[179718]

[179719]18645.

[179720]Terkelsen, O. B. F.; Jahnen–Dechent, W.; Nielsen, H.; Moos,
T.;

[179721]Fink, E.; Nawratil, P.; Muller–Esterl, W.; Mollgard, K.: Rat

fetuin:

[179722]distribution of protein and mRNA in embryonic and neonatal rat tissues. Anat.

[179723]Embryol. 197: 125–133, 1998.

[179724]

[179725]

[179726]18646.

[179727]Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha-2-HS-glycoprotein

[179728]typing by isoelectric focusing. Hum. Genet. 67: 70–71, 1984.

[179729]

[179730]

[179731]18647.

[179732]Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha-2-HS-glycoprotein

[179733]allele (AHS*5*) in two Japanese families. Hum. Genet. 68: 264–265,

[179734]1984.

[179735]

[179736]

[179737]18648.

[179738]Umetsu, K.; Yuasa, I.; Ikuta, M.; Suzuki, T.: Alpha-
2-HS-glycoprotein

[179739](A2HS) polymorphism in a Japanese population: existence
of two new

[179740]variants. Jpn. J. Hum. Genet. 32: 263-266, 1987.

[179741]

[179742]

[179743]18649.

[179744]Umetsu, K.; Yuasa, I.; Suzuki, T.: The polymorphism of
desialyzed

[179745]alpha-2-HS-glycoprotein (AHS): isoelectric focusing in
2.5M urea as

[179746]a method for identification of genetic variants. Hum.
Genet. 73:

[179747]372-373, 1986.

[179748]

[179749]

[179750]18650.

[179751]Yoshioka, Y.; Gejyo, F.; Marti, T.; Rickli, E. E.; Burgi, W.;

[179752]Offner, G. D.; Troxler, R. F.; Schmid, K.: The complete
amino acid

[179753]sequence of the A-chain of human plasma alpha-
2HS-glycoprotein. J.

[179754]Biol. Chem. 261: 1665-1676, 1986.

[179755]

[179756]

[179757]18651.

[179758]Zelinski, T.; Kaita, H.; Lewis, M.; Coghlan, G.; Craig, D.:
The

[179759]sequence of chromosome 3 loci AHSG:TF:CHE1. Hum.
Hered. 37: 1-6,

[179760]1987.

[179761]

[179762]

[179763]18652.

[179764]Elpeleg, O. N.; Shaag, A.; Glustein, J. Z.; Anikster, Y.;
Joseph,

[179765]A.; Saada, A.: Lipoamide dehydrogenase deficiency in

Ashkenazi Jews:

[179766]an insertion mutation in the mitochondrial leader sequence. Hum.

[179767]Mutat. 10: 256–257, 1997.

[179768]

[179769]

[179770]18653.

[179771]Feigenbaum, A. S.; Robinson, B. H.: The structure of the human

[179772]dihydrolipoamide dehydrogenase gene (DLD) and its upstream elements. Genomics 17:

[179773]376–381, 1993.

[179774]

[179775]

[179776]18654.

[179777]Hong, Y. S.; Kerr, D. S.; Craigen, W. J.; Tan, J.; Pan, Y.; Lusk,

[179778]M.; Patel, M. S.: Identification of two mutations in a compound heterozygous

[179779]child with dihydrolipoamide dehydrogenase deficiency.

Hum. Molec.

[179780]Genet. 5: 1925–1930, 1996.

[179781]

[179782]

[179783]18655.

[179784]Johnson, M.; Yang, H.–S.; Johanning, G. L.; Patel, M. S.:

Characterization

[179785]of the mouse dihydrolipoamide dehydrogenase (Dld)

gene: genomic structure,

[179786]promoter sequence, and chromosomal localization. Ge-

nomics 41: 320–326,

[179787]1997.

[179788]

[179789]

[179790]18656.

[179791]Liu, T.–C.; Kim, H.; Arizmendi, C.; Kitano, A.; Patel, M. S.:

[179792]Identification of two missense mutations in a dihy-

drolipoamide dehydrogenase-deficient

[179793]patient. Proc. Nat. Acad. Sci. 90: 5186–5190, 1993.

[179794]

[179795]

[179796]18657.

[179797]Matalon, R.; Michals, K.; Stumpf, D.; Goodman, S.; Parks, J.:

[179798]Lactic acidosis due to lipoamide dehydrogenase (LAD) deficiency: improvement

[179799]after oral lipoic acid. (Abstract) Am. J. Hum. Genet. 33: 48A, 1981.

[179800]

[179801]

[179802]18658.

[179803]Munnich, A.; Saudubray, J. M.; Taylor, J.; Charpentier, C.; Marsac,

[179804]C.; Rocchiccioli, F.; Amedee-Manesme, O.; Coude, F. X.; Frezal, J.;

[179805]Robinson, B. H.: Congenital lactic acidosis, alpha-ketoglutaric aciduria

[179806]and variant form of maple syrup urine disease due to a single enzyme

[179807]defect: dihydrolipoyl dehydrogenase deficiency. Acta Pae-

diat. Scand. 71:

[179808]167–171, 1982.

[179809]

[179810]

[179811]18659.

[179812]Olson, S.; Song, B. J.; Huh, T.–L.; Chi, Y.–T.; Veech, R. L.;

[179813]McBride, O. W.: Three genes for enzymes of the pyruvate
dehydrogenase

[179814]complex map to human chromosomes 3, 7, and X. Am. J.
Hum. Genet. 46:

[179815]340–349, 1990.

[179816]

[179817]

[179818]18660.

[179819]Otulakowski, G.; Robinson, B. H.: Isolation and sequence
determination

[179820]of cDNA clones for porcine and human lipoamide dehy-
drogenase. J.

[179821]Biol. Chem. 262: 17313–17318, 1987.

[179822]

[179823]

[179824]18661.

[179825]Otulakowski, G.; Robinson, B. H.; Willard, H. F.: Gene for
lipoamide

[179826]dehydrogenase maps to human chromosome 7. Somat.
Cell Molec. Genet. 14:

[179827]411–414, 1988.

[179828]

[179829]

[179830]18662.

[179831]Pons, G.; Raefsky–Estrin, C.; Carothers, D. J.; Pepin, R. A.;

[179832]Javed, A. A.; Jesse, B. W.; Ganapathi, M. K.; Samols, D.; Pa-
tel, M.

[179833]S.: Cloning and cDNA sequence of the dihydrolipoamide
dehydrogenase

[179834]component of human alpha–ketoacid dehydrogenase
complexes. Proc.

[179835]Nat. Acad. Sci. 85: 1422–1426, 1988.

[179836]

[179837]

[179838]18663.

[179839]Robinson, B. H.; Taylor, J.; Kahler, S. G.; Kirkman, H. N.:

Lactic

[179840]acidemia, neurologic deterioration and carbohydrate dependence in

[179841]a girl with dihydrolipoyl dehydrogenase deficiency. Europ. J. Pediat. 136:

[179842]35–39, 1981.

[179843]

[179844]

[179845]18664.

[179846]Robinson, B. H.; Taylor, J.; Sherwood, W. G.: Deficiency of dihydrolipoyl

[179847]dehydrogenase (a component of the pyruvate and alpha-ketoglutarate

[179848]dehydrogenase complexes): a cause of congenital lactic acidosis in

[179849]infancy. Pediat. Res. 11: 1198–1202, 1977.

[179850]

[179851]

[179852]18665.

[179853]Sakaguchi, Y.; Yoshino, M.; Aramaki, S.; Yoshida, I.; Yamashita,

[179854]F.; Kuhara, T.; Matsumoto, I.; Hayashi, T.: Dihydrolipoyl dehydrogenase

[179855]deficiency: a therapeutic trial with branched-chain amino acid restriction. Europ.

[179856]J. Pediat. 145: 271-274, 1986.

[179857]

[179858]

[179859]18666.

[179860]Scherer, S. W.; Otulakowski, G.; Robinson, B. H.; Tsui, L.-C.

[179861]: Localization of the human dihydrolipoamide dehydrogenase gene (DLD)

[179862]to 7q31-q32. Cytogenet. Cell Genet. 56: 176-177, 1991.

[179863]

[179864]

[179865]18667.

[179866]Shaag, A.; Saada, A.; Berger, I.; Mandel, H.; Joseph, A.;
Feigenbaum,

[179867]A.; Elpeleg, O. N.: Molecular basis of lipoamide dehydro-
genase deficiency

[179868]in Ashkenazi Jews. Am. J. Med. Genet. 82: 177–182, 1999.

[179869]

[179870]

[179871]18668.

[179872]Martin-Subero, J. I.; Gesk, S.; Harder, L.; Sonoki, T.;
Tucker,

[179873]P. W.; Schlegelberger, B.; Grote, W.; Novo, F. J.; Calasanz,
M. J.;

[179874]Hansmann, M. L.; Dyer, M. J. S.; Siebert, R.: Recurrent in-
volvement

[179875]of the REL and BCL11A loci in classical Hodgkin lym-
phoma. Blood 99:

[179876]1474–1477, 2002.

[179877]

[179878]

[179879]18669.

[179880]Hankinson, S. E.; Willett, W. C.; Colditz, G. A.; Hunter, D. J.;

[179881]Michaud, D. S.; Deroo, B.; Rosner, B.; Speizer, F. E.; Pollak, M.

[179882]: Circulating concentrations of insulin-like growth factor-I and risk

[179883]of breast cancer. Lancet 351: 1393–1396, 1998.

[179884]

[179885]

[179886]18670.

[179887]Jones, C.; Slijepcevic, P.; Marsh, S.; Baker, E.; Langdon, W. Y.;

[179888]Richards, R. I.; Tunncliffe, A.: Physical linkage of the fragile

[179889]site FRA11B and a Jacobsen syndrome chromosome deletion breakpoint

[179890]in 11q23.3. Hum. Molec. Genet. 3: 2123–2130, 1994.

[179891]

[179892]

[179893]18671.

[179894]Esumi, N.; Budarf, M.; Ciccarelli, L.; Sellinger, B.; Kozak, C.

[179895]A.; Wistow, G.: Conserved gene structure and genomic linkage for

[179896]D-dopachrome tautomerase (DDT) and MIF. Mammalian Genome 9: 753–757,

[179897]1998.

[179898]

[179899]

[179900]18672.

[179901]Kleemann, R.; Hausser, A.; Geiger, G.; Mischke, R.; Burger–Kentischer,

[179902]A.; Flieger, O.; Johannes, F.–J.; Roger, T.; Calandra, T.; Kapurniotu,

[179903]A.; Grell, M.; Finkelmeier, D.; Brunner, H.; Bernhagen, J.: Intracellular

[179904]action of the cytokine MIF to modulate AP–1 activity and the cell

[179905]cycle through Jab1. Nature 408: 211–216, 2000.

[179906]

[179907]

[179908]18673.

[179909]Kozak, C. A.; Adamson, M. C.; Buckler, C. E.; Segovia, L.;
Paralkar,

[179910]V.; Wistow, G.: Genomic cloning of mouse MIF
(macrophage inhibitory

[179911]factor) and genetic mapping of the human and mouse ex-
pressed gene

[179912]and nine mouse pseudogenes. Genomics 27: 405–411,
1995.

[179913]

[179914]

[179915]18674.

[179916]Paralkar, V.; Wistow, G.: Cloning the human gene for
macrophage

[179917]migration inhibitory factor (MIF). Genomics 19: 48–51,
1994.

[179918]

[179919]

[179920]18675.

[179921]Pastrana, D. V.; Raghavan, N.; Fitzgerald, P.; Eisinger, S.
W.;

[179922]Metz, C.; Bucala, R.; Schleimer, R. P.; Bickel, C.; Scott, A.
L.:

[179923]Filarial nematode parasites secrete a homologue of the
human cytokine

[179924]macrophage migration inhibitory factor. Infect. Immun.
66: 5955–5963,

[179925]1998.

[179926]

[179927]

[179928]18676.

[179929]Roger, T.; David, J.; Glauser, M. P.; Calandra, T.: MIF regu-
lates

[179930]innate immune response through modulation of Toll-like
receptor 4. Nature 414:

[179931]920–924, 2001.

[179932]

[179933]

[179934]18677.

[179935]Weiser, W. Y.; Temple, P. A.; Witek-Giannotti, J. S.; Re-
mold,

[179936]H. G.; Clark, S. C.; David, J. R.: Molecular cloning of a
cDNA encoding

[179937]a human macrophage migration inhibitory factor. Proc.
Nat. Acad.

[179938]Sci. 86: 7522–7526, 1989.

[179939]

[179940]

[179941]18678.

[179942]Emi, M.; Asaoka, H.; Matsumoto, A.; Itakura, H.; Kurihara,
Y.;

[179943]Wada, Y.; Kanamori, H.; Yazaki, Y.; Takahashi, E.; Lepert,
M.; Lalouel,

[179944]J.–M.; Kodama, T.; Mukai, T.: Structure, organization, and
chromosomal

[179945]mapping of the human macrophage scavenger receptor
gene. J. Biol.

[179946]Chem. 268: 2120–2125, 1993.

[179947]

[179948]

[179949]18679.

[179950]Latil, A.; Lidereau, R.: Genetic aspects of prostate cancer.

Virchows

[179951]Arch. 432: 389–406, 1998.

[179952]

[179953]

[179954]18680.

[179955]Matsumoto, A.; Naito, M.; Itakura, H.; Ikemoto, S.; Asaoka,
H.;

[179956]Hayakawa, I.; Kanamori, H.; Aburatani, H.; Takaku, F.;
Suzuki, H.;

[179957]Kobari, Y.; Miyai, T.; Takahashi, K.; Cohen, E. H.; Wydro,
R.; Housman,

[179958]D. E.; Kodama, T.: Human macrophage scavenger recep-
tors: primary

[179959]structure, expression, and localization in atherosclerotic
lesions. Proc.

[179960]Nat. Acad. Sci. 87: 9133–9137, 1990.

[179961]

[179962]

[179963]18681.

[179964]Xu, J.; Zheng, S. L.; Hawkins, G. A.; Faith, D. A.; Kelly, B.;

[179965]Isaacs, S. D.; Wiley, K. E.; Chang, B.; Ewing, C. M.; Bu-
jnovszky,

[179966]P.; Carpten, J. D.; Bleecker, E. R.; Walsh, P. C.; Trent, J. M.;
Meyers,

[179967]D. A.; Isaacs, W. B.: Linkage and association studies of
prostate

[179968]cancer susceptibility: evidence for linkage at 8p22–23.
Am. J. Hum.

[179969]Genet. 69: 341–350, 2001.

[179970]

[179971]

[179972]18682.

[179973]Xu, J.; Zheng, S. L.; Komiya, A.; Mychaleckyj, J. C.; Isaacs,
S.

[179974]D.; Hu, J. J.; Sterling, D.; Lange, E. M.; Hawkins, G. A.;
Turner,

[179975]A.; Ewing, C. M.; Faith, D. A.; and 19 others: Germline mu-
tations

[179976]and sequence variants of the macrophage scavenger re-
ceptor 1 gene

[179977]are associated with prostate cancer risk. Nature Genet. 32:
321–325,
[179978]2002.

[179979]

[179980]

[179981]18683.

[179982]Holness, C. L.; Simmons, D. L.: Molecular cloning of CD68,
a human
[179983]macrophage marker related to lysosomal glycoproteins.

Blood 81:
[179984]1607–1613, 1993.

[179985]

[179986]

[179987]18684.

[179988]Jones, E.; Quinn, C. M.; See, C. G.; Montgomery, D. S.;
Ford, M.

[179989]J.; Kolble, K.; Gordon, S.; Greaves, D. R.: The linked human
elongation

[179990]initiation factor 4A1 (EIF4A1) and CD68 genes map to
chromosome 17p13. Genomics 53:

[179991]248–250, 1998.

[179992]

[179993]

[179994]18685.

[179995]Chavrier, P.; Janssen–Timmen, U.; Mattei, M.–G.; Zerial, M.; Bravo,

[179996]R.; Charnay, P.: Structure, chromosome location, and expression of

[179997]the mouse zinc finger gene Krox–20: multiple gene products and coregulation

[179998]with the proto–oncogene c–fos. Molec. Cell. Biol. 9: 787–797, 1989.

[179999]

[180000]

[180001]18686.

[180002]Le Beau, M. M.; Espinosa, R., III; Neuman, W. L.; Stock, W.; Roulston,

[180003]D.; Larson, R. A.; Keinanen, M.; Westbrook, C. A.: Cytogenetic and

[180004]molecular delineation of the smallest commonly deleted region of chromosome

[180005]5 in malignant myeloid diseases. Proc. Nat. Acad. Sci. 90:
5484–5488,
[180006]1993.

[180007]

[180008]

[180009]18687.

[180010]Barrow, L. L.; Simin, K.; Jones, J. M.; Lee, D. C.; Meisler, M.

[180011]H.: Conserved linkage of early growth response 4, annexin
4, and

[180012]transforming growth factor alpha on mouse chromosome
6. Genomics 19:

[180013]388–390, 1994.

[180014]

[180015]

[180016]18688.

[180017]Crosby, S. D.; Veile, R. A.; Donis–Keller, H.; Baraban, J. M.;

[180018]Bhat, R. V.; Simburger, K. S.; Milbrandt, J.: Neural–specific
expression,

[180019]genomic structure, and chromosomal localization of the

gene encoding

[180020]the zinc-finger transcription factor NGFI-C. Proc. Nat. Acad. Sci. 89:

[180021]4739-4743, 1992.

[180022]

[180023]

[180024]18689.

[180025]Boerkoel, C. F.; Takashima, H.; Bacino, C. A.; Daentl, D.; Lupski,

[180026]J. R.: EGR2 mutation R359W causes a spectrum of Dejerine-Sottas neuropathy. Neurogenetics 3:

[180027]153-157, 2001.

[180028]

[180029]

[180030]18690.

[180031]Willcutt, E. G.; Pennington, B. F.; Smith, S. D.; Cardon, L. R.;

[180032]Gayan, J.; Knopik, V. S.; Olson, R. K.; DeFries, J. C.: Quantitative

[180033]trait locus for reading disability on chromosome 6p is

pleiotropic

[180034]for attention-deficit/hyperactivity disorder. Am. J. Med.

Genet. 114:

[180035]260–268, 2002.

[180036]

[180037]

[180038]18691.

[180039]Inazu, A.; Brown, M. L.; Hesler, C. B.; Agellon, L. B.;

Koizumi,

[180040]J.; Takata, K.; Maruhama, Y.; Mabuchi, H.; Tall, A. R.: In-

creased

[180041]high-density lipoprotein levels caused by a common

cholesteryl-ester

[180042]transfer protein gene mutation. New Eng. J. Med. 323:

1234–1238,

[180043]1990.

[180044]

[180045]

[180046]18692.

[180047]Saito, F.: A pedigree of homozygous familial hyperal-

phalipoproteinemia. Metabolism 33:

[180048]629–633, 1984.

[180049]

[180050]

[180051]18693.

[180052]Andres, D. A.; Milatovich, A.; Ozcelik, T.; Wenzlau, J. M.;
Brown,

[180053]M. S.; Goldstein, J. L.; Francke, U.: cDNA cloning of the two
subunits

[180054]of human CAAX farnesyltransferase and chromosomal
mapping of FNTA

[180055]and FNTB loci and related sequences. Genomics 18:
105–112, 1993.

[180056]

[180057]

[180058]18694.

[180059]Epstein, W. W.; Lever, D.; Leining, L. M.; Bruenger, E.;
Rilling,

[180060]H. C.: Quantitation of prenylcysteines by a selective cleav-
age reaction. Proc.

[180061]Nat. Acad. Sci. 88: 9668–9670, 1991.

[180062]

[180063]

[180064]18695.

[180065]Long, S. B.; Casey, P. J.; Beese, L. S.: Reaction path of protein
tein

[180066]farnesyltransferase at atomic resolution. Nature 419:
645–650, 2002.

[180067]

[180068]

[180069]18696.

[180070]Porter, J. C.; Messer, A.: Genetic mapping of farnesyltrans-
ferase

[180071]alpha(Fnta) to mouse chromosome 8. Mammalian Genome
7: 622–623,

[180072]1996.

[180073]

[180074]

[180075]18697.

[180076]Schafer, W. R.; Rine, J.: Protein prenylation: genes, en-

zymes,

[180077]targets, and functions. Annu. Rev. Genet. 26: 209–237,
1992.

[180078]

[180079]

[180080]18698.

[180081]Wang, T.; Danielson, P. D.; Li, B.; Shah, P. C.; Kim, S. D.;
Donahoe,

[180082]P. K.: The p21ras farnesyltransferase alpha subunit in
TGF-beta and

[180083]activin signaling. Science 271: 1120–1122, 1996.

[180084]

[180085]

[180086]18699.

[180087]Zhang, F. L.; Diehl, R. E.; Kohl, N. E.; Gibbs, J. B.; Giros, B.;

[180088]Casey, P. J.; Omer, C. A.: cDNA cloning and expression of
rat and

[180089]human protein geranylgeranyltransferase type-I. J. Biol.
Chem. 269:

[180090]3175–3180, 1994.

[180091]

[180092]

[180093]18700.

[180094]Roifman, C. M.; Zhang, J.; Chitayat, D.; Sharfe, N.: A partial

[180095]deficiency of interleukin-7R-alpha is sufficient to abrogate T-cell

[180096]development and cause severe combined immunodeficiency. Blood 96:

[180097]2803-2807, 2000.

[180098]

[180099]

[180100]18701.

[180101]Abraham, R.; Myers, A.; Wavrant-DeVrieze, F.; Hamshere, M. L.;

[180102]Thomas, H. V.; Marshall, H.; Compton, D.; Spurlock, G.; Turic, D.;

[180103]Hoogendoorn, B.; Kwon, J. M.; Petersen, R. C.; and 12 others: Substantial

[180104]linkage disequilibrium across the insulin-degrading enzyme locus but

[180105]no association with late-onset Alzheimer's disease. Hum.

Genet. 109:

[180106]646–652, 2001.

[180107]

[180108]

[180109]18702.

[180110]Affholter, J. A.; Fried, V. A.; Roth, R. A.: Human insulin-degrading

[180111]enzyme shares structural and functional homologies with E. coli protease

[180112]III. Science 242: 1415–1418, 1988.

[180113]

[180114]

[180115]18703.

[180116]Affholter, J. A.; Hsieh, C.-L.; Francke, U.; Roth, R. A.: Insulin-degrading

[180117]enzyme: stable expression of the human complementary DNA, characterization

[180118]of its protein product, and chromosomal mapping of the human and mouse

[180119]genes. Molec. Endocr. 4: 1125–1135, 1990.

[180120]

[180121]

[180122]18704.

[180123]Edbauer, D.; Willem, M.; Lammich, S.; Steiner, H.; Haass, C.:

[180124]Insulin-degrading enzyme rapidly removes the beta-amyloid precursor

[180125]protein intracellular domain (AICD). J. Biol. Chem. 277: 13389–13393,

[180126]2002.

[180127]

[180128]

[180129]18705.

[180130]Espinosa, R., III; Lemons, R. S.; Perlman, R. K.; Kuo, W.–L.; Rosner,

[180131]M. R.; Le Beau, M. M.: Localization of the gene encoding insulin-degrading

[180132]enzyme to human chromosome 10, bands q23–q25. Cytogenet. Cell Genet. 57:

[180133]184–186, 1991.

[180134]

[180135]

[180136]18706.

[180137]Qiu, W. Q.; Walsh, D. M.; Ye, Z.; Vekrellis, K.; Zhang, J.;
Podlisny,

[180138]M. B.; Rosner, M. R.; Safavi, A.; Hersh, L. B.; Selkoe, D. J.:
Insulin-degrading

[180139]enzyme regulates extracellular levels of amyloid beta-
protein by degradation. J.

[180140]Biol. Chem. 273: 32730–32738, 1998.

[180141]

[180142]

[180143]18707.

[180144]Qiu, W. Q.; Ye, Z.; Kholodenko, D.; Seubert, P.; Selkoe, D.
J.

[180145]: Degradation of amyloid beta-protein by a metallopro-
tease secreted

[180146]by microglia and other neural and non-neural cells. J. Biol.
Chem. 272:

[180147]6641–6646, 1997.

[180148]

[180149]

[180150]18708.

[180151]Glasgow, B. J.; Heinzmann, C.; Kojis, T.; Sparkes, R. S.;
Mohandas,

[180152]T.; Bateman, J. B.: Assignment of tear lipocalin gene to
human chromosome

[180153]9q34–9qter. Curr. Eye Res. 12: 1019–1023, 1993.

[180154]

[180155]

[180156]18709.

[180157]Herzog, H.; Baumgartner, M.; Holzfeind, P.; Redl, B.: Con-
firmation

[180158]of 9q34 as the chromosomal site of the human lipocalin
LCN1 gene. Cytogenet.

[180159]Cell Genet. 69: 108–109, 1995.

[180160]

[180161]

[180162]18710.

[180163]Holzfeind, P.; Merschak, P.; Rogatsch, H.; Culig, Z.; Fe-
ichtinger,

[180164]H.; Klocker, H.; Redl, B.: Expression of the gene for tear
lipocalin/von

[180165]Ebner's gland protein in human prostate. FEBS Lett. 395:
95-98,

[180166]1996.

[180167]

[180168]

[180169]18711.

[180170]Lacazette, E.; Pitiot, G.; Jobert, S.; Mallet, J.; Gachon, A. M.

[180171]F.: Fine genetic mapping of LCN1/D9S1826 within 9q34.
Ann. Hum.

[180172]Genet. 61: 449-455, 1997.

[180173]

[180174]

[180175]18712.

[180176]Lassagne, H.; Gachon, A. M. F.: Cloning of a human

lacrimal lipocalin

[180177]secreted in tears. Exp. Eye Res. 56: 605–609, 1993.

[180178]

[180179]

[180180]18713.

[180181]Lassagne, H.; Nguyen, V. C.; Mattei, M. G.; Gachon, A. M. F.:

[180182]Assignment of LCN1 to human chromosome 9 is confirmed. Cytogent.

[180183]Cell Genet. 71: 104, 1995.

[180184]

[180185]

[180186]18714.

[180187]Lassagne, H.; Ressot, C.; Mattei, M. G.; Gachon, A. M. F.:
Assignment

[180188]of the human tear lipocalin gene (LCN1) to 9q34 by in situ
hybridization. Genomics 18:

[180189]160–161, 1993.

[180190]

[180191]

[180192]18715.

[180193]Nguyen Van Cong; Weil, D.; Finaz, C.; Cohen-Haguenaer,
O.; Gross,

[180194]M.-S.; Jegou-Foubert, C.; de Tand, M.-F.; Cochet, C.; de
Grouchy,

[180195]J.; Frezal, J.: Panel of twenty-five independent man-rodent
hybrids

[180196]for human genetic marker mapping. Ann. Genet. 29:
20-26, 1986.

[180197]

[180198]

[180199]18716.

[180200]Pervaiz, S.; Brew, K.: Homology and structure-function
correlations

[180201]between alpha-1-acid glycoprotein and serum retinol-
binding protein

[180202]and its relatives. FASEB J. 1: 209-214, 1987.

[180203]

[180204]

[180205]18717.

[180206]Redl, B.; Holzfeind, P.; Lottspeich, F.: cDNA cloning and sequencing

[180207]reveals human tear prealbumin to be a member of the lipophilic-ligand

[180208]carrier protein superfamily. J. Biol.Chem. 28: 20282-20287, 1992.

[180209]

[180210]

[180211]18718.

[180212]Redl, B.; Wojnar, P.; Ellemunter, H.; Feichtinger, H.: Identification

[180213]of a lipocalin in mucosal glands of the human tracheo-bronchial tree

[180214]and its enhanced secretion in cystic fibrosis. Lab. Invest. 78:

[180215]1121-1129, 1998.

[180216]

[180217]

[180218]18719.

[180219]van't Hof, W.; Blankenvoorde, M. F. J.; Veerman, E. C. I.;
Nieuw

[180220]Amerongen, A. V.: The salivary lipocalin Von Ebner's Gland
protein

[180221]is a cysteine proteinase inhibitor. J. Biol. Chem. 272:
1837-1841,

[180222]1997.

[180223]

[180224]

[180225]18720.

[180226]Huebner, K.; Cannizzaro, L. A.; Croce, C. M.; Frey, A. Z.;
Wallner,

[180227]B. P.; Hecht, B. K.; Hecht, F.: Chromosome localization of
the human

[180228]genes for lipocortin I and the lipocortin II family.
(Abstract) Cytogenet.

[180229]Cell Genet. 46: 631 only, 1987.

[180230]

[180231]

[180232]18721.

[180233]Huebner, K.; Cannizzaro, L. A.; Frey, A. Z.; Hecht, B. K.;
Hecht,

[180234]F.; Croce, C. M.; Wallner, B. P.: Chromosomal localization
of the

[180235]human genes for lipocortin I and lipocortin II. Oncogene
Res. 2:

[180236]299–310, 1988.

[180237]

[180238]

[180239]18722.

[180240]Spano, F.; Raugei, G.; Palla, E.; Colella, C.; Melli, M.: Char-
acterization

[180241]of the human lipocortin–2–encoding multigene family: its
structure

[180242]suggests the existence of a short amino acid unit under-
going duplication. Gene 95:

[180243]243–251, 1990.

[180244]

[180245]

[180246]18723.

[180247]Takahashi, S.; Reddy, S. V.; Chirgwin, J. M.; Devlin, R.;
Haipok,

[180248]C.; Anderson, J.; Roodman, G. D.: Cloning and identifica-
tion of annexin

[180249]II as an autocrine/paracrine factor that increases osteo-
clast formation

[180250]and bone resorption. J. Biol. Chem. 269: 28696–28701,
1994.

[180251]

[180252]

[180253]18724.

[180254]Harman, S. M.; Metter, E. J.; Blackman, M. R.; Landis, P. K.;

[180255]Carter, H. B.: Serum levels of insulin-like growth factor I
(IGF-I),

[180256]IGF-II, IGF-binding protein-3, and prostate-specific anti-
gen as predictors

[180257]of clinical prostate cancer. J. Clin. Endocr. Metab. 85:
4258–4265,

[180258]2000.

[180259]

[180260]

[180261]18725.

[180262]Holly, J.: Insulin-like growth factor-I and new opportunities

[180263]for cancer prevention. Lancet 351: 1373-1375, 1998.

[180264]

[180265]

[180266]18726.

[180267]Hoppener, J. W. M.; de Pagter-Holthuizen, P.; Geurts van Kessel,

[180268]A. H. M.; Jansen, M.; Kittur, S. D.; Antonarakis, S. E.; Lips, C.

[180269]J. M.; Sussenbach, J. S.: The human gene encoding insulin-like growth

[180270]factor I is located on chromosome 12. Hum. Genet. 69: 157-160, 1985.

[180271]

[180272]

[180273]18727.

[180274]Le Bouc, Y.; Dreyer, D.; Jaeger, F.; Binoux, M.; Sondermeyer,

[180275]P.: Complete characterization of the human IGF-I nucleotide sequence

[180276]isolated from a newly constructed adult liver cDNA library.
FEBS

[180277]Lett. 196: 108–112, 1986.

[180278]

[180279]

[180280]18728.

[180281]Hellstrom, A.; Perruzzi, C.; Ju, M.; Engstrom, E.; Hard, A.-L.;

[180282]Liu, J.-L.; Albertsson-Wikland, K.; Carlsson, B.; Niklasson, A.; Sjodell,

[180283]L.; LeRoith, D.; Senger, D. R.; Smith, L. E. H.: Low IGF-I suppresses

[180284]VEGF-survival signaling in retinal endothelial cells: direct correlation

[180285]with clinical retinopathy of prematurity. Proc. Nat. Acad. Sci. 98:

[180286]5804–5808, 2001.

[180287]

[180288]

[180289]18729.

[180290]Lembo, G.; Rockman, H. A.; Hunter, J. J.; Steinmetz, H.;
Koch,

[180291]W. J.; Ma, L.; Printz, M. P.; Ross, J., Jr.; Chien, K. R.; Powell–
Braxton,

[180292]L.: Elevated blood pressure and enhanced myocardial con-
tractility

[180293]in mice with severe IGF–1 deficiency. J. Clin. Invest. 98:
2648–2655,

[180294]1996.

[180295]

[180296]

[180297]18730.

[180298]Li, C. H.; Yamashiro, D.; Gospodarowicz, D.; Kaplan, S. L.;
Van

[180299]Vliet, G.: Total synthesis of insulin–like growth factor I
(somatomedin

[180300]C). Proc. Nat. Acad. Sci. 80: 2216–2220, 1983.

[180301]

[180302]

[180303]18731.

[180304]Liu, J.-P.; Baker, J.; Perkins, A. S.; Robertson, E. J.; Efstra-
tiadis,

[180305]A.: Mice carrying null mutations of the genes encoding in-
sulin-like

[180306]growth factor I (Igf-1) and type 1 IGF receptor (Igf1r). Cell
75:

[180307]59-72, 1993.

[180308]

[180309]

[180310]18732.

[180311]Mathews, L. S.; Norstedt, G.; Palmiter, R. D.: Regulation of

[180312]insulin-like growth factor I gene expression by growth
hormone. Proc.

[180313]Nat. Acad. Sci. 83: 9343-9347, 1986.

[180314]

[180315]

[180316]18733.

[180317]Morton, C.; Rall, L.; Bell, G.; Shows, T.: Human insulin-like

[180318]growth factor-1 (IGF1) is encoded at 12q22-q24.1, and
insulin-like

[180319]growth factor-2 (IGF2) is at 11p15. (Abstract) Cytogenet.
Cell Genet. 40:

[180320]703 only, 1985.

[180321]

[180322]

[180323]18734.

[180324]Musaro, A.; McCullagh, K.; Paul, A.; Houghton, L.; Do-
browolny,

[180325]G.; Molinaro, M.; Barton, E. R.; Sweeney, H. L.; Rosenthal,
N.: Localized

[180326]Igf-1 transgene expression sustains hypertrophy and re-
generation in

[180327]senescent skeletal muscle. Nature Genet. 27: 195-200,
2001.

[180328]

[180329]

[180330]18735.

[180331]Musaro, A.; McCullagh, K. J. A.; Naya, F. J.; Olson, E. N.;
Rosenthal,

[180332]N.: IGF-1 induces skeletal myocyte hypertrophy through calcineurin

[180333]in association with GATA-2 and NF-ATc1. Nature 400: 581-585, 1999.

[180334]

[180335]

[180336]18736.

[180337]Playford, M. P.; Bicknell, D.; Bodmer, W. F.; Macaulay, V. M.

[180338]: Insulin-like growth factor 1 regulates the location, stability,

[180339]and transcriptional activity of beta-catenin. Proc. Nat. Acad. Sci. 97:

[180340]12103-12108, 2000.

[180341]

[180342]

[180343]18737.

[180344]Powell-Braxton, L.; Hollingshead, P.; Warburton, C.; Dowd, M.;

[180345]Pitts-Meek, S.; Dalton, D.; Gillett, N.; Stewart, T. A.: IGF-I is

[180346]required for normal embryonic growth in mice. Genes
Dev. 7: 2609–2617,
[180347]1993.

[180348]

[180349]

[180350]18738.

[180351]Rapp, R.; Deger, A.; Blum, W.; Koch, R.; Weber, U.: Charac-
terization

[180352]of the protein which binds insulin-like growth factor in
human serum. Europ.

[180353]J. Biochem. 172: 421–425, 1988.

[180354]

[180355]

[180356]18739.

[180357]Rotwein, P.: Two insulin-like growth factor I messenger
RNAs

[180358]are expressed in human liver. Proc. Nat. Acad. Sci. 83:
77–81, 1986.

[180359]

[180360]

[180361]18740.

[180362]Schoenle, E. J.; Zenobi, P. D.; Torresani, T.; Werder, E. A.;

[180363]Zachmann, M.; Froesch, E. R.: Recombinant human insulin-like growth

[180364]factor I (rhIGF I) reduces hyperglycaemia in patients with extreme

[180365]insulin resistance. Diabetologia 34: 675–679, 1991.

[180366]

[180367]

[180368]18741.

[180369]Semsarian, C.; Wu, M.-J.; Ju, Y.-K.; Marciniec, T.; Yeoh, T.;

[180370]Allen, D. G.; Harvey, R. P.; Graham, R. M.: Skeletal muscle hypertrophy

[180371]is mediated by a Ca^{2+} -dependent calcineurin signalling pathway. Nature 400:

[180372]576–581, 1999.

[180373]

[180374]

[180375]18742.

[180376]Svoboda, M. E.; Van Wyk, J. J.; Klapper, D. G.; Fellows, R.
E.;

[180377]Grissom, F. E.; Schleuter, R. J.: Purification of somatomedin-C from

[180378]human plasma: chemical and biological properties, partial
sequence

[180379]analysis, and relationship to other somatomedins. Bio-
chemistry 19:

[180380]790-797, 1980.

[180381]

[180382]

[180383]18743.

[180384]Taylor, B. A.; Grieco, D.: Localization of the gene encoding

[180385]insulin-like growth factor I on mouse chromosome 10.

Cytogenet. Cell

[180386]Genet. 56: 57–58, 1991.

[180387]

[180388]

[180389]18744.

[180390]Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T.

[180391]B.: Insulin-like growth factor genes: chromosome organization and

[180392]association with disease. (Abstract) Am. J. Hum. Genet.

36: 121S

[180393]only, 1984.

[180394]

[180395]

[180396]18745.

[180397]Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T.

[180398]B.: Localization of insulin-like growth factor genes to human chromosomes

[180399]11 and 12. Nature 310: 784–786, 1984.

[180400]

[180401]

[180402]18746.

[180403]Ullrich, A.; Berman, C. H.; Dull, T. J.; Gray, A.; Lee, J. M.

[180404]: Isolation of the human insulin-like growth factor I gene using a

[180405]single synthetic DNA probe. EMBO J. 3: 361–364, 1984.

[180406]

[180407]

[180408]18747.

[180409]Gough, N. M.; Rakar, S.; Harpur, A.; Wilks, A. F.: Localization

[180410]of genes for two members of the JAK family of protein tyrosine kinases

[180411]to murine chromosomes 4 and 19. Mammalian Genome 6: 247–248, 1995.

[180412]

[180413]

[180414]18748.

[180415]Ihle, J. N.: Cytokine receptor signalling. Nature 377:
591–594,

[180416]1995.

[180417]

[180418]

[180419]18749.

[180420]Pritchard, M. A.; Baker, E.; Callen, D. F.; Sutherland, G. R.;

[180421]Wilks, A. F.: Two members of the JAK family of protein ty-
rosine kinases

[180422]map to chromosomes 1p31.3 and 9p24. Mammalian
Genome 3: 36–38, 1992.

[180423]

[180424]

[180425]18750.

[180426]Campbell, G. S.; Argetsinger, L. S.; Ihle, J. N.; Kelly, P. A.;

[180427]Rillema, J. A.; Carter–Su, C.: Activation of JAK2 tyrosine ki-
nase

[180428]by prolactin receptors in Nb2 cells and mouse mammary gland explants. Proc.

[180429]Nat. Acad. Sci. 91: 5232–5236, 1994.

[180430]

[180431]

[180432]18751.

[180433]Huang, L. J.; Constantinescu, S. N.; Lodish, H. F.: The N-terminal

[180434]domain of Janus kinase 2 is required for Golgi processing and cell

[180435]surface expression of erythropoietin receptor. Molec. Cell 8: 1327–1338,

[180436]2001.

[180437]

[180438]

[180439]18752.

[180440]Neubauer, H.; Cumano, A.; Muller, M.; Wu, H.; Huffstadt, U.; Pfeffer,

[180441]K.: Jak2 deficiency defines an essential developmental checkpoint

[180442]in definitive hematopoiesis. Cell 93: 397–409, 1998.

[180443]

[180444]

[180445]18753.

[180446]Parganas, E.; Wang, D.; Stravopodis, D.; Topham, D. J.;

Marine,

[180447]J.–C.; Teglund, S.; Vanin, E. F.; Bodner, S.; Colamonici, O.

R.; van

[180448]Deursen, J. M.; Grosveld, G.; Ihle, J. N.: Jak2 is essential for

[180449]signaling through a variety of cytokine receptors. Cell 93:

385–395,

[180450]1998.

[180451]

[180452]

[180453]18754.

[180454]Saltzman, A.; Stone, M.; Franks, C.; Searfoss, G.; Munro,

R.; Jaye,

[180455]M.; Ivashchenko, Y.: Cloning and characterization of hu-

man Jak–2

[180456]kinase: high mRNA expression in immune cells and mus–

cle tissue. Biochem.

[180457]Biophys. Res. Commun. 246: 627–633, 1998.

[180458]

[180459]

[180460]18755.

[180461]Schwaller, J.; Parganas, E.; Wang, D.; Cain, D.; Aster, J. C.;

[180462]Williams, I. R.; Lee, C.-K.; Gerthner, R.; Kitamura, T.;

Frantsve,

[180463]J.; Anastasiadou, E.; Loh, M. L.; Levy, D. E.; Ihle, J. N.;

Gilliland,

[180464]D. G.: Stat5 is essential for the myelo- and lymphoprolif-
erative

[180465]disease induced by TEL/JAK2. Molec. Cell 6: 693–704,
2000.

[180466]

[180467]

[180468]18756.

[180469]Watling, D.; Guschin, D.; Muller, M.; Silvennoinen, O.; Wit-
thuhn,

[180470]B. A.; Quelle, F. W.; Rogers, N. C.; Schindler, C.; Stark, G.

R.;

[180471] Ihle, J. N.; Kerr, I. M.: Complementation by the protein tyrosine

[180472] kinase JAK2 of a mutant cell line defective in the interferon-gamma

[180473] signal transduction pathway. Nature 366: 166–170, 1993.

[180474]

[180475]

[180476] 18757.

[180477] Cihakova, D.; Trebusak, K.; Heino, M.; Fadeyev, V.; Tiulpakov,

[180478] A.; Battelino, T.; Tar, A.; Halasz, Z.; Blumel, P.; Tawfik, S.; Krohn,

[180479] K.; Lebl, J.; Peterson, P.; MEWPE–APECED Study Group: Novel AIRE

[180480] mutations and P450 cytochrome autoantibodies in Central and Eastern

[180481] European patients with APECED. Hum. Mutat. 18: 225–232, 2001.

[180482]

[180483]

[180484]18758.

[180485]Clemente, M. G.; Obermayer–Straub, P.; Meloni, A.; Strass–
burg,

[180486]C. P.; Arangino, V.; Tukey, R. H.; de Virgiliis, S.; Manns, M.
P.

[180487]: Cytochrome P450 1A2 is a hepatic autoantigen in au–
toimmune polyglandular

[180488]syndrome type 1. J. Clin. Endocr. Metab. 82: 1353–1361,
1997.

[180489]

[180490]

[180491]18759.

[180492]Craig, J. M.; Schiff, L. H.; Boone, J. E.: Chronic moniliasis

[180493]associated with Addison's disease. Am. J. Dis. Child. 89:
669–684,

[180494]1955.

[180495]

[180496]

[180497]18760.

[180498]Ekwall, O.; Hedstrand, H.; Haavik, J.; Perheentupa, J.; Bet–

terle,

[180499]C.; Gustafsson, J.; Husebye, E.; Rorsman, F.; Kampe, O.:

Pteridin-dependent

[180500]hydroxylases as autoantigens in autoimmune polyen-

docrine syndrome

[180501]type I. J. Clin. Endocr. Metab. 85: 2944-2950, 2000.

[180502]

[180503]

[180504]18761.

[180505]Cetani, F.; Barbesino, G.; Borsari, S.; Pardi, E.; Cianferotti,

[180506]L.; Pinchera, A.; Marcocci, C.: A novel mutation of the au-
toimmune

[180507]regulator gene in an Italian kindred with autoimmune
polyendocrinopathy-candidiasis-ectodermal

[180508]dystrophy, acting in a dominant fashion and strongly
cosegregating

[180509]with hypothyroid autoimmune thyroiditis. J. Clin. Endocr.
Metab. 86:

[180510]4747-4752, 2001.

[180511]

[180512]

[180513]18762.

[180514]Finnish–German APECED Consortium: An autoimmune disease, APECED,

[180515]caused by mutations in a novel gene featuring two PHD–type zinc–finger

[180516]domains. Nature Genet. 17: 399–403, 1997.

[180517]

[180518]

[180519]18763.

[180520]Foz, M.; Mirada, A.; Guardia, J.: Endocrine disorders in a family.

[180521](Letter) Lancet II: 269 only, 1970.

[180522]

[180523]

[180524]18764.

[180525]Gass, J. D. M.: The syndrome of keratoconjunctivitis, superficial

[180526]moniliasis, idiopathic hypoparathyroidism and Addison's disease. Am.

[180527]J. Ophthal. 54: 660–674, 1962.

[180528]

[180529]

[180530]18765.

[180531]Gylling, M.; Tuomi, T.; Bjorses, P.; Kontiainen, S.; Partanen,

[180532]J.; Christie, M. R.; Knip, M.; Perheentupa, J.; Miettinen, A.:
Beta-cell

[180533]autoantibodies, human leukocyte antigen II alleles, and
type 1 diabetes

[180534]in autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy. J.

[180535]Clin. Endocr. Metab. 85: 4434–4440, 2000.

[180536]

[180537]

[180538]18766.

[180539]Hedstrand, H.; Ekwall, O.; Haavik, J.; Landgren, E.; Betterle,

[180540]C.; Perheentupa, J.; Gustafsson, J.; Husebye, E.; Rorsman,
F.; Kampe,

[180541]O.: Identification of tyrosine hydroxylase as an autoantigen in autoimmune

[180542]polyendocrine syndrome type I. Biochem. Biophys. Res.

Commun. 267:

[180543]456–461, 2000.

[180544]

[180545]

[180546]18767.

[180547]Heino, M.; Peterson, P.; Kudoh, J.; Shimizu, N.; Antonarakis,

[180548]S. E.; Scott, H. S.; Krohn, K.: APECED mutations in the autoimmune

[180549]regulator (AIRE) gene. Hum. Mutat. 18: 205–211, 2001.

[180550]

[180551]

[180552]18768.

[180553]Heino, M.; Scott, H. S.; Chen, Q.; Peterson, P.; Maenpaa, U.;

[180554]Papasavvas, M.–P.; Mittaz, L.; Barras, C.; Rossier, C.; Chrousos,

[180555]G. P.; Stratakis, C. A.; Nagamine, K.; Kudoh, J.; Shimizu, N.; Maclaren,

[180556]N.; Antonarakis, S. E.; Krohn, K.: Mutation analyses of
North American

[180557]APS-1 patients. Hum. Mutat. 13: 69-74, 1999.

[180558]

[180559]

[180560]18769.

[180561]Hendrix, T. R.: Personal Communication. Baltimore, Md.
1/14/1985.

[180562]

[180563]

[180564]18770.

[180565]Hiekkala, H.: Idiopathic hypoparathyroidism, adrenal in-
sufficiency

[180566]and moniliasis in children. Ann. Paediat. Fenn. 10:
213-222, 1964.

[180567]

[180568]

[180569]18771.

[180570]Hung, W.; Migeon, C. J.; Parrott, R. H.: A possible autoim-

mune

[180571]basis for Addison's disease in three siblings, one with idiopathic

[180572]hypoparathyroidism, pernicious anemia and superficial moniliasis. New

[180573]Eng. J. Med. 269: 658–663, 1963.

[180574]

[180575]

[180576]18772.

[180577]Husebye, E. S.; Gebre-Medhin, G.; Tuomi, T.; Perheentupa, J.;

[180578]Landin-Olsson, M.; Gustafsson, J.; Rorsman, F.; Kampe, O.: Autoantibodies

[180579]against aromatic L-amino acid decarboxylase in autoimmune polyendocrine

[180580]syndrome type I. J. Clin. Endocr. Metab. 82: 147–150, 1997.

[180581]

[180582]

[180583]18773.

[180584]Ishii, T.; Suzuki, Y.; Ando, N.; Matsuo, N.; Ogata, T.: Novel

[180585]mutations of the autoimmune regulator gene in two siblings with autoimmune

[180586]polyendocrinopathy–candidiasis–ectodermal dystrophy. J. Clin. Endocr.

[180587]Metab. 85: 2922–2926, 2000.

[180588]

[180589]

[180590]18774.

[180591]Kenny, F. M.; Holliday, M. D.: Hypoparathyroidism, moniliasis,

[180592]Addison's and Hashimoto's disease. Hypercalcemia treated with intravenously

[180593]administered sodium sulfate. New Eng. J. Med. 271: 708–713, 1964.

[180594]

[180595]

[180596]18775.

[180597]Krohn, K.; Uibo, R.; Aavik, E.; Peterson, P.; Savilahti, K.:

[180598]Identification by molecular cloning of an autoantigen associated with

[180599]Addison's disease as steroid 17-alpha-hydroxylase.

Lancet 339: 770-773,

[180600]1992.

[180601]

[180602]

[180603]18776.

[180604]Kunin, A. S.; MacKay, B. R.; Burns, S. L.; Halberstam, M. J.:

[180605]The syndrome of hypoparathyroidism and adrenocortical
insufficiency,

[180606]a possible sequel of hepatitis: case report and review of
the literature. Am.

[180607]J. Med. 34: 856-866, 1963.

[180608]

[180609]

[180610]18777.

[180611]Louria, D. B.; Shannon, D. C.; Johnson, G.; Caroline, L.;
Okas,

[180612]A.; Taschdjian, C.: The susceptibility to moniliasis in chil-
dren

[180613]with endocrine hypofunction. Trans. Assoc. Am. Phys. 80:

236–249,

[180614]1967.

[180615]

[180616]

[180617]18778.

[180618]Maghnie, M.; Cosi, G.; Genovese, E.; Manca-Bitti, M. L.;
Cohen,

[180619]A.; Zecca, S.; Tinelli, C.; Gallucci, M.; Bernasconi, S.;
Boscherini,

[180620]B.; Severi, F.; Arico, M.: Central diabetes insipidus in chil-
dren

[180621]and young adults. New Eng. J. Med. 343: 998–1007, 2000.

[180622]

[180623]

[180624]18779.

[180625]Marieb, N. J.; Melby, J. C.; Lyall, S. S.: Isolated hypoaldos-
teronism

[180626]associated with idiopathic hypoparathyroidism. Arch. In-
tern. Med. 134:

[180627]424–429, 1974.

[180628]

[180629]

[180630]18780.

[180631]McKusick, V. A.: Personal Communication. Baltimore, Md.
1985.

[180632]

[180633]

[180634]18781.

[180635]Meloni, A.; Perniola, R.; Faa, V.; Corvaglia, E.; Cao, A.;
Rosatelli,

[180636]M. C.: Delineation of the molecular defects in the AIRE
gene in autoimmune

[180637]polyendocrinopathy–candidiasis–ectodermal dystrophy
patients from

[180638]Southern Italy. J. Clin. Endocr. Metab. 87: 841–846, 2002.

[180639]

[180640]

[180641]18782.

[180642]Nagamine, K.; Peterson, P.; Scott, H. S.; Kudoh, J.; Mi–

noshima,

[180643]S.; Heino, M.; Krohn, K. J. E.; Lalioti, M. D.; Mullis, P. E.;

Antonarakis,

[180644]S. E.; Kawasaki, K.; Asakawa, S.; Ito, F.; Shimizu, N.: Posi-

tional

[180645]cloning of the APECED gene. Nature Genet. 17: 393-398,

1997.

[180646]

[180647]

[180648]18783.

[180649]Neufeld, M.; Maclaren, N.; Blizzard, R.: Autoimmune

polyglandular

[180650]syndrome. Pediat. Ann. 9: 154-162, 1980.

[180651]

[180652]

[180653]18784.

[180654]Neufeld, M.; Maclaren, N. K.; Blizzard, R. M.: Two types of

autoimmune

[180655]Addison's disease associated with different polyglandular

autoimmune

[180656](PGA) syndromes. Medicine 60: 355-362, 1981.

[180657]

[180658]

[180659]18785.

[180660]Nithiyananthan, R.; Heward, J. M.; Allahabadia, A.; Barnett, A.

[180661]H.; Franklyn, J. A.; Gough, S. C. L.: A heterozygous deletion of

[180662]the autoimmune regulator (AIRE1) gene, autoimmune thyroid disease,

[180663]and type 1 diabetes: no evidence for association. J. Clin. Endocr.

[180664]Metab. 85: 1320–1322, 2000.

[180665]

[180666]

[180667]18786.

[180668]Okano, M.; Bell, D. W.; Haber, D. A.; Li, E.: DNA methyltransferases

[180669]Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian

[180670]development. Cell 99: 247–257, 1999.

[180671]

[180672]

[180673]18787.

[180674]Wijmenga, C.; van den Heuvel, L. P. W. J.; Strengman, E.;
Luyten,

[180675]J. A. F. M.; van der Burgt, I. J. A. M.; de Groot, R.; Smeets,
D.

[180676]F. C. M.; Draaisma, J. M. T.; van Dongen, J. J.; De Abreu, R.
A.;

[180677]Pearson, P. L.; Sandkuijl, L. A.; Weemaes, C. M. R.: Local-
ization

[180678]of the ICF syndrome to chromosome 20 by homozygosity
mapping. Am.

[180679]J. Hum. Genet. 63: 803–809, 1998.

[180680]

[180681]

[180682]18788.

[180683]Xu, G.–L.; Bestor, T. H.; Bourc'his, D.; Hsieh, C.–L.; Tom-
merup,

[180684]N.; Bugge, M.; Hulten, M.; Qu, X.; Russo, J. J.; Viegas-
Pequignot,

[180685]E.: Chromosome instability and immunodeficiency syndrome caused by

[180686]mutations in a DNA methyltransferase gene. Nature 402: 187–191,

[180687]1999.

[180688]

[180689]

[180690]18789.

[180691]Chuang, D. T.; Shih, V. E.: Disorders of branched chain amino

[180692]acid and keto acid metabolism.In: Scriver, C. R.; Beaudet, A. L.;

[180693]Sly, W. S.; Valle, D.: Metabolic and Molecular Bases of Inherited

[180694]Disease. New York: McGraw–Hill (pub.) (7th ed.): 1995. Pp. 1239–1277.

[180695]

[180696]

[180697]18790.

[180698]Smeitink, J.; van den Heuvel, L.: Human mitochondrial complex

[180699]I in health and disease. Am. J. Hum. Genet. 64:

1505–1510, 1999.

[180700]

[180701]

[180702]18791.

[180703]Pizarro, T. T.; Michie, M. H.; Bentz, M.; Woraratanadharm,
J.;

[180704]Smith, M. F., Jr.; Foley, E.; Moskaluk, C. A.; Bickston, S. J.;
Cominelli,

[180705]F.: IL–18, a novel immunoregulatory cytokine, is up–
regulated in

[180706]Crohn's disease: expression and localization in intestinal
mucosal

[180707]cells. J. Immun. 162: 6829–6835, 1999.

[180708]

[180709]

[180710]18792.

[180711]Bailey, W. J.; Slightom, J. L.; Goodman, M.: Rejection of the

[180712]'flying primate' hypothesis by phylogenetic evidence from
the epsilon–globin

[180713]gene. Science 256: 86–89, 1992.

[180714]

[180715]

[180716]18793.

[180717]Baralle, F. E.; Shoulders, C. C.; Proudfoot, N. J.: The primary
mary

[180718]structure of the human epsilon–globin gene. Cell 21:
621–626, 1980.

[180719]

[180720]

[180721]18794.

[180722]Fritsch, E. F.; Lawn, R. M.; Maniatis, T.: Molecular cloning
and

[180723]characterization of the human beta–like globin gene cluster. Cell 19:

[180724]959–972, 1980.

[180725]

[180726]

[180727]18795.

[180728]Gale, R. E.; Clegg, J. B.; Huehns, E. R.: Human embryonic haemoglobins

[180729]Gower 1 and Gower 2. Nature 280: 162–164, 1979.

[180730]

[180731]

[180732]18796.

[180733]Gilman, J. G.; Smithies, O.: Fetal hemoglobin variants in mice. Science 160:

[180734]885–886, 1968.

[180735]

[180736]

[180737]18797.

[180738]He, Z.; Russell, J. E.: A human embryonic hemoglobin inhibits

[180739]Hb S polymerization in vitro and restores a normal phenotype to mouse

[180740]models of sickle cell disease. Proc. Nat. Acad. Sci. 99: 10635–10640,

[180741]2002.

[180742]

[180743]

[180744]18798.

[180745]Huehns, E. R.; Dance, N.; Beaven, G. H.; Hecht, F.; Motulsky, A.

[180746]G.: Human embryonic hemoglobin. Nature 201: 1095–1097, 1964.

[180747]

[180748]

[180749]18799.

[180750]Huehns, E. R.; Flynn, F. V.; Butler, E. A.; Beaven, G. H.: Two
[180751]new hemoglobin variants in a very young human embryo.
Nature 189:

[180752]496–497, 1961.

[180753]

[180754]

[180755]18800.

[180756]Raich, N.; Papayannopoulou, T.; Stamatoyannopoulos, G.;
Enver,

[180757]T.: Demonstration of a human epsilon-globin gene si-

lencer with studies

[180758]in transgenic mice. Blood 79: 861–864, 1992.

[180759]

[180760]

[180761]18801.

[180762]Ramsay, M.; Thomson, J. A.; Jenkins, T.: A new epsilon
globin

[180763]HincII variant fragment length in a South African Negroid
family. J.

[180764]Med. Genet. 23: 145–150, 1986.

[180765]

[180766]

[180767]18802.

[180768]Shen, S.–H.; Smithies, O.: Human globin psi–beta–2 is not
a globin–related

[180769]sequence. Nucleic Acids Res. 10: 7809–7818, 1982.

[180770]

[180771]

[180772]18803.

[180773]Le Coniat, M.; Kinet, J. P.; Berger, R.: The human genes for the

[180774]alpha and gamma subunits of the mast cell receptor for immunoglobulin

[180775]E are located on human chromosome band 1q23. Immunogenetics 32:

[180776]183–186, 1990.

[180777]

[180778]

[180779]18804.

[180780]'t Hart, L. M.; Stolk, R. P.; Heine, R. J.; Grobbee, D. E.; van

[180781]der Does, F. E. E.; Maassen, J. A.: Association of the insulin-receptor

[180782]variant met-985 with hyperglycemia and non-insulin-dependent diabetes

[180783]mellitus in the Netherlands: a population-based study. Am. J. Hum.

[180784]Genet. 59: 1119–1125, 1996.

[180785]

[180786]

[180787]18805.

[180788]Accili, D.; Drago, J.; Lee, E. J.; Johnson, M. D.; Cool, M. H.;

[180789]Salvatore, P.; Asico, L. D.; Jose, P. A.; Taylor, S. I.; West-
phal,

[180790]H.: Early neonatal death in mice homozygous for a null al-
lele of

[180791]the insulin receptor gene. Nature Genet. 12: 106–109,
1996.

[180792]

[180793]

[180794]18806.

[180795]Accili, D.; Frapier, C.; Mosthaf, L.; McKeon, C.; Elbein, S.
C.;

[180796]Permutt, M. A.; Ramos, E.; Lander, E.; Ullrich, A.; Taylor, S.
I.

[180797]: A mutation in the insulin receptor gene that impairs
transport of

[180798]the receptor to the plasma membrane and causes insulin-
resistant diabetes. EMBO

[180799]J. 8: 2509–2517, 1989.

[180800]

[180801]

[180802]18807.

[180803]Al-Gazali, L. I.; Khalil, M.; Devadas, K.: A syndrome of insulin

[180804]resistance resembling leprechaunism in five sibs of consanguineous

[180805]parents. J. Med. Genet. 30: 470–475, 1993.

[180806]

[180807]

[180808]18808.

[180809]Bar, R. S.; Muggeo, M.; Roth, J.; Kahn, C. R.; Havrankova, J.;

[180810]Imperato-McGinley, J.: Insulin resistance, acanthosis nigricans,

[180811]and normal insulin receptors in a young woman: evidence for a postreceptor

[180812]defect. J. Clin. Endocr. Metab. 47: 620–625, 1978.

[180813]

[180814]

[180815]18809.

[180816]Barbetti, F.; Gejman, P. V.; Taylor, S. I.; Raben, N.; Cama, A.;

[180817]Bonora, E.; Pizzo, P.; Moghetti, P.; Muggeo, M.; Roth, J.:
Detection

[180818]of mutations in insulin receptor gene by denaturing gradient gel electrophoresis. Diabetes 41:

[180819]408–415, 1992.

[180820]

[180821]

[180822]18810.

[180823]Barnes, N. D.; Palumbo, P. J.; Hayles, A. B.; Folgar, H.: Insulin

[180824]resistance, skin changes, and virilization: a recessively inherited

[180825]syndrome possibly due to pineal gland dysfunction. Diabetologia 10:

[180826]285–289, 1974.

[180827]

[180828]

[180829]18811.

[180830]Belke, D. D.; Betuing, S.; Tuttle, M. J.; Gravelleau, C.;
Young,

[180831]M. E.; Pham, M.; Zhang, D.; Cooksey, R. C.; McClain, D. A.;
Litwin,

[180832]S. E.; Taegtmeyer, H.; Severson, D.; Kahn, C. R.; Abel, E.
D.: Insulin

[180833]signaling coordinately regulates cardiac size, metabolism,
and contractile

[180834]protein isoform expression. J. Clin. Invest. 109: 629–639,
2002.

[180835]

[180836]

[180837]18812.

[180838]Benecke, H.; Flier, J. S.; Moller, D. E.: Alternatively spliced

[180839]variants of the insulin receptor protein: expression in nor-
mal and

[180840]diabetic human tissues. J. Clin. Invest. 89: 2066–2070,
1992.

[180841]

[180842]

[180843]18813.

[180844]Bruning, J. C.; Gautam, D.; Burks, D. J.; Gillette, J.; Schubert,

[180845]M.; Orban, P. C.; Klein, R.; Krone, W.; Muller-Wieland, D.; Kahn,

[180846]C. R.: Role of brain insulin receptor in control of body weight and

[180847]reproduction. Science 289: 2122–2125, 2000.

[180848]

[180849]

[180850]18814.

[180851]Bruning, J. C.; Michael, M. D.; Winnay, J. N.; Hayashi, T.; Horsch,

[180852]D.; Accili, D.; Goodyear, L. J.; Kahn, C. R.: A muscle-specific insulin

[180853]receptor knockout exhibits features of the metabolic syndrome of NIDDM

[180854]without altering glucose tolerance. Molec. Cell 2: 559–569, 1998.

[180855]

[180856]

[180857]18815.

[180858]Cama, A.; de la Luz Sierra, M.; Ottini, L.; Kadowaki, T.;

Gorden,

[180859]P.; Imperato-McGinley, J.; Taylor, S. I.: A mutation in the
tyrosine

[180860]kinase domain of the insulin receptor associated with in-
sulin resistance

[180861]in an obese woman. J. Clin. Endocr. Metab. 73: 894-901,
1991.

[180862]

[180863]

[180864]18816.

[180865]Henry, I.; Humphries, S. E.; Tata, F.; Barichard, F.; Holm,
M.;

[180866]Williamson, R.; Junien, C.: The gene for HMG CoA reduc-
tase (HMGCR)

[180867]is on human chromosome 5. (Abstract) Cytogenet. Cell
Genet. 40:

[180868]649-650, 1985.

[180869]

[180870]

[180871]18817.

[180872]Humphries, S. E.; Tata, F.; Henry, I.; Barichard, F.; Holm, M.;

[180873]Junien, C.; Williamson, R.: The isolation, characterisation, and

[180874]chromosomal assignment of the gene for human 3-hydroxy-3-methylglutaryl

[180875]coenzyme A reductase, (HMG-CoA reductase). Hum. Genet. 71: 254-258,

[180876]1985.

[180877]

[180878]

[180879]18818.

[180880]Julier, C.; Delepine, M.; Keavney, B.; Terwilliger, J.; Davis,

[180881]S.; Weeks, D. E.; Bui, T.; Jeunemaitre, X.; Velho, G.; Froguel, P.;

[180882]Ratcliffe, P.; Corvol, P.; Soubrier, F.; Lathrop, G. M.: Genetic

[180883]susceptibility for human familial essential hypertension in a region

[180884]of homology with blood pressure linkage on rat chromosome 10. Hum.

[180885]Molec. Genet. 6: 2077–2085, 1997.

[180886]

[180887]

[180888]18819.

[180889]Rutherford, S.; Johnson, M. P.; Curtain, R. P.; Griffiths, L.

[180890]R.: Chromosome 17 and the inducible nitric oxide synthase gene in

[180891]human essential hypertension. Hum. Genet. 109: 408–415, 2001.

[180892]

[180893]

[180894]18820.

[180895]Aprelikova, O.; Pajusola, K.; Partanen, J.; Armstrong, E.; Alitalo,

[180896]R.; Bailey, S. K.; McMahon, J.; Wasmuth, J.; Huebner, K.; Alitalo,

[180897]K.: FLT4, a novel class III receptor tyrosine kinase in chromosome

[180898]5q33–qter. Cancer Res. 52: 746–748, 1992.

[180899]

[180900]

[180901]18821.

[180902]Dumont, D. J.; Jussila, L.; Taipale, J.; Lymboussaki, A.; Mu-
stonen,

[180903]T.; Pajusola, K.; Breitman, M.; Alitalo, K.: Cardiovascular
failure

[180904]in mouse embryos deficient in VEGF receptor-3. Science
282: 946-949,

[180905]1998.

[180906]

[180907]

[180908]18822.

[180909]Evans, A. L.; Brice, G.; Sotirova, V.; Mortimer, P.; Beninson,

[180910]J.; Burnand, K.; Rosbotham, J.; Child, A.; Sarfarazi, M.:
Mapping

[180911]of primary congenital lymphedema to the 5q35.3 region.
Am. J. Hum.

[180912]Genet. 64: 547-555, 1999.

[180913]

[180914]

[180915]18823.

[180916]Ferrell, R. E.; Levinson, K. L.; Esman, J. H.; Kimak, M. A.;
Lawrence,

[180917]E. C.; Barmada, M. M.; Finegold, D. N.: Hereditary lym-
phedema: evidence

[180918]for linkage and genetic heterogeneity. Hum. Molec. Genet.
7: 2073–2078,

[180919]1998.

[180920]

[180921]

[180922]18824.

[180923]Galland, F.; Karamysheva, A.; Mattei, M.–G.; Rosnet, O.;
Marchetto,

[180924]S.; Birnbaum, D.: Chromosomal localization of FLT4, a
novel receptor–type

[180925]tyrosine kinase gene. Genomics 13: 475–478, 1992.

[180926]

[180927]

[180928]18825.

[180929]Irrthum, A.; Karkkainen, M. J.; Devriendt, K.; Alitalo, K.;

Vikkula,

[180930]M.: Congenital hereditary lymphedema caused by a mutation that inactivates

[180931]VEGFR3 tyrosine kinase. Am. J. Hum. Genet. 67: 295–301, 2000.

[180932]

[180933]

[180934]18826.

[180935]Kaipainen, A.; Korhonen, J.; Mustonen, T.; van Hinsbergh, V. W.

[180936]M.; Fang, G.–H.; Dumont, D.; Breitman, M.; Alitalo, K.: Expression

[180937]of the fms–like tyrosine kinase 4 gene becomes restricted to lymphatic

[180938]endothelium during development. Proc. Nat. Acad. Sci. 92: 3566–3570,

[180939]1995.

[180940]

[180941]

[180942]18827.

[180943]Karkkainen, M. J.; Ferrell, R. E.; Lawrence, E. C.; Kimak, M.

A.;

[180944]Levinson, K. L.; McTigue, M. A.; Alitalo, K.; Finegold, D. N.:

Missense

[180945]mutations interfere with VEGFR-3 signalling in primary

lymphoedema. Nature

[180946]Genet. 25: 153-159, 2000.

[180947]

[180948]

[180949]18828.

[180950]Karkkainen, M. J.; Saaristo, A.; Jussila, L.; Karila, K. A.;

Lawrence,

[180951]E. C.; Pajusola, K.; Bueler, H.; Eichmann, A.; Kauppinen, R.;

Kettunen,

[180952]M. I.; Yla-Herttuala, S.; Finegold, D. N.; Ferrell, R. E.; Ali-

talo,

[180953]K.: A model for gene therapy of human hereditary lym-

phedema. Proc.

[180954]Nat. Acad. Sci. 98: 12677-12682, 2001.

[180955]

[180956]

[180957]18829.

[180958]Lee, J.; Gray, A.; Yuan, J.; Luoh, S.-M.; Avraham, H.; Wood, W.

[180959]I.: Vascular endothelial growth factor-related protein: a ligand

[180960]and specific activator of the tyrosine kinase receptor Flt4. Proc.

[180961]Nat. Acad. Sci. 93: 1988-1992, 1996.

[180962]

[180963]

[180964]18830.

[180965]Lyon, M. F.; Glenister, P. H.: New Mutants. Mouse Newsletter 71:

[180966]26 only, 1984.

[180967]

[180968]

[180969]18831.

[180970]Lyon, M. F.; Glenister, P. H.: Gene order of Chy-vt-Re on chromosome

[180971]11. Mouse Newsletter 74: 96 only, 1986.

[180972]

[180973]

[180974]18832.

[180975]Milroy, W. F.: An undescribed variety of hereditary
oedema. New

[180976]York Med. J. 56: 505–508, 1892.

[180977]

[180978]

[180979]18833.

[180980]Offori, T. W.; Platt, C. C.; Stephens, M.; Hopkinson, G. B.:

[180981]Angiosarcoma in congenital hereditary lymphoedema
(Milroy's disease):

[180982]diagnostic beacons and a review of the literature. Clin.
Exp. Derm. 18:

[180983]174–177, 1993.

[180984]

[180985]

[180986]18834.

[180987]Pajusola, K.; Aprelikova, O.; Korhonen, J.; Kaipainen, A.;
Pertovaara,

[180988]L.; Alitalo, R.; Alitalo, K.: FLT4 receptor tyrosine kinase contains

[180989]seven immunoglobulin-like loops and is expressed in multiple human

[180990]tissues and cell lines. Cancer Res. 52: 5738–5743, 1992.

[180991]

[180992]

[180993]18835.

[180994]Pajusola, K.; Aprelikova, O.; Pelicci, G.; Weich, H.; Claesson-Welsh,

[180995]L.; Alitalo, K.: Signalling properties of FLT4, a proteolytically

[180996]processed receptor tyrosine kinase related to two VEGF receptors. Oncogene 9:

[180997]3545–3555, 1994.

[180998]

[180999]

[181000]18836.

[181001]Walter, J. W.; North, P. E.; Waner, M.; Mizeracki, A.; Blei, F.;

[181002]Walker, J. W. T.; Reinisch, J. F.; Marchuk, D. A.: Somatic

mutation

[181003]of vascular endothelial growth factor receptors in juvenile hemangioma. Genes

[181004]Chromosomes Cancer 33: 295–303, 2002.

[181005]

[181006]

[181007]18837.

[181008]Ragoussis, J.; Senger, G.; Trowsdale, J.; Campbell, I. G.:
Genomic

[181009]organization of the human folate receptor genes on chromosome 11q13. Genomics 14:

[181010]423–430, 1992.

[181011]

[181012]

[181013]18838.

[181014]Ratnam, M.; Marquardt, H.; Duhring, J. L.; Freisheim, J. H.:
Homologous

[181015]membrane folate binding proteins in human placenta:
cloning and sequence

[181016]of a cDNA. Biochemistry 28: 8249–8254, 1989.

[181017]

[181018]

[181019]18839.

[181020]Barber, R. C.; Shaw, G. M.; Lammer, E. J.; Greer, K. A.;
Biela,

[181021]T. A.; Lacey, S. W.; Wasserman, C. R.; Finnell, R. H.: Lack of
association

[181022]between mutations in the folate receptor-alpha gene and
spina bifida. Am.

[181023]J. Med. Genet. 76: 310-317, 1998.

[181024]

[181025]

[181026]18840.

[181027]Bowcock, A. M.; Lacey, S.; Saltman, D.; Mohandas, T. K.;
Kamen,

[181028]B. A.; Taggart, R. T.: Localization of the folate receptor
gene to

[181029]chromosome 11q13. (Abstract) Cytogenet. Cell Genet. 58:
1955 only,

[181030]1991.

[181031]

[181032]

[181033]18841.

[181034]Campbell, I. G.; Jones, T. A.; Foulkes, W. D.; Trowsdale, J.:

[181035]Folate-binding protein is a marker for ovarian cancer.

Cancer Res. 51:

[181036]5329–5338, 1991.

[181037]

[181038]

[181039]18842.

[181040]Kranes, A.; Balogh, K., Jr.: Liver disease in a patient with

[181041]von Hippel–Lindau disease. New Eng. J. Med. 275:

950–959, 1966.

[181042]

[181043]

[181044]18843.

[181045]Lamiell, J. M.: Personal Communication. San Francisco,

Calif.

[181046]9/1987.

[181047]

[181048]

[181049]18844.

[181050]Lamiell, J. M.; Salazar, F. G.; Hsia, Y. E.: Von Hippel–Lindau

[181051]disease affecting 43 members of a single kindred.

Medicine 68: 1–29,

[181052]1989.

[181053]

[181054]

[181055]18845.

[181056]Latif, F.; Tory, K.; Gnarr, J.; Yao, M.; Duh, F.–M.; Orcutt,

[181057]M. L.; Stackhouse, T.; Kuzmin, I.; Modi, W.; Geil, L.;

Schmidt, L.;

[181058]Zhou, F.; Li, H.; Wei, M. H.; Chen, F.; Glenn, G.; Choyke, P.;

Walther,

[181059]M. M.; Weng, Y.; Duan, D.–S. R.; Dean, M.; Glavac, D.;

Richards, F.

[181060]M.; Crossey, P. A.; Ferguson–Smith, M. A.; Le Paslier, D.;

Chumakov,

[181061]I.; Cohen, D.; Chinault, A. C.; Maher, E. R.; Linehan, W. M.;

Zbar,

[181062]B.; Lerman, M. I.: Identification of the von Hippel–Lindau disease

[181063]tumor suppressor gene. Science 260: 1317–1320, 1993.

[181064]

[181065]

[181066]18846.

[181067]Lee, S.; Chen, D. Y. T.; Humphrey, J. S.; Gnarr, J. R.; Linehan,

[181068]W. M.; Klausner, R. D.: Nuclear/cytoplasmic localization of the von

[181069]Hippel–Lindau tumor suppressor gene product is determined by cell

[181070]density. Proc. Nat. Acad. Sci. 93: 1770–1775, 1996.

[181071]

[181072]

[181073]18847.

[181074]Lenz, T.; Thiede, H. M.; Nussberger, J.; Atlas, S. A.; Distler,

[181075]A.; Schulte, K. L.: Hyperreninemia and secondary hyperaldosteronism

[181076]in a patient with pheochromocytoma and von Hippel–Lindau disease. Nephron 62:

[181077]345–350, 1992.

[181078]

[181079]

[181080]18848.

[181081]Lindau, A.: Zur Frage der Angiomatosis Retinae und Ihrer Hirncomplication. Acta

[181082]Ophthal. 4: 193–226, 1927.

[181083]

[181084]

[181085]18849.

[181086]Loeb, D. B.; Pericak–Vance, M. A.; Stajich, J. M.; Vance, J. M.

[181087]: A novel mutation in the von Hippel–Lindau gene. Hum. Molec. Genet. 3:

[181088]1423–1424, 1994.

[181089]

[181090]

[181091]18850.

[181092]Lui, W. O.; Chen, J.; Glasker, S.; Bender, B. U.; Madura, C.;

[181093]Khoo, S. K.; Kort, E.; Larsson, C.; Neumann, H. P. H.; Teh,
B. T.

[181094]: Selective loss of chromosome 11 in pheochromocytomas
associated

[181095]with the VHL syndrome. *Oncogene* 21: 1117–1122, 2002.

[181096]

[181097]

[181098]18851.

[181099]Maddock, I. R.; Moran, A.; Maher, E. R.; Teare, M. D.; Nor-
man,

[181100]A.; Payne, S. J.; Whitehouse, R.; Dodd, C.; Lavin, M.; Hart-
ley, N.;

[181101]Super, M.; Evans, D. G. R.: A genetic register for von Hip-
pel–Lindau

[181102]disease. *J. Med. Genet.* 33: 120–127, 1996.

[181103]

[181104]

[181105]18852.

[181106]Maher, E. R.; Bentley, E.; Yates, J. R. W.; Barton, D.; Jennings,

[181107]A.; Fellows, I. W.; Ponder, M. A.; Ponder, B. A. J.; Benjamin, C.;

[181108]Harris, R.; Ferguson-Smith, M. A.: Mapping of von Hippel-Lindau disease

[181109]to chromosome 3p confirmed by genetic linkage analysis. J. Neurol.

[181110]Sci. 100: 27-30, 1990.

[181111]

[181112]

[181113]18853.

[181114]Maher, E. R.; Bentley, E.; Yates, J. R. W.; Latif, F.; Lerman,

[181115]M.; Zbar, B.; Affara, N. A.; Ferguson-Smith, M. A.: Mapping of the

[181116]von Hippel-Lindau disease locus to a small region of chromosome 3p

[181117]by genetic linkage analysis. Genomics 10: 957-960, 1991.

[181118]

[181119]

[181120]18854.

[181121]Maher, E. R.; Iselius, L.; Yates, J. R. W.; Littler, M.; Benjamin,

[181122]C.; Harris, R.; Sampson, J.; Williams, A.; Ferguson-Smith, M. A.;

[181123]Morton, N.: Von Hippel-Lindau disease: a genetic study. J. Med.

[181124]Genet. 28: 443-447, 1991.

[181125]

[181126]

[181127]18855.

[181128]Maher, E. R.; Yates, J. R. W.; Ferguson-Smith, M. A.: Statistical

[181129]analysis of the two stage mutation model in von Hippel-Lindau disease,

[181130]and in sporadic cerebellar haemangioblastoma and renal cell carcinoma. J.

[181131]Med. Genet. 27: 311-314, 1990.

[181132]

[181133]

[181134]18856.

[181135]Mahon, P. C.; Hirota, K.; Semenza, G. L.: FIH-1: a novel protein

[181136]that interacts with HIF-1-alpha and VHL to mediate repression of HIF-1

[181137]transcriptional activity. Genes Dev. 15: 2675-2686, 2001.

[181138]

[181139]100. Manski, T. J.; Heffner, D. K.; Glenn, G. M.; Patronas, N. J.;

[181140]Pikus, A. T.; Katz, D.; Lebovics, R.; Sledjeski, K.; Choyke, P. L.;

[181141]Zbar, B.; Linehan, W. M.; Oldfield, E. H.: Endolymphatic sac tumors:

[181142]a source of morbid hearing loss in von Hippel-Lindau disease. J.A.M.A. 277:

[181143]1461-1466, 1997.

[181144]

[181145]101. Maxwell, P. H.; Wiesener, M. S.; Chang, G.-W.; Clifford, S. C.;

[181146]Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.; Maher, E.

[181147]R.; Ratcliffe, P. J.: The tumour suppressor protein VHL targets hypoxia-inducible

[181148]factors for oxygen-dependent proteolysis. Nature 399: 271-275, 1999.

[181149]

[181150]102. McCabe, C. M.; Flynn, H. W., Jr.; Shields, C. L.; Shields, J.

[181151]A.; Regillo, C. D.; McDonald, H. R.; Berrocal, M. H.; Gass, J. D.

[181152]M.; Mieler, W. F.: Juxtapapillary capillary hemangiomas: clinical

[181153]features and visual acuity outcomes. Ophthalmology 107: 2240-2249,

[181154]2000.

[181155]

[181156]103. McKusick, V. A.; Abbey, H.; Bettersby, E. J.; Borhani, N. O.;

[181157]Boyer, S. H., IV; Cohen, B. H.; Ferguson-Smith, M. A.; Franke, F.

[181158]R.; Gordon, H.; Handmaker, S. D.; Harris, W. S.; Hawkins, M. R.;

[181159]and 13 others: Medical genetics 1960. J. Chronic Dis. 14:

1-198,

[181160]1961. Fig. 71.

[181161]

[181162]104. Melmon, K. L.; Rosen, S. W.: Lindau's disease: review
of the

[181163]literature and study of a large kindred. Am. J. Med. 36:
595-617,

[181164]1964.

[181165]

[181166]105. Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W.
G., Jr.;

[181167]Pavletich, N. P.: Structure of an HIF-1-alpha-pVHL com-
plex: hydroxyproline

[181168]recognition in signaling. Science 296: 1886-1889, 2002.

[181169]

[181170]106. Mukhopadhyay, D.; Knebelmann, B.; Cohen, H. T.;
Ananth, S.; Sukhatme,

[181171]V. P.: The von Hippel-Lindau tumor suppressor gene
product interacts

[181172]with Sp1 to repress vascular endothelial growth factor
promoter activity. Molec

[181173]. Cell. Biol. 17: 5629–5639, 1997.

[181174]

[181175]107. Neumann, H. P. H.; Berger, D. P.; Sigmund, G.; Blum, U.; Schmidt,

[181176]D.; Parmer, R. J.; Volk, B.; Kirste, G.: Pheochromocytomas, multiple

[181177]endocrine neoplasia type 2, and von Hippel–Lindau disease. New Eng.

[181178]J. Med. 329: 1531–1538, 1993.

[181179]

[181180]108. Neumann, H. P. H.; Eng, C.; Mulligan, L. M.; Glavac, D.; Zauner,

[181181]I.; Ponder, B. A. J.; Crossey, P. A.; Maher, E. R.; Brauch, H.: Consequences

[181182]of direct genetic testing for germline mutations in the clinical management

[181183]of families with multiple endocrine neoplasia, type II. J.A.M.A. 274:

[181184]1149–1151, 1995.

[181185]

[181186]109. Neumann, H. P. H.; Wiestler, O. D.: Clustering of fea–

tures of

[181187] von Hippel–Lindau syndrome: evidence for a complex genetic locus. *Lancet* 337:

[181188] 1052–1054, 1991.

[181189]

[181190] 110. Nibbelink, D. W.; Peters, B. H.; McCormick, W. F.: On the association

[181191] of pheochromocytoma and cerebellar hemangioblastoma. *Neurology* 19:

[181192] 455–460, 1969.

[181193]

[181194] 111. Oberstrass, J.; Reifenberger, G.; Reifenberger, J.; Wechsler,

[181195] W.; Collins, V. P.: Mutation of the von Hippel–Lindau tumour suppressor

[181196] gene in capillary haemangioblastomas of the central nervous system. *J.*

[181197] *Path.* 179: 151–156, 1996.

[181198]

[181199] 112. Ohh, M.; Yauch, R. L.; Lonergan, K. M.; Whaley, J. M.; Stemmer–Rachamimov,

[181200]A. O.; Louis, D. N.; Gavin, B. J.; Kley, N.; Kaelin, W. G., Jr.;
Iliopoulos,

[181201]O.: The von Hippel–Lindau tumor suppressor protein is
required for

[181202]proper assembly of an extracellular fibronectin matrix.
Molec. Cell 1:

[181203]959–968, 1998.

[181204]

[181205]113. Olschwang, S.; Richard, S.; Boisson, C.; Giraud, S.;
Laurent–Puig,

[181206]P.; Resche, F.; Thomas, G.: Germline mutation profile of
the VHL

[181207]gene in von Hippel–Lindau disease and in sporadic he–
mangioblastoma. Hum.

[181208]Mutat. 12: 424–430, 1998.

[181209]

[181210]114. Otenasek, F. J.; Silver, M. L.: Spinal hemangioma
(hemangioblastoma)

[181211]in Lindau's disease: report of six cases in a single family.
J. Neurosurg. 18:

[181212]295–300, 1961.

[181213]

[181214]115. Pack, S. D.; Zbar, B.; Pak, E.; Ault, D. O.; Humphrey, J. S.;

[181215]Pham, T.; Hurley, K.; Weil, R. J.; Park, W.-S.; Kuzmin, I.; Stolle,

[181216]C.; Glenn, G.; Liotta, L. A.; Lerman, M. I.; Klausner, R. D.; Linehan,

[181217]W. M.; Zhuang, Z.: Constitutional von Hippel-Lindau (VHL) gene deletions

[181218]detected in VHL families by fluorescence in situ hybridization. Cancer

[181219]Res. 59: 5560-5564, 1999.

[181220]

[181221]116. Pause, A.; Lee, S.; Lonergan, K. M.; Klausner, R. D.: The von

[181222]Hippel-Lindau tumor suppressor gene is required for cell cycle exit

[181223]upon serum withdrawal. Proc. Nat. Acad. Sci. 95: 993-998, 1998.

[181224]

[181225]117. Price, E. B., Jr.: Papillary cystadenoma of the epididymis:

[181226]a clinicopathologic analysis of 20 cases. Arch. Path. 91:
456–470,

[181227]1971.

[181228]

[181229]118. Probst, A.; Lotz, M.; Heitz, P.: Von Hippel–Lindau's
disease,

[181230]syringomyelia and multiple endocrine tumors: a complex
neuroendocrinopathy. Virchows

[181231]Arch. Path. Anat. Histol. 378: 265–272, 1978.

[181232]

[181233]119. Prowse, A. H.; Webster, A. R.; Richards, F. M.;
Richard, S.;

[181234]Olschwang, S.; Resche, F.; Affara, N. A.; Maher, E. R.: So-
matic inactivation

[181235]of the VHL gene in Von Hippel–Lindau disease tumors.
Am. J. Hum.

[181236]Genet. 60: 765–771, 1997.

[181237]

[181238]120. Rho, Y. M.: Von Hippel–Lindau's disease: a report of
5 cases. Canad.

[181239]Med. Assoc. J. 101: 135–142, 1969.

[181240]

[181241]121. Richard, S.; Croisille, L.; Yvart, J.; Casadeval, N.; Eschwege,

[181242]P.; Aghakhani, N.; David, P.; Gaudric, A.; Scigalla, P.; Hermine,

[181243]O.: Paradoxical secondary polycythemia in von Hippel-Lindau patients

[181244]treated with anti-vascular endothelial growth factor receptor therapy. Blood 99:

[181245]3851-3853, 2002.

[181246]

[181247]122. Richards, F. M.; Crossey, P. A.; Phipps, M. E.; Foster, K.; Latif,

[181248]F.; Evans, G.; Sampson, J.; Lerman, M. I.; Zbar, B.; Affara, N. A.;

[181249]Ferguson-Smith, M. A.; Maher, E. R.: Detailed mapping of germline

[181250]deletions of the von Hippel-Lindau disease tumour suppressor gene. Hum.

[181251]Molec. Genet. 3: 595-598, 1994.

[181252]

[181253]123. Richards, F. M.; Maher, E. R.; Latif, F.; Phipps, M. E.;
Tory,
[181254]K.; Lush, M.; Crossey, P. A.; Oostra, B.; Gustavson, K. H.;
Green,
[181255]J.; Turner, G.; Yates, J. R. W.; Linehan, W. M.; Affara, N. A.;
Lerman,
[181256]M.; Zbar, B.; Ferguson-Smith, M. A.: Detailed genetic
mapping of
[181257]the von Hippel-Lindau disease tumour suppressor gene. J.
Med. Genet. 30:
[181258]104-107, 1993.

[181259]

[181260]124. Richards, F. M.; Phipps, M. E.; Latif, F.; Yao, M.;
Crossey,
[181261]P. A.; Foster, K.; Linehan, W. M.; Affara, N. A.; Lerman, M.
I.; Zbar,
[181262]B.; Ferguson-Smith, M. A.; Maher, E. R.: Mapping the von
Hippel-Lindau
[181263]disease tumour suppressor gene: identification of
germline deletions
[181264]by pulsed field gel electrophoresis. Hum. Molec. Genet. 2:
879-882,

[181265]1993.

[181266]

[181267]125. Richards, F. M.; Schofield, P. N.; Fleming, S.; Maher, E. R.

[181268]: Expression of the von Hippel–Lindau disease tumour suppressor gene

[181269]during human embryogenesis. Hum. Molec. Genet. 5: 639–644, 1996.

[181270]

[181271]126. Rubenstein, J. L.; Yaari, H.: von Hippel–Lindau and the genetics

[181272]of astrocytoma. J. Nat. Cancer Inst. 86: 142–143, 1994.

[181273]

[181274]127. Sander, S.; Normann, T.; Mathisen, W.: Pheochromocytoma associated

[181275]with von–Hippel–Lindau's disease in a family. Scand. J. Urol. Nephrol. 4:

[181276]259–263, 1970.

[181277]

[181278]128. Schechterman, L.: Lindau's disease: report of an unusual case

[181279]and two additional cases in a Negro family. Med. Ann.
D.C. 30: 64–76,
[181280]1961.

[181281]

[181282]129. Schimke, R. N.; Collins, D. L.; Rothberg, P. G.: Func-
tioning

[181283]carotid paraganglioma in the von Hippel–Lindau syn-
drome. (Letter) Am.

[181284]J. Med. Genet. 80: 533–534, 1998.

[181285]

[181286]130. Schoenfeld, A.; Davidowitz, E. J.; Burk, R. D.: A sec-
ond major

[181287]native von Hippel–Lindau gene product, initiated from an
internal

[181288]translation start site, functions as a tumor suppressor.

Proc. Nat.

[181289]Acad. Sci. 95: 8817–8822, 1998.

[181290]

[181291]131. Seizinger, B. R.; Farmer, G.; Haines, J.; Anderson, K.;
Whaley,

[181292]J.; Hettlich, C.; Decker, J.; Rouleau, G.; Smith, D.; Drabkin,

H.;

[181293]Filling-Katz, M.; Neumann, H.; Collins, D.; Hsia, E.; Green, J.; Waziri,

[181294]M.; Gusella, J.; Li, F.: Isolating the gene(s) for von Hippel-Lindau

[181295]disease and renal cell carcinoma. (Abstract) Am. J. Hum. Genet. 45

[181296](suppl.): A32, 1989.

[181297]

[181298]132. Seizinger, B. R.; Rouleau, G. A.; Ozelius, L. J.; Lane,

[181299]A. H.; Farmer, G. E.; Lamiell, J. M.; Haines, J.; Yuen, J.

[181300]W.; Collins, D.; Majoor-Krakauer, D.; et al.: Von Hippel-Lindau

[181301]disease maps to the region of chromosome 3 associated with renal cell

[181302]carcinoma. Nature 332: 268-269, 1988.

[181303]

[181304]133. Seizinger, B. R.; Smith, D. I.; Filling-Katz, M. R.; Neumann,

[181305]H.; Green, J. S.; Choyke, P. L.; Anderson, K. M.; Freiman, R. N.;

[181306]Klauck, S. M.; Whaley, J.; Decker, H.-J. H.; Hsia, Y. E.;

Collins,

[181307]D.; Halperin, J.; Lamiell, J. M.; Oostra, B.; Waziri, M. H.;

Gorin,

[181308]M. B.; Scherer, G.; Drabkin, H. A.; Aronin, N.; Schinzel, A.;

Martuza,

[181309]R. L.; Gusella, J. F.; Haines, J. L.: Genetic flanking markers
refine

[181310]diagnostic criteria and provide insights into the genetics
of Von

[181311]Hippel Lindau disease. Proc. Nat. Acad. Sci. 88:
2864–2868, 1991.

[181312]

[181313]134. Sgambati, M. T.; Stolle, C.; Choyke, P. L.; Walther, M.
M.; Zbar,

[181314]B.; Linehan, W. M.; Glenn, G. M.: Mosaicism in von Hippel–
Lindau

[181315]disease: lessons from kindreds with germline mutations
identified

[181316]in offspring with mosaic parents. Am. J. Hum. Genet. 66:
84–91,

[181317]2000.

[181318]

[181319]135. Sharp, W. V.; Platt, R. L.: Familial pheochromocytoma:
association

[181320]with von-Hippel-Lindau's disease. Angiology 22:
141-146, 1971.

[181321]

[181322]136. Shokeir, M. H. K.: Von Hippel-Lindau syndrome: a
report on three

[181323]kindreds. J. Med. Genet. 7: 155-157, 1970.

[181324]

[181325]137. Silver, M. L.: Hereditary vascular tumors of the ner-
vous system. J.A.M.A.

[181326]156: 1053-1056, 1954.

[181327]

[181328]138. Stolle, C.; Glenn, G.; Zbar, B.; Humphrey, J. S.;
Choyke, P.;

[181329]Walther, M.; Pack, S.; Hurley, K.; Audrey, C.; Klausner, R.;
Linehan,

[181330]W. M.: Improved detection of germline mutations in the
von Hippel-Lindau

[181331]disease tumor suppressor gene. Hum. Mutat. 12:

417-423, 1998.

[181332]

[181333]139. Thomas, M.; Burnside, R. M.: Von Hippel-Lindau disease. Am.

[181334]J. Ophthal. 51: 140-146, 1961.

[181335]

[181336]140. Tisherman, S. E.; Gregg, F. J.; Danowski, T. S.: Familial pheochromocytoma

[181337]. J.A.M.A. 182: 152-156, 1962.

[181338]

[181339]141. Tisherman, S. E.; Tisherman, B. G.; Tisherman, S. A.; Dunmire,

[181340]S.; Levey, G. S.; Mulvihill, J. J.: Three-decade investigation of

[181341]familial pheochromocytoma: an allele of von Hippel-Lindau disease? Arch.

[181342]Int. Med. 153: 2550-2556, 1993.

[181343]

[181344]142. Tory, K.; Brauch, H.; Linehan, M.; Barba, D.; Oldfield, E.; Filling-Katz,

[181345]M.; Seizinger, B.; Nakamura, Y.; White, R.; Marshall, F. F.;

Lerman,

[181346]M. I.; Zbar, B.: Specific genetic change in tumors associated with

[181347]von Hippel–Lindau disease. J. Nat. Cancer Inst. 81: 1097–1101, 1989.

[181348]

[181349]143. Tsuda, H.; Fukushima, S.; Takahashi, M.; Hikosaka, Y.; Hayashi,

[181350]K.: Familial bilateral papillary cystadenoma of the epididymis: report

[181351]of three cases in siblings. Cancer 37: 1831–1839, 1976.

[181352]

[181353]144. Tyers, M.; Willems, A. R.: One ring to rule a superfamily of

[181354]E3 ubiquitin ligases. Science 284: 602–604, 1999.

[181355]

[181356]145. Vance, J. M.; Small, K. W.; Jones, M. A.; Stajich, J. M.; Yamaoka,

[181357]L. H.; Roses, A. D.; Hung, W.–Y.; Pericak–Vance, M. A.: Confirmation

[181358]of linkage in von Hippel–Lindau disease. Genomics 6:

565–567, 1990.

[181359]

[181360]146. van der Harst, E.; de Krijger, R. R.; Dinjens, W. N. M.;

Weeks,

[181361]L. E.; Bonjer, H. J.; Bruining, H. A.; Lamberts, S. W. J.;

Koper,

[181362]J. W.: Germline mutations in the vhl gene in patients presenting

[181363]with pheochromocytomas. Int. J. Cancer. 77: 337–340, 1998.

[181364]

[181365]147. Versteeg, R.: Aberrant methylation in cancer.

(Editorial) Am.

[181366]J. Hum. Genet. 60: 751–754, 1997.

[181367]

[181368]148. Vogelstein, B.: Personal Communication. Baltimore, Md. 1/6/1995.

[181369]

[181370]149. von Hippel, E.: Ueber eine sehr seltene Erkrankung der Netzhaut. Albrecht

[181371]von Graefes Arch. Ophthal. 59: 83–106, 1904.

[181372]

[181373]150. Webster, A. R.; Maher, E. R.; Bird, A. C.; Moore, A. T.:
Risk

[181374]of multisystem disease in isolated ocular angioma
(haemangioblastoma). J.

[181375]Med. Genet. 37: 62–63, 2000.

[181376]

[181377]151. Webster, A. R.; Richards, F. M.; MacRonal, F. E.;
Moore, A.

[181378]T.; Maher, E. R.: An analysis of phenotypic variation in the
familial

[181379]cancer syndrome von Hippel–Lindau disease: evidence for
modifier effects. Am.

[181380]J. Hum. Genet. 63: 1025–1035, 1998.

[181381]

[181382]152. Wells, R. A.; Reeders, S. T.; Green, J.; Johnson, G. J.;
Robson,

[181383]K. J. H.: Linkage studies of von Hippel–Lindau syndrome
using multiple–locus

[181384]hypervariable DNA probes. (Abstract) Cytogenet. Cell
Genet. 46:

[181385]714, 1987.

[181386]

[181387]153. Wesolowski, D. P.; Ellwood, R. A.; Schwab, R. E.;
Farah, J.:

[181388]Hippel–Lindau syndrome in identical twins. Brit. J. Radiol.
54:

[181389]982–986, 1981.

[181390]

[181391]154. Wiesener, M. S.; Seyfarth, M.; Warnecke, C.; Jur-
gensen, J. S.;

[181392]Rosenberger, C.; Morgan, N. V.; Maher, E. R.; Frei, U.;
Eckardt, K.–U.

[181393]: Paraneoplastic erythrocytosis associated with an inacti-
vating point

[181394]mutation of the von Hippel–Lindau gene in a renal cell
carcinoma. Blood 99:

[181395]3562–3565, 2002.

[181396]

[181397]155. Wise, K. S.; Gibson, J. A.: Von Hippel–Lindau's dis-
ease and

[181398]phaeochromocytoma. Brit. Med. J. 1: 441, 1971.

[181399]

[181400]156. Wyburn-Mason, R.: Arteriovenous aneurysm of mid-brain and retina,

[181401]facial naevi and mental changes. Brain 66: 163–203,
1943.

[181402]

[181403]157. Zatyka, M.; da Silva, N. F.; Clifford, S. C.; Morris, M.
R.;

[181404]Wiesener, M. S.; Eckardt, K.–U.; Houlston, R. S.; Richards,
F. M.;

[181405]Latif, F.; Maher, E. R.: Identification of cyclin D1 and other
novel

[181406]targets for the von Hippel–Lindau tumor suppressor gene
by expression

[181407]array analysis and investigation of cyclin D1 genotype as a
modifier

[181408]in von Hippel–Lindau disease. Cancer Res. 62:
3803–3811, 2002.

[181409]

[181410]158. Zbar, B.; Kishida, T.; Chen, F.; Schmidt, L.; Maher, E.
R.; Richards,

[181411]F. M.; Crossey, P. A.; Webster, A. R.; Affara, N. A.; Fergu-
son–Smith,

[181412]M. A.; Brauch, H.; Glavac, D.; and 14 others: Germline mu-
tations

[181413]in the Von Hippel–Lindau disease (VHL) gene in families
from North

[181414]America, Europe, and Japan. Hum. Mutat. 8: 348–357,
1996.

[181415]

[181416]159. Zhuang, Z.; Emmert–Buck, M. R.; Roth, M. J.; Gnarr,
J.; Linehan,

[181417]W. M.; Liotta, L. A.; Lubensky, I. A.: Von Hippel–Lindau
disease

[181418]gene deletion detected in microdissected sporadic human
colon carcinoma

[181419]specimens. Hum. Path. 27: 152–156, 1996.

[181420]

[181421]

[181422]18857.

[181423]Thomas, J. T.; Kilpatrick, M. W.; Lin, K.; Erlacher, L.; Lem–
bessis,

[181424]P.; Costa, T.; Tsipouras, P.; Luyten, F. P.: Disruption of hu–
man

[181425]limb morphogenesis by a dominant negative mutation in
CDMP1. Nature

[181426]Genet. 58–64, 1997.

[181427]

[181428]

[181429]18858.

[181430]Devlin, R. H.; Deeb, S.; Brunzell, J.; Hayden, M. R.: Partial

[181431]gene duplication involving exon–Alu interchange results
in lipoprotein

[181432]lipase deficiency. Am. J. Hum. Genet. 46: 112–119, 1990.

[181433]

[181434]

[181435]18859.

[181436]Dichek, H. L.; Fojo, S. S.; Beg, O. U.; Skarlatos, S. I.; Brun-
zell,

[181437]J. D.; Cutler, G. B., Jr.; Brewer, H. B., Jr.: Identification of

[181438]2 separate allelic mutations in the lipoprotein lipase gene
of a patient

[181439]with the familial hyperchylomicronemia syndrome. J. Biol.
Chem. 266:

[181440]473–477, 1991.

[181441]

[181442]

[181443]18860.

[181444]Eckel, R. H.: Lipoprotein lipase: a multifunctional enzyme
relevant

[181445]to common metabolic diseases. New Eng. J. Med. 320:
1060–1068, 1989.

[181446]

[181447]

[181448]18861.

[181449]Emi, M.; Hata, A.; Robertson, M.; Iverius, P.–H.; Hegele, R.;

[181450]Lalouel, J.–M.: Lipoprotein lipase deficiency resulting from
a nonsense

[181451]mutation in exon 3 of the lipoprotein lipase gene. Am. J.
Hum. Genet. 47:

[181452]107–111, 1990.

[181453]

[181454]

[181455]18862.

[181456]Emi, M.; Wilson, D. E.; Iverius, P.-H.; Wu, L.; Hata, A.;
Hegele,

[181457]R.; Williams, R. R.; Lalouel, J.-M.: Missense mutation
(gly-to-glu188)

[181458]of human lipoprotein lipase imparting functional defi-
ciency. J. Biol.

[181459]Chem. 265: 5910-5916, 1990.

[181460]

[181461]

[181462]18863.

[181463]Faustinella, F.; Chang, A.; Van Biervliet, J. P.; Rosseneu, M.;

[181464]Vinaimont, N.; Smith, L. C.; Chen, S.-H.; Chan, L.: Catalytic
triad

[181465]residue mutation (asp156-to-gly) causing familial
lipoprotein lipase

[181466]deficiency: co-inheritance with a nonsense mutation
(ser447-to-ter)

[181467]in a Turkish family. J. Biol. Chem. 266: 14418-14424,
1991.

[181468]

[181469]

[181470]18864.

[181471]Feoli-Fonseca, J. C.; Levy, E.; Godard, M.; Lambert, M.: Familial

[181472]lipoprotein lipase deficiency in infancy: clinical, biochemical, and

[181473]molecular study. J. Pediatr. 133: 417–423, 1998.

[181474]

[181475]

[181476]18865.

[181477]Fisher, R. M.; Humphries, S. E.; Talmud, P. J.: Common variation

[181478]in the lipoprotein lipase gene: effects on plasma lipids and risk

[181479]of atherosclerosis. Atherosclerosis 135: 145–159, 1997.

[181480]

[181481]

[181482]18866.

[181483]Franklin, S. M.: Splenomegaly with lipaemia. Proc. Roy. Soc.

[181484]Med. 30: 711, 1937.

[181485]

[181486]

[181487]18867.

[181488]Funke, H.; Klug, J.; Assmann, G.: Hind III RFLP in the
lipoprotein

[181489]lipase gene, (LPL). Nucleic Acids Res. 15: 9102, 1987.

[181490]

[181491]

[181492]18868.

[181493]Funke, H.; Reckwerth, A.; Stapenhorst, D.; Schulze Beier-
ing, M.;

[181494]Jansen, M.; Assmann, G.: Bst NI (Eco RII) RFLP in the
lipoprotein

[181495]lipase gene (LPL). Nucleic Acids Res. 16: 2741, 1988.

[181496]

[181497]

[181498]18869.

[181499]Gagne, C.; Brun, L. D.; Julien, P.; Moorjani, S.; Lupien, P. J.

[181500]: Primary lipoprotein lipase activity deficiency: clinical in-

vestigation

[181501]of a French Canadian population. Canad. Med. Assoc. J.
140: 405–411,

[181502]1989.

[181503]

[181504]

[181505]18870.

[181506]Gilbert, B.; Rouis, M.; Griglio, S.; de Lumley, L.; Laplaud,
P.–M.

[181507]: Lipoprotein lipase (LPL) deficiency: a new patient ho-
mozygote for

[181508]the preponderant mutation gly88–to–glu in the human
LPL gene and review

[181509]of reported mutations: 75% are clustered in exons 5 and
6. Ann. Genet. 44:

[181510]25–32, 2001.

[181511]

[181512]

[181513]18871.

[181514]Ginzinger, D. G.; Lewis, M. E. S.; Ma, Y.; Jones, B. R.; Liu,

[181515]G.; Jones, S. D.; Hayden, M. R.: A mutation in the lipoprotein lipase

[181516]gene is the molecular basis of chylomicronemia in a colony of domestic

[181517]cats. J. Clin. Invest. 97: 1257–1266, 1996.

[181518]

[181519]

[181520]18872.

[181521]Gotoda, T.; Yamada, N.; Kawamura, M.; Kozaki, K.; Mori, N.; Ishibashi,

[181522]S.; Shimano, H.; Takaku, F.; Yazaki, Y.; Furuichi, Y.; Murase, T.

[181523]: Heterogeneous mutations in the human lipoprotein lipase gene in

[181524]patients with familial lipoprotein lipase deficiency. J. Clin. Invest. 88:

[181525]1856–1864, 1991.

[181526]

[181527]

[181528]18873.

[181529]Gotoda, T.; Yamada, N.; Murase, T.; Miyake, S.; Murakami, R.;

[181530]Kawamura, M.; Kozaki, K.; Mori, N.; Shimano, H.; Shimada, M.; Yazaki,

[181531]Y.: A newly identified null allelic mutation in the human lipoprotein

[181532]lipase (LPL) gene of a compound heterozygote with familial LPL deficiency. Biochim.

[181533]Biophys. Acta 1138: 353–356, 1992.

[181534]

[181535]

[181536]18874.

[181537]Hata, A.; Emi, M.; Luc, G.; Basdevant, A.; Gambert, P.; Iverius,

[181538]P.–H.; Lalouel, J.–M.: Compound heterozygote for lipoprotein lipase

[181539]deficiency: ser-to-thr(244) and transition in 3-prime splice site

[181540]of intron 2 (AG-to-AA) in the lipoprotein lipase gene. Am. J. Hum.

[181541]Genet. 47: 721–726, 1990.

[181542]

[181543]

[181544]18875.

[181545]Hata, A.; Ridinger, D. N.; Sutherland, S. D.; Emi, M.;

Kwong,

[181546]L. K.; Shuhua, J.; Lubbers, A.; Guy-Grand, B.; Basdevant,

A.; Iverius,

[181547]P. H.; Wilson, D. E.; Lalouel, J.-M.: Missense mutations in

exon

[181548]5 of the human lipoprotein lipase gene: inactivation cor-

relates with

[181549]loss of dimerization. J. Biol. Chem. 267: 20132-20139,

1992.

[181550]

[181551]

[181552]18876.

[181553]Haubenwallner, S.; Horl, G.; Shachter, N. S.; Presta, E.;

Fried,

[181554]S. K.; Hofler, G.; Kostner, G. M.; Breslow, J. L.; Zechner, R.:

A

[181555]novel missense mutation in the gene for lipoprotein lipase

resulting

[181556]in a highly conservative amino acid substitution

(asp180-to-glu) causes

[181557]familial chylomicronemia (type I hyperlipoproteinemia).

Genomics 18:

[181558]392–396, 1993.

[181559]

[181560]

[181561]18877.

[181562]Havel, R. J.; Gordon, R. S.: Idiopathic hyperlipemia:

metabolic

[181563]studies in an affected family. J. Clin. Invest. 39:

1777–1790, 1960.

[181564]

[181565]

[181566]18878.

[181567]Heaney, A. P.; Sharer, N.; Rameh, B.; Braganza, J. M.; Dur-

rington,

[181568]P. N.: Prevention of recurrent pancreatitis in familial

lipoprotein

[181569]lipase deficiency with high-dose antioxidant therapy. J.

Clin. Endocr.

[181570]Metab. 84: 1203–1205, 1999.

[181571]

[181572]

[181573]18879.

[181574]Heinzmann, C.; Ladas, J.; Antonarakis, S.; Kirchgessner, T.;

[181575]Schotz, M.; Lusis, A. J.: RFLP for the human lipoprotein lipase (LPL)

[181576]gene: HindIII. Nucleic Acids Res. 15: 6763, 1987.

[181577]

[181578]

[181579]18880.

[181580]Heizmann, C.; Kirchgessner, T.; Kwiterovich, P. O.; Ladas, J.

[181581]A.; Derby, C.; Antonarakis, S. E.; Lusis, A. J.: DNA polymorphism

[181582]haplotypes of the human lipoprotein lipase gene: possible association

[181583]with high density lipoprotein levels. Hum. Genet. 86:

578–584, 1991.

[181584]

[181585]

[181586]18881.

[181587]Henderson, H.; Ma, Y.; Kastelein, J.; Roederer, G.; Julien, P.;

[181588]Brunzell, J.; Hayden, M. R.: Identification of the molecular defects

[181589]underlying chylomicronemia in the majority of 75 separate probands

[181590]with LPL deficiency.(Abstract) Clin. Res. 39: 336A, 1991.

[181591]

[181592]

[181593]18882.

[181594]Henderson, H. E.; Bijvoet, S. M.; Mannens, M. A. M. M.; Bruin,

[181595]T.; Erkelens, D. W.; Hayden, M. R.; Kastelein, J. J. P.: Ile225-to-thr

[181596]loop mutation in the lipoprotein lipase (LPL) gene is a de novo event. Am.

[181597]J. Med. Genet. 78: 313–316, 1998.

[181598]

[181599]

[181600]18883.

[181601]Sjalander, A.; Birgander, R.; Rannug, A.; Alexandrie, A.-K.;
Tornling,

[181602]G.; Beckman, G.: Association between the p21 codon
31A1 (arg) allele

[181603]and lung cancer. Hum. Hered. 46: 221-225, 1996.

[181604]

[181605]

[181606]18884.

[181607]Fryns, J. P.; van den Berghe, K.; van Assche, A.; van den
Berghe,

[181608]H.: Prenatal diagnosis of campomelic dwarfism. Clin.
Genet. 19:

[181609]199-201, 1981.

[181610]

[181611]

[181612]18885.

[181613]Mellows, H. J.; Pryse-Davies, J.; Bennett, M. J.; Carter, C. O.

[181614]: The camptomelic syndrome in two female siblings. Clin.

Genet. 18:

[181615]137–141, 1980.

[181616]

[181617]

[181618]18886.

[181619]Millauer, B.; Shawver, L. K.; Plate, K. H.; Risau, W.; Ullrich,

[181620]A.: Glioblastoma growth inhibited in vivo by a dominant-negative

[181621]Flk-1 mutant. Nature 367: 576–579, 1994.

[181622]

[181623]

[181624]18887.

[181625]Cho, K. R.; Oliner, J. D.; Simons, J. W.; Hedrick, L.; Fearon,

[181626]E. R.; Preisinger, A. C.; Hedge, P.; Silverman, G. A.; Vogelstein,

[181627]B.: The DCC gene: structural analysis and mutations in colorectal

[181628]carcinomas. Genomics 19: 525–531, 1994.

[181629]

[181630]

[181631]18888.

[181632]Fazeli, A.; Dickinson, S. L.; Hermiston, M. L.; Tighe, R. V.;
Steen,

[181633]R. G.; Small, C. G.; Stoeckli, E. T.; Keino–Masu, K.; Masu,
M.; Rayburn,

[181634]H.; Simons, J.; Bronson, R. T.; Gordon, J. I.; Tessier–Lavi–
gne, M.;

[181635]Weinberg, R. A.: Phenotype of mice lacking functional
Deleted in

[181636]colorectal cancer (Dcc) gene. Nature 386: 796–804, 1997.

[181637]

[181638]

[181639]18889.

[181640]Fearon, E. R.; Cho, K. R.; Nigro, J. M.; Kern, S. E.; Simons, J.

[181641]W.; Ruppert, J. M.; Hamilton, S. R.; Preisinger, A. C.;
Thomas, G.;

[181642]Kinzler, K. W.; Vogelstein, B.: Identification of a chromo–

some 18q

[181643]gene that is altered in colorectal cancers. Science 247:
49–56,

[181644]1990.

[181645]

[181646]

[181647]18890.

[181648]Gotley, D. C.; Reeder, J. A.; Fawcett, J.; Walsh, M. D.; Bates,
[181649]P.; Simmons, D. L.; Antalis, T. M.: The deleted in colon
cancer (DCC)

[181650]gene is consistently expressed in colorectal cancers and
metastases. Oncogene 13:

[181651]787–795, 1996.

[181652]

[181653]

[181654]18891.

[181655]Hohne, M. W.; Halatsch, M.–E.; Kahl, G. F.; Weinel, R. J.:
Frequent

[181656]loss of expression of the potential tumor suppressor gene
DCC in ductal

[181657]pancreatic adenocarcinoma. Cancer Res. 52: 2616–2619,
1992.

[181658]

[181659]

[181660]18892.

[181661]Jen, J.; Kim, H.; Piantadosi, S.; Liu, Z.–F.; Levitt, R. C.; Sis-
tonen,

[181662]P.; Kinzler, K. W.; Vogelstein, B.; Hamilton, S. R.: Allelic
loss

[181663]of chromosome 18q and prognosis in colorectal cancer.
New Eng. J.

[181664]Med. 331: 213–221, 1994.

[181665]

[181666]

[181667]18893.

[181668]Keino–Masu, K.; Masu, M.; Hinck, L.; Leonardo, E. D.;
Chan, S.

[181669]S.–Y.; Culotti, J. G.; Tessier–Lavigne, M.: Deleted in col-
orectal

[181670]cancer (DCC) encodes a netrin receptor. Cell 87: 175–185,
1996.

[181671]

[181672]

[181673]18894.

[181674]Keynes, R.; Cook, G. M. W.: Axon guidance molecules. Cell
83:

[181675]161–169, 1995.

[181676]

[181677]

[181678]18895.

[181679]Kolodziej, P. A.; Timpe, L. C.; Mitchell, K. J.; Fried, S. R.;

[181680]Goodman, C. S.; Jan, L. Y.; Jan, Y. N.: Frazzled encodes a
Drosophila

[181681]member of the DCC immunoglobulin subfamily and is re-
quired for CNS

[181682]and motor axon guidance. Cell 87: 197–204, 1996.

[181683]

[181684]

[181685]18896.

[181686]Maesawa, C.; Tamura, G.; Ogasawara, S.; Suzuki, Y.;

Sakata, K.;

[181687]Sugimura, J.; Nishizuka, S.; Sato, N.; Ishida, K.; Saito, K.;

Satodate,

[181688]R.: Loss of heterozygosity at the DCC gene locus is not
crucial for

[181689]the acquisition of metastatic potential in oesophageal
squamous cell

[181690]carcinoma. *Europ. J. Cancer.* 32A: 896–898, 1996.

[181691]

[181692]

[181693]18897.

[181694]Mehlen, P.; Rabizadeh, S.; Snipas, S. J.; Assa-Munt, N.;
Salvesen,

[181695]G. S.; Bredesen, D. E.: The DCC gene product induces
apoptosis by

[181696]a mechanism requiring receptor proteolysis. *Nature* 395:
801–804,

[181697]1998.

[181698]

[181699]

[181700]18898.

[181701]Miyake, S.; Nagai, K.; Yoshino, K.; Oto, M.; Endo, M.;
Yuasa,

[181702]Y.: Point mutations and allelic deletion of tumor suppres-
sor gene

[181703]DCC in human esophageal squamous cell carcinomas and
their relation

[181704]to metastasis. Cancer Res. 54: 3007–3010, 1994.

[181705]

[181706]

[181707]18899.

[181708]Nigro, J. M.; Cho, K. R.; Fearon, E. R.; Kern, S. E.; Ruppert,

[181709]J. M.; Oliner, J. D.; Kinzler, K. W.; Vogelstein, B.: Scrambled
exons. Cell 64:

[181710]607–613, 1991.

[181711]

[181712]

[181713]18900.

[181714]Shibata, D.; Reale, M. A.; Lavin, P.; Silverman, M.; Fearon,
E.

[181715]R.; Steele, G., Jr.; Jessup, J. M.; Loda, M.; Summerhayes, I.

C.:

[181716]The DCC protein and prognosis in colorectal cancer. New
Eng. J. Med. 335:

[181717]1727–1732, 1996.

[181718]

[181719]

[181720]18901.

[181721]Stein, E.; Tessier–Lavigne, M.: Hierarchical organization of
[181722]guidance receptors: silencing of netrin attraction by Slit
through

[181723]a Robo/DCC receptor complex. Science 291: 1928–1938,
2001.

[181724]

[181725]

[181726]18902.

[181727]Tanaka, K.; Oshimura, M.; Kikuchi, R.; Seki, M.; Hayashi,
T.;

[181728]Miyaki, M.: Suppression of tumorigenicity in human colon
carcinoma

[181729]cells by introduction of normal chromosome 5 or 18. Na–

ture 349:

[181730]340–342, 1991.

[181731]

[181732]

[181733]18903.

[181734]Uchino, S.; Tsuda, H.; Noguchi, M.; Yokota, J.; Terada, M.;
Saito,

[181735]T.; Kobayashi, M.; Sugimura, T.; Hirohashi, S.: Frequent
loss of

[181736]heterozygosity at the DCC locus in gastric cancer. Cancer
Res. 52:

[181737]3099–3102, 1992.

[181738]

[181739]

[181740]18904.

[181741]Vogelstein, B.: Personal Communication. Baltimore, Md.
11/30/1995.

[181742]

[181743]

[181744]18905.

[181745]Zetter, B. R.: Adhesion molecules in tumor metastasis.
Semin.

[181746]Cancer Biol. 4: 219–229, 1993.

[181747]

[181748]

[181749]18906.

[181750]Devor, E. J.; Grandy, D. K.; Civelli, O.; Litt, M.; Burgess, A.

[181751]K.; Isenberg, K. E.; van de Wetering, B. J. M.; Oostra, B.:

Genetic

[181752]linkage is excluded for the D2–dopamine receptor

lambda–HD2G1 and

[181753]flanking loci on chromosome 11q22–q23 in Tourette syn-
drome. Hum.

[181754]Hered. 40: 105–108, 1990.

[181755]

[181756]

[181757]18907.

[181758]Urbanek, M.; Legro, R. S.; Driscoll, D. A.; Azziz, R.;

Ehrmann,

[181759]D. A.; Norman, R. J.; Strauss, J. F., III; Spielman, R. S.;

Dunaif,

[181760]A.: Thirty-seven candidate genes for polycystic ovary syndrome: strongest

[181761]evidence for linkage is with follistatin. Proc. Nat. Acad. Sci. 96:

[181762]8573–8578, 1999.

[181763]

[181764]

[181765]18908.

[181766]Freemantle, S. J.; Taylor, S. M.; Krystal, G.; Moran, R. G.: Upstream

[181767]organization of and multiple transcripts from the human folylpoly-gamma-glutamate

[181768]synthetase gene. J. Biol. Chem. 270: 9579–9584, 1995.

[181769]

[181770]

[181771]18909.

[181772]Garrow, T. A.; Admon, A.; Shane, B.: Expression cloning of

a human

[181773]cDNA encoding folylpoly(gamma-glutamate) synthetase
and determination

[181774]of its primary structure. Proc. Nat. Acad. Sci. 89:
9151-9155, 1992.

[181775]

[181776]

[181777]18910.

[181778]Jones, C.; Kao, F.-T.: Regional mapping of the folylpolyg-
lutamate

[181779]synthetase gene (FPGS) to 9cen-q34. (Abstract) Cytogenet.
Cell Genet. 37:

[181780]499-500, 1984.

[181781]

[181782]

[181783]18911.

[181784]Jones, C.; Kao, F.-T.; Taylor, R. T.: Chromosomal assign-
ment of

[181785]the gene for folylpolyglutamate synthetase to human
chromosome 9. Cytogenet.

[181786]Cell Genet. 28: 181-194, 1980.

[181787]

[181788]

[181789]18912.

[181790]Jones, C.; Kao, F. T.: Assignment of the human gene complementing

[181791]the auxotrophic marker GAT-minus to chromosome 9.
(Abstract) Cytogenet.

[181792]Cell Genet. 25: 168, 1979.

[181793]

[181794]

[181795]18913.

[181796]Kao, F. T.; Puck, T. T.: Genetics of somatic mammalian cells.

[181797]VII. Induction and isolation of nutritional mutants in Chinese hamster

[181798]cells. Proc. Nat. Acad. Sci. 60: 1275-1281, 1968.

[181799]

[181800]

[181801]18914.

[181802]Sussman, D. J.; Milman, G.; Shane, B.: Characterization of human

[181803]folylpolyglutamate synthetase expressed in Chinese hamster ovary cells. Somat.

[181804]Cell Molec. Genet. 12: 531–540, 1986.

[181805]

[181806]

[181807]18915.

[181808]Taylor, R. T.; Hanna, M. L.: Folate-dependent enzymes in cultured

[181809]Chinese hamster cells: folylpolyglutamate synthetase and its absence

[181810]in mutants auxotrophic for glycine, adenosine and thymidine. Arch.

[181811]Biochem. Biophys. 181: 331–344, 1977.

[181812]

[181813]

[181814]18916.

[181815]Taylor, S. M.; Freemantle, S. J.; Moran, R. G.: Structural organization

[181816]of the human folypoly-gamma-glutamate synthetase
gene: evidence for

[181817]a single genomic locus. Cancer Res. 55: 6030–6034,
1995.

[181818]

[181819]

[181820]18917.

[181821]Matsuo, K.; Owens, J. M.; Tonko, M.; Elliott, C.; Chambers,
T.

[181822]J.; Wagner, E. F.: Fos1 is a transcriptional target of c-Fos
during

[181823]osteoclast differentiation. Nature Genet. 24: 184–187,
2000.

[181824]

[181825]

[181826]18918.

[181827]Buters, J. T. M.; Tang, B.–K.; Pineau, T.; Gelboin, H. V.;
Kimura,

[181828]S.; Gonzalez, F. J.: Role of CYP1A2 in caffeine pharmacoki-
netics

[181829]and metabolism: studies using mice deficient in CYP1A2.

Pharmacogenetics 6:

[181830]291–296, 1996.

[181831]

[181832]

[181833]18919.

[181834]Butler, M. A.; Iwasaki, M.; Guengerich, F. P.; Kadlubar, F. F.

[181835]: Human cytochrome P–450(PA) (P–450IA2), the
phenacetin O–deethylase,

[181836]is primarily responsible for the hepatic 3–demethylation
of caffeine

[181837]and N–oxidation of carcinogenic arylamines. Proc. Nat.
Acad. Sci. 86:

[181838]7696–7700, 1989.

[181839]

[181840]

[181841]18920.

[181842]Christiansen, L.; Bygum, A.; Jensen, A.; Thomsen, K.;
Brandrup,

[181843]F.; Horder, M.; Petersen, N. E.: Association between
CYP1A2 polymorphism

[181844]and susceptibility to porphyria cutanea tarda. Hum. Genet.

107:

[181845]612–614, 2000.

[181846]

[181847]

[181848]18921.

[181849]Devonshire, H. W.; Kong, I.; Cooper, M.; Sloan, T. P.; Idle, J.

[181850]R.; Smith, R. L.: The contribution of genetically determined oxidation

[181851]status to inter-individual variation in phenacetin disposition. Brit.

[181852]J. Clin. Pharm. 16: 157–166, 1983.

[181853]

[181854]

[181855]18922.

[181856]Ikeya, K.; Jaiswal, A. K.; Owens, R. A.; Jones, J. E.; Nebert,

[181857]D. W.; Kimura, S.: Human CYP1A2: sequence, gene structure, comparison

[181858]with the mouse and rat orthologous gene, and differences

in liver

[181859]1A2 mRNA expression. Molec. Endocr. 3: 1399–1408, 1989.

[181860]

[181861]

[181862]18923.

[181863]Jaiswal, A. K.; Nebert, D. W.; McBride, O. W.; Gonzalez, F. J.

[181864]: Human P(3)450: cDNA and complete protein sequence, repetitive Alu

[181865]sequences in the 3–prime nontranslated region, and localization of

[181866]gene to chromosome 15. J. Exp. Path. 3: 1–17, 1987.

[181867]

[181868]

[181869]18924.

[181870]Liang, H.–C. L.; Li, H.; McKinnon, R. A.; Duffy, J. J.; Potter,

[181871]S. S.; Puga, A.; Nebert, D. W.: Cyp1a2(–/–) null mutant mice develop

[181872]normally but show deficient drug metabolism. Proc. Nat. Acad. Sci. 93:

[181873]1671–1676, 1996.

[181874]

[181875]

[181876]18925.

[181877]Nebert, D. W.: Personal Communication. Bethesda, Md.
2/3/1988.

[181878]

[181879]

[181880]18926.

[181881]Sesardic, D.; Boobis, A. R.; Edwards, R. J.; Davies, D. S.: A

[181882]form of cytochrome P450 in man, orthologous to form d
in the rat,

[181883]catalyses the O-deethylation of phenacetin and is in-
ducible by cigarette

[181884]smoking. Brit. J. Clin. Pharm. 26: 363–372, 1988.

[181885]

[181886]

[181887]18927.

[181888]Tantcheva-Poor, I.; Zaigler, M.; Rietbrock, S.; Fuhr, U.: Es-

timation

[181889]of cytochrome P-450 CYP1A2 activity in 863 healthy Caucasians using

[181890]a saliva-based caffeine test. Pharmacogenetics 9: 131-144, 1999.

[181891]

[181892]

[181893]18928.

[181894]Wooding, S. P.; Watkins, W. S.; Bamshad, M. J.; Dunn, D. M.; Weiss,

[181895]R. B.; Jorde, L. B.: DNA sequence variation in a 3.7-kb noncoding

[181896]sequence 5-prime of the CYP1A2 gene: implications for human population

[181897]history and natural selection. Am. J. Hum. Genet. 71: 528-542, 2002.

[181898]

[181899]

[181900]18929.

[181901]Bale, A. E.; Mitchell, A. L.; Gonzalez, F. J.; McBride, O. W.:

[181902]Localization of CYP2F1 by multipoint linkage analysis and

pulsed-field

[181903]gel electrophoresis. Genomics 10: 284–286, 1991.

[181904]

[181905]

[181906]18930.

[181907]Mitchell, A. L.; Bale, A. E.; Gonzalez, F.; McBride, O. W.:

Mapping

[181908]of cytochrome p450 IIA, IIB, and IIF subfamilies on chromosome 19

[181909]by linkage analysis. (Abstract) Cytogenet. Cell Genet. 51: 1045,

[181910]1989.

[181911]

[181912]

[181913]18931.

[181914]Nhamburo, P. T.; Kimura, S.; McBride, O. W.; Kozak, C. A.;

Gelboin,

[181915]H. V.; Gonzalez, F. J.: The human CYP2F gene subfamily: identification

[181916]of a cDNA encoding a new cytochrome P450, cDNA-di-

rected expression,

[181917]and chromosome mapping. Biochemistry 29: 5491–5499,
1990.

[181918]

[181919]

[181920]18932.

[181921]Imaoka, S.; Yoneda, Y.; Sugimoto, T.; Hiroi, T.; Yamamoto,
K.;

[181922]Nakatani, T.; Funae, Y.: CYP4B1 is a possible risk factor
for bladder

[181923]cancer in humans. Biochem. Biophys. Res. Commun. 277:
776–780, 2000.

[181924]

[181925]

[181926]18933.

[181927]Nhamburo, P. T.; Gonzalez, F. J.; McBride, O. W.; Gelboin,
H. V.;

[181928]Kimura, S.: Identification of a new P450 expressed in hu-
man lung:

[181929]complete cDNA sequence, cDNA-directed expression, and
chromosome mapping. Biochemistry 28:

[181930]8060–8066, 1989.

[181931]

[181932]

[181933]18934.

[181934]Thum, T.; Borlak, J.: Gene expression in distinct regions of
the

[181935]heart. Lancet 355: 979–983, 2000.

[181936]

[181937]

[181938]18935.

[181939]Xu, L.; Xia, J.; Jiang, H.; Zhou, J.; Li, H.; Wang, D.; Pan, Q.;

[181940]Long, Z.; Fan, C.; Deng, H.–X.: Mutation analysis of hereditary multiple
itary multiple

[181941]exostoses in the Chinese. Hum. Genet. 105: 45–50, 1999.

[181942]

[181943]

[181944]18936.

[181945]Joutel, A.; Favrole, P.; Labauge, P.; Chabriat, H.; Lescoat,

C.;

[181946]Andreux, F.; Domenga, V.; Cecillon, M.; Vahedi, K.;

Ducros, A.; Cave-Riant,

[181947]F.; Bousser, M. G.; Tournier-Lasserre, E.: Skin biopsy immunostaining

[181948]with a Notch3 monoclonal antibody for CADASIL diagnosis. Lancet 358:

[181949]2049–2051, 2001.

[181950]

[181951]

[181952]18937.

[181953]Barinaga, M.: An intriguing new lead on Huntington's disease. Science 271:

[181954]1233–1234, 1996.

[181955]

[181956]

[181957]18938.

[181958]Burke, J. R.; Enghild, J. J.; Martin, M. E.; Jou, Y.-S.; Myers,

[181959]R. M.; Roses, A. D.; Vance, J. M.; Strittmatter, W. J.: Huntingtin

[181960]and DRPLA proteins selectively interact with the enzyme
GAPDH. Nature

[181961]Med. 2: 347–350, 1996.

[181962]

[181963]

[181964]18939.

[181965]Eubanks, J. H.; Djabali, M.; Selleri, L.; Grandy, D. K.; Civelli,

[181966]O.; McElligott, D. L.; Evans, G. A.: Structure and linkage of
the

[181967]D2 dopamine receptor and neural cell adhesion molecule
genes on human

[181968]chromosome 11q23. Genomics 14: 1010–1018, 1992.

[181969]

[181970]

[181971]18940.

[181972]Gejman, P. V.; Ram, A.; Gelernter, J.; Friedman, E.; Cao, Q.;

[181973]Pickar, D.; Blum, K.; Noble, E. P.; Kranzler, H. R.; O'Malley,
S.;

[181974]Hamer, D. H.; Whitsitt, F.; Rao, P.; DeLisi, L. E.; Virkkunen,

M.;

[181975]Linnoila, M.; Goldman, D.; Gershon, E. S.: No structural mutation

[181976]in the dopamine D2 receptor gene in alcoholism or schizophrenia: analysis

[181977]using denaturing gradient gel electrophoresis. J.A.M.A. 271: 204–208,

[181978]1994.

[181979]

[181980]

[181981]18941.

[181982]Gelernter, J.; Grandy, D. K.; Bunzow, J.; Pakstis, A. J.; Civelli,

[181983]O.; Retief, A. E.; Litt, M.; Kidd, K. K.: D(2) dopamine receptor

[181984]locus (probe hD2G1) maps close to D11S29 (probe L7) and is also linked

[181985]to PBGD (probe PBGD0.0) and D11S84 (probe p-2-7-ID6) on 11q. (Abstract) Cytogenet.

[181986]Cell Genet. 51: 1002 only, 1989.

[181987]

[181988]

[181989]18942.

[181990]Gelernter, J.; Pakstis, A. J.; Grandy, D.; Litt, M.; Retief, A.

[181991]E.; Kennedy, J. L.; Hing-Loh, A.; Schoolfield, G.; Civelli, O.;
Kidd,

[181992]K. K.: Linkage map of eight human chromosome 11q
markers, including

[181993]DRD2, spanning 60 cM. Cytogenet. Cell Genet. 60: 26–28,
1992.

[181994]

[181995]

[181996]18943.

[181997]Gelernter, J.; Pakstis, A. J.; Pauls, D. L.; Kurlan, R.;
Gancher,

[181998]S. T.; Civelli, O.; Grandy, D.; Kidd, K. K.: Gilles de la
Tourette

[181999]syndrome is not linked to D2-dopamine receptor. Arch.
Gen. Psychiat. 47:

[182000]1073–1077, 1990.

[182001]

[182002]

[182003]18944.

[182004]Giros, B.; Sokoloff, P.; Martres, M.-P.; Riou, J.-F.; Emorine,

[182005]L. J.; Schwartz, J.-C.: Alternative splicing directs the expression
pression

[182006]of two D(2) dopamine receptor isoforms. Nature 342:
923-926, 1989.

[182007]

[182008]

[182009]18945.

[182010]Grandy, D. K.; Litt, M.; Allen, L.; Bunzow, J. R.; Marchionni,

[182011]M.; Makam, H.; Reed, L.; Magenis, R. E.; Civelli, O.: The
human dopamine

[182012]D(2) receptor gene is located on chromosome 11 at
q22-q23 and identifies

[182013]a TaqI RFLP. Am. J. Hum. Genet. 45: 778-785, 1989.

[182014]

[182015]

[182016]18946.

[182017]Grandy, D. K.; Marchionni, M. A.; Makam, H.; Stofko, R. E.;
Alfano,

[182018]M.; Frothingham, L.; Fischer, J. B.; Burke–Howie, K. J.; Bun-
zow, J.

[182019]R.; Server, A. C.; Civelli, O.: Cloning of the cDNA and gene
for

[182020]a human D2 dopamine receptor. Proc. Nat. Acad. Sci. 86:
9762–9766,

[182021]1989.

[182022]

[182023]

[182024]18947.

[182025]Klein, C.; Brin, M. F.; Kramer, P.; Sena–Esteves, M.; de
Leon,

[182026]D.; Doheny, D.; Bressman, S.; Fahn, S.; Breakefield, X. O.;
Ozelius,

[182027]L. J.: Association of a missense change in the D2
dopamine receptor

[182028]with myoclonus dystonia. Proc. Nat. Acad. Sci. 96:
5173–5176, 1999.

[182029]

[182030]

[182031]18948.

[182032]Maldonado, R.; Salardi, A.; Valverde, O.; Samad, T. A.;
Roques,

[182033]B. P.; Borrelli, E.: Absence of opiate rewarding effects in
mice

[182034]lacking dopamine D2 receptors. Nature 388: 586–589,
1997.

[182035]

[182036]

[182037]18949.

[182038]Milligan, G.: Receptors as kissing cousins. Science 288:
65–67,

[182039]2000.

[182040]

[182041]

[182042]18950.

[182043]Monsma, F. J., Jr.; McVittie, L. D.; Gerfen, C. R.; Mahan, L.

[182044]C.; Sibley, D. R.: Multiple D(2) dopamine receptors pro–

duced by alternative

[182045]RNA splicing. Nature 342: 926–929, 1989.

[182046]

[182047]

[182048]18951.

[182049]Oliveri, R. L.; Annesi, G.; Zappia, M.; Civitelli, D.; Monte-
santi,

[182050]R.; Branca, D.; Nicoletti, G.; Spadafora, P.; Pasqua, A. A.;
Cittadella,

[182051]R.; Andreoli, V.; Gambardella, A.; Aguglia, U.; Quattrone,
A.: Dopamine

[182052]D2 receptor gene polymorphism and the risk of levodopa-
induced dyskinesias

[182053]in PD. Neurology 53: 1425–1430, 1999.

[182054]

[182055]

[182056]18952.

[182057]Rocheville, M.; Lange, D. C.; Kumar, U.; Patel, S. C.; Patel,

[182058]R. C.; Patel, Y. C.: Receptors for dopamine and somato-
statin: formation

[182059]of hetero-oligomers with enhanced functional activity.

Science 288:

[182060]154–157, 2000.

[182061]

[182062]

[182063]18953.

[182064]Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.;
Grandy,

[182065]D.; Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an
11q22.3–9p22

[182066]translocation with affective disorder: proximity of the
dopamine D2

[182067]receptor gene relative to the translocation breakpoint.
(Abstract) Am.

[182068]J. Hum. Genet. 45 (suppl.): A220 only, 1989.

[182069]

[182070]

[182071]18954.

[182072]St Clair, D.; Blackwood, D.; Muir, W.; Carothers, A.; Walker,

[182073]M.; Spowart, G.; Gosden, C.; Evans, H. J.: Association

within a family

[182074]of a balanced autosomal translocation with major mental illness. Lancet 336:

[182075]13–16, 1990.

[182076]

[182077]

[182078]18955.

[182079]Suarez, B. K.; Parsian, A.; Hampe, C. L.; Todd, R. D.; Reich,
[182080]T.; Cloninger, C. R.: Linkage disequilibria at the D2
dopamine receptor

[182081]locus (DRD2) in alcoholics and controls. Genomics 19:
12–20, 1994.

[182082]

[182083]

[182084]18956.

[182085]Usiello, A.; Baik, J.–H.; Rouge–Pont, F.; Picetti, R.; Dierich,

[182086]A.; LeMeur, M.; Piazza, P. V.; Borrelli, E.: Distinct functions
of

[182087]the two isoforms of dopamine D2 receptors. Nature 408:
199–203,

[182088]2000.

[182089]

[182090]

[182091]18957.

[182092]Wang, J.; Liu, Z.-L.; Chen, B.: Association study of
dopamine

[182093]D2, D3 receptor gene polymorphisms with motor fluctua-
tions in PD. Neurology 56:

[182094]1757–1759, 2001.

[182095]

[182096]

[182097]18958.

[182098]Accili, D.; Fishburn, C. S.; Drago, J.; Steiner, H.; Lachowicz,

[182099]J. E.; Park, B.-H.; Gauda, E. B.; Lee, E. J.; Cool, M. H.; Sib-
ley,

[182100]D. R.; Gerfen, C. R.; Westphal, H.; Fuchs, S.: A targeted
mutation

[182101]of the D3 dopamine receptor gene is associated with hy-
peractivity

[182102]in mice. Proc. Nat. Acad. Sci. 93: 1945–1949, 1996.

[182103]

[182104]

[182105]18959.

[182106]Asico, L. D.; Ladines, C.; Fuchs, S.; Accili, D.; Carey, R. M.;

[182107]Semeraro, C.; Pocchiari, F.; Felder, R. A.; Eisner, G. M.;
Jose, P.

[182108]A.: Disruption of the dopamine D3 receptor gene pro-
duces renin-dependent

[182109]hypertension. J. Clin. Invest. 102: 493–498, 1998.

[182110]

[182111]

[182112]18960.

[182113]Crocq, M.–A.; Mant, R.; Asherson, P.; Williams, J.; Hode, Y.;
Mayerova,

[182114]A.; Collier, D.; Lannfelt, L.; Sokoloff, P.; Schwartz, J.–C.;
Gill,

[182115]M.; Macher, J.–P.; McGuffin, P.; Owen, M. J.: Association
between

[182116]schizophrenia and homozygosity at the dopamine D3 re-
ceptor gene. J.

[182117]Med. Genet. 29: 858–860, 1992.

[182118]

[182119]

[182120]18961.

[182121]Guillin, O.; Diaz, J.; Carroll, P.; Griffon, N.; Schwartz, J.–C.;

[182122]Sokoloff, P.: BDNF controls dopamine D3 receptor expression and triggers

[182123]behavioural sensitization. Nature 411: 86–89, 2001.

[182124]

[182125]

[182126]18962.

[182127]Ilani, T.; Ben–Shachar, D.; Strous, R. D.; Mazor, M.;
Sheinkman,

[182128]A.; Kotler, M.; Fuchs, S.: A peripheral marker for
schizophrenia:

[182129]increased levels of D3 dopamine receptor mRNA in blood
lymphocytes. Proc.

[182130]Nat. Acad. Sci. 98: 625–628, 2001.

[182131]

[182132]

[182133]18963.

[182134]Le Coniat, M.; Sokoloff, P.; Hillion, J.; Martres, M.-P.;
Giros,

[182135]B.; Pilon, C.; Schwartz, J.-C.; Berger, R.: Chromosomal lo-
calization

[182136]of the human D-3 dopamine receptor gene. Hum. Genet.
87: 618-620,

[182137]1991.

[182138]

[182139]

[182140]18964.

[182141]Nanko, S.; Sasaki, T.; Fukuda, R.; Hattori, M.; Dai, X. Y.;
Kazamatsuri,

[182142]H.; Kuwata, S.; Juji, T.; Gill, M.: A study of the association
between

[182143]schizophrenia and the dopamine D3 receptor gene. Hum.
Genet. 92:

[182144]336-338, 1993.

[182145]

[182146]

[182147]18965.

[182148]Nothen, M. M.; Cichon, S.; Propping, P.; Fimmers, R.;
Schwab, S.

[182149]G.; Wildenauer, D. B.: Excess of homozygosity at the
dopamine D3

[182150]receptor gene in schizophrenia not confirmed. (Letter) J.
Med. Genet. 30:

[182151]708–712, 1993.

[182152]

[182153]

[182154]18966.

[182155]Bevilacqua, M.; Butcher, E.; Furie, B.; Furie, B.; Gallatin, M.;

[182156]Gimbrone, M.; Harlan, J.; Kishimoto, K.; Lasky, L.; McEver,
R.; Paulson,

[182157]J.; Rosen, S.; Seed, B.; Siegelman, M.; Springer, T.; Stool-
man, L.;

[182158]Tedder, T.; Varki, A.; Wagner, D.; Weissman, I.; Zimmer-
man, G.: Selectins:

[182159]a family of adhesion receptors. (Letter) Cell 67: 233 only,
1991.

[182160]

[182161]

[182162]18967.

[182163]Bevilacqua, M. P.; Stengelin, S.; Gimbrone, M. A., Jr.; Seed, B.

[182164]: Endothelial leukocyte adhesion molecule 1: an inducible receptor

[182165]for neutrophils related to complement regulatory proteins and lectins. Science 243:

[182166]1160–1165, 1989.

[182167]

[182168]

[182169]18968.

[182170]Collins, T.; Williams, A.; Johnston, G. I.; Kim, J.; Eddy, R.;

[182171]Shows, T.; Gimbrone, M. A., Jr.; Bevilacqua, M. P.: Structure and

[182172]chromosomal location of the gene for endothelial-leukocyte adhesion

[182173]molecule 1. J. Biol. Chem. 266: 2466–2473, 1991.

[182174]

[182175]

[182176]18969.

[182177]DeLisser, H. M.; Christofidou-Solomidou, M.; Sun, J.;
Nakada, M.

[182178]T.; Sullivan, K. E.: Loss of endothelial surface expression
of E-selectin

[182179]in a patient with recurrent infections. Blood 94: 884–894,
1999.

[182180]

[182181]

[182182]18970.

[182183]Wang, N.; Chintala, S. K.; Fini, M. E.; Schuman, J. S.: Acti-
vation

[182184]of a tissue-specific stress response in the aqueous out-
flow pathway

[182185]of the eye defines the glaucoma disease phenotype. Na-
ture Med. 7:

[182186]304–309, 2001.

[182187]

[182188]

[182189]18971.

[182190]Watson, M. L.; Kingsmore, S. F.; Johnston, G. I.; Siegelman, M.

[182191]H.; Le Beau, M. M.; Lemons, R. S.; Bora, N. S.; Howard, T. A.; Weissman,

[182192]J. L.; McEver, R. P.; Seldin, M. F.: Genomic organization of the

[182193]selectin family of leukocyte adhesion molecules on human and mouse

[182194]chromosome 1. J. Exp. Med. 172: 263–272, 1990.

[182195]

[182196]

[182197]18972.

[182198]Wenzel, K.; Felix, S.; Kleber, F. X.; Brachold, R.; Menke, T.;

[182199]Schattke, S.; Schulte, K. L.; Glaser, C.; Rohde, K.; Baumann, G.;

[182200]Speer, A.: E-selectin polymorphism and atherosclerosis: an association

[182201]study. Hum. Molec. Genet. 3: 1935–1937, 1994.

[182202]

[182203]

[182204]18973.

[182205]Zheng, F.; Chevalier, J. A.; Zhang, L. Q.; Virgil, D.; Ye, S. Q.;

[182206]Kwiterovich, P. O.: An HphI polymorphism in the E-selectin gene is

[182207]associated with premature coronary artery disease. Clin. Genet. 59:

[182208]58–64, 2001.

[182209]

[182210]

[182211]18974.

[182212]Burgess, W. H.; Mehlman, T.; Marshak, D. R.; Fraser, B. A.; Maciag,

[182213]T.: Structural evidence that endothelial cell growth factor beta

[182214]is the precursor of both endothelial cell growth factor alpha and

[182215]acidic fibroblast growth factor. Proc. Nat. Acad. Sci. 83: 7216–7220,

[182216]1986.

[182217]

[182218]

[182219]18975.

[182220]Eckenstein, F. P.: Fibroblast growth factors in the nervous system. J.

[182221]Neurobiol. 25: 1467–1480, 1994.

[182222]

[182223]

[182224]18976.

[182225]Gautschi–Sova, P.; Muller, T.; Bohlen, P.: Amino acid sequence

[182226]of human acidic fibroblast growth factor. Biochem. Biophys. Res.

[182227]Commun. 140: 874–880, 1986.

[182228]

[182229]

[182230]18977.

[182231]Jaye, M.; Howk, R.; Burgess, W.; Ricca, G. A.; Chiu, I.–M.; Ravera,

- [182232]M. W.; O'Brien, S. J.; Modi, W. S.; Maciag, T.; Drohan, W. N.:
Human
[182233]endothelial cell growth factor: cloning, nucleotide se-
quence, and
[182234]chromosome localization. Science 233: 541–545, 1986.
- [182235]
- [182236]
- [182237]18978.
- [182238]Jung, J.; Zheng, M.; Goldfarb, M.; Zaret, K. S.: Initiation of
[182239]mammalian liver development from endoderm by fibrob-
last growth factors. Science 284:
[182240]1998–2003, 1999.
- [182241]
- [182242]
- [182243]18979.
- [182244]Mergia, A.; Eddy, R.; Abraham, J. A.; Fiddes, J. C.; Shows,
T.
[182245]B.: The genes for basic and acidic fibroblast growth fac-
tors are
[182246]on different human chromosomes. Biochem. Biophys. Res.

Commun. 138:

[182247]644–651, 1986.

[182248]

[182249]

[182250]18980.

[182251]Pellegrini, L.; Burke, D. F.; von Delft, F.; Mulloy, B.; Blundell,

[182252]T. L.: Crystal structure of fibroblast growth factor receptor ectodomain

[182253]bound to ligand and heparin. Nature 407: 1029–1034, 2000.

[182254]

[182255]

[182256]18981.

[182257]Plotnikov, A. N.; Hubbard, S. R.; Schlessinger, J.; Mohammadi,

[182258]M.: Crystal structures of two FGF–FGFR complexes reveal the determinants

[182259]of ligand–receptor specificity. Cell 101: 413–424, 2000.

[182260]

[182261]

[182262]18982.

[182263]Wang, W.-P.; Lehtoma, K.; Varban, M. L.; Krishnan, I.;
Chiu, I.-M.

[182264]: Cloning of the gene coding for human class 1 heparin-
binding growth

[182265]factor and its expression in fetal tissues. Molec. Cell. Biol.
9:

[182266]2387-2395, 1989.

[182267]

[182268]

[182269]18983.

[182270]Gamez, J.; Ferreiro, C.; Accarino, M. L.; Guarner, L.;
Tadesse,

[182271]S.; Marti, R. A.; Andreu, A. L.; Raguer, N.; Cervera, C.; Hi-
rano,

[182272]M.: Phenotypic variability in a Spanish family with MNGIE.
Neurology 59:

[182273]455-457, 2002.

[182274]

[182275]

[182276]18984.

[182277]Hagiwara, K.; Stenman, G.; Honda, H.; Sahlin, P.; Anders-
son, A.;

[182278]Miyazono, K.; Heldin, C. H.; Ishikawa, F.; Takaku, F.: Orga-
nization

[182279]and chromosomal localization of the human platelet-de-
rived endothelial

[182280]cell growth factor gene. Molec. Cell. Biol. 11: 2125–2132,
1991.

[182281]

[182282]

[182283]18985.

[182284]Nishino, I.; Spinazzola, A.; Hirano, M.: Thymidine phos-
phorylase

[182285]gene mutations in MNGIE, a human mitochondrial disor-
der. Science 283:

[182286]689–692, 1999.

[182287]

[182288]

[182289]18986.

[182290]Stenman, G.; Sahlin, P.; Dumanski, J. P.; Hagiwara, K.;

Ishikawa,

[182291]F.; Miyazono, K.; Collins, V. P.; Heldin, C.-H.: Regional localization

[182292]of the human platelet-derived endothelial cell growth factor (ECGF1)

[182293]gene to chromosome 22q13. Cytogenet. Cell Genet. 59: 22-23, 1992.

[182294]

[182295]

[182296]18987.

[182297]Stenman, G.; Sahlin, P.; Hagiwara, K.; Dumanski, J.;

Collins, V.;

[182298]Heldin, C.-H.: Mapping of the human platelet-derived endothelial

[182299]cell growth factor (PD-ECGF) gene to chromosome 22q13. (Abstract) Cytogenet.

[182300]Cell Genet. 58: 2051 only, 1991.

[182301]

[182302]

[182303]18988.

[182304]Kayes–Wandover, K.; White, P. C.: Steroidogenic enzyme
gene expression

[182305]in the human heart. J. Clin. Endocr. Metab. 85:
2519–2525, 2000.

[182306]

[182307]

[182308]18989.

[182309]Carrero–Valenzuela, R. D.; Quan, F.; Lightowlers, R.; Ken-
naway,

[182310]N. G.; Litt, M.; Forte, M.: Human cytochrome c oxidase
subunit VIb:

[182311]characterization and mapping of a multigene family. Gene
102: 229–236,

[182312]1991.

[182313]

[182314]

[182315]18990.

[182316]Taanman, J.–W.; Schrage, C.; Ponne, N.; Bolhuis, P.; de
Vries,

[182317]H.; Agsteribbe, E.: Nucleotide sequence of cDNA encoding subunit

[182318]VIb of human cytochrome c oxidase. Nucleic Acids Res. 17: 1766

[182319]only, 1989.

[182320]

[182321]

[182322]18991.

[182323]Taanman, J.-W.; van der Veen, A. Y.; Schrage, C.; de Vries, H.;

[182324]Buys, C. H. C. M.: Assignment of the gene coding for human cytochrome

[182325]c oxidase subunit VIb to chromosome 19, band q13.1, by fluorescence

[182326]in situ hybridisation. Hum. Genet. 87: 325-327, 1991.

[182327]

[182328]

[182329]18992.

[182330]Sinke, R. J.; Tanigami, A.; Nakamura, Y.; Geurts van Kessel, A.

[182331]: Reverse mapping of the gene encoding the human fos-related antigen-1

[182332](fra-1) within chromosome band 11q13. Genomics 18: 165 only, 1993.

[182333]

[182334]

[182335]18993.

[182336]Cox, D. R.; Epstein, L. B.; Epstein, C. J.: Genes coding for sensitivity

[182337]to interferon (IfRec) and soluble superoxide dismutase (SOD-1) are

[182338]linked in mouse and man and map to mouse chromosome 16. Proc. Nat.

[182339]Acad. Sci. 77: 2168-2172, 1980.

[182340]

[182341]

[182342]18994.

[182343]De Clercq, E.; Edy, V. G.; Cassiman, J.-J.: Chromosome 21 does

[182344]not code for an interferon receptor. Nature 264: 249-251, 1976.

[182345]

[182346]

[182347]18995.

[182348]Epstein, L. B.; Epstein, C. J.: Localization of the gene AVG
for

[182349]the antiviral expression of immune and classical interferon
to the

[182350]distal portion of the long arm of chromosome 21. J. Infect.
Dis. 133

[182351](suppl.): A56–A62, 1976.

[182352]

[182353]

[182354]18996.

[182355]Faltynek, C. R.; Branca, A. A.; McCandless, S.; Baglioni, C.:

[182356]Characterization of an interferon receptor on human lym-
phoblastoid

[182357]cells. Proc. Nat. Acad. Sci. 80: 3269–3273, 1983.

[182358]

[182359]

[182360]18997.

[182361]Fournier, A.; Zhang, Z. Q.; Tan, Y. H.: Human beta:alpha
but not

[182362]gamma interferon binding site is a product of the chro-
mosome 21 interferon

[182363]action gene. Somat. Cell Molec. Genet. 11: 291–295,
1985.

[182364]

[182365]

[182366]18998.

[182367]Lutfalla, G.; Gardiner, K.; Proudhon, D.; Vielh, E.; Uze, G.:

[182368]The structure of the human interferon alpha/beta receptor
gene. J.

[182369]Biol. Chem. 267: 2802–2809, 1992.

[182370]

[182371]

[182372]18999.

[182373]Lutfalla, G.; Roeckel, N.; Mogensen, K. E.; Mattei, M. G.;
Uze,

[182374]G.: Assignment of the human interferon–alpha receptor

gene to chromosome

[182375]21q22.1 by in situ hybridization. J. Interferon Res. 10:
515–517,

[182376]1990.

[182377]

[182378]

[182379]19000.

[182380]Maroun, L. E.: Interferon action and chromosome 21 tri-
somy. (Letter) J.

[182381]Theor. Biol. 86: 603–606, 1980.

[182382]

[182383]

[182384]19001.

[182385]Novick, D.; Cohen, B.; Rubinstein, M.: The human inter-
feron alpha/beta

[182386]receptor: characterization and molecular cloning. Cell 77:
391–400,

[182387]1994.

[182388]

[182389]

[182390]19002.

[182391]Raziuddin, A.; Sarkar, F. H.; Dutkowski, R.; Shulman, L.;
Ruddle,

[182392]F. H.; Gupta, S. L.: Receptors for human alpha and beta in-
terferon

[182393]but not for gamma interferon are specified by human
chromosome 21. Proc.

[182394]Nat. Acad. Sci. 81: 5504–5508, 1984.

[182395]

[182396]

[182397]19003.

[182398]Revel, M.; Bash, D.; Ruddle, F. H.: Antibodies to a cell-
surface

[182399]component coded by human chromosome 21 inhibit ac-
tion of interferon. Nature 260:

[182400]139–141, 1976.

[182401]

[182402]

[182403]19004.

[182404]Topham, M. K.; Prescott, S. M.: Mammalian diacylglycerol kinases,

[182405]a family of lipid kinases with signaling functions. J. Biol. Chem. 274:

[182406]11447–11450, 1999.

[182407]

[182408]

[182409]19005.

[182410]Calabrese, G.; Crescenzi, C.; Morizio, E.; Palka, G.; Guerra, E.;

[182411]Alberti, S.: Assignment of TACSTD1 (alias TROP1, M4S1) to human chromosome

[182412]2p21 and refinement of mapping of TACSTD2 (alias TROP2, M1S1) to human

[182413]chromosome 1p32 by in situ hybridization. Cytogenet. Cell Genet. 92:

[182414]164–165, 2001.

[182415]

[182416]

[182417]19006.

[182418]Fornaro, M.; Dell'Arciprete, R.; Stella, M.; Bucci, C.; Nutini,
[182419]M.; Capri, M. G.; Alberti, S.: Cloning of the gene encoding
Trop-2,

[182420]a cell-surface glycoprotein expressed by human carcinomas. Int. J.

[182421]Cancer 62: 610-618, 1995.

[182422]

[182423]

[182424]19007.

[182425]Linnenbach, A. J.; Seng, B. A.; Wu, S.; Robbins, S.; Scollon, M.;

[182426]Pyrce, J. J.; Druck, T.; Huebner, K.: Retroposition in a family of

[182427]carcinoma-associated antigen genes. Molec. Cell Biol. 13: 1507-1515,

[182428]1993.

[182429]

[182430]

[182431]19008.

[182432]Linnenbach, A. J.; Wojciorowski, J.; Wu, S.; Pyrc, J. J.; Ross,

[182433]A. H.; Dietzschold, B.; Speicher, D.; Koprowski, H.: Sequence investigation

[182434]of the major gastrointestinal tumor-associated antigen

gene family,

[182435]GA733. Proc. Nat. Acad. Sci. 86: 27–31, 1989.

[182436]

[182437]

[182438]19009.

[182439]Ren, Z.; Lin, P.-Y.; Klintworth, G. K.; Iwata, F.; Munier, F.
L.;

[182440]Schorderet, D. F.; El Matri, L.; Theendakara, V.; Basti, S.;
Reddy,

[182441]M.; Hejtmancik, J. F.: Allelic and locus heterogeneity in au-
tosomal

[182442]recessive gelatinous drop-like corneal dystrophy. Hum.
Genet. 110:

[182443]568–577, 2002.

[182444]

[182445]

[182446]19010.

[182447]TsujiKawa, M.; Kurahashi, H.; Tanaka, T.; Nishida, K.; Shi-
momura,

[182448]Y.; Tano, Y.; Nakamura, Y.: Identification of the gene re-

sponsible

[182449]for gelatinous drop-like corneal dystrophy. Nature Genet.
21: 420–423,

[182450]1999.

[182451]

[182452]

[182453]19011.

[182454]Willecke, K.; Jungbluth, S.; Dahl, E.; Hennemann, H.;
Heynkes,

[182455]R.; Grzeschik, K.–H.: Six genes of the human connexin
gene family

[182456]coding for gap junctional proteins are assigned to four
different

[182457]human chromosomes. Europ. J. Cell Biol. 53: 275–280,
1990.

[182458]

[182459]

[182460]19012.

[182461]Reed, K. E.; Westphale, E. M.; Larson, D. M.; Wang, H.–Z.;
Veenstra,

[182462]R. D.; Beyer, E. C.: Molecular cloning and functional ex–

pression

[182463]of human connexin37, an endothelial cell gap junction protein. J.

[182464]Clin. Invest. 91: 997–1004, 1993.

[182465]

[182466]

[182467]19013.

[182468]Van Camp, G.; Coucke, P.; Speleman, F.; Van Roy, N.; Beyer, E.

[182469]C.; Oostra, B. A.; Willems, P. J.: The gene for human gap junction

[182470]protein connexin37 (GJA4) maps to chromosome 1p35.1, in the vicinity

[182471]of D1S195. Genomics 30: 402–403, 1995.

[182472]

[182473]

[182474]19014.

[182475]Gelb, B. D.; Zhang, J.; Cotter, P. D.; Gershin, I. F.; Desnick,

[182476]R. J.: Physical mapping of the human connexin 40 (GJA5) flavin-containing

[182477]monooxygenase 5, and natriuretic peptide receptor A
genes on 1q21. Genomics 39:

[182478]409–411, 1997.

[182479]

[182480]

[182481]19015.

[182482]Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Cardiac myocytes

[182483]express multiple gap junction proteins. Circ. Res. 70:
438–444,

[182484]1992.

[182485]

[182486]

[182487]19016.

[182488]Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Molecular cloning

[182489]of two human cardiac gap junction proteins, connexin40
and connexin45. J.

[182490]Molec. Cell. Cardiol. 26: 861–868, 1994.

[182491]

[182492]

[182493]19017.

[182494]Oviedo-Orta, E.; Hoy, T.; Evans, W. H.: Intercellular communication

[182495]in the immune system: differential expression of connexin40 and 43,

[182496]and perturbation of gap junction channel functions in peripheral blood

[182497]and tonsil human lymphocyte subpopulations. Immunology 99: 578-590,

[182498]2000.

[182499]

[182500]

[182501]19018.

[182502]Seul, K. H.; Tadros, P. N.; Beyer, E. C.: Mouse connexin40: gene

[182503]structure and promoter analysis. Genomics 46: 120-126, 1997.

[182504]

[182505]

[182506]19019.

[182507]Britz–Cunningham, S. H.; Shah, M. M.; Zuppan, C. W.;
Fletcher,

[182508]W. H.: Mutations of the connexin43 gap–junction gene in
patients

[182509]with heart malformations and defects of laterality. New
Eng. J. Med. 332:

[182510]1323–1329, 1995.

[182511]

[182512]

[182513]19020.

[182514]Brueckner, M.; D'Eustachio, P.; Horwich, A. L.: Linkage
mapping

[182515]of a mouse gene, iv, that controls left–right asymmetry of
the heart

[182516]and viscera. Proc. Nat. Acad. Sci. 86: 5035–5038, 1989.

[182517]

[182518]

[182519]19021.

[182520]Burdine, R. D.; Schier, A. F.: Conserved and divergent
mechanisms

[182521]in left-right axis formation. Genes Dev. 14: 763–776,
2000.

[182522]

[182523]

[182524]19022.

[182525]Casey, B.; Ballabio, A.: Connexin43 mutations in sporadic
and

[182526]familial defects of laterality. (Letter) New Eng. J. Med. 333:
941,

[182527]1995.

[182528]

[182529]

[182530]19023.

[182531]Corcos, I. A.; Meese, E. U.; Loch-Caruso, R.: Human con-
nexin 43

[182532]gene locus, GJA1, sublocalized to band 6q21–q23.2. Cy-
togenet. Cell

[182533]Genet. 64: 31–32, 1993.

[182534]

[182535]

[182536]19024.

[182537]Debrus, S.; Tuffery, S.; Matsuoka, R.; Galal, O.; Sarda, P.;
Sauer,

[182538]U.; Bozio, A.; Tanman, B.; Toutain, A.; Claustres, M.; Le
Paslier,

[182539]D.; Bouvagnet, P.: Lack of evidence for connexin 43 gene
mutations

[182540]in human autosomal recessive lateralization defects. J.
Molec. Cell.

[182541]Cardiol. 29: 1423–1431, 1997.

[182542]

[182543]

[182544]19025.

[182545]Wijnen, J. T.; Oldenburg, M.; Bloemendal, H.; Meera Khan,
P.:

[182546]GS(gamma-S)–crystallin (CRYGS) assignment to chromo-
some 3. (Abstract) Cytogenet.

[182547]Cell Genet. 51: 1108 only, 1989.

[182548]

[182549]

[182550]19026.

[182551]Hai, T.; Liu, F.; Coukos, W. J.; Green, M. R.: Transcription factor

[182552]ATF cDNA clones: an extensive family of leucine zipper proteins able

[182553]to selectively form DNA-binding heterodimers. Genes Dev. 3: 2083–2090,

[182554]1989.

[182555]

[182556]

[182557]19027.

[182558]Zucman, J.; Delattre, O.; Desmaze, C.; Epstein, A. L.; Stenman,

[182559]G.; Speleman, F.; Fletchers, C. D. M.; Aurias, A.; Thomas, G.: EWS

[182560]and ATF-1 gene fusion induced by t(12;22) translocation in malignant

[182561]melanoma of soft parts. Nature Genet. 4: 341–345, 1993.

[182562]

[182563]

[182564]19028.

[182565]Barco, A.; Alarcon, J. M.; Kandel, E. R.: Expression of constitutively

[182566]active CREB protein facilitates the late phase of long-term potentiation

[182567]by enhancing synaptic capture. Cell 108: 689–703, 2002.

[182568]

[182569]

[182570]19029.

[182571]Barton, C. H.; Ajioka, J. W.; Roach, T. I. A.; Blackwell, J. M.

[182572]: Mapping Creb-1 to chromosome 1 in the mouse. Genomics 14: 790–792,

[182573]1992.

[182574]

[182575]

[182576]19030.

[182577]Barton, K.; Muthusamy, N.; Chanyangam, M.; Fischer, C.; Clendenin,

[182578]C.; Leiden, J. M.: Defective thymocyte proliferation and IL–

2 production

[182579]in transgenic mice expressing a dominant-negative form
of CREB. Nature 379:

[182580]81–85, 1996.

[182581]

[182582]

[182583]19031.

[182584]Carlezon, W. A., Jr.; Thome, J; Olson, V. G.; Lane-Ladd, S.
B.;

[182585]Brodkin, E. S.; Hiroi, N.; Duman, R. S.; Neve, R. L.; Nestler,
E.

[182586]J.: Regulation of cocaine reward by CREB. Science 282:
2272–2275,

[182587]1998.

[182588]

[182589]

[182590]19032.

[182591]Cole, T. J.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.;
Schutz,

[182592]G.; Ruppert, S.: The mouse CREB (cAMP responsive ele–

ment binding

[182593]protein) gene: structure, promoter analysis, and chromosomal localization. Genomics 13:

[182594]974–982, 1992.

[182595]

[182596]

[182597]19033.

[182598]Fentzke, R. C.; Korcarz, C. E.; Lang, R. M.; Lin, H.; Leiden, J.

[182599]M.: Dilated cardiomyopathy in transgenic mice expressing a dominant-negative

[182600]CREB transcription factor in the heart. J. Clin. Invest. 101: 2415–2426,

[182601]1998.

[182602]

[182603]

[182604]19034.

[182605]Herzig, S.; Long, F.; Jhala, U. S.; Hedrick, S.; Quinn, R.; Bauer,

[182606]A.; Rudolph, D.; Schutz, G.; Yoon, C.; Puigserver, P.;

Spiegelman,

[182607]B.; Montminy, M.: CREB regulates hepatic gluconeogenesis through

[182608]the coactivator PGC-1. Nature 413: 179-183, 2001.

[182609]

[182610]

[182611]19035.

[182612]Benson, A. M.; Hunkeler, M. J.; Talalay, P.: Increase of NAD(P)H:quinone

[182613]reductase by dietary antioxidants: possible role in protection against

[182614]carcinogenesis and toxicity. Proc. Nat. Acad. Sci. 77: 5216-5220,

[182615]1980.

[182616]

[182617]

[182618]19036.

[182619]Chen, L. Z.; Harris, P. C.; Apostolou, S.; Baker, E.; Holman, K.;

[182620]Lane, S. A.; Nancarrow, J. K.; Whitmore, S. A.; Stallings, R.

L.;

[182621]Hildebrand, C. E.; Richards, R. I.; Sutherland, G. R.; Callen, D.

[182622]F.: A refined physical map of the long arm of human chromosome 16. Genomics 10:

[182623]308–312, 1991.

[182624]

[182625]

[182626]19037.

[182627]Edwards, Y. H.; Hopkinson, D. A.; Carritt, B.: A genetic characterization

[182628]of the human diaphorase–4 deficiency. Ann. Hum. Genet. 47: 97–105,

[182629]1983.

[182630]

[182631]

[182632]19038.

[182633]Grzeschik, K.–H.: Assignment of a structural gene for a fourth

[182634]human diaphorase (DIA–4) to chromosome 16 in man–

mouse somatic cell

[182635]hybrids. Hum. Genet. 53: 189–193, 1980.

[182636]

[182637]

[182638]19039.

[182639]Jaiswal, A. K.; McBride, O. W.; Adesnik, M.; Nebert, D. W.:
Human

[182640]dioxin-inducible cytosolic NAD(P)H:menadione oxidore-
ductase: cDNA

[182641]sequence and localization of gene to chromosome 16. J.
Biol. Chem. 263:

[182642]13572–13578, 1988.

[182643]

[182644]

[182645]19040.

[182646]Kelsey, K. T.; Ross, D.; Traver, R. D.; Christiani, D. C.; Zuo,

[182647]Z. F.; Spitz, M. R.; Wang, M.; Xu, X.; Lee, B. K.; Schwartz, B.
S.;

[182648]Wiencke, J. K.: Ethnic variation in the prevalence of a com-
mon NAD(P)H

[182649]quinone oxidoreductase polymorphism and its implications for anti-cancer

[182650]chemotherapy. Brit. J. Cancer 76: 852–854, 1997.

[182651]

[182652]

[182653]19041.

[182654]Lavinha, J.; Morrison, N.; Glasgow, L.; Ferguson-Smith, M.
A.:

[182655]Further evidence for the regional localization of human
APRT and DIA4

[182656]on chromosome 16. (Abstract) Cytogenet. Cell Genet. 37:
517 only,

[182657]1984.

[182658]

[182659]

[182660]19042.

[182661]Moran, J. L.; Siegel, D.; Ross, D.: A potential mechanism
underlying

[182662]the increased susceptibility of individuals with a polymor-
phism in

[182663]NAD(P)H:quinone oxidoreductase 1 (NQO1) to benzene toxicity. Proc.

[182664]Nat. Acad. Sci. 96: 8150–8155, 1999.

[182665]

[182666]

[182667]19043.

[182668]Povey, S.; Wilson, D.; Edwards, Y. H.: Assignment of a human diaphorase

[182669](DIA-4) to chromosome 16. Ann. Hum. Genet. 43: 349–353, 1980.

[182670]

[182671]

[182672]19044.

[182673]Radjendirane, V.; Joseph, P.; Lee, Y.-H.; Kimura, S.; Klein-Szanto,

[182674]A. J. P.; Gonzalez, F. J.; Jaiswal, A. K.: Disruption of the DT diaphorase

[182675](NQO1) gene in mice leads to increased menadione toxicity. J. Biol.

[182676]Chem. 273: 7382–7389, 1998.

[182677]

[182678]

[182679]19045.

[182680]Rothman, N.; Smith, M. T.; Hayes, R. B.; Traver, R. D.;
Hoener,

[182681]B.; Campleman, S.; Li, G. L.; Dosemeci, M.; Linet, M.;
Zhang, L.;

[182682]Xi, L.; Wacholder, S.; Lu, W.; Meyer, K. B.; Titenko-Holland,
N.;

[182683]Stewart, J. T.; Yin, S.; Ross, D.: Benzene poisoning, a risk
factor

[182684]for hematological malignancy, is associated with the
NQO1 609C-T mutation

[182685]and rapid fractional excretion of chlorzoxazone. Cancer
Res. 57:

[182686]2839-2842, 1997.

[182687]

[182688]

[182689]19046.

[182690]Smith, M. T.: Benzene, NQO1, and genetic susceptibility to
cancer.

[182691](Commentary) Proc. Nat. Acad. Sci. 96: 7624–7626, 1999.

[182692]

[182693]

[182694]19047.

[182695]Traver, R. D.; Horikoshi, T.; Danenberg, K. D.; Stadlbauer, T.

[182696]H.; Danenberg, P. V.; Ross, D.; Gibson, N. W.:

NAD(P)H:quinone oxidoreductase

[182697]gene expression in human colon carcinoma cells: characterization of

[182698]a mutation which modulates DT-diaphorase activity and mitomycin sensitivity. Cancer

[182699]Res. 52: 797–802, 1992.

[182700]

[182701]

[182702]19048.

[182703]Traver, R. D.; Siegel, D.; Beall, H. D.; Phillips, R. M.; Gibson, son,

[182704]N. W.; Franklin, W. A.; Ross, D.: Characterization of a polymorphism

[182705]in NAD(P)H: quinone oxidoreductase (DT-diaphorase).

Brit. J. Cancer 75:

[182706]69–75, 1997.

[182707]

[182708]

[182709]19049.

[182710]Zhang, L.; Rothman, N.; Wang, Y.; Hayes, R. B.; Li, G.;

Dosemeci,

[182711]M.; Yin, S.; Kolachana, P.; Titenko–Holland, N.; Smith, M.

T.: Increased

[182712]aneusomy and long arm deletion of chromosomes 5 and 7
in the lymphocytes

[182713]of Chinese workers exposed to benzene. Carcinogenesis
19: 1955–1961,

[182714]1998.

[182715]

[182716]

[182717]19050.

[182718]Zhang, L.; Wang, Y.; Shang, N.; Smith, M. T.: Benzene
metabolites

[182719]induce the loss of long arm deletion of chromosomes 5
and 7 in human

[182720]lymphocytes. Leukemia Res. 22: 105–113, 1998.

[182721]

[182722]

[182723]19051.

[182724]Roychoudhury, A. K.; Nei, M.: Human Polymorphic Genes:
World Distribution.

[182725]New York: Oxford Univ. Press (pub.) 1988.

[182726]

[182727]

[182728]19052.

[182729]Anholt, R. R. H.; Pederson, P. L.; Desouza, E. B.; Synder, S.
H.

[182730]: The peripheral-type benzodiazepine receptor: localiza-
tion to the

[182731]mitochondrial outer membrane. J. Biol. Chem. 261:
576–583, 1986.

[182732]

[182733]

[182734]19053.

[182735]DeBernardi, M. A.; Crowe, R. R.; Mocchetti, I.; Shows, T. B.;
Eddy,

[182736]R. L.; Costa, E.: Chromosomal localization of the human
diazepam

[182737]binding inhibitor gene. Proc. Nat. Acad. Sci. 85:
6561–6565, 1988.

[182738]

[182739]

[182740]19054.

[182741]Campanelli, J. T.; Hoch, W.; Rupp, F.; Kreiner, T.; Scheller,
R.

[182742]H.: Agrin mediates cell contact–induced acetylcholine re–
ceptor clustering. Cell 67:

[182743]909–916, 1991.

[182744]

[182745]

[182746]19055.

[182747]DeChiara, T. M.; Bowen, D. C.; Valenzuela, D. M.; Sim–
mons, M. V.;

[182748] Poueymirou, W. T.; Thomas, S.; Kinetz, E.; Compton, D. L.;
Rojas,
[182749] E.; Park, J. S.; Smith, C.; DiStefano, P. S.; Glass, D. J.; Bur-
den,
[182750] S. J.; Yancopoulos, G. D.: The receptor tyrosine kinase
MuSK is required
[182751] for neuromuscular junction formation in vivo. Cell 85:
501–512,
[182752] 1996.
[182753]
[182754]
[182755] 19056.
[182756] Anonymous: Freely associating. (Editorial) Nature Genet.
22:
[182757] 1–2, 1999.
[182758]
[182759]
[182760] 19057.
[182761] Bell, G. I.; Rall, L. B.; Sanchez–Pescador, R.; Merryweather,
J.

[182762]P.; Scott, J.; Eddy, R. L.; Shows, T. B.: Human alpha-
2-macroglobulin

[182763]gene is located on chromosome 12. Somat. Cell Molec.
Genet. 11:

[182764]285-289, 1985.

[182765]

[182766]

[182767]19058.

[182768]Bergqvist, D.; Nilsson, I. M.: Hereditary alpha-
2-macroglobulin

[182769]deficiency. Scand. J. Haemat. 23: 433-436, 1979.

[182770]

[182771]

[182772]19059.

[182773]Blacker, D.; Crystal, A. S.; Wilcox, M. A.; Laird, N. M.;
Tanzi,

[182774]R. E.: Reply to Rudrasingham et al. Nature Genet. 22:
21-22, 1999.

[182775]

[182776]

[182777]19060.

[182778]Blacker, D.; Wilcox, M. A.; Laird, N. M.; Rodes, L.; Horvath, S.

[182779]M.; Go, R. C. P.; Perry, R.; Watson, B., Jr.; Bassett, S. S.; McInnis,

[182780]M. G.; Albert, M. S.; Hyman, B. T.; Tanzi, R. E.: Alpha-2 macroglobulin

[182781]is genetically associated with Alzheimer disease. Nature Genet. 19:

[182782]357-360, 1998.

[182783]

[182784]

[182785]19061.

[182786]David, F.; Kan, C. C.; Lucotte, G.: Two Taq I RFLPs for human

[182787]alpha-2 macroglobulin (alpha-2M) using a full length cDNA probe. Nucleic

[182788]Acids Res. 15: 374 only, 1987.

[182789]

[182790]

[182791]19062.

[182792]Chan, S. Y.; Empig, C. J.; Welte, F. J.; Speck, R. F.;

Schmaljohn,

[182793]A.; Kreisberg, J. F.; Goldsmith, M. A.: Folate receptor- α is

[182794]a cofactor for cellular entry by Marburg and Ebola viruses.

Cell 106:

[182795]117–126, 2001.

[182796]

[182797]

[182798]19063.

[182799]Czeizel, A. E.; Dudas, I.: Prevention of the first occurrence

[182800]of neural-tube defects by periconceptional vitamin supplementation. New

[182801]Eng. J. Med. 327: 1832–1835, 1992.

[182802]

[182803]

[182804]19064.

[182805]De Marco, P.; Moroni, A.; Merello, E.; de Franchis, R.; An-

dreussi,

[182806]L.; Finnell, R. H.; Barber, R. C.; Cama, A.; Capra, V.: Folate pathway

[182807]gene alterations in patients with neural tube defects. Am. J. Med.

[182808]Genet. 95: 216–223, 2000.

[182809]

[182810]

[182811]19065.

[182812]Elwood, P. C.: Molecular cloning and characterization of the human

[182813]folate-binding protein cDNA from placenta and malignant tissue culture

[182814](KB) cells. J. Biol. Chem. 264: 14893–14901, 1989.

[182815]

[182816]

[182817]19066.

[182818]Elwood, P. C.; Nachmanoff, K.; Saikawa, Y.; Page, S. T.; Pacheco,

[182819]P.; Roberts, S.; Chung, K.–N.: The divergent 5–prime ter–

mini of the

[182820]alpha human folate receptor (hFR) mRNAs originate from
two tissue-specific

[182821]promoters and alternative splicing: characterization of the
alpha

[182822]hFR gene structure. Biochemistry 36: 1467–1478, 1997.

[182823]

[182824]

[182825]19067.

[182826]Boman, B. M.; Wildrick, D. M.; Alfaro, S. R.: Chromosome
18 allele

[182827]loss at the D18S6 locus in human colorectal carcinomas.
Biochem.

[182828]Biophys. Res. Commun. 155: 463–469, 1988.

[182829]

[182830]

[182831]19068.

[182832]Chan, S. S.–Y.; Zheng, H.; Su, M.–W.; Wilk, R.; Killeen, M.
T.;

[182833]Hedgecock, E. M.; Culotti, J. G.: UNC–40, a C. elegans ho–

molog of

[182834]the DCC (deleted in colorectal cancer), is required in
motile cells

[182835]responding to UNC-6 netrin cues. Cell 87: 187-195,
1996.

[182836]

[182837]

[182838]19069.

[182839]Sarmanova, J.; Benesova, K.; Gut, I.; Nedelcheva-Kris-
tensen, V.;

[182840]Tynkova, L.; Soucek, P.: Genetic polymorphisms of bio-
transformation

[182841]enzymes in patients with Hodgkin's and non-Hodgkin's
lymphomas. Hum.

[182842]Molec. Genet. 10: 1265-1273, 2001.

[182843]

[182844]

[182845]19070.

[182846]Nadeau, J. H.; Eicher, E. M.: Conserved linkage of soluble
aconitase

[182847]and galactose-1-phosphate uridyl transferase in mouse

and man: assignment

[182848]of these genes to mouse chromosome 4. Cytogenet. Cell
Genet. 34:

[182849]271–281, 1982.

[182850]

[182851]

[182852]19071.

[182853]Albrecht, F. E.; Drago, J.; Felder, R. A.; Printz, M. P.; Eisner,

[182854]G. M.; Robillard, J. E.; Sibley, D. R.; Westphal, H. J.; Jose, P.

[182855]A.: Role of the D–1A dopamine receptor in the pathogene-
sis of genetic

[182856]hypertension. J. Clin. Invest. 97: 2283–2288, 1996.

[182857]

[182858]

[182859]19072.

[182860]Bermak, J. C.; Li, M.; Bullock, C.; Zhou, Q.–Y.: Regulation
of

[182861]transport of the dopamine D1 receptor by a new mem-
brane-associated

[182862]ER protein. Nature Cell Biol. 3: 492–498, 2001.

[182863]

[182864]

[182865]19073.

[182866]Boulton, J.; Lewis, M. S.; Wainscoat, J. S.: Physical linkage
age

[182867]of glucocorticoid receptor (GR) and dopamine D1 receptor (DRD1) on

[182868]the long arm of chromosome 5.(Abstract) Cytogenet. Cell
Genet. 58:

[182869]1894 only, 1991.

[182870]

[182871]

[182872]19074.

[182873]Castner, S. A.; Williams, G. V.; Goldman-Rakic, P. S.: Re-
versal

[182874]of antipsychotic-induced working memory deficits by
short-term dopamine

[182875]D1 receptor stimulation. Science 287: 2020–2022, 2000.

[182876]

[182877]

[182878]19075.

[182879]Dearry, A.; Gingrich, J. A.; Falardeau, P.; Fremeau, R. T.,
Jr.;

[182880]Bates, M. D.; Caron, M. G.: Molecular cloning and expres-
sion of the

[182881]gene for a human D(1) dopamine receptor. Nature 347:
72-76, 1990.

[182882]

[182883]

[182884]19076.

[182885]Grandy, D. K.; Zhou, Q.-Y.; Allen, L.; Litt, R.; Magenis, R.
E.;

[182886]Civelli, O.; Litt, M.: A human D(1) dopamine receptor gene
is located

[182887]on chromosome 5 at q35.1 and identifies an EcoRI RFLP.
Am. J. Hum.

[182888]Genet. 47: 828-834, 1990.

[182889]

[182890]

[182891]19077.

[182892]Krushkal, J.; Xiong, M.; Ferrell, R.; Sing, C. F.; Turner, S. T.;

[182893]Boerwinkle, E.: Linkage and association of adrenergic and
dopamine

[182894]receptor genes in the distal portion of the long arm of
chromosome

[182895]5 with systolic blood pressure variation. Hum. Molec.
Genet. 7:

[182896]1379–1383, 1998.

[182897]

[182898]

[182899]19078.

[182900]Mayerhofer, A.; Hemmings, H. C., Jr.; Snyder, G. L.; Green-
gard,

[182901]P.; Boddien, S.; Berg, U.; Brucker, C.: Functional
dopamine-1 receptors

[182902]and DARPP-32 are expressed in human ovary and granu-
losa luteal cells

[182903]in vitro. J. Clin. Endocr. Metab. 84: 257–264, 1999.

[182904]

[182905]

[182906]19079.

[182907]Sunahara, R. K.; Niznik, H. B.; Weiner, D. M.; Stormann, T.
M.;

[182908]Brann, M. R.; Kennedy, J. L.; Gelernter, J. E.; Rozmahel, R.;
Yang,

[182909]Y.; Israel, Y.; Seeman, P.; O'Dowd, B. F.: Human dopamine
D1 receptor

[182910]encoded by an intronless gene on chromosome 5. Nature
347: 80–83,

[182911]1990.

[182912]

[182913]

[182914]19080.

[182915]Wilkie, T. M.; Chen, Y.; Gilbert, D. J.; Moore, K. J.; Yu, L.;

[182916]Simon, M. I.; Copeland, N. G.; Jenkins, N. A.: Identification,
chromosomal

[182917]location, and genome organization of mammalian G–
protein–coupled receptors. Genomics 18:

[182918]175–184, 1993.

[182919]

[182920]

[182921]19081.

[182922]Xu, M.; Moratalla, R.; Gold, L. H.; Hiroi, N.; Koob, G. F.;
Graybiel,

[182923]A. M.; Tonegawa, S.: Dopamine D1 receptor mutant mice
are deficient

[182924]in striatal expression of dynorphin and in dopamine-mediated behavioral

[182925]responses. Cell 79: 729–742, 1994.

[182926]

[182927]

[182928]19082.

[182929]Zhou, Q.-Y.; Grandy, D. K.; Thambi, L.; Kushner, J. A.; Van
Tol,

[182930]H. H. M.; Cone, R.; Pribnow, D.; Salon, J.; Bunzow, J. R.;
Civelli,

[182931]O.: Cloning and expression of human and rat D(1)
dopamine receptors. Nature 347:

[182932]76–80, 1990.

[182933]

[182934]

[182935]19083.

[182936]Balk, J.-H.; Picetti, R.; Salardi, A.; Thirlet, G.; Dierich, A.;

[182937]Depaulls, A.; Le Meur, M.; Borrelli, E.: Parkinsonian-like
locomotor

[182938]impairment in mice lacking dopamine D2 receptors. Na-
ture 377: 424-428,

[182939]1995.

[182940]

[182941]

[182942]19084.

[182943]Basu, S.; Nagy, J. A.; Pal, S.; Vasile, E.; Eckelhoefer, I. A.;

[182944]Bliss, V. S.; Manseau, E. J.; Dasgupta, P. S.; Dvorak, H. F.;
Mukhopadhyay,

[182945]D.: The neurotransmitter dopamine inhibits angiogenesis
induced by

[182946]vascular permeability factor/vascular endothelial growth
factor. Nature

[182947]Med. 7: 569-574, 2001.

[182948]

[182949]

[182950]19085.

[182951]Blum, K.; Noble, E. P.; Sheridan, P. J.; Montgomery, A.;
Ritchie,

[182952]T.; Jagadeeswaran, P.; Nogami, H.; Briggs, A. H.; Cohn, J.
B.: Allelic

[182953]association of human dopamine D(2) receptor gene in al-
coholism. J.A.M.A. 263:

[182954]2055–2060, 1990.

[182955]

[182956]

[182957]19086.

[182958]Bolos, A. M.; Dean, M.; Lucas–Derse, S.; Ramsburg, M.;
Brown, G.

[182959]L.; Goldman, D.: Population and pedigree studies reveal a
lack of

[182960]association between the dopamine D(2) receptor gene and
alcoholism. J.A.M.A. 264:

[182961]3156–3160, 1990.

[182962]

[182963]

[182964]19087.

[182965]Bunzow, J. R.; Van Tol, H. H.; Grandy, D. K.; Albert, P.; Sallan,
lon,

[182966]J.; Christie, M.; Machida, C. A.; Neve, K. A.; Civelli, O.:
Cloning

[182967]and expression of a rat D2 dopamine receptor cDNA. Na-
ture 336: 783–787,

[182968]1988.

[182969]

[182970]

[182971]19088.

[182972]Dal Toso, R.; Sommer, B.; Ewert, M.; Herb, A.; Pritchett, D.
B.;

[182973]Bach, A.; Shivers, B. D.; Seeburg, P. H.: The dopamine D(2)
receptor:

[182974]two molecular forms generated by alternative splicing.
EMBO J. 8:

[182975]4025–4034, 1989.

[182976]

[182977]

[182978]19089.

[182979]Seeman, P.; Guan, H.-C.; Van Tol, H. H. M.: Dopamine D4
receptors

[182980]elevated in schizophrenia. Nature 365: 441–445, 1993.

[182981]

[182982]

[182983]19090.

[182984]Seeman, P.; Ulpian, C.; Chouinard, G.; Van Tol, H. H. M.;
Dwosh,

[182985]H.; Lieberman, J. A.; Siminovitch, K.; Liu, I. S. C.; Waye, J.;
Voruganti,

[182986]P.; Hudson, C.; Serjeant, G. R.; Masibay, A. S.; Seeman, M.
V.: Dopamine

[182987]D4 receptor variant, D4GLYCINE194, in Africans, but not
in Caucasians:

[182988]no association with schizophrenia. Am. J. Med. Genet. 54:
384–390,

[182989]1994.

[182990]

[182991]

[182992]19091.

[182993]Smalley, S. L.; Bailey, J. N.; Palmer, C. G.; Cantwell, D. P.;

[182994]McGough, J. J.; Del'Homme, M. A.; Asarnow, J. R.; Woodward, J. A.;

[182995]Ramsey, C.; Nelson, S. F.: Evidence that the dopamine D4 receptor

[182996]is a susceptibility gene in attention deficit hyperactivity disorder. Molec.

[182997]Psychiat. 3: 427–430, 1998.

[182998]

[182999]

[183000]19092.

[183001]Meera Khan, P.: Personal Communication. Leiden, The Netherlands

[183002]1977.

[183003]

[183004]

[183005]19093.

[183006]Takai, T.; Li, M.; Sylvestre, D.; Clynes, R.; Ravetch, J. V.:

[183007]FcR gamma chain deletion results in pleiotrophic effector
cell defects. Cell 76:

[183008]519–529, 1994.

[183009]

[183010]

[183011]19094.

[183012]Garman, S. C.; Kinet, J.–P.; Jardetzky, T. S.: Crystal struc-
ture

[183013]of the human high–affinity IgE receptor. Cell 95: 951–961,
1998.

[183014]

[183015]

[183016]19095.

[183017]Garman, S. C.; Wurzburg, B. A.; Tarchevskaya, S. S.; Kinet,
J.–P.;

[183018]Jardetzky, T. S.: Structure of the Fc fragment of human IgE
bound

[183019]to its high–affinity receptor Fc–epsilon–RI–alpha. Nature

406: 259–266,
[183020]2000.

[183021]

[183022]

[183023]19096.

[183024]Liu, F.–T.; Albrandt, K.; Robertson, M. W.: cDNA hetero-
geneity

[183025]suggests structural variants related to the high–affinity IgE
receptor. Proc.

[183026]Nat. Acad. Sci. 85: 5639–5643, 1988.

[183027]

[183028]

[183029]19097.

[183030]Shimizu, A.; Tepler, I.; Benfey, P. N.; Berenstein, E. H.; Sir-
aganian,

[183031]R. P.; Leder, P.: Human and rat mast cell high–affinity im-
munoglobulin

[183032]E receptors: characterization of putative alpha–chain gene
products. Proc.

[183033]Nat. Acad. Sci. 85: 1907–1911, 1988.

[183034]

[183035]

[183036]19098.

[183037]Miyajima, I.; Dombrowicz, D.; Martin, T. R.; Ravetch, J. V.;
Kinet,

[183038]J.-P.; Galli, S. J.: Systemic anaphylaxis in the mouse can be
mediated

[183039]largely through IgG1 and Fc-gamma-RIII: assessment of
the cardiopulmonary

[183040]changes, mast cell degranulation, and death associated
with active

[183041]or IgE- or IgG1-dependent passive anaphylaxis. J. Clin.
Invest. 99:

[183042]901-914, 1997.

[183043]

[183044]

[183045]19099.

[183046]Tepler, I.; Morton, C. C.; Shimizu, A.; Holcombe, R. F.;
Eddy,

[183047]R.; Shows, T. B.; Leder, P.: The gene for the human mast
cell high-affinity

[183048]IgE receptor alpha chain: chromosomal localization to
1q21-q23 and

[183049]RFLP analysis. Am. J. Hum. Genet. 45: 761-765, 1989.

[183050]

[183051]

[183052]19100.

[183053]Zhu, D.; Kepley, C. L.; Zhang, M.; Zhang, K.; Saxon, A.: A
novel

[183054]human immunoglobulin Fc-gamma-Fc-epsilon bifunc-
tional fusion protein

[183055]inhibits Fc-epsilon-RI-mediated degranulation. Nature
Med. 8: 518-521,

[183056]2002.

[183057]

[183058]

[183059]19101.

[183060]Bain, G.; Robanus Maandag, E. C.; Izon, D. J.; Amsen, D.;
Kruisbeek,

[183061]A. M.; Weintraub, B. C.; Krop, I.; Schlissel, M. S.; Feeney, A.
J.;

[183062]van Roon, M.; van der Valk, M.; te Riele, H. P. J.; Berns, A.;
Murre,

[183063]C.: E2A proteins are required for proper B cell develop-
ment and initiation

[183064]of immunoglobulin gene rearrangements. Cell 79:
885–892, 1994.

[183065]

[183066]

[183067]19102.

[183068]El Ghouzzi, V.; Legeai–Mallet, L.; Aresta, S.; Benoist, C.;
Munnich,

[183069]A.; de Gunzburg, J.; Bonaventure, J.: Saethre–Chotzen mu-
tations cause

[183070]TWIST protein degradation or impaired nuclear location.
Hum. Molec.

[183071]Genet. 9: 813–819, 2000.

[183072]

[183073]

[183074]19103.

[183075]Henthorn, P.; Kiledjian, M.; Kadesch, T.: Two distinct tran-
scription

[183076]factors that bind the immunoglobulin enhancer mu-
E5/kappa-E2 motif. Science 247:

[183077]467-470, 1990.

[183078]

[183079]

[183080]19104.

[183081]Kamps, M. P.; Murre, C.; Sun, X.; Baltimore, D.: A new
homeobox

[183082]gene contributes the DNA binding domain of the t(1;19)
translocation

[183083]protein in pre-B ALL. Cell 60: 547-555, 1990.

[183084]

[183085]

[183086]19105.

[183087]Mellentin, J. D.; Murre, C.; Donlon, T. A.; McCaw, P. S.;
Smith,

[183088]S. D.; Carroll, A. J.; McDonald, M. E.; Baltimore, D.; Cleary,
M.

[183089]L.: The gene for enhancer binding proteins E12/E47 lies at
the t(1;19)

[183090]breakpoint in acute leukemias. Science 246: 379–382,
1989.

[183091]

[183092]

[183093]19106.

[183094]Murre, C.; McCaw, P. S.; Baltimore, D. :Cell 56: 777–783,
1989.

[183095]

[183096]

[183097]19107.

[183098]Nourse, J.; Mellentin, J. D.; Galili, N.; Wilkinson, J.; Stan-
bridge,

[183099]E.; Smith, S. D.; Cleary, M. L.: Chromosomal translocation
t(1;19)

[183100]results in synthesis of a homeobox fusion mRNA that
codes for a potential

[183101]chimeric transcription factor. Cell 60: 535–545, 1990.

[183102]

[183103]

[183104]19108.

[183105]Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[183106]M.; Kayano, T.; Shows, T. B.; Bell, G. I.: Assignment of
genes encoding

[183107]three human glucose transporter/transporter-like pro-
teins (GLUT4,

[183108]GLUT5 and GLUT6) to chromosomes 17, 1 and 5, respec-
tively. (Abstract) Cytogenet.

[183109]Cell Genet. 51: 997 only, 1989.

[183110]

[183111]

[183112]19109.

[183113]Gould, G. W.; Holman, G. D.: The glucose transporter fam-
ily: structure,

[183114]function and tissue-specific expression. Biochem. J. 295:
329-341,

[183115]1993.

[183116]

[183117]

[183118]19110.

[183119]Hauguel-de Mouzon, S.; Challier, J. C.; Kacemi, A.;

Cauzac, M.;

[183120]Malek, A.; Girard, J.: The GLUT3 glucose transporter isoform is differentially

[183121]expressed within human placental cell types. J. Clin. Endocr. Metab. 82:

[183122]2689–2694, 1997.

[183123]

[183124]

[183125]19111.

[183126]Kayano, T.; Burant, C. F.; Fukumoto, H.; Gould, G. W.; Fan, Y.

[183127]S.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Seino, S.; Bell, G. I.

[183128]: Human facilitative glucose transporters: isolation, functional characterization,

[183129]and gene localization of cDNAs encoding an isoform (GLUT5) expressed

[183130]in small intestine, kidney, muscle, and adipose tissue and an unusual

[183131]glucose transporter pseudogene-like sequence (GLUT6). J. Biol. Chem. 265:

[183132]13276–13282, 1990.

[183133]

[183134]

[183135]19112.

[183136]Kayano, T.; Fukumoto, H.; Eddy, R. L.; Fan, Y.-S.; Byers, M. G.;

[183137]Shows, T. B.; Bell, G. I.: Evidence for a family of human glucose

[183138]transporter-like proteins: sequence and gene localization of a protein

[183139]expressed in fetal skeletal muscle and other tissues. J. Biol. Chem. 263:

[183140]15245–15248, 1988.

[183141]

[183142]

[183143]19113.

[183144]Aitken, D. A.; Ferguson-Smith, M. A.: Gene dosage evidence for

[183145]the regional assignment of the GOT-S structural gene locus to 10q24–10q25. Cytogenet.

[183146]Cell Genet. 22: 468–471, 1978.

[183147]

[183148]

[183149]19114.

[183150]Creagan, R.; Tischfield, J.; McMorris, F. A.; Chen, S.-H.;
Hirschi,

[183151]M.; Chen, T.-T.; Ricciuti, F.; Ruddle, F. H.: Assignment of
the genes

[183152]for human peptidase A to chromosome 18 and cytoplas-
mic glutamic oxaloacetate

[183153]transaminase to chromosome 10 using somatic-cell hy-
brids. Cytogenet.

[183154]Cell Genet. 12: 187-198, 1973.

[183155]

[183156]

[183157]19115.

[183158]Ford, G. C.; Eichele, G.; Jansonius, J. N.: Three-di-
mensional

[183159]structure of a pyridoxal-phosphate-dependent enzyme,
mitochondrial

[183160]aspartate aminotransferase. Proc. Nat. Acad. Sci. 77:
2559-2563,

[183161]1980.

[183162]

[183163]

[183164]19116.

[183165]Gitelman, B. J.; Tomkins, D. J.; Partington, M. W.; Roberts,
M.

[183166]H.; Simpson, N. E.: Gene dosage studies of glutamic ox-
aloacetic transaminase

[183167](GOT) and hexokinase (HK) in two patients with possible
partial trisomy

[183168]10q. (Abstract) Am. J. Hum. Genet. 32: 41A only, 1980.

[183169]

[183170]

[183171]19117.

[183172]Junien, C.; Despoisse, S.; Turleau, C.; de Grouchy, J.;
Bucher,

[183173]T.; Fundele, R.: Assignment of phosphoglycerate mutase
(PGAMA) to

[183174]human chromosome 10: regional mapping of GOT1 and
PGAMA to subbands

[183175]10q26.1 (or q25.3). Ann. Genet. 25: 25–27, 1982.

[183176]

[183177]

[183178]19118.

[183179]Overhauser, J.; Mewar, R.; Rojas, K.; Lia, K.; Kline, A. D.;
Silverman,

[183180]G. A.: STS map of genes and anonymous DNA fragments
on human chromosome

[183181]18 using a panel of somatic cell hybrids. Genomics 15:
387–391,

[183182]1993.

[183183]

[183184]

[183185]19119.

[183186]Abdalla, S. A.; Pece–Barbara, N.; Vera, S.; Tapia, E.; Paez,
E.;

[183187]Bernabeu, C.; Letarte, M.: Analysis of ALK–1 and endoglin
in newborns

[183188]from families with hereditary hemorrhagic telangiectasia
type 2. Hum.

[183189]Molec. Genet. 9: 1227–1237, 2000.

[183190]

[183191]

[183192]19120.

[183193]Bode–Lesniewska, B.; Dours–Zimmermann, M. T.; Oder–
matt, B. F.;

[183194]Briner, J.; Heitz, P. U.; Zimmermann, D. R.: Distribution of
the

[183195]large aggregating proteoglycan versican in adult human
tissues. J.

[183196]Histochem. Cytochem. 44: 303–312, 1996.

[183197]

[183198]

[183199]19121.

[183200]Dours–Zimmermann, M. T.; Zimmermann, D. R.: A novel
glycosaminoglycan

[183201]attachment domain identified in two alternative splice
variants of

[183202]human versican. J. Biol. Chem. 269: 32992–32998, 1994.

[183203]

[183204]

[183205]19122.

[183206]Iozzo, R. V.; Naso, M. F.; Cannizzaro, L. A.; Wasmuth, J. J.;
McPherson,

[183207]J. D.: Mapping of the versican proteoglycan gene (CSPG2)
to the long

[183208]arm of human chromosome 5 (5q12–5q14). *Genomics* 14:
845–851, 1992.

[183209]

[183210]

[183211]19123.

[183212]Kjellen, L.; Lindahl, U.: Proteoglycans: structures and in-
teractions. *Annu.*

[183213]*Rev. Biochem.* 60: 443–475, 1991.

[183214]

[183215]

[183216]19124.

[183217]Naso, M. F.; Morgan, J. L.; Buchberg, A. M.; Siracusa, L. D.;
Iozzo,

[183218]R. V.: Expression pattern and mapping of the murine ver-

sican gene

[183219](Cspg2) to chromosome 13. Genomics 29: 297–300,
1995.

[183220]

[183221]

[183222]19125.

[183223]Naso, M. F.; Zimmermann, D. R.; Iozzo, R. V.: Characteri-
zation

[183224]of the complete genomic structure of the human versican
gene and functional

[183225]analysis of its promoter. J. Biol. Chem. 269:
32999–33008, 1994.

[183226]

[183227]

[183228]19126.

[183229]Zako, M.; Shinomura, T.; Ujita, M.; Ito, K.; Kimata, K.: Ex-
pression

[183230]of PG-M(V3), an alternatively spliced form of PG-M with-
out a chondroitin

[183231]sulfate attachment region in mouse and human tissues. J.
Biol. Chem. 270:

[183232]3914–3918, 1995.

[183233]

[183234]

[183235]19127.

[183236]Zimmermann, D. R.; Ruoslahti, E.: Multiple domains of the
large

[183237]fibroblast proteoglycan, versican. EMBO J. 8: 2975–2981,
1989.

[183238]

[183239]

[183240]19128.

[183241]Grunder, S.; Geisler, H.–S.; Rainer, S.; Fink, J. K.: Acid–
sensing

[183242]ion channel (ASIC) 4 gene: physical mapping, genomic or–
ganisation,

[183243]and evaluation as a candidate for paroxysmal dystonia.
Europ. J.

[183244]Hum. Genet. 9: 672–676, 2001.

[183245]

[183246]

[183247]19129.

[183248]Koch, G.; Lalley, P. A.; McAvoy, M.; Shows, T. B.: Assign-
ment

[183249]of LIPA, associated with human acid lipase deficiency, to
human chromosome

[183250]10 and comparative assignment to mouse chromosome
19. Somat. Cell

[183251]Genet. 7: 345–358, 1981.

[183252]

[183253]

[183254]19130.

[183255]Kadotani, H.; Hirano, T.; Masugi, M.; Nakamura, K.; Nakao,
K.;

[183256]Katsuki, M.; Nakanishi, S.: Motor discoordination results
from combined

[183257]gene disruption of the NMDA receptor NR2A and NR2C
subunits, but not

[183258]from single disruption of the NR2A or NR2C subunit. J.
Neurosci. 16:

[183259]7859–7867, 1996.

[183260]

[183261]

[183262]19131.

[183263]Lin, Y. J.; Bovetto, S.; Carver, J. M.; Giordano, T.: Cloning

[183264]of the cDNA for the human NMDA receptor NR2C subunit
and its expression

[183265]in the central nervous system and periphery. Molec. Brain
Res. 43:

[183266]57–64, 1996.

[183267]

[183268]

[183269]19132.

[183270]Islam, M. Q.; Platz, A.; Szpirer, J.; Szpirer, C.; Levan, G.;
Mannervik,

[183271]B.: Chromosomal localization of human glutathione trans-
ferase genes

[183272]of classes alpha, mu and pi. Hum. Genet. 82: 338–342,
1989.

[183273]

[183274]

[183275]19133.

[183276]Albert, L. J.; Inman, R. D.: Molecular mimicry and autoimmunity. New

[183277]Eng. J. Med. 341: 2068–2074, 1999.

[183278]

[183279]

[183280]19134.

[183281]Asada, H.; Kawamura, Y.; Maruyama, K.; Kume, H.; Ding, R.-G.; Ji,

[183282]F. Y.; Kanbara, N.; Kuzume, H.; Sanbo, M.; Yagi, T.; Obata, K.: Mice

[183283]lacking the 65 kDa isoform of glutamic acid decarboxylase (GAD65)

[183284]maintain normal levels of GAD67 and GABA in their brains but are susceptible

[183285]to seizures. Biochem. Biophys. Res. Commun. 229: 891–895, 1996.

[183286]

[183287]

[183288]19135.

[183289]Baekkeskov, S.; Aanstoot, H.-J.; Christgau, S.; Reetz, A.; Solimena,

[183290]M.; Cascalho, M.; Folli, F.; Richter-Olesen, H.; Camilli, P.-D.:

[183291]Identification of the 64K autoantigen in insulin-dependent diabetes

[183292]as the GABA-synthesizing enzyme glutamic acid decarboxylase. Nature 347:

[183293]151-156, 1990.

[183294]

[183295]

[183296]19136.

[183297]Bu, D.-F.; Erlander, M. G.; Hitz, B. C.; Tillakaratne, N. J. K.;

[183298]Kaufman, D. L.; Wagner-McPherson, C. B.; Evans, G. A.; Tobin, A. J.

[183299]: Two human glutamate decarboxylases, 65-kDa GAD and 67-kDa GAD, are

[183300]each encoded by a single gene. Proc. Nat. Acad. Sci. 89: 2115-2119,

[183301]1992.

[183302]

[183303]

[183304]19137.

[183305]Cram, D. S.; Barnett, L. D.; Joseph, J. L.; Harrison, L. C.:

Cloning

[183306]and partial nucleotide sequence of human glutamic acid
decarboxylase

[183307]cDNA from brain and pancreatic islets. Biochem. Biophys.

Res. Commun. 176:

[183308]1239–1244, 1991.

[183309]

[183310]

[183311]19138.

[183312]De Aizpurua, H. J.; Wilson, Y. M.; Harrison, L. C.: Glutamic

acid

[183313]decarboxylase autoantibodies in preclinical insulin-
dependent diabetes. Proc.

[183314]Nat. Acad. Sci. 89: 9841–9845, 1992.

[183315]

[183316]

[183317]19139.

[183318]Edelhoff, S.; Grubin, C. E.; Karlsen, A. E.; Adler, D. A.; Fos-

ter,

[183319]D.; Disteché, C. M.; Lernmark, A.: Mapping of glutamic acid decarboxylase

[183320](GAD) genes. Genomics 17: 93–97, 1993.

[183321]

[183322]

[183323]19140.

[183324]Jones, D. B.; Armstrong, N. W.: Coxsackie virus and diabetes revisited.

[183325](Letter) Nature Med. 1: 284 only, 1995.

[183326]

[183327]

[183328]19141.

[183329]Karlsen, A. E.; Hagopian, W. A.; Grubin, C. E.; Dube, S.; Disteché,

[183330]C. M.; Adler, D. A.; Barmeier, H.; Mathewes, S.; Grant, F. J.; Foster,

[183331]D.; Lernmark, A.: Cloning and primary structure of a human islet

[183332]isoform of glutamic acid decarboxylase from chromosome

10. Proc.

[183333]Nat. Acad. Sci. 88: 8337–8341, 1991.

[183334]

[183335]

[183336]19142.

[183337]Kash, S. F.; Johnson, R. S.; Tecott, L. H.; Noebels, J. L.;
Mayfield,

[183338]R. D.; Hanahan, D.; Baekkeskov, S.: Epilepsy in mice deficient in

[183339]the 65–kDa isoform of glutamic acid decarboxylase. Proc.
Nat. Acad.

[183340]Sci. 94: 14060–14065, 1997.

[183341]

[183342]

[183343]19143.

[183344]Kaufman, D. L.; Erlander, M. G.; Clare-Salzler, M.; Atkinson,
son,

[183345]M. A.; Maclaren, N. K.; Tobin, A. J.: Autoimmunity to two
forms of

[183346]glutamate decarboxylase in insulin–dependent diabetes

mellitus. J.

[183347]Clin. Invest. 89: 283–292, 1992.

[183348]

[183349]

[183350]19144.

[183351]Lohmann, T.; Hawa, M.; Leslie, R. D. G.; Lane, R.; Picard, J.;

[183352]Londei, M.: Immune reactivity to glutamic acid decarboxy-
lase 65 in

[183353]stiff-man syndrome and type 1 diabetes mellitus. Lancet
356: 31–35,

[183354]2000.

[183355]

[183356]

[183357]19145.

[183358]Richter, W.; Endl, J.; Eiermann, T. H.; Brandt, M.; Kientsch-En-
gel,

[183359]R.; Thivolet, C.; Jungfer, H.; Scherbaum, W. A.: Human
monoclonal

[183360]islet cell antibodies from a patient with insulin-dependent
diabetes

[183361]mellitus reveal glutamate decarboxylase as the target antigen. Proc.

[183362]Nat. Acad. Sci. 89: 8467–8471, 1992.

[183363]

[183364]

[183365]19146.

[183366]Tian, J.; Lehmann, P. V.; Kaufman, D. L.: T cell cross-reactivity

[183367]between coxsackievirus and glutamate decarboxylase is associated with

[183368]a murine diabetes susceptibility allele. J. Exp. Med. 180: 1979–1984,

[183369]1994.

[183370]

[183371]

[183372]19147.

[183373]McKusick, V. A.: The morbid anatomy of the human genome: a review

[183374]of gene mapping in clinical medicine (part 1). Medicine 65: 1–33,

[183375]1986.

[183376]

[183377]

[183378]19148.

[183379]Panteghini, M.: Aspartate aminotransferase isoenzymes.
Clin.

[183380]Biochem. 23: 311–319, 1990.

[183381]

[183382]

[183383]19149.

[183384]Pol, S.; Bousquet–Lemerrier, B.; Pave–Preux, M.; Bulle, F.;
Passage,

[183385]E.; Hanoune, J.; Mattei, M. G.; Barouki, R.: Chromosomal
localization

[183386]of human aspartate aminotransferase genes by in situ hy-
bridization. Hum.

[183387]Genet. 83: 159–164, 1989.

[183388]

[183389]

[183390]19150.

[183391]Scott, E. M.; Wright, R. C.: An alternate method for
demonstration

[183392]of erythrocytic aminotransferases on starch gels. Am. J.
Hum. Genet. 33:

[183393]561–563, 1981.

[183394]

[183395]

[183396]19151.

[183397]Spritz, R. A.; Emanuel, B. S.; Chern, C. J.; Mellman, W. J.:

[183398]Gene dosage effect: intraband mapping of human soluble
glutamic oxaloacetic

[183399]transaminase. Cytogenet. Cell Genet. 23: 149–156, 1979.

[183400]

[183401]

[183402]19152.

[183403]Tomkins, D. J.; Gitelman, B. J.; Roberts, M. H.: Confirma-
tion

[183404]of a de novo duplication, dup(10)(q24–q26), by GOT1

gene dosage studies. Hum.

[183405]Genet. 63: 369–373, 1983.

[183406]

[183407]

[183408]19153.

[183409]Wang, C.–Y.; Huang, Y.–Q.; Shi, J.–O.; Marron, M. P.; Ruan, Q.–G.;

[183410]Hawkins–Lee, B.; Ochoa, B.; She, J.–X.: Genetic homogeneity, high–resolution

[183411]mapping, and mutation analysis of the urofacial (Ochoa) syndrome and

[183412]exclusion of the glutamate oxaloacetate transaminase gene (GOT1) in

[183413]the critical region as the disease gene. Am. J. Med. Genet. 84:

[183414]454–459, 1999.

[183415]

[183416]

[183417]19154.

[183418]Wurzinger, K. H.; Mohrenweiser, H. W.: Studies on the ge–

netic

[183419]and non-genetic (physiological) variation of human erythrocyte glutamic

[183420]oxaloacetic transaminase. Ann. Hum. Genet. 46: 191–201, 1982.

[183421]

[183422]

[183423]19155.

[183424]Abel, E. D.; Kaulbach, H. C.; Tian, R.; Hopkins, J. C. A.; Duffy,

[183425]J.; Doetschman, T.; Minnemann, T.; Boers, M.–E.; Hadro, E.; Oberste–Berghaus,

[183426]C.; Quist, W.; Lowell, B. B.; Ingwall, J. S.; Kahn, B. B.: Cardiac

[183427]hypertrophy with preserved contractile function after selective deletion

[183428]of GLUT4 from the heart. J. Clin. Invest. 104: 1703–1714, 1999.

[183429]

[183430]

[183431]19156.

[183432]Abel, E. D.; Peroni, O.; Kim, J. K.; Kim, Y.-B.; Boss, O.;
Hadro,

[183433]E.; Minnemann, T.; Shulman, G. I.; Kahn, B. B.: Adipose-
selective

[183434]targeting of the GLUT4 gene impairs insulin action in
muscle and liver. Nature 409:

[183435]729–733, 2001.

[183436]

[183437]

[183438]19157.

[183439]Bell, G. I.; Kayano, T.; Buse, J. B.; Burant, C. F.; Takeda, J.;

[183440]Lin, D.; Fukumoto, H.; Seino, S.: Molecular biology of
mammalian

[183441]glucose transporters. Diabetes Care 13: 198–208, 1990.

[183442]

[183443]

[183444]19158.

[183445]Bell, G. I.; Murray, J. C.; Nakamura, Y.; Kayano, T.; Eddy, R.

[183446]L.; Fan, Y.–S.; Byers, M. G.; Shows, T. B.: Polymorphic hu-
man insulin-responsive

[183447]glucose-transporter gene on chromosome 17p13. Dia-
betes 38: 1072–1075,

[183448]1989.

[183449]

[183450]

[183451]19159.

[183452]Birnbaum, M. J.: Identification of a novel gene encoding an
insulin-responsive

[183453]glucose transporter protein. Cell 57: 305–315, 1989.

[183454]

[183455]

[183456]19160.

[183457]Chiang, S.-H.; Baumann, C. A.; Kanzaki, M.; Thurmond, D.
C.; Watson,

[183458]R. T.; Neudauer, C. L.; Macara, I. G.; Pessin, J. E.; Saltiel, A.

[183459]R.: Insulin-stimulated GLUT4 translocation requires the
CAP-dependent

[183460]activation of TC10. Nature 410: 944-948, 2001.

[183461]

[183462]

[183463]19161.

[183464]Garvey, W. T.; Maianu, L.; Zhu, J.-H.; Brechtel-Hook, G.;
Wallace,

[183465]P.; Baron, A. D.: Evidence for defects in the trafficking and
translocation

[183466]of GLUT4 glucose transporters in skeletal muscle as a
cause of human

[183467]insulin resistance. J. Clin. Invest. 101: 2377-2386, 1998.

[183468]

[183469]

[183470]19162.

[183471]Ikemoto, S.; Thompson, K. S.; Itakura, H.; Lane, M. D.;
Ezaki,

[183472]O.: Expression of an insulin-responsive glucose trans-
porter (GLUT4)

[183473]minigene in transgenic mice: effect of exercise and role in
glucose

[183474]homeostasis. Proc. Nat. Acad. Sci. 92: 865–869, 1995.

[183475]

[183476]

[183477]19163.

[183478]Katz, E. B.; Stenbit, A. E.; Hatton, K.; DePinho, R.; Charron,

[183479]M. J.: Cardiac and adipose tissue abnormalities but not di-
abetes

[183480]in mice deficient in GLUT4. Nature 377: 151–155, 1995.

[183481]

[183482]

[183483]19164.

[183484]Kusari, J.; Verma, U. S.; Buse, J. B.; Henry, R. R.; Olefsky,
[183485]J. M.: Analysis of the gene sequences of the insulin recep-
tor and
[183486]the insulin-sensitive glucose transporter (GLUT-4) in pa-
tients with
[183487]common-type non-insulin-dependent diabetes mellitus. J.
Clin. Invest. 88:
[183488]1323-1330, 1991.

[183489]

[183490]

[183491]19165.

[183492]Muraoka, A.; Sakura, H.; Kim, K.; Kishimoto, M.; Akanuma,
Y.;

[183493]Buse, J. B.; Yasuda, K.; Seino, S.; Bell, G. I.; Yazaki, Y.; Ka-
suga,

[183494]M.; Kadowaki, T.: Polymorphism in exon 4a of the human
GLUT4/muscle-fat

[183495]facilitative glucose transporter gene detected by SSCP.
Nucleic Acids

[183496]Res. 19: 4313 only, 1991.

[183497]

[183498]

[183499]19166.

[183500]Ribon, V.; Printen, J. A.; Hoffman, N. G.; Kay, B. K.; Saltiel,

[183501]A. R.: A novel, multifunctional c-Cbl binding protein in insulin

[183502]receptor signaling in 3T3-L1 adipocytes. Molec. Cell. Biol. 18:

[183503]872-879, 1998.

[183504]

[183505]

[183506]19167.

[183507]Zisman, A.; Peroni, O. D.; Abel, E. D.; Michael, M. D.; Mauvais-Jarvis,

[183508]F.; Lowell, B. B.; Wojtaszewski, J. F. P.; Hirshman, M. F.; Virkamaki,

[183509]A.; Goodyear, L. J.; Kahn, C. R.; Kahn, B. B.: Targeted disruption

[183510]of the glucose transporter 4 selectively in muscle causes insulin

[183511]resistance and glucose intolerance. Nature Med. 6: 924-928, 2000.

[183512]

[183513]

[183514]19168.

[183515]Astrin, K. H.; Arredondo-Vega, F. X.; Desnick, R. J.; Smith, M.

[183516]: Assignment of the gene for cytosolic alanine aminotransferase (AAT1)

[183517]to human chromosome 8. Ann. Hum. Genet. 46: 125-133, 1982.

[183518]

[183519]

[183520]19169.

[183521]Chen, S.-H.; Giblett, E. R.: Polymorphism of soluble glutamic-pyruvic

[183522]transaminase: a new genetic marker in man. Science 173: 148-149,

[183523]1971.

[183524]

[183525]

[183526]19170.

[183527]Cook, P. J. L.; Jeremiah, S. J.; Buckton, K. E.: Exclusion mapping

[183528]of GPT. (Abstract) Cytogenet. Cell Genet. 32: 258 only, 1982.

[183529]

[183530]

[183531]19171.

[183532]Mombaerts, P.; Wang, F.; Dulac, C.; Vassar, R.; Chao, S. K.; Nemes,

[183533]A.; Mendelsohn, M.; Edmondson, J.; Axel, R.: The molecular biology

[183534]of olfactory perception. Cold Spring Harbor Symp. Quant. Biol. 61:

[183535]135–145, 1996.

[183536]

[183537]

[183538]19172.

[183539]Reed, R. R.: Genetic approaches to mammalian olfaction. Cold

[183540]Spring Harbor Symp. Quant. Biol. 61: 165–172, 1996.

[183541]

[183542]

[183543]19173.

[183544]Ronnett, G. V.; Snyder, S. H.: Molecular messengers of olfaction. Trends

[183545]Neurosci. 15: 508–513, 1992.

[183546]

[183547]

[183548]19174.

[183549]Schwab, S. G.; Hallmayer, J.; Lerer, B.; Albus, M.; Bornmann, M.;

[183550]Honig, S.; Strauss, M.; Segman, R.; Lichtermann, D.; Knapp, M.; Trixler,

[183551]M.; Maier, W.; Wildenauer, D. B.: Support for a chromosome 18p locus

[183552]conferring susceptibility to functional psychoses in families with

[183553]schizophrenia, by association and linkage analysis. Am. J. Hum. Genet. 63:

[183554]1139–1152, 1998.

[183555]

[183556]

[183557]19175.

[183558]Davignon, I.; Barnard, M.; Gavrilova, O.; Sweet, K.; Wilkie, T.

[183559]M.: Gene structure of murine Gna11 and Gna15: tandemly duplicated

[183560]Gq class G protein alpha subunit genes. Genomics 31: 359–366, 1996.

[183561]

[183562]

[183563]19176.

[183564]Jiang, M.; Pandey, S.; Tran, V. T.; Fong, H. K.: Guanine nucleotide-binding

[183565]regulatory proteins in retinal pigment epithelial cells. Proc. Nat.

[183566]Acad. Sci. 88: 3907–3911, 1991.

[183567]

[183568]

[183569]19177.

[183570]Offermanns, S.; Zhao, L.-P.; Gohla, A.; Sarosi, I.; Simon, M. I.;

[183571]Wilkie, T. M.: Embryonic cardiomyocyte hypoplasia and craniofacial

[183572]defects in G-alpha-q/G-alpha-11-mutant mice. EMBO J. 17: 4304-4312,

[183573]1998.

[183574]

[183575]

[183576]19178.

[183577]Strathmann, M. P.; Simon, M. I.: G-alpha-12 and G-alpha-13 subunits

[183578]define a fourth class of G protein alpha subunits. Proc. Nat. Acad.

[183579]Sci. 88: 5582-5586, 1991.

[183580]

[183581]

[183582]19179.

[183583]Ahmed, S. F.; Barr, D. G. D.; Bonthron, D. T.: GNAS1 mutations

[183584]and progressive osseous heteroplasia. (Letter) New Eng. J. Med. 346:

[183585]1669–1670, 2002.

[183586]

[183587]

[183588]19180.

[183589]Ahmed, S. F.; Dixon, P. H.; Bonthron, D. T.; Stirling, H. F.; Barr,

[183590]D. G. D.; Kelnar, C. J. H.; Thakker, R. V.: GNAS1 mutational analysis

[183591]in pseudohypoparathyroidism. Clin. Endocr. 49: 525–531, 1998.

[183592]

[183593]

[183594]19181.

[183595]Ahrens, W.; Hiort, O.; Staedt, P.; Kirschner, T.; Marschke, C.;

[183596]Kruse, K.: Analysis of the GNAS1 gene in Albright's hereditary osteodystrophy. J.

[183597]Clin. Endocr. Metab. 86: 4630–4634, 2001.

[183598]

[183599]

[183600]19182.

[183601]Aldred, M. A.; Trembath, R. C.: Activating and inactivating mutations

[183602]in the human GNAS1 gene. Hum. Mutat. 16: 183–189, 2000.

[183603]

[183604]

[183605]19183.

[183606]Ashley, P. L.; Ellison, J.; Sullivan, K. A.; Bourne, H. R.; Cox,

[183607]D. R.: Chromosomal assignment of the murine Gi and Gs genes. (Abstract) Am.

[183608]J. Hum. Genet. 41: A155 only, 1987.

[183609]

[183610]

[183611]19184.

[183612]Ballare, E.; Mantovani, S.; Lania, A.; Di Blasio, A. M.; Vallar,

[183613]L.; Spada, A.: Activating mutations of the GS-alpha gene

are associated

[183614]with low levels of GS- α protein in growth hormone-secreting tumors. J.

[183615]Clin. Endocr. Metab. 83: 4386–4390, 1998.

[183616]

[183617]

[183618]19185.

[183619]Bastepe, M; Pincus, J. E.; Sugimoto, T.; Tojo, K. Kanatani, M.;

[183620]Azuma, Y.; Kruse, K.; Rosenbloom, A. L.; Koshiyama, H.; Juppner, H.

[183621]: Positional dissociation between the genetic mutation responsible

[183622]for pseudohypoparathyroidism type Ib and the associated methylation

[183623]defect at exon A/B: evidence for a long-range regulatory element within

[183624]the imprinted GNAS1 locus. Hum. Molec. Genet. 10: 1231–1241, 2001.

[183625]

[183626]

[183627]19186.

[183628]Bastepe, M.; Juppner, H.: GNAS1 mutations and progressive osseous

[183629]heteroplasia. (Letter) New Eng. J. Med. 346: 1671 only, 2002.

[183630]

[183631]

[183632]19187.

[183633]Bastepe, M.; Lane, A. H.; Juppner, H.: Parental uniparental isodisomy

[183634]of chromosome 20q--and the resulting changes in GNAS1 methylation--as

[183635]a plausible cause of pseudohypoparathyroidism. Am. J. Hum. Genet. 68:

[183636]1283-1289, 2001.

[183637]

[183638]

[183639]19188.

[183640]Bianco, P.; Riminucci, M.; Majolagbe, A.; Kuznetsov, S. A.; Collins,

[183641]M. T.; Mankani, M. H.; Corsi, A.; Bone, H. G.; Wientroub, S.; Spiegel,

[183642]A. M.; Fisher, L. W.; Robey, P. G.: Mutations of the GNAS1 gene,

[183643]stromal cell dysfunction, and osteomalacic changes in non-McCune-Albright

[183644]fibrous dysplasia of bone. J. Bone Miner. Res. 15: 120-128, 2000.

[183645]

[183646]

[183647]19189.

[183648]Billestrup, N.; Swanson, L. W.; Vale, W.: Growth hormone-releasing

[183649]factor stimulates proliferation of somatotrophs in vitro. Proc. Nat.

[183650]Acad. Sci. 83: 6854-6857, 1986.

[183651]

[183652]

[183653]19190.

[183654]Candelieri, G. A.; Glorieux, F. H.; Prud'Homme, J.; St.-Arnaud,

[183655]R.: Increased expression of the c-fos proto-oncogene in bone from

[183656]patients with fibrous dysplasia. New Eng. J. Med. 332: 1546–1551,

[183657]1995.

[183658]

[183659]

[183660]19191.

[183661]Candelieri, G. A.; Roughley, P. J.; Glorieux, F. H.: Polymerase

[183662]chain reaction–based technique for the selective enrichment and analysis

[183663]of mosaic arg201 mutations in G alpha s from patients with fibrous

[183664]dysplasia of bone. Bone 21: 201–206, 1997.

[183665]

[183666]

[183667]19192.

[183668]Carel, J. C.; Le Stunff, C.; Condamine, L.; Mallet, E.; Chaus-sain,

[183669]J. L.; Adnot, P.; Garabedian, M.; Bougneres, P.: Resistance to the

[183670]lipolytic action of epinephrine: a new feature of protein GS deficiency. J.

[183671]Clin. Endocr. Metab. 84: 4127–4131, 1999.

[183672]

[183673]

[183674]19193.

[183675]Carter, A.; Bardin, C.; Collins, R.; Simons, C.; Bray, P.; Spiegel,

[183676]A.: Reduced expression of multiple forms of the alpha subunit of

[183677]the stimulatory GTP-binding protein in pseudohypoparathyroidism type

[183678]Ia. Proc. Nat. Acad. Sci. 84: 7266–7269, 1987.

[183679]

[183680]

[183681]19194.

[183682]Cattanach, B. M.; Kirk, M.: Differential activity of maternally

[183683]and paternally derived chromosome regions in mice. Na-
ture 315: 496–498,
[183684]1985.

[183685]

[183686]

[183687]19195.

[183688]Coutant, R.; Lumbroso, S.; Rey, R.; Lahlou, N.; Venara, M.;
Rouleau,

[183689]S.; Sultan, C.; Limal, J.–M.: Macroorchidism due to au-
tonomous hyperfunction

[183690]of Sertoli cells and GS-alpha gene mutation: an unusual
expression

[183691]of McCune–Albright syndrome in a prepubertal boy. J.
Clin. Endocr.

[183692]Metab. 86: 1778–1781, 2001.

[183693]

[183694]

[183695]19196.

[183696]Kim, H.–S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T.,

[183697]Jr.; Rosenfeld, R. G.: Identification of a family of low–

affinity

[183698]insulin-like growth factor binding proteins (IGFBPs): characterization

[183699]of connective tissue growth factor as a member of the IGFBP superfamily. Proc.

[183700]Nat. Acad. Sci. 94: 12981–12986, 1997.

[183701]

[183702]

[183703]19197.

[183704]Martinerie, C.; Viegas-Pequignot, E.; Guenard, I.; Dutrillaux,

[183705]B.; Nguyen, V. C.; Bernheim, A.; Perbal, B.: Physical mapping of

[183706]human loci homologous to the chicken nov proto-oncogene. Oncogene 7:

[183707]2529–2534, 1992.

[183708]

[183709]

[183710]19198.

[183711]Nakanishi, T.; Yamaai, T.; Asano, M.; Nawachi, K.; Suzuki,

M.;

[183712]Sugimoto, T.; Takigawa, M.: Overexpression of connective tissue growth

[183713]factor/hypertrophic chondrocyte-specific gene product 24 decreases

[183714]bone density in adult mice and induces dwarfism.

Biochem. Biophys.

[183715]Res. Commun. 281: 678–681, 2001.

[183716]

[183717]

[183718]19199.

[183719]Castor, C. W.; Furlong, A. M.; Carter-Su, C.: Connective tissue

[183720]activation. XXIX. Stimulation of glucose transport by connective tissue

[183721]activating peptide-III. Biochemistry 24: 1762–1767, 1985.

[183722]

[183723]

[183724]19200.

[183725]Castor, C. W.; Miller, J. W.; Walz, D. A.: Structural and bio-

logical

[183726]characteristics of connective tissue activating peptide
(CTAP-III),

[183727]a major human platelet derived growth factor. Proc. Nat.
Acad. Sci. 80:

[183728]765–769, 1983.

[183729]

[183730]

[183731]19201.

[183732]Krijgsveld, J.; Zaat, S. A. J.; Meeldijk, J.; van Veelen, P. A.;

[183733]Fang, G.; Poolman, B.; Brandt, E.; Ehlert, J. E.; Kuijpers, A.
J.;

[183734]Engbers, G. H. M.; Feijen, J.; Dankert, J.: Thrombocidins,
microbicidal

[183735]proteins from human blood platelets, are C-terminal dele-
tion products

[183736]of CXC chemokines. J. Biol. Chem. 275: 20374–20381,
2000.

[183737]

[183738]

[183739]19202.

[183740]Majumdar, S.; Gonder, D.; Koutsis, B.; Poncz, M.: Characterization

[183741]of the human beta-thromboglobulin gene: comparison with the gene for

[183742]platelet factor 4. J. Biol. Chem. 266: 5785–5789, 1991.

[183743]

[183744]

[183745]19203.

[183746]O'Donovan, N.; Galvin, M.; Morgan, J. G.: Physical mapping of

[183747]the CXC chemokine locus on human chromosome 4. Cytogenet. Cell Genet. 84:

[183748]39–42, 1999.

[183749]

[183750]

[183751]19204.

[183752]Tunnacliffe, A.; Majumdar, S.; Yan, B.; Poncz, M.: Genes for beta-thromboglobulin

[183753]and platelet factor 4 are closely linked and form part of a cluster

[183754]of related genes on chromosome 4. Blood 79: 2896–2900,
1992.

[183755]

[183756]

[183757]19205.

[183758]Wenger, R. H.; Hameister, H.; Clemetson, K. J.: Human
platelet

[183759]basic protein/connective tissue activating peptide–III
maps in a gene

[183760]cluster on chromosome 4q12–q13 along with other genes
of the beta–thromboglobulin

[183761]superfamily. Hum. Genet. 87: 367–368, 1991.

[183762]

[183763]

[183764]19206.

[183765]McGill, J. R.; Boyd, D.; Barrett, K. J.; Drysdale, J. W.; Moore,

[183766]C. M.: Localization of human ferritin H (heavy) and L (light)
subunits

[183767]by in situ hybridization. (Abstract) Am. J. Hum. Genet. 36:
146S

[183768]only, 1984.

[183769]

[183770]

[183771]19207.

[183772]Murray, M. T.; White, K.; Munro, H. N.: Conservation of
ferritin

[183773]heavy subunit gene structure: implications for the regula-
tion of ferritin

[183774]gene expression. Proc. Nat. Acad. Sci. 84: 7438–7442,
1987.

[183775]

[183776]

[183777]19208.

[183778]Papadopoulos, P.; Bhavsar, D.; Zappone, E.; David, V.;
Jones,

[183779]C.; Worwood, M.; Drysdale, J.: A second human ferritin H
locus on

[183780]chromosome 11. Cytogenet. Cell Genet. 61: 107–108,
1992.

[183781]

[183782]

[183783]19209.

[183784]Richard, C. W.; Withers, D. A.; Meeker, T. C.; Myers, R. M.:

[183785]A radiation hybrid map of the proximal long arm of human chromosome

[183786]11 containing the MEN-1 and bcl-1 disease locus.

(Abstract) Cytogenet.

[183787]Cell Genet. 58: 1970 only, 1991.

[183788]

[183789]

[183790]19210.

[183791]Worwood, M.; Brook, J. D.; Cragg, S. J.; Hellkuhl, B.; Jones,

[183792]B. M.; Perera, P.; Roberts, S. H.; Shaw, D. J.: Assignment of human

[183793]ferritin genes to chromosomes 11 and 19q13.3-19qter.

Hum. Genet. 69:

[183794]371-374, 1985.

[183795]

[183796]

[183797]19211.

[183798]Wu, K.-J.; Polack, A.; Dalla-Favera, R.: Coordinated regulation

[183799]of iron-controlling genes, H-ferritin and IRP2, by c-MYC. Science 283:

[183800]676-679, 1999.

[183801]

[183802]

[183803]19212.

[183804]Yachou, A.; Mattei, M. G.; Roeckel, N.; Grandchamp, B.; Beaumont,

[183805]C.: Mouse ferritin H sequences map to chromosomes 3, 6, and 19. Genomics 9:

[183806]204-206, 1991.

[183807]

[183808]

[183809]19213.

[183810]Yachou, A.-K.; Renaudie, F.; Guenet, J.-L.; Simon-Chazottes, D.;

[183811]Jones, R.; Grandchamp, B.; Beaumont, C.: Mouse ferritin H

multigene

[183812]family is polymorphic and contains a single multiallelic
functional

[183813]gene located on chromosome 19. Genomics 10: 531–538,
1991.

[183814]

[183815]

[183816]19214.

[183817]Yousoufian, H.; Chance, P.; Tuck–Muller, C. M.; Jabs, E.
W.:

[183818]Association of a new chromosomal deletion
[del(1)(q32q42)] with diaphragmatic

[183819]hernia: assignment of a human ferritin gene. Hum. Genet.
78: 267–270,

[183820]1988.

[183821]

[183822]

[183823]19215.

[183824]Aguilar–Martinez, P.; Biron, C.; Masméjean, C.; Jeanjean,
P.; Schved,

[183825]J.–F.: A novel mutation in the iron responsive element of

ferritin

[183826]L-subunit gene as a cause for hereditary hyperferritine-
mia-cataract

[183827]syndrome. (Letter) Blood 88: 1895-1903, 1996.

[183828]

[183829]

[183830]19216.

[183831]Beaumont, C.; Leneuve, P.; Devaux, I.; Scoazec, J.-Y.;
Berthier,

[183832]M.; Loiseau, M.-N.; Grandchamp, B.; Bonneau, D.: Muta-
tion in the

[183833]iron responsive element of the L ferritin mRNA in a family
with dominant

[183834]hyperferritinaemia and cataract. Nature Genet. 11:
444-446, 1995.

[183835]

[183836]

[183837]19217.

[183838]Brown, A. J. P.; Leibold, E. A.; Munro, H. N.: Isolation of
cDNA

[183839]clones for the light subunit of rat liver ferritin: evidence

that

[183840]the light subunit is encoded by a multigene family. Proc.
Nat. Acad.

[183841]Sci. 80: 1265–1269, 1983.

[183842]

[183843]

[183844]19218.

[183845]Camaschella, C.; Zecchina, G.; Lockitch, G.; Roetto, A.;
Campanella,

[183846]A.; Arosio, P.; Levi, S.: A new mutation (G51C) in the iron-
responsive

[183847]element (IRE) of L-ferritin associated with hyperferriti-
naemia-cataract

[183848]syndrome decreases the binding affinity of the mutated
IRE for iron-regulatory

[183849]proteins. Brit. J. Haemat. 108: 480–482, 2000.

[183850]

[183851]

[183852]19219.

[183853]Cazzola, M.; Bergamaschi, G.; Tonon, L.; Arbustini, E.;

Grasso,

[183854]M.; Vercesi, E.; Barosi, G.; Bianchi, P. E.; Cairo, G.; Arosio, P.

[183855]: Hereditary hyperferritinemia–cataract syndrome: relationship between

[183856]phenotypes and specific mutations in the iron–responsive element of

[183857]ferritin light–chain mRNA. Blood 90: 814–821, 1997.

[183858]

[183859]

[183860]19220.

[183861]Cazzola, M.; Foglieni, B.; Bergamaschi, G.; Levi, S.; Lazzarino,

[183862]M.; Arosio, P.: A novel deletion of the L–ferritin iron–responsive

[183863]element responsible for severe hereditary hyperferritinaemia–cataract

[183864]syndrome. Brit. J. Haemat. 116: 667–670, 2002.

[183865]

[183866]

[183867]19221.

[183868]Xu, W.; Gorman, P. A.; Rider, S. H.; Hedge, P. J.; Moore, G.;
[183869]Prichard, C.; Sheer, D.; Solomon, E.: Construction of a ge-
netic map
[183870]of human chromosome 17 by use of chromosome-medi-
ated gene transfer. Proc.
[183871]Nat. Acad. Sci. 85: 8563–8567, 1988.
[183872]
[183873]
[183874]19222.
[183875]Aarts, H. J. M.; Den Dunnen, J. T.; Lubsen, N. H.; Schoen-
makers,
[183876]J. G. G.: Linkage between the beta-B2 and beta-B3 crys-
tallin genes
[183877]in man and rat: a remnant of an ancient beta-crystallin
gene cluster. Gene 59:
[183878]127–135, 1987.
[183879]
[183880]
[183881]19223.
[183882]Fishman, G. I.; Eddy, R. L.; Shows, T. B.; Rosenthal, L.;

Leinwand,

[183883]L. A.: The human connexin gene family of gap junction proteins: distinct

[183884]chromosomal locations but similar structures. Genomics 10: 250–256,

[183885]1991.

[183886]

[183887]

[183888]19224.

[183889]Campbell, H. D.; Webb, G. C.; Kono, T.; Taniguchi, T.; Ford, J.

[183890]H.; Young, I. G.: Assignment of the interleukin–2 receptor beta chain

[183891]gene (Il–2rb) to band E on mouse chromosome 15. Genomics 12: 179–180,

[183892]1992.

[183893]

[183894]

[183895]19225.

[183896]Bonneau, D.; Huret, J. L.; Godeau, G.; Couet, D.; Putter–

man, M.;

[183897]Tanzer, J.; Babin, P.; Larregue, M.: Recurrent ctb(7)(q31.3)

and

[183898]possible laminin involvement in a neonatal cutis laxa with

a Marfan

[183899]phenotype. Hum. Genet. 87: 317–319, 1991.

[183900]

[183901]

[183902]19226.

[183903]Buchanan, R.; Wyatt, G. P.: Marfan's syndrome presenting

as an

[183904]intrapartum death. Arch. Dis. Child. 60: 1074–1076,

1985.

[183905]

[183906]

[183907]19227.

[183908]Burgeson, R. E.; Chiquet, M.; Deutzmann, R.; Ekblom, P.;

Engel,

[183909]J.; Kleinman, H.; Martin, G. R.; Meneguzzi, G.; Paulsson,

M.; Sanes,

[183910]J.; Timpl, R.; Tryggvason, K.; Yamada, Y.; Yurchenco, P. D.:

A new

[183911]nomenclature for the laminins. Matrix Biol. 14: 209–211,
1994.

[183912]

[183913]

[183914]19228.

[183915]Day, D. L.; Burke, B. A.: Pulmonary emphysema in a
neonate with

[183916]Marfan syndrome. Pediat. Radiol. 16: 518–521, 1986.

[183917]

[183918]

[183919]19229.

[183920]Elliott, R. W.; Barlow, D.; Hogan, B. L. M.: Linkage of genes

[183921]for laminin B1 and B2 subunits on chromosome 1 in
mouse. In Vitro

[183922]Cell Dev. Biol. 21: 477–484, 1985.

[183923]

[183924]

[183925]19230.

[183926]Gross, D. M.; Robinson, L. K.; Smith, L. T.; Glass, N.;
Rosenberg,

[183927]H.; Duvic, M.: Severe perinatal Marfan syndrome. *Pedi-*
atrics 84:

[183928]83–89, 1989.

[183929]

[183930]

[183931]19231.

[183932]Hohn, A. R.; Webb, H. M.: Cardiac studies of infant twins
with

[183933]Marfan's syndrome. *Am. J. Dis. Child.* 122: 526–528,
1971.

[183934]

[183935]

[183936]19232.

[183937]Hopker, V. H.; Shewan, D.; Tessier–Lavigne, M.; Poo, M.;
Holt,

[183938]C.: Growth–cone attraction to netrin–1 is converted to re–
pulsion

[183939]by laminin–1. *Nature* 401: 69–73, 1999.

[183940]

[183941]

[183942]19233.

[183943]Huret, J.; Bonneau, D.; Godeau, G.; Leheup, B.; Larregue, M.:

[183944]Neonatal cutis laxa with Marfan habitus: this syndrome maps with B1

[183945]laminin at the junction 7q31.3–q32. (Abstract) Cytogenet. Cell Genet. 58:

[183946]1922 only, 1991.

[183947]

[183948]

[183949]19234.

[183950]Iwamoto, Y.; Robey, F. A.; Graf, J.; Sasaki, M.; Kleinman, H.

[183951]K.; Yamada, Y.; Martin, G. R.: YIGSR, a synthetic laminin pentapeptide,

[183952]inhibits experimental metastasis formation. Science 238: 1132–1134,

[183953]1987.

[183954]

[183955]

[183956]19235.

[183957]Jaye, M.; Modi, W. S.; Ricca, G. A.; Mudd, R.; Chiu, I.-M.;
O'Brien,

[183958]S. J.; Drohan, W. N.: Isolation of a cDNA clone for the hu-
man laminin-B1

[183959]chain and its gene localization. Am. J. Hum. Genet. 41:
605-615,

[183960]1987.

[183961]

[183962]

[183963]19236.

[183964]Kleinman, H. K.: Personal Communication. Bethesda, Md.
1/7/1982.

[183965]

[183966]

[183967]19237.

[183968]Lababidi, Z.; Monzon, C.: Early cardiac manifestations of
Marfan's

[183969]syndrome in the newborn. Am. Heart J. 102: 943-945,

1981.

[183970]

[183971]

[183972]19238.

[183973]Modi, W. S.; Jaye, M.; O'Brien, S. J.: Chromosomal localization
tion

[183974]of a cDNA clone for the human B1 laminin chain.

(Abstract) Cytogenet.

[183975]Cell Genet. 46: 663 only, 1987.

[183976]

[183977]

[183978]19239.

[183979]Neimann, N.; Rauber, G.; Marchal, C.; Vidailhet, M.; Fall,
M.

[183980]: Maladie de Marfan chez un nouveau-ne avec atteintes
polyviscerales:

[183981]etude anatomo-clinique. Ann. Paediat. 15: 619-624,
1968.

[183982]

[183983]

[183984]19240.

[183985]Pikkarainen, T.; Eddy, R.; Fukushima, Y.; Byers, M.; Shows, T.;

[183986]Pihlajaniemi, T.; Saraste, M.; Tryggvason, K.: Human laminin B1 chain:

[183987]a multidomain protein with gene (LAMB1) locus in the q22 region of

[183988]chromosome 7. J. Biol. Chem. 262: 10454–10462, 1987.

[183989]

[183990]

[183991]19241.

[183992]Pikkarainen, T.; Savolainen, E.-R.; Tryggvason, K.: Nhe I and

[183993]Hinc II polymorphisms in the human laminin B1 chain gene on 7q22. Nucleic

[183994]Acids Res. 17: 4424 only, 1989.

[183995]

[183996]

[183997]19242.

[183998]Sasaki, M.; Kato, S.; Kohno, K.; Martin, G. R.; Yamada, Y.:

Sequence

[183999]of the cDNA encoding the laminin B1 chain reveals a multidomain protein

[184000]containing cysteine-rich repeats. Proc. Nat. Acad. Sci. 84: 935-939,

[184001]1987.

[184002]

[184003]

[184004]19243.

[184005]Vuolteenaho, R.; Chow, L. T.; Tryggvason, K.: Structure of the

[184006]human laminin B1 chain gene. J. Biol. Chem. 265: 15611-15616, 1990.

[184007]

[184008]

[184009]19244.

[184010]Gremlich, S.; Porret, A.; Hani, E. H.; Cherif, D.; Vionnet, N.;

[184011]Froguel, P.; Thorens, B.: Cloning, functional expression, and chromosomal

[184012]localization of the human pancreatic islet glucose-de-

pendent insulintropic

[184013]polypeptide receptor. Diabetes 44: 1202–1208, 1995.

[184014]

[184015]

[184016]19245.

[184017]Miyawaki, K.; Yamada, Y.; Ban, N.; Ihara, Y.; Tsukiyama, K.;
Zhou,

[184018]H.; Fujimoto, S.; Oku, A.; Tsuda, K.; Toyokuni, S.; Hiai, H.;
Mizunoya,

[184019]W.; and 9 others: Inhibition of gastric inhibitory polypep-
tide

[184020]signaling prevents obesity. Nature Med. 8: 738–742,
2002.

[184021]

[184022]

[184023]19246.

[184024]Stoffel, M.; Fernald, A. A.; Le Beau, M. M.; Bell, G. I.: As-
signment

[184025]of the gastric inhibitory polypeptide receptor gene (GIPR)
to chromosome

[184026]bands 19q13.2–q13.3 by fluorescence in situ hybridiza-

tion. Genomics 28:

[184027]607–609, 1995.

[184028]

[184029]

[184030]19247.

[184031]Usdin, T. B.; Mezey, E.; Button, D. C.; Brownstein, M. J.;
Bonner,

[184032]T. I.: Gastric inhibitory polypeptide receptor, a member of
the secretin–vasoactive

[184033]intestinal peptide receptor family, is widely distributed in
peripheral

[184034]organs and the brain. Endocrinology 133: 2861–2870,
1993.

[184035]

[184036]

[184037]19248.

[184038]Volz, A.; Groke, R.; Lankat–Buttgereit, B.; Fehmann, H.–C.;
Bode,

[184039]H. P.; Goke, B.: Molecular cloning, functional expression,
and signal

[184040]transduction of the GIP–receptor cloned from a human in–

sulinoma. FEBS

[184041]Lett. 373: 23–29, 1995.

[184042]

[184043]

[184044]19249.

[184045]Yamada, Y.; Hayami, T.; Nakamura, K.; Kaisaki, P. J.;
Someya, Y.;

[184046]Wang, C.–Z.; Seino, S.; Seino, Y.: Human gastric inhibitory
polypeptide

[184047]receptor: cloning of the gene (GIPR) and cDNA. Genomics
29: 773–776,

[184048]1995.

[184049]

[184050]

[184051]19250.

[184052]Miyawaki, K.; Yamada, Y.; Yano, H.; Niwa, H.; Ban, N.;
Ihara, Y.;

[184053]Kubota, A.; Fujimoto, S.; Kajikawa, M.; Kuroe, A.; Tsuda,
K.; Hashimoto,

[184054]H.; Yamashita, T.; Jomori, T.; Tashiro, F.; Miyazaki, J.;

Seino, Y.

[184055] Glucose intolerance caused by a defect in the entero-insular axis:

[184056] a study in gastric inhibitory polypeptide receptor knock-out mice. Proc.

[184057] Nat. Acad. Sci. 96: 14843–14847, 1999.

[184058]

[184059]

[184060] 19251.

[184061] Fukushige, S.; Murotsu, T.; Matsubara, K.: Chromosomal assignment

[184062] of human genes for gastrin, thyrotropin (TSH)–beta subunit and C-erb-2

[184063] by chromosome sorting combined with velocity sedimentation and southern

[184064] hybridization. Biochem. Biophys. Res. Commun. 134: 477–483, 1986.

[184065]

[184066]

[184067] 19252.

[184068] Justice, M. J.; Gilbert, D. J.; Kinzler, K. W.; Vogelstein, B.;

[184069]Buchberg, A. M.; Ceci, J. D.; Matsuda, Y.; Chapman, V. M.;
Patriotis,

[184070]C.; Makris, A.; Tsiichlis, P. N.; Jenkins, N. A.; Copeland, N.
G.:

[184071]A molecular genetic linkage map of mouse chromosome
18 reveals extensive

[184072]linkage conservation with human chromosomes 5 and 18.
Genomics 13:

[184073]1281–1288, 1992.

[184074]

[184075]

[184076]19253.

[184077]Lebacqz–Verheyden, A.–M.; Bertness, V.; Kirsch, I.; Hollis,
G. F.;

[184078]McBride, O. W.; Battey, J.: Human gastrin–releasing pep-
tide gene

[184079]maps to chromosome band 18q21. Somat. Cell Molec.
Genet. 13: 81–86,

[184080]1987.

[184081]

[184082]

[184083]19254.

[184084]Naylor, S. L.; Sakaguchi, A. Y.; Spindel, E.; Chin, W. W.:

Human

[184085]gastrin-releasing peptide gene is located on chromosome
18. Somat.

[184086]Cell Molec. Genet. 13: 87-91, 1987.

[184087]

[184088]

[184089]19255.

[184090]Naylor, S. L.; Spindel, E.; Chin, W. W.; Sakaguchi, A. Y.:

Gastrin

[184091]releasing peptide gene is located on human chromosome
18. (Abstract) Cytogenet.

[184092]Cell Genet. 40: 711 only, 1985.

[184093]

[184094]

[184095]19256.

[184096]Sausville, E. A.; Lebacqz-Verheyden, A.-M.; Spindel, E. R.;

Cuttitta,

[184097]F.; Gazdar, A. F.; Battey, J. F.: Expression of the gastrin-

releasing

[184098]peptide gene in human small cell lung cancer: evidence
for alternative

[184099]processing resulting in three distinct mRNAs. J. Biol.
Chem. 261:

[184100]2451–2457, 1986.

[184101]

[184102]

[184103]19257.

[184104]Spindel, E. R.; Chin, W. W.; Price, J.; Rees, L. H.; Besser, G.

[184105]M.; Habener, J. F.: Cloning and characterization of cDNAs
encoding

[184106]human gastrin–releasing peptide. Proc. Nat. Acad. Sci. 81:
5699–5703,

[184107]1984.

[184108]

[184109]

[184110]19258.

[184111]Spindel, E. R.; Zilberberg, M. D.; Habener, J. F.; Chin, W. W.

[184112]: Two prohormones for gastrin-releasing peptide are encoded by two

[184113]mRNAs differing by 19 nucleotides. Proc. Nat. Acad. Sci. 83: 19–23,

[184114]1986.

[184115]

[184116]

[184117]19259.

[184118]Kudo, S.; Chagnovich, D.; Rearden, A.; Mattei, M. G.; Fukuda, M.

[184119]: Molecular analysis of a hybrid gene encoding human gly-cophorin variant

[184120]Miltnerberger V-like molecule. J. Biol. Chem. 265: 13825–13829, 1990.

[184121]

[184122]

[184123]19260.

[184124]Kudo, S.; Fukuda, M.: Identification of a novel human gly-cophorin,

[184125]glycophorin E, by isolation of genomic clones and complementary DNA

[184126]clones utilizing polymerase chain reaction. J. Biol. Chem.
265:

[184127]1102–1110, 1990.

[184128]

[184129]

[184130]19261.

[184131]Onda, M.; Fukuda, M.: Detailed physical mapping of the
genes encoding

[184132]glycophorins A, B, and E, as revealed by P1 plasmids con-
taining human

[184133]genomic DNA. Gene 159: 225–230, 1995.

[184134]

[184135]

[184136]19262.

[184137]Onda, M.; Kudo, S.; Rearden, A.; Mattei, M.–G.; Fukuda,
M.: Identification

[184138]of a precursor genomic segment that provided a sequence
unique to

[184139]glycophorin B and E genes. Proc. Nat. Acad. Sci. 90:
7220–7224,

[184140]1993.

[184141]

[184142]

[184143]19263.

[184144]Vignal, A.; Rahuel, C.; London, J.; Cherif Zahar, B.; Schaff, S.;

[184145]Hattab, C.; Okubo, Y.; Cartron, J.-P.: A novel gene member of the

[184146]human glycophorin A and B gene family: molecular cloning and expression. Europ.

[184147]J. Biochem. 191: 619–625, 1990.

[184148]

[184149]

[184150]19264.

[184151]Board, P. G.; Jones, I. M.; Bentley, A. K.: Molecular cloning

[184152]and nucleotide sequence of human alpha-1 acid glycoprotein cDNA. Gene 44:

[184153]127–131, 1986.

[184154]

[184155]

[184156]19265.

[184157]Cox, D. W.; Francke, U.: Direct assignment of orosomucoid to human

[184158]chromosome 9 and alpha-2-HS-glycoprotein to chromosome 3 using human

[184159]fetal liver x rat hepatoma hybrids. Hum. Genet. 70: 109-115, 1985.

[184160]

[184161]

[184162]19266.

[184163]Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Orosomucoid.

[184164]Washington: National Biomedical Research Foundation (pub.) 5: 1972.

[184165]Pp. D310-D316.

[184166]

[184167]

[184168]19267.

[184169]Dente, L.; Ciliberto, G.; Cortese, R.: Structure of the hu-

man

[184170]alpha-1-acid glycoprotein gene: sequence homology with
other human

[184171]acute phase protein genes. Nucleic Acids Res. 13:
3941-3952, 1985.

[184172]

[184173]

[184174]19268.

[184175]Dente, L.; Pizza, M. G.; Metspalu, A.; Cortese, R.: Structure

[184176]and expression of the genes coding for human alpha-
1-acid glycoprotein. EMBO

[184177]J. 6: 2289-2296, 1987.

[184178]

[184179]

[184180]19269.

[184181]Dente, L.; Ruther, U.; Tripodi, M.; Wagner, E. F.; Cortese,
R.

[184182]: Expression of human alpha-1-acid glycoprotein genes in
cultured

[184183]cells and in transgenic mice. Genes Dev. 2: 259-266,

1988.

[184184]

[184185]

[184186]19270.

[184187]Eap, C. B.; Cuendet, C.; Baumann, P.: Orosomucoid

(alpha-1 acid

[184188]glycoprotein) phenotyping by use of immobilized pH gradients with

[184189]8 M urea and immunoblotting: a new variant encountered in a population

[184190]study. Hum. Genet. 80: 183-185, 1988.

[184191]

[184192]

[184193]19271.

[184194]Eiberg, H.; Mohr, J.; Nielsen, L. S.: Linkage of orosomucoid (ORM)

[184195]to ABO and AK1. (Abstract) Cytogenet. Cell Genet. 32: 272 only,

[184196]1982.

[184197]

[184198]

[184199]19272.

[184200]Johnson, A. M.; Schmid, K.; Alper, C. A.: Inheritance of human
man

[184201]alpha(1)-acid glycoprotein (orosomucoid) variants. J. Clin.
Invest. 48:

[184202]2293-2299, 1969.

[184203]

[184204]

[184205]19273.

[184206]Luckenbach, C.; Kompf, J.; Ritter, H.: Orosomucoid (ORM1)
subtyping

[184207]and formal genetics. Hum. Genet. 87: 429-432, 1991.

[184208]

[184209]

[184210]19274.

[184211]Rocchi, M.; Roncuzzi, L.; Santamaria, R.; Archidiacono, N.;
Dente,

[184212]L.; Romeo, G.: Mapping through somatic cell hybrids and

cDNA probes

[184213]of protein C to chromosome 2, factor X to chromosome 13, and alpha-1-acid

[184214]glycoprotein to chromosome 9. Hum. Genet. 74: 30-33, 1986.

[184215]

[184216]

[184217]19275.

[184218]Schmid, K.; Tokita, K.; Yoshizaki, H.: The alpha-1-acid glycoprotein

[184219]variants of normal Caucasian and Japanese individuals. J. Clin. Invest. 44:

[184220]1394-1401, 1965.

[184221]

[184222]

[184223]19276.

[184224]Tokita, K.; Schmid, K.: Variants of alpha-1-acid glycoprotein. Nature 200:

[184225]266 only, 1963.

[184226]

[184227]

[184228]19277.

[184229]Tomei, L.; Eap, C. B.; Baumann, P.; Dente, L.: Use of trans-genic

[184230]mice for the characterization of human alpha-1-acid glycoprotein (orosomucoid)

[184231]variants. Hum. Genet. 84: 89-91, 1989.

[184232]

[184233]

[184234]19278.

[184235]Umetsu, K.; Ikeda, N.; Kashimura, S.; Suzuki, T.: Orosomucoid

[184236](ORM) typing by print lectinofixation: a new technique for isoelectric

[184237]focusing--two common alleles in Japan. Hum. Genet. 71: 223-224,

[184238]1985.

[184239]

[184240]

[184241]19279.

[184242]Umetsu, K.; Yuasa, I.; Nishimura, H.; Sasaki, H.; Suzuki, T.:

[184243]Genetic polymorphisms of orosomucoid and alpha-
2-HS-glycoprotein in

[184244]a Philippine population. Hum. Hered. 38: 287-290, 1988.

[184245]

[184246]

[184247]19280.

[184248]Webb, G. C.; Earle, M. E.; Merritt, C.; Board, P. G.: Local-
ization

[184249]of human alpha-1 glycoprotein genes to 9q31-q34.1. Cy-
togenet. Cell

[184250]Genet. 47: 18-21, 1988.

[184251]

[184252]

[184253]19281.

[184254]Weidinger, S.; Muller, T.; Schwarzfischer, F.; Cleve, H.:
Three

[184255]new orosomucoid (ORM) variants revealed by isoelectric
focusing and

[184256]print immunofixation. Hum. Genet. 77: 286-288, 1987.

[184257]

[184258]

[184259]19282.

[184260]Yuasa, I.; Umetsu, K.; Vogt, U.; Nakamura, H.; Nanba, E.;
Tamaki,

[184261]N.; Irizawa, Y.: Human orosomucoid polymorphism:
molecular basis

[184262]of the three common ORM1 alleles, ORM1*F1, ORM1*F2,
and ORM1*S. Hum.

[184263]Genet. 99: 393–398, 1997.

[184264]

[184265]

[184266]19283.

[184267]Yuasa, I.; Weidinger, S.; Umetsu, K.; Suenaga, K.; Ishimoto,
G.;

[184268]Eap, B. C.; Duche, J.–C.; Baumann, P.: Orosomucoid sys-
tem: 17 additional

[184269]orosomucoid variants and proposal for a new nomencla-
ture. Vox Sang. 64:

[184270]47–55, 1993.

[184271]

[184272]

[184273]19284.

[184274]Triantafilou, K.; Triantafilou, M.; Dedrick, R. L.: A
CD14-independent

[184275]LPS receptor cluster. Nature Immun. 2: 338–345, 2001.

[184276]

[184277]

[184278]19285.

[184279]Saito, S.; Okui, K.; Tokino, T.; Oshimura, M.; Nakamura,
Y.: Isolation

[184280]and mapping of 68 RFLP markers on human chromosome
6. Am. J. Hum.

[184281]Genet. 50: 65–70, 1992.

[184282]

[184283]

[184284]19286.

[184285]Takahashi, I.; Tanuma, R.; Hirata, M.; Hashimoto, K.: A
cosmid

[184286]clone at the D6S182 locus on human chromosome 6p12
contains the 90-kDa

[184287]heat shock protein beta-gene (HSP90-beta). Mammalian
Genome 5: 121-122,

[184288]1994.

[184289]

[184290]

[184291]19287.

[184292]Durkin, A. S.; Maglott, D. R.; Vamvakopoulos, N. C.;
Zoghbi, H.

[184293]Y.; Nierman, W. C.: Assignment of an intron-containing
human heat-shock

[184294]protein gene (hsp90-beta, HSPCB) to chromosome 6 near
TCTE1 (6p21)

[184295]and two intronless pseudogenes to chromosomes 4 and
15 by polymerase

[184296]chain reaction amplification from a panel of hybrid cell
lines. Genomics 18:

[184297]452-454, 1993.

[184298]

[184299]

[184300]19288.

[184301]Klinger, H. P.: Suppression of tumorigenicity in somatic cell

[184302]hybrids. I. Suppression and reexpression of tumorigenicity in diploid

[184303]human x D98/AH2 hybrids and independent segregation of tumorigenicity

[184304]from other cell phenotypes. Cytogenet. Cell Genet. 27: 254–266,

[184305]1980.

[184306]

[184307]

[184308]19289.

[184309]Lichy, J. H.; Modi, W. S.; Seuanez, H. N.; Howley, P. M.: Identification

[184310]of a human chromosome 11 gene which is differentially regulated in

[184311]tumorigenic and nontumorigenic somatic cell hybrids of HeLa cells. Cell

[184312]Growth Differ. 3: 541–548, 1992.

[184313]

[184314]

[184315]19290.

[184316]Stanbridge, E. J.: Suppression of malignancy in human cells. Nature 260:

[184317]17–20, 1976.

[184318]

[184319]

[184320]19291.

[184321]Corrigall, V. M.; Arastu, M.; Khan, S.; Shah, C.; Fife, M.; Smeets,

[184322]T.; Tak, P.–P.; Panayi, G. S.: Functional IL–2 receptor beta (CD122)

[184323]and gamma (CD132) chains are expressed by fibroblast–like synoviocytes:

[184324]activation by IL–2 stimulates monocyte chemoattractant protein–1 production. J.

[184325]Immun. 166: 4141–4147, 2001.

[184326]

[184327]

[184328]19292.

[184329]Gnarra, J. R.; Otani, H.; Wang, M. G.; McBride, O. W.;

Sharon,

[184330]M.; Leonard, W. J.: Human interleukin 2 receptor beta-chain gene:

[184331]chromosomal localization and identification of 5-prime regulatory

[184332]sequences. Proc. Nat. Acad. Sci. 87: 3440–3444, 1990.

[184333]

[184334]

[184335]19293.

[184336]Hatakeyama, M.; Tsudo, M.; Minamoto, S.; Kono, T.; Doi, T.; Miyata,

[184337]T.; Miyasaka, M.; Taniguchi, T.: Interleukin–2 receptor beta chain

[184338]gene: generation of three receptor forms by cloned human alpha and

[184339]beta chain cDNA's. Science 244: 551–556, 1989.

[184340]

[184341]

[184342]19294.

[184343]Lamaze, C.; Dujeancourt, A.; Baba, T.; Lo, C. G.; Ben-
merah, A.;

[184344]Dautry-Varsat, A.: Interleukin 2 receptors and detergent-
resistant

[184345]membrane domains define a clathrin-independent endo-
cytic pathway. Molec.

[184346]Cell 7: 661-671, 2001.

[184347]

[184348]

[184349]19295.

[184350]Shibuya, H.; Yoneyama, M.; Nakamura, Y.; Harada, H.;
Hatakeyama,

[184351]M.; Minamoto, S.; Kono, T.; Doi, T.; White, R.; Taniguchi,
T.: The

[184352]human interleukin-2 receptor beta-chain gene: genomic
organization,

[184353]promoter analysis and chromosomal assignment. Nucleic
Acids Res. 18:

[184354]3697-3703, 1990.

[184355]

[184356]

[184357]19296.

[184358]Suzuki, H.; Kundig, T. M.; Furlonger, C.; Wakeham, A.;
Timms, E.;

[184359]Matsuyama, T.; Schmits, R.; Simard, J. J. L.; Ohashi, P. S.;
Griesser,

[184360]H.; Taniguchi, T.; Paige, C. J.; Mak, T. W.: Deregulated T
cell activation

[184361]and autoimmunity in mice lacking interleukin-2 receptor
beta. Science 268:

[184362]1472-1476, 1995.

[184363]

[184364]

[184365]19297.

[184366]Tsudo, M.; Kitamura, F.; Miyasaka, M.: Characterization of
the

[184367]interleukin 2 receptor beta chain using three distinct
monoclonal

[184368]antibodies. Proc. Nat. Acad. Sci. 86: 1982-1986, 1989.

[184369]

[184370]

[184371]19298.

[184372]Alitalo, T.; Koistinen, R.; Aalto-Setälä, K.; Kontula, K.;
Julkunen,

[184373]M.; Janne, O.; de la Chapelle, A.: Gene for the human in-
sulin-like

[184374]growth factor-binding protein/placental protein 12: map-
ping to 7p12-p13

[184375]and demonstration of a BglII RFLP. (Abstract) Cytogenet.
Cell Genet. 51:

[184376]950 only, 1989.

[184377]

[184378]

[184379]19299.

[184380]Alitalo, T.; Kontula, K.; Koistinen, R.; Aalto-Setälä, K.;
Julkunen,

[184381]M.; Janne, O. A.; Seppälä, M.; de la Chapelle, A.: The gene
encoding

[184382]human low-molecular weight insulin-like growth-factor
binding protein

[184383](IGF-BP25): regional localization to 7p12-p13 and de-
scription of a

[184384]DNA polymorphism. Hum. Genet. 83: 335–338, 1989.

[184385]

[184386]

[184387]19300.

[184388]Ballard, F. J.; Baxter, R. C.; Binoux, M.; Clemmons, D. R.;
Drop,

[184389]S. L. S.; Hall, K.; Hintz, R. L.; Rechler, M. M.; Rutanen, E.
M.;

[184390]Schwander, J. C.: Report on the nomenclature of the IGF
binding proteins. J.

[184391]Clin. Endocr. Metab. 70: 817–818, 1990.

[184392]

[184393]

[184394]19301.

[184395]Brewer, M. T.; Stetler, G. L.; Squires, C. H.; Thompson, R.
C.;

[184396]Busby, W. H.; Clemmons, D. R.: Cloning, characterization,
and expression

[184397]of a human insulin–like growth factor binding protein.
Biochem. Biophys.

[184398]Res. Commun. 152: 1289–1297, 1988.

[184399]

[184400]

[184401]19302.

[184402]Brinkman, A.; Groffen, C.; Kortleve, D. J.; Geurts van
Kessel,

[184403]A.; Drop, S. L. S.: Isolation and characterization of a cDNA
encoding

[184404]the low molecular weight insulin-like growth factor bind-
ing protein

[184405](IBP-1). EMBO J. 7: 2417–2423, 1988.

[184406]

[184407]

[184408]19303.

[184409]Brinkman, A.; Groffen, C. A. H.; Kortleve, D. J.; Drop, S. L.
S.

[184410]: Organization of the gene encoding the insulin-like
growth factor

[184411]binding protein IBP-1. Biochem. Biophys. Res. Commun.
157: 898–907,

[184412]1988.

[184413]

[184414]

[184415]19304.

[184416]Ekstrand, J.; Ehrenborg, E.; Stern, I.; Stellan, B.; Zech, L.;

[184417]Luthman, H.: The gene for insulin-like growth factor
binding protein-1

[184418]is localized to human chromosomal region 7p14-p12.

Genomics 6: 413-418,

[184419]1990.

[184420]

[184421]

[184422]19305.

[184423]Popovici, R. M.; Lu, M.; Bhatia, S.; Faessen, G. H.; Giaccia,
A.

[184424]J.; Giudice, L. C.: Hypoxia regulates insulin-like growth
factor-binding

[184425]protein 1 in human fetal hepatocytes in primary culture:
suggestive

[184426]molecular mechanisms for in utero fetal growth restriction

caused

[184427]by uteroplacental insufficiency. J. Clin. Endocr. Metab. 86:
2653–2659,

[184428]2001.

[184429]

[184430]

[184431]19306.

[184432]Agarwal, N.; Hsieh, C. L.; Sills, D.; Swaroop, M.; Desai, B.;
Francke,

[184433]U.; Swaroop, A.: Sequence analysis, expression and chro-
mosomal localization

[184434]of a gene, isolated from a subtracted human retina cDNA
library, that

[184435]encodes an insulin-like growth factor binding protein
(IGFBP2). Exp.

[184436]Eye Res. 52: 549–561, 1991.

[184437]

[184438]

[184439]19307.

[184440]Allander, S. V.; Larsson, C.; Ehrenborg, E.; Suwanichkul,

A.; Weber,

[184441]G.; Morris, S. L.; Bajalica, S.; Kiefer, M. C.; Luthman, H.;
Powell,

[184442]D. R.: Characterization of the chromosomal gene and pro-
moter for

[184443]human insulin-like growth factor binding protein-5. J.
Biol. Chem. 269:

[184444]10891-10898, 1994.

[184445]

[184446]

[184447]19308.

[184448]Ehrenborg, E.; Vilhelmsdotter, S.; Bajalica, S.; Larsson, C.;
Stern,

[184449]I.; Koch, J.; Brondum-Nielsen, K.; Luthman, H.: Structure
and localization

[184450]of the human insulin-like growth factor-binding protein 2
gene. Biochem.

[184451]Biophys. Res. Commun. 176: 1250-1255, 1991.

[184452]

[184453]

[184454]19309.

[184455]Wood, W. I.; Cachianes, G.; Henzel, W. J.; Winslow, G. A.;
Spencer,

[184456]S. A.; Hellmiss, R.; Martin, J. L.; Baxter, R. C.: Cloning and
expression

[184457]of the growth hormone-dependent insulin-like growth
factor-binding

[184458]protein. Molec. Endocr. 2: 1176–1185, 1988.

[184459]

[184460]

[184461]19310.

[184462]Eklund, L.; Piuholo, J.; Komulainen, J.; Sormunen, R.; Ong-
varrasopone,

[184463]C.; Fassler, R.; Muona, A.; Ilves, M.; Ruskoaho, H.; Takala,
T. E.

[184464]S.; Pihlajaniemi, T.: Lack of type XV collagen causes a
skeletal

[184465]myopathy and cardiovascular defects in mice. Proc. Nat.
Acad. Sci. 98:

[184466]1194–1199, 2001.

[184467]

[184468]

[184469]19311.

[184470]Hagg, P. M.; Hagg, P. O.; Peltonen, S.; Autio-Harmanen,
H.; Pihlajaniemi,

[184471]T.: Location of type XV collagen in human tissues and its
accumulation

[184472]in the interstitial matrix of the fibrotic kidney. Am. J. Path.

150:

[184473]2075–2086, 1997.

[184474]

[184475]

[184476]19312.

[184477]Hagg, P. M.; Horelli-Kuitunen, N.; Eklund, L.; Palotie, A.;
Pihlajaniemi,

[184478]T.: Cloning of mouse type XV collagen sequences and
mapping of the

[184479]corresponding gene to 4B1–3: comparison of mouse and
human alpha–1(XV)

[184480]collagen sequences indicates divergence in the number of
small collagenous

[184481]domains. Genomics 45: 31–41, 1997.

[184482]

[184483]

[184484]19313.

[184485]Hagg, P. M.; Muona, A.; Lietard, J.; Kivirikko, S.; Pihla-
janiemi,

[184486]T.: Complete exon–intron organization of the human gene
for the alpha–1

[184487]chain of type XV collagen (COL15A1) and comparison with
the homologous

[184488]Col18a1 gene. J. Biol. Chem. 273: 17824–17831, 1998.

[184489]

[184490]

[184491]19314.

[184492]Huebner, K.; Cannizzaro, L. A.; Jabs, E. W.; Kivirikko, S.;
Manzone,

[184493]H.; Pihlajaniemi, T.; Myers, J. C.: Chromosomal assignment
of a gene

[184494]encoding a new collagen type (COL15A1) to 9q21–q22.

Genomics 14:

[184495]220–224, 1992.

[184496]

[184497]

[184498]19315.

[184499]Kivirikko, S.; Heinamaki, P.; Rehn, M.; Honkanen, N.; My-
ers, J.

[184500]C.; Pihlajaniemi, T.: Primary structure of the alpha-1 chain
of human

[184501]type XV collagen and exon-intron organization in the
3-prime region

[184502]of the corresponding gene. J. Biol. Chem. 269:
4773-4779, 1994.

[184503]

[184504]

[184505]19316.

[184506]Muragaki, Y.; Abe, N.; Ninomiya, Y.; Olsen, B. R.; Ooshima,
A.

[184507]: The human alpha-1(XV) collagen chain contains a large
amino-terminal

[184508]non-triple helical domain with a tandem repeat structure
and homology

[184509]to alpha-1(XVIII) collagen. J. Biol. Chem. 269: 4042-4046,
1994.

[184510]

[184511]

[184512]19317.

[184513]Myers, J. C.; Dion, A. S.; Abraham, V.; Amenta, P. S.: Type

XV

[184514]collagen exhibits a widespread distribution in human tissues but a

[184515]distinct localization in basement membrane zones. Cell Tissue Res. 286:

[184516]493–505, 1996.

[184517]

[184518]

[184519]19318.

[184520]Myers, J. C.; Kivirikko, S.; Gordon, M. K.; Dion, A. S.; Pihlajaniemi,

[184521]T.: Identification of a previously unknown human collagen chain,

[184522]alpha-1(XV), characterized by extensive interruptions in the triple-helical

[184523]region. Proc. Nat. Acad. Sci. 89: 10144–10148, 1992.

[184524]

[184525]

[184526]19319.

[184527]Ramchandran, R.; Dhanabal, M.; Volk, R.; Waterman, M. J.

F.; Segal,

[184528]M.; Lu, H.; Knebelmann, B.; Sukhatme, V. P.: Antiangio-
genic activity

[184529]of restin, NC10 domain of human collagen XV: compari-
son to endostatin. Biochem.

[184530]Biophys. Res. Commun. 255: 735–739, 1999.

[184531]

[184532]

[184533]19320.

[184534]Rehn, M.; Hintikka, E.; Pihlajaniemi, T.: Primary structure
of

[184535]the alpha 1 chain of mouse type XVIII collagen, partial
structure

[184536]of the corresponding gene, and comparison of the alpha
1(XVIII) chain

[184537]with its homologue, the alpha 1(XV) collagen chain. J. Biol.
Chem. 269:

[184538]13929–13935, 1994.

[184539]

[184540]

[184541]19321.

[184542]Sasaki, T.; Larsson, H.; Tisi, D.; Claesson-Welsh, L.; Hohenester,

[184543]E.; Timpl, R.: Endostatins derived from collagens XV and XVIII differ

[184544]in structural and binding properties, tissue distribution and anti-angiogenic

[184545]activity. J. Molec. Biol. 301: 1179–1190, 2000.

[184546]

[184547]

[184548]19322.

[184549]Pan, T.-C.; Zhang, R.-Z.; Mattei, M.-G.; Timpl, R.; Chu, M.-L.

[184550]: Cloning and chromosomal location of human alpha-1(XVI) collagen.

[184551]Proc. Nat. Acad. Sci. 89: 6565–6569, 1992.

[184552]

[184553]

[184554]19323.

[184555]Yamaguchi, N.; Kimura, S.; McBride, O. W.; Hori, H.; Yamada, Y.;

[184556]Kanamori, T.; Yamakoshi, H.; Nagai, Y.: Molecular cloning and partial

[184557]characterization of a novel collagen chain, alpha-1(XVI), consisting

[184558]of repetitive collagenous domains and cysteine-containing non-collagenous

[184559]segments. J. Biochem. 112: 856-863, 1992.

[184560]

[184561]

[184562]19324.

[184563]Curtis, A. R. J.; Fey, C.; Morris, C. M.; Bindoff, L. A.; Ince,

[184564]P. G.; Chinnery, P. F.; Coulthard, A.; Jackson, M. J.; Jackson, A.

[184565]P.; McHale, D. P.; Hay, D.; Barker, W. A.; Markham, A. F.; Bates,

[184566]D.; Curtis, A.; Burn, J.: Mutation in the gene encoding ferritin

[184567]light polypeptide causes dominant adult-onset basal ganglia disease. Nature

[184568]Genet. 28: 350-354, 2001.

[184569]

[184570]

[184571]19325.

[184572]Eiberg, H.; Bisgaard, M. L.; Mohr, J.: Linkage between alpha-1-B-glycoprotein

[184573](A1BG) and Lutheran (LU) red blood group system: assignment to chromosome

[184574]19: new genetic variants of A1BG. Clin. Genet. 36: 415-418, 1989.

[184575]

[184576]

[184577]19326.

[184578]Eiberg, H.; Nielsen, L. S.; Gahne, B.; Juneja, R. K.; Mohr, J.

[184579]: Exclusion data for the alpha-1-B glycoprotein (A1BG) polymorphism.

[184580](Abstract) Cytogenet. Cell Genet. 51: 994 only, 1989.

[184581]

[184582]

[184583]19327.

[184584]Gahne, B.; Juneja, R. K.; Stratil, A.: Genetic polymorphism

of

[184585]human plasma alpha-1-B-glycoprotein: phenotyping by immunoblotting

[184586]or by a simple method of 2-D electrophoresis. Hum.

Genet. 76: 111-115,

[184587]1987.

[184588]

[184589]

[184590]19328.

[184591]Ishioka, N.; Takahashi, N.; Putnam, F. W.: Amino acid sequence

[184592]of human plasma alpha-1B-glycoprotein: homology to the immunoglobulin

[184593]supergene family. Proc. Nat. Acad. Sci. 83: 2363-2367, 1986.

[184594]

[184595]

[184596]19329.

[184597]Juneja, R. K.; Weitkamp, L. R.; Stratil, A.; Gahne, B.; Gut-tormsen,

[184598]S. A.: Further studies of the plasma alpha-

1-B-glycoprotein polymorphism:

[184599]two new alleles and allele frequencies in Caucasians and
in American

[184600]blacks. Hum. Hered. 38: 267-272, 1988.

[184601]

[184602]

[184603]19330.

[184604]Anderson, L.; Anderson, N. G.: High resolution two-
dimensional

[184605]electrophoresis of human plasma proteins. Proc. Nat.
Acad. Sci. 74:

[184606]5421-5425, 1977.

[184607]

[184608]

[184609]19331.

[184610]Boutin, B.; Feng, S. H.; Arnaud, P.: The genetic polymor-
phism

[184611]of alpha(2)-HS glycoprotein: study by ultrathin-layer iso-
electric

[184612]focusing and immunoblot. Am. J. Hum. Genet. 37:

1098–1105, 1985.

[184613]

[184614]

[184615]19332.

[184616]Cox, D. W.; Andrews, B. J.: Silver stain immunofixation for
alpha-2-HS-glycoprotein:

[184617]a new method for detection of protein heterogeneity.In:
Stathakos,

[184618]D.: Electrophoresis '82. Berlin: Walter de Gruyter (pub.)
1983.

[184619]Pp. 243–247.

[184620]

[184621]

[184622]19333.

[184623]Cox, D. W.; Andrews, B. J.; Wills, D. E.: Genetic polymor-
phism

[184624]of alpha-2-HS-glycoprotein. Am. J. Hum. Genet. 38:
699–706, 1986.

[184625]

[184626]

[184627]19334.

[184628]Cox, D. W.; Francke, U.; Allderdice, P. W.; McAlpine, P. J.:
Gene

[184629]mapping of human serum proteins using hepatoma hy-
brids and human chromosome

[184630]deletions and duplications. Genetics 107 (suppl.):
s22-s23, 1984.

[184631]

[184632]

[184633]19335.

[184634]Eiberg, H.; Mohr, J.; Nielsen, L. S.: A2HS: new methods of
phenotyping

[184635]and analysis of linkage relations: assignment to chromo-
some 3. (Abstract) Cytogenet.

[184636]Cell Genet. 37: 461 only, 1984.

[184637]

[184638]

[184639]19336.

[184640]Gejyo, F.; Chang, J.-L.; Burgi, W.; Schmid, K.; Offner, G. D.;

[184641]Troxler, R. F.; Van Halbeek, H.; Dorland, L.; Gerwig, G. J.;

Vliegenthart,

[184642]J. F. G.: Characterization of the B-chain of human plasma
alpha(2)HS-glycoprotein:

[184643]the complete amino acid sequence and primary structure
of its heteroglycan. J.

[184644]Biol. Chem. 258: 4966–4971, 1983.

[184645]

[184646]

[184647]19337.

[184648]Jahnen–Dechent, W.: Personal Communication. Mainz,
Germany

[184649]7/22/1998.

[184650]

[184651]

[184652]19338.

[184653]Jahnen–Dechent, W.; Schinke, T.; Trindl, A.; Muller–Esterl,
W.;

[184654]Sablitzky, F.; Kaiser, S.; Blessing, M.: Cloning and targeted
deletion

[184655]of the mouse fetuin gene. J. Biol. Chem. 272:

31496–31503, 1997.

[184656]

[184657]

[184658]19339.

[184659]Lee, C.-C.; Bowman, B. H.; Yang, F.: Human alpha-
2-HS-glycoprotein:

[184660]the A and B chains with a connecting sequence are en-
coded by a single

[184661]mRNA transcript. Proc. Nat. Acad. Sci. 84: 4403–4407,
1987.

[184662]

[184663]

[184664]19340.

[184665]Hulsebos, T. J. M.; Bijlsma, E. K.; Geurts van Kessel, A. H.
M.;

[184666]Brakenhoff, R. H.; Westerveld, A.: Direct assignment of the
human

[184667]beta-B2 and beta-B3 and crystallin genes to
22q11.2–q12: markers for

[184668]neurofibromatosis 2. Cytogenet. Cell Genet. 56: 171–175,
1991.

[184669]

[184670]

[184671]19341.

[184672]Hulsebos, T. J. M.; Jenkins, N. A.; Gilbert, D. J.; Copeland,
N.

[184673]G.: The beta crystallin genes on human chromosome 22
define a new

[184674]region of homology with mouse chromosome 5. Genomics
25: 574–576,

[184675]1995.

[184676]

[184677]

[184678]19342.

[184679]Lampi, K. J.; Ma, Z.; Shih, M.; Shearer, T. R.; Smith, J. B.;
Smith,

[184680]D. L.; David, L. L.: Sequence analysis of beta-A3, beta-B3,
and beta-A4

[184681]crystallins completes the identification of the major pro-
teins in

[184682]young human lens. J. Biol. Chem. 272: 2268–2275, 1997.

[184683]

[184684]

[184685]19343.

[184686]van Rens, G.; Geurts van Kessel, A.; Bloemendal, H.: Local-
ization

[184687]of the beta-A4 crystallin (CRYBA4) gene on human chro-
mosome 22 in

[184688]the region q11.2-q13.1. (Abstract) Cytogenet. Cell Genet.
58: 2052

[184689]only, 1991.

[184690]

[184691]

[184692]19344.

[184693]van Rens, G. L. M.; Geurts van Kessel, A. H. M.; Bloemen-
dal, H.

[184694]: Localization of the beta-A4-crystallin gene (CRYBA4) on
human chromosome

[184695]22 in the region q11.2-q13.1. Cytogenet. Cell Genet. 61:
180-183,

[184696]1992.

[184697]

[184698]

[184699]19345.

[184700]Brakenhoff, R. H.; Henskens, H. A. M.; van Rossum, M. W.
P. C.;

[184701]Lubsen, N. H.; Schoenmakers, J. G. G.: Activation of the
gamma-E-crystallin

[184702]pseudogene in the human hereditary Coppock-like
cataract. Hum. Molec.

[184703]Genet. 3: 279–283, 1994.

[184704]

[184705]

[184706]19346.

[184707]den Dunnen, J. T.; Jongbloed, R. J. E.; Geurts van Kessel, A.
H.

[184708]M.; Schoenmakers, J. G. G.: Human lens gamma-crystallin
sequences

[184709]are located in the p12-qter region of chromosome 2.
Hum. Genet. 70:

[184710]217–221, 1985.

[184711]

[184712]

[184713]19347.

[184714]Hejtmancik, J. F.: The genetics of cataract: our vision be-
comes

[184715]clearer. (Editorial) Am. J. Hum. Genet. 62: 520–525, 1998.

[184716]

[184717]

[184718]19348.

[184719]Heon, E.; Priston, M.; Schorderet, D. F.; Billingsley, G. D.;
Girard,

[184720]P. O.; Lubsen, N.; Munier, F. L.: The gamma–crystallins
and human

[184721]cataracts: a puzzle made clearer. Am. J. Hum. Genet. 65:
1261–1267,

[184722]1999. Note: Erratum: Am. J. Hum. Genet. 66: 753 only,
2000.

[184723]

[184724]

[184725]19349.

[184726]Klopp, N.; Favor, J.; Loster, J.; Lutz, R. B.; Neuhauser-Klaus,

[184727]A.; Prescott, A.; Pretsch, W.; Quinlan, R. A.; Sandilands, A.; Vrensen,

[184728]G. F. J. M.; Graw, J.: Three murine cataract mutants (Cat2) are defective

[184729]in different gamma-crystallin genes. Genomics 52: 152–158, 1998.

[184730]

[184731]

[184732]19350.

[184733]Piatigorsky, J.: Lens crystallins and their gene families. Cell 38:

[184734]620–621, 1984.

[184735]

[184736]

[184737]19351.

[184738]Shiloh, Y.; Donlon, T.; Bruns, G.; Breitman, M. L.; Tsui, L.-C.

[184739]: Assignment of the human gamma-crystallin gene cluster (CRYG) to

[184740]the long arm of chromosome 2, region q33–36. Hum.

Genet. 73: 17–19,

[184741]1986.

[184742]

[184743]

[184744]19352.

[184745]Siezen, R. J.; Thomson, J. A.; Kaplan, E. D.; Benedek, G. B.:

[184746]Human lens gamma–crystallins: isolation, identification,
and characterization

[184747]of the expressed gene products. Proc. Nat. Acad. Sci. 84:
6088–6092,

[184748]1987.

[184749]

[184750]

[184751]19353.

[184752]Skow, L. C.; Donner, M. E.; Huang, S.–M.; Gardner, J. M.;
Taylor,

[184753]B. A.; Beamer, W. G.; Lalley, P. A.: Mapping of mouse
gamma crystallin

[184754]genes on chromosome 1. Biochem. Genet. 26: 557–570,

1988.

[184755]

[184756]

[184757]19354.

[184758]Tsui, L.-C.; Breitman, M. L.; Meakin, S. O.; Willard, H. F.;
Shiloh,

[184759]Y.; Donlon, T.; Bruns, G.: Localization of the human
gamma-crystallin

[184760]gene cluster (CRYG) to the long arm of chromosome 2, re-
gion q33-q35.

[184761](Abstract) Cytogenet. Cell Genet. 40: 763-764, 1985.

[184762]

[184763]

[184764]19355.

[184765]Vidal, S. M.; Epstein, D. J.; Malo, D.; Weith, A.; Vekemans,
M.;

[184766]Gros, P.: Identification and mapping of six microdissected
genomic

[184767]DNA probes to the proximal region of mouse chromo-
some 1. Genomics 14:

[184768]32-37, 1992.

[184769]

[184770]

[184771]19356.

[184772]Fajans, S. S.; Bell, G. I.; Polonsky, K. S.: Molecular mechanisms

[184773]and clinical pathophysiology of maturity-onset diabetes of the young. New

[184774]Eng. J. Med. 345: 971–980, 2001.

[184775]

[184776]

[184777]19357.

[184778]Lindner, T.; Gragnoli, C.; Furuta, H.; Cockburn, B. N.; Petzold,

[184779]C.; Rietzsch, H.; Weiss, U.; Schulze, J.; Bell, G. I.: Hepatic function

[184780]in a family with a nonsense mutation (R154X) in the hepatocyte nuclear

[184781]factor-4-alpha/MODY1 gene. J. Clin. Invest. 100: 1400–1405, 1997.

[184782]

[184783]

[184784]19358.

[184785]Moller, A. M.; Dalgaard, L. T.; Ambye, L.; Hansen, L.;
Schmitz,

[184786]O.; Hansen, T.; Pedersen, O.: A novel Phe75fsdelT mutation in the

[184787]hepatocyte nuclear factor-4-alpha gene in a Danish pedigree with maturity-onset

[184788]diabetes of the young. J. Clin. Endocr. Metab. 84:
367-369, 1999.

[184789]

[184790]

[184791]19359.

[184792]Thomas, H.; Jaschowitz, K.; Bulman, M.; Frayling, T. M.;
Mitchell,

[184793]S. M. S.; Roosen, S.; Lingott-Frieg, A.; Tack, C. J.; Ellard, S.;

[184794]Ryffel, G. U.; Hattersley, A. T.: A distant upstream promoter of

[184795]the HNF-4-alpha gene connects the transcription factors involved in

[184796]maturity-onset diabetes of the young. Hum. Molec. Genet.

10: 2089–2097,

[184797]2001.

[184798]

[184799]

[184800]19360.

[184801]Joseph, L. J.; Le Beau, M. M.; Jamieson, G. A., Jr.; Acharya, S.;

[184802]Shows, T. B.; Rowley, J. D.; Sukhatme, V. P.: Molecular cloning,

[184803]sequencing, and mapping of EGR2, a human early growth response gene

[184804]encoding a protein with 'zinc-binding finger' structure.

Proc. Nat.

[184805]Acad. Sci. 85: 7164–7168, 1988.

[184806]

[184807]

[184808]19361.

[184809]Nagarajan, R.; Svaren, J.; Le, N.; Araki, T.; Watson, M.; Milbrandt,

[184810]J.: EGR2 mutations in inherited neuropathies dominant–

negatively

[184811]inhibit myelin gene expression. Neuron 30: 355–368,
2001.

[184812]

[184813]

[184814]19362.

[184815]Timmerman, V.; De Jonghe, P.; Ceuterick, C.; De Vriendt,
E.; Lofgren,

[184816]A.; Nelis, E.; Warner, L. E.; Lupski, J. R.; Martin, J.-J.; Van
Broeckhoven,

[184817]C.: Novel missense mutation in the early growth response
2 gene associated

[184818]with Dejerine–Sottas syndrome phenotype. Neurology 52:
1827–1832,

[184819]1999.

[184820]

[184821]

[184822]19363.

[184823]Warner, L. E.; Mancias, P.; Butler, I.; Lupski, J. R.: Mutation

[184824]in the early growth response 2 (EGR2) transcription factor

associated

[184825]with recessive congenital hypomyelinating neuropathy
(CHN). (Abstract) Am.

[184826]J. Hum. Genet. 61 (suppl.): A350 only, 1997.

[184827]

[184828]

[184829]19364.

[184830]Warner, L. E.; Mancias, P.; Butler, I. J.; McDonald, C. M.;
Keppen,

[184831]L.; Koob, K. G.; Lupski, J. R.: Mutations in the early growth
response

[184832]2 (EGR2) gene are associated with hereditary
myelinopathies. Nature

[184833]Genet. 18: 382–384, 1998.

[184834]

[184835]

[184836]19365.

[184837]Warner, L. E.; Svaren, J.; Milbrandt, J.; Lupski, J. R.: Func-
tional

[184838]consequences of mutations in the early growth response 2

gene (EGR2)

[184839]correlate with severity of human myelinopathies. Hum.

Molec. Genet. 8:

[184840]1245–1251, 1999.

[184841]

[184842]

[184843]19366.

[184844]Wu, J.; Joseph, L.; Sukhatme, V. P.; Kidd, K. K.: A HindIII
polymorphism

[184845]identified by the human early growth response gene 2
(EGR2) on chromosome

[184846]

[184847]19367.

[184848]Nucleic Acids Res. 16: 11855 only, 1988.

[184849]

[184850]

[184851]19368.

[184852]Kaneda, Y.; Hayes, H.; Uchida, T.; Yoshida, M. C.; Okada,
Y.:

[184853]Regional assignment of five genes on human chromosome

19. Chromosoma 95:

[184854]8–12, 1987.

[184855]

[184856]

[184857]19369.

[184858]Hoeffler, J. P.; Meyer, T. E.; Yun, Y.; Jameson, J. L.;

Habener,

[184859]J. F.: Cyclic AMP–responsive DNA–binding protein: structure based

[184860]on a cloned placental cDNA. Science 242: 1430–1433, 1988.

[184861]

[184862]

[184863]19370.

[184864]Chan, E. F.; Gat, U.; McNiff, J. M.; Fuchs, E.: A common human

[184865]skin tumour is caused by activating mutations in beta–catenin. Nature

[184866]Genet. 21: 410–413, 1999.

[184867]

[184868]

[184869]19371.

[184870]Horelli-Kuitunen, N.; Kvist, A.-P.; Helaakoski, T.; Kivirikko,

[184871]K.; Pihlajaniemi, T.; Palotie, A.: The order and transcriptional

[184872]orientation of the human COL13A1 and P4HA genes on chromosome 10 long

[184873]arm determined by high-resolution FISH. Genomics 46: 299-302, 1997.

[184874]

[184875]

[184876]19372.

[184877]Pajunen, L.; Tamminen, M.; Solomon, E.; Pihlajaniemi, T.:

Assignment

[184878]of the gene coding for the alpha 1 chain of collagen type XIII (COL13A1)

[184879]to human chromosome region 10q11-qter. Cytogenet. Cell Genet. 52:

[184880]190-193, 1989.

[184881]

[184882]

[184883]19373.

[184884]Shows, T. B.; Tikka, L.; Byers, M. G.; Eddy, R. L.; Haley, L.
L.;

[184885]Henry, W. M.; Prockop, D. J.; Tryggvason, K.: Assignment
of the human

[184886]collagen alpha-1(XIII) chain gene (COL13A1) to the q22
region of chromosome

[184887]10. Genomics 5: 128-133, 1989.

[184888]

[184889]

[184890]19374.

[184891]Tikka, L.; Pihlajaniemi, T.; Henttu, P.; Prockop, D. J.; Tryg-
gvason,

[184892]K.: Gene structure for the alpha-1 chain of a human
short-chain collagen

[184893](type XIII) with alternatively spliced transcripts and trans-
lation

[184894]termination codon at the 5-prime end of the last exon.
Proc. Nat.

[184895]Acad. Sci. 85: 7491–7495, 1988.

[184896]

[184897]

[184898]19375.

[184899]Cunningham, B. A.; Hemperly, J. J.; Murray, B. A.; Prediger, E.

[184900]A.; Brackenbury, R.; Edelman, G. M.: Neural cell adhesion molecule:

[184901]structure, immunoglobulin-like domains, cell surface modulation, and

[184902]alternative RNA splicing. Science 236: 799–806, 1987.

[184903]

[184904]

[184905]19376.

[184906]D'Eustachio, P.; Davisson, M. T.: Resolution of the staggerer

[184907](sg) mutation from the neural cell adhesion molecule locus (Ncam)

[184908]on mouse chromosome 9. Mammalian Genome 4: 278–280, 1993.

[184909]

[184910]

[184911]19377.

[184912]D'Eustachio, P.; Owens, G. C.; Edelman, G. M.; Cunningham, B. A.

[184913]: Chromosomal location of the gene encoding the neural cell adhesion

[184914]molecule (N-CAM) in the mouse. Proc. Nat. Acad. Sci. 82: 7631-7635,

[184915]1985.

[184916]

[184917]

[184918]19378.

[184919]Lin, D. M.; Fetter, R. D.; Kopczynski, C.; Grenningloh, G.; Goodman,

[184920]C. S.: Genetic analysis of fasciclin II in Drosophila: defasciculation,

[184921]refasciculation, and altered fasciculation. Neuron 13: 1055-1069,

[184922]1994.

[184923]

[184924]

[184925]19379.

[184926]Altare, F.; Jouanguy, E.; Lamhamedi-Cherradi, S.; Fondaneche, M.-C.;

[184927]Fizame, C.; Ribierre, F.; Merlin, G.; Dembic, Z.; Schreiber, R.; Lisowska-Grospierre,

[184928]B.; Fischer, A.; Seboun, E.; Casanova, J.-L.: A causative relationship

[184929]between mutant IFN γ R1 alleles and impaired cellular response to IFN- γ

[184930]in a compound heterozygous child. (Letter) Am. J. Hum. Genet. 62:

[184931]723-726, 1998.

[184932]

[184933]

[184934]19380.

[184935]Dalton, D. K.; Pitts-Meek, S.; Keshav, S.; Figari, I. S.; Bradley,

[184936]A.; Stewart, T. A.: Multiple defects of immune cell function in mice

[184937]with disrupted interferon-gamma genes. Science 259:
1739-1742, 1993.

[184938]

[184939]

[184940]19381.

[184941]Dessein, A. J.; Hillaire, D.; Elwali, N. E. M. A.; Marquet, S.;

[184942]Mohamed-Ali, Q.; Mirghani, A.; Henri, S.; Abdelhameed, A.
A.; Saeed,

[184943]O. K.; Magzoub, M. M. A.; Abel, L.: Severe hepatic fibrosis
in Schistosoma

[184944]mansoni infection is controlled by a major locus that is
closely linked

[184945]to the interferon-gamma receptor gene. Am. J. Hum.
Genet. 65: 709-721,

[184946]1999.

[184947]

[184948]

[184949]19382.

[184950]Fellous, M.: Personal Communication. Paris, France
10/24/1986.

[184951]

[184952]

[184953]19383.

[184954]Fellous, M.; Couillin, P.; Rosa, F.; Metezeau, P.; Foubert, C.;

[184955]Gross, M. S.; Frezal, J.; Van Cong, N.: Receptor for human gamma

[184956]interferon is specified by human chromosome 18.

(Abstract) Cytogenet.

[184957]Cell Genet. 40: 627–628, 1985.

[184958]

[184959]

[184960]19384.

[184961]Jouanguy, E.; Altare, F.; Lamhamedi, S.; Revy, P.; Emile, J.–F.;

[184962]Newport, M.; Levin, M.; Blanche, S.; Seboun, E.; Fischer, A.; Casanova,

[184963]J.–L.: Interferon–gamma–receptor deficiency in an infant with fatal

[184964]bacille Calmette–Guerin infection. New Eng. J. Med. 335: 1956–1961,

[184965]1996.

[184966]

[184967]

[184968]19385.

[184969]Jouanguy, E.; Dupuis, S.; Pallier, A.; Doffinger, R.; Fon-
daneche,

[184970]M.-C.; Fieschi, C.; Lamhamedi-Cherradi, S.; Altare, F.;
Emile, J.-F.;

[184971]Lutz, P.; Bordigoni, P.; Cokugras, H.; Akcakaya, N.; Land-
man-Parker,

[184972]J.; Donnadieu, J.; Camcioglu, Y.; Casanova, J.-L.: In a novel
form

[184973]of IFN-gamma receptor 1 deficiency, cell surface recep-
tors fail to

[184974]bind IFN-gamma. J. Clin. Invest. 105: 1429-1436, 2000.

[184975]

[184976]

[184977]19386.

[184978]Jouanguy, E.; Lamhamedi-Cherradi, S.; Lammas, D.; Dor-
man, S. E.;

[184979]Fondaneche, M.-C.; Dupuis, S.; Doffinger, R.; Altare, F.;

Girdlestone,

[184980]J.; Emile, J.-F.; Ducoulombier, H.; Edgar, D.; and 10 others:

A

[184981]human IFNGR1 small deletion hotspot associated with
dominant susceptibility

[184982]to mycobacterial infection. Nature Genet. 21: 370–378,
1999.

[184983]

[184984]

[184985]19387.

[184986]Kaplan, D. H.; Shankaran, V.; Dighe, A. S.; Stockert, E.;

Aguet,

[184987]M.; Old, L. J.; Schreiber, R. D.: Demonstration of an inter-
feron

[184988]gamma-dependent tumor surveillance system in immuno-
competent mice. Proc.

[184989]Nat. Acad. Sci. 95: 7556–7561, 1998.

[184990]

[184991]

[184992]19388.

[184993]Le Coniat, M.; Alcaide-Loridan, C.; Fellous, M.; Berger, R.:

[184994]Human interferon gamma receptor 1 (IFNGR1) gene maps
to chromosome

[184995]region 6q23–6q24. Hum. Genet. 84: 92–94, 1989.

[184996]

[184997]

[184998]19389.

[184999]Lekstrom-Himes, J. A.; Gallin, J. I.: Immunodeficiency dis-
eases

[185000]caused by defects in phagocytes. New Eng. J. Med. 343:
1703–1714,

[185001]2000.

[185002]

[185003]

[185004]19390.

[185005]Levin, M.; Newport, M. J.; D'Souza, S.; Kalabalikis, P.;
Brown,

[185006]I. N.; Lenicker, H. M.; Agius, P. V.; Davies, E. G.; Thrasher,
A.;

[185007]Klein, N.; Blackwell, J. M.: Familial disseminated atypical

mycobacterial

[185008]infection in childhood: a human mycobacterial susceptibility gene? Lancet 345:

[185009]79–83, 1995.

[185010]

[185011]

[185012]19391.

[185013]Mariano, T. M.; Kozak, C. A.; Langer, J. A.; Pestka, S.: The

[185014]mouse immune interferon receptor gene is located on chromosome 10. J.

[185015]Biol. Chem. 262: 5812–5814, 1987.

[185016]

[185017]

[185018]19392.

[185019]Deloukas, P.; Schuler, G. D.; Gyapay, G.; Beasley, E. M.; Soderlund,

[185020]C.; Rodriguez–Tome, P.; Hui, L.; Matise, T. C.; McKusick, K. B.; Beckmann,

[185021]J. S.; Bentolila, S.; Bihoreau, M.–T.; and 53 others: A physical cal

[185022]map of 30,000 human genes. Science 282: 744–746,
1998.

[185023]

[185024]

[185025]19393.

[185026]Wu, W. J.; Erickson, J. W.; Lin, R.; Cerione, R. A.: The
gamma-subunit

[185027]of the coatamer complex binds Cdc42 to mediate trans-
formation. Nature 405:

[185028]800–804, 2000.

[185029]

[185030]

[185031]19394.

[185032]Davis, S. T.; Benson, B. G.; Bramson, H. N.; Chapman, D.
E.; Dickerson,

[185033]S. H.; Dold, K. M.; Eberwein, D. J.; Edelstein, M.; Frye, S. V.;
Gampe,

[185034]R. T., Jr.; Griffin, R. J.; Harris, P. A.; and 14 others: Preven-
tion

[185035]of chemotherapy-induced alopecia in rats by CDK in-
hibitors. Science 291:

[185036]134–137, 2001.

[185037]

[185038]

[185039]19395.

[185040]De Bondt, H. L.; Rosenblatt, J.; Jancarik, J.; Jones, H. D.;
Morgan,

[185041]D. O.; Kim, S.–H.: Crystal structure of cyclin–dependent
kinase 2. Nature 363:

[185042]595–602, 1993.

[185043]

[185044]

[185045]19396.

[185046]Basu, S.; Binder, R. J.; Ramalingam, T.; Srivastava, P. K.:
CD91

[185047]is a common receptor for heat shock proteins gp96,
hsp90, hsp70, and

[185048]calreticulin. Immunity 14: 303–313, 2001.

[185049]

[185050]

[185051]19397.

[185052]Beisiegel, U.; Weber, W.; Ihrke, G.; Herz, J.; Stanley, K. K.:

[185053]The LDL-receptor-related protein, LRP, is an apolipoprotein E-binding

[185054]protein. Nature 341: 162-164, 1989.

[185055]

[185056]

[185057]19398.

[185058]Binder, R. J.; Han, D. K.; Srivastava, P. K.: CD91: a receptor

[185059]for heat shock protein gp96. Nature Immun. 1: 151-155, 2000.

[185060]

[185061]

[185062]19399.

[185063]Forus, A.; Maelandsmo, G. M.; Fodstad, Y.; Myklebost, O.:

The

[185064]genes for the alpha-2-macroglobulin receptor/LDL receptor-related

[185065]protein and GLI are located within a chromosomal seg-

ment of about

[185066]300 kilobases and are coamplified in a rhabdomyosarcoma cell line.

[185067](Abstract) Cytogenet. Cell Genet. 58: 1977 only, 1991.

[185068]

[185069]

[185070]19400.

[185071]Forus, A.; Myklebost, O.: A physical map of a 1.3-Mb region on

[185072]the long arm of chromosome 12, spanning the GLI and LRP loci. Genomics 14:

[185073]117-120, 1992.

[185074]

[185075]

[185076]19401.

[185077]Hilliker, C.; Van Leuven, F.; Van Den Berghe, H.: Assignment of

[185078]the gene coding for the alpha(2)-macroglobulin receptor to mouse chromosome

[185079]15 and to human chromosome 12q13-q14 by isotopic

and nonisotopic in

[185080]situ hybridization. Genomics 13: 472–474, 1992.

[185081]

[185082]

[185083]19402.

[185084]Kristensen, T.; Moestrup, S. K.; Gliemann, J.; Bendtsen, L.;
Sand,

[185085]O.; Sottrup-Jensen, L.: Evidence that the newly cloned
low-density-lipoprotein

[185086]receptor related protein (LRP) is the alpha-
2-macroglobulin receptor. FEBS

[185087]Lett. 276: 151–155, 1990.

[185088]

[185089]

[185090]19403.

[185091]Lendon, C. L.; Talbot, C. J.; Craddock, N. J.; Han, S. W.;
Wragg,

[185092]M.; Morris, J. C.; Goate, A. M.: Genetic association studies
between

[185093]dementia of the Alzheimer's type and three receptors for

apolipoprotein

[185094]E in a Caucasian population. Neurosci. Lett. 222:
187–190, 1997.

[185095]

[185096]

[185097]19404.

[185098]Myklebost, O.; Arheden, K.; Rogne, S.; Geurts van Kessel,
A.;

[185099]Mandahl, N.; Herz, J.; Stanley, K.; Heim, S.; Mitelman, F.:
The gene

[185100]for the human putative apoE receptor is on chromosome
12 in the segment

[185101]q13–14. Genomics 5: 65–69, 1989.

[185102]

[185103]

[185104]19405.

[185105]Pericak–Vance, M. A.; Bass, M. P.; Yamaoka, L. H.; Gaskell,
P.

[185106]C.; Scott, W. K.; Terwedow, H. A.; Menold, M. M.; Con-
neally, P. M.;

[185107]Small, G. W.; Vance, J. M.; Saunders, A. M.; Roses, A. D.;

Haines,

[185108]J. L.: Complete genomic screen in late-onset familial
Alzheimer disease:

[185109]evidence for a new locus on chromosome 12. J.A.M.A.
278: 1237–1241,
[185110]1997.

[185111]

[185112]

[185113]19406.

[185114]Scott, W. K.; Yamaoka, L. H.; Bass, M. P.; Gaskell, P. C.;
Conneally,

[185115]P. M.; Small, G. W.; Farrer, L. A.; Auerbach, S. A.; Saunders,
A.

[185116]M.; Roses, A. D.; Haines, J. L.; Pericak–Vance, M. A.: No
genetic

[185117]association between the LRP receptor and sporadic or
late-onset familial

[185118]Alzheimer disease. Neurogenetics 1: 179–183, 1998.

[185119]

[185120]

[185121]19407.

[185122]Strickland, D. K.; Ashcom, J. D.; Williams, S.; Burgess, W.
H.;

[185123]Migliorini, M.; Argraves, W. S.: Sequence identity between
the alpha-2-macroglobulin

[185124]receptor and low density lipoprotein receptor-related pro-
tein suggests

[185125]that this molecule is a multifunctional receptor. J. Biol.
Chem. 265:

[185126]17401-17404, 1990.

[185127]

[185128]

[185129]19408.

[185130]Yochem, J.; Greenwald, I.: A gene for a low density
lipoprotein

[185131]receptor-related protein in the nematode *Caenorhabditis*
elegans. Proc.

[185132]Nat. Acad. Sci. 90: 4572-4576, 1993.

[185133]

[185134]

[185135]19409.

[185136]Ladiaz, J. A. A.; Karathanasis, S. K.: Regulation of the
apolipoprotein

[185137]AI gene by ARP-1, a novel member of the steroid receptor
superfamily. Science 251:

[185138]561-565, 1991.

[185139]

[185140]

[185141]19410.

[185142]Modi, W. S.; Seuanez, H.; Mietus-Snyder, M.; O'Brien, S. J.;
Karathanasis,

[185143]S. K.: Chromosomal localization of the ARP-1 gene to
15q26. (Abstract) Cytogenet.

[185144]Cell Genet. 58: 1995 only, 1991.

[185145]

[185146]

[185147]19411.

[185148]Pereira, F. A.; Qiu, Y.; Zhou, G.; Tsai, M.-J.; Tsai, S. Y.: The

[185149]orphan nuclear receptor COUP-TFII is required for angio-
genesis and

[185150]heart development. Genes Dev. 13: 1037-1049, 1999.

[185151]

[185152]

[185153]19412.

[185154]Widom, R. L.; Ladas, J. A. A.; Kouidou, S.; Karathanasis, S.
K.

[185155]: Synergistic interactions between transcription factors
control expression

[185156]of the apolipoprotein AI gene in liver cells. Molec. Cell.
Biol. 11:

[185157]677–687, 1991.

[185158]

[185159]

[185160]19413.

[185161]Ikeda, H.; Yamaguchi, M.; Sugai, S.; Aze, Y.; Narumiya, S.;
Kakizuka,

[185162]A.: Expanded polyglutamine in the Machado–Joseph dis-
ease protein

[185163]induces cell death in vitro and in vivo. Nature Genet. 13:
196–202,

[185164]1996.

[185165]

[185166]

[185167]19414.

[185168]Kang, H.; Sun, L. D.; Atkins, C. M.; Soderling, T. R.; Wilson,

[185169]M. A.; Tonegawa, S.: An important role of neural activity-
dependent

[185170]CaMKIV signaling in the consolidation of long-term mem-
ory. Cell 106:

[185171]771–783, 2001.

[185172]

[185173]

[185174]19415.

[185175]Raman, V.; Blaeser, F.; Ho, N.; Engle, D. L.; Williams, C. B.;

[185176]Chatila, T. A.: Requirement for
Ca(2+)/calmodulin-dependent kinase

[185177]type IV/Gr in setting the thymocyte selection threshold. J.
Immun. 167:

[185178]6270–6278, 2001.

[185179]

[185180]

[185181]19416.

[185182]Santisteban, I.; Arredondo-Vega, F. X.; Kelly, S.; Mary, A.;
Fischer,

[185183]A.; Hummell, D. S.; Lawton, A.; Sorensen, R. U.; Stiehm, E.
R.; Uribe,

[185184]L.; Weinberg, K.; Hershfield, M. S.: Novel splicing, mis-
sense, and

[185185]deletion mutations in seven adenosine deaminase-defi-
cient patients

[185186]with late/delayed onset of combined immunodeficiency
disease: contribution

[185187]of genotype to phenotype. J. Clin. Invest. 92: 2291-2302,
1993.

[185188]

[185189]100. Schmalstieg, F. C.; Mills, G. C.; Tsuda, H.; Goldman,
A. S.:

[185190]Severe combined immunodeficiency in a child with a
healthy adenosine

[185191]deaminase deficient mother. Pediat. Res. 17: 935-940,
1983.

[185192]

[185193]101. Schrader, W. P.; Pollara, B.; Meuwissen, H. J.: Characterization

[185194]of the residual adenosine deaminating activity in the spleen of a

[185195]patient with combined immunodeficiency disease and adenosine deaminase

[185196]deficiency. Proc. Nat. Acad. Sci. 75: 446–450, 1978.

[185197]

[185198]102. Scott, C. R.; Chen, S.–H.; Giblett, E. R.: Deletion of the carrier

[185199]state in combined immunodeficiency disease associated with deaminase

[185200]deficiency. J. Clin. Invest. 53: 1194–1196, 1974.

[185201]

[185202]103. Shovlin, C. L.; Hughes, J. M. B.; Simmonds, H. A.; Fairbanks,

[185203]L.; Deacock, S.; Lechler, R.; Roberts, I.; Webster, A. D. B.: Adult

[185204]presentation of adenosine deaminase deficiency. (Letter) Lancet 341:

[185205]1471, 1993.

[185206]

[185207]104. Shovlin, C. L.; Simmonds, H. A.; Fairbanks, L. D.;
Deacock, S.

[185208]J.; Hughes, J. M. B.; Lechler, R. I.; Webster, A. D. B.; Sun,
X.-M.;

[185209]Webb, J. C.; Soutar, A. K.: Adult onset immunodeficiency
caused by

[185210]inherited adenosine deaminase deficiency. J. Immun. 153:
2331-2339,

[185211]1994.

[185212]

[185213]105. Spencer, N.; Hopkinson, D. A.; Harris, H.: Adenosine
deaminase

[185214]polymorphism in man. Ann. Hum. Genet. 32: 9-14, 1968.

[185215]

[185216]106. Stephan, V.; Wahn, V.; Le Deist, F.; Dirksen, U.; Bro-
ker, B.;

[185217]Muller-Fleckenstein, I.; Horneff, G.; Schroten, H.; Fischer,
A.; de

[185218]Saint Basile, G.: Atypical X-linked severe combined im-
munodeficiency

[185219]due to possible spontaneous reversion of the genetic defect in T cells. New

[185220]Eng. J. Med. 335: 1563–1567, 1996.

[185221]

[185222]107. Tariverdian, G.; Ritter, H.: Adenosine deaminase polymorphism

[185223](EC 3.5.4.4): formal genetics and linkage relations. Humangenetik 7:

[185224]176–178, 1969.

[185225]

[185226]108. Tischfield, J. A.; Creagan, R. P.; Nichols, E. A.; Ruddle, F.

[185227]H.: Assignment of a gene for adenosine deaminase to human chromosome

[185228]20. Hum. Hered. 24: 1–11, 1974.

[185229]

[185230]109. Tzall, S.; Ellenbogen, A.; Eng, F.; Hirschhorn, R.: Identification

[185231]and characterization of nine RFLPs at the adenosine deaminase (ADA)

[185232]locus. Am. J. Hum. Genet. 44: 864–875, 1989.

[185233]

[185234]110. Umetsu, D. T.; Schlossman, C. M.; Ochs, H. D.; Her-
shfield, M.

[185235]S.: Heterogeneity of phenotype in two siblings with
adenosine deaminase

[185236]deficiency. J. Allergy Clin. Immun. 93: 543–550, 1994.

[185237]

[185238]111. Valerio, D.; Dekker, B. M. M.; Duyvesteyn, M. G. C.;
van der

[185239]Voorn, L.; Berkvens, T. M.; van Ormondt, H.; van der Eb, A.
J.: One

[185240]adenosine deaminase allele in a patient with severe com-
bined immunodeficiency

[185241]contains a point mutation abolishing enzyme activity.
EMBO J. 5:

[185242]113–119, 1986.

[185243]

[185244]112. Valerio, D.; Duyvesteyn, M. G. C.; Dekker, B. M. M.;
Weeda, G.;

[185245]Berkvens, T. M.; van der Voorn, L.; van Ormondt, H.; van
der Eb, A.

[185246]J.: Adenosine deaminase: characterization and expression
of a gene

[185247]with a remarkable promoter. EMBO J. 4: 437–443, 1985.

[185248]

[185249]113. Valerio, D.; Duyvesteyn, M. G. C.; Meera Khan, P.;
Pearson, P.

[185250]L.; Geurts van Kessel, A.; van Ormondt, H.: Direct assign-
ment of

[185251]ADA gene to chromosome 20. (Abstract) Cytogenet. Cell
Genet. 37:

[185252]599, 1984.

[185253]

[185254]114. Valerio, D.; Duyvesteyn, M. G. C.; van Ormondt, H.;
Meera Khan,

[185255]P.; van der Eb, A. J.: Adenosine deaminase (ADA) defi-
ciency in cells

[185256]derived from humans with severe combined immunodefi-
ciency is due to

[185257]an aberration of the ADA protein. Nucleic Acids Res. 12:
1015–1024,

[185258]1984.

[185259]

[185260]115. Valerio, D.; McIvor, R. S.; Williams, S. R.; Duyvesteyn, M. G.

[185261]C.; van Ormondt, H.; van der Eb, A. J.; Martin, D. W., Jr.:
Cloning

[185262]of human adenosine deaminase cDNA and expression in
mouse cells. Gene 31:

[185263]147–153, 1984.

[185264]

[185265]116. Van der Weyden, M. B.; Kelley, W. N.: Adenosine
deaminase deficiency

[185266]in severe combined immunodeficiency: evidence for a
posttranslational

[185267]defect. (Abstract) J. Clin. Invest. 53: 81A–82A, 1974.

[185268]

[185269]117. Wakamiya, M.; Blackburn, M. R.; Jurecic, R.;
McArthur, M. J.;

[185270]Geske, R. S.; Cartwright, J., Jr.; Mitani, K.; Vaishnav, S.;
Belmont,

[185271]J. W.; Kellems, R. E.; Finegold, M. J.; Montgomery, Jr., C.
A.; Bradley,

[185272]A.; Caskey, C. T.: Disruption of the adenosine deaminase gene causes

[185273]hepatocellular impairment and perinatal lethality in mice. Proc.

[185274]Nat. Acad. Sci. 92: 3673–3677, 1995.

[185275]

[185276]118. Weitkamp, L. R.: Further data on the genetic linkage relations

[185277]of the adenosine deaminase locus. Hum. Hered. 21: 351–356, 1971.

[185278]

[185279]119. Weitkamp, L. R.: Genetic linkage relationships of the ADA and

[185280]6–PGD loci in 'Humangenetik.' (Letter) Humangenetik 15: 359–360,

[185281]1972.

[185282]

[185283]120. Wiginton, D. A.; Adrian, G. S.; Friedman, R. L.; Suttle, D. P.;

[185284]Hutton, J. J.: Cloning of cDNA sequences of human adenosine deaminase. Proc.

[185285]Nat. Acad. Sci. 80: 7481–7485, 1983.

[185286]

[185287]121. Wiginton, D. A.; Adrian, G. S.; Hutton, J. J.: Sequence
of human

[185288]adenosine deaminase cDNA including the coding region
and a small intron. Nucleic

[185289]Acids Res. 12: 2439–2446, 1984.

[185290]

[185291]122. Wiginton, D. A.; Hutton, J. J.: Immunoreactive protein
in adenosine

[185292]deaminase deficient human lymphoblast cell lines. J. Biol.
Chem. 257:

[185293]3211–3217, 1982.

[185294]

[185295]123. Wiginton, D. A.; Kaplan, D. J.; States, J. C.; Akeson, A.
L.;

[185296]Perme, C. M.; Bilyk, I. J.; Vaughn, A. J.; Lattier, D. L.; Hut-
ton,

[185297]J. J.: Complete sequence and structure of the gene for hu-
man adenosine

[185298]deaminase. Biochemistry 25: 8234–8244, 1986.

[185299]

[185300]124. Yokoyama, S.; Hayashi, T.; Yoshimura, Y.; Irimada, K.; Saito,

[185301]T.; Akiba, T.; Tsuchiya, S.: Severe combined immunodeficiency disease

[185302]with adenosine deaminase deficiency. Tohoku J. Exp. Med. 129: 197–202,

[185303]1979.

[185304]

[185305]125. Yount, J.; Nichols, P.; Ochs, H. D.; Hammar, S. P.; Scott, C.

[185306]R.; Chen, S.–H.; Giblett, E. R.; Wedgwood, R. J.: Absence of erythrocyte

[185307]adenosine deaminase associated with severe combined immunodeficiency. J.

[185308]Pediat. 84: 173–177, 1974.

[185309]

[185310]126. Ziegler, J. B.; Lee, C. H.; Van Der Weyden, M. B.; Bag-nara, A.

[185311]S.; Beveridge, J.: Severe combined immunodeficiency and adenosine

[185312]deaminase deficiency: failure of enzyme replacement therapy. Arch.

[185313]Dis. Child. 55: 452–457, 1980.

[185314]

[185315]127. Ziegler, J. B.; Van Der Weyden, M. B.; Lee, C. H.; Daniel, A.

[185316]: Prenatal diagnosis for adenosine deaminase deficiency. J. Med.

[185317]Genet. 18: 154–156, 1981.

[185318]

[185319]

[185320]19417.

[185321]Sikela, J. M.; Adamson, M. C.; Wilson–Shaw, D.; Kozak, C. A.:

[185322]Genetic mapping of the gene for Ca(2+)/calmodulin–dependent protein

[185323]kinase IV (Camk–4) to mouse chromosome 18. Genomics 8: 579–582,

[185324]1990.

[185325]

[185326]

[185327]19418.

[185328]Sikela, J. M.; Law, M. L.; Kao, F.-T.; Hartz, J. A.; Wei, Q.;
Hahn,

[185329]W. E.: Chromosomal localization of the human gene for
brain Ca(2+)/calmodulin-dependent

[185330]protein kinase type IV. Genomics 4: 21-27, 1989.

[185331]

[185332]

[185333]19419.

[185334]Wei, F.; Qiu, C.-S.; Liauw, J.; Robinson, D. A.; Ho, N.;
Chatila,

[185335]T.; Zhuo, M.: Calcium-calmodulin-dependent protein ki-
nase IV is required

[185336]for fear memory. Nature Neurosci. 5: 573-579, 2002.

[185337]

[185338]

[185339]19420.

[185340]Wu, H.; Kanatous, S. B.; Thurmond, F. A.; Gallardo, T.;

Isotani,

[185341]E.; Bassel-Duby, R.; Williams, R. S.: Regulation of mitochondrial

[185342]biogenesis in skeletal muscle by CaMK. Science 296:
349–352, 2002.

[185343]

[185344]

[185345]19421.

[185346]Wu, J. Y.; Ribar, T. J.; Cummings, D. E.; Burton, K. A.; McKnight,

[185347]G. S.; Means, A. R.: Spermiogenesis and exchange of basic nuclear

[185348]proteins are impaired in male germ cells lacking Camk4.
Nature Genet. 25:

[185349]448–452, 2000.

[185350]

[185351]

[185352]19422.

[185353]Dooley, T. P.: Personal Communication. San Antonio, Tex.
7/21/1992.

[185354]

[185355]

[185356]19423.

[185357]Dooley, T. P.; Weiland, K. L.; Simon, M.: cDNA sequence of
human

[185358]p11 calpactin I light chain. Genomics 13: 866–868, 1992.

[185359]

[185360]

[185361]19424.

[185362]Harder, T.; Kube, E.; Gerke, V.: Cloning and characteriza-
tion

[185363]of the human gene encoding p11: structural similarity to
other members

[185364]of the S-100 gene family. Gene 113: 269–274, 1992.

[185365]

[185366]

[185367]19425.

[185368]Kube, E.; Weber, K.; Gerke, V.: Primary structure of human,
chicken,

[185369]and *Xenopus laevis* p11, a cellular ligand of the Src-kinase

substrate,

[185370]annexin II. Gene 102: 255–259, 1991.

[185371]

[185372]

[185373]19426.

[185374]Okuse, K.; Malik–Hall, M.; Baker, M. D.; Poon, W.–Y. L.;

Kong,

[185375]H.; Chao, M. V.; Wood, J. N.: Annexin II light chain regulates sensory

[185376]neuron–specific sodium channel expression. Nature 417: 653–656,

[185377]2002.

[185378]

[185379]

[185380]19427.

[185381]Saris, C. J. M.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.;

[185382]Noonan, D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and tissue

[185383]distribution of the mRNA for bovine and murine p11, the S100–related

[185384]light chain of the protein-tyrosine kinase substrate p36
(calpactin

[185385]l). J. Biol. Chem. 262: 10663–10671, 1987.

[185386]

[185387]

[185388]19428.

[185389]Inazawa, J.; Nakagawa, H.; Misawa, S.; Abe, T.; Minoshima,
S.;

[185390]Fukuyama, R.; Maki, M.; Murachi, T.; Hatanaka, M.;
Shimizu, N.: Assignment

[185391]of the human calpastatin gene (CAST) to chromosome 5 at
region q14–q22.

[185392]Cytogenet. Cell Genet. 54: 156–158, 1990.

[185393]

[185394]

[185395]19429.

[185396]Inazawa, J.; Nakagawa, H.; Misawa, S.; Abe, T.; Minoshima,
S.;

[185397]Fukuyama, R.; Maki, M.; Murachi, T.; Hatanaka, M.;
Shimizu, N.: Assignment

[185398]of the human calpastatin gene (CAST) to chromosome 5 at
region q15–q21.

[185399](Abstract) Cytogenet. Cell Genet. 58: 1898 only, 1991.

[185400]

[185401]

[185402]19430.

[185403]Mimori, T.; Suganuma, K.; Tanami, Y.; Nojima, T.; Mat-
sumura, M.;

[185404]Fujii, T.; Yoshizawa, T.; Suzuki, K.; Akizuki, M.: Autoanti-
bodies

[185405]to calpastatin (an endogenous inhibitor for calcium-
dependent neutral

[185406]protease, calpain) in systemic rheumatic diseases. Proc.
Nat. Acad.

[185407]Sci. 92: 7267–7271, 1995.

[185408]

[185409]

[185410]19431.

[185411]Pontremoli, S.; Salamino, F.; Sparatore, B.; De Tullio, R.;
Pontremoli,

[185412]R.; Melloni, E.: Characterization of the calpastatin defect in erythrocytes

[185413]from patients with essential hypertension. Biochem. Biophys. Res.

[185414]Commun. 157: 867–874, 1988.

[185415]

[185416]

[185417]19432.

[185418]Matthijs, G.; Schollen, E.; Pirard, M.; Budarf, M. L.; Van Schaftingen,

[185419]E.; Cassiman, J.-J.: PMM (PMM1), the human homologue of SEC53 or

[185420]yeast phosphomannomutase, is localized on chromosome 22q13. Genomics 40:

[185421]41–47, 1997.

[185422]

[185423]

[185424]19433.

[185425]Carritt, B.: Somatic cell genetic evidence for the presence of

[185426]a gene for citrullinemia on human chromosome 9. Cyto-
genet. Cell Genet. 19:

[185427]44–48, 1977.

[185428]

[185429]

[185430]19434.

[185431]Carritt, B.; Goldfarb, P. S. G.; Hooper, M. L.; Slack, C.:
Chromosome

[185432]assignment of a human gene for argininosuccinate syn-
thetase expression

[185433]in Chinese hamster–human somatic cell hybrids. Exp. Cell
Res. 106:

[185434]71–78, 1977.

[185435]

[185436]

[185437]19435.

[185438]Daiger, S. P.; Hoffman, N. S.; Wildin, R. S.; Su, T.–S.: Multi-
ple,

[185439]independent restriction site polymorphisms in human
DNA detected with

[185440]a cDNA probe to argininosuccinate synthetase (AS). Am. J. Hum. Genet. 36:

[185441]736–749, 1984.

[185442]

[185443]

[185444]19436.

[185445]Daiger, S. P.; Wildin, R. S.; Su, T.–S.: Polymorphic variants

[185446]of restriction fragments of human DNA detected with a probe to argininosuccinate

[185447]synthetase. (Abstract) Am. J. Hum. Genet. 33: 136A only, 1981.

[185448]

[185449]

[185450]19437.

[185451]Elshourbagy, N. A.; Walker, D. W.; Boguski, M. S.; Gordon, J. I.;

[185452]Taylor, J. M.: The nucleotide and derived amino acid sequence of

[185453]human apolipoprotein A–IV mRNA and the close linkage of its gene to

[185454]the genes of apolipoproteins A-I and C-III. J. Biol. Chem.

261:

[185455]1998–2002, 1986.

[185456]

[185457]

[185458]19438.

[185459]Elshourbagy, N. A.; Walker, D. W.; Paik, Y.-K.; Boguski, M.
S.;

[185460]Freeman, M.; Gordon, J. I.; Taylor, J. M.: Structure and ex-
pression

[185461]of the human apolipoprotein A-IV gene. J. Biol. Chem.
262: 7973–7981,

[185462]1987.

[185463]

[185464]

[185465]19439.

[185466]Green, P. H. R.; Glickman, R. M.; Riley, J. W.; Quinet, E.:
Human

[185467]apolipoprotein A-IV: intestinal origin and distribution in
plasma. J.

[185468]Clin. Invest. 65: 911–919, 1980.

[185469]

[185470]

[185471]19440.

[185472]Kamboh, M. I.; Ferrell, R. E.: Genetic studies of human
apolipoproteins.

[185473]I. Polymorphism of apolipoprotein A–IV. Am. J. Hum.
Genet. 41: 119–127,

[185474]1987.

[185475]

[185476]

[185477]19441.

[185478]Kamboh, M. I.; Williams, E. R.; Law, J. C.; Aston, C. E.;
Bunker,

[185479]C. H.; Ferrell. R. E.; Pollitzer, W. S.: Molecular basis of a
unique

[185480]African variant (A-IV 5) of human apolipoprotein A-IV and
its significance

[185481]in lipid metabolism. Genet. Epidemiol. 9: 379-388, 1992.

[185482]

[185483]

[185484]19442.

[185485]Karathanasis, S. K.: Apolipoprotein multigene family: tan-
dem organization

[185486]of human apolipoprotein AI, CIII, and AIV genes. Proc. Nat.
Acad.

[185487]Sci. 82: 6374-6378, 1985.

[185488]

[185489]

[185490]19443.

[185491]Lohse, P.; Kindt, M. R.; Rader, D. J.; Brewer, H. B., Jr.: Ge-

netic

[185492]polymorphism of human plasma apolipoprotein A-IV is
due to nucleotide

[185493]substitutions in the apolipoprotein A-IV gene. J. Biol.
Chem. 265:

[185494]10061-10064, 1990.

[185495]

[185496]

[185497]19444.

[185498]Lohse, P.; Kindt, M. R.; Rader, D. J.; Brewer, H. B., Jr.: Hu-
man

[185499]plasma apolipoproteins A-IV-0 and A-IV-3: molecular ba-
sis for two

[185500]rare variants of apolipoprotein A-IV-1. J. Biol. Chem. 265:
12734-12739,

[185501]1990.

[185502]

[185503]

[185504]19445.

[185505]Menzel, H.-J.; Boerwinkle, E.; Schrangl-Will, S.; Utermann,

G.

[185506]: Human apolipoprotein A-IV polymorphism: frequency and effect on

[185507]lipid and lipoprotein levels. Hum. Genet. 79: 368-372, 1988.

[185508]

[185509]

[185510]19446.

[185511]Menzel, H.-J.; Kovary, P. M.; Assmann, G.: Apolipoprotein A-IV

[185512]polymorphism in man. Hum. Genet. 62: 349-352, 1982.

[185513]

[185514]

[185515]19447.

[185516]Menzel, H.-J.; Sigurdsson, G.; Boerwinkle, E.; Schrangl-Will,

[185517]S.; Dieplinger, H.; Utermann, G.: Frequency and effect of human apolipoprotein

[185518]A-IV polymorphism on lipid and lipoprotein levels in an Icelandic

[185519]population. Hum. Genet. 84: 344-346, 1990.

[185520]

[185521]

[185522]19448.

[185523]Rogne, S.; Myklebost, O.; Olaisen, B.; Gedde-Dahl, T., Jr.;
Prydz,

[185524]H.: Confirmation of the close linkage between the loci for
human

[185525]apolipoproteins AI and AIV by the use of a cloned cDNA
probe and two

[185526]restriction site polymorphisms. Hum. Genet. 72: 68–71,
1986.

[185527]

[185528]

[185529]19449.

[185530]Schamaun, O.; Olaisen, B.; Mevag, B.; Gedde-Dahl, T., Jr.;
Ehnholm,

[185531]C.; Teisberg, P.: The two apolipoprotein loci apoA-I and
apoA-IV

[185532]are closely linked in man. Hum. Genet. 68: 181–184,
1984.

[185533]

[185534]

[185535]19450.

[185536]Tracy, R. P.; Currie, R. M.; Young, D. S.: Two-dimensional
gel

[185537]electrophoresis of serum specimens from a normal population. Clin.

[185538]Chem. 28: 890–899, 1982.

[185539]

[185540]

[185541]19451.

[185542]von Eckardstein, A.; Funke, H.; Schulte, M.; Erren, M.;
Schulte,

[185543]H.; Assmann, G.: Nonsynonymous polymorphic sites in the
apolipoprotein

[185544](apo) A–IV gene are associated with changes in the concentration of

[185545]apo B– and apo A–I–containing lipoproteins in a normal
population. Am.

[185546]J. Hum. Genet. 50: 1115–1128, 1992.

[185547]

[185548]

[185549]19452.

[185550]Gautier, T.; Masson, D.; Jong, M. C.; Duverneuil, L.; Le
Guern,

[185551]N.; Deckert, V.; Pais de Barros, J.-P.; Dumont, L.; Bataille,
A.;

[185552]Zak, Z.; Jiang, X.-C.; Tall, A. R.; Havekes, L. M.; Lagrost, L.:

[185553]Apolipoprotein CI deficiency markedly augments plasma
lipoprotein

[185554]changes mediated by human cholesteryl ester transfer
protein (CETP)

[185555]in CETP transgenic/ApoCI-knocked out mice. J. Biol.
Chem. 277: 31354–31363,

[185556]2002.

[185557]

[185558]

[185559]19453.

[185560]Brown, S.; Wiebel, F. J.; Gelboin, H. V.; Minna, J. D.: As-
signment

[185561]of a locus required for flavoprotein-linked monooxyge-
nase expression

[185562]to human chromosome 2. Proc. Nat. Acad. Sci. 73:
4628–4632, 1976.

[185563]

[185564]

[185565]19454.

[185566]Chen, Y. T.; Tukey, R. H.; Swan, D. C.; Negishi, N.; Nebert,
D.

[185567]W.: Characterization of the human P1–450 genomic gene.
(Abstract) Clin.

[185568]Res. 31: 456A, 1983.

[185569]

[185570]

[185571]19455.

[185572]Corchero, J.; Pimprale, S.; Kimura, S.; Gonzalez, F. J.: Or-
ganization

[185573]of the CYP1A cluster on human chromosome 15: implica-
tions for gene

[185574]regulation. Pharmacogenetics 11: 1–6, 2001.

[185575]

[185576]

[185577]19456.

[185578]Gorman, C.; Padmanabhan, R.; Howard, B. H.: High efficiency DNA-mediated

[185579]transformation of primate cells. Science 221: 551–553, 1983.

[185580]

[185581]

[185582]19457.

[185583]Gorman, C. M.: CAT: an easy assay for gene expression (citation

[185584]classic). Current Contents (Life Sciences) 36(22): 8, 1993.

[185585]

[185586]

[185587]19458.

[185588]Hildebrand, C. E.; Gonzalez, F. J.; Kozak, C. A.; Nebert, D. W.

[185589]: Regional linkage analysis of the dioxin-inducible P-450 gene family

[185590]on mouse chromosome 9. Biochem. Biophys. Res. Commun. 130: 396–406,

[185591]1985.

[185592]

[185593]

[185594]19459.

[185595]Lee, S.; Wu, X.; Reid, M. E.; Zelinski, T.; Redman, C. M.:

Molecular

[185596]basis of the Kell (K1) phenotype. Blood 85: 912–916,

1995.

[185597]

[185598]

[185599]19460.

[185600]Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.: The

human

[185601]Kell blood group gene maps to chromosome 7q33 and its

expression is

[185602]restricted to erythroid cells. Blood 81: 2804–2809, 1993.

[185603]

[185604]

[185605]19461.

[185606]Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.:

Molecular

[185607]cloning and primary structure of Kell blood group protein.

Proc.

[185608]Nat. Acad. Sci. 88: 6353–6357, 1991.

[185609]

[185610]

[185611]19462.

[185612]Marsh, W. L.: Molecular biology of blood groups: cloning
of the

[185613]Kell gene. Transfusion 32: 98–101, 1992.

[185614]

[185615]

[185616]19463.

[185617]Morton, N. E.; Krieger, H.; Steinberg, A. G.; Rosenfield, R.
E.

[185618]: Genetic evidence confirming the localization of Sutter in
the Kell

[185619]blood-group system. Vox Sang. 10: 608–613, 1965.

[185620]

[185621]

[185622]19464.

[185623]Murphy, M. T.; Morrison, N.; Miles, J. S.; Fraser, R. H.;
Spurr,

[185624]N. K.; Boyd, E.: Regional chromosomal assignment of the
Kell blood

[185625]group locus (KEL) to chromosome 7q33-q35 by fluores-
cence in situ hybridization:

[185626]evidence for the polypeptide nature of antigenic variation.
Hum.

[185627]Genet. 91: 585-588, 1993.

[185628]

[185629]

[185630]19465.

[185631]Parke, J. T.; Riccardi, V. M.; Lewis, R. A.; Ferrell, R. E.:

[185632]A syndrome of microcephaly and retinal pigmentary ab-
normalities without

[185633]mental retardation in a family with coincidental autosomal
dominant

[185634]hyperreflexia. Am. J. Med. Genet. 17: 585-594, 1984.

[185635]

[185636]

[185637]19466.

[185638]Purohit, K. R.; Weber, J. L.; Ward, L. J.; Keats, B. J. B.: The

[185639]Kell blood group locus is close to the cystic fibrosis locus
on chromosome

[185640]7. Hum. Genet. 89: 457–458, 1992.

[185641]

[185642]

[185643]19467.

[185644]Russo, D.; Wu, X.; Redman, C. M.; Lee, S.: Expression of
Kell

[185645]blood group protein in nonerythroid tissues Blood 96:
340–346, 2000.

[185646]

[185647]

[185648]19468.

[185649]Stroup, M.; MacIlroy, M.; Walker, R.; Aydelotte, J. V.: Evi-
dence

[185650]that Sutter belongs to the Kell blood group system. Trans-
fusion 5:

[185651]309–314, 1965.

[185652]

[185653]

[185654]19469.

[185655]Wagner, T.; Berer, A.; Lanzer, G.; Geissler, K.: Kell is not

[185656]restricted to the erythropoietic lineage but is also ex-
pressed on

[185657]myeloid progenitor cells. Brit. J. Haemat. 110: 409–411,
2000.

[185658]

[185659]

[185660]19470.

[185661]Weitkamp, L. R.; Townes, P. L.; Johnston, E.: Linkage data
on

[185662]urinary pepsinogen and the Kell blood group. Birth De-
fects Orig.

[185663]Art. Ser. 11(3): 281–282, 1975. Note: Alternate: Cyto-
genet. Cell

[185664]Genet. 14: 451–452, 1975.

[185665]

[185666]

[185667]19471.

[185668]Wimer, B. M.; Marsh, W. L.; Taswell, H. F.: Clinical characteristics

[185669]of the McLeod blood group phenotype. (Abstract) 19th Annual Meeting

[185670]of the Am. Soc. Hemat., Boston , 1976.

[185671]

[185672]

[185673]19472.

[185674]Yu, L.–C.; Twu, Y.–C.; Chang, C.–Y.; Lin, M.: Molecular basis

[185675]of the Kell–null phenotype: a mutation at the splice site of human

[185676]KEL gene abolishes the expression of Kell blood group antigens. J.

[185677]Biol. Chem. 276: 10247–10252, 2001.

[185678]

[185679]

[185680]19473.

[185681]Zelinski, T. A.; Coghlan, G. E.; Myal, Y.; White, L. J.;

Philipps,

[185682]S. E.: Assignment of the Kell blood group locus to chromosome 7q.

[185683](Abstract) Cytogenet. Cell Genet. 58: 1927 only, 1991.

[185684]

[185685]

[185686]19474.

[185687]Field, L. L.; Marazita, M. L.; Spence, M. A.; Crandall, B. F.;

[185688]Sparkes, R. S.: Is JK linked to IGK on chromosome 2?

(Abstract) Cytogenet.

[185689]Cell Genet. 40: 628–629, 1985.

[185690]

[185691]

[185692]19475.

[185693]Gedde–Dahl, T.: Personal Communication. Oslo, Norway
9/26/1986.

[185694]

[185695]

[185696]19476.

[185697]Geitvik, G. A.; Hoyheim, B.; Gedde-Dahl, T.; Grzeschik, K.
H.;

[185698]Lothe, R.; Tomter, H.; Olaisen, B.: The Kidd (JK) blood
group locus

[185699]assigned to chromosome 18 by close linkage to a DNA-
RFLP. Hum. Genet. 77:

[185700]205-209, 1987.

[185701]

[185702]

[185703]19477.

[185704]Hulten, M.; Lindsten, J.; Pen-Ming, L. M.; Fraccaro, M.;
Mannini,

[185705]A.; Trepolo, L.; Robson, E. B.; Heiken, A.; Tellingén, K. G.:
Possible

[185706]localization of the genes for the Kidd blood group on an
autosome

[185707]involved in a reciprocal translocation. Nature 211:
1067-1068, 1968.

[185708]

[185709]

[185710]19478.

[185711]Leppert, M.; Ferrell, R.; Kamboh, M. I.; Beasley, J.;

O'Connell,

[185712]P.; Lathrop, M.; Lalouel, J.-M.; White, R.: Linkage of the
polymorphic

[185713]protein markers F13B, C1S, C1R, and blood group antigen
Kidd in CEPH

[185714]reference families. (Abstract) Cytogenet. Cell Genet. 46:
647, 1987.

[185715]

[185716]

[185717]19479.

[185718]Lucien, N.; Chiaroni, J.; Cartron, J.-P.; Bailly, P.: Partial

[185719]deletion in the JK locus causing a Jk(null) phenotype.

Blood 99:

[185720]1079-1081, 2002.

[185721]

[185722]

[185723]19480.

[185724]Lucien, N.; Sidoux–Walter, F.; Olives, B.; Moulds, J.; Le
Pennec,

[185725]P.–Y.; Cartron, J.–P.; Bailly, P.: Characterization of the gene
encoding

[185726]the human Kidd blood group/urea transporter protein:
evidence for

[185727]splice site mutations in Jk–null individuals. J. Biol. Chem.
273:

[185728]12973–12980, 1998.

[185729]

[185730]

[185731]19481.

[185732]Olives, B.; Mattei, M.–G.; Huet, M.; Neau, P.; Martial, S.;
Cartron,

[185733]J.–P.; Bailly, P.: Kidd blood group and urea transport func-
tion of

[185734]human erythrocytes are carried by the same protein. J.
Biol. Chem. 270:

[185735]15607–15610, 1995.

[185736]

[185737]

[185738]19482.

[185739]Muramatsu, T.; Zu-Cheng, W.; Yi-Ru, F.; Kou-Bao, H.;
Heqin, Y.;

[185740]Yamada, K.; Higuchi, S.; Harada, S.; Kono, H.: Alcohol and
aldehyde

[185741]dehydrogenase genotypes and drinking behavior of Chi-
nese living in

[185742]Shanghai. Hum. Genet. 96: 151-154, 1995.

[185743]

[185744]

[185745]19483.

[185746]Womack, J. E.: Personal Communication. College Station,
Texas

[185747]2/26/1990.

[185748]

[185749]

[185750]19484.

[185751]Hiraoka, L. R.; Hsu, L.; Hsieh, C.-L.: Assignment of ALDH3

to

[185752]human chromosome 17p11.2 and ALDH5 to human chromosome 9p13. Genomics 25:

[185753]323–325, 1995.

[185754]

[185755]

[185756]19485.

[185757]Hsu, L. C.; Chang, W.-C.; Shibuya, A.; Yoshida, A.: Human stomach

[185758]aldehyde dehydrogenase cDNA and genomic cloning, primary structure,

[185759]and expression in Escherichia coli. J. Biol. Chem. 267: 3030–3037,

[185760]1992.

[185761]

[185762]

[185763]19486.

[185764]Kays, W. T.; Piatigorsky, J.: Aldehyde dehydrogenase class 3 expression:

[185765]identification of a cornea-preferred gene promoter in

transgenic mice. Proc.

[185766]Nat. Acad. Sci. 94: 13594–13599, 1997.

[185767]

[185768]

[185769]19487.

[185770]Rogers, G. R.; Markova, N. G.; De Laurenzi, V.; Rizzo, W.
B.; Compton,

[185771]J. G.: Genomic organization and expression of the human
fatty aldehyde

[185772]dehydrogenase gene (FALDH). Genomics 39: 127–135,
1997.

[185773]

[185774]

[185775]19488.

[185776]Santisteban, I.; Povey, S.; West, L. F.; Parrington, J. M.;
Hopkinson,

[185777]D. A.: Chromosome assignment, biochemical and im-
munological studies

[185778]on a human aldehyde dehydrogenase, ALDH3. Ann. Hum.
Genet. 49: 87–100,

[185779]1985.

[185780]

[185781]

[185782]19489.

[185783]Teng, Y.-S.: Stomach aldehyde dehydrogenase: report of a new locus. Hum.

[185784]Hered. 31: 74-77, 1981.

[185785]

[185786]

[185787]19490.

[185788]Vasiliou, V.; Bairoch, A.; Tipton, K. F.; Nebert, D. W.: Eukaryotic

[185789]aldehyde dehydrogenase (ALDH) genes: human polymorphisms, and recommended

[185790]nomenclature based on divergent evolution and chromosomal mapping. Pharmacogenetics 9:

[185791]421-434, 1999.

[185792]

[185793]

[185794]19491.

[185795]Wunderle, V. M.; Critcher, R.; Hastie, N.; Goodfellow, P. N.;

[185796]Schedl, A.: Deletion of long-range regulatory elements
upstream of

[185797]SOX9 causes campomelic dysplasia. Proc. Nat. Acad. Sci.
95: 10649–10654,

[185798]1998.

[185799]

[185800]

[185801]19492.

[185802]Young, I. D.; Zuccollo, J. M.; Maltby, E. L.; Broderick, N. J.

[185803]: Campomelic dysplasia associated with a de novo 2q;17q
reciprocal

[185804]translocation. J. Med. Genet. 29: 251–252, 1992.

[185805]

[185806]

[185807]19493.

[185808]Fornerod, M.; van Duersen, J.; van Baal, S.; Reynolds, A.;
Davis,

[185809]D.; Murti, K. G.; Fransen, J.; Grosveld, G.: The human ho–

mologue

[185810]of yeast CRM1 is in a dynamic subcomplex with CAN/
Nup214 and a novel

[185811]nuclear pore component Nup88. EMBO J. 16: 807–816,
1997.

[185812]

[185813]

[185814]19494.

[185815]Kraemer, D.; Wozniak, R. W.; Blobel, G.; Radu, A.: The hu-
man CAN

[185816]protein, a putative oncogene product associated with
myeloid leukemogenesis,

[185817]is a nuclear pore complex protein that faces the cyto-
plasm. Proc.

[185818]Nat. Acad. Sci. 91: 1519–1523, 1994.

[185819]

[185820]

[185821]19495.

[185822]van Deursen, J.; Boer, J.; Kasper, L.; Grosveld, G.: G(2) ar-
rest

[185823]and impaired nucleocytoplasmic transport in mouse em-

bryos lacking

[185824]the proto-oncogene CAN/Nup214. EMBO J. 15:
5574–5583, 1996.

[185825]

[185826]

[185827]19496.

[185828]von Lindern, M.; Poustka, A.; Lehrach, H.; Grosveld, G.:
The (6;9)

[185829]chromosome translocation, associated with a specific
subtype of acute

[185830]nonlymphocytic leukemia, leads to aberrant transcription
of a target

[185831]gene on 9q34. Molec. Cell. Biol. 10: 4016–4026, 1990.

[185832]

[185833]

[185834]19497.

[185835]Lynch, H. T.; Schuelke, G. S.; Kimberling, W. J.; Albano, W.
A.;

[185836]Lynch, J. F.; Biscione, K. A.; Lipkin, M. L.; Deschner, E. E.;
Mikol,

[185837]Y. B.; Sandberg, A. A.; Elston, R. C.; Bailey–Wilson, J. E.;

Danes,

[185838]B. S.: Hereditary nonpolyposis colorectal cancer (Lynch syndromes

[185839]I and II). II. Biomarker studies. Cancer 56: 939–951, 1985.

[185840]

[185841]

[185842]19498.

[185843]Hanauer, A.; Heilig, R.; Levin, M.; Moisan, J. P.; Grzeschik, K.

[185844]H.; Mandel, J. L.: The actin gene family in man: assignment of the

[185845]gene for skeletal muscle alpha-actin to chromosome 1, and presence

[185846]of actin sequences on autosomes 2 and 3, and on the X and Y chromosomes.

[185847](Abstract) Cytogenet. Cell Genet. 37: 487–488, 1984.

[185848]

[185849]

[185850]19499.

[185851]Ilkovski, B.; Cooper, S. T.; Nowak, K.; Ryan, M. M.; Yang,

N.;

[185852]Schnell, C.; Durling, H. J.; Roddick, L. G.; Wilkinson, I.; Kornberg,

[185853]A. J.; Collins, K. J.; Wallace, G.; Gunning, P.; Hardeman, E. C.;

[185854]Laing, N. G.; North, K. N.: Nemaline myopathy caused by mutations

[185855]in the muscle alpha-skeletal-actin gene. Am. J. Hum. Genet. 68:

[185856]1333–1343, 2001.

[185857]

[185858]

[185859]19500.

[185860]Nowak, K. J.; Wattanasirichaigoon, D.; Goebel, H. H.; Wilce, M.;

[185861]Pelin, K.; Donner, K.; Jacob, R. L.; Hubner, C.; Oexle, K.; Anderson,

[185862]J. R.; Verity, C. M.; North, K. N.; and 13 others: Mutations in

[185863]the skeletal muscle alpha-actin gene in patients with actin myopathy

[185864]and nemaline myopathy. Nature Genet. 23: 208–212,

1999.

[185865]

[185866]

[185867]19501.

[185868]Kedes, L.; Ng, S.-Y.; Lin, C.-S.; Gunning, P.; Eddy, R.;

Shows,

[185869]T.; Leavitt, J.: The human beta-actin multigene family.

Trans. Assoc.

[185870]Am. Phys. 98: 42-46, 1985.

[185871]

[185872]

[185873]19502.

[185874]Leavitt, J.; Bushar, G.; Kakunaga, T.; Hamada, H.; Hi-

rakawa, T.;

[185875]Goldman, D.; Merrill, C.: Variations in expression of mu-

tant beta-actin

[185876]accompanying incremental increases in human fibroblast

tumorigenicity. Cell 28:

[185877]259-268, 1982.

[185878]

[185879]

[185880]19503.

[185881]Nakajima-Iijima, S.; Hamada, H.; Reddy, P.; Kakunaga, T.:

Molecular

[185882]structure of the human cytoplasmic beta-actin gene; interspecies homology

[185883]of sequences in the introns. Proc. Nat. Acad. Sci. 82:
6133-6137,

[185884]1985.

[185885]

[185886]

[185887]19504.

[185888]Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.;
Kedes,

[185889]L.; Shows, T.: Chromosome 7 assignment of the human
beta-actin functional

[185890]gene (ACTB) and the chromosomal dispersion of pseudogenes. (Abstract) Cytogenet.

[185891]Cell Genet. 40: 712 only, 1985.

[185892]

[185893]

[185894]19505.

[185895]Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.;

Shows,

[185896]T.; Kedes, L.: Evolution of the functional human beta-actin
gene

[185897]and its multi-pseudogene family: conservation of the
noncoding regions

[185898]and chromosomal dispersion of pseudogenes. Molec. Cell.
Biol. 5:

[185899]2720-2732, 1985.

[185900]

[185901]

[185902]19506.

[185903]Toyama, S.; Toyama, S.: A variant form of beta-actin in a
mutant

[185904]of KB cells resistant to cytochalasin B. Cell 37: 609-614,
1984.

[185905]

[185906]

[185907]19507.

[185908]Vandekerckhove, J.; Weber, K.: Mammalian cytoplasmic
actins are

[185909]the products of at least two genes and differ in primary
structure

[185910]in at least 25 identified positions from skeletal muscle
actins. Proc.

[185911]Nat. Acad. Sci. 75: 1106–1110, 1978.

[185912]

[185913]

[185914]19508.

[185915]Allderdice, P. W.; Kaita, H.; Lewis, M.; McAlpine, P. J.;
Wong,

[185916]P.; Anderson, J.; Giblett, E. R.: Segregation of marker loci
in families

[185917]with an inherited paracentric insertion of chromosome 9.
Am. J. Hum.

[185918]Genet. 39: 612–617, 1986.

[185919]

[185920]

[185921]19509.

[185922]Dayhoff, M. O.: Atlas of Protein Sequence and Structure.
Dehydrogenases.

[185923]Washington: National Biomedical Research Foundation
(pub.) 5: 1972.

[185924]Pp. D141–D144.

[185925]

[185926]

[185927]19510.

[185928]Edenberg, H. J.: Regulation of the mammalian alcohol de-
hydrogenase

[185929]genes. Prog. Nucleic Acids Res. Molec. Biol. 64: 295–341,
2000.

[185930]

[185931]

[185932]19511.

[185933]Osier, M. V.; Pakstis, A. J.; Soodyall, H.; Comas, D.; Gold-
man,

[185934]D.; Odunsi, A.; Okonofua, F.; Parnas, J.; Schulz, L. O.;
Bertranpetit,

[185935]J.; Bonne-Tamir, B.; Lu, R.-B.; Kidd, J. R.; Kidd, K. K.: A
global

[185936]perspective on genetic variation at the ADH genes reveals
unusual

[185937]patterns of linkage disequilibrium and diversity. Am. J.
Hum. Genet. 71:

[185938]84–99, 2002.

[185939]

[185940]

[185941]19512.

[185942]Smith, M.; Hopkinson, D. A.; Harris, H.: Studies on the
subunit

[185943]structure and molecular size of the human dehydrogenase
isozymes determined

[185944]by the different loci, ADH(1), ADH(2), and ADH(3). Ann.
Hum. Genet. 36:

[185945]401–414, 1973.

[185946]

[185947]

[185948]19513.

[185949]Comings, D. E.; Gade-Andavolu, R.; Gonzalez, N.; Blake, H.; Wu,

[185950]S.; MacMurray, J. P.: Additive effect of three noradrenergic genes

[185951](ADRA2A, ADRA2C, DBH) on attention-deficit hyperactivity disorder

[185952]and learning disabilities in Tourette syndrome subjects. Clin. Genet. 55:

[185953]160-172, 1999.

[185954]

[185955]

[185956]19514.

[185957]Halperin, J. M.; Newcorn, J. H.; Koda, V. H.; Pick, L.; McKay,

[185958]K. E.; Knott, P.: Noradrenergic mechanisms in ADHD children with

[185959]and without reading disabilities: a replication and extension. J.

[185960]Am. Acad. Child Adolesc. Psychiat. 36: 1688-1697, 1997.

[185961]

[185962]

[185963]19515.

[185964]Hein, L.; Altman, J. D.; Kobilka, B. K.: Two functionally distinct
tinct

[185965]alpha-2-adrenergic receptors regulate sympathetic neurotransmission. Nature 402:

[185966]181-184, 1999.

[185967]

[185968]

[185969]19516.

[185970]Hoehe, M.; Berrettini, W.; Leppert, M.; Lalouel, J.-M.; Byerley,
ley,

[185971]W.; Gershon, E.; White, R.: Genetic mapping of adrenergic
receptor

[185972]genes. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A143
only, 1989.

[185973]

[185974]

[185975]19517.

[185976]Hoehe, M. R.; Berrettini, W. H.; Lentes, K.-U.: Draft identifies
fies

[185977]a two allele DNA polymorphism in the human alpha-
2-adrenergic receptor

[185978]gene (ADRAR), using a 5.5 kb probe (p ADRAR). Nucleic
Acids Res. 16:

[185979]9070 only, 1988.

[185980]

[185981]

[185982]19518.

[185983]Kobilka, B. K.; Matsui, H.; Kobilka, T. S.; Yang-Feng, T. L.;
Francke,

[185984]U.; Caron, M. G.; Lefkowitz, R. J.; Regan, J. W.: Cloning, se-
quencing,

[185985]and expression of the gene coding for the human platelet
alpha-2-adrenergic

[185986]receptor. Science 238: 650-656, 1987.

[185987]

[185988]

[185989]19519.

[185990]Oakey, R. J.; Caron, M. G.; Lefkowitz, R. J.; Seldin, M. F.:
Genomic

[185991]organization of adrenergic and serotonin receptors in the mouse: linkage

[185992]mapping of sequence-related genes provides a method for examining

[185993]mammalian chromosome evolution. Genomics 10: 338–344, 1991.

[185994]

[185995]

[185996]19520.

[185997]Philipp, M.; Brede, M. E.; Hadamek, K.; Gessler, M.; Lohse, M.

[185998]J.; Hein, L.: Placental alpha-2-adrenoceptors control vascular development

[185999]at the interface between mother and embryo. Nature Genet. 31: 311–315,

[186000]2002.

[186001]

[186002]

[186003]19521.

[186004]Surprenant, A.; Horstman, D. A.; Akbarali, H.; Limbird, L. E.:

[186005]A point mutation of the alpha-2-adrenoceptor that blocks coupling

[186006]to potassium but not calcium currents. Science 257: 977-980, 1992.

[186007]

[186008]

[186009]19522.

[186010]Yang-Feng, T. L.; Kobilka, B. K.; Caron, M. G.; Lefkowitz, R.

[186011]J.; Francke, U.: Chromosomal assignment of genes for an alpha-adrenergic

[186012]receptor (ADRAR) and for another member of this receptor family coupled

[186013]to guanine nucleotide regulatory proteins (RG21). (Abstract) Cytogenet.

[186014]Cell Genet. 46: 722-723, 1987.

[186015]

[186016]

[186017]19523.

[186018]Bruno, J. F.; Whittaker, J.; Song, J.; Berelowitz, M.: Molecular
lar

[186019]cloning and sequencing of a cDNA encoding a human al-
pha-1A adrenergic

[186020]receptor. Biochem. Biophys. Res. Commun. 179:
1485-1490, 1991.

[186021]

[186022]

[186023]19524.

[186024]Esbenshade, T. A.; Hirasawa, A.; Tsujimoto, G.; Tanaka, T.;
Yano,

[186025]J.; Minneman, K. P.; Murphy, T. J.: Cloning of the human
alpha-1D-adrenergic

[186026]receptor and inducible expression of three subtypes in
SK-N-MC cells. Molec.

[186027]Pharm. 47: 977-985, 1995.

[186028]

[186029]

[186030]19525.

[186031]Forray, C.; Bard, J. A.; Wetzel, J. M.; Chiu, G.; Shapiro, E.;

[186032]Tang, R.; Lepor, H.; Hartig, P. R.; Weinshank, R. L.;
Branchek, T.

[186033]A.; Gluchowski, C.: The alpha-1-adrenergic receptor that mediates

[186034]smooth muscle contraction in human prostate has the pharmacological

[186035]properties of the cloned human alpha-1C subtype. Molec. Pharm. 45:

[186036]703-708, 1994.

[186037]

[186038]

[186039]19526.

[186040]Loftus, S. K.; Shiang, R.; Warrington, J. A.; Bengtsson, U.; McPherson,

[186041]J. D.; Wasmuth, J. J.: Genes encoding adrenergic receptors are not

[186042]clustered on the long arm of human chromosome 5. Cytogenet. Cell

[186043]Genet. 67: 69-74, 1994.

[186044]

[186045]

[186046]19527.

[186047]Lomasney, J. W.; Cotecchia, S.; Lorenz, W.; Leung, W.-Y.;
Schwinn,

[186048]D. A.; Yang-Feng, T. L.; Brownstein, M.; Lefkowitz, R. J.;
Caron,

[186049]M. G.: Molecular cloning and expression of the cDNA for
the alpha-1A-adrenergic

[186050]receptor: the gene for which is located on human chromo-
some 5. J.

[186051]Biol. Chem. 266: 6365-6369, 1991.

[186052]

[186053]

[186054]19528.

[186055]Schwinn, D. A.; Johnston, G. I.; Page, S. O.; Mosley, M. J.;
Wilson,

[186056]K. H.; Worman, N. P.; Campbell, S.; Fidock, M. D.; Furness,
L. M.;

[186057]Parry-Smith, D. J.; Peter, B.; Bailey, D. S.: Cloning and
pharmacological

[186058]characterization of human alpha-1 adrenergic receptors:
sequence corrections

[186059]and direct comparison with other species homologues. J.
Pharm. Exp.

[186060]Ther. 272: 134–142, 1995.

[186061]

[186062]

[186063]19529.

[186064]Schwinn, D. A.; Lomasney, J. W.: Pharmacologic characterization

[186065]of cloned alpha-1-adrenoceptor subtypes: selective antagonists suggest

[186066]the existence of a fourth subtype. Europ. J. Pharm. 227: 433–436,

[186067]1992.

[186068]

[186069]

[186070]19530.

[186071]Weinberg, D. H.; Trivedi, P.; Tan, C. P.; Mitra, S.; Perkins-Barrow,

[186072]A.; Borkowski, D.; Strader, C. D.; Bayne, M.: Cloning, expression

[186073]and characterization of human alpha adrenergic receptors alpha-1A,

[186074]alpha-1B, and alpha-1C. Biochem. Biophys. Res. Commun. 201: 1296-1304,
[186075]1994.

[186076]

[186077]

[186078]19531.

[186079]Yang-Feng, T. L.; Han, H.; Lomasney, J. W.; Caron, M. G.:
Localization

[186080]of the cDNA for an alpha-1-adrenergic receptor subtype
(ADRA1D) to

[186081]chromosome band 20p13. Cytogenet. Cell Genet. 66:
170-171, 1994.

[186082]

[186083]

[186084]19532.

[186085]Allen, L. F.; Lefkowitz, R. J.; Caron, M. G.; Cotecchia, S.: G-
protein-coupled

[186086]receptor genes as protooncogenes: constitutively activating
mutation

[186087]of the alpha-1B-adrenergic receptor enhances mitogenesis
and tumorigenicity. Proc.

[186088]Nat. Acad. Sci. 88: 11354–11358, 1991.

[186089]

[186090]

[186091]19533.

[186092]Markert, M. L.; Hershfield, M. S.; Wiginton, D. A.; States, J.

[186093]C.; Ward, F. E.; Bigner, S. H.; Buckley, R. H.; Kaufman, R.
E.; Hutton,

[186094]J. J.: Identification of a deletion in the adenosine deami-
nase gene

[186095]in a child with severe combined immunodeficiency. J. Im-
mun. 138:

[186096]3203–3206, 1987.

[186097]

[186098]

[186099]19534.

[186100]Markert, M. L.; Hutton, J. J.; Wiginton, D. A.; States, J. C.;

[186101]Kaufman, R. E.: Adenosine deaminase (ADA) deficiency
due to deletion

[186102]of the ADA gene promoter and first exon by homologous

recombination

[186103]between two Alu elements. J. Clin. Invest. 81: 1323–1327,
1988.

[186104]

[186105]

[186106]19535.

[186107]Markert, M. L.; Norby–Slycord, C.; Ward, F. E.: A high pro-
portion

[186108]of ADA point mutations associated with a specific alanine-
to-valine

[186109]substitution. Am. J. Hum. Genet. 45: 354–361, 1989.

[186110]

[186111]

[186112]19536.

[186113]Meuwissen, H. J.; Pollara, B.; Pickering, R. J.: Combined
immunodeficiency

[186114]disease associated with adenosine deaminase deficiency
(report on

[186115]a workshop held in Albany, New York, October 1, 1973). J.
Pediat. 86:

[186116]169–181, 1975.

[186117]

[186118]

[186119]19537.

[186120]Migchielsen, A. A. J.; Breuer, M. L.; van Roon, M. A.; te Riele,

[186121]H.; Zurcher, C.; Ossendorp, F.; Toutain, S.; Hershfield, M. S.; Berns,

[186122]A.; Valerio, D.: Adenosine–deaminase–deficient mice die perinatally

[186123]and exhibit liver–cell degeneration, atelectasis and small intestinal

[186124]cell death. Nature Genet. 10: 279–287, 1995.

[186125]

[186126]

[186127]19538.

[186128]Mitchell, B. S.; Mejias, E.; Daddona, P. E.; Kelley, W. N.: Purinogenic

[186129]immunodeficiency disease: selective toxicity of deoxyri–bonucleosides

[186130]for T–cells. Proc. Nat. Acad. Sci. 75: 5011–5014, 1978.

[186131]

[186132]

[186133]19539.

[186134]Moen, R. C.; Horowitz, S. D.; Sondel, P. M.; Borcharding,
W. R.;

[186135]Trigg, M. E.; Billing, R.; Hong, R.: Immunologic reconstitu-
tion after

[186136]haploidentical bone marrow transplantation for immune
deficiency disorders:

[186137]treatment of bone marrow cells with monoclonal antibody
CT-2 and complement. Blood 70:

[186138]664-669, 1987.

[186139]

[186140]

[186141]19540.

[186142]Mohandas, T.; Sparkes, R. S.; Suh, E. J.; Hershfield, M. S.:

[186143]Regional localization of the human genes for S-
adenosylhomocysteine

[186144]hydrolase (cen-q131) and adenosine deaminase
(q131-qter) on chromosome

[186145]20. Hum. Genet. 66: 292–295, 1984.

[186146]

[186147]

[186148]19541.

[186149]Nielsen, K. B.; Tommerup, N.; Jespersen, B.; Nygaard, P.;
Kleif,

[186150]L.: Segregation of a t(3;20) translocation through three
generations

[186151]resulting in unbalanced karyotypes in six persons. J. Med.
Genet. 23:

[186152]446–451, 1986.

[186153]

[186154]

[186155]19542.

[186156]Orkin, S. H.; Daddona, P. E.; Shewach, D. S.; Markham, A.
F.;

[186157]Bruns, G. A.; Goff, S. C.; Kelley, W. N.: Molecular cloning of
human

[186158]adenosine deaminase gene sequences. J. Biol. Chem. 258:
12753–12756,

[186159]1983.

[186160]

[186161]

[186162]19543.

[186163]Ozsahin, H.; Arredondo-Vega, F. X.; Santisteban, I.;

Fuhrer, H.;

[186164]Tuchschmid, P.; Jochum, W.; Aguzzi, A.; Lederman, H. M.;

Fleischman,

[186165]A.; Winkelstein, J. A.; Seger, R. A.; Hershfield, M. S.:

Adenosine

[186166]deaminase deficiency in adults. Blood 89: 2849–2855,

1997.

[186167]

[186168]

[186169]19544.

[186170]Onodera, M.; Ariga, T.; Kawamura, N.; Kobayashi, I.;

Ohtsu, M.;

[186171]Yamada, M.; Tame, A.; Furuta, H.; Okano, M.; Matsumoto,

S.; Kotani,

[186172]H.; McGarrity, G. J.; Blaese, R. M.; Sakiyama, Y.: Successful

peripheral

[186173]T-lymphocyte-directed gene transfer for a patient with
severe combined

[186174]immune deficiency caused by adenosine deaminase defi-
ciency. Blood 91:

[186175]30-36, 1998.

[186176]

[186177]

[186178]19545.

[186179]Palmer, T. D.; Hock, R. A.; Osborne, W. R. A.; Miller, A. D.:

[186180]Efficient retrovirus-mediated transfer and expression of a
human adenosine

[186181]deaminase gene in diploid skin fibroblasts from an
adenosine deaminase-deficient

[186182]human. Proc. Nat. Acad. Sci. 84: 1055-1059, 1987.

[186183]

[186184]

[186185]19546.

[186186]Parkman, R.; Gelfand, E. W.; Rosen, F. S.; Sanderson, A.;
Hirschhorn,

[186187]R.: Severe combined immunodeficiency and adenosine

deaminase deficiency. New

[186188]Eng. J. Med. 292: 714–719, 1975.

[186189]

[186190]

[186191]19547.

[186192]Petersen, M. B.; Tranebjaerg, L.; Tommerup, N.; Nygaard, P.; Edwards,

[186193]H.: New assignment of the adenosine deaminase gene locus to chromosome

[186194]20q13.11 by study of a patient with interstitial deletion 20q. J.

[186195]Med. Genet. 24: 93–96, 1987.

[186196]

[186197]

[186198]19548.

[186199]Polmar, S. H.; Stern, R. C.; Schwartz, A. L.; Wetzler, E. M.;

[186200]Chase, P. A.; Hirschhorn, R.: Enzyme replacement therapy for adenosine

[186201]deaminase deficiency and severe combined immunodeficiency. New Eng.

[186202]J. Med. 295: 1337–1343, 1976.

[186203]

[186204]

[186205]19549.

[186206]Ratech, H.; Greco, M. A.; Gallo, G.; Rimoin, D. L.; Kamino, H.;

[186207]Hirschhorn, R.: Pathologic findings in adenosine–deaminase–deficient

[186208]severe combined immunodeficiency. I. Kidney, adrenal, and chondro–osseous

[186209]tissue alterations. Am. J. Path. 120: 157–169, 1985.

[186210]

[186211]

[186212]19550.

[186213]Ritter, H.; Wendt, G. G.; Tariverdian, G.; Zelch, J.; Rube, M.;

[186214]Kirchberg, G.: Genetics and linkage analysis of adenosine deaminase. Humangenetik 14:

[186215]69–71, 1971.

[186216]

[186217]

[186218]19551.

[186219]Rogers, M. H.; Lwin, R.; Fairbanks, L.; Gerritsen, B.; Gaspar,

[186220]H. B.: Cognitive and behavioral abnormalities in adenosine deaminase

[186221]deficient severe combined immunodeficiency. J. Pediat. 139: 44–50,

[186222]2001.

[186223]

[186224]

[186225]19552.

[186226]Rothschild, C. B.; Akots, G.; Hayworth, R.; Pettenati, M. J.;

[186227]Rao, P. N.; Wood, P.; Stolz, F.–M.; Hansmann, I.; Serino, K.; Keith,

[186228]T. P.; Fajans, S. S.; Bowden, D. W.: A genetic map of chromosome

[186229]20q12–q13.1: multiple highly polymorphic microsatellite and RFLP markers

[186230]linked to the maturity–onset diabetes of the young (MODY) locus. Am.

[186231]J. Hum. Genet. 52: 110–123, 1993.

[186232]

[186233]

[186234]19553.

[186235]Rubinstein, A.; Hirschhorn, R.; Sicklick, M.; Murphy, R. A.:

[186236]In vivo and in vitro effects of thymosin and adenosine
deaminase on

[186237]adenosine–deaminase–deficient lymphocytes. New Eng. J.
Med. 300:

[186238]387–392, 1979.

[186239]

[186240]

[186241]19554.

[186242]Rudd, N. L.; Bain, H. W.; Giblett, E.; Chen, S.–H.; Worton, R.

[186243]G.: Partial trisomy 20 confirmed by gene dosage studies.
Am. J.

[186244]Med. Genet. 4: 357–364, 1979.

[186245]

[186246]

[186247]19555.

[186248]Koch, G.; Shows, T. B.: Somatic cell genetics of adenosine deaminase

[186249]expression and severe combined immune deficiency disease in man. Proc.

[186250]Nat. Acad. Sci. 77: 4211–4215, 1980.

[186251]

[186252]

[186253]19556.

[186254]Gautam, M.; Noakes, P. G.; Moscoso, L.; Rupp, F.; Scheller, R.

[186255]H.; Merlie, J. P.; Sanes, J. R.: Defective neuromuscular synaptogenesis

[186256]in agrin-deficient mutant mice. Cell 85: 525–535, 1996.

[186257]

[186258]

[186259]19557.

[186260]Glass, D. J.; Bowen, D. C.; Stitt, T. N.; Radziejewski, C.; Bruno,

[186261]J.; Ryan, T. E.; Gies, D. R.; Shah, S.; Mattsson, K.; Burden,

S. J.;

[186262]DiStefano, P. S.; Valenzuela, D. M.; DeChiara, T. M.; Yan-
copoulos,

[186263]G. D.: Agrin acts via a MuSK receptor complex. Cell 85:
513–523,

[186264]1996.

[186265]

[186266]

[186267]19558.

[186268]Khan, A. A.; Bose, C.; Yam, L. S.; Soloski, M. J.; Rupp, F.:
Physiological

[186269]regulation of the immunological synapse by agrin. Science
292: 1681–1686,

[186270]2001.

[186271]

[186272]

[186273]19559.

[186274]Lin, W.; Burgess, R. W.; Dominguez, B.; Pfaff, S. L.; Sanes,
J.

[186275]R.; Lee, K.–F.: Distinct roles of nerve and muscle in post–

synaptic

[186276]differentiation of the neuromuscular synapse. Nature 410:
1057–1064,

[186277]2001.

[186278]

[186279]

[186280]19560.

[186281]McMahan, U. J.: The agrin hypothesis Cold Spring Harb.
Symp.

[186282]Quant. Biol. 50: 407–418, 1990.

[186283]

[186284]

[186285]19561.

[186286]Rupp, F.; Ozcelik, T.; Linial, M.; Peterson, K.; Francke, U.;
Scheller,

[186287]R.: Structure and chromosomal localization of the mam-
malian agrin

[186288]gene. J. Neurosci. 12: 3535–3544, 1992.

[186289]

[186290]

[186291]19562.

[186292]Rupp, F.; Payan, D. G.; Magill-Solc, C.; Cowan, D. M.;
Scheller,

[186293]R. H.: Structure and expression of a rat agrin. Neuron 6:
811-823,

[186294]1991.

[186295]

[186296]

[186297]19563.

[186298]Bard, L. A.: Heterogeneity in Waardenburg's syndrome: re-
port of

[186299]a family with ocular albinism. Arch. Ophthal. 96:
1193-1198, 1978.

[186300]

[186301]

[186302]19564.

[186303]Morell, R.; Spritz, R. A.; Ho, L.; Pierpont, J.; Guo, W.; Fried-
man,

[186304]T. B.; Asher, J. H., Jr.: Apparent digenic inheritance of

Waardenburg

[186305]syndrome type 2 (WS2) and autosomal recessive ocular albinism (AROA). Hum.

[186306]Molec. Genet. 6: 659–664, 1997.

[186307]

[186308]

[186309]19565.

[186310]Smith, S. D.; Kelley, P. M.; Kenyon, J. B.; Hoover, D.: Tietz

[186311]syndrome (hypopigmentation/deafness) caused by mutation of MITF. J.

[186312]Med. Genet. 37: 446–448, 2000.

[186313]

[186314]

[186315]19566.

[186316]Tietz, W.: A syndrome of deaf–mutism associated with albinism

[186317]showing dominant autosomal inheritance. Am. J. Hum. Genet. 15: 259–264,

[186318]1963.

[186319]

[186320]

[186321]19567.

[186322]Umans, L.; Serneels, L.; Overbergh, L.; Lorent, K.; Van
Leuven,

[186323]F.; Van den Berghe, H.: Targeted inactivation of the mouse
alpha-2-macroglobulin

[186324]gene. J. Biol. Chem. 270: 19778–19785, 1995.

[186325]

[186326]

[186327]19568.

[186328]D'Alfonso, S.; Richiardi, P. M.: A polymorphic variation in a

[186329]putative regulation box of the TNFA promoter region. Im-
munogenetics 39:

[186330]150–154, 1994.

[186331]

[186332]

[186333]19569.

[186334]Mietus-Snyder, M.; Charmley, P.; Korf, B.; Ladas, J. A. A.:

Gatti,

[186335]R. A. and Karathanasis, S. K.: Genetic linkage of the human apolipoprotein

[186336]AI-CIII-AIV gene cluster and the neural cell adhesion molecule (NCAM)

[186337]gene. Genomics 7: 633-637, 1990.

[186338]

[186339]

[186340]19570.

[186341]Mietus-Snyder, M.; Korf, B.; Ladias, J. A.; Karathanasis, S. K.

[186342]: Linkage of the human apolipoproteins A1, C3, A4 and the neural cell

[186343]adhesion molecule (NCAM) genes. (Abstract) Cytogenet. Cell Genet. 51:

[186344]1044 only, 1989.

[186345]

[186346]

[186347]19571.

[186348]Nguyen, C.; Mattei, M. G.; Goridis, C.; Mattei, J. F.; Jordan,

[186349]B. R.: Localization of the human N-CAM gene to chromosome 11 by in

[186350]situ hybridization with a murine N-CAM cDNA probe.

(Abstract) Cytogenet.

[186351]Cell Genet. 40: 713 only, 1985.

[186352]

[186353]

[186354]19572.

[186355]Nguyen, C.; Mattei, M. G.; Mattei, J.-F.; Santoni, M.-J.; Goridis,

[186356]C.; Jordan, B. R.: Localization of the human NCAM gene to band q23

[186357]of chromosome 11: the third gene coding for a cell interaction molecule

[186358]mapped to the distal portion of the long arm of chromosome 11. J.

[186359]Cell Biol. 102: 711-715, 1986.

[186360]

[186361]

[186362]19573.

[186363]Rabinowitz, J. E.; Rutishauser, U.; Magnuson, T.: Targeted mutation

[186364]of Ncam to produce a secreted molecule results in a dominant embryonic

[186365]lethality. Proc. Nat. Acad. Sci. 93: 6421–6424, 1996.

[186366]

[186367]

[186368]19574.

[186369]Rutishauser, U.; Acheson, A.; Hall, A. K.; Mann, D. M.; Sunshine,

[186370]J.: The neural cell adhesion molecule (NCAM) as a regulator of cell-cell

[186371]interactions. Science 240: 53–57, 1988.

[186372]

[186373]

[186374]19575.

[186375]Rutishauser, U.; Goridis, C.: NCAM: the molecule and its genetics. Trends

[186376]Genet. 2: 72–76, 1986.

[186377]

[186378]

[186379]19576.

[186380]Telatar, M.; Lange, E.; Uhrhammer, N.; Gatti, R. A.: New
localization

[186381]of NCAM, proximal to DRD2 at chromosome 11q23. Mam-
malian Genome 6:

[186382]59–60, 1995.

[186383]

[186384]

[186385]19577.

[186386]Agellon, L. B.; Drover, V. A. B.; Cheema, S. K.; Gbaguidi, G.
F.;

[186387]Walsh, A.: Dietary cholesterol fails to stimulate the human
cholesterol

[186388]7- α -hydroxylase gene (CYP7A1) in transgenic mice. J.
Biol. Chem. 277:

[186389]20131–20134, 2002.

[186390]

[186391]

[186392]19578.

[186393] Angelin, B.; Einarsson, K.; Hellstrom, K.; Leijd, B.: Bile acid
[186394] kinetics in relation to endogenous triglyceride metabolism
in various
[186395] types of hyperlipoproteinemia. J. Lipid Res. 19:
1004–1016, 1978.

[186396]

[186397]

[186398] 19579.

[186399] Angelin, B.; Hershon, K. S.; Brunzell, J. D.: Bile acid
metabolism
[186400] in hereditary forms of hypertriglyceridemia: evidence for
an increased
[186401] synthesis rate in monogenic familial hypertriglyceridemia.
Proc.

[186402] Nat. Acad. Sci. 84: 5434–5438, 1987.

[186403]

[186404]

[186405] 19580.

[186406] Cohen, J. C.; Cali, J. J.; Jelinek, D. F.; Mehrabian, M.;
Sparkes,

[186407]R. S.; Lusis, A. J.; Russell, D. W.; Hobbs, H. H.: Cloning of
the

[186408]human cholesterol 7- α -hydroxylase gene (CYP7) and
localization

[186409]to chromosome 8q11-q12. Genomics 14: 153-161, 1992.

[186410]

[186411]

[186412]19581.

[186413]Drover, V. A. B.; Wong, N. C. W.; Agellon, L. B.: A distinct
thyroid

[186414]hormone response element mediates repression of the
human cholesterol

[186415]7- α -hydroxylase (CYP7A1) gene promoter. Molec. En-
docr. 16: 14-23,

[186416]2002.

[186417]

[186418]

[186419]19582.

[186420]Goodwin, B.; Jones, S. A.; Price, R. R.; Watson, M. A.; Mc-
Kee,

[186421]D. D.; Moore, L. B.; Galardi, C.; Wilson, J. G.; Lewis, M. C.;
Roth,

[186422]M. E.; Maloney, P. R.; Willson, T. M.; Kliewer, S. A.: A regu-
latory

[186423]cascade of the nuclear receptors FXR, SHP-1, and LRH-1
represses bile

[186424]acid biosynthesis. Molec. Cell 6: 517-526, 2000.

[186425]

[186426]

[186427]19583.

[186428]Molowa, D. T.; Chen, W. S.; Cimis, G. M.; Tan, C. P.: Tran-
scriptional

[186429]regulation of the human cholesterol 7 alpha-hydroxylase
gene. Biochemistry 31:

[186430]2539-2544, 1992.

[186431]

[186432]

[186433]19584.

[186434]Nitta, M.; Ku, S.; Brown, C.; Okamoto, A. Y.; Shan, B.: CPF:
an

[186435] orphan nuclear receptor that regulates liver-specific expression of

[186436] the human cholesterol 7- α -hydroxylase gene. Proc. Nat. Acad.

[186437] Sci. 96: 6660–6665, 1999.

[186438]

[186439]

[186440] 19585.

[186441] Noshiro, M.; Okuda, K.: Molecular cloning and sequence analysis

[186442] of cDNA encoding human cholesterol 7 α -hydroxylase. FEBS Lett. 268:

[186443] 137–140, 1990.

[186444]

[186445]

[186446] 19586.

[186447] Paumgartner, G.; Sauerbruch, T.: Gallstones: pathogenesis. Lancet 338:

[186448] 1117–1121, 1991.

[186449]

[186450]

[186451]19587.

[186452]Wang, J.; Freeman, D. J.; Grundy, S. M.; Levine, D. M.;

Guerra,

[186453]R.; Cohen, J. C.: Linkage between cholesterol

7- α -hydroxylase

[186454]and high plasma low-density lipoprotein cholesterol concentrations. J.

[186455]Clin. Invest. 101: 1283-1291, 1998.

[186456]

[186457]

[186458]19588.

[186459]Akita, H.; Chiba, H.; Tsuchihashi, K.; Tsuji, M.; Kumagai, M.;

[186460]Matsuno, K.; Kobayashi, K.: Cholesteryl ester transfer protein gene:

[186461]two common mutations and their effect on plasma high-density lipoprotein

[186462]cholesterol content. J. Clin. Endocr. Metab. 79: 1615-1618, 1994.

[186463]

[186464]

[186465]19589.

[186466]Altshuler, D.; Kruglyak, L.; Lander, E.: Genetic polymorphisms

[186467]and disease. (Letter) New Eng. J. Med. 338: 1626 only, 1998.

[186468]

[186469]

[186470]19590.

[186471]Brown, M. L.; Inazu, A.; Hesler, C. B.; Agellon, L. B.; Mann, C.;

[186472]Whitlock, M. E.; Marcel, Y. L.; Milne, R. W.; Koizumi, J.; Mabuchi,

[186473]H.; Takeda, R.; Tall, A. R.: Molecular basis of lipid transfer protein

[186474]deficiency in a family with increased high-density lipoproteins. Nature 342:

[186475]448–451, 1989.

[186476]

[186477]

[186478]19591.

[186479]Drayna, D.; Jarnagin, A. S.; McLean, J.; Henzel, W.; Kohr, W.;

[186480]Fielding, C.; Lawn, R.: Cloning and sequencing of human cholesteryl

[186481]ester transfer protein cDNA. Nature 327: 632–634, 1987.

[186482]

[186483]

[186484]19592.

[186485]Drayna, D.; Lawn, R. M.: Multiple RFLPs at the human cholesteryl

[186486]ester transfer protein (CETP) locus. Nucleic Acids Res. 15: 4698

[186487]only, 1987.

[186488]

[186489]

[186490]19593.

[186491]Haefliger, J.–A.; Bruzzone, R.; Jenkins, N. A.; Gilbert, D. J.;

[186492]Copeland, N. G.; Paul, D. L.: Four novel members of the

connexin

[186493]family of gap junction proteins: molecular cloning, expression, and

[186494]chromosome mapping. J. Biol. Chem. 267: 2057–2064, 1992.

[186495]

[186496]

[186497]19594.

[186498]Hsieh, C.-L.; Kumar, N. M.; Gilula, N. B.; Francke, U.: Dis-
tribution

[186499]of genes for gap junction membrane channel proteins on
human and mouse

[186500]chromosomes. Somat. Cell Molec. Genet. 17: 191-200,
1991.

[186501]

[186502]

[186503]19595.

[186504]Spielman, R. S.; McGinnis, R. E.; Ewens, W. J.: Transmission

[186505]test for linkage disequilibrium: the insulin gene region
and insulin-dependent

[186506]diabetes mellitus (IDDM). Am. J. Hum. Genet. 52:
506-516, 1993.

[186507]

[186508]

[186509]19596.

[186510]Altshuler, D.; Hirschhorn, J. N.; Klannemark, M.; Lindgren,

C.

[186511]M.; Vohl, M.-C.; Nemesh, J.; Lane, C. R.; Schaffner, S. F.;
Bolk,

[186512]S.; Brewer, C.; Tuomi, T.; Gaudet, D.; Hudson, T. J.; Daly,
M.; Groop,

[186513]L.; Lander, E. S.: The common PPAR-gamma pro12ala
polymorphism is

[186514]associated with decreased risk of type 2 diabetes. Nature
Genet. 76-80,

[186515]2000.

[186516]

[186517]

[186518]19597.

[186519]Fakhrai-Rad, H.; Nikoshkov, A.; Kamel, A.; Fernstrom, M.;
Zierath,

[186520]J. R.; Norgren, S.; Luthman, H.; Galli, J.: Insulin-degrading
enzyme

[186521]identified as a candidate diabetes susceptibility gene in
GK rats. Hum.

[186522]Molec. Genet. 9: 2149-2158, 2000.

[186523]

[186524]

[186525]19598.

[186526]Gersuk, V. H.; Rose, T. M.; Todaro, G. J.: Molecular cloning
and

[186527]chromosomal localization of a pseudogene related to the
human acyl-CoA

[186528]binding protein/diazepam binding inhibitor. Genomics 25:
469–476,

[186529]1995.

[186530]

[186531]

[186532]19599.

[186533]DeStefano, A. L.; Baldwin, C. T.; Burzstyn, M.; Gavras, I.;
Handy,

[186534]D. E.; Joost, O.; Martel, T.; Nicolaou, M.; Schwartz, F.;
Streeten,

[186535]D. H. P.; Farrer, L. A.; Gavras, H.: Autosomal dominant or-
thostatic

[186536]hypotensive disorder maps to chromosome 18q. Am. J.
Hum. Genet. 63:

[186537]1425–1430, 1998.

[186538]

[186539]

[186540]19600.

[186541]Barnett, T.; Zimmermann, W.: Workshop report: proposed
nomenclature

[186542]for the carcinoembryonic antigen (CEA) gene family. Tu-
mor Biol. 11:

[186543]59–63, 1990.

[186544]

[186545]

[186546]19601.

[186547]Boulton, I. C.; Gray-Owen, S. D.: Neisserial binding to
CEACAM1

[186548]arrests the activation and proliferation of CD4+ T lym-
phocytes. Nature

[186549]Immun. 3: 229–236, 2002.

[186550]

[186551]

[186552]19602.

[186553]Ergun, S.; Kilic, N.; Ziegeler, G.; Hansen, A.; Nollau, P.;

Gotze,

[186554]J.; Wurmbach, J.-H.; Horst, A.; Weil, J.; Fernando, M.; Wagner, C.

[186555]: CEA-related cell adhesion molecule 1: a potent angiogenic factor

[186556]and a major effector of vascular endothelial growth factor. Molec.

[186557]Cell 5: 311-320, 2000.

[186558]

[186559]

[186560]19603.

[186561]Hinoda, Y.; Neumaier, M.; Hefta, S. A.; Drzeniek, Z.; Wagner,

[186562]C.; Shively, L.; Hefta, L. J. F.; Shively, J. E.; Paxton, R. J.:

[186563]Molecular cloning of a cDNA coding biliary glycoprotein I: primary

[186564]structure of a glycoprotein immunologically crossreactive with carcinoembryonic

[186565]antigen. Proc. Nat. Acad. Sci. 85: 6959-6963, 1988.

[186566]

[186567]

[186568]19604.

[186569]Neumaier, M.; Paululat, S.; Chan, A.; Matthaes, P.; Wanger, C.

[186570]: Biliary glycoprotein, a potential human cell adhesion molecule,

[186571]is down-regulated in colorectal carcinomas. Proc. Nat. Acad. Sci. 90:

[186572]10744–10748, 1993.

[186573]

[186574]

[186575]19605.

[186576]Poy, M. N.; Yang, Y.; Rezaei, K.; Fernstrom, M. A.; Lee, A. D.;

[186577]Kido, Y.; Erickson, S. K.; Najjar, S. M.: CEACAM1 regulates insulin

[186578]clearance in liver. Nature Genet. 30: 270–276, 2002.

[186579]

[186580]

[186581]19606.

[186582]Robbins, J.; Robbins, P. F.; Kozak, C. A.; Callahan, R.: The
[186583]mouse biliary glycoprotein gene (Bgp): partial nucleotide
sequence,
[186584]expression, and chromosomal assignment. Genomics 10:
583–587, 1991.

[186585]

[186586]

[186587]19607.

[186588]Thompson, J. A.; Grunert, F.; Zimmermann, W.: Carci-
noembryonic

[186589]antigen gene family: molecular biology and clinical per-
spectives. J.

[186590]Clin. Lab. Anal. 5: 344–366, 1991.

[186591]

[186592]

[186593]19608.

[186594]Willard, H. F.; Meakin, S. O.; Tsui, L.-C.; Breitman, M. L.:

[186595]Assignment of human gamma crystallin multigene family
to chromosome

[186596]2. Somat. Cell Molec. Genet. 11: 511–516, 1985.

[186597]

[186598]

[186599]19609.

[186600]Wistow, G.; Piatigorsky, J.: Recruitment of enzymes as lens structural

[186601]proteins. Science 236: 1554–1556, 1987.

[186602]

[186603]

[186604]19610.

[186605]Zneimer, S. M.; Womack, J. E.: Regional localization of the fibronectin

[186606]and gamma crystallin genes to mouse chromosome 1 by in situ hybridization. Cytogenet.

[186607]Cell Genet. 48: 238–241, 1988.

[186608]

[186609]

[186610]19611.

[186611]Daniels, G.; King, M.-J.; Avent, N. D.; Khalid, G.; Reid, M.;

[186612]Mallinson, G.; Symthe, J.; Cedergren, B.: A point mutation

in the

[186613]GYPC gene results in the expression of the blood group
An(a) antigen

[186614]on glycophorin D but not on glycophorin C: further evi-
dence that glycophorin

[186615]D is a product of the GYPC gene. Blood 82: 3198–3203,
1993.

[186616]

[186617]

[186618]19612.

[186619]El-Maliki, B.; Blanchard, D.; Dahr, W.; Beyreuther, K.;
Cartron,

[186620]J.-P.: Structural homology between glycophorins C and D
of human

[186621]erythrocytes. Europ. J. Biochem. 183: 639–643, 1989.

[186622]

[186623]

[186624]19613.

[186625]Jorgensen, J.; Drachmann, O.; Gavin, J.: Duch, Dh(a), a low
frequency

[186626]red cell antigen. Hum. Hered. 32: 73–75, 1982.

[186627]

[186628]

[186629]19614.

[186630]King, M. J.; Avent, N. D.; Mallinson, G.; Reid, M. E.: Point

[186631]mutation in the glycophorin C gene results in the expression of the

[186632]blood group antigen Dh(a). Vox Sang. 63: 56–58, 1992.

[186633]

[186634]

[186635]19615.

[186636]Le Van Kim, C.; Colin, Y.; Blanchard, D.; Dahr, W.; London, J.;

[186637]Cartron, J.-P.: Gerbich blood group deficiency of the Ge:–1,–2,–3

[186638]and Ge:–1,–2,3 types: immunochemical study and genomic analysis with

[186639]cDNA probes. Europ. J. Biochem. 165: 571–579, 1987.

[186640]

[186641]

[186642]19616.

[186643]Mattei, M. G.; Colin, Y.; Le Van Kim, C.; Mattei, J. F.;

Cartron,

[186644]J. P.: Localization of the gene for human erythrocyte gly-
cophorin

[186645]C to chromosome 2, q14-q21. Hum. Genet. 74: 420-422,
1986.

[186646]

[186647]

[186648]19617.

[186649]Pasvol, G.; Anstee, D. J.; Tanner, M. J. A.: Glycophorin C
and

[186650]the invasion of red cells by Plasmodium falciparum. Lancet
I: 907-908,

[186651]1984.

[186652]

[186653]

[186654]19618.

[186655]Race, R. R.; Sanger, R.: Blood Groups in Man. Oxford:
Blackwell

[186656]Sci. Publ. (pub.) (6th ed.): 1975. Pp. 416–421.

[186657]

[186658]

[186659]19619.

[186660]Reid, M. E.: The Gerbich blood group antigens: a review.
Med.

[186661]Lab. Sci. 43: 177–182, 1972.

[186662]

[186663]

[186664]19620.

[186665]Reid, M. E.; Sullivan, C.; Taylor, M.; Anstee, D. J.: Inheri-
tance

[186666]of human–erythrocyte Gerbich blood group antigens. Am.
J. Hum. Genet. 41:

[186667]1117–1123, 1987.

[186668]

[186669]

[186670]19621.

[186671]Simmons, R. T.; Albrey, J. A.: A 'new' blood group antigen

Webb

[186672](Wb) of low frequency found in two Australian families.

Med. J. Aust. I:

[186673]8–10, 1963.

[186674]

[186675]

[186676]19622.

[186677]Sondag, D.; Alloisio, N.; Blanchard, D.; Ducluzeau, M.–T.;

Colonna,

[186678]P.; Bachir, D.; Bloy, C.; Cartron, J.–P.; Delaunay, J.: Gerbich

reactivity

[186679]in 4.1(–) hereditary elliptocytosis and protein 4.1 level in

blood

[186680]group Gerbich deficiency. Brit. J. Haemat. 65: 43–50,

1987.

[186681]

[186682]

[186683]19623.

[186684]Spring, F. A.: Immunochemical characterisation of the

low-incidence

[186685]antigen, Dh(a). Vox Sang. 61: 65–68, 1991.

[186686]

[186687]

[186688]19624.

[186689]Telen, M. J.; Le Van Kim, C.; Guizzo, M. L.; Cartron, J.-P.;
Colin,

[186690]Y.: Erythrocyte Webb-type glycophorin C variant lacks N-
glycosylation

[186691]due to an asparagine to serine substitution. Am. J. Hemat.
37: 51-52,

[186692]1991.

[186693]

[186694]

[186695]19625.

[186696]Winardi, R.; Reid, M.; Conboy, J.; Mohandas, N.: Molecular
analysis

[186697]of glycophorin C deficiency in human erythrocytes. Blood
81: 2799-2803,

[186698]1993.

[186699]

[186700]

[186701]19626.

[186702]Bierhuizen, M. F. A.; Mattei, M.-G.; Fukuda, M.: Expression
of

[186703]the developmental I antigen by a cloned human cDNA en-
coding a member

[186704]of a beta-1,6-N-acetylglucosaminyltransferase gene fam-
ily. Genes

[186705]Dev. 7: 468-478, 1993.

[186706]

[186707]

[186708]19627.

[186709]Lin-Chu, M.; Broadberry, R. E.; Okubo, Y.; Tanaka, M.: The
i phenotype

[186710]and congenital cataracts among Chinese in Taiwan (Letter)
Transfusion 31:

[186711]676-677, 1991.

[186712]

[186713]

[186714]19628.

[186715]Ogata, H.; Okubo, Y.; Akabane, T.: Phenotype i associated

with

[186716]congenital cataract in Japanese. Transfusion 19: 166–168,
1979.

[186717]

[186718]

[186719]19629.

[186720]Yeh, J.–C.; Ong, E.; Fukuda, M.: Molecular cloning and ex-
pression

[186721]of a novel beta–1,6–N–acetylglucosaminyltransferase that
forms core

[186722]2, core 4, and I branches. J. Biol. Chem. 274: 3215–3221,
1999.

[186723]

[186724]

[186725]19630.

[186726]Yu, L.–C.; Twu, Y.–C.; Chang, C.–Y.; Lin, M.: Molecular ba-
sis

[186727]of the adult i phenotype and the gene responsible for the
expression

[186728]of the human blood group I antigen. Blood 98:
3840–3845, 2001.

[186729]

[186730]

[186731]19631.

[186732]Allen, F. H., Jr.; Krabbe, S. M.; Corcoran, P. A.: A new phenotype
notype

[186733](McLeod) in the Kell blood-group system. Vox Sang. 6:
555-560, 1961.

[186734]

[186735]

[186736]19632.

[186737]Avent, N. D.; Martin, P. G.: Kell typing by allele-specific
PCR

[186738](ASP). Brit. J. Haemat. 93: 728-730, 1996.

[186739]

[186740]

[186741]19633.

[186742]Chown, B.; Lewis, M.; Kaita, K.: A 'new' Kell blood-group
phenotype. Nature 180:

[186743]711 only, 1957.

[186744]

[186745]

[186746]19634.

[186747]Conneally, P. M.; Nance, W. E.; Huntzinger, R. S.: Linkage
analysis

[186748]of Kell–Sutter and PTC loci. (Abstract) Am. J. Hum. Genet.
26: 22A

[186749]only, 1974.

[186750]

[186751]

[186752]19635.

[186753]Abdelhak, S.; Kalatzis, V.; Heilig, R.; Compain, S.; Samson,
D.;

[186754]Vincent, C.; Weil, D.; Cruaud, C.; Sahly, I.; Leibovici, M.;
Bitner–Glindzicz,

[186755]M.; Francis, M.; Lacombe, D.; Vigneron, J.; Charachon, R.;
Boven,

[186756]K.; Bedbeder, P.; Van Regemorter, N.; Weissenbach, J.; Pe-
tit, C.:

[186757]A human homologue of the Drosophila eyes absent gene
underlies branchio–oto–renal

[186758](BOR) syndrome and identifies a novel gene family. Nature
Genet. 15:
[186759]157–164, 1997.

[186760]

[186761]

[186762]19636.

[186763]Tommerup, N.; Schempp, W.; Meinecke, P.; Pedersen, S.;
Bolund,
[186764]L.; Brandt, C.; Goodpasture, C.; Guldberg, P.; Held, K.;
Reinwein,
[186765]H.; Saugstad, O. D.; Scherer, G.; Skjeldal, O.; Toder, R.;
Westvik,
[186766]J.; van der Hagen, C. B.; Wolf, U.: Assignment of an auto-
somal sex
[186767]reversal locus (SRA1) and campomelic dysplasia (CMPD1)
to 17q24.3–q25.1. Nature
[186768]Genet. 4: 170–174, 1993.

[186769]

[186770]

[186771]19637.

[186772]Azuma, N.; Nishina, S.; Yanagisawa, H.; Okuyama, T.; Yamada, M.

[186773]: PAX6 missense mutation in isolated foveal hypoplasia.
(Letter) Nature

[186774]Genet. 13: 141–142, 1996.

[186775]

[186776]

[186777]19638.

[186778]Curran, R. E.; Robb, R. M.: Isolated foveal hypoplasia.
Arch.

[186779]Ophthal. 94: 48–50, 1976.

[186780]

[186781]

[186782]19639.

[186783]O'Donnell, F. E., Jr.; Pappas, H. R.: Autosomal dominant
foveal

[186784]hypoplasia and presenile cataracts: a new syndrome. Arch.
Ophthal. 100:

[186785]279–281, 1982.

[186786]

[186787]

[186788]19640.

[186789]Anderson, M. J.; Shelton, G. D.; Cavenee, W. K.; Arden, K.
C.:

[186790]Embryonic expression of the tumor-associated
PAX3-FKHR fusion protein

[186791]interferes with the developmental functions of Pax3. Proc.
Nat. Acad.

[186792]Sci. 98: 1589–1594, 2001.

[186793]

[186794]

[186795]19641.

[186796]Anderson, M. J.; Viars, C. S.; Czekay, S.; Cavenee, W. K.;
Arden,

[186797]K. C.: Cloning and characterization of three human fork-
head genes

[186798]that comprise an FKHR-like gene subfamily. Genomics 47:
187–199,

[186799]1998.

[186800]

[186801]

[186802]19642.

[186803]Barr, F. G.; Galili, N.; Holick, J.; Biegel, J. A.; Rovera, G.;

[186804]Emanuel, B. S.: Rearrangement of the PAX3 paired box
gene in the

[186805]paediatric solid tumor alveolar rhabdomyosarcoma. Na-
ture Genet. 3:

[186806]113–117, 1993.

[186807]

[186808]

[186809]19643.

[186810]Davis, R. J.; Barr, F. G.: Fusion genes resulting from alter-
native

[186811]chromosomal translocations are overexpressed by gene-
specific mechanisms

[186812]in alveolar rhabdomyosarcoma. Proc. Nat. Acad. Sci. 94:
8047–8051,

[186813]1997.

[186814]

[186815]

[186816]19644.

[186817]Fredericks, W. J.; Galili, N.; Mukhopadhyay, S.; Rovera, G.;

Bennicelli,

[186818]J.; Barr, F. G.; Rauscher, F. J., III: The PAX3–FKHR fusion
protein

[186819]created by the t(2;13) translocation in alveolar rhab-
domyosarcomas

[186820]is a more potent transcriptional activator than PAX3.

Molec. Cell.

[186821]Biol. 15: 1522–1535, 1995.

[186822]

[186823]

[186824]19645.

[186825]Galili, N.; Davis, R. J.; Fredericks, W. J.; Mukhopadhyay, S.;

[186826]Rauscher, F. J., III; Emanuel, B. S.; Rovera, G.; Barr, F. G.:

Fusion

[186827]of a fork head domain gene to PAX3 in the solid tumour
alveolar rhabdomyosarcoma. Nature

[186828]Genet. 5: 230–235, 1993.

[186829]

[186830]

[186831]19646.

[186832]Khan, J.; Bittner, M. L.; Saal, L. H.; Teichmann, U.; Azorsa,
D.

[186833]O.; Gooden, G. C.; Pavan, W. J.; Trent, J. M.; Meltzer, P. S.:
cDNA

[186834]microarrays detect activation of a myogenic transcription
program

[186835]by the PAX3–FKHR fusion oncogene. Proc. Nat. Acad. Sci.
96: 13264–13269,

[186836]1999.

[186837]

[186838]

[186839]19647.

[186840]Angle, C. R.: Congenital bowing and angulation of the
long bones. Pediatrics 13:

[186841]257–268, 1954.

[186842]

[186843]

[186844]19648.

[186845]Bain, A. D.; Barrett, H. S.: Congenital bowing of the long bones:

[186846]report of a case. Arch. Dis. Child. 34: 516–524, 1959.

[186847]

[186848]

[186849]19649.

[186850]Bell, D. M.; Leung, K. K. H.; Wheatley, S. C.; Ng, L. J.; Zhou,

[186851]S.; Ling, K. W.; Sham, M. H.; Koopman, P.; Tam, P. P. L.;
Cheah, K.

[186852]S. E.: SOX9 directly regulates the type-II collagen gene.
Nature

[186853]Genet. 16: 174–178, 1997.

[186854]

[186855]

[186856]19650.

[186857]Bi, W.; Huang, W.; Whitworth, D. J.; Deng, J. M.; Zhang, Z.;
Behringer,

[186858]R. R.; de Crombrughe, B.: Haploinsufficiency of Sox9 re-
sults in

[186859]defective cartilage primordia and premature skeletal min-

eralization. Proc.

[186860]Nat. Acad. Sci. 98: 6698–6703, 2001.

[186861]

[186862]

[186863]19651.

[186864]Bishop, C. E.; Whitworth, D. J.; Qin, Y.; AgoulNIK, A. I.;
AgoulNIK,

[186865]I. U.; Harrison, W. R.; Behringer, R. R.; Overbeek, P. A.: A
transgenic

[186866]insertion upstream of Sox9 is associated with dominant
XX sex reversal

[186867]in the mouse. Nature Genet. 26: 490–494, 2000.

[186868]

[186869]

[186870]19652.

[186871]Caffey, J. P.: Prenatal bowing and thickening of tubular
bones,

[186872]with multiple cutaneous dimples in arms and legs: a con-
genital syndrome

[186873]of mechanical origin. Am. J. Dis. Child. 74: 543–562,

1947.

[186874]

[186875]

[186876]19653.

[186877]Cameron, F. J.; Hageman, R. M.; Cooke–Yarborough, C.;
Kwok, C.;

[186878]Goodwin, L. L.; Sillence, D. O.; Sinclair, A. H.: A novel germ
line

[186879]mutation in SOX9 causes familial campomelic dysplasia
and sex reversal. Hum.

[186880]Molec. Genet. 5: 1625–1630, 1996.

[186881]

[186882]

[186883]19654.

[186884]Cameron, F. J.; Sinclair, A. H.: Mutations in SRY and SOX9:
testis–determining

[186885]genes. Hum. Mutat. 9: 388–395, 1997.

[186886]

[186887]

[186888]19655.

[186889]Cooke, C. T.; Mulcahy, M. T.; Cullity, G. J.; Watson, M.;
Sprague,

[186890]P.: Campomelic dysplasia with sex reversal: morphological
and cytogenetic

[186891]studies of a case. Pathology 17: 526–529, 1985.

[186892]

[186893]

[186894]19656.

[186895]Cremin, B. J.; Orsmond, G.; Beighton, P.: Autosomal recessive
sive

[186896]inheritance in camptomelic dwarfism.(Letter) Lancet I:
488–489,

[186897]1973.

[186898]

[186899]

[186900]19657.

[186901]Dagna Bricarelli, F.; Fraccaro, M.; Lindsten, J.; Muller, U.;

[186902]Baggio, P.; Carbone, L. D. L.; Hjerpe, A.; Lindgren, F.;

Mayerova,

[186903]A.; Ringertz, H.; Ritzen, E. M.; Rovetta, D. C.; Sicchero, C.;

Wolf,

[186904]U.: Sex-reversed XY females with campomelic dysplasia

are H-Y negative. Hum.

[186905]Genet. 57: 15-22, 1981.

[186906]

[186907]

[186908]19658.

[186909]Ebensperger, C.; Jager, R. J.; Lattermann, U.; Dagna

Bricarelli,

[186910]F.; Keutel, J.; Lindsten, J.; Rehder, H.; Muller, U.; Wolf, U.:

No

[186911]evidence of mutations in four candidate genes for male

sex determination/differentiation

[186912]in sex-reversed XY females with campomelic dysplasia.

Ann. Genet. 34:

[186913]233-238, 1991.

[186914]

[186915]

[186916]19659.

[186917]Fontaine, G.; Walbaum, R.; Farriaux, J. P.; Tilmont, P.;

Peuzin,

[186918]F.; Delecour, M.: Le conseil genetique dans la dysplasie
campomelique

[186919](a propos de deux observations). J. Genet. Hum. 28:
267–279, 1980.

[186920]

[186921]

[186922]19660.

[186923]Foster, J. W.; Dominguez–Steglich, M. A.; Guioli, S.; Kwok,
C.;

[186924]Weller, P. A.; Stevanovic, M.; Weissenbach, J.; Mansour, S.;

Young,

[186925]I. D.; Goodfellow, P. N.; Brook, J. D.; Schafer, A. J.: Cam–
pomelic

[186926]dysplasia and autosomal sex reversal caused by mutations
in an SRY–related

[186927]gene. Nature 372: 525–530, 1994.

[186928]

[186929]

[186930]19661.

[186931]Friedrich, U.; Schaefer, E.; Meinecke, P.: Campomelic dys-
plasia

[186932]without overt campomelia. Clin. Dysmorph. 1: 172–178,
1992.

[186933]

[186934]

[186935]19662.

[186936]Gasca, S.; Canizares, J.; de Santa Barbara, P.; Mejean, C.;
Poulat,

[186937]F.; Berta, P.; Boizet–Bonhoure, B.: A nuclear export signal
within

[186938]the high mobility group domain regulates the nucleocyto-
plasmic translocation

[186939]of SOX9 during sexual determination. Proc. Nat. Acad. Sci.
99: 11199–11204,

[186940]2002.

[186941]

[186942]

[186943]19663.

[186944]Glass, R. B. J.; Rosenbaum, K. N.: Acampomelic cam-
pomelic dysplasia:

[186945]further radiographic variations. Am. J. Med. Genet. 69:
29–32, 1997.

[186946]

[186947]

[186948]19664.

[186949]Hall, B.; Spranger, J. W.: Campomelic dysplasia: further
elucidation

[186950]of a distinct entity. Am. J. Dis. Child. 134: 285–289, 1980.

[186951]

[186952]

[186953]19665.

[186954]Hoefnagel, D.; Wurster–Hill, D. H.; Dupree, W. B.;
Benirschke,

[186955]K.; Fuld, G. L.: Camptomelic dwarfism associated with XY–
gonadal

[186956]dysgenesis and chromosome anomalies. Clin. Genet. 13:
489–499, 1978.

[186957]

[186958]

[186959]19666.

[186960]Houston, C. S.; Opitz, J. M.; Spranger, J. W.; Macpherson, R.

[186961]I.; Reed, M. H.; Gilbert, E. F.; Herrmann, J.; Schinzel, A.:
The

[186962]campomelic syndrome: review, report of 17 cases, and
follow-up on

[186963]the currently 17-year-old boy first reported by Maroteaux
et al in

[186964]1971. Am. J. Med. Genet. 15: 3–28, 1983.

[186965]

[186966]

[186967]19667.

[186968]Hovmoller, M. L.; Osuna, A.; Eklof, O.; Fredga, K.; Hjerpe, A.;

[186969]Lindsten, J.; Ritzen, M.; Stanescu, V.; Svenningsen, N.:
Camptomelic

[186970]dwarfism. A genetically determined mesenchymal disorder
combined with

[186971]sex reversal. Hereditas 86: 51–62, 1977.

[186972]

[186973]

[186974]19668.

[186975]Huang, W.; Chung, U.; Kronenberg, H. M.; de Crombrughe, B.:

[186976]The chondrogenic transcription factor Sox9 is a target of signaling

[186977]by the parathyroid hormone-related peptide in the growth plate of

[186978]endochondral bones. Proc. Nat. Acad. Sci. 98: 160–165, 2001.

[186979]

[186980]

[186981]19669.

[186982]Kanai, Y.; Koopman, P.: Structural and functional characterization

[186983]of the mouse Sox9 promoter: implications for campomelic dysplasia. Hum.

[186984]Molec. Genet. 8: 691–696, 1999.

[186985]

[186986]

[186987]19670.

[186988]Medema, R. H.; Kops, G. J. P. L.; Bos, J. L.; Burgering, B. M.

[186989]T.: AFX-like forkhead transcription factors mediate cell-cycle regulation

[186990]by Ras and PKB through p27(kip1). Nature 404: 782–787, 2000.

[186991]

[186992]

[186993]19671.

[186994]Eicher, E. M.; Womack, J. E.: Chromosomal location of soluble

[186995]glutamic-pyruvic transaminase-1 (Gpt-1) in the mouse. Biochem. Genet. 15:

[186996]1–8, 1977.

[186997]

[186998]

[186999]19672.

[187000]Falk, C. T.; Huss, J.: Linkage data on the chromosome 16 markers

[187001]HP and PGP: no additional support for or against the mapping of GPT.

[187002](Abstract) Cytogenet. Cell Genet. 40: 626 only, 1985.

[187003]

[187004]

[187005]19673.

[187006]Ferrell, R. E.: Personal Communication. Pittsburgh, Pa.
1/22/1988.

[187007]

[187008]

[187009]19674.

[187010]Jeremiah, S. J.; Kielty, C.; Povey, S.; McLellan, T.: Immuno-
logical

[187011]characterization of human GPT in hybrids supports as-
signment to chromosome

[187012]

[187013]19675.

[187014](Abstract) Cytogenet. Cell Genet. 37: 498 only, 1984.

[187015]

[187016]8. Kalimanovska, V.; Majkic-Singh, N.; Stojanov, M.; Groz-
danic, V.;

[187017]Vucetic, G.; Andelic, M.; Gligorovic, V.; Tomasevic, R.: Human red

[187018]cell glutamic-pyruvic transaminase polymorphism in Serbia, Yugoslavia. Hum.

[187019]Hered. 33: 319-321, 1983.

[187020]

[187021]

[187022]19676.

[187023]Kielty, C. M.; Povey, S.; Hopkinson, D. A.: Regulation of expression

[187024]of liver-specific enzymes: II. Activation and chromosomal localization

[187025]of soluble glutamate-pyruvate transaminase. Ann. Hum. Genet. 46:

[187026]135-143, 1982.

[187027]

[187028]

[187029]19677.

[187030]King, M.-C.; Go, R. C. P.; Elston, R. C.; Lynch, H. T.; Petrakis,

[187031]N. L.: Allele increasing susceptibility to human breast cancer may

[187032]be linked to the glutamate-pyruvate transaminase locus.

Science 208:

[187033]406-408, 1980.

[187034]

[187035]

[187036]19678.

[187037]Kompf, J.: Population genetics of soluble glutamic-pyruvic transaminase

[187038](EC:2.6.1.2): gene frequencies in southwestern Germany.

Humangenetik 14:

[187039]76-77, 1971.

[187040]

[187041]

[187042]19679.

[187043]Lahav, M.; Szeinberg, A.: A red-cell glutamic-pyruvic transaminase

[187044]polymorphism in several population groups in Israel.

Hum. Hered. 22:

[187045]533–538, 1972.

[187046]

[187047]

[187048]19680.

[187049]Marazita, M. L.; Spence, M. A.; Sparkes, R. S.; Field, L. L.;

[187050]Crandall, B. F.; Sparkes, M. C.; Crist, M.: Linkage relations
of

[187051]GPT (glutamic–pyruvate transaminase). (Abstract) Cyto–
genet. Cell

[187052]Genet. 40: 690 only, 1985.

[187053]

[187054]

[187055]19681.

[187056]McLellan, T.: Two previously undetected variants of glu–
tamic–pyruvic

[187057]transaminase found by acidic polyacrylamide gel elec–
trophoresis. Am.

[187058]J. Hum. Genet. 34: 623–628, 1982.

[187059]

[187060]

[187061]19682.

[187062]McLellan, T.; Cannon, L. A.; Bishop, D. T.; Skolnick, M. H.:

[187063]The cumulative lod score between a breast cancer susceptibility locus

[187064]and GPT is -3.86 . (Abstract) Cytogenet. Cell Genet. 37: 536-537,

[187065]1984.

[187066]

[187067]

[187068]19683.

[187069]Mithal, Y.; Lane, A. B.; Jenkins, T.: Absence of red cell glutamic-pyruvate

[187070]transaminase: discovery of a 'silent' allele homozygote. Am. J. Hum.

[187071]Genet. 32: 42-46, 1980.

[187072]

[187073]

[187074]19684.

[187075]O'Connell, P.; Nakamura, Y.; Lathrop, G. M.; Leppert, M.;
Cartwright,

[187076]P.; Lalouel, J.-M.; White, R.: Three genetic linkage groups
on chromosome

[187077]8. (Abstract) Cytogenet. Cell Genet. 46: 672 only, 1987.

[187078]

[187079]

[187080]19685.

[187081]Olaisen, B.: Genetics of the GPT system: family, mother-
child

[187082]and association studies. Clin. Genet. 7: 245-254, 1975.

[187083]

[187084]

[187085]19686.

[187086]Pelzer, C. F.; Norum, R. A.: Identification of human red cell

[187087]glutamate-pyruvate transaminase (GPT) phenotypes by
isoelectric focusing. Am.

[187088]J. Hum. Genet. 37: 147-152, 1985.

[187089]

[187090]

[187091]19687.

[187092]Rocha, J.; Amorim, A.; Almeida, V. M.; Oliveira, J. P.; Leao,

[187093]M.; Tavares, M. C.; Pereira, M. S.; Vidal-Pinheiro, L.: Gene
dosage

[187094]evidence for the regional assignment of GPT

(glutamate-pyruvate transaminase;

[187095]E.C. 2.6.1.2) locus to 8q24.2-8qter. Hum. Genet. 80:
299-300, 1988.

[187096]

[187097]

[187098]19688.

[187099]Sanders, M. F.; King, M. C.; Lattanzio, D.; Crandall, J.; Le-
ung,

[187100]R.: Absence of linkage between HP and GPT. (Abstract)
Cytogenet.

[187101]Cell Genet. 37: 536-537, 1984.

[187102]

[187103]

[187104]19689.

[187105]Santachiara Benerecetti, A. S.; Beretta, M.; Pampiglione, S.:

[187106]Red cell glutamic–pyruvic transaminase polymorphism in
a sample of

[187107]the Italian population: a new variant allele: GPT(8). Hum.
Hered. 25:

[187108]276–278, 1975.

[187109]

[187110]

[187111]19690.

[187112]Sohocki, M. M.; Sullivan, L. S.; Harrison, W. R.; Sodergren,
E.

[187113]J.; Elder, F. F. B.; Weinstock, G.; Tanase, S.; Daiger, S. P.:
Human

[187114]glutamate pyruvate transaminase (GPT): localization to
8q24.3, cDNA

[187115]and genomic sequences, and polymorphic sites. Genomics
40: 247–252,

[187116]1997.

[187117]

[187118]

[187119]19691.

[187120]Sparkes, M. C.; Crist, M.; Sparkes, R. S.: Glutamate pyruvate

[187121]transaminase null allele in seven new families. Hum. Genet. 65:

[187122]147–148, 1983.

[187123]

[187124]

[187125]19692.

[187126]Wijnen, L. M. M.; Meera Khan, P.: Assignment of GPT to human

[187127]chromosome 16. (Abstract) Cytogenet. Cell Genet. 32: 327 only, 1982.

[187128]

[187129]

[187130]19693.

[187131]Burant, C. F.; Takeda, J.; Brot-Laroche, E.; Bell, G. I.; Davidson,

[187132]N. O.: Fructose transporter in human spermatozoa and small intestine

[187133]is GLUT5. J. Biol. Chem. 267: 14523–14526, 1992.

[187134]

[187135]

[187136]19694.

[187137]Davidson, N. O.; Hausman, A. M. L.; Ifkovits, C. A.; Buse, J. B.;

[187138]Gould, G. W.; Burant, C. F.; Bell, G. I.: Human intestinal glucose

[187139]transporter expression and localization of GLUT5. Am. J. Physiol. 262:

[187140]C795–C800, 1992.

[187141]

[187142]

[187143]19695.

[187144]Cremonini, N.; Graziano, E.; Chiarini, V.; Sforza, A.; Zampa,

[187145]G. A.: Atypical McCune–Albright syndrome associated with growth hormone–prolactin

[187146]pituitary adenoma: natural history, long-term follow-up, and SMS 201–995--bromocriptine

[187147]combined treatment results. J. Clin. Endocr. Metab. 75:
1166–1169,
[187148]1992.

[187149]

[187150]

[187151]19696.

[187152]DeChiara, T.; Robertson, E. J.; Efstratiadis, A.: Parental im-
printing

[187153]of the mouse insulin–like growth factor II gene. Cell 64:
849–859,

[187154]1991.

[187155]

[187156]

[187157]19697.

[187158]Eddy, M. C.; Jan de Beur, S. M.; Yandow, S. M.; McAlister,
W.

[187159]H.; Shore, E. M.; Kaplan, F. S.; Whyte, M. P.; Levine, M. A.:
Deficiency

[187160]of the alpha–subunit of the stimulatory G protein and se-
vere extraskeletal

[187161]ossification. J. Bone Miner. Res. 15: 2074–2083, 2000.

[187162]

[187163]

[187164]19698.

[187165]Falconer, M. A.; Cope, C. L.; Robb–Smith, A. H. T.: Fibrous
dysplasia

[187166]of bone with endocrine disorders and cutaneous pigmen-
tation (Albright's

[187167]disease). Quart. J. Med. 11: 121–154, 1942.

[187168]

[187169]

[187170]19699.

[187171]Albertsen, H. M.; Smith, S. A.; Mazoyer, S.; Fujimoto, E.;
Stevens,

[187172]J.; Williams, B.; Rodriguez, P.; Cropp, C. S.; Slijepcevic, P.;
Carlson,

[187173]M.; Robertson, M.; Bradley, P.; Lawrence, E.; Harrington,
T.; Mei

[187174]Sheng, Z.; Hoopes, R.; Sternberg, N.; Brothman, A.; Calla-
han, R.;

[187175]Ponder, B. A. J.; White, R.: A physical map and candidate genes in

[187176]the BRCA1 region on chromosome 17q12–21. Nature Genet. 7: 472–479,

[187177]1994.

[187178]

[187179]

[187180]19700.

[187181]Khatib, Z. A.; Inaba, T.; Valentine, M.; Look, A. T.: Chromosomal

[187182]localization and cDNA cloning of the human DBP and TEF genes. Genomics 23:

[187183]344–351, 1994.

[187184]

[187185]

[187186]19701.

[187187]Pilla, M.; Perachon, S.; Sautel, F.; Garrido, F.; Mann, A.; Wermuth,

[187188]C. G.; Schwartz, J.–C.; Everitt, B. J.; Sokoloff, P.: Selective inhibition

[187189]of cocaine-seeking behaviour by a partial dopamine D3
receptor agonist. Nature 400:

[187190]371–375, 1999.

[187191]

[187192]

[187193]19702.

[187194]Demange, P.; Voges, D.; Benz, J.; Liemann, S.; Gottig, P.;
Berendes,

[187195]R.; Burger, A.; Huber, R.: Annexin V: the key to under-
standing ion

[187196]selectivity and voltage regulation? Trends Biochem. Sci.
19: 272–276,

[187197]1994.

[187198]

[187199]

[187200]19703.

[187201]Funakoshi, T.; Heimark, R. L.; Hendrickson, L. E.; Mc-
Mullen, B.

[187202]A.; Fujikawa, K.: Human placental anticoagulant protein:
isolation

[187203]and characterization. Biochemistry 26: 5572–5578, 1987.

[187204]

[187205]

[187206]19704.

[187207]Grundmann, U.; Abel, K.-J.; Bohn, H.; Lobermann, H.;
Lottspeich,

[187208]F.; Kupper, H.: Characterization of cDNA encoding human
placental

[187209]anticoagulant protein (PP4): homology with the lipocortin
family. Proc.

[187210]Nat. Acad. Sci. 85: 3708–3712, 1988.

[187211]

[187212]

[187213]19705.

[187214]Kaplan, R.; Jaye, M.; Burgess, W. H.; Schlaepfer, D. D.;
Haigler,

[187215]H. T.: Cloning and expression of cDNA for human en-
donexin II, a Ca(2+)

[187216]and phospholipid binding protein. J. Biol. Chem. 263:
8037–8043,

[187217]1988.

[187218]

[187219]

[187220]19706.

[187221]Modi, W. S.; Seuanez, H.; Jaye, M.; Kaplan, R.; Haigler, H.;
O'Brien,

[187222]S. J.: Chromosomal mapping of the endonexin II gene.
(Abstract) Cytogenet.

[187223]Cell Genet. 51: 1046, 1989.

[187224]

[187225]

[187226]19707.

[187227]Modi, W. S.; Seuanez, H. N.; Jaye, M.; Haigler, H. J.; Kaplan,

[187228]R.; O'Brien, S. J.: The human endonexin II (ENX2) gene is
located

[187229]at 4q28–q32. Cytogenet. Cell Genet. 52: 167–169, 1989.

[187230]

[187231]

[187232]19708.

[187233]Rodriguez–Garcia, M. I.; Kozak, C. A.; Morgan, R. O.; Fernandez,

[187234]M. P.: Mouse annexin V chromosomal localization, cDNA sequence conservation,

[187235]and molecular evolution. *Genomics* 31: 151–157, 1996.

[187236]

[187237]

[187238]19709.

[187239]Tait, J. F.; Frankenberry, D. A.; Shiang, R.; Murray, J. C.; Adler,

[187240]D. A.; Disteché, C. M.: Chromosomal localization of the human gene

[187241]for annexin V (placental anticoagulant protein I) to 4q26–q28. *Cytogenet.*

[187242]*Cell Genet.* 57: 187–192, 1991.

[187243]

[187244]

[187245]19710.

[187246]Arinami, T.; Ishikawa, M.; Inoue, A.; Yanagisawa, M.; Masaki, T.;

[187247]Yoshida, M. C.; Hamaguchi, H.: Chromosomal assignments of the human

[187248]endothelin family genes: the endothelin-1 gene (EDN1) to 6p23-p24,

[187249]the endothelin-2 gene (EDN2) to 1p34, and the endothelin-3 gene (EDN3)

[187250]to 20q13.2-q13.3. Am. J. Hum. Genet. 48: 990-996, 1991.

[187251]

[187252]

[187253]19711.

[187254]Benatti, L.; Bonecchi, L.; Cozzi, L.; Sarmientos, P.: Two preproendothelin

[187255]1 mRNAs transcribed by alternative promoters. J. Clin. Invest. 91:

[187256]1149-1156, 1993.

[187257]

[187258]

[187259]19712.

[187260]Berge, K. E.; Berg, K.: No effect of a TaqI polymorphism in DNA

[187261]at the endothelin I (EDN1) locus on normal blood pressure level or

[187262]variability. Clin. Genet. 41: 90–95, 1992.

[187263]

[187264]

[187265]19713.

[187266]Bloch, K. D.; Friedrich, S. P.; Lee, M.–E.; Eddy, R. L.; Shows,

[187267]T. B.; Quertermous, T.: Structural organization and chromosomal assignment

[187268]of the gene encoding endothelin. J. Biol. Chem. 264: 10851–10857,

[187269]1989.

[187270]

[187271]

[187272]19714.

[187273]Bourgeois, C.; Robert, B.; Rebourcet, R.; Mondon, F.; Mignot, T.–M.;

[187274]Duc–Goiran, P.; Ferre, F.: Endothelin–1 and ET(A) receptor expression

[187275]in vascular smooth muscle cells from human placenta: a

new ET(A) receptor

[187276] messenger ribonucleic acid is generated by alternative
splicing of

[187277] exon 3. J. Clin. Endocr. Metab. 82: 3116–3123, 1997.

[187278]

[187279]

[187280] 19715.

[187281] Clouthier, D. E.; Hosoda, K.; Richardson, J. A.; Williams, S.
C.;

[187282] Yanagisawa, H.; Kuwaki, T.; Kumada, M.; Hammer, R. E.;
Yanagisawa,

[187283] M.: Cranial and cardiac neural crest defects in endothelin–
A receptor–deficient

[187284] mice. Development 125: 813–824, 1998.

[187285]

[187286]

[187287] 19716.

[187288] Giaid, A.; Gibson, S. J.; Ibrahim, N. B. N.; Legon, S.; Bloom,

[187289] S. R.; Yanagisawa, M.; Masaki, T.; Varndell, I. M.; Polak, J.
M.:

[187290]Endothelin 1, an endothelium-derived peptide, is expressed in neurons

[187291]of the human spinal cord and dorsal root ganglia. Proc. Nat. Acad.

[187292]Sci. 86: 7634–7638, 1989.

[187293]

[187294]

[187295]19717.

[187296]Inoue, A.; Yanagisawa, M.; Kimura, S.; Kasuya, Y.; Miyauchi, T.;

[187297]Goto, K.; Masaki, T.: The human endothelin family: three structurally

[187298]and pharmacologically distinct isopeptides predicted by three separate

[187299]genes. Proc. Nat. Acad. Sci. 86: 2863–2867, 1989.

[187300]

[187301]

[187302]19718.

[187303]Inoue, A.; Yanagisawa, M.; Takuwa, Y.; Mitsui, Y.; Kobayashi, M.;

[187304]Masaki, T.: The human preproendothelin-1 gene: complete nucleotide

[187305]sequence and regulation of expression. J. Biol. Chem. 264: 14954-14959,

[187306]1989.

[187307]

[187308]

[187309]19719.

[187310]Shaikh, S.; Ball, D.; Craddock, N.; Castle, D.; Hunt, N.; Mant,

[187311]R.; Owen, M.; Collier, D.; Gill, M.: The dopamine D3 receptor gene:

[187312]no association with bipolar affective disorder. J. Med. Genet. 30:

[187313]308-309, 1993.

[187314]

[187315]

[187316]19720.

[187317]Sokoloff, P.; Giros, B.; Martres, M.-P.; Bouthenet, M.-L.; Schwartz,

[187318]J.-C.: Molecular cloning and characterization of a novel dopamine

[187319]receptor (D-3) as a target for neuroleptics. Nature 347: 146-151,

[187320]1990.

[187321]

[187322]

[187323]19721.

[187324]Spurlock, G.; Williams, J.; McGuffin, P.; Aschauer, H. N.; Lenzinger,

[187325]E.; Fuchs, K.; Sieghart, W. C.; Meszaros, K.; Fathi, N.; Laurent,

[187326]C.; Mallet, J.; Macciardi, F.; Pedrini, S.; Gill, M.; Hawi, Z.; Gibson,

[187327]S.; Jazin, E. E.; Yang, H.-T.; Adolfsson, R.; Pato, C. N.; Dourado,

[187328]A. M.; Owen, M. J.: European Multicentre Association Study of Schizophrenia:

[187329]a study of the DRD2 ser311-to-cys and DRD3 ser9-to-gly polymorphisms. Am.

[187330]J. Med. Genet. 81: 24-28, 1998.

[187331]

[187332]

[187333]19722.

[187334]Asghari, V.; Sanyal, S.; Buchwaldt, S.; Paterson, A.; Jo-
vanovic,

[187335]V.; Van Tol, H. H.: Modulation of intracellular cyclic AMP
levels

[187336]by different human dopamine D4 receptor variants. J.
Neurochem. 65:

[187337]1157–1165, 1995.

[187338]

[187339]

[187340]19723.

[187341]Benjamin, J.; Li, L.; Patterson, C.; Greenberg, B. D.; Mur-
phy,

[187342]D. L.; Hamer, D. H.: Population and familial association
between

[187343]the D4 dopamine receptor gene and measures of novelty
seeking. Nature

[187344]Genet. 12: 81–84, 1996.

[187345]

[187346]

[187347]19724.

[187348]Castellanos, F. X.; Lau, E.; Tayebi, N.; Lee, P.; Long, R. E.;

[187349]Giedd, J. N.; Sharp, W.; Marsh, W. L.; Walter, J. M.; Ham-
burger, S.

[187350]D.; Ginns, E. I.; Rapoport, J. L.; Sidransky, E.: Lack of an
association

[187351]between a dopamine-4 receptor polymorphism and atten-
tion-deficit/hyperactivity

[187352]disorder: genetic and brain morphometric analyses.

Molec. Psychiat. 3:

[187353]431-434, 1998.

[187354]

[187355]

[187356]19725.

[187357]Chang, F.-M.; Kidd, J. R.; Livak, K. J.; Pakstis, A. J.; Kidd,

[187358]K. K.: The world-wide distribution of allele frequencies at
the human

[187359]dopamine D4 receptor locus. Hum. Genet. 98: 91-101,

1996.

[187360]

[187361]

[187362]19726.

[187363]Cloninger, C. R.; Adolfsson, R.; Svrakic, N. M.: Mapping
genes

[187364]for human personality. Nature Genet. 12: 3–4, 1996.

[187365]

[187366]

[187367]19727.

[187368]Cloninger, C. R.; Svrakic, D. M.; Przybeck, T. R.: A psy-
chobiological

[187369]model of temperament and character. Arch. Gen. Psychiat.
50: 975–990,

[187370]1993.

[187371]

[187372]

[187373]19728.

[187374]Cook, E. H.; Stein, M. A.; Krasowski, M. D.; Cox, N. J.;

Olkon,

[187375]D. M.; Kieffer, J. E.; Leventhal, B. L.: Association of attention-deficit

[187376]disorder and the dopamine transporter gene. Am. J. Hum. Genet. 56:

[187377]993–998, 1995.

[187378]

[187379]

[187380]19729.

[187381]De Luca, A.; Rizzardi, M.; Torrente, I.; Alessandroni, R.; Salvioli,

[187382]G. P.; Filograsso, N.; Dallapiccola, B.; Novelli, G.:

Dopamine D4

[187383]receptor (DRD4) polymorphism and adaptability trait during infancy:

[187384]a longitudinal study in 1- to 5-month-old neonates. Neurogenetics 3:

[187385]79–82, 2001.

[187386]

[187387]

[187388]19730.

[187389]Ding, Y.-C.; Chi, H.-C.; Grady, D. L.; Morishima, A.; Kidd, J.

[187390]R.; Kidd, K. K.; Flodman, P.; Spence, M. A.; Schuck, S.; Swanson,

[187391]J. M.; Zhang, Y.-P.; Moyzis, R. K.: Evidence of positive selection

[187392]acting at the human dopamine receptor D4 gene locus. Proc. Nat. Acad.

[187393]Sci. 99: 309–314, 2002.

[187394]

[187395]

[187396]19731.

[187397]Ebstein, R. P.; Novick, O.; Umansky, R.; Pirelli, B.; Osher, Y.;

[187398]Blaine, D.; Bennett, E. R.; Nemanov, L.; Katz, M.; Belmaker, R. H.

[187399]: Dopamine D4 receptor exon III polymorphism associated with the human

[187400]personality trait of novelty seeking. Nature Genet. 12: 78–80, 1996.

[187401]

[187402]

[187403]19732.

[187404]Eisenberg, J.; Zohar, A.; Mei-Tal, G.; Steinberg, A.; Tar-
takovsky,

[187405]E.; Gritsenko, I.; Nemanov, L.; Ebstein, R. P.: A haplotype
relative

[187406]risk study of the dopamine D4 receptor (DRD4) exon III
repeat polymorphism

[187407]and attention deficit hyperactivity disorder (ADHD). Am. J.
Med.

[187408]Genet. (Neuropsychiat. Genet.) 96: 258–261, 2000.

[187409]

[187410]

[187411]19733.

[187412]Gelernter, J.; Kennedy, J. L.; Van Tol, H. H. M.; Civelli, O.;

[187413]Kidd, K. K.: The D4 dopamine receptor (DRD4) maps to
distal 11p close

[187414]to HRAS. Genomics 13: 208–210, 1992.

[187415]

[187416]

[187417]19734.

[187418]Gelernter, J.; Kennedy, J. L.; Van Tol, H. H. M.; Niznik, H. B.;

[187419]Civelli, O.; Kidd, K. K.: The D4 dopamine receptor (DRD4) maps to

[187420]distal 11p close to HRAS. (Abstract) Cytogenet. Cell Genet. 58:

[187421]1960, 1991.

[187422]

[187423]

[187424]19735.

[187425]Gelernter, J.; Kranzler, H.; Coccaro, E.; Siever, L.; New, A.;

[187426]Mulgrew, C. L.: D4 dopamine-receptor (DRD4) alleles and novelty seeking

[187427]in substance-dependent, personality-disorder, and control subjects. Am.

[187428]J. Hum. Genet. 61: 1144–1152, 1997.

[187429]

[187430]

[187431]19736.

[187432]Harpending, H.; Cochran, G.: In our genes. Proc. Nat. Acad.

[187433]Sci. 99: 10–12, 2002.

[187434]

[187435]

[187436]19737.

[187437]Jovanovic, V.; Guan, H.–C.; Van Tol, H. H. M.: Comparative pharmacological

[187438]and functional analysis of the human dopamine D4.2 and D4.10 receptor

[187439]variants. Pharmacogenetics 9: 561–568, 1999.

[187440]

[187441]

[187442]19738.

[187443]LaHoste, G. J.; Swanson, J. M.; Wigal, S. B.; Glabe, C.; Wi-gal,

[187444]T.; King, N.; Kennedy, J. L.: Dopamine D4 receptor gene polymorphism

[187445]is associated with attention deficit hyperactivity disorder. Molec.

[187446]Psychiat. 1: 121–124, 1996.

[187447]

[187448]

[187449]19739.

[187450]Liu, I. S. C.; Seeman, P.; Sanyal, S.; Ulpian, C.; Rodgers–
Johnson,

[187451]P. E. B.; Serjeant, G. R.; Van Tol, H. H. M.: Dopamine D4
receptor

[187452]variant in Africans, D4(valine194glycine), is insensitive to
dopamine

[187453]and clozapine: report of a homozygous individual. Am. J.
Med. Genet. 61:

[187454]277–282, 1996.

[187455]

[187456]

[187457]19740.

[187458]Malhotra, A. K.; Virkkunen, M.; Rooney, W.; Eggert, M.;
Linnoila,

[187459]M.; Goldman, D.: The association between the dopamine
D(4) receptor

[187460](D4DR) 16 amino acid repeat polymorphism and novelty seeking. Molec.

[187461]Psychiat. 1: 388–391, 1996.

[187462]

[187463]

[187464]19741.

[187465]Muramatsu, T.; Higuchi, S.; Murayama, M.; Matsushita, S.; Hayashida,

[187466]M.: Association between alcoholism and the dopamine D4 receptor gene. J.

[187467]Med. Genet. 33: 113–115, 1996.

[187468]

[187469]

[187470]19742.

[187471]Nothen, M. M.; Cichon, S.; Hemmer, S.; Hebebrand, J.; Remschmidt,

[187472]H.; Lehmkuhl, G.; Poustka, F.; Schmidt, M.; Catalano, M.; Fimmers,

[187473]R.; Korner, J.; Rietschel, M.; Propping, P.: Human dopamine D4 receptor

[187474]gene: frequent occurrence of a null allele and observation of homozygosity. Hum.

[187475]Molec. Genet. 3: 2207–2212, 1994.

[187476]

[187477]

[187478]19743.

[187479]Petronis, A.; Van Tol, H. H. M.; Lichter, J. B.; Livak, K. J.;

[187480]Kennedy, J. L.: The D4 dopamine receptor gene maps on 11p proximal

[187481]to HRAS. Genomics 18: 161–163, 1993.

[187482]

[187483]

[187484]19744.

[187485]Rowe, D. C.; Stever, C.; Giedinghagen, L. N.; Gard, J. M. C.;

[187486]Cleveland, H. H.; Terris, S. T.; Mohr, J. H.; Sherman, S.; Abramowitz,

[187487]A.; Waldman, I. D.: Dopamine DRD4 receptor polymorphism and attention

[187488]deficit hyperactivity disorder. Molec. Psychiat. 3: 419–426,

1998.

[187489]

[187490]

[187491]19745.

[187492]Rubinstein, M.; Phillips, T. J.; Bunzow, J. R.; Falzone, T. L.;

[187493]Dziewczapolski, G.; Zhang, G.; Fang, Y.; Larson, J. L.; Mc-
Dougall,

[187494]J. A.; Chester, J. A.; Saez, C.; Pugsley, T. A.; Gershanik, O.;
Low,

[187495]M. J.; Grandy, D. K.: Mice lacking dopamine D4 receptors
are supersensitive

[187496]to ethanol, cocaine, and methamphetamine. Cell 90:
991–1001, 1997.

[187497]

[187498]

[187499]19746.

[187500]Barsh, G. S.; Seeburg, P. H.; Gelinas, R. E.: The human
growth

[187501]hormone gene family:structure and evolution of the chro-
mosomal locus. Nucleic

[187502]Acids Res. 11: 3939–3958, 1983.

[187503]

[187504]

[187505]19747.

[187506]Cremers, F. P. M.; Molloy, C. M.; van de Pol, D. J. R.; van
den

[187507]Hurk, J. A. J. M.; Bach, I.; Geurts van Kessel, A. H. M.; Rop-
ers,

[187508]H.-H.: An autosomal homologue of the choroideremia
gene colocalizes

[187509]with the Usher syndrome type II locus on the distal part of
chromosome

[187510]1q. Hum. Molec. Genet. 1: 71–75, 1992.

[187511]

[187512]

[187513]19748.

[187514]Halford, S.; Freedman, M. S.; Bellingham, J.; Inglis, S. L.;
Poopalasundaram,

[187515]S.; Soni, B. G.; Foster, R. G.; Hunt, D. M.: Characterization
of

[187516]a novel human opsin gene with wide tissue expression
and identification

[187517]of embedded and flanking genes on chromosome 1q43.

Genomics 72:

[187518]203–208, 2001.

[187519]

[187520]

[187521]19749.

[187522]van Bokhoven, H.; van Genderen, C.; Molloy, C. M.; van de
Pol,

[187523]D. J. R.; Cremers, C. W. R. J.; van Aarem, A.; Schwartz, M.;
Rosenberg,

[187524]T.; Geurts van Kessel, A. H. M.; Ropers, H.–H.; Cremers, F.
P. M.

[187525]: Mapping of the choroideremia–like (CHML) gene at
1q42–qter and mutation

[187526]analysis in patients with Usher syndrome type II. Ge-
nomics 19: 385–387,

[187527]1994.

[187528]

[187529]

[187530]19750.

[187531]Amato, F.; Warnes, G. M.; Kirby, C. A.; Norman, R. J.: Infer-
tility

[187532]caused by hCG autoantibody. J. Clin. Endocr. Metab. 87:
993-997,

[187533]2002.

[187534]

[187535]

[187536]19751.

[187537]Boothby, M.; Ruddon, R. W.; Anderson, C.; McWilliams, D.;
Boime,

[187538]I.: A single gonadotropin alpha-subunit gene in normal
tissue and

[187539]tumor-derived cell lines. J. Biol. Chem. 256: 5121-5127,
1981.

[187540]

[187541]

[187542]19752.

[187543]Bordelon, M.; Kohler, P. O.: Synthesis of human glycoprotein hormone

[187544]in somatic cell hybrids. (Abstract) J. Cell Biol. 67: 37A only,

[187545]1975.

[187546]

[187547]

[187548]19753.

[187549]Bordelon-Riser, M. E.; Siciliano, M. J.; Kohler, P. O.: Necessity

[187550]for two human chromosomes for human chorionic gonadotropin production

[187551]in human-mouse hybrids. Somat. Cell Genet. 5: 597-613, 1979.

[187552]

[187553]

[187554]19754.

[187555]Chin, W. W.: Personal Communication. Boston, Mass. 2/15/1982.

[187556]

[187557]

[187558]19755.

[187559]Fiddes, J. C.; Goodman, H. M.: Isolation, cloning and sequence

[187560]analysis of the cDNA for the alpha-subunit of human chorionic gonadotropin. Nature 281:

[187561]351-356, 1979.

[187562]

[187563]

[187564]19756.

[187565]Fiddes, J. C.; Goodman, H. M.: The gene encoding the common alpha

[187566]subunit of the four human glycoprotein hormones. J. Molec. Appl.

[187567]Genet. 1: 3-18, 1981.

[187568]

[187569]

[187570]19757.

[187571]Hardin, J. W.; Riser, M. E.; Trent, J. M.; Kohler, P. O.: The

[187572]chorionic gonadotropin alpha-subunit gene is on human
chromosome 18

[187573]in JEG cells. Proc. Nat. Acad. Sci. 80: 6282–6285, 1983.

[187574]

[187575]

[187576]19758.

[187577]Heitz, P. U.; Kasper, M.; Kloppel, G.; Polak, J. M.;
Vaitukaitis,

[187578]J. L.: Glycoprotein-hormone alpha-chain production by
pancreatic

[187579]endocrine tumors: a specific marker for malignancy–
–immunocytochemical

[187580]analysis of tumors of 155 patients. Cancer 51: 277–282,
1983.

[187581]

[187582]

[187583]19759.

[187584]Hoshina, M.; Boothby, M. R.; Hussa, R. D.; Pattillo, R. A.;
Camel,

[187585]H. M.; Boime, I.: Segregation patterns of polymorphic re-
striction

[187586]sites of the gene encoding the alpha subunit of human chorionic gonadotropin

[187587]in trophoblastic disease. Proc. Nat. Acad. Sci. 81: 2504–2507, 1984.

[187588]

[187589]

[187590]19760.

[187591]Lebo, R. V.: Personal Communication. San Francisco, Calif.

[187592]1/14/1980.

[187593]

[187594]

[187595]19761.

[187596]Morgan, F. J.; Birken, S.; Canfield, R. E.: The amino acid sequence

[187597]of human chorionic gonadotropin: the alpha subunit and beta subunit. J.

[187598]Biol. Chem. 250: 5247–5258, 1975.

[187599]

[187600]

[187601]19762.

[187602]Naylor, S. L.; Chin, W. W.; Goodman, H. M.; Lalley, P. A.;
Grzeschik,

[187603]K.-H.; Sakaguchi, A. Y.: Chromosome assignment of the
genes encoding

[187604]the alpha and beta subunits of the glycoprotein hormones
in man and

[187605]mouse. Somat. Cell Genet. 9: 757–770, 1983.

[187606]

[187607]

[187608]19763.

[187609]Ruddon, R. W.; Hanson, C. A.; Addison, N. J.: Synthesis
and processing

[187610]of human chorionic gonadotropin subunits in cultured
choriocarcinoma

[187611]cells. Proc. Nat. Acad. Sci. 76: 5143–5147, 1979.

[187612]

[187613]

[187614]19764.

[187615]Trent, J.: Personal Communication. Tucson, Ariz.

11/23/1982.

[187616]

[187617]

[187618]19765.

[187619]Hart, A. W.; Baeza, N.; Apelqvist, A.; Edlund, H.: Attenua-
tion

[187620]of FGF signalling in mouse beta-cells leads to diabetes.

Nature 408:

[187621]864–868, 2000.

[187622]

[187623]

[187624]19766.

[187625]Mammarella, S.; Romano, F.; Di Valerio, A.; Creati, B.; Es-
posito,

[187626]D. L.; Palmirotta, R.; Capani, F.; Vitullo, P.; Volpe, G.; Bat-
tista,

[187627]P.; Della Loggia, F.; Mariani–Costantini, R.; Cama, A.: In-
teraction

[187628]between the G1057D variant of IRS–2 and overweight in
the pathogenesis

[187629]of type 2 diabetes. Hum. Molec. Genet. 9: 2517–2521,

2000.

[187630]

[187631]

[187632]19767.

[187633]Triggs–Raine, B. L.; Kirkpatrick, R. D.; Kelly, S. L.; Norquay,

[187634]L. D.; Cattini, P. A.; Yamagata, K.; Hanley, A. J. G.; Zinman,
B.;

[187635]Harris, S. B.; Barrett, P. H.; Hegele, R. A.: HNF1–alpha
G319S, a

[187636]transactivation–deficient mutant, is associated with al–
tered dynamics

[187637]of diabetes onset in an Oji–Cree community. Proc. Nat.
Acad. Sci. 99:

[187638]4614–4619, 2002.

[187639]

[187640]

[187641]19768.

[187642]Tuomi, T.; Carlsson, A.; Li, H.; Isomaa, B.; Miettinen, A.;
Nilsson,

[187643]A.; Nissen, M.; Ehrnstrom, B.–O.; Forsen, B.; Snickars, B.;

Lahti,

[187644]K.; Forsblom, C.; Saloranta, C.; Taskinen, M.-R.; Groop, L.

C.: Clinical

[187645]and genetic characteristics of type 2 diabetes with and

without GAD

[187646]antibodies. Diabetes 48: 150–157, 1999.

[187647]

[187648]

[187649]19769.

[187650]Yuan, M.; Konstantopoulos, N.; Lee, J.; Hansen, L.; Li,

Z.-W.;

[187651]Karin, M.; Shoelson, S. E.: Reversal of obesity- and diet-

induced

[187652]insulin resistance with salicylates or targeted disruption of

Ikk-beta. Science 293:

[187653]1673–1677, 2001.

[187654]

[187655]

[187656]19770.

[187657]Hart, T. C.; Champagne, C.; Zhou, J.; Van Dyke, T. E.: As-

signment

[187658]of the gene for diacylglycerol kinase (DAGK) to human
chromosome 12. Mammalian

[187659]Genome 5: 123–124, 1994.

[187660]

[187661]

[187662]19771.

[187663]Hart, T. C.; Zhou, J.; Champagne, C.; Van Dyke, T. E.; Rao,
P.

[187664]N.; Pettenati, M. J.: Assignment of the human diacylglyc-
erol kinase

[187665]gene (DAGK) to 12q13.3 using fluorescence in situ hy-
bridization analysis. Genomics 22:

[187666]246–247, 1994.

[187667]

[187668]

[187669]19772.

[187670]Pilz, A.; Schaap, D.; Hunt, D.; Fitzgibbon, J.: Chromosomal
localization

[187671]of three mouse diacylglycerol kinase (DAGK) genes: genes

sharing sequence

[187672]homology to the Drosophila retinal degeneration A (rdgA) gene. Genomics 26:

[187673]599–601, 1995.

[187674]

[187675]

[187676]19773.

[187677]Schaap, D.; de Widt, J.; van der Wal, J.; Vandekerckhove, J.;
van

[187678]Damme, J.; Gussow, D.; Ploegh, H. L.; van Blitterswijk, W.
J.; van

[187679]der Bend, R. L.: Purification, cDNA-cloning and expression
of human

[187680]diacylglycerol kinase. FEBS Lett. 275: 151–158, 1990.

[187681]

[187682]

[187683]19774.

[187684]Barker, P. E.; Shipp, M. A.; D'Adamio, L.; Masteller, E. L.;
Reinherz,

[187685]E. L.: The common acute lymphoblastic leukemia antigen

gene maps

[187686]to chromosomal region 3(q21-q27). J. Immun. 142:
283-287, 1989.

[187687]

[187688]

[187689]19775.

[187690]D'Adamio, L.; Shipp, M. A.; Masteller, E. L.; Reinherz, E. L.:

[187691]Organization of the gene encoding common acute lym-
phoblastic leukemia

[187692]antigen (neutral endopeptidase 24.11): multiple miniex-
ons and separate

[187693]5-prime untranslated regions. Proc. Nat. Acad. Sci. 86:
7103-7107,

[187694]1989.

[187695]

[187696]

[187697]19776.

[187698]Debiec, H.; Guigonis, V.; Mougnot, B.; Decobert, F.; Hay-
mann,

[187699]J.-P.; Bensman, A.; Deschenes, G.; Ronco, P. M.: Antenatal

membranous

[187700]glomerulonephritis due to anti-neutral endopeptidase antibodies. New

[187701]Eng. J. Med. 346: 2053–2060, 2002.

[187702]

[187703]

[187704]19777.

[187705]Letarte, M.; Vera, S.; Tran, R.; Addis, J. B. L.; Onizuka, R. J.;

[187706]Quackenbush, E. J.; Jongeneel, C. V.; McInnes, R. R.: Common acute

[187707]lymphocytic leukemia antigen is identical to neutral endopeptidase. J.

[187708]Exp. Med. 168: 1247–1253, 1988.

[187709]

[187710]

[187711]19778.

[187712]Shipp, M. A.; Vijayaraghavan, J.; Schmidt, E. V.; Masteller, E.

[187713]L.; D'Adamio, L.; Hersh, L. B.; Reinherz, E. L.: Common acute lymphoblastic

[187714]leukemia antigen (CALLA) is active neutral endopeptidase

24.11 ('enkephalinase'):

[187715]direct evidence by cDNA transfection analysis. Proc. Nat.

Acad. Sci. 86:

[187716]297–301, 1989.

[187717]

[187718]

[187719]19779.

[187720]Tran–Paterson, R.; Willard, H. F.; Letarte, M.: The common
acute

[187721]lymphoblastic leukemia antigen (neutral endopeptidase–
–3.4.24.11)

[187722]gene is located on human chromosome 3. Cancer Genet.
Cytogenet. 42:

[187723]129–134, 1989.

[187724]

[187725]

[187726]19780.

[187727]Lahn, B. T.; Page, D. C.: Four evolutionary strata on the
human

[187728]X chromosome. Science 286: 964–967, 1999.

[187729]

[187730]

[187731]19781.

[187732]Lee, F. A.; Issacs, H.; Strauss, J.: The 'camptomelic' syndrome.

[187733]Short life-span dwarfism with respiratory distress, hypotonia, peculiar

[187734]facies, and multiple skeletal and cartilaginous deformities. Am.

[187735]J. Dis. Child. 124: 485–496, 1972.

[187736]

[187737]

[187738]19782.

[187739]Lynch, S. A.; Gaunt, M. L.; Minford, A. M. B.: Campomelic dysplasia:

[187740]evidence of autosomal dominant inheritance. J. Med. Genet. 30: 683–686,

[187741]1993.

[187742]

[187743]

[187744]19783.

[187745]Macpherson, R. I.; Skinner, S. A.; Donnenfeld, A. E.: Acampomelic

[187746]campomelic dysplasia. *Pediat. Radiol.* 20: 90–93, 1989.

[187747]

[187748]

[187749]19784.

[187750]Mansour, S.; Hall, C. M.; Pembrey, M. E.; Young, I. D.: A clinical

[187751]and genetic study of campomelic dysplasia. *J. Med. Genet.* 32: 415–420,

[187752]1995.

[187753]

[187754]

[187755]19785.

[187756]Maraia, R.; Saal, H. M.; Wangsa, D.: A chromosome 17q de novo

[187757]paracentric inversion in a patient with campomelic dyspla-

sia; case

[187758]report and etiologic hypothesis. Clin. Genet. 39: 401–408,
1991.

[187759]

[187760]

[187761]19786.

[187762]Maroteaux, P.; Spranger, J. W.; Opitz, J. M.; Kucera, J.;
Lowry,

[187763]R. B.; Schimke, R. N.; Kagan, S. M.: Le syndrome cam-
pomelique. Presse

[187764]Med. 22: 1157–1162, 1971.

[187765]

[187766]

[187767]19787.

[187768]Meyer, J.; Sudbeck, P.; Held, M.; Wagner, T.; Schmitz, M.
L.;

[187769]Bricarelli, F. D.; Eggermont, E.; Friedrich, U.; Haas, O. A.;
Kobelt,

[187770]A.; Leroy, J. G.; van Maldergem, L.; Michel, E.; Mitulla, B.;
Pfeiffer,

[187771]R. A.; Schinzel, A.; Schmidt, H.; Scherer, G.: Mutational

analysis

[187772]of the SOX9 gene in campomelic dysplasia and autosomal sex reversal:

[187773]lack of genotype/phenotype correlations. Hum. Molec. Genet. 6: 91–98, [187774]1997.

[187775]

[187776]

[187777]19788.

[187778]Moedjono, S. J.; Crandall, B. F.; Sparkes, R. S.; Feldman, G.

[187779]M.; Austin, G. E.; Perry, S.: The campomelic syndrome in a singleton

[187780]and monozygotic twins. Clin. Genet. 18: 397–401, 1980.

[187781]

[187782]

[187783]19789.

[187784]Moog, U.; Jansen, N. J. G.; Scherer, G.; Schrandt–Stumpel, C.

[187785]T. R. M.: Acampomelic campomelic syndrome. Am. J. Med. Genet. 104:

[187786]239–245, 2001.

[187787]

[187788]

[187789]19790.

[187790]Morais da Silva, S.; Hacker, A.; Harley, V.; Goodfellow, P.;
Swain,

[187791]A.; Lovell–Badge, R.: Sox9 expression during gonadal de-
velopment

[187792]implies a conserved role for the gene in testis differentia-
tion in

[187793]mammals and birds. Nature Genet. 14: 62–68, 1996.

[187794]

[187795]

[187796]19791.

[187797]Murakami, S.; Kan, M.; McKeehan, W. L.; de Crombrugghe,
B.: Up–regulation

[187798]of the chondrogenic Sox9 gene by fibroblast growth fac-
tors is mediated

[187799]by the mitogen–activated protein kinase pathway. Proc.
Nat. Acad.

[187800]Sci. 97: 1113–1118, 2000.

[187801]

[187802]

[187803]19792.

[187804]Ninomiya, S.; Isomura, M.; Narahara, K.; Seino, Y.; Nakamura,

[187805]Y.: Isolation of a testis-specific cDNA on chromosome 17q from a

[187806]region adjacent to the breakpoint of t(12;17) observed in a patient

[187807]with acampomelic campomelic dysplasia and sex reversal. Hum. Molec.

[187808]Genet. 5: 69–72, 1996.

[187809]

[187810]

[187811]19793.

[187812]Ninomiya, S.; Yokoyama, Y.; Teraoka, M.; Mori, R.; Inoue, C.;

[187813]Yamashita, S.; Tamai, H.; Funato, M.; Seino, Y.: A novel mutation

[187814](296 del G) of the SOX9 gene in a patient with campomelic syndrome

[187815]and sex reversal. Clin. Genet. 58: 224–227, 2000.

[187816]

[187817]

[187818]19794.

[187819]Olney, P. N.; Kean, L. S.; Graham, D.; Elsas, L. J.; May, K. M.

[187820]: Campomelic syndrome and deletion of SOX9. Am. J. Med. Genet. 84:

[187821]20–24, 1999.

[187822]

[187823]

[187824]19795.

[187825]Ozkilic, A.; Seven, M.; Yuksel, A.: A case of acampomelic campomelic

[187826]dysplasia. Genet. Counsel. 13: 23–28, 2002.

[187827]

[187828]

[187829]19796.

[187830]Patel, M.; Dorman, K. S.; Zhang, Y.-H.; Huang, B.-L.;
Arnold,

[187831]A. P.; Sinsheimer, J. S.; Vilain, E.; McCabe, E. R. B.: Primate
DAX1,

[187832]SRY, and SOX9: evolutionary stratification of sex-
determination pathway. Am.

[187833]J. Hum. Genet. 68: 275–280, 2001.

[187834]

[187835]

[187836]19797.

[187837]Pfeifer, D.; Kist, R.; Dewar, K.; Devon, K.; Lander, E. S.; Bir-
ren,

[187838]B.; Korniszewski, L.; Back, E.; Scherer, G.: Campomelic
dysplasia

[187839]translocation breakpoints are scattered over 1 Mb proxi-
mal to SOX9:

[187840]evidence for an extended control region. Am. J. Hum.
Genet. 65:

[187841]111–124, 1999.

[187842]

[187843]

[187844]19798.

[187845]Puck, S. M.; Haseltine, F. P.; Francke, U.: Absence of H-Y
antigen

[187846]in an XY female with campomelic dysplasia. Hum. Genet.
57: 23-27,

[187847]1981.

[187848]

[187849]

[187850]19799.

[187851]Rimoin, D. L.: Personal Communication. Torrance, Calif.
8/12/1976.

[187852]

[187853]

[187854]19800.

[187855]Rodriguez, J. I.: Vascular anomalies in campomelic syn-
drome. Am.

[187856]J. Med. Genet. 46: 185-192, 1993.

[187857]

[187858]

[187859]19801.

[187860]Savarirayan, R.; Bankier, A.: Acampomelic campomelic
dysplasia

[187861]with de novo 5q;17q reciprocal translocation and severe
phenotype. J.

[187862]Med. Genet. 35: 597–599, 1998.

[187863]

[187864]

[187865]19802.

[187866]Schimke, R. N.: XY sex-reversed campomelia--possibly an
X-linked

[187867]disorder? (Letter) Clin. Genet. 16: 62–63, 1979.

[187868]

[187869]

[187870]19803.

[187871]Shafai, T.; Schwartz, L.: Camptomelic syndrome in sib-
lings. J.

[187872]Pediat. 89: 512–513, 1976.

[187873]

[187874]

[187875]19804.

[187876]Spranger, J.: Advances in bone dysplasias. (Abstract) Sixth

[187877]Int. Cong. Hum. Genet. Jerusalem , 1981.

[187878]

[187879]

[187880]19805.

[187881]Stuve, A.; Wiedemann, H.-R.: Congenital bowing of the
long bones

[187882]in two sisters. (Letter) Lancet I: 495, 1971.

[187883]

[187884]

[187885]19806.

[187886]Sudbeck, P.; Schmitz, M. L.; Baeuerle, P. A.; Scherer, G.:
Sex

[187887]reversal by loss of the C-terminal transactivation domain
of human

[187888]SOX9. Nature Genet. 13: 230–232, 1996.

[187889]

[187890]

[187891]19807.

[187892]Thong, M.-K.; Scherer, G.; Kozlowski, K.; Haan, E.; Morris, L.

[187893]: Acampomelic campomelic dysplasia with SOX9 mutation. Am. J. Med.

[187894]Genet. 93: 421-425, 2000.

[187895]

[187896]

[187897]19808.

[187898]Thurmon, T. F.; De Fraites, E. B.; Anderson, E. E.: Familial

[187899]campomelic dwarfism. J. Pediat. 83: 841-843, 1973.

[187900]

[187901]

[187902]19809.

[187903]Vidal, V. P. I.; Chaboissier, M.-C.; de Rooij, D. G.; Schedl,

[187904]A.: Sox9 induces testis development in XX transgenic

mice. Nature

[187905]Genet. 28: 216–217, 2001.

[187906]

[187907]

[187908]19810.

[187909]Wagner, T.; Wirth, J.; Meyer, J.; Zabel, B.; Held, M.; Zimmer,
mer,

[187910]J.; Pasantes, J.; Dagna Bricarelli, F.; Keutel, J.; Hustert, E.;
Wolf,

[187911]U.; Tommerup, N.; Schempp, W.; Scherer, G.: Autosomal
sex reversal

[187912]and campomelic dysplasia are caused by mutations in and
around the

[187913]SRY-related gene SOX9. Cell 79: 1111–1120, 1994.

[187914]

[187915]

[187916]19811.

[187917]Weller, S. D. V.: Hypophosphatasia with congenital dim-
ples. Proc.

[187918]Roy. Soc. Med. 52: 637, 1959.

[187919]

[187920]

[187921]19812.

[187922]Wirth, J.; Wagner, T.; Meyer, J.; Pfeiffer, R. A.; Tietze, H.-U.;

[187923]Schempp, W.; Scherer, G.: Translocation breakpoints in three patients

[187924]with campomelic dysplasia and autosomal sex reversal map more than

[187925]130 kb from SOX9. Hum. Genet. 97: 186–193, 1996.

[187926]

[187927]

[187928]19813.

[187929]Wright, E.; Hargrave, M. R.; Christiansen, J.; Cooper, L.; Kun,

[187930]J.; Evans, T.; Gangadharan, U.; Greenfield, A.; Koopman, P.: The

[187931]Sry-related gene Sox9 is expressed during chondrogenesis in mouse

[187932]embryos. Nature Genet. 9: 15–20, 1995.

[187933]

[187934]

[187935]19814.

[187936]Albritton, L. M.; Bowcock, A. M.; Eddy, R. L.; Morton, C. C.;
Tseng,

[187937]L.; Farrer, L. A.; Cavalli-Sforza, L. L.; Shows, T. B.; Cun-
ningham,

[187938]J. M.: The human cationic amino acid transporter (ATRC1):
physical

[187939]and genetic mapping to 13q12-q14. Genomics 12:
430-434, 1992.

[187940]

[187941]

[187942]19815.

[187943]Bowcock, A. M.; Gerken, S. C.; Barnes, R. I.; Shiang, R.;
Jabs,

[187944]E. W.; Warren, A. C.; Antonarakis, S.; Retief, A. E.;
Vergnaud, G.;

[187945]Leppert, M.; Lalouel, J.-M.; White, R. L.; Cavalli-Sforza, L.
L.:

[187946]The CEPH consortium linkage map of human chromosome

13. Genomics 16:

[187947]486–496, 1993.

[187948]

[187949]

[187950]19816.

[187951]Kim, J. W.; Closs, E. I.; Albritton, L. M.; Cunningham, J. M.:

[187952]Transport of cationic amino acids by the mouse ecotropic
retrovirus

[187953]receptor. *Nature* 352: 725–728, 1991.

[187954]

[187955]

[187956]19817.

[187957]Kozak, C. A.; Albritton, L. M.; Cunningham, J. M.: Genetic
mapping

[187958]of a cloned sequence responsible for susceptibility to
ecotropic murine

[187959]leukemia viruses. *J. Virol.* 64: 3119–3121, 1990.

[187960]

[187961]

[187962]19818.

[187963]Oie, H. K.; Gazdar, A. F.; Lalley, P. A.; Russell, E. K.; Minna,

[187964]J. D.; DeLarco, J.; Todaro, G. J.; Francke, U.: Mouse chromosome

[187965]5 codes for ecotropic murine leukaemia virus cell-surface receptor. Nature 274:

[187966]60-62, 1978.

[187967]

[187968]

[187969]19819.

[187970]Ruddle, N. H.; Conta, B. S.; Leinwand, L.; Kozak, C.; Ruddle, F.;

[187971]Besmer, P.; Baltimore, D.: Assignment of the receptor for ecotropic

[187972]murine leukemia virus to mouse chromosome 5. J. Exp. Med. 148: 451-465,

[187973]1978.

[187974]

[187975]

[187976]19820.

[187977]Boldog, F.; Erlandsson, R.; Klein, G.; Sumegi, J.: Long-range

[187978]restriction enzyme maps of DNF15S2, D3S2 and c-raf1 loci on the short

[187979]arm of human chromosome 3. Cancer Genet. Cytogenet. 42: 295–306,

[187980]1989.

[187981]

[187982]

[187983]19821.

[187984]Cook, R. M.; Burke, B. J.; Buchhagen, D. L.; Minna, J. D.; Miller,

[187985]Y. E.: Human aminoacylase-1: cloning, sequence, and expression analysis

[187986]of a chromosome 3p21 gene inactivated in small cell lung cancer. J.

[187987]Biol. Chem. 268: 17010–17017, 1993.

[187988]

[187989]

[187990]19822.

[187991]Gemmill, R. M.; Varella-Garcia, M.; Smith, D. I.; Erickson, P.;

[187992]Golembieski, W.; Miller, Y.; Coyle-Morris, J.; Tommerup, N.; Drabkin,

[187993]H. A.: A 2.5 Mb physical map within 3p21.1 spans the breakpoint associated

[187994]with Greig cephalopolysyndactyly syndrome. Genomics 11: 93-102,

[187995]1991.

[187996]

[187997]

[187998]19823.

[187999]Ginzinger, D. G.; Shridhar, V.; Baldini, A.; Taggart, R. T.; Miller,

[188000]O. J.; Smith, D. I.: The human loci DNF15S2 and D3S94 have a high

[188001]degree of sequence similarity to acyl-peptide hydrolase and are located

[188002]at 3p21.3. Am. J. Hum. Genet. 50: 826-833, 1992.

[188003]

[188004]

[188005]19824.

[188006]Jones, W. M.; Scaloni, A.; Bossa, F.; Popowicz, A. M.;

Schneewind,

[188007]O.; Manning, J. M.: Genetic relationship between acylpep-
tide hydrolase

[188008]and acylase, two hydrolytic enzymes with similar binding
but different

[188009]catalytic specificities. Proc. Nat. Acad. Sci. 88: 2194–2198,
1991.

[188010]

[188011]

[188012]19825.

[188013]Miller, Y. E.; Drabkin, H.; Jones, C.; Fisher, J. H.: Aminoacylase-1:

[188014]cDNA isolation, regional assignment to chromosome
3p21.1 and identification

[188015]of a cross-hybridizing sequence on chromosome 18.
(Abstract) Am.

[188016]J. Hum. Genet. 45 (suppl.): A28 only, 1989.

[188017]

[188018]

[188019]19826.

[188020]Miller, Y. E.; Drabkin, H.; Jones, C.; Fisher, J. H.: Human
aminoacylase-1:

[188021]cloning, regional assignment to distal chromosome
3p21.1, and identification

[188022]of a cross-hybridizing sequence on chromosome 18. Ge-
nomics 8: 149-154,

[188023]1990.

[188024]

[188025]

[188026]19827.

[188027]Miller, Y. E.; Minna, J. D.; Gazdar, A. F.: Lack of expression

[188028]of aminoacylase-1 in small cell lung cancer: evidence for
inactivation

[188029]of genes encoded by chromosome 3p. J. Clin. Invest. 83:
2120-2124,

[188030]1989.

[188031]

[188032]

[188033]19828.

[188034]Nadeau, J. H.: A chromosomal segment conserved since divergence

[188035]of lineages leading to man and mouse: the gene order of aminoacylase-1,

[188036]transferrin, and beta-galactosidase on mouse chromosome 9. Genet.

[188037]Res. 48: 175-178, 1986.

[188038]

[188039]

[188040]19829.

[188041]Naylor, S. L.; Elliott, R. W.; Brown, J. A.; Shows, T. B.: Mapping ping

[188042]of aminoacylase-1 and beta-galactosidase-A to homologous regions of

[188043]human chromosome 3 and mouse chromosome 9 suggests location of additional

[188044]genes. Am. J. Hum. Genet. 34: 235-244, 1982.

[188045]

[188046]

[188047]19830.

[188048]Naylor, S. L.; Shows, T. B.; Klebe, R. J.: Bioautographic visualization

[188049]of aminoacylase-1: assignment of the structural gene ACY-1 to chromosome

[188050]3 in man. Somat. Cell Genet. 5: 11-21, 1979.

[188051]

[188052]

[188053]19831.

[188054]Voss, R.; Lerer, I.; Povey, S.; Solomon, E.; Bobrow, M.: Confirmation

[188055]and further regional assignment of aminoacylase 1 (ACY-1) on human

[188056]chromosome 3 using a simplified detection method. Ann. Hum. Genet. 44:

[188057]1-10, 1980.

[188058]

[188059]

[188060]19832.

[188061]Cook, P. W.; Piepkorn, M.; Clegg, C. H.; Plowman, G. D.; DeMay,

[188062]J. M.; Brown, J. R.; Pittelkow, M. R.: Transgenic expression
of the

[188063]human amphiregulin gene induces a psoriasis-like phe-
notype. J. Clin.

[188064]Invest. 100: 2286–2294, 1997.

[188065]

[188066]

[188067]19833.

[188068]Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly,
J.;

[188069]Bowen–Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of
the amphiregulin

[188070]and the platelet–growth factor receptor alpha genes to the
proximal

[188071]long arm of chromosome 4. (Abstract) Cytogenet. Cell
Genet. 51:

[188072]990 only, 1989.

[188073]

[188074]

[188075]19834.

[188076]Kimura, H.; Fischer, W. H.; Schubert, D.: Structure, expression

[188077]and function of a schwannoma-derived growth factor. Nature 348:

[188078]257–260, 1990.

[188079]

[188080]

[188081]19835.

[188082]Gillesen–Kaesbach, G.; Demuth, S.; Thiele, H.; Theile, U.; Lich,

[188083]C.; Horsthemke, B.: A previously unrecognised phenotype characterised

[188084]by obesity, muscular hypotonia, and ability to speak in patients with

[188085]Angelman syndrome caused by an imprinting defect. Europ. J. Hum.

[188086]Genet. 7: 638–644, 1999.

[188087]

[188088]

[188089]19836.

[188090]Cockayne, D. A.; Muchamuel, T.; Grimaldi, J. C.; Muller-Steffner,

[188091]H.; Randall, T. D.; Lund, F. E.; Murray, R.; Schuber, F.; Howard,

[188092]M. C.: Mice deficient for the ecto-nicotinamide adenine dinucleotide

[188093]glycohydrolase CD38 exhibit altered humoral immune responses. Blood 92:

[188094]1324–1333, 1998.

[188095]

[188096]

[188097]19837.

[188098]Ferrero, E.; Malavasi, F.: Human CD38, a leukocyte receptor and

[188099]ectoenzyme, is a member of a novel eukaryotic gene family of nicotinamide

[188100]adenine dinucleotide +/-converting enzymes: extensive structure homology

[188101]with the genes for murine bone marrow stromal cell antigen 1 and aplysian

[188102]ADP-ribosyl cyclase. J. Immun. 159: 3858–3865, 1997.

[188103]

[188104]

[188105]19838.

[188106]Ferrero, E.; Saccucci, F.; Malavasi, F.: The human CD38 gene:

[188107]polymorphism, CpG island, and linkage to the CD157 (BST-1) gene. Immunogenetics 49:

[188108]597–604, 1999.

[188109]

[188110]

[188111]19839.

[188112]Fukushi, Y.; Kato, I.; Takasawa, S.; Sasaki, T.; Ong, B. H.; Sato,

[188113]M.; Ohsaga, A.; Sato, K.; Shirato, K.; Okamoto, H.; Maruyama, Y.:

[188114]Identification of cyclic ADP-ribose-dependent mechanisms in pancreatic

[188115]muscarinic Ca(2+) signaling using CD38 knockout mice. J. Biol. Chem. 276:

[188116]649–655, 2001.

[188117]

[188118]

[188119]19840.

[188120]Jackson, D. G.; Bell, J. I.: Isolation of a cDNA encoding the

[188121]human CD38 (T10) molecule, a cell surface glycoprotein
with an unusual

[188122]discontinuous pattern of expression during lymphocyte
differentiation. J.

[188123]Immun. 144: 2811–2815, 1990.

[188124]

[188125]

[188126]19841.

[188127]Kaisho, T.; Ishikawa, J.; Oritani, K.; Inazawa, J.; Tomizawa,
H.;

[188128]Muraoka, O.; Ochi, T.; Hirano, T.: BST–1, a surface
molecule of bone

[188129]marrow stromal cell lines that facilitates pre–B–cell
growth. Proc.

[188130]Nat. Acad. Sci. 91: 5325–5329, 1994.

[188131]

[188132]

[188133]19842.

[188134]Nakagawara, K.; Mori, M.; Takasawa, S.; Nata, K.; Takamura, T.;

[188135]Berlova, A.; Tohgo, A.; Karasawa, T.; Yonekura, H.; Takeuchi, T.;

[188136]Okamoto, H.: Assignment of CD38, the gene encoding human leukocyte

[188137]antigen CD38 (ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase), to

[188138]chromosome 4p15. Cytogenet. Cell Genet. 69: 38-39, 1995.

[188139]

[188140]

[188141]19843.

[188142]Takasawa, S.; Nata, K.; Yonekura, H.; Okamoto, H.: Cyclic ADP-ribose

[188143]in insulin secretion from pancreatic beta cells. Science 259: 370-373,

[188144]1993.

[188145]

[188146]

[188147]19844.

[188148]Takasawa, S.; Tohgo, A.; Noguchi, N.; Koguma, T.; Nata, K.; Sugimoto,

[188149]T.; Yonekura, H.; Okamoto, H.: Synthesis and hydrolysis of cyclic

[188150]ADP-ribose by human leukocyte antigen CD38 and inhibition of the hydrolysis

[188151]by ATP. J. Biol. Chem. 268: 26052-26054, 1993.

[188152]

[188153]

[188154]19845.

[188155]Acosta, J.; Hettinga, J.; Fluckiger, R.; Krumrei, N.; Goldfine,

[188156]A.; Angarita, L.; Halperin, J.: Molecular basis for a link between

[188157]complement and the vascular complications of diabetes. Proc. Nat.

[188158]Acad. Sci. 97: 5450-5455, 2000.

[188159]

[188160]

[188161]19846.

[188162]Bickmore, W. A.; Longbottom, D.; Oghene, K.; Fletcher, J.
M.; van

[188163]Heyningen, V.: Colocalization of the human CD59 gene to
11p13 with

[188164]the MIC11 cell surface antigen. Genomics 17: 129–135,
1993.

[188165]

[188166]

[188167]19847.

[188168]Davies, A.; Simmons, D. L.; Hale, G.; Harrison, R. A.; Tighe,
H.;

[188169]Lachmann, P. J.; Waldmann, H.: CD59, an LY-6-like pro-
tein expressed

[188170]in human lymphoid cells, regulates the action of the com-
plement membrane

[188171]attack complex on homologous cells. J. Exp. Med. 170:
637–654, 1989.

[188172]

[188173]

[188174]19848.

[188175]Forsberg, U. H.; Bazil, V.; Stefanova, I.; Schroder, J.: Gene
[188176]for human CD59 (likely Ly-6 homologue) is located on the
short arm
[188177]of chromosome 11. Immunogenetics 30: 188–193, 1989.
[188178]
[188179]
[188180]19849.
[188181]Harada, R.; Okada, N.; Fujita, T.; Okada, H.: Purification of
[188182]1F5 antigen that prevents complement attack on homolo-
gous cell membranes. J.
[188183]Immun. 144: 1823–1828, 1990.
[188184]
[188185]
[188186]19850.
[188187]Heckl–Ostreicher, B.; Ragg, S.; Drechsler, M.; Scherthan,
H.; Royer–Pokora,
[188188]B.: Localization of the human CD59 gene by fluorescence
in situ hybridization
[188189]and pulsed–field gel electrophoresis. Cytogenet. Cell

Genet. 63:

[188190]144–146, 1993.

[188191]

[188192]

[188193]19851.

[188194]Holt, D. S.; Botto, M.; Bygrave, A. E.; Hanna, S. M.; Walport,

[188195]M. J.; Morgan, B. P.: Targeted deletion of the CD59 gene
causes spontaneous

[188196]intravascular hemolysis and hemoglobinuria. Blood 98:
442–449, 2001.

[188197]

[188198]

[188199]19852.

[188200]Holt, D. S.; Powell, M. B.; Rushmere, N. K.; Morgan, B. P.:
Genomic

[188201]structure and chromosome location of the gene encoding
mouse CD59. Cytogenet.

[188202]Cell Genet. 89: 264–267, 2000.

[188203]

[188204]

[188205]19853.

[188206]Huppi, K.; Duncan, R.; Potter, M.: Myc-1 is centromeric to
the

[188207]linkage group Ly-6-Sis-Gdc-1 on mouse chromosome 15.

Immunogenetics 27:

[188208]215-219, 1988.

[188209]

[188210]

[188211]19854.

[188212]Kamiura, S.; Nolan, C. M.; Meruelo, D.: Long-range physi-
cal map

[188213]of the Ly-6 complex: mapping the Ly-6 multigene family
by field-inversion

[188214]and two-dimensional gel electrophoresis. Genomics 12:
89-105, 1992.

[188215]

[188216]

[188217]19855.

[188218]Low, M. G.; Saltiel, A. R.: Structural and functional roles of

[188219]glycosyl-phosphatidylinositol in membranes. Science 239:
268-275,
[188220]1988.

[188221]

[188222]

[188223]19856.

[188224]Mahoney, J. F.; Urakaze, M.; Hall, S.; DeGasperi, R.; Chang,
H.-M.;

[188225]Sugiyama, E.; Warren, C. D.; Borowitz, M.; Nicholson-
Weller, A.; Rosse,

[188226]W. F.; Yeh, E. T. H.: Defective glycosylphosphatidylinositol
anchor

[188227]synthesis in paroxysmal nocturnal hemoglobinuria granu-
locytes. Blood 79:

[188228]1400-1403, 1992.

[188229]

[188230]

[188231]19857.

[188232]Mao, M.; Yu, M.; Tong, J.-H.; Ye, J.; Zhu, J.; Huang, Q.-H.;
Fu,

[188233]G.; Yu, L.; Zhao, S.-Y.; Waxman, S.; Lanotte, M.; Wang,
Z.-Y.; Tan,
[188234]J.-Z.; Chan, S.-J.; Chen, Z.: RIG-E, a human homolog of
the murine
[188235]Ly-6 family, is induced by retinoic acid during the differ-
entiation
[188236]of acute promyelocytic leukemia cell. Proc. Nat. Acad. Sci.
93:
[188237]5910-5914, 1996.

[188238]

[188239]

[188240]19858.

[188241]Meri, S.; Morgan, B. P.; Davies, A.; Daniels, R. H.;
Olavesen,
[188242]M. G.; Waldmann, H.; Lachmann, P. J.: Human protectin
(CD59), an
[188243]18,000-20,000 MW complement lysis restricting factor,
inhibits C5b-8
[188244]catalysed insertion of C9 into lipid bilayers. Immunology
71: 1-9,
[188245]1990.

[188246]

[188247]

[188248]19859.

[188249]Meri, S.; Morgan, B. P.; Wing, M.; Jones, J.; Davies, A.; Po-
dack,

[188250]E.; Lachmann, P. J.: Human protectin (CD59), an 18–20–kD
homologous

[188251]complement restriction factor, does not restrict perforin-
mediated

[188252]lysis. J. Exp. Med. 172: 367–370, 1990.

[188253]

[188254]

[188255]19860.

[188256]Motoyama, N.; Okada, N.; Yamashina, M.; Okada, H.:
Paroxysmal

[188257]nocturnal hemoglobinuria due to hereditary nucleotide
deletion in

[188258]the HRF20 (CD59) gene. Europ. J. Immun. 22: 2669–2673,
1992.

[188259]

[188260]

[188261]19861.

[188262]Okada, N.; Harada, R.; Fujiita, T.; Okada, H.: A novel
membrane

[188263]glycoprotein capable of inhibiting membrane attack by
homologous complement. Int.

[188264]Immun. 1: 205–208, 1989.

[188265]

[188266]

[188267]19862.

[188268]Anderson, L.; Anderson, N. G.: High resolution two–
dimensional

[188269]electrophoresis of human plasma proteins. Proc. Nat.
Acad. Sci. 12:

[188270]5421–5425, 1977.

[188271]

[188272]

[188273]19863.

[188274]Cohen, R. D.; Castellani, L. W.; Qiao, J.–H.; Van Lenten, B.

J.;

[188275]Lusis, A. J.; Reue, K.: Reduced aortic lesions and elevated high

[188276]density lipoprotein levels in transgenic mice overexpressing mouse

[188277]apolipoprotein A-IV. J. Clin. Invest. 99: 1906-1916, 1997.

[188278]

[188279]

[188280]19864.

[188281]Duverger, N.; Tremp, G.; Caillaud, J.-M.; Emmanuel, F.; Castro,

[188282]G.; Fruchart, J.-C.; Steinmetz, A.; Deneffe, P.: Protection against

[188283]atherogenesis in mice mediated by human apolipoprotein A-IV. Science 273:

[188284]966-968, 1996.

[188285]

[188286]

[188287]19865.

[188288]Stubbs, L.; Carver, E.; Ashworth, L.; Lopez-Molina, L.: Lo-

cation

[188289]of the DBP transcription factor gene in human and mouse.

Mammalian

[188290]Genome 7: 65–67, 1996.

[188291]

[188292]

[188293]19866.

[188294]Szpirer, C.; Riviere, M.; Cortese, R.; Nakamura, T.; Islam, M.

[188295]Q.; Levan, G.; Szpirer, J.: Chromosomal localization in man and rat

[188296]of the genes encoding the liver–enriched transcription factors C/EBP,

[188297]DBP, and HNF1/LFB–1 (CEBP, DBP, and transcription factor 1, TCF1,

[188298]respectively) and of the hepatocyte growth factor/scatter factor gene

[188299](HGF). Genomics 13: 293–300, 1992.

[188300]

[188301]

[188302]19867.

[188303]Boorstein, W. R.; Vamvakopoulos, N. C.; Fiddes, J. C.: Human chorionic

[188304]gonadotropin beta-subunit is encoded by at least eight genes arranged

[188305]in tandem and inverted pairs. Nature 300: 419–422, 1982.

[188306]

[188307]

[188308]19868.

[188309]Fiddes, J. C.; Goodman, H. M.: The cDNA for the beta-subunit of

[188310]human chorionic gonadotropin suggests evolution of a gene by readthrough

[188311]into the 3-prime-untranslated region. Nature 286: 684–687, 1980.

[188312]

[188313]

[188314]19869.

[188315]Graham, M. Y.; Otani, T.; Boime, I.; Olson, M. V.; Carle, G. F.;

[188316]Chaplin, D. D.: Cosmid mapping of the human chorionic

gonadotropin

[188317]beta subunit genes by field-inversion gel electrophoresis.

Nucleic

[188318]Acids Res. 15: 4437-4448, 1987.

[188319]

[188320]

[188321]19870.

[188322]Julier, C.; Weil, D.; Couillin, P.; Cote, J. C.; Boue, A.;

Thirion,

[188323]J. P.; Kaplan, J. C.; Junien, C.: Confirmation of the assign-

ment

[188324]of the genes coding for human chorionic gonadotropin

beta subunit

[188325]to chromosome 19. (Abstract) Cytogenet. Cell Genet. 37:

501-502,

[188326]1984.

[188327]

[188328]

[188329]19871.

[188330]Gersten, K. M.; Natsuka, S.; Trinchera, M.; Petryniak, B.;

Kelly,

[188331]R. J.; Hiraiwa, N.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.;

[188332]Lowe, J. B.: Molecular cloning, expression, chromosomal assignment,

[188333]and tissue-specific expression of a murine alpha-(1,3)-fucosyltransferase

[188334]locus corresponding to the human ELAM-1 ligand fucosyl transferase. J.

[188335]Biol. Chem. 270: 25047-25056, 1995.

[188336]

[188337]

[188338]19872.

[188339]Julier, C.; Weil, D.; Couillin, P.; Cote, J. C.; Van Cong, N.;

[188340]Foubert, C.; Boue, A.; Thirion, J. P.; Kaplan, J. C.; Junien, C.:

[188341]The beta chorionic gonadotropin-beta luteinizing gene cluster maps

[188342]to human chromosome 19. Hum. Genet. 67: 174-177, 1984.

[188343]

[188344]

[188345]19873.

[188346]Lunardi-Iskander, Y.; Bryant, J. L.; Zeman, R. A.; Lam, V.
H.;

[188347]Samaniego, F.; Besnier, J. M.; Hermans, P.; Thierry, A. R.;
Gill,

[188348]P.; Gallo, R. C.: Tumorigenesis and metastasis of neoplas-
tic Kaposi's

[188349]sarcoma cell line in immunodeficient mice blocked by a
human pregnancy

[188350]hormone. Nature 375: 64-68, 1995.

[188351]

[188352]

[188353]19874.

[188354]Policastro, P.; Ovitt, C. E.; Hoshina, M.; Fukuoka, H.;
Boothby,

[188355]M. R.; Biome, I.: The beta-subunit of human chorionic go-
nadotropin

[188356]is encoded by multiple genes. J. Biol. Chem. 258:
11492-11499, 1983.

[188357]

[188358]

[188359]19875.

[188360]Policastro, P. F.; Daniels–McQueen, S.; Carle, G.; Boime, I.:

[188361]A map of the hCG–beta–LH–beta gene cluster. J. Biol.

Chem. 261:

[188362]5907–5916, 1986.

[188363]

[188364]

[188365]19876.

[188366]Talmadge, K.; Vamvakopoulos, N. C.; Fiddes, J. C.: Evolu-
tion

[188367]of the genes for the beta subunits of human chorionic go-
nadotropin

[188368]and luteinizing hormone. Nature 307: 37–40, 1984.

[188369]

[188370]

[188371]19877.

[188372]Warburton, D.; Gersen, S.; Yu, M.–T.; Jackson, C.; Han-
delin, B.;

[188373]Housman, D.: Monochromosomal rodent-human hybrids
from microcell

[188374]fusion of human lymphoblastoid cells containing an in-
serted dominant

[188375]selectable marker. Genomics 6: 358-366, 1990.

[188376]

[188377]

[188378]19878.

[188379]Reseland, J. E.; Larsen, F.; Solheim, J.; Eriksen, J. A.;
Hanssen,

[188380]L. E.; Prydz, H.: A novel human chymotrypsin-like diges-
tive enzyme. J.

[188381]Biol. Chem. 272: 8099-8104, 1997.

[188382]

[188383]

[188384]19879.

[188385]Honey, N. K.; Sakaguchi, A. Y.; Quinto, C.; MacDonald, R.
J.; Bell,

[188386]G. I.; Craik, C.; Rutter, W. J.; Naylor, S. L.: Chromosomal
assignment

[188387]of human genes for serine proteases trypsin, chymotrypsin B and elastase. Somat.

[188388]Cell Molec. Genet. 10: 369–376, 1984.

[188389]

[188390]

[188391]19880.

[188392]Natt, E.; Magenis, R. E.; Zimmer, J.; Mansouri, A.; Scherer, G.

[188393]: Regional assignment of the loci for uvomorulin (UVO) and chymotrypsinogen

[188394]B (CTRB) with the help of two overlapping deletions on the long arm

[188395]of chromosome 16. Cytogenet. Cell Genet. 50: 145–148, 1989.

[188396]

[188397]

[188398]19881.

[188399]Sakaguchi, A. Y.; Naylor, S. L.; Quinto, C.; Rutter, W. J.; Shows,

[188400]T. B.: The chymotrypsinogen B gene (CTRB) is on human chromosome

[188401]16. Cytogenet. Cell Genet. 32: 313 only, 1982.

[188402]

[188403]

[188404]19882.

[188405]Tomita, N.; Izumoto, Y.; Horii, A.; Doi, S.; Yokouchi, H.;
Ogawa,

[188406]M.; Mori, T.; Matsubara, K.: Molecular cloning and nu-
cleotide sequence

[188407]of human pancreatic prechymotrypsinogen cDNA.

Biochem. Biophys. Res.

[188408]Commun. 158: 569–575, 1989.

[188409]

[188410]

[188411]19883.

[188412]Westphal, E.–M.; Burmeister, M.; Wienker, T. F.; Lehrach,
H.; Bender,

[188413]K.; Scherer, G.: Tyrosine aminotransferase and chy-
motrypsinogen B

[188414]are linked to haptoglobin on human chromosome 16q:
comparison of genetic

[188415]and physical distances. Genomics 1: 313–319, 1987.

[188416]

[188417]

[188418]19884.

[188419]Angeletti, R. H.: Chromogranins and neuroendocrine secretion.

[188420](Editorial) Lab. Invest. 55: 387–390, 1986.

[188421]

[188422]

[188423]19885.

[188424]Bhargava, G.; Russell, J.; Sherwood, L. M.: Phosphorylation of

[188425]parathyroid secretory protein. Proc. Nat. Acad. Sci. 80: 878–881,

[188426]1983.

[188427]

[188428]

[188429]19886.

[188430]Cetin, Y.; Aunis, D.; Bader, M.–F.; Galindo, E.; Jorns, A.;

Bargsten,

[188431]G.; Grube, D.: Chromostatin, a chromogranin A-derived
bioactive peptide,

[188432]is present in human pancreatic insulin (beta) cells. Proc.
Nat. Acad.

[188433]Sci. 90: 2360–2364, 1993.

[188434]

[188435]

[188436]19887.

[188437]Lu, Q.; Lemke, G.: Homeostatic regulation of the immune
system

[188438]by receptor tyrosine kinases of the Tyro 3 family. Science
293:

[188439]306–311, 2001.

[188440]

[188441]

[188442]19888.

[188443]Abrahamson, M.: Human cysteine proteinase inhibitors:
isolation,

[188444]physiological importance, inhibitory mechanism, gene

structure and

[188445]relation to hereditary cerebral hemorrhage. Scand. J. Clin.
Lab.

[188446]Invest. 48 (suppl. 191): 21–31, 1988.

[188447]

[188448]

[188449]19889.

[188450]Abrahamson, M.; Grubb, A.: Increased body temperature
accelerates

[188451]aggregation of the leu68-to-gln mutant cystatin C, the
amyloid-forming

[188452]protein in hereditary cystatin C amyloid angiopathy. Proc.
Nat. Acad.

[188453]Sci. 91: 1416–1420, 1994.

[188454]

[188455]

[188456]19890.

[188457]Abrahamson, M.; Grubb, A.; Olafsson, I.; Lundwall, A.:
Molecular

[188458]cloning and sequence analysis of cDNA coding for the

precursor of

[188459]the human cysteine proteinase inhibitor cystatin C. FEBS
Lett. 216:

[188460]229–233, 1987.

[188461]

[188462]

[188463]19891.

[188464]Abrahamson, M.; Islam, M. Q.; Szpirer, J.; Szpirer, C.;
Levan,

[188465]G.: The human cystatin C gene (CST3), mutated in heredi-
tary cystatin

[188466]C amyloid angiopathy, is located on chromosome 20.

Hum. Genet. 82:

[188467]223–226, 1989.

[188468]

[188469]

[188470]19892.

[188471]Abrahamson, M.; Jonsdottir, S.; Olafsson, I.; Jensson, O.;
Grubb,

[188472]A.: Hereditary cystatin C amyloid angiopathy: identifica-

tion of the

[188473]disease-causing mutation and specific diagnosis by polymerase chain

[188474]reaction based analysis. Hum. Genet. 89: 377–380, 1992.

[188475]

[188476]

[188477]19893.

[188478]Cohen, D. H.; Feiner, H.; Jensson, O.; Frangione, B.: Amyloid

[188479]fibril in hereditary cerebral hemorrhage with amyloidosis (HCHWA)

[188480]is related to the gastroentero–pancreatic neuroendocrine protein,

[188481]gamma trace. J. Exp. Med. 158: 623–628, 1983.

[188482]

[188483]

[188484]19894.

[188485]Ghiso, J.; Jensson, O.; Frangione, B.: Amyloid fibrils in hereditary

[188486]cerebral hemorrhage with amyloidosis of Icelandic type is

a variant

[188487]of gamma-trace basic protein (cystatin C). Proc. Nat.

Acad. Sci. 83:

[188488]2974–2978, 1986.

[188489]

[188490]

[188491]19895.

[188492]Ghiso, J.; Pons-Estel, B.; Frangione, B.: Hereditary cerebral

[188493]amyloid angiopathy: the amyloid fibrils contain a protein
which is

[188494]a variant of cystatin C, an inhibitor of lysosomal cysteine
proteases. Biochem.

[188495]Biophys. Res. Commun. 136: 548–554, 1986.

[188496]

[188497]

[188498]19896.

[188499]Grubb, A.; Jensson, O.; Gudmundsson, G.; Arnason, A.;
Lofberg,

[188500]H.; Malm, J.: Abnormal metabolism of gamma-trace alka-
line microprotein:

[188501]the basic defect in hereditary cerebral hemorrhage with amyloidosis. New

[188502]Eng. J. Med. 311: 1547–1549, 1984.

[188503]

[188504]

[188505]19897.

[188506]Galbraith, G. M. P.; Pandey, J. P.: Tumor necrosis factor alpha

[188507](TNF-alpha) gene polymorphism in alopecia areata. Hum. Genet. 96:

[188508]433–436, 1995.

[188509]

[188510]

[188511]19898.

[188512]Tazi-Ahnini, R.; di Giovine, F. S.; McDonagh, A. J. G.; Messenger,

[188513]A. G.; Amadou, C.; Cox, A.; Duff, G. W.; Cork, M. J.: Structure and

[188514]polymorphism of the human gene for the interferon-induced p78 protein

[188515](MX1): evidence of association with alopecia areata in the
Down syndrome

[188516]region. Hum. Genet. 106: 639–645, 2000.

[188517]

[188518]

[188519]19899.

[188520]Webb, D. J.; Wen, J.; Lysiak, J. J.; Umans, L.; Van Leuven, F.;

[188521]Gonias, S. L.: Murine alpha-macroglobulins demonstrate
divergent

[188522]activities as neutralizers of transforming growth factor-
beta and

[188523]as inducers of nitric oxide synthesis: a possible mecha-
nism for the

[188524]endotoxin insensitivity of the alpha-2-macroglobulin
gene knock-out

[188525]mouse. J. Biol. Chem. 271: 24982–24988, 1996.

[188526]

[188527]

[188528]19900.

[188529]Wilson, A. G.; di Giovine, F. S.; Blakemore, A. I. F.; Duff, G.

[188530]W.: Single base polymorphism in the human tumour
necrosis factor

[188531]alpha (TNF-alpha) gene detectable by NcoI restriction of
PCR product. Hum.

[188532]Molec. Genet. 1: 353 only, 1992.

[188533]

[188534]

[188535]19901.

[188536]Belanger, L.; Roy, S.; Allard, D.: New albumin gene
3-prime adjacent

[188537]to the alpha-1-fetoprotein locus. J. Biol. Chem. 269:
5481-5484,

[188538]1994.

[188539]

[188540]

[188541]19902.

[188542]Lichenstein, H. S.; Lyons, D. E.; Wurfel, M. M.; Johnson, D.
A.;

[188543]McGinley, M. D.; Leidli, J. C.; Trollinger, D. B.; Mayer, J. P.;
Wright,

[188544]S. D.; Zukowski, M. M.: Afamin is a new member of the al-
bumin, alpha-fetoprotein,

[188545]and vitamin D-binding protein gene family. J. Biol. Chem.
269: 18149-18154,

[188546]1994.

[188547]

[188548]

[188549]19903.

[188550]Nishio, H.; Dugaiczyk, A.: Complete structure of the human alpha-albumin

[188551]gene, a new member of the serum albumin multigene family. Proc. Nat.

[188552]Acad. Sci. 93: 7557–7561, 1996.

[188553]

[188554]

[188555]19904.

[188556]Song, Y.–H.; Naumova, A. K.; Liebhaber, S. A.; Cooke, N. E.: Physical

[188557]and meiotic mapping of the region of human chromosome 4q11–q13 encompassing

[188558]the vitamin D binding protein DBP/Gc-globulin and albumin multigene

[188559]cluster. Genome Res. 9: 581–587, 1999.

[188560]

[188561]

[188562]19905.

[188563]Beattie, W. G.; Dugaiczyk, A.: Structure and evolution of human

[188564]alpha-fetoprotein deduced from partial sequence of cloned cDNA. Gene 20:

[188565]415–422, 1982.

[188566]

[188567]

[188568]19906.

[188569]Belayew, A.; Tilghman, S. M.: Genetic analysis of alpha-fetoprotein

[188570]synthesis in mice. Molec. Cell. Biol. 2: 1427–1435, 1982.

[188571]

[188572]

[188573]19907.

[188574]D'Eustachio, P.; Ingram, R. S.; Tilghman, S. M.; Ruddle, F. H.

[188575]: Murine alpha-fetoprotein and albumin: two evolutionarily linked

[188576]proteins encoded on the same mouse chromosome. Somat. Cell Genet. 7:

[188577]289–294, 1981.

[188578]

[188579]

[188580]19908.

[188581]Eiferman, F. A.; Young, P. R.; Scott, R. W.; Tilghman, S. M.:

[188582]Intragenic amplification and divergence in the mouse al-
pha-fetoprotein

[188583]gene. *Nature* 294: 713–718, 1981.

[188584]

[188585]

[188586]19909.

[188587]Faucett, W. A.; Greenberg, F.; Rose, E.; Alpert, E.; Ban-
calari,

[188588]L.; Kardon, N. B.; Mizjewski, G.; Knight, G.; Haddow, J. E.:
Congenital

[188589]deficiency of alpha-fetoprotein. (Abstract) *Am. J. Hum.*
Genet. 45

[188590](suppl.): A259, 1989.

[188591]

[188592]

[188593]19910.

[188594]Ferguson-Smith, M. A.; May, H. M.; Aitken, D. A.; O'Hare,
E.; Yates,

[188595]J. R. W.; Gallagher, J.; Krumlauf, R.; Tilghman, S. M.:
Hereditary

[188596]persistence of alphafetoprotein (HPAFP); linkage studies
with chromosome

[188597]4 markers. (Abstract) Cytogenet. Cell Genet. 37: 469,
1984.

[188598]

[188599]

[188600]19911.

[188601]Ferguson-Smith, M. A.; Yates, J. R. W.; Kelly, D.; Aitken, D.
A.;

[188602]May, H. M.; Krumlauf, R.; Tilghman, S. M.: Hereditary per-
sistence

[188603]of alphafetoprotein maps to the long arm of chromosome
4. (Abstract) Cytogenet.

[188604]Cell Genet. 40: 628, 1985.

[188605]

[188606]

[188607]19912.

[188608]Gabant, P.; Forrester, L.; Nichols, J.; Van Reeth, T.; De
Mees,

[188609]C.; Pajack, B.; Watt, A.; Smitz, J.; Alexandre, H.; Szpirer, C.;
Szpirer,

[188610]J.: Alpha-fetoprotein, the major fetal serum protein, is not
essential

[188611]for embryonic development but is required for female fer-
tility. Proc.

[188612]Nat. Acad. Sci. 99: 12865–12870, 2002.

[188613]

[188614]

[188615]19913.

[188616]Gibbs, P. E. M.; Zielinski, R.; Boyd, C.; Dugaiczky, A.:
Structure,

[188617]polymorphism, and novel repeated DNA elements revealed
by a complete

[188618]sequence of the human alpha-fetoprotein gene. Biochem-
istry 26: 1332–1343,

[188619]1987.

[188620]

[188621]

[188622]19914.

[188623]Seri, M.; Celli, I.; Betsos, N.; Claudiani, F.; Camera, G.;
Romeo,

[188624]G.: A cys634gly substitution of the RET proto-oncogene in
a family

[188625]with recurrence of multiple endocrine neoplasia type 2A
and cutaneous

[188626]lichen amyloidosis. Clin. Genet. 51: 86–90, 1997.

[188627]

[188628]

[188629]19915.

[188630]Kamal, A.; Almenar-Queralt, A.; LeBlanc, J. F.; Roberts, E.
A.;

[188631]Goldstein, L. S. B.: Kinesin-mediated axonal transport of a
membrane

[188632]compartment containing beta-secretase and presenilin-1
requires APP. Nature 414:

[188633]643–648, 2001.

[188634]

[188635]

[188636]19916.

[188637]Kang, D. E.; Soriano, S.; Xia, X.; Eberhart, C. G.; De
Strooper,

[188638]B.; Zheng, H.; Koo, E. H.: Presenilin couples the paired
phosphorylation

[188639]of beta-catenin independent of Axin: implications for
beta-catenin

[188640]activation in tumorigenesis. Cell 110: 751–762, 2002.

[188641]

[188642]

[188643]19917.

[188644]Hofstra, R. M. W.; Sijmons, R. H.; Stelwagen, T.; Stulp, R.
P.;

[188645]Kousseff, B. G.; Lips, C. J. M.; Steijlen, P. M.; Van Voorst
Vader,

[188646]P. C.; Buys, C. H. C. M.: RET mutation screening in familial
cutaneous

[188647]lichen amyloidosis and in skin amyloidosis associated with
multiple

[188648]endocrine neoplasia. J. Invest. Derm. 107: 215–218, 1996.

[188649]

[188650]

[188651]19918.

[188652]Bordet, T.; Lesbordes, J.–C.; Rouhani, S.; Castelnau–
Ptakhine,

[188653]L.; Schmalbruch, H.; Haase, G.; Kahn, A.: Protective effects
of cardiotrophin–1

[188654]adenoviral gene transfer on neuromuscular degeneration
in transgenic

[188655]ALS mice. Hum. Molec. Genet. 10: 1925–1933, 2001.

[188656]

[188657]

[188658]19919.

[188659]Fantes, J. A.; Bickmore, W. A.; Fletcher, J. M.; Ballesta, F.;

[188660]Hanson, I. M.; van Heyningen, V.: Submicroscopic dele-
tions at the

[188661]WAGR locus, revealed by nonradioactive in situ hybridiza-
tion. Am.

[188662]J. Hum. Genet. 51: 1286–1294, 1992.

[188663]

[188664]

[188665]19920.

[188666]Glaser, T.; Jepeal, L.; Edwards, J. G.; Young, S. R.; Favor, J.;

[188667]Maas, R. L.: PAX6 gene dosage effect in a family with congenital

[188668]cataracts, aniridia, anophthalmia and central nervous system defects. Nature

[188669]Genet. 7: 463–471, 1994.

[188670]

[188671]

[188672]19921.

[188673]Glaser, T.; Lane, J.; Housman, D.: A mouse model of the aniridia–Wilms

[188674]tumor deletion syndrome. Science 250: 823–827, 1990.

[188675]

[188676]

[188677]19922.

[188678]Gronskov, K.; Olsen, J. H.; Sand, A.; Pedersen, W.; Carlsen,

N.;

[188679]Jylling, A. M. B.; Lyngbye, T.; Brondum-Nielsen, K.; Rosenberg, T.

[188680]: Population-based risk estimates of Wilms tumor in sporadic aniridia:

[188681]a comprehensive mutation screening procedure of PAX6 identifies 80%

[188682]of mutations in aniridia. Hum. Genet. 109: 11–18, 2001.

[188683]

[188684]

[188685]19923.

[188686]Hanson, I. M.; Seawright, A.; Hardman, K.; Hodgson, S.; Zaletayev,

[188687]D.; Fekete, G.; van Heyningen, V.: PAX6 mutations in aniridia. Hum.

[188688]Molec. Genet. 2: 915–920, 1993.

[188689]

[188690]

[188691]19924.

[188692]Hill, R. E.; Favor, J.; Hogan, B. L. M.; Ton, C. C. T.; Saun-

ders,

[188693]G. F.; Hanson, I. M.; Prosser, J.; Jordan, T.; Hastie, N. D.;

van

[188694]Heyningen, V.: Mouse small eye results from mutations in

a paired-like

[188695]homeobox-containing gene. Nature 354: 522–525, 1991.

[188696]

[188697]

[188698]19925.

[188699]Jordan, T.; Hanson, I.; Zaletayev, D.; Hodgson, S.; Prosser,

J.;

[188700]Seawright, A.; Hastie, N.; van Heyningen, V.: The human

PAX6 gene

[188701]is mutated in two patients with aniridia. Nature Genet. 1:

328–332,

[188702]1992.

[188703]

[188704]

[188705]19926.

[188706]Karpen, G. H.: Position effect variegation and the new bi–

ology

[188707]of heterochromatin. Curr. Opin. Genet. Dev. 4: 281–291,
1994.

[188708]

[188709]

[188710]19927.

[188711]Lyon, M. F.: Personal Communication. Harwell, England
6/9/1988.

[188712]

[188713]

[188714]19928.

[188715]Martha, A.; Strong, L. C.; Ferrell, R. E.; Saunders, G. F.:
Three

[188716]novel aniridia mutations in the human PAX6 gene. Hum.
Mutat. 6:

[188717]44–49, 1995.

[188718]

[188719]

[188720]19929.

[188721]Matsuo, T.; Osumi-Yamashita, N.; Noji, S.; Ohuchi, H.;

Koyama,

[188722]E.; Myokai, F.; Matsuo, N.; Taniguchi, S.; Doi, H.; Iseki, S.;

Ninomiya,

[188723]Y.; Fujiwara, M.; Watanabe, T.; Eto, K.: A mutation in the

Pax-6

[188724]gene in rat small eye is associated with impaired migra-

tion of midbrain

[188725]crest cells. Nature Genet. 3: 299-304, 1993.

[188726]

[188727]

[188728]19930.

[188729]Oliver, M. D.; Dotan, S. A.; Chemke, J.; Abraham, F. A.:

Isolated

[188730]foveal hypoplasia. Brit. J. Ophthal. 71: 926-930, 1987.

[188731]

[188732]

[188733]19931.

[188734]Prosser, J.; van Heyningen, V.: PAX6 mutations reviewed.

Hum.

[188735]Mutat. 11: 93–108, 1998.

[188736]

[188737]

[188738]19932.

[188739]Quiring, R.; Walldorf, U.; Kloter, U.; Gehring, W. J.: Homology
ogy

[188740]of the eyeless gene of Drosophila to the small eye gene in
mice and

[188741]aniridia in humans. Science 265: 785–789, 1994.

[188742]

[188743]

[188744]19933.

[188745]Salvini-Plawen, L.; Mayr, E.: On the evolution of photore-
ceptors

[188746]and eyes.In: Hecht, M. K.; Steere, W.; Wallace, B.: Evolu-
tionary

[188747]Biology. New York: Plenum Pub. (pub.) 10: 1977. Pp.
207–263.

[188748]

[188749]

[188750]19934.

[188751]Schedl, A.; Ross, A.; Lee, M.; Engelkamp, D.; Rashbass, P.;
van

[188752]Heyningen, V.; Hastie, N. D.: Influence of PAX6 gene
dosage on development:

[188753]overexpression causes severe eye abnormalities. Cell 86:
71–82,

[188754]1996.

[188755]

[188756]

[188757]19935.

[188758]Stone, D. L.; Kenyon, K. R.; Green, W. R.; Ryan, S. J.: Con-
genital

[188759]central corneal leukoma (Peters' anomaly). Am. J. Ophthal.
81: 173–193,

[188760]1976.

[188761]

[188762]

[188763]19936.

[188764]van der Meer–de Jong, R.; Dickinson, M. E.; Woychik, R. P.;

Stubbs,

[188765]L.; Hetherington, C.; Hogan, B. L. M.: Location of the gene involving

[188766]the small eye mutation on mouse chromosome 2 suggests homology with

[188767]human aniridia 2 (AN2). Genomics 7: 270–275, 1990.

[188768]

[188769]

[188770]19937.

[188771]Newport, M. J.; Huxley, C. M.; Huston, S.; Hawrylowicz, C. M.;

[188772]Oostra, B. A.; Williamson, R.; Levin, M.: A mutation in the interferon–gamma–receptor

[188773]gene and susceptibility to mycobacterial infection. New Eng. J. Med. 335:

[188774]1941–1949, 1996.

[188775]

[188776]

[188777]19938.

[188778]Cowan, C. A.; Yokoyama, N.; Bianchi, L. M.; Henkemeyer,

M.; Fritzschn,

[188779]B.: EphB2 guides axons at the midline and is necessary for normal

[188780]vestibular function. Neuron 26: 417–430, 2000.

[188781]

[188782]

[188783]19939.

[188784]Ton, C. C. T.; Hirvonen, H.; Miwa, H.; Weil, M. M.; Monaghan,

[188785]P.; Jordan, T.; van Heyningen, V.; Hastie, N. D.; Meijers-Heijboer,

[188786]H.; Drechsler, M.; Royer-Pokora, B.; Collins, F.; Swaroop, A.; Strong,

[188787]L. C.; Saunders, G. F.: Positional cloning and characterization of

[188788]a paired box- and homeobox-containing gene from the aniridia region. Cell 67:

[188789]1059–1074, 1991.

[188790]

[188791]

[188792]19940.

[188793]Novick, D.; Orchansky, P.; Revel, M.; Rubinstein, M.: The human

[188794]interferon-gamma receptor: purification, characterization, and preparation

[188795]of antibodies. J. Biol. Chem. 262: 8483-8487, 1987.

[188796]

[188797]

[188798]19941.

[188799]Orchansky, P.; Rubinstein, M.; Fischer, D. G.: The interferon-gamma

[188800]receptor in human monocytes is different from the one in nonhematopoietic

[188801]cells. J. Immun. 136: 169-173, 1986.

[188802]

[188803]

[188804]19942.

[188805]Papanicolaou, G. J.; Parsa, N. Z.; Meltzer, P. S.; Trent, J. M.

[188806]: Assignment of interferon gamma receptor (IFNGR1) to human chromosome

[188807]bands 6q24.1-q24.2 by in situ hybridization. Cytogenet.

Cell Genet. 76:

[188808]181–182, 1997. Note: Erratum: Cytogenet. Cell Genet. 78:
132 only,

[188809]1997.

[188810]

[188811]

[188812]19943.

[188813]Pierre–Audigier, C.; Jouanguy, E.; Lamhamedi, S.; Altare,
F.;

[188814]Rauzier, J.; Vincent, V.; Canioni, D.; Emile, J. F.; Fischer, A.;

[188815]Blanche, S.; Gaillard, J. L.; Casanova, J. L.: Fatal dissemi-
nated

[188816]Mycobacterium smegmatis infection in a child with inher-
ited interferon

[188817]gamma receptor deficiency. Clin. Infect. Dis. 24: 982–984,
1997.

[188818]

[188819]

[188820]19944.

[188821]Rashidbaigi, A.; Langer, J. A.; Jung, V.; Jones, C.; Morse, H.

[188822]G.; Tischfield, J. A.; Trill, J. J.; Kung, H.-F.; Pestka, S.: The

[188823]gene for the human immune interferon receptor is located
on chromosome

[188824]6. Proc. Nat. Acad. Sci. 83: 384–388, 1986.

[188825]

[188826]

[188827]19945.

[188828]Rettig, W. J.; Grzeschik, K.-H.; Yenamandra, A. K.; Garcia,
E.;

[188829]Old, L. J.: Definition of selectable cell surface markers for
human

[188830]chromosomes and chromosome segments in rodent–hu-
man hybrids. Somat.

[188831]Cell Molec. Genet. 14: 223–231, 1988.

[188832]

[188833]

[188834]19946.

[188835]Shankaran, V.; Ikeda, H.; Bruce, A. T.; White, J. M.; Swan-
son,

[188836]P. E.; Old, L. J.; Schreiber, R. D.: IFN–gamma and lympho–

cytes prevent

[188837]primary tumour development and shape tumour immuno-
genicity. Nature 410:

[188838]1107–1111, 2001.

[188839]

[188840]

[188841]19947.

[188842]Bauer, R.; Imhof, A.; Pscherer, A.; Kopp, H.; Moser, M.;
Seegers,

[188843]S.; Kerscher, M.; Tainsky, M. A.; Hofstaedter, F.; Buettner,
R.:

[188844]The genomic structure of the human AP-2 transcription
factor. Nucleic

[188845]Acids Res. 22: 1413–1420, 1994.

[188846]

[188847]

[188848]19948.

[188849]Buettner, R.; Kannan, P.; Imhof, A.; Bauer, R.; Yim, S. O.;
Glockshuber,

[188850]R.; Van Dyke, M. W.; Tainsky, M. A.: An alternatively

spliced mRNA

[188851]from the AP-2 gene encodes a negative regulator of transcriptional

[188852]activation by AP-2. Molec. Cell. Biol. 13: 4174-4185, 1993.

[188853]

[188854]

[188855]19949.

[188856]Ahmad, I.: Mash-1 is expressed during ROD photoreceptor differentiation

[188857]and binds an E-box, E(opsin-1), in the rat opsin gene. Brain Res.

[188858]Dev. Brain Res. 90: 184-189, 1995.

[188859]

[188860]

[188861]19950.

[188862]Ball, D. W.; Azzoli, C. G.; Baylin, S. B.; Chi, D.; Dou, S.; Donis-Keller,

[188863]H.; Cumaraswamy, A.; Borges, M.; Nelkin, B. D.: Identification of

[188864]a human achaete-scute homolog highly expressed in neu-

roendocrine tumors. Proc.

[188865]Nat. Acad. Sci. 90: 5648–5652, 1993.

[188866]

[188867]

[188868]19951.

[188869]Guillemot, F.; Lo, L.–C.; Johnson, J. E.; Auerbach, A.; Anderson,

[188870]D. J.; Joyner, A. L.: Mammalian achaete–scute homolog 1 is required

[188871]for the early development of olfactory and autonomic neurons. Cell 75:

[188872]463–476, 1993.

[188873]

[188874]

[188875]19952.

[188876]Renault, B.; Lieman, J.; Ward, D.; Krauter, K.; Kucherlapati, R.

[188877]: Localization of the human achaete–scute homolog gene (ASCL1) distal

[188878]to phenylalanine hydroxylase (PAH) and proximal to tumor

rejection

[188879]antigen (TRA1) on chromosome 12q22–q23. Genomics
30: 81–83, 1995.

[188880]

[188881]

[188882]19953.

[188883]Yang, B.; Ma, T.; Verkman, A. S.: Erythrocyte water perme-
ability

[188884]and renal function in double knockout mice lacking aqua-
porin–1 and

[188885]aquaporin–3. J. Biol. Chem. 276: 624–628, 2001.

[188886]

[188887]

[188888]19954.

[188889]Bunz, F.; Kobayashi, R.; Stillman, B.: cDNAs encoding the
large

[188890]subunit of human replication factor C. Proc. Nat. Acad.
Sci. 90:

[188891]11014–11018, 1993.

[188892]

[188893]

[188894]1995.

[188895]Lossie, A. C.; Haugen, B. R.; Wood, W. M.; Camper, S. A.;
Gordon,

[188896]D. F.: Chromosomal localization of the large subunit of
mouse replication

[188897]factor C in the mouse and human. Mammalian Genome 6:
58–59, 1995.

[188898]

[188899]

[188900]19956.

[188901]Luckow, B.; Bunz, F.; Stillman, B.; Lichter, P.; Schutz, G.:
Cloning,

[188902]expression, and chromosomal localization of the
140–kilodalton subunit

[188903]of replication factor C from mice and humans. Molec. Cell.
Biol. 14:

[188904]1626–1634, 1994.

[188905]

[188906]

[188907]19957.

[188908]Pennaneach, V.; Salles-Passador, I.; Munshi, A.; Brickner, H.;

[188909]Regazzoni, K.; Dick, F.; Dyson, N.; Chen, T.-T.; Wang, J. Y. J.; Fotedar,

[188910]R.; Fotedar, A.: The large subunit of replication factor C promotes

[188911]cell survival after DNA damage in an LxCxE motif- and Rb-dependent

[188912]manner. Molec. Cell 7: 715-727, 2001.

[188913]

[188914]

[188915]19958.

[188916]Uchiumi, F.; Ohta, T.; Tanuma, S.: Replication factor C recognizes

[188917]5-prime-phosphate ends of telomeres. Biochem. Biophys. Res. Commun. 229:

[188918]310-315, 1996.

[188919]

[188920]

[188921]19959.

[188922]Schmickel, R. D.: Contiguous gene syndromes: a component of recognizable

[188923]syndromes. J. Pediat. 109: 231–241, 1986.

[188924]

[188925]

[188926]19960.

[188927]Olson, E.; Srivastava, D.: Molecular pathways controlling heart

[188928]development. Science 272: 671–676, 1996.

[188929]

[188930]

[188931]19961.

[188932]Lamballe, F.; Klein, R.; Barbacid, M.: TRKC, a new member of the

[188933]TRK family of tyrosine protein kinases, is a receptor for neurotrophin-3. Cell 66:

[188934]967–979, 1991.

[188935]

[188936]

[188937]19962.

[188938]McGregor, L. M.; Baylin, S. B.; Griffin, C. A.; Hawkins, A. L.;

[188939]Nelkin, B. D.: Molecular cloning of the cDNA for human
TrkC (NTRK3),

[188940]chromosomal assignment, and evidence for a splice variant. Genomics 22:

[188941]267–272, 1994.

[188942]

[188943]

[188944]19963.

[188945]Segal, R. A.; Goumnerova, L. C.; Kwon, Y. K.; Stiles, C. D.;
Pomeroy,

[188946]S. L.: Expression of the neurotrophin receptor TrkC is
linked to

[188947]a favorable outcome in medulloblastoma. Proc. Nat. Acad.
Sci. 91:

[188948]12867–12871, 1994.

[188949]

[188950]

[188951]19964.

[188952]Valent, A.; Danglot, G.; Bernheim, A.: Mapping of the tyrosine

[188953]kinase receptors trkA (NTRK1), trkB (NTRK2) and trkC (NTRK3) to human

[188954]chromosomes 1q22, 9q22 and 15q25 by fluorescence in situ hybridization. Europ.

[188955]J. Hum. Genet. 5: 102–104, 1997.

[188956]

[188957]

[188958]19965.

[188959]Kielkopf, C. L.; Rodionova, N. A.; Green, M. R.; Burley, S. K.

[188960]: A novel peptide recognition mode revealed by the X-ray structure

[188961]of a core U2AF35/U2AF65 heterodimer. Cell 106: 595–605, 2001.

[188962]

[188963]

[188964]19966.

[188965]Lalioti, M. D.; Gos, A.; Green, M. R.; Rossier, C.; Morris, M.

[188966]A.; Antonarakis, S. E.: The gene for human U2 snRNP auxiliary factor

[188967]small 35-kDa subunit (U2AF1) maps to the progressive myoclonus epilepsy

[188968](EPM1) critical region on chromosome 21q22.3. Genomics 33: 298–300,

[188969]1996.

[188970]

[188971]

[188972]19967.

[188973]Hayashizaki, Y.; Shibata, H.; Hirotsune, S.; Sugino, H.; Okazaki,

[188974]Y.; Sasaki, N.; Hirose, K.; Imoto, H.; Okuizumi, H.; Muramatsu, M.;

[188975]Komatsubara, H.; Shiroishi, T.; Moriwaki, K.; Katsuki, M.; Hatano,

[188976]N.; Sasaki, H.; Ueda, T.; Mise, N.; Takagi, N.; Plass, C.; Chapman,

[188977]V. M.: Identification of an imprinted U2af binding protein related

[188978]sequence on mouse chromosome 11 using the RLGS method. Nature Genet. 6:

[188979]33–40, 1994.

[188980]

[188981]

[188982]19968.

[188983]Zhang, M.; Zamore, P. D.; Carmo–Fonseca, M.; Lamond, A.
I.; Green,

[188984]M. R.: Cloning and intracellular localization of the U2
small nuclear

[188985]ribonucleoprotein auxiliary factor small subunit. Proc. Nat.
Acad.

[188986]Sci. 89: 8769–8773, 1992.

[188987]

[188988]

[188989]19969.

[188990]Baker, R. T.; Board, P. G.: The human ubiquitin–52 amino
acid

[188991]fusion protein gene shares several structural features with
mammalian

[188992]ribosomal protein genes. Nucleic Acids Res. 19:
1035–1040, 1991.

[188993]

[188994]

[188995]19970.

[188996]Lund, P. K.; Moats–Staats, B. M.; Simmons, J. G.; Hoyt, E.;
D'Ercole,

[188997]A. J.; Martin, F.; Van Wyk, J. J.: Nucleotide sequence analysis of

[188998]a cDNA encoding human ubiquitin reveals that ubiquitin is
synthesized

[188999]as a precursor. J. Biol. Chem. 260: 7609–7613, 1985.

[189000]

[189001]

[189002]19971.

[189003]Webb, G. C.; Baker, R. T.; Coggan, M.; Board, P. G.: Localization
ization

[189004]of the human UBA52 ubiquitin fusion gene to chromosome band 19p13.1–p12. Genomics 19:

[189005]567–569, 1994.

[189006]

[189007]

[189008]19972.

[189009]Duncan, A. M. V.; Anderson, L.; Duff, C.; Ozawa, T.;
Suzuki, H.;

[189010]Worton, R.; Rozen, R.: Assignment of the gene (UQCRFS1)
for the Rieske

[189011]iron-sulfur protein subunit of the mitochondrial cy-
tochrome bc-1 complex

[189012]to the 22q13 and 19q12-q13.1 regions of the human
genome. Genomics 21:

[189013]281-283, 1994.

[189014]

[189015]

[189016]19973.

[189017]Pennacchio, L. A.; Bergmann, A.; Fukushima, A.; Okubo,
K.; Salemi,

[189018]A.; Lennon, G. G.: Structure, sequence and location of the
UQCRFS1

[189019]gene for the human Rieske Fe-S protein. Gene 155:
207-211, 1995.

[189020]

[189021]

[189022]19974.

[189023]Cunningham, J. M.; Vanin, E. F.; Tran, N.; Valentine, M.;
Jane,

[189024]S. M.: The human transcription factor CP2 (TFCP2), a com-
ponent of

[189025]the human gamma-globin stage selector protein, maps to
chromosome

[189026]region 12q13 and is within 250 kb of the NF-E2 gene. Ge-
nomics 30:

[189027]398-399, 1995.

[189028]

[189029]

[189030]19975.

[189031]Jane, S. M.; Nienhuis, A. W.; Cunningham, J. M.:
Hemoglobin switching

[189032]in man and chicken is mediated by a heteromeric complex
between the

[189033]ubiquitous transcription factor CP2 and a developmentally
specific

[189034]protein. EMBO J. 13: 197-105, 1995.

[189035]

[189036]

[189037]19976.

[189038]Lambert, J.-C.; Goumidi, L.; Wavrant-De Vrieze, F.;
Frigard, B.;

[189039]Harris, J. M.; Cummings, A.; Coates, J.; Pasquier, F.; Cotel,
D.;

[189040]Gaillac, M.; St. Clair, D.; Mann, D. M. A.; Hardy, J.; Lendon,
C.

[189041]L.; Amouyel, P.; Chartier-Harlin, M.-C.: The transcriptional
factor

[189042]LBP-1c/CP2/LSF gene on chromosome 12 is a genetic de-
terminant of Alzheimer's

[189043]disease. Hum. Molec. Genet. 9: 2275-2280, 2000.

[189044]

[189045]

[189046]19977.

[189047]Swendeman, S. L.; Spielholz, C.; Jenkins, N. A.; Gilbert, D.
J.;

[189048]Copeland, N. G.; Sheffery, M.: Characterization of the ge-
nomic structure,

[189049]chromosomal location, promoter, and developmental ex-

pression of the

[189050]alpha-globin transcription factor CP2. J. Biol. Chem. 269:
11663–11671,

[189051]1994.

[189052]

[189053]

[189054]19978.

[189055]Taylor, A. E.; Yip, A.; Brayne, C.; Easton, D.; Evans, J. G.;
Xuereb,

[189056]J.; Cairns, N.; Esiri, M. M.; Rubinsztein, D. C.: Genetic as-
sociation

[189057]of an LBP-1c/CP2/LSF gene polymorphism with late onset
Alzheimer's

[189058]disease. J. Med. Genet. 38: 232–233, 2001.

[189059]

[189060]

[189061]19979.

[189062]Chen, C.-R.; Kang, Y.; Siegel, P. M.; Massague, J.: E2F4/5
and

[189063]p107 as Smad cofactors linking the TGF-beta receptor to

c-myc repression. Cell 110:

[189064]19–32, 2002.

[189065]

[189066]

[189067]19980.

[189068]Zhang, Y.; Venkatraj, V. S.; Fischer, S. G.; Warburton, D.;
Chellappan,

[189069]S. P.: Genomic cloning and chromosomal assignment of
the E2F dimerization

[189070]partner TFDP gene family. Genomics 39: 95–98, 1997.

[189071]

[189072]

[189073]19981.

[189074]Moloney, D. J.; Panin, V. M.; Johnston, S. H.; Chen, J.; Shao,

[189075]L.; Wilson, R.; Wang, Y.; Stanley, P.; Irvine, K. D.; Halti-
wanger,

[189076]R. S.; Vogt, T. F.: Fringe is a glycosyltransferase that mod-
ifies

[189077]Notch. Nature 406: 369–375, 2000.

[189078]

[189079]

[189080]19982.

[189081]Sestan, N.; Artavanis-Tsakonas, S.; Rakic, P.: Contact-dependent

[189082]inhibition of cortical neurite growth mediated by Notch signaling. Science 286:

[189083]741–746, 1999.

[189084]

[189085]

[189086]19983.

[189087]Tanigaki, K.; Nogaki, F.; Takahashi, J.; Tashiro, K.; Kurooka,

[189088]H.; Honjo, T.: Notch1 and Notch3 instructively restrict bFGF-responsive

[189089]multipotent neural progenitor cells to an astroglial fate. Neuron 29:

[189090]45–55, 2001.

[189091]

[189092]

[189093]19984.

[189094]Barton, D. E.; Foellmer, B. E.; Du, J.; Tamm, J.; Derynck, R.;

[189095]Francke, U.: Chromosomal mapping of genes for trans-
forming growth

[189096]factors beta-2 and beta-3 in man and mouse: dispersion
of TGF-beta

[189097]gene family. Oncogene Res. 3: 323-331, 1988.

[189098]

[189099]

[189100]19985.

[189101]de Martin, R.; Haendler, B.; Hofer-Warbinek, R.; Gaugi-
tisch, H.;

[189102]Wrann, M.; Schusener, H.; Seifert, J. M.; Bodmer, S.;

Fontana, A.;

[189103]Hofer, E.: Complementary DNA for human glioblastoma-de-
rived T cell

[189104]suppressor factor, a novel member of the transforming
growth factor-beta

[189105]gene family. EMBO J. 6: 3673-3677, 1987.

[189106]

[189107]

[189108]19986.

[189109]Hanks, S. K.; Armour, R.; Baldwin, J. H.; Maldonado, F.;

Spiess,

[189110]J.; Holley, W.: Amino acid sequence of the BSC-1 cell

growth inhibitor

[189111](polyergin) deduced from the nucleotide sequence of the
cDNA. Proc.

[189112]Nat. Acad. Sci. 85: 79-82, 1988.

[189113]

[189114]

[189115]19987.

[189116]Madisen, L.; Webb, N. R.; Rose, T. M.; Marquardt, H.;

Ikeda, T.;

[189117]Twardzik, D.; Seyedin, S.; Purchio, A. F.: Transforming

growth factor-beta-2:

[189118]cDNA cloning and sequence analysis. DNA 7: 1-8, 1988.

[189119]

[189120]

[189121]19988.

[189122]Dean, M.; Park, M.; Vande Woude, G. F.: Characterization
of the

[189123]rearranged TPR–MET oncogene breakpoint. Molec. Cell.
Biol. 7: 921–924,

[189124]1987.

[189125]

[189126]

[189127]19989.

[189128]Robinson, B. H.; Taylor, J.; Sherwood, W. G.: The genetic
heterogeneity

[189129]of lactic acidosis: occurrence of recognisable inborn errors
of metabolism

[189130]in a pediatric population with lactic acidosis. Pediat. Res.
14:

[189131]950–962, 1980.

[189132]

[189133]

[189134]19990.

[189135]Hosokawa, Y.; Suzuki, H.; Nishikimi, M.; Matsukage, A.;
Yoshida,

- [189136]M. C.; Ozawa, T.: Chromosomal assignment of the gene
for the ubiquinone-binding
[189137]protein of human mitochondrial cytochrome bc₁ complex.
Biochem. Int. 21:
[189138]41–44, 1990.
- [189139]
- [189140]
- [189141]19991.
- [189142]Suzuki, H.; Hosokawa, Y.; Toda, H.; Nishikimi, M.; Ozawa,
T.:
[189143]Cloning and sequencing of a cDNA for human mitochon-
drial ubiquinone-binding
[189144]protein of complex III. Biochem. Biophys. Res. Commun.
156: 987–994,
[189145]1988.
- [189146]
- [189147]
- [189148]19992.
- [189149]van Leeuwen, F. W.; de Kleijn, D. P. V.; van den Hurk, H.
H.; Neubauer,

[189150]A.; Sonnemans, M. A. F.; Sluijs, J. A.; Koycu, S.; Ramdjielal, R.

[189151]D. J.; Salehi, A.; Martens, G. J. M.; Grosveld, F. G.; Burbach, J.

[189152]P. H.; Hol, E. M.: Frameshift mutants of beta-amyloid precursor protein

[189153]and ubiquitin-B in Alzheimer's and Down patients. Science 279: 242-247,

[189154]1998.

[189155]

[189156]

[189157]19993.

[189158]Webb, G. C.; Baker, R. T.; Fagan, K.; Board, P. G.: Localization

[189159]of the human UbB polyubiquitin gene to chromosome band 17p11.1-17p12. Am.

[189160]J. Hum. Genet. 46: 308-315, 1990.

[189161]

[189162]

[189163]19994.

[189164]Day, I. N. M.; Hinks, L. J.; Thompson, R. J.: The structure of
[189165]the gene encoding protein gene product 9.5 (PGP9.5), a
neuron-specific
[189166]ubiquitin C-terminal hydrolase. Biochem. J. 268: 521–524,
1990.

[189167]

[189168]

[189169]19995.

[189170]Day, I. N. M.; Thompson, R. J.: Molecular cloning of cDNA
coding
[189171]for human PGP 9.5 protein: a novel cytoplasmic marker for
neurones

[189172]and neuroendocrine cells. FEBS Lett. 210: 157–160, 1987.

[189173]

[189174]

[189175]19996.

[189176]Doran, J. F.; Jackson, P.; Kynoch, P.; Thompson, R. J.: Iso-
lation

[189177]of PGP 9.5, a new human neurone-specific protein de-
tected by high

[189178]resolution two-dimensional electrophoresis. J. Neurochem. 40: 1542–1547,
[189179]1983.

[189180]

[189181]

[189182]19997.

[189183]Edwards, Y. H.; Fox, M. F.; Povey, S.; Hinks, L. J.; Day, I. N.

[189184]M.; Thompson, R. J.: The gene for human neuron specific
ubiquitin

[189185]C-terminal hydrolase maps to chromosome 4p14.
(Abstract) Cytogenet.

[189186]Cell Genet. 58: 1886–1887, 1991.

[189187]

[189188]

[189189]19998.

[189190]Kurihara, L. J.; Kikuchi, T.; Wada, K.; Tilghman, S. M.: Loss

[189191]of Uch-L1 and Uch-L3 leads to neurodegeneration, posterior
paralysis

[189192]and dysphagia. Hum. Molec. Genet. 10: 1963–1970,

2001.

[189193]

[189194]

[189195]1999.

[189196]Kurihara, L. J.; Semenova, E.; Levorse, J. M.; Tilghman, S.

M.

[189197]: Expression and functional analysis of Uch-L3 during mouse development. Molec.

[189198]Cell. Biol. 20: 2498–2504, 2000.

[189199]

[189200]

[189201]20000.

[189202]Leroy, E.; Boyer, R.; Auburger, G.; Leube, B.; Ulm, G.;

Mezey,

[189203]E.; Harta, G.; Brownstein, M. J.; Jonnalagada, S.; Chernova,

T.; Dehejia,

[189204]A.; Lavedan, C.; Gasser, T.; Steinbach, P. J.; Wilkinson, K.

D.; Polymeropoulos,

[189205]M. H.: The ubiquitin pathway in Parkinson's disease.

(Letter) Nature 395:

[189206]451–452, 1998.

[189207]

[189208]

[189209]20001.

[189210]Lincoln, S.; Vaughan, J.; Wood, N.; Baker, M.; Adamson, J.;
Gwinn-Hardy,

[189211]K.; Lynch, T.; Hardy, J.; Farrer, M.: Low frequency of
pathogenic

[189212]mutations in the ubiquitin carboxy-terminal hydrolase
gene in familial

[189213]Parkinson's disease. Neuroreport 10: 427-429, 1999.

[189214]

[189215]

[189216]20002.

[189217]MacDonald, M. E.: Gadzooks! Nature Genet. 23: 10-11,
1999.

[189218]

[189219]

[189220]20003.

[189221]Saigoh, K.; Wang, Y.-L.; Suh, J.-G.; Yamanishi, T.; Sakai,

Y.;

[189222]Kiyosawa, H.; Harada, T.; Ichihara, N.; Wakana, S.; Kikuchi, T.; Wada,

[189223]K.: Intragenic deletion in the gene encoding ubiquitin carboxy-terminal

[189224]hydrolase in gad mice. Nature Genet. 23: 47–51, 1999.

[189225]

[189226]

[189227]20004.

[189228]Suh, J. G.; Yamanishi, T.; Matsui, K.; Tanaka, K.; Wada, K.:

[189229]Mapping of the gracile axonal dystrophy (gad) gene to a region between

[189230]D5Mit197 and D5Mit113 on proximal mouse chromosome 5. Genomics 27:

[189231]549–551, 1995.

[189232]

[189233]

[189234]20005.

[189235]Wilkinson, K. D.; Lee, K. M.; Deshpande, S.; Duerksen-Hughes,

[189236]P.; Boss, J. M.; Pohl, J.: The neuron-specific protein PGP
9.5 is

[189237]a ubiquitin carboxyl-terminal hydrolase. Science 246:
670-672, 1989.

[189238]

[189239]

[189240]20006.

[189241]Yamazaki, K.; Wakasugi, N.; Tomita, T.; Kikuchi, T.;
Mukoyama,

[189242]M.; Ando, K.: Gracile axonal dystrophy (GAD), a new neu-
rological

[189243]mutant in the mouse. Proc. Soc. Exp. Biol. Med. 187:
209-215, 1988.

[189244]

[189245]

[189246]20007.

[189247]Vidal, R.; Frangione, B.; Rostagno, A.; Mead, S.; Revesz, T.;
Plant,

[189248]G.; Ghiso, J.: A stop-codon mutation in the BRI gene asso-
ciated with

[189249]familial British dementia. Nature 399: 776-781, 1999.

[189250]

[189251]

[189252]20008.

[189253]Adler, D. A.; Tseng, B. Y.; Wang, T. S.-F.; Disteché, C. M.:
Physical

[189254]mapping of the genes for three components of the mouse
DNA replication

[189255]complex: polymerase alpha to the X chromosome, pri-
mase p49 subunit

[189256]to chromosome 10, and primase p58 subunit to chromo-
some 1. Genomics 9:

[189257]642-646, 1991.

[189258]

[189259]

[189260]20009.

[189261]Ogilvie, A. D.; Battersby, S.; Bubb, V. J.; Fink, G.; Harmar,

[189262]A. J.; Goodwin, G. M.; Smith, C. A. D.: Polymorphism in
serotonin

[189263]transporter gene associated with susceptibility to major
depression. Lancet 347:

[189264]731–733, 1996.

[189265]

[189266]

[189267]20010.

[189268]Doniger, J.; DiPaolo, J. A.: Coordinate N-RAS mRNA up-regulation

[189269]with mutational activation in tumorigenic guinea pig cells.
Nucleic

[189270]Acids Res. 16: 969–980, 1988.

[189271]

[189272]

[189273]20011.

[189274]Jeffers, M.; Paciucci, R.; Pellicer, A.: Characterization of
UNR:

[189275]a gene closely linked to N-RAS. Nucleic Acids Res. 18:
4891–4899,

[189276]1990.

[189277]

[189278]

[189279]20012.

[189280]Mirnics, K.; Middleton, F. A.; Marquez, A.; Lewis, D. A.;
Levitt,

[189281]P.: Molecular characterization of schizophrenia viewed by
microarray

[189282]analysis of gene expression in prefrontal cortex. Neuron
28: 53–67,

[189283]2000.

[189284]

[189285]

[189286]20013.

[189287]Engelkamp, D.; Schafer, B. W.; Mattei, M. G.; Erne, P.;
Heizmann,

[189288]C. W.: Six S100 genes are clustered on human chromo-
some 1q21: identification

[189289]of two genes coding for the two previously unreported
calcium-binding

[189290]proteins S100D and S100E. Proc. Nat. Acad. Sci. 90:
6547–6551, 1993.

[189291]

[189292]

[189293]20014.

[189294]Herrera, G. A.; Turbat-Herrera, E. A.; Lott, R. L.: S-100
protein

[189295]expression by primary and metastatic adenocarcinomas.
Am. J. Clin.

[189296]Path. 89: 168-176, 1988.

[189297]

[189298]

[189299]20015.

[189300]Isobe, T.; Tsugira, A.; Okuyama, T.: Amino acid sequence
of the

[189301]subunit structure of bovine brain S-100 protein (PAP 1-b).
J. Neurochem. 30:

[189302]921-923, 1978.

[189303]

[189304]

[189305]20016.

[189306]Moore, B. W.: A soluble protein characteristic of the ner-
vous

[189307]system. Biochem. Biophys. Res. Commun. 19: 739-744,

1965.

[189308]

[189309]

[189310]20017.

[189311]Morii, K.; Tanaka, R.; Takahashi, Y.; Minoshima, S.;

Fukuyama,

[189312]R.; Shimizu, N.; Kuwano, R.: Structure and chromosome
assignment

[189313]of human S100 alpha and beta subunit genes. Biochem.
Biophys. Res.

[189314]Commun. 175: 185–191, 1991.

[189315]

[189316]

[189317]20018.

[189318]Most, P.; Bernotat, J.; Ehlermann, P.; Pleger, S. T.; Reppel,
M.;

[189319]Borries, M.; Niroomand, F.; Pieske, B.; Janssen, P. M. L.;

Eschenhagen,

[189320]T.; Karczewski, P.; Smith, G. L.; Koch, W. J.; Katus, H. A.;

Remppis,

[189321]A.: S100A1: a regulator of myocardial contractility. Proc.

Nat.

[189322]Acad. Sci. 98: 13889–13894, 2001.

[189323]

[189324]

[189325]20019.

[189326]Asada, Y.; Nadeau, J. H.: Fert is on mouse chromosome
11, not

[189327]chromosome 17. Mammalian Genome 5: 830 only, 1994.

[189328]

[189329]

[189330]20020.

[189331]Hao, Q.-L.; Heisterkamp, N.; Groffen, J.: Isolation and se-
quence

[189332]analysis of a novel human tyrosine kinase gene. Molec.
Cell. Biol. 9:

[189333]1587–1593, 1989.

[189334]

[189335]

[189336]20021.

- [189337]Dryja, T. P.; Grondin, V. J.; Ringens, P.; Cotran, P.; Berson, E. L.; Travis, G.: Isolation of human retinal cDNA fragments homologous to the murine rds gene transcript. (Abstract) Invest. Ophthalmol. Vis. Sci. 30 (suppl.): 43 only, 1989.
- [189340]
- [189341]
- [189342]
- [189343]20022.
- [189344]Farrar, G. J.; Jordan, S. A.; Kenna, P.; Humphries, M. M.; Kumar-Singh, R.; McWilliam, P.; Allamand, V.; Sharp, E.; Humphries, P.: Autosomal dominant retinitis pigmentosa: localization of a disease gene (RP6) to the short arm of chromosome 6. Genomics 11: 870-874, 1991.
- [189345]
- [189346]
- [189347]20023.

[189351]Farrar, G. J.; Kenna, P.; Jordan, S. A.; Kumar–Singh, R.;
Humphries,

[189352]M. M.; Sharp, E. M.; Sheils, D. M.; Humphries, P.: A three–
base–pair

[189353]deletion in the peripherin–RDS gene in one form of retini–
tis pigmentosa. Nature 354:

[189354]478–480, 1991.

[189355]

[189356]

[189357]20024.

[189358]Feist, R. M.; White, M. F., Jr.; Skalka, H.; Stone, E. M.:
Choroidal

[189359]neovascularization in a patient with adult foveomacular
dystrophy

[189360]and a mutation in the retinal degeneration slow gene
(pro210–to–arg). Am.

[189361]J. Ophthal. 118: 259–260, 1994.

[189362]

[189363]

[189364]20025.

[189365]Felbor, U.; Schilling, H.; Weber, B. H. F.: Adult vitelliform

[189366]macular dystrophy is frequently associated with mutations
in the peripherin/RDS

[189367]gene. Hum. Mutat. 10: 301–309, 1997.

[189368]

[189369]

[189370]20026.

[189371]Gass, J. D. M.: A clinicopathologic study of a peculiar
foveomacular

[189372]dystrophy. Trans. Am. Ophthal. Soc. 72: 139–155, 1974.

[189373]

[189374]

[189375]20027.

[189376]Jackson, K. E.; Mitchell, E. B.; Stone, E. M.; Ferrell, R. E.;

[189377]Gorin, M. B.: The identification of an exon–2 peripherin
mutation

[189378]in a family with heterogeneous manifestations of a butter-
fly pattern

[189379]macular dystrophy. (Abstract) Am. J. Hum. Genet. 53

(suppl.): 1177

[189380]only, 1993.

[189381]

[189382]

[189383]20028.

[189384]Jordan, S. A.; Farrar, G. J.; Kumar–Singh, R.; Kenna, P.;
Humphries,

[189385]M. M.; Allamand, V.; Sharp, E. M.; Humphries, P.: Autosomal dominant
mal dominant

[189386]retinitis pigmentosa (adRP; RP6): cosegregation of RP6
and the peripherin–RDS

[189387]locus in a late–onset family of Irish origin. Am. J. Hum.
Genet. 50:

[189388]634–639, 1992.

[189389]

[189390]

[189391]20029.

[189392]Okamoto, H.; Yonemori, F.; Wakitani, K.; Minowa, T.;
Maeda, K.;

[189393]Shinkai, H.: A cholesteryl ester transfer protein inhibitor

attenuates

[189394]atherosclerosis in rabbits. Nature 406: 203–207, 2000.

[189395]

[189396]

[189397]20030.

[189398]Oliveira, H. C. F.; Chouinard, R. A.; Agellon, L. B.; Bruce, C.;

[189399]Ma, L.; Walsh, A.; Breslow, J. L.; Tall, A. R.: Human cholesteryl

[189400]ester transfer protein gene proximal promoter contains dietary cholesterol

[189401]positive responsive elements and mediates expression in small intestine

[189402]and periphery while predominant liver and spleen expression is controlled

[189403]by 5–prime–distal sequences. J. Biol. Chem. 271: 31831–31838, 1996.

[189404]

[189405]

[189406]20031.

[189407]Sakai, N.; Yamashita, S.; Hirano, K.; Menju, M.; Arai, T.;

Kobayashi,

[189408]K.; Ishigami, M.; Yoshida, Y.; Hoshino, T.; Nakajima, N.;

Kameda-Takemura,

[189409]K.; Matsuzawa, Y.: Frequency of exon 15 missense mutation (442D:G)

[189410]in cholesteryl ester transfer protein gene in hyperlipoproteinemic

[189411]Japanese subjects. *Atherosclerosis* 114: 139–145, 1995.

[189412]

[189413]

[189414]20032.

[189415]Sparkes, R. S.; Drayna, D.; Mohandas, T.; Klisak, I.; Heinzmann,

[189416]C.; Lawn, R.; Lusis, A. J.: Assignment of cholesterol ester transfer

[189417]protein (CETP) gene to human 16q21. (Abstract) *Cytogenet. Cell Genet.* 46:

[189418]696 only, 1987.

[189419]

[189420]

[189421]20033.

[189422]Takahashi, K.; Jiang, X.-C.; Sakai, N.; Yamashita, S.; Hirano,

[189423]K.; Bujo, H.; Yamazaki, H.; Kusunoki, J.; Miura, T.; Kussie, P.; Matsuzawa,

[189424]Y.; Saito, Y.; Tall, A.: A missense mutation in the cholesteryl ester

[189425]transfer protein gene with possible dominant effects on plasma high

[189426]density lipoproteins. J. Clin. Invest. 92: 2060–2064, 1993.

[189427]

[189428]

[189429]20034.

[189430]Yamashita, S.; Hui, D. Y.; Sprecher, D. L.; Matsuzawa, Y.; Sakai,

[189431]N.; Tarui, S.; Kaplan, D.; Wetterau, J. R.; Harmony, J. A.: Total

[189432]deficiency of plasma cholesteryl ester transfer protein in subjects

[189433]homozygous and heterozygous for the intron 14 splicing defect. Biochem.

[189434]Biophys. Res. Commun. 170: 1346–1351, 1990.

[189435]

[189436]

[189437]20035.

[189438]Zhong, S.; Sharp, D. S.; Grove, J. S.; Bruce, C.; Yano, K.;
Curb,

[189439]J. D.; Tall, A. R.: Increased coronary heart disease in
Japanese-American

[189440]men with mutation in the cholesteryl ester transfer protein
gene despite

[189441]increased HDL levels. J. Clin. Invest. 97: 2917–2923,
1996.

[189442]

[189443]

[189444]20036.

[189445]Chung, B.-C.; Matteson, K. J.; Voutilainen, R.; Mohandas,
T. K.;

[189446]Miller, W. L.: Human cholesterol side-chain cleavage en-
zyme, P450scc:

[189447]cDNA cloning, assignment of the gene to chromosome 15,
and expression

[189448]in the placenta. Proc. Nat. Acad. Sci. 83: 8962–8966,

1986.

[189449]

[189450]

[189451]20037.

[189452]Durocher, F.; Morissette, J.; Simard, J.: Genetic linkage mapping

[189453]of the CYP11A1 gene encoding the cholesterol side-chain cleavage P450(scc)

[189454]close to the CYP1A1 gene and D15S204 in the chromosome 15q22.33-q23

[189455]region. Pharmacogenetics 8: 49-53, 1998.

[189456]

[189457]

[189458]20038.

[189459]Billingsley, G. D.; Walter, M. A.; Hammond, G. L.; Cox, D. W.:

[189460]Physical mapping of four serpin genes: alpha-1-antitrypsin, alpha-1-antichymotrypsin,

[189461]corticosteroid-binding globulin, and protein C inhibitor, within a

[189462]280-kb region on chromosome 14q32.1. Am. J. Hum.

Genet. 52: 343–353,

[189463]1993.

[189464]

[189465]

[189466]20039.

[189467]Wang, J.; Hannon, G. J.; Beach, D. H.: Risky immortaliza-
tion

[189468]by telomerase. (Letter) Nature 405: 755–756, 2000.

[189469]

[189470]

[189471]20040.

[189472]Wu, K.-J.; Grandori, C.; Amacker, M.; Simon-Vermot, N.;
Polack,

[189473]A.; Lingner, J.; Dalla-Favera, R.: Direct activation of TERT
transcription

[189474]by c-MYC. Nature Genet. 21: 220–224, 1999.

[189475]

[189476]

[189477]20041.

[189478]Greco, A.; Ittmann, M.; Barletta, C.; Basilico, C.; Croce, C.
M.;

[189479]Cannizzaro, L. A.; Huebner, K.: Chromosomal localization
of human

[189480]genes required for G(1) progression in mammalian cells.
Genomics 4:

[189481]240–245, 1989.

[189482]

[189483]

[189484]20042.

[189485]Ittmann, M.; Greco, A.; Basilico, C.: Isolation of the human
gene

[189486]that complements a temperature-sensitive cell cycle mu-
tation in BHK

[189487]cells. Molec. Cell. Biol. 7: 3386–3393, 1987.

[189488]

[189489]

[189490]20043.

[189491]Ip, N. Y.; Stitt, T. N.; Tapley, P.; Klein, R.; Glass, D. J.;
Fandl,

[189492]J.; Greene, L. A.; Barbacid, M.; Yancopoulos, G. D.: Similarities

[189493]and differences in the way neurotrophins interact with the Trk receptors

[189494]in neuronal and nonneuronal cells. Neuron 10: 137–149, 1993.

[189495]

[189496]

[189497]20044.

[189498]Knezevich, S. R.; McFadden, D. E.; Tao, W.; Lim, J. F.; Sorensen,

[189499]P. H. B.: A novel ETV6–NTRK3 gene fusion in congenital fibrosarcoma. Nature

[189500]Genet. 18: 184–187, 1998.

[189501]

[189502]

[189503]20045.

[189504]Bonnetfont, J.–P.; Chretien, D.; Rustin, P.; Robinson, B.; Vassault,

[189505]A.; Aupetit, J.; Charpentier, C.; Rabier, D.; Saudubray, J.–M.; Munnich,

[189506]A.: Alpha-ketoglutarate dehydrogenase deficiency presenting as congenital

[189507]lactic acidosis. J. Pediat. 121: 255–258, 1992.

[189508]

[189509]

[189510]20046.

[189511]Guffon, N.; Lopez-Mediavilla, C.; Dumoulin, R.; Mousson, B.; Godinot,

[189512]C.; Carrier, H.; Collombet, J. M.; Divry, P.; Mathieu, M.; Guibaud,

[189513]P.: 2-Ketoglutarate dehydrogenase deficiency, a rare cause of primary

[189514]hyperlactataemia: report of a new case. J. Inherit. Metab. Dis. 16:

[189515]821–830, 1993.

[189516]

[189517]

[189518]20047.

[189519]Kohlschutter, A.; Behbehani, A.; Langenbeck, U.; Albani, M.; Heidemann,

[189520]P.; Hoffmann, G.; Kleineke, J.; Lehnert, W.; Wendel, U.: A
familial

[189521]progressive neurodegenerative disease with 2-oxoglutaric
aciduria. Europ.

[189522]J. Pediat. 138: 32-37, 1982.

[189523]

[189524]

[189525]20048.

[189526]Koike, K.: The gene encoding human 2-oxoglutarate de-
hydrogenase:

[189527]structural organization and mapping to chromosome
7p13-p14. Gene 159:

[189528]261-266, 1995.

[189529]

[189530]

[189531]20049.

[189532]Koike, K.; Urata, Y.; Goto, S.: Cloning and nucleotide se-
quence

[189533]of the cDNA encoding human 2-oxoglutarate dehydroge-
nase (lipoamide). Proc.

[189534]Nat. Acad. Sci. 89: 1963–1967, 1992.

[189535]

[189536]

[189537]20050.

[189538]Szabo, P.; Cai, X.; Ali, G.; Blass, J. P.: Localization of the

[189539]gene (OGDH) coding for the E1 α component of the alpha-ketoglutarate

[189540]dehydrogenase complex to chromosome 7p13–p11.2.

Genomics 20: 324–326,

[189541]1994.

[189542]

[189543]

[189544]20051.

[189545]Kury, S.; Dreno, B.; Bezieau, S.; Giraudet, S.; Kharfi, M.;

Kamoun,

[189546]R.; Moisan, J.–P.: Identification of SLC39A4, a gene involved in

[189547]acrodermatitis enteropathica. Nature Genet. 31: 239–240, 2002.

[189548]

[189549]

[189550]20052.

[189551]Rogers, E. E.; Eide, D. J.; Guerinot, M. L.: Altered selectivity

[189552]in an Arabidopsis metal transporter. Proc. Nat. Acad. Sci.

97: 12356–12360,

[189553]2000.

[189554]

[189555]

[189556]20053.

[189557]Wang, K.; Pugh, E. W.; Griffen, S.; Doheny, K. F.; Mostafa, W.

[189558]Z.; al-Aboosi, M. M.; el-Shanti, H.; Gitschier, J.: Homozygosity

[189559]mapping places the acrodermatitis enteropathica gene on chromosomal

[189560]region 8q24.3. Am. J. Hum. Genet. 68: 1055–1060, 2001.

[189561]

[189562]

[189563]20054.

[189564]Haberle, J.; Pauli, S.; Linnebank, M.; Kleijer, W. J.; Bakker,

[189565]H. D.; Wanders, R. J. A.; Harms, E.; Koch, H. G.: Structure of the

[189566]human argininosuccinate synthetase gene and an improved system for

[189567]molecular diagnostics in patients with classical and mild

citrullinemia. Hum.

[189568]Genet. 110: 327–333, 2002.

[189569]

[189570]

[189571]20055.

[189572]Barbosa, J.; Rich, S.; Dunsworth, T.; Swanson, J.: Linkage disequilibrium

[189573]between insulin-dependent diabetes and the Kidd blood group Jk(b)

[189574]allele. J. Clin. Endocr. Metab. 55: 193–195, 1982.

[189575]

[189576]

[189577]20056.

[189578]Harper, P. A.; Healy, P. J.; Dennis, J. A.; O'Brien, J. J.; Rayward,

[189579]D. H.: Citrullinaemia as a cause of neurological disease in neonatal

[189580]Friesian calves. Aust. Vet. J. 63: 378–379, 1986.

[189581]

[189582]

[189583]20057.

[189584]Harper, P. A. W.; Healy, P. J.; Dennis, J. A.: Animal model of
[189585]human disease: citrullinemia (argininosuccinate syn-
thetase deficiency). Am.

[189586]J. Path. 135: 1213–1215, 1989.

[189587]

[189588]

[189589]20058.

[189590]Jackson, M. J.; Surh, L. C.; O'Brien, W. E.; Beaudet, A. L.:

[189591]Assignment of the structural gene for argininosuccinate
synthetase

[189592]to proximal mouse chromosome 2. Genomics 6: 545–547,
1990.

[189593]

[189594]

[189595]20059.

[189596]Kobayashi, K.; Ichiki, H.; Saheki, T.; Tatsuno, M.;
Uchiyama,

[189597]C.; Nukada, O.; Yoda, T.: Structure of an abnormal messenger RNA

[189598]for argininosuccinate synthetase in citrullinemia. Hum. Genet. 76:

[189599]27-32, 1987.

[189600]

[189601]

[189602]20060.

[189603]Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.; Beaudet,

[189604]A. L.: Characterization of nine mutant alleles causing citrullinemia.

[189605](Abstract) Am. J. Hum. Genet. 45 (suppl.): A201 only, 1989.

[189606]

[189607]

[189608]20061.

[189609]Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.; Beaudet,

[189610]A. L.: Heterogeneity of mutations in argininosuccinate synthetase

[189611]causing human citrullinemia. J. Biol. Chem. 265:
11361–11367, 1990.

[189612]

[189613]

[189614]20062.

[189615]Kobayashi, K.; Kakinoki, H.; Fukushige, T.; Shaheen, N.;
Terazono,

[189616]H.; Saheki, T.: Nature and frequency of mutations in the
argininosuccinate

[189617]synthetase gene that cause classical citrullinemia. Hum.
Genet. 96:

[189618]454–463, 1995.

[189619]

[189620]

[189621]20063.

[189622]Kobayashi, K.; Rosenbloom, C.; Beaudet, A. L.; O'Brien, W.
E.

[189623]: Additional mutations in argininosuccinate synthetase
causing citrullinemia. Molec.

[189624]Biol. Med. 8: 95–100, 1991.

[189625]

[189626]

[189627]20064.

[189628]Kobayashi, K.; Saheki, T.; Imamura, Y.; Noda, T.; Inoue, I.;
Matuo,

[189629]S.; Hagihara, S.; Nomiyama, H.; Jinno, Y.; Shimada, K.:
Messenger

[189630]RNA coding for argininosuccinate synthetase in citrulline-
mia. Am.

[189631]J. Hum. Genet. 38: 667-680, 1986.

[189632]

[189633]

[189634]20065.

[189635]Kobayashi, K.; Shaheen, N.; Terazono, H.; Saheki, T.: Mu-
tations

[189636]in argininosuccinate synthetase mRNA of Japanese pa-
tients, causing

[189637]classical citrullinemia. Am. J. Hum. Genet. 55: 1103-1112,
1994.

[189638]

[189639]

[189640]20066.

[189641]Li, C.-M.; Chao, H.-K.; Liu, Y.-F.; Su, T.-S.: A nonsense mutation

[189642]is responsible for the RNA-negative phenotype in human citrullinaemia. Europ.

[189643]J. Hum. Genet. 9: 685-689, 2001.

[189644]

[189645]

[189646]20067.

[189647]Northrup, H.; Lathrop, M.; Lu, S.-Y.; Daiger, S. P.; Beaudet,

[189648]A. L.; O'Brien, W. E.: Multilocus linkage analysis with the human

[189649]argininosuccinate synthetase gene. Genomics 5: 442-444, 1989.

[189650]

[189651]

[189652]20068.

[189653]Sase, M.; Kobayashi, K.; Imamura, Y.; Saheki, T.; Nakano,

K.;

[189654]Miura, S.; Mori, M.: Level of translatable messenger RNA coding for

[189655]argininosuccinate synthetase in the liver of the patients with quantitative-type

[189656]citrullinemia. Hum. Genet. 69: 130–134, 1985.

[189657]

[189658]

[189659]20069.

[189660]Todd, S.; Naylor, S. L.: New chromosomal mapping assignments

[189661]for argininosuccinate synthetase pseudogene 1, interferon-beta-3 gene,

[189662]and the diazepam binding inhibitor gene. Somat. Cell Molec. Genet. 18:

[189663]381–385, 1992.

[189664]

[189665]

[189666]20070.

[189667]Rousseau-Merck, M. F.; Pizon, V.; Tavitian, A.; Berger, R.:

Chromosome

[189668]mapping of the human RAS related RAP1A, RAP1B and RAP2 genes to chromosomes

[189669]1p13-12, 12q14 and 13q34, respectively. (Abstract) Cytogenet. Cell

[189670]Genet. 51: 1070 only, 1989.

[189671]

[189672]

[189673]20071.

[189674]Rousseau-Merck, M. F.; Pizon, V.; Tavitian, A.; Berger, R.:

Chromosome

[189675]mapping of the human RAS-related RAP1A, RAP1B, and RAP2 genes to chromosomes

[189676]1p12-p13, 12q14, and 13q34, respectively. Cytogenet. Cell Genet. 53:

[189677]2-4, 1990.

[189678]

[189679]

[189680]20072.

[189681]Sebzda, E.; Bracke, M.; Tugal, T.; Hogg, N.; Cantrell, D. A.:

[189682]Rap1a positively regulates T cells via integrin activation rather

[189683]than inhibiting lymphocyte signaling. Nature Immun. 3: 251–258,

[189684]2002.

[189685]

[189686]

[189687]20073.

[189688]Takai, S.; Nishino, N.; Kitayama, H.; Ikawa, Y.; Noda, M.: Mapping

[189689]of the KREV1 transformation suppressor gene and its pseudogene (KREV1P)

[189690]to human chromosome 1p13.3 and 14q24.3, respectively, by fluorescence

[189691]in situ hybridization. Cytogenet. Cell Genet. 63: 59–61, 1993.

[189692]

[189693]

[189694]20074.

[189695]Zhu, J. J.; Qin, Y.; Zhao, M.; Van Aelst, L.; Malinow, R.: Ras

[189696]and Rap control AMPA receptor trafficking during synaptic plasticity. Cell 110: [189697]443–455, 2002.

[189698]

[189699]

[189700]20075.

[189701]Ahmad, F.; Goldstein, B. J.: Functional association between the

[189702]insulin receptor and the transmembrane protein–tyrosine phosphatase

[189703]LAR in intact cells. J. Biol. Chem. 272: 448–457, 1997.

[189704]

[189705]

[189706]20076.

[189707]Disteche, C. M.; Adler, D. A.; Tedder, T. F.; Saito, H.: Map–ping

[189708]of the genes for LYAM1, a new lymphocyte adhesion molecule, and for

[189709]LAR, a new receptor–linked protein tyrosine phosphatase, to human

[189710]chromosome 1 (Abstract) Cytogenet. Cell Genet. 51: 990
only, 1989.

[189711]

[189712]

[189713]20077.

[189714]Harder, K. W.; Saw, J.; Miki, N.; Jirik, F.: Coexisting amplifi-
cations

[189715]of the chromosome 1p32 genes (PTPRF and MYCL1) en-
coding protein tyrosine

[189716]phosphatase LAR and L-myc in a small cell lung cancer
line. Genomics 27:

[189717]552–553, 1995.

[189718]

[189719]

[189720]20078.

[189721]Jirik, F. R.; Harder, K. W.; Melhado, I. G.; Anderson, L. L.;
Duncan,

[189722]A. M. V.: The gene for leukocyte antigen-related tyrosine
phosphatase

[189723](LAR) is localized to human chromosome 1p32, a region
frequently deleted

[189724]in tumors of neuroectodermal origin. Cytogenet. Cell

Genet. 61:

[189725]266–268, 1992.

[189726]

[189727]

[189728]20079.

[189729]Nam, H.-J.; Poy, F.; Krueger, N. X.; Saito, H.; Frederick, C.

A.

[189730]: Crystal structure of the tandem phosphatase domains of

RPTP LAR. Cell 97:

[189731]449–457, 1999.

[189732]

[189733]

[189734]20080.

[189735]O'Grady, P.; Krueger, N. X.; Streuli, M.; Saito, H.: Genomic

organization

[189736]of the human LAR protein tyrosine phosphatase gene and

alternative

[189737]splicing in the extracellular fibronectin type-III domains. J.

Biol.

[189738]Chem. 269: 25193–25199, 1994.

[189739]

[189740]

[189741]20081.

[189742]Schaapveld, R. Q.; Schepens, J. T.; Robinson, G. W.; At-
tema, J.;

[189743]Oerlemans, F. T.; Fransen, J. A.; Streuli, M.; Wieringa, B.;
Hennighausen,

[189744]L.; Hendriks, W. J.: Impaired mammary gland development
and function

[189745]in mice lacking LAR receptor–like tyrosine phosphatase
activity. Dev.

[189746]Biol. 188: 134–146, 1997.

[189747]

[189748]

[189749]20082.

[189750]Schaapveld, R. Q. J.; van den Maagdenberg, A. M. J. M.;
Schepens,

[189751]J. T. G.; Olde Weghuis, D.; Geurts van Kessel, A.; Wieringa,
B.; Hendriks,

[189752]W. J. A. J.: The mouse gene Ptprf encoding the leukocyte common antigen-related

[189753]molecule LAR: cloning, characterization, and chromosomal localization. Genomics 27:

[189754]124–130, 1995.

[189755]

[189756]

[189757]20083.

[189758]Tsujioka, K.; Kawakami, N.; Uchino, Y.; Ichijo, T.; Furukawa,

[189759]T.; Saito, H.; Yamamoto, H.: Distinct functions of the two protein

[189760]tyrosine phosphatase domains of LAR (leukocyte common antigen-related)

[189761]on tyrosine dephosphorylation of insulin receptor. Molec. Endocr. 15:

[189762]271–280, 2001.

[189763]

[189764]

[189765]20084.

[189766]Ali, R. R.; Sarra, G.-M.; Stephens, C.; de Alwis, M.; Bain-
bridge,

[189767]J. W. B.; Munro, P. M.; Fauser, S.; Reichell, M. B.; Kinnon,
C.; Hunt,

[189768]D. M.; Bhattacharya, S. S.; Thrasher, A. J.: Restoration of
photoreceptor

[189769]ultrastructure and function in retinal degeneration slow
mice by gene

[189770]therapy. Nature Genet. 25: 306-310, 2000.

[189771]

[189772]

[189773]20085.

[189774]Bascom, R. A.; Connell, G.; Garcia-Heras, J.; Collins, L.;
Ledbetter,

[189775]D.; Molday, R. S.; Kalnins, V.; McInnes, R. R.: Molecular
and ultrastructural

[189776]characterization of the products of the human retinopathy
candidate

[189777]genes ROM1 and RDS. (Abstract) Am. J. Hum. Genet. 47
(suppl.): A101

[189778]only, 1990.

[189779]

[189780]

[189781]20086.

[189782]Connell, G.; Bascom, R.; Molday, L.; Reid, D.; McInnes, R.
R.;

[189783]Molday, R. S.: Photoreceptor peripherin is the normal
product of

[189784]the gene responsible for retinal degeneration in the rds
mouse. Proc.

[189785]Nat. Acad. Sci. 88: 723–726, 1991.

[189786]

[189787]

[189788]20087.

[189789]Demant, P.; Ivanyi, D.; van Nie, R.: The map position of the
rds

[189790]gene on the 17th chromosome of the mouse. Tissue Anti-
gens 13: 53–55,

[189791]1979.

[189792]

[189793]

[189794]20088.

[189795]Makela, T. P.; Kere, J.; Winqvist, R.; Alitalo, K.: Intrachro-
mosomal

[189796]rearrangements fusing L-myc and rlf in small-cell lung
cancer. Molec.

[189797]Cell. Biol. 11: 4015-4021, 1991.

[189798]

[189799]

[189800]20089.

[189801]Makela, T. P.; Saksela, K.; Evan, G.; Alitalo, K.: A fusion
protein

[189802]formed by L-myc and a novel gene in SCLC. EMBO J. 10:
1331-1335,

[189803]1991.

[189804]

[189805]

[189806]20090.

[189807]Brody, L. C.; Abel, K. J.; Castilla, L. H.; Couch, F. J.; McKin-
ley,

[189808]D. R.; Yin, G.-Y.; Ho, P. P.; Merajver, S.; Chan-

drasekharappa, S.

[189809]C.; Xu, J.; Cole, J. L.; Struewing, J. P.; Valdes, J. M.; Collins,

[189810]F. S.; Weber, B. L.: Construction of a transcription map
surrounding

[189811]the BRCA1 locus of human chromosome 17. Genomics 25:
238–247, 1995.

[189812]

[189813]

[189814]20091.

[189815]Hloch, P.; Schiedner, G.; Stahl, H.: Complete cDNA se-
quence of

[189816]the human p68 protein. Nucleic Acids Res. 18: 3045,
1990.

[189817]

[189818]

[189819]20092.

[189820]Iggo, R.; Gough, A.; Xu, W.; Lane, D. P.; Spurr, N. K.: Chro-
mosome

[189821]mapping of the human gene encoding the 68–kDa nuclear
antigen (p68)

[189822]by using the polymerase chain reaction. Proc. Nat. Acad.

Sci. 86:

[189823]6211–6214, 1989.

[189824]

[189825]

[189826]20093.

[189827]Iggo, R. D.; Xu, W.; Lane, D. P.; Gough, A.; Spurr, N. K.:

Assignment

[189828]of P68 to chromosome 17 using PCR. (Abstract) Cyto-
genet. Cell Genet. 51:

[189829]1017, 1989.

[189830]

[189831]

[189832]20094.

[189833]Acker, J.; Mattei, M.–G.; Wintzerith, M.; Roeckel, N.; De-
petris,

[189834]D.; Vigneron, M.; Keding, C.: Chromosomal localization
of human

[189835]RNA polymerase II subunit genes. Genomics 20: 496–499,
1994.

[189836]

[189837]

[189838]20095.

[189839]Bourquin, J.-P.; Stagljar, I.; Meier, P.; Moosmann, P.; Silke,

[189840]J.; Baechi, T.; Georgiev, O.; Schaffner, W.: A serine/arginine-rich

[189841]nuclear matrix cyclophilin interacts with the C-terminal domain of

[189842]RNA polymerase II. Nucleic Acids Res. 25: 2055–2061, 1997.

[189843]

[189844]

[189845]20096.

[189846]Cowburn, A. S.; Sladek, K.; Soja, J.; Adamek, L.; Nizankowska,

[189847]E.; Szczeklik, A.; Lam, B. K.; Penrose, J. F.; Austen, K. F.; Holgate,

[189848]S. T.; Sampson, A. P.: Overexpression of leukotriene C(4) synthase

[189849]in bronchial biopsies from patients with aspirin-intolerant asthma. J.

[189850]Clin. Invest. 101: 834–846, 1998.

[189851]

[189852]

[189853]20097.

[189854]Schulze, A.; Hansen, C.; Skakkebaek, N. E.; Brondum-Nielsen, K.;

[189855]Ledbetter, D. H.; Tommerup, N.: Exclusion of SNRPN as a major determinant

[189856]of Prader-Willi syndrome by a translocation breakpoint. Nature Genet. 12:

[189857]452-454, 1996.

[189858]

[189859]

[189860]20098.

[189861]Schweizer, J.; Zynger, D.; Francke, U.: In vivo nuclease hypersensitivity

[189862]studies reveal multiple sites of parental origin-dependent differential

[189863]chromatin conformation in the 150 kb SNRPN transcription unit. Hum.

[189864]Molec. Genet. 8: 555-566, 1999.

[189865]

[189866]

[189867]2009.

[189868]Shemer, R.; Hershko, A. Y.; Perk, J.; Mostoslavsky, R.;
Tsuberi,

[189869]B.; Cedar, H.; Buiting, K.; Razin, A.: The imprinting box of
the

[189870]Prader-Willi/Angelman syndrome domain. Nature Genet.
26: 440-443,

[189871]2000.

[189872]

[189873]

[189874]2010.

[189875]Sun, Y.; Nicholls, R. D.; Butler, M. G.; Saitoh, S.; Hainline,

[189876]B. E.; Palmer, C. G.: Breakage in the SNRPN locus in a bal-
anced 46,XY,t(15;19)

[189877]Prader-Willi syndrome patient. Hum. Molec. Genet. 5:
517-524, 1996.

[189878]

[189879]

[189880]20101.

[189881]Sutcliffe, J. S.; Nakao, M.; Christian, S.; Orstavik, K. H.;

Tommerup,

[189882]N.; Ledbetter, D. H.; Beaudet, A. L.: Deletions of a differentially

[189883]methylated CpG island at the SNRPN gene define a putative imprinting

[189884]control region. Nature Genet. 8: 52–58, 1994.

[189885]

[189886]

[189887]20102.

[189888]Wirth, J.; Back, E.; Huttenhofer, A.; Nothwang, H.–G.; Lich, C.;

[189889]Gross, S.; Menzel, C.; Schinzel, A.; Kioschis, P.; Tommerup, N.; Ropers,

[189890]H.–H.; Horsthemke, B.; Buiting, K.: A translocation breakpoint cluster

[189891]disrupts the newly defined 3–prime end of the SNURF–SNRPN transcription

[189892]unit on chromosome 15. Hum. Molec. Genet. 10: 201–210, 2001.

[189893]

[189894]

[189895]20103.

[189896]Guan, G.; Jiang, G.; Koch, R. L.; Shechter, I.: Molecular
cloning

[189897]and functional analysis of the promoter of the human
squalene synthase

[189898]gene. J. Biol. Chem. 270: 21958–21965, 1995.

[189899]

[189900]

[189901]20104.

[189902]Jiang, G.; McKenzie, T. L.; Conrad, D. G.; Shechter, I.:
Transcriptional

[189903]regulation by lovastatin and 25–hydroxycholesterol in
HepG2 cells

[189904]and molecular cloning and expression of the cDNA for the
human hepatic

[189905]squalene synthase. J. Biol. Chem. 268: 12818–12824,
1993.

[189906]

[189907]

[189908]20105.

[189909]Shechter, I.; Conrad, D. G.; Hart, I.; Berger, R. C.; McKenzie,
zie,

[189910]T. L.; Bleskan, J.; Patterson, D.: Localization of the squalene synthase

[189911]gene (FDFT1) to human chromosome 8p22–p23.1. Genomics 20: 116–118,

[189912]1994.

[189913]

[189914]

[189915]20106.

[189916]Stevanovic, M.; Zuffardi, O.; Collignon, J.; Lovell–Badge, R.;

[189917]Goodfellow, P.: The cDNA sequence and chromosomal location of the

[189918]human SOX2 gene. Mammalian Genome 5: 640–642,
1994.

[189919]

[189920]

[189921]20107.

[189922]Cricher, R.; Stitson, R. N. M.; Wade-Martins, R.; Easty, D. J.;

[189923]Farr, C. J.: Assignment of Sox4 to mouse chromosome 13 bands A3–A5

[189924]by fluorescence in situ hybridization; refinement of the human SOX4

[189925]location to 6p22.3 and of SOX20 to chromosome 17p12.3. Cytogenet.

[189926]Cell Genet. 81: 294–295, 1998.

[189927]

[189928]

[189929]20108.

[189930]Denny, P.; Swift, S.; Connor, F.; Ashworth, A.: An SRY-related

[189931]gene expressed during spermatogenesis in the mouse encodes a sequence-specific

[189932]DNA-binding protein. EMBO J. 11: 3705–3712, 1992.

[189933]

[189934]

[189935]20109.

[189936]Farr, C. J.; Easty, D. J.; Ragoussis, J.; Collignon, J.; Lovell-Badge,

[189937]R.; Goodfellow, P. N.: Characterization and mapping of the human

[189938]SOX4 gene. Mammalian Genome 4: 577–584, 1993.

[189939]

[189940]

[189941]20110.

[189942]Hansen, G. M.; Skapura, D.; Justice, M. J.: Genetic profile of

[189943]insertion mutations in mouse leukemias and lymphomas. Genome Res. 10:

[189944]237–243, 2000.

[189945]

[189946]

[189947]20111.

[189948]Li, J.; et al; et al: Leukaemia disease genes: large-scale cloning

[189949]and pathway predictions. Nature Genet. 23: 348–353, 1999.

[189950]

[189951]

[189952]20112.

[189953]Lund, A. H.; Turner, G.; Trubetskoy, A.; Verhoeven, E.;
Wientjens,

[189954]E.; Hulsman, D.; Russell, R.; DePinho, R. A.; Lenz, J.; van
Lohuizen,

[189955]M.: Genome-wide retroviral insertional tagging of genes
involved

[189956]in cancer in Cdkn2a-deficient mice. Nature Genet. 32:
160–165, 2002.

[189957]

[189958]

[189959]20113.

[189960]Suzuki, T.; Shen, H.; Akagi, K.; Morse, H. C., III; Malley, J.

[189961]D.; Naiman, D. Q.; Jenkins, N. A.; Copeland, N. G.: New
genes involved

[189962]in cancer identified by retroviral tagging. Nature Genet.
32: 166–174,

[189963]2002. Note: Erratum: Nature Genet. 32: 331 only, 2002.

[189964]

[189965]

[189966]20114.

[189967]van de Wetering, M.; Oosterwegel, M.; van Norren, K.;
Clevers,

[189968]H.: Sox-4, an Sry-like HMG box protein, is a transcrip-
tional activator

[189969]in lymphocytes. EMBO J. 12: 3847-3854, 1993.

[189970]

[189971]

[189972]20115.

[189973]Shoyab, M.; McDonald, V. L.; Bradley, J. G.; Todaro, G. J.:
Amphiregulin:

[189974]a bifunctional growth-modulating glycoprotein produced
by the phorbol

[189975]12-myristate 13-acetate-treated human breast adenocar-
cinoma cell line

[189976]MCF-7. Proc. Nat. Acad. Sci. 85: 6528-6532, 1988.

[189977]

[189978]

[189979]20116.

[189980]Shoyab, M.; Plowman, G. D.; McDonald, V. L.; Bradley, J.

G.; Todaro,

[189981]G. J.: Structure and function of human amphiregulin: a
member of

[189982]the epidermal growth factor family. Science 243:
1074–1076, 1989.

[189983]

[189984]

[189985]20117.

[189986]Brock, A.; Mortensen, P. B.; Mortensen, B. B.; Roge, H. R.:

Familial

[189987]occurrence of diminished pancreatic amylase in serum--a
'silent' Amy-2

[189988]allelic variant? Clin. Chem. 34: 1516–1517, 1988.

[189989]

[189990]

[189991]20118.

[189992]Carfagna, M.; Gaudio, L.; Patricolo, M. R.; Spadacenti, F.:

Pancreatic

[189993]amylase polymorphism: another example of a distinctive
gene frequency

[189994]among Sardinians. Hum. Hered. 26: 59–65, 1976.

[189995]

[189996]

[189997]20119.

[189998]Donahue, R. P.; Bias, W. B.; Renwick, J. H.; McKusick, V. A.:

[189999]Probable assignment of the Duffy blood group locus to
chromosome 1

[190000]in man. Proc. Nat. Acad. Sci. 61: 949–955, 1968.

[190001]

[190002]

[190003]20120.

[190004]Groot, P. C.; Bleeker, M. J.; Pronk, J. C.; Arwert, F.; Mager,

[190005]W. H.; Planta, R. J.; Eriksson, A. W.; Frants, R. R.: Human
pancreatic

[190006]amylase is encoded by two different genes. Nucleic Acids
Res. 16:

[190007]4724 only, 1988.

[190008]

[190009]

[190010]20121.

[190011]Groot, P. C.; Bleeker, M. J.; Pronk, J. C.; Arwert, F.; Mager,

[190012]W. H.; Planta, R. J.; Eriksson, A. W.; Frants, R. R.: The human alpha-amylase

[190013]multigene family consists of haplotypes with variable numbers of genes. Genomics 5:

[190014]29–42, 1989.

[190015]

[190016]

[190017]20122.

[190018]Groot, P. C.; Mager, W. H.; Frants, R. R.: Interpretation of polymorphic

[190019]DNA patterns in the human alpha-amylase multigene family. Genomics 10:

[190020]779–785, 1991.

[190021]

[190022]

[190023]20123.

[190024]Groot, P. C.; Mager, W. H.; Henriquez, N. V.; Pronk, J. C.;
Arwert,

[190025]F.; Planta, R. J.; Eriksson, A. W.; Frants, R. R.: Evolution of
the

[190026]human alpha-amylase multigene family through unequal,
homologous,

[190027]and inter- and intrachromosomal crossovers. Genomics 8:
97-105,

[190028]1990.

[190029]

[190030]

[190031]20124.

[190032]Hill, C. J.; Rowe, S. I.; Lovrien, E. W.: Probable genetic link-
age

[190033]between human serum amylase (AMY-2) and Duffy blood
groups. Nature 235:

[190034]162-163, 1972.

[190035]

[190036]

[190037]20125.

[190038]Hjorth, J. P.; Lusi, A. J.; Nielsen, J. T.: Multiple structural
[190039]genes for mouse amylase. Biochem. Genet. 18: 281–302,
1980.

[190040]

[190041]

[190042]20126.

[190043]Jorgensen, H. R.; Kristensen, B.; Mortensen, P. B.: Familial
[190044]incidence of reduced activity of pancreas correlated with
amylase–isoenzyme
[190045]in the serum. Ugeskr. Laeger 146: 657–659, 1984.

[190046]

[190047]

[190048]20127.

[190049]Kamaryt, J.; Adamek, R.; Vrba, M.: Possible linkage be-
tween uncoiler

[190050]chromosome Un 1 and amylase polymorphism Amy 2 loci.
Humangenetik 11:

[190051]213–220, 1971.

[190052]

[190053]

[190054]20128.

[190055]Merritt, A. D.; Lovrien, E. W.; Rivas, M. L.; Conneally, P. M.

[190056]: Human amylase loci: genetic linkage with the Duffy
blood group locus

[190057]and assignment to linkage group I. Am. J. Hum. Genet. 25:
523–538,

[190058]1973.

[190059]

[190060]

[190061]20129.

[190062]Merritt, A. D.; Rivas, M. L.; Bixler, D.; Newell, R.: Salivary

[190063]and pancreatic amylase: electrophoretic characterizations
and genetic

[190064]studies. Am. J. Hum. Genet. 25: 510–522, 1973.

[190065]

[190066]

[190067]20130.

[190068]Merritt, A. D.; Rivas, M. L.; Ward, J. C.: Evidence for close
[190069]linkage of human amylase loci. Nature N.B. 239: 243–244,
1972.

[190070]

[190071]

[190072]20131.

[190073]Pronk, J. C.; Frants, R. R.; Jansen, W.; Eriksson, A. W.;
Tonino,

[190074]G. J. M.: Evidence for duplication of the human salivary
amylase

[190075]gene. Hum. Genet. 60: 32–35, 1982.

[190076]

[190077]

[190078]20132.

[190079]Rosenblum, B. B.; Merritt, A. D.: Human pancreatic alpha-
amylase:

[190080]phenotypic codominance and new electrophoretic vari-
ants. Am. J. Hum.

[190081]Genet. 30: 434–441, 1978.

[190082]

[190083]

[190084]20133.

[190085]Sjolund, K.; Haggmark, A.; Ihse, I.; Skude, G.; Karnstrom, U.;

[190086]Wikander, M.: Selective deficiency of pancreatic amylase.

Gut 32:

[190087]546–548, 1991.

[190088]

[190089]

[190090]20134.

[190091]Spence, M. A.; Sparkes, R. S.; Heckenlively, J. R.; Pearlman,

[190092]J. T.; Zedalis, D.; Sparkes, M.; Crist, M.; Tideman, S.: Prob-
able

[190093]genetic linkage between autosomal dominant retinitis pig-
mentosa (RP)

[190094]and amylase (AMY-2): evidence of an RP locus on chro-
mosome 1. Am.

[190095]J. Hum. Genet. 29: 397–404, 1977.

[190096]

[190097]

[190098]20135.

[190099]Tricoli, J. V.; Shows, T. B.: Regional assignment of human
amylase

[190100](AMY) to p22–p21 of chromosome 1. Somat. Cell Molec.
Genet. 10:

[190101]205–210, 1984.

[190102]

[190103]

[190104]20136.

[190105]Young, R. A.; Hagenbuchle, O.; Schibler, U.: A single
mouse alpha–amylase

[190106]gene specifies two different tissue–specific mRNAs. Cell
23: 451–458,

[190107]1981.

[190108]

[190109]

[190110]20137.

[190111]de Soyza, K.: Polymorphism of human salivary amylase: a
preliminary

[190112]communication. Hum. Genet. 45: 189–192, 1978.

[190113]

[190114]

[190115]20138.

[190116]Dracopoli, N. C.; Meisler, M. H.: Mapping the human amy-
lase gene

[190117]cluster on the proximal short arm of chromosome 1 using
a highly informative

[190118](CA)_n repeat. Genomics 7: 97–102, 1990.

[190119]

[190120]

[190121]20139.

[190122]Gumucio, D. L.; Wiebauer, K.; Caldwell, R. M.; Samuelson,
L. C.;

[190123]Meisler, M. H.: Concerted evolution of human amylase
genes. Molec.

[190124]Cell. Biol. 8: 1197–1205, 1988.

[190125]

[190126]

[190127]20140.

[190128]Ishizaki, K.; Noda, A.; Ikenaga, M.; Ida, K.; Omoto, K.;
Nakamura,

[190129]Y.; Matsubara, K.: Restriction fragment length polymor-
phism detected

[190130]by human salivary amylase cDNA. Hum. Genet. 71:
261–262, 1985.

[190131]

[190132]

[190133]20141.

[190134]Kamaryt, J.; Laxova, R.: Amylase heterogeneity: some ge-
netic and

[190135]clinical aspects. Humangenetik 1: 579–586, 1965.

[190136]

[190137]

[190138]20142.

[190139]Kamaryt, J.; Laxova, R.: Amylase heterogeneity variants in
man. Humangenetik 3:

[190140]41–45, 1966.

[190141]

[190142]

[190143]20143.

[190144]McGeachin, R. L.: Multiple molecular forms of amylase.
Ann. N.Y.

[190145]Acad. Sci. 151: 208–212, 1968.

[190146]

[190147]

[190148]20144.

[190149]Muenke, M.; Lindgren, V.; de Martinville, B.; Francke, U.:
Comparative

[190150]analysis of mouse–human hybrids with rearranged chromosomes 1 by in

[190151]situ hybridization and Southern blotting: high resolution
mapping

[190152]of NRAS, NGFB, and AMY on chromosome 1. Somat. Cell
Molec. Genet. 10:

[190153]589–599, 1984.

[190154]

[190155]

[190156]20145.

[190157]Nishide, T.; Nakamura, Y.; Emi, M.; Yamamoto, T.; Ogawa,

M.; Mori,

[190158]T.; Matsubara, K.: Primary structure of human salivary al-
pha-amylase

[190159]gene. Gene 41: 299–304, 1986.

[190160]

[190161]

[190162]20146.

[190163]Knoll, J. H. M.; Nicholls, R. D.; Magenis, R. E.; Glatt, K.;
Graham,

[190164]J. M., Jr.; Kaplan, L.; Lalande, M.: Angelman syndrome:
three molecular

[190165]classes identified with chromosome 15q11q13-specific
DNA markers. Am.

[190166]J. Hum. Genet. 47: 149–155, 1990.

[190167]

[190168]

[190169]20147.

[190170]Ono, H.; Kuno, Y.; Tanaka, H.; Yamashina, M.; Tsuyoshi,
T.; Kondo,

[190171]N.; Orii, T.: A case of paroxysmal nocturnal hemoglobin–

uria without

[190172]deficiency of decay-accelerating factor on erythrocytes.

Blood 75:

[190173]1746–1747, 1990.

[190174]

[190175]

[190176]20148.

[190177]Petranka, J. G.; Fleenor, D. E.; Sykes, K.; Kaufman, R. E.;
Rosse,

[190178]W. F.: Structure of the CD59–encoding gene: further evidence of a

[190179]relationship to murine lymphocyte antigen Ly–6 protein.

Proc. Nat.

[190180]Acad. Sci. 89: 7876–7879, 1992.

[190181]

[190182]

[190183]20149.

[190184]Rosse, W. F.: Personal Communication. Durham, N. C.

6/3/1993.

[190185]

[190186]

[190187]20150.

[190188]Rosse, W. F.; Parker, C. J.: Paroxysmal nocturnal
hemoglobinuria. Clin.

[190189]Haemat. 14: 105–125, 1985.

[190190]

[190191]

[190192]20151.

[190193]Rother, R. P.; Rollins, S. A.; Mennone, J.; Chodera, A.; Fi-
del,

[190194]S. A.; Bessler, M.; Hillmen, P.; Squinto, S. P.: Expression of
recombinant

[190195]transmembrane CD59 in paroxysmal nocturnal
hemoglobinuria B cells

[190196]confers resistance to human complement. Blood 84:
2604–2611, 1994.

[190197]

[190198]

[190199]20152.

[190200]Tone, M.; Walsh, L. A.; Waldmann, H.: Gene structure of

human

[190201]CD59 and demonstration that discrete mRNAs are generated by alternative

[190202]polyadenylation. J. Molec. Biol. 227: 971–976, 1992.

[190203]

[190204]

[190205]20153.

[190206]Walsh, L. A.; Tone, M.; Thiru, S.; Waldmann, H.: The CD59 antigen--a

[190207]multifunctional molecule. Tissue Antigens 40: 213–220, 1992.

[190208]

[190209]

[190210]20154.

[190211]Yamashina, M.; Ueda, E.; Kinoshita, T.; Takami, T.; Ojima, A.;

[190212]Ono, H.; Tanaka, H.; Kondo, N.; Orii, T.; Okada, N.; Okada, H.; Inoue,

[190213]K.; Kitani, T.: Inherited complete deficiency of 20–kilodalton homologous

[190214]restriction factor (CD59) as a cause of paroxysmal noctur–

nal hemoglobinuria. New

[190215]Eng. J. Med. 323: 1184–1189, 1990.

[190216]

[190217]

[190218]20155.

[190219]Kumanogoh, A.; Watanabe, C.; Lee, I.; Wang, X.; Shi, W.;
Araki,

[190220]H.; Hirata, H.; Iwahori, K.; Uchida, J.; Yasui, T.; Matsumoto,
M.;

[190221]Yoshida, K.; Yakura, H.; Pan, C.; Parnes, J. R.; Kikutani, H.:
Identification

[190222]of CD72 as a lymphocyte receptor for the class IV
semaphorin CD100:

[190223]a novel mechanism for regulating B cell signaling. Immu-
nity 13:

[190224]621–631, 2000.

[190225]

[190226]

[190227]20156.

[190228]Von Hoegen, I.; Hsieh, C.–L.; Scharing, R.; Francke, U.;

Parnes,

[190229]J. R.: Identity of human Lyb-2 and CD72 and localization
of the gene

[190230]to chromosome 9. Europ. J. Immun. 21: 1425-1431,
1991.

[190231]

[190232]

[190233]20157.

[190234]Cambiaggi, C.; Scupoli, M. T.; Cestari, T.; Gerosa, F.;
Carra,

[190235]G.; Tridente, G.; Accolla, R. S.: Constitutive expression of
CD69

[190236]in interspecies T-cell hybrids and locus assignment to hu-
man chromosome

[190237]12. Immunogenetics 36: 117-120, 1992.

[190238]

[190239]

[190240]20158.

[190241]Lopez-Cabrera, M.; Santis, A. G.; Fernandez-Ruiz, E.;
Blacher,

[190242]R.; Esch, F.; Sanchez-Mateos, P.; Sanchez-Madrid, F.:

Molecular cloning,
[190243]expression, and chromosomal localization of the human
earliest lymphocyte
[190244]activation antigen AIM/CD69, a new member of the C-
type animal lectin
[190245]superfamily of signal-transmitting receptors. J. Exp. Med.
178:
[190246]537-547, 1993.

[190247]

[190248]

[190249]20159.

[190250]Chandra, T.; Stackhouse, R.; Kidd, V. J.; Robson, K. J. H.;
Woo,
[190251]S. L. C.: Sequence homology between human alpha-
1-antichymotrypsin,
[190252]alpha-1-antitrypsin, and antithrombin III. Biochemistry
22: 5055-5061,
[190253]1983.

[190254]

[190255]

[190256]20160.

[190257]Eriksson, S.; Lindmark, B.; Lilia, H.: Familial alpha-
1-antichymotrypsin

[190258]deficiency. Acta Med. Scand. 220: 447-453, 1986.

[190259]

[190260]

[190261]20161.

[190262]Gilfix, B. M.; Briones, L.: Absence of the A1252G mutation
in

[190263]alpha 1-antichymotrypsin in a North American population
suffering

[190264]from dementia. J. Cereb. Blood Flow Metab. 17: 233-235,
1997.

[190265]

[190266]

[190267]20162.

[190268]Haines, J. L.; Pritchard, M. L.; Saunders, A. M.; Schildkraut,

[190269]J. M.; Growdon, J. H.; Gaskell, P. C.; Farrer, L. A.; Auer-
bach, S.

[190270]A.; Gusella, J. F.; Locke, P. A.; Rosi, B. L.; Yamaoka, L.;
Small,

[190271]G. W.; Conneally, P. M.; Roses, A. D.; Pericak-Vance, M. A.:

No genetic

[190272]effect of alpha-1-antichymotrypsin in Alzheimer disease.

Genomics 33:

[190273]53-56, 1996.

[190274]

[190275]

[190276]20163.

[190277]Haines, J. L.; Scott, W. K.; Pericak-Vance, M. A.: Reply to

'Genetic

[190278]effect of alpha-1-antichymotrypsin on the risk of

Alzheimer disease.'

[190279](Letter) Genomics 40: 384-385, 1997.

[190280]

[190281]

[190282]20164.

[190283]Kamboh, M. I.; Aston, C. E.; Ferrell, R. E.; Dekosky, S. T.:

Genetic

[190284]effect of alpha-1-antichymotrypsin on the risk of

Alzheimer disease.

[190285](Letter) Genomics 41: 382–385, 1997.

[190286]

[190287]

[190288]20165.

[190289]Kamboh, M. I.; Sanghera, D. K.; Ferrell, R. E.; DeKosky, S.
T.

[190290]: APOE*4-associated Alzheimer's disease risk is modified
by alpha-1-antichymotrypsin

[190291]polymorphism. Nature Genet. 10: 486–488, 1995.

[190292]

[190293]

[190294]20166.

[190295]Kelsey, G. D.; Abeliovich, D.; McMahon, C. J.; Whitehouse,
D.;

[190296]Corney, G.; Povey, S.; Hopkinson, D. A.; Wolfe, J.; Mieli-
Vergani,

[190297]G.; Mowat, A. P.: Cloning of the human alpha-1 antichy-
motrypsin gene

[190298]and genetic analysis of the gene in relation to alpha-1 an-
titrypsin

[190299]deficiency. J. Med. Genet. 25: 361–368, 1988.

[190300]

[190301]

[190302]20167.

[190303]Morgan, K.; Licastro, F.; Tilley, L.; Ritchie, A.; Morgan, L.;

[190304]Pedrini, S.; Kalsheker, N.: Polymorphism in the alpha-
1-antichymotrypsin

[190305](ACT) gene promoter: effect on expression in transfected
glial and

[190306]liver cell lines and plasma ACT concentrations. Hum.
Genet. 109:

[190307]303–310, 2001.

[190308]

[190309]

[190310]20168.

[190311]Morgan, K.; Morgan, L.; Carpenter, K.; Lowe, J.; Lam, L.;
Cave,

[190312]S.; Xuereb, J.; Wischik, C.; Harrington, C.; Kalsheker, N. A.:
Microsatellite

[190313]polymorphism of the alpha-1-antichymotrypsin gene lo-

cus associated

[190314]with sporadic Alzheimer's disease. Hum. Genet. 99:
27–31, 1997.

[190315]

[190316]

[190317]20169.

[190318]Munoz, E.; Obach, V.; Oliva, R.; Marti, M. J.; Ezquerra, M.;
Pastor,

[190319]P.; Ballesta, F.; Tolosa, E.: Alpha–1–antichymotrypsin gene
polymorphism

[190320]and susceptibility to Parkinson's disease. Neurology 52:
297–301,

[190321]1999.

[190322]

[190323]

[190324]20170.

[190325]Poller, W.; Faber, J.–P.; Scholz, S.; Weidinger, S.;
Bartholome,

[190326]K.; Olek, K.; Eriksson, S.: Mis–sense mutation of alpha–
1–antichymotrypsin

[190327]gene associated with chronic lung disease. (Letter) Lancet

339:

[190328]1538, 1992.

[190329]

[190330]

[190331]20171.

[190332]Poller, W.; Faber, J.-P.; Weidinger, S.; Tief, K.; Scholz, S.;

[190333]Fischer, M.; Olek, K.; Kirchgesser, M.; Heidtmann, H.-H.: A
leucine-to-proline

[190334]substitution causes a defective alpha-1-antichymotrypsin
allele associated

[190335]with familial obstructive lung disease. Genomics 17:
740-743, 1993.

[190336]

[190337]

[190338]20172.

[190339]Rabin, M.; Watson, M.; Breg, W. R.; Kidd, V.; Woo, S. L. C.;
Ruddle,

[190340]F. H.: Human alpha-1-antichymotrypsin and alpha-
1-antitrypsin (PI)

[190341]genes map to the same region on chromosome 14.

(Abstract) Cytogenet.

[190342]Cell Genet. 40: 728, 1985.

[190343]

[190344]

[190345]20173.

[190346]Rabin, M.; Watson, M.; Kidd, V.; Woo, S. L. C.; Breg, W. R.;
Ruddle,

[190347]F. H.: Regional location of alpha-1-antichymotrypsin and
alpha-1-antitrypsin

[190348]genes on human chromosome 14. Somat. Cell Molec.
Genet. 12: 209-214,

[190349]1986.

[190350]

[190351]

[190352]20174.

[190353]Han, P.; Fletcher, C. F.; Copeland, N. G.; Jenkins, N. A.;
Yaremko,

[190354]L. M.; Michaeli, T.: Assignment of the mouse Pde7A gene
to the proximal

[190355]region of chromosome 3 and of the human PDE7A gene to

chromosome 8q13. Genomics 48:

[190356]275–276, 1998.

[190357]

[190358]

[190359]20175.

[190360]Michaeli, T.; Bloom, T. J.; Martins, T.; Loughney, K.; Ferguson,
son,

[190361]K.; Riggs, M.; Rodgers, L.; Beavo, J. A.; Wigler, M.: Isolation
and

[190362]characterization of a previously undetected human cAMP
phosphodiesterase

[190363]by complementation of cAMP phosphodiesterase–deficient
Saccharomyces

[190364]cerevisiae. J. Biol. Chem. 268: 12925–12932, 1993.

[190365]

[190366]

[190367]20176.

[190368]Milatovich, A.; Bolger, G.; Michaeli, T.; Francke, U.: Chro-
mosome

[190369]localizations of genes for five cAMP–specific phosphodi-

esterases in

[190370]man and mouse. Somat. Cell Molec. Genet. 20: 75–86,
1994.

[190371]

[190372]

[190373]20177.

[190374]Loughney, K.; Martins, T. J.; Harris, E. A. S.; Sadhu, K.;
Hicks,

[190375]J. B.; Sonnenburg, W. K.; Beavo, J. A.; Ferguson, K.: Isola-
tion and

[190376]characterization of cDNAs corresponding to two human
calcium, calmodulin-regulated,

[190377]3-prime,5-prime-cyclic nucleotide phosphodiesterases. J.
Biol. Chem. 271:

[190378]796–806, 1996.

[190379]

[190380]

[190381]20178.

[190382]Wilson, D. E.; McKenna, L.: Assignment of the human gene
for phosphodiesterase

[190383]1A to chromosome 4. (Abstract) Am. J. Hum. Genet. 43:

A162 only,

[190384]1988.

[190385]

[190386]

[190387]20179.

[190388]Ritter, J. K.; Chen, F.; Sheen, Y. Y.; Tran, H. M.; Kimura, S.;

[190389]Yeatman, M. T.; Owens, I. S.: A novel complex locus UGT1
encodes

[190390]human bilirubin, phenol, and other UDP-
glucuronosyltransferase isozymes

[190391]with identical carboxyl termini. J. Biol. Chem. 267:
3257-3261,

[190392]1992.

[190393]

[190394]

[190395]20180.

[190396]Tukey, R. H.; Strassburg, C. P.: Human UDP-
glucuronosyltransferases:

[190397]metabolism, expression, and disease. Annu. Rev. Pharm.
Toxicol. 40:

[190398]581–616, 2000.

[190399]

[190400]

[190401]20181.

[190402]James, P. F.; Grupp, I. L.; Grupp, G.; Woo, A. L.; Askew, G. R.;

[190403]Croyle, M. L.; Walsh, R. A.; Lingrel, J. B.: Identification of a

[190404]specific role for the Na,K–ATPase alpha–2 isoform as a
regulator of

[190405]calcium in the heart. Molec. Cell 3: 555–563, 1999.

[190406]

[190407]

[190408]20182.

[190409]Kawakami, K.; Ohta, T.; Nojima, H.; Nagano, K.: Primary
structure

[190410]of the alpha–subunit of human Na,K–ATPase deduced
from cDNA sequence. J.

[190411]Biochem. 100: 389–397, 1986.

[190412]

[190413]

[190414]20183.

[190415]Kent, R. B.; Fallows, D. A.; Geissler, E.; Glaser, T.; Emanuel,

[190416]J. R.; Lalley, P. A.; Levenson, R.; Housman, D. E.: Genes
encoding

[190417]alpha and beta subunits of Na,K-ATPase are located on
three different

[190418]chromosomes in the mouse. Proc. Nat. Acad. Sci. 84:
5369-5373, 1987.

[190419]

[190420]

[190421]20184.

[190422]Ruiz-Opazo, N.; Barany, F.; Hirayama, K.; Herrera, V. L. M.:

[190423]Confirmation of mutant alpha-1 Na,K-ATPase gene and
transcript in

[190424]Dahl salt-sensitive/JR rats. Hypertension 24: 260-270,
1994.

[190425]

[190426]

[190427]20185.

[190428]Ovchinnikov, Y. A.; Monastyrskaya, G. S.; Broude, N. E.;
Allikmets,

[190429]R. L.; Ushkaryov, Y. A.; Melkov, A. M.; Smirnov, Y. V.;
Malyshev,

[190430]I. V.; Dulubova, I. E.; Petrukhin, K. E.; Gryshin, A. V.;
Sverdlov,

[190431]V. E.; Kiyatkin, N. I.; Kostina, M. B.; Modyanov, N. N.;
Sverdlov,

[190432]E. D.: The family of human Na⁺,K⁺-ATPase genes: a par-
tial nucleotide

[190433]sequence related to the alpha-subunit. FEBS Lett. 213:
73-80, 1987.

[190434]

[190435]

[190436]20186.

[190437]Shull, M. M.; Lingrel, J. B.: Multiple genes encode the hu-
man

[190438]Na⁺,K⁺-ATPase catalytic subunit. Proc. Nat. Acad. Sci. 84:
4039-4043,

[190439]1987.

[190440]

[190441]

[190442]20187.

[190443]Sverdlov, E. D.; Broude, N. E.; Sverdlov, V. E.;

Monastyrskaya,

[190444]G. S.; Grishin, A. V.; Petrukhin, K. E.; Akopyanz, N. S.;

Modyanov,

[190445]N. N.; Ovchinnikov, Y. A.: Family of Na⁺,K⁺-ATPase

genes: intra-individual

[190446]tissue-specific restriction fragment length polymorphism.

FEBS Lett. 221:

[190447]129-133, 1987.

[190448]

[190449]

[190450]20188.

[190451]Yang-Feng, T. L.; Schneider, J. W.; Lindgren, V.; Shull, M.

M.;

[190452]Benz, E. J., Jr.; Lingrel, J. B.; Francke, U.: Chromosomal lo-

calization

[190453]of human Na⁺,K⁺-ATPase alpha- and beta-subunit

genes. Genomics 2:

[190454]128-138, 1988.

[190455]

[190456]

[190457]20189.

[190458]Gloor, S.; Antonicek, H.; Sweadner, K. J.; Pagliusi, S.;
Frank,

[190459]R.; Moos, M.; Schachner, M.: The adhesion molecule on
glia (AMOG)

[190460]is a homologue of the beta subunit of the Na,K-ATPase. J.
Cell Biol. 110:

[190461]165-174, 1990.

[190462]

[190463]

[190464]20190.

[190465]Hsieh, C.-L.; Cheng-Deutsch, A.; Gloor, S.; Schachner, M.;
Francke,

[190466]U.: Assignment of Amog (adhesion molecule on glia) gene
to mouse

[190467]chromosome 11 near Zfp-3 and Asgr-1,2 and to human
chromosome 17. Somat.

[190468]Cell Molec. Genet. 16: 401-405, 1990.

[190469]

[190470]

[190471]20191.

[190472]Malo, D.; Schurr, E.; Levenson, R.; Gros, P.: Assignment of
Na,K-ATPase

[190473]beta-(2)-subunit gene (Atpb-2) to mouse chromosome

11. Genomics 6:

[190474]697-699, 1990.

[190475]

[190476]

[190477]20192.

[190478]Martin-Vasallo, P.; Dackowski, P.; Emanuel, J. R.; Leven-
son, R.

[190479]: Identification of a putative isoform of the Na,K-ATPase
beta subunit:

[190480]primary structure and tissue-specific expression. J. Biol.
Chem. 164:

[190481]4613-4618, 1989.

[190482]

[190483]

[190484]20193.

[190485]Pagliusi, S.; Antonicek, H.; Gloor, S.; Frank, R.; Moos, M.;
Schachner,

[190486]M.: Identification of a cDNA clone specific for the neural
adhesion

[190487]molecule AMOG. J. Neurosci. Res. 22: 113–119, 1989.

[190488]

[190489]

[190490]20194.

[190491]Gallango, M. L.; Muller, A.; Suinaga, R.: Biochemical char-
acterization

[190492]of a red cell UMP kinase variant found in the Warao Indi-
ans of Venezuela. Biochem.

[190493]Genet. 16: 1085–1093, 1978.

[190494]

[190495]

[190496]20195.

[190497]Gallango, M. L.; Suinaga, R.: Uridine monophosphate ki-
nase polymorphism

[190498]in two Venezuelan populations. Am. J. Hum. Genet. 30:

215–218, 1978.

[190499]

[190500]

[190501]20196.

[190502]Giblett, E. R.; Anderson, J. E.; Chen, S.–H.; Teng, Y.–S.; Cohen,
hen,

[190503]F.: Uridine monophosphate kinase: a new genetic polymorphism with
morphism with

[190504]possible clinical implications. Am. J. Hum. Genet. 26:
627–635,

[190505]1974.

[190506]

[190507]

[190508]20197.

[190509]Giblett, E. R.; Anderson, J. E.; Lewis, M.; Kaita, H.: A new
polymorphic

[190510]enzyme, uridine monophosphate kinase: gene frequencies
and a linkage

[190511]analysis. Birth Defects Orig. Art. Ser. 11(3): 159–161,
1975. Note:

[190512]Alternate: Cytogenet. Cell Genet. 14: 329–331, 1975.

[190513]

[190514]

[190515]20198.

[190516]Medrano, L.; Green, H.: A uridine kinase deficient mutant
of 3T3

[190517]and a selective method for cells containing the enzyme.

Cell 1:

[190518]23–26, 1974.

[190519]

[190520]

[190521]20199.

[190522]Petersen, G. M.; Silimperi, D. R.; Scott, E. M.; Hall, D. B.;
Rotter,

[190523]J. I.; Ward, J. I.: Uridine monophosphate kinase 3: a ge-
netic marker

[190524]for susceptibility to Haemophilus influenzae type B dis-
ease. Lancet II:

[190525]417–418, 1985.

[190526]

[190527]

[190528]20200.

[190529]Ranzani, G.; Bertolotti, E.; Santachiara-Benerecetti, A. S.:

The

[190530]polymorphism of the red cell uridine monophosphate kinase in two samples

[190531]of the Italian population. Hum. Hered. 27: 332–335, 1977.

[190532]

[190533]

[190534]20201.

[190535]Ruddle, F. H.; Creagan, R. P.: Parasexual approaches to the genetics

[190536]of man. Ann. Rev. Genet. 9: 407–486, 1975.

[190537]

[190538]

[190539]20202.

[190540]Satlin, A.; Kucherlapati, R. S.; Ruddle, F. H.: Assignment of

[190541]the gene for human uridine monophosphate kinase to chromosome 1 using

[190542]somatic cell hybrid clone panels. Cytogenet. Cell Genet.

15: 146–152,

[190543]1975.

[190544]

[190545]

[190546]20203.

[190547]Anderson, J. A.; Teng, Y.–S.; Giblett, E. R.: Stains for six
enzymes

[190548]potentially applicable to chromosomal assignment by cell
hybridization. Cytogenet.

[190549]Cell Genet. 14: 295–299, 1975.

[190550]

[190551]

[190552]20204.

[190553]Hoglund, L.; Reichard, P. :J. Biol. Chem. 265: 6589–6595,
1990.

[190554]

[190555]

[190556]20205.

[190557]Paglia, D. E.; Valentine, W. N.; Brockway, R. A.: Identifica–

tion

[190558]of thymidine nucleotidase and deoxyribonucleotidase activities among

[190559]normal isozymes of 5-prime-nucleotidase in human erythrocytes. Proc.

[190560]Nat. Acad. Sci. 81: 588–592, 1984.

[190561]

[190562]

[190563]20206.

[190564]Rampazzo, C.; Gallinaro, L.; Milanesi, E.; Frigimelica, E.; Reichard,

[190565]P.; Bianchi, V.: A deoxyribonucleotidase in mitochondria: involvement

[190566]in regulation of dNTP pools and possible link to genetic disease. Proc.

[190567]Nat. Acad. Sci. 97: 8239–8244, 2000.

[190568]

[190569]

[190570]20207.

[190571]Rampazzo, C.; Johansson, M.; Gallinaro, L.; Ferraro, P.;

Hellman,

[190572]U.; Karlsson, A.; Reichard, P.; Bianchi, V.: Mammalian
5-prime(3-prime)-deoxyri

[190573]bonucleotidase, cDNA cloning, and overexpression of the
enzyme in

[190574]Escherichia coli and mammalian cells. J. Biol. Chem. 275:
5409-5415,

[190575]2000.

[190576]

[190577]

[190578]20208.

[190579]Swallow, D. M.; Turner, V. S.; Hopkinson, D. A.: Isozymes
of rodent

[190580]5-prime-nucleotidase: evidence for two independent
structural loci

[190581]UMPH-1 and UMPH-2. Ann. Hum. Genet. 47: 9-17, 1983.

[190582]

[190583]

[190584]20209.

[190585]Tjernshaugen, H.; Fritzson, P. :Biochem. J. 154: 77-80,
1976.

[190586]

[190587]

[190588]20210.

[190589]Wilson, D. E.; Swallow, D. M.; Povey, S.: Assignment of the
human

[190590]gene for uridine 5-prime-monophosphate phosphohydro-
lase (UMPH2) to

[190591]the long arm of chromosome 17. Ann. Hum. Genet. 50:
223-227, 1986.

[190592]

[190593]

[190594]20211.

[190595]Wilson, D. E.; Woodard, D.; Sandler, A.; Erickson, J.; Gur-
ney,

[190596]A.: Provisional assignment of the gene for uridine
monophosphatase-2

[190597](Umph-2) to mouse chromosome 11. Biochem. Genet. 25:
1-6, 1987.

[190598]

[190599]

[190600]20212.

[190601]Xiong, J.-P.; Stehle, T.; Diefenbach, B.; Zhang, R.; Dunker, R.;

[190602]Scott, D. L.; Joachimiak, A.; Goodman, S. L.; Arnaout, M. A.: Crystal

[190603]structure of the extracellular segment of integrin alpha-V-beta-3. Science 294:

[190604]339-345, 2001.

[190605]

[190606]

[190607]20213.

[190608]Xiong, J.-P.; Stehle, T.; Zhang, R.; Joachimiak, A.; Frech, M.;

[190609]Goodman, S. L.; Arnaout, M. A.: Crystal structure of the extracellular

[190610]segment of integrin alpha-V-beta-3 in complex with an Arg-Gly-Asp

[190611]ligand. Science 296: 151-155, 2002.

[190612]

[190613]

[190614]20214.

[190615]Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly, J.;

[190616]Bowen-Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of the amphiregulin

[190617]and the platelet-growth factor receptor alpha genes to the proximal

[190618]long arm of chromosome 4. (Abstract) Cytogenet. Cell Genet. 51:

[190619]990, 1989.

[190620]

[190621]

[190622]20215.

[190623]Gronwald, R. G. K.; Adler, D. A.; Kelly, J. D.; Disteche, C. M.;

[190624]Bowen-Pope, D. F.: The human PDGF receptor alpha-subunit gene maps

[190625]to chromosome 4 in close proximity to c-kit. Hum. Genet. 85: 383-385,

[190626]1990.

[190627]

[190628]

[190629]20216.

[190630]Hol, F. A.; Geurds, M. P. A.; Chatkupt, S.; Shugart, Y. Y.;

Balling,

[190631]R.; Schrandt-Stumpel, C. T. R. M.; Johnson, W. G.; Hamel,

B. C. J.;

[190632]Mariman, E. C. M.: PAX genes and human neural tube defects: an amino

[190633]acid substitution in PAX1 in a patient with spina bifida. J. Med.

[190634]Genet. 8: 655–660, 1996.

[190635]

[190636]

[190637]20217.

[190638]Hsieh, C.-L.; Navankasattusas, S.; Escobedo, J. A.;

Williams, L.

[190639]T.; Francke, U.: Chromosomal localization of the gene for AA-type

[190640]platelet-derived growth factor receptor (PDGFRA) in humans and mice. Cytogenet.

[190641]Cell Genet. 56: 160–163, 1991.

[190642]

[190643]

[190644]20218.

[190645]Ikuno, Y.; Kazlauskas, A.: TGF-beta-1-dependent contraction of

[190646]fibroblasts is mediated by th PDGF-alpha receptor. Invest. Ophthal.

[190647]Vis. Sci. 43: 41-46, 2002.

[190648]

[190649]

[190650]20219.

[190651]Joosten, P. H. L. J.; Hol, F. A.; van Beersum, S. E. C.; Peters,

[190652]H.; Hamel, B. C. J.; Afink, G. B.; van Zoelen, E. J. J.; Mari-man,

[190653]E. C. M.: Altered regulation of platelet-derived growth factor receptor-alpha

[190654]gene-transcription in vitro by spina bifida-associated mutant Pax1

[190655]proteins. Proc. Nat. Acad. Sci. 95: 14459-14463, 1998.

[190656]

[190657]

[190658]20220.

[190659]Sanke, T.; Bell, G. I.; Sample, C.; Rubenstein, A. H.; Steiner,

[190660]D. F.: An islet amyloid peptide is derived from an
89-amino acid

[190661]precursor by proteolytic processing. J. Biol. Chem. 263:
17243-17246,

[190662]1988.

[190663]

[190664]

[190665]20221.

[190666]Verchere, C. B.; D'Alessio, D. A.; Palmiter, R. D.; Weir, G.
C.;

[190667]Bonner-Weir, S.; Baskin, D. G.; Kahn, S. E.: Islet amyloid
formation

[190668]associated with hyperglycemia in transgenic mice with
pancreatic beta

[190669]cell expression of human islet amyloid polypeptide. Proc.
Nat. Acad.

[190670]Sci. 93: 3492-3496, 1996.

[190671]

[190672]

[190673]20222.

[190674]Westermarck, P.; Engstrom, U.; Johnson, K. H.; Westermarck, G. T.;

[190675]Betsholtz, C.: Islet amyloid polypeptide: pinpointing amino acid

[190676]residues linked to amyloid fibril formation. Proc. Nat. Acad. Sci. 87:

[190677]5036–5040, 1990.

[190678]

[190679]

[190680]20223.

[190681]Lin, F.; Worman, H. J.: Structural organization of the human gene

[190682](LMNB1) encoding nuclear lamin B1. Genomics 27: 230–236, 1995.

[190683]

[190684]

[190685]20224.

[190686]Maeno, H.; Sugimoto, K.; Nakajima, N.: Genomic structure
of the

[190687]mouse gene (Lmnb1) encoding nuclear lamin B1. Ge-
nomics 30: 342–346,

[190688]1995.

[190689]

[190690]

[190691]20225.

[190692]Bignon, C.; Roux–Dosseto, M.; Zeigler, M. E.; Mattei,
M.–G.; Lissitzky,

[190693]J.–C.; Wicha, M. S.; Martin, P.–M.: Genomic analysis of the
67–kDa

[190694]laminin receptor in normal and pathological tissues: cir-
cumstantial

[190695]evidence for retroposon features. Genomics 10: 481–485,
1991.

[190696]

[190697]

[190698]20226.

[190699]Gehlsen, K. R.; Dillner, L.; Engvall, E.; Ruoslahti, E.: The
human

[190700]laminin receptor is a member of the integrin family of cell adhesion

[190701]receptors. Science 241: 1228–1229, 1988. Note: Erratum: Science

[190702]245:342–343, 1989.

[190703]

[190704]

[190705]20227.

[190706]Jackers, P.; Minoletti, F.; Belotti, D.; Clausse, N.; Sozzi, G.;

[190707]Sobel, M. E.; Castronovo, V.: Isolation from a multigene family of

[190708]the active human gene of the metastasis–associated multifunctional

[190709]protein 37LRP/p40 at chromosome 3p21.3. Oncogene 13: 495–503, 1996.

[190710]

[190711]

[190712]20228.

[190713]Lafreniere, R. G.; Brown, C. J.; Rider, S.; Chelly, J.; Taillon–Miller,

[190714]P.; Chinault, A. C.; Monaco, A. P.; Willard, H. F.: 2.6 Mb
YAC contig

[190715]of the human X inactivation center region in Xq13: physical linkage

[190716]of the RPS4X, PHKA1, XIST and DXS128E genes. Hum.
Molec. Genet. 2:

[190717]1105–1115, 1993.

[190718]

[190719]

[190720]20229.

[190721]Montuori, N.; Muller, F.; De Riu, S.; Fenzi, G.; Sobel, M. E.;

[190722]Rossi, G.; Vitale, M.: Laminin receptors in differentiated
thyroid

[190723]tumors: restricted expression of the 67–kilodalton laminin
receptor

[190724]in follicular carcinoma cells. J. Clin. Endocr. Metab. 84:
2086–2092,

[190725]1999.

[190726]

[190727]

[190728]20230.

[190729]Satoh, K.; Narumi, K.; Sakai, T.; Abe, T.; Kikuchi, T.; Matsushima,

[190730]K.; Sindoh, S.; Motomiya, M.: Cloning of 67-kDa laminin receptor

[190731]cDNA and gene expression in normal and malignant cell lines of the

[190732]human lung. Cancer Lett. 62: 199–203, 1992.

[190733]

[190734]

[190735]20231.

[190736]Tohgo, A.; Takasawa, S.; Munakata, H.; Yonekura, H.; Hayashi, N.;

[190737]Okamoto, H.: Structural determination and characterization of a 40

[190738]kDa protein isolated from rat 40 S ribosomal subunit. FEBS Lett. 340:

[190739]133–138, 1994.

[190740]

[190741]

[190742]20232.

[190743]Yow, H. K.; Wong, J. M.; Chen, H. S.; Lee, C. G.; Davis, S.;
Steele,

[190744]G. D., Jr.; Chen, L. B.: Increased mRNA expression of a
laminin-binding

[190745]protein in human colon carcinoma: complete sequence of
a full-length

[190746]cDNA encoding the protein. Proc. Nat. Acad. Sci. 85:
6394-6398,

[190747]1988. Note: Erratum: Proc. Nat. Acad. Sci. 86: 7032 only,
1989.

[190748]

[190749]

[190750]20233.

[190751]Kanzaki, T.; Olofsson, A.; Moren, A.; Wernstedt, C.; Hell-
man, U.;

[190752]Miyazono, K.; Claesson-Welsh, L.; Heldin, C. H.: TGF-beta
1 binding

[190753]protein: a component of the large latent complex of TGF-
beta 1 with

[190754]multiple repeat sequences. Cell 61: 1051-1061, 1990.

[190755]

[190756]

[190757]20234.

[190758]Oklu, R.; Hesketh, R.: The latent transforming growth factor beta

[190759]binding protein (LTBP) family. Biochem. J. 352: 601–610, 2000.

[190760]

[190761]

[190762]20235.

[190763]Stenman, G.; Sahlin, P.; Olofsson, A.; Geurts van Kessel, A.; Miyazono,

[190764]K.: Assignment of the gene encoding the latent TGF-beta-1-binding

[190765]protein (LTBP1) to human chromosome 2, region p12-q22. Cytogenet.

[190766]Cell Genet. 66: 117–119, 1994.

[190767]

[190768]

[190769]20236.

[190770]Brodsky, G. L.; Muntoni, F.; Miocic, S.; Sinagra, G.; Sewry, C.;

[190771]Mestroni, L.: Lamin A/C gene mutation associated with dilated cardiomyopathy

[190772]with variable skeletal muscle involvement. Circulation 101: 473–476,

[190773]2000.

[190774]

[190775]

[190776]20237.

[190777]Cao, H.; Hegele, R. A.: Nuclear lamin A/C R482Q mutation in Canadian

[190778]kindreds with Dunnigan-type familial partial lipodystrophy. Hum.

[190779]Molec. Genet. 9: 109–112, 2000.

[190780]

[190781]

[190782]20238.

[190783]De Sandre-Giovannoli, A.; Chaouch, M.; Kozlov, S.; Vallat, J.–M.;

- [190784]Tazir, M.; Kassouri, N.; Szepetowski, P.; Hammadouche, T.; Vandenberghe, A.; Stewart, C. L.; Grid, D.; Levy, N.: Homozygous defects in LMNA, encoding lamin A/C nuclear-envelope proteins, cause autosomal recessive axonal neuropathy in human (Charcot-Marie-Tooth disorder type 2) and mouse. Am. J. Hum. Genet. 70: 726-736, 2002. Note: Erratum: Am. J. Hum. Genet. 70: 1075 only, 2002.
- [190790]
- [190791]
- [190792]20239.
- [190793]Fatkin, D.; MacRae, C.; Sasaki, T.; Wolff, M. R.; Porcu, M.; Frenneaux, M.; Atherton, J.; Vidaillet, H. J., Jr.; Spudich, S.; De Girolami, U.; Seidman, J. G.; Seidman, C. E.: Missense mutations in the rod domain of the lamin A/C gene as causes of dilated cardiomyopathy and

[190797]conduction-system disease. New Eng. J. Med. 341:
1715-1724, 1999.

[190798]

[190799]

[190800]20240.

[190801]Fisher, D. Z.; Chaudhary, N.; Blobel, G.: cDNA sequencing
of nuclear

[190802]lamins A and C reveals primary and secondary structural
homology to

[190803]intermediate filament proteins. Proc. Nat. Acad. Sci. 83:
6450-6454,

[190804]1986.

[190805]

[190806]

[190807]20241.

[190808]Flier, J. S.: Pushing the envelope on lipodystrophy. Nature
Genet. 24:

[190809]103-104, 2000.

[190810]

[190811]

[190812]20242.

[190813]Garg, A.; Vinaitheerthan, M.; Weatherall, P. T.; Bowcock, A. M.

[190814]: Phenotypic heterogeneity in patients with familial partial lipodystrophy

[190815](Dunnigan variety) related to the site of missense mutations in lamin

[190816]A/C gene. J. Clin. Endocr. Metab. 86: 59–65, 2001.

[190817]

[190818]

[190819]20243.

[190820]Genschel, J.; Schmidt, H. H.–J.: Mutations in the LMNA gene encoding

[190821]lamin A/C. Hum. Mutat. 16: 451–459, 2000.

[190822]

[190823]

[190824]20244.

[190825]Guilly, M. N.; Bensussan, A.; Bourge, J. F.; Bornens, M.; Courvalin,

[190826]J. C.: A human T lymphoblastic cell line lacks lamins A and

C. EMBO

[190827]J. 6: 3795–3799, 1987.

[190828]

[190829]

[190830]20245.

[190831]Hegele, R. A.; Cao, H.; Harris, S. B.; Zinman, B.; Hanley, A.

[190832]J.; Anderson, C. M.: Genetic variation in LMNA modulates
plasma leptin

[190833]and indices of obesity in aboriginal Canadians. *Physiol.*

Genomics 3:

[190834]39–44, 2000.

[190835]

[190836]

[190837]20246.

[190838]Hegele, R. A.; Cao, H.; Huff, M. W.; Anderson, C. M.: LMNA
R482Q

[190839]mutation in partial lipodystrophy associated with reduced
plasma leptin

[190840]concentration. *J. Clin. Endocr. Metab.* 85: 3089–3093,
2000.

[190841]

[190842]

[190843]20247.

[190844]Hegele, R. A.; Huff, M. W.; Young, T. K.: Common genomic
variation

[190845]in LMNA modulates indexes of obesity in Inuit. J. Clin. En-
docr. Metab. 86:

[190846]2747–2751, 2001.

[190847]

[190848]

[190849]20248.

[190850]Krohne, G.; Benavente, R.: The nuclear lamins: a multigene
family

[190851]of proteins in evolution and differentiation. Exp. Cell Res.
162:

[190852]1–10, 1986.

[190853]

[190854]

[190855]20249.

[190856]Lebel, S.; Raymond, Y.: Lamin A is not synthesized as a
larger

[190857]precursor polypeptide. Biochem. Biophys. Res. Commun.
149: 417–423,

[190858]1987.

[190859]

[190860]

[190861]20250.

[190862]Lin, F.; Worman, H. J.: Structural organization of the hu-
man

[190863]gene encoding nuclear lamin A and nuclear lamin C. J.
Biol. Chem. 268:

[190864]16321–16326, 1993.

[190865]

[190866]

[190867]20251.

[190868]Lloyd, D. J.; Trembath, R. C.; Shackleton, S.: A novel inter-
action

[190869]between lamin A and SREBP1: implications for partial
lipodystrophy

[190870]and other laminopathies. Hum. Molec. Genet. 11:
769–777, 2002.

[190871]

[190872]

[190873]20252.

[190874]McKeon, F. D.; Kirschner, M. W.; Caput, D.: Homologies in
both

[190875]primary and secondary structure between nuclear enve-
lope and intermediate

[190876]filament proteins. Nature 319: 463–468, 1986.

[190877]

[190878]

[190879]20253.

[190880]Muchir, A.; Bonne, G.; van der Kooi, A. J.; van Meegen, M.;
Baas,

[190881]F.; Bolhuis, P. A.; de Visser, M.; Schwartz, K.: Identification
of

[190882]mutations in the gene encoding lamins A/C in autosomal
dominant limb

[190883]girdle muscular dystrophy with atrioventricular conduction
disturbances

[190884](LGMD1B). Hum. Molec. Genet. 9: 1453–1459, 2000.

[190885]

[190886]

[190887]20254.

[190888]Novelli, G.; Muchir, A.; Sangiuolo, F.; Helbling–Leclerc, A.;

[190889]D'Apice, M. R.; Massart, C.; Capon, F.; Sbraccia, P.; Federici, M.;

[190890]Lauro, R.; Tudisco, C.; Pallotta, R.; Scarano, G.; Dallapiccola, B.;

[190891]Merlini, L.; Bonne, G.: Mandibuloacral dysplasia is caused by a mutation

[190892]in LMNA–encoding lamin A/C. Am. J. Hum. Genet. 71: 426–431, 2002.

[190893]

[190894]

[190895]20255.

[190896]Raffaele di Barletta, M.; Ricci, E.; Galluzzi, G.; Tonali, P.;

[190897]Mora, M.; Morandi, L.; Romorini, A.; Voit, T.; Orstavik, K. H.; Merlini,

[190898]L.; Trevisan, C.; Biancalana, V.; Housmanowa–Petrusewicz,

I.; Bione,

[190899]S.; Ricotti, R.; Schwartz, K.; Bonne, G.; Toniolo, D.: Differ-
ent

[190900]mutations in the LMNA gene cause autosomal dominant
and autosomal

[190901]recessive Emery–Dreifuss muscular dystrophy. Am. J.

Hum. Genet. 66:

[190902]1407–1412, 2000.

[190903]

[190904]

[190905]20256.

[190906]Schmidt, H. H.–J.; Genschel, J.; Baier, P.; Schmidt, M.; Ock-
enga,

[190907]J.; Tietge, U. J. F.; Propsting, M.; Buttner, C.; Manns, M. P.;
Lochs,

[190908]H.; Brabant, G.: Dyslipemia in familial partial lipodystrophy
caused

[190909]by and R482W mutation in the LMNA gene. J. Clin. Endocr.
Metab. 86:

[190910]2289–2295, 2001.

[190911]

[190912]

[190913]20257.

[190914]Shackleton, S.; Lloyd, D. J.; Jackson, S. N. J.; Evans, R.;

Niermeijer,

[190915]M. F.; Singh, B. M.; Schmidt, H.; Brabant, G.; Kumar, S.;

Durrington,

[190916]P. N.; Gregory, S.; O'Rahilly, S.; Trembath, R. C.: LMNA,

encoding

[190917]lamin A/C, is mutated in partial lipodystrophy. Nature

Genet. 24:

[190918]153–156, 2000.

[190919]

[190920]

[190921]20258.

[190922]Wydner, K. L.; McNeil, J. A.; Lin, F.; Worman, H. J.;

Lawrence,

[190923]J. B.: Chromosomal assignment of human nuclear enve-

lope protein genes

[190924]LMNA, LMNB1, and LBR by fluorescence in situ hybridiza-

tion. Genomics 32:

[190925]474–478, 1996.

[190926]

[190927]

[190928]20259.

[190929]Furukawa, K.; Hotta, Y.: cDNA cloning of a germ cell specific

[190930]lamin B3 from mouse spermatocytes and analysis of its function by

[190931]ectopic expression in somatic cells. EMBO J. 12: 97–106, 1993.

[190932]

[190933]

[190934]20260.

[190935]Furukawa, K.; Inagaki, H.; Hotta, Y.: Identification and cloning

[190936]of an mRNA coding for a germ cell-specific A-type lamin in mice.

[190937]Exp. Cell Res. 212: 426–430, 1994.

[190938]

[190939]

[190940]20261.

[190941]Garbarz, M.; Devaux, I.; Bournier, O.; Grandchamp, B.;
Dhermy,

[190942]D.: Protein 4.1 Lille, a novel mutation in the downstream
initiation

[190943]codon of protein 4.1 gene associated with heterozygous
4,1(-) hereditary

[190944]elliptocytosis. Hum. Mutat. 5: 339–340, 1995.

[190945]

[190946]

[190947]20262.

[190948]Garbarz, M.; Dhermy, D.; Lecomte, M. C.; Feo, C.; Chave-
roche,

[190949]I.; Galand, C.; Bournier, O.; Bertrand, O.; Boivin, P.: A vari-
ant

[190950]of erythrocyte membrane skeletal protein band 4.1 asso-
ciated with

[190951]hereditary elliptocytosis. Blood 64: 1006–1015, 1984.

[190952]

[190953]

[190954]20263.

[190955]Geerdink, R. A.; Nijenhuis, L. E.; Huizinga, J.: Hereditary
elliptocytosis:

[190956]linkage data in man. Ann. Hum. Genet. 30: 363–378,
1967.

[190957]

[190958]

[190959]20264.

[190960]Jensson, O.; Jonasson, T.; Olafsson, O.: Hereditary ellipto-
cytosis

[190961]in Iceland. Brit. J. Haemat. 13: 844–854, 1967.

[190962]

[190963]

[190964]20265.

[190965]Kan, Y.–W.: Personal Communication. San Francisco, Calif.

[190966]2/28/1986.

[190967]

[190968]

[190969]20266.

[190970]Kuroda, S.; Takeuchi, T.; Nagamori, H.: Data on the link–

age between

[190971]elliptocytosis and Rh blood type. Jpn. J. Hum. Genet. 5:
112–118,

[190972]1960.

[190973]

[190974]

[190975]20267.

[190976]Lambert, S.; Conboy, J.; Zail, S.: A molecular study of het-
erozygous

[190977]protein 4.1 deficiency in hereditary elliptocytosis. Blood
72: 1926–1929,

[190978]1988.

[190979]

[190980]

[190981]20268.

[190982]Lambert, S.; Zail, S.: Partial deficiency of protein 4.1 in
hereditary

[190983]elliptocytosis. Am. J. Hemat. 26: 263–272, 1987.

[190984]

[190985]

[190986]20269.

[190987]Lipton, E. L.: Elliptocytosis with hemolytic anemia: the effects

[190988]of splenectomy. Pediatrics 15: 67–82, 1955.

[190989]

[190990]

[190991]20270.

[190992]Lux, S. E.; Wolfe, L. C.: Inherited disorders of the red cell

[190993]membrane skeleton. Pediat. Clin. N. Am. 27: 463–486, 1980.

[190994]

[190995]

[190996]20271.

[190997]Marchesi, S. L.; Conboy, J.; Agre, P.; Letsinger, J. T.;

Marchesi,

[190998]V. T.; Speicher, D. W.; Mohandas, N.: Molecular analysis of insertion/deletion

[190999]mutations in protein 4.1 in elliptocytosis. I. Biochemical

identification

[191000]of rearrangements in the spectrin/actin binding domain
and functional

[191001]characterizations. J. Clin. Invest. 86: 516–523, 1990.

[191002]

[191003]

[191004]20272.

[191005]McGuire, M.; Agre, P.: Three distinct variants of protein
4.1

[191006]in Caucasian hereditary elliptocytosis. (Abstract) Clin. Res.
35:

[191007]428A, 1987.

[191008]

[191009]

[191010]20273.

[191011]McGuire, M.; Smith, B. L.; Agre, P.: Distinct variants of ery-
throcyte

[191012]protein 4.1 inherited in linkage with elliptocytosis and Rh
type in

[191013]three white families. Blood 72: 287–293, 1988.

[191014]

[191015]

[191016]20274.

[191017]Morle, L.; Garbarz, M.; Alloisio, N.; Girot, R.; Chaveroche,
I.;

[191018]Boivin, P.; Delaunay, J.: The characterization of protein 4.1
Presles,

[191019]a shortened variant of RBC membrane protein 4.1. Blood
65: 1511–1517,

[191020]1985.

[191021]

[191022]

[191023]20275.

[191024]Morle, L.; Pothier, B.; Alloisio, N.; Ducluzeau, M.–T.; Mar-
ques,

[191025]S.; Olim, G.; Martins e Silva, J.; Feo, C.; Garbarz, M.;
Chaveroche,

[191026]I.; Boivin, P.; Delaunay, J.: Red cell membrane alteration
involving

[191027]protein 4.1 and protein 3 in a case of recessively inherited
haemolytic

[191028]anemia. Europ. J. Haemat. 38: 447–455, 1987.

[191029]

[191030]

[191031]20276.

[191032]Morton, N. E.: The detection and estimation of linkage between
tween

[191033]the genes for elliptocytosis and the Rh blood type. Am. J.
Hum. Genet. 8:

[191034]80–96, 1956.

[191035]

[191036]

[191037]20277.

[191038]Nielsen, J. A.; Strunk, K. W.: Homozygous hereditary ellip-
tocytes

[191039]as the cause of haemolytic anemia in infancy. Scand. J.
Haemat. 5:

[191040]486–496, 1968.

[191041]

[191042]

[191043]20278.

[191044]Parra, M.; Gascard, P.; Walensky, L. D.; Snyder, S. H.; Mo-
handas,

[191045]N.; Conboy, J. G.: Cloning and characterization of 4.1G
(EPB41L2),

[191046]a new member of the skeletal protein 4.1 (EPB41) gene
family. Genomics 49:

[191047]298–306, 1998.

[191048]

[191049]

[191050]20279.

[191051]Peters, J. C.; Rowland, M.; Israels, L. G.; Zipursky, A.: Ery-
throcyte

[191052]sodium transport in hereditary elliptocytosis. Canad. J.
Physiol.

[191053]Pharm. 44: 817–827, 1966.

[191054]

[191055]

[191056]20280.

[191057]Roberts, J. A. F.: Genetic linkage in man, with particular

reference

[191058]to the usefulness of very small bodies of data. Quart. J.

Med. 14:

[191059]27–33, 1945.

[191060]

[191061]

[191062]20281.

[191063]Shi, Z.–T.; Afzal, V.; Coller, B.; Patel, D.; Chasis, J. A.;

Parra,

[191064]M.; Lee, G.; Paszty, C.; Stevens, M.; Walensky, L.; Peters, L.

L.;

[191065]Mohandas, N.; Rubin, E.; Conboy, J. G.: Protein

4.1R–deficient mice

[191066]are viable but have erythroid membrane skeleton abnormalities. J.

[191067]Clin. Invest. 103: 331–340, 1999.

[191068]

[191069]

[191070]20282.

[191071]Takakuwa, Y.; Tchernia, G.; Rossi, M.; Benabadji, M.; Mo–

handas,

[191072]N.: Restoration of normal membrane stability to unstable protein

[191073]4.1-deficient erythrocyte membranes by incorporation of purified protein

[191074]4.1. J. Clin. Invest. 78: 80-85, 1986.

[191075]

[191076]

[191077]20283.

[191078]Tang, C.-J. C.; Tang, T. K.: Rapid localization of membrane skeletal

[191079]protein 4.1 (EL1) to human chromosome 1p33-p34.2 by nonradioactive

[191080]in situ hybridization. Cytogenet. Cell Genet. 57: 119, 1991.

[191081]

[191082]

[191083]20284.

[191084]Tang, T. K.; Leto, T. L.; Correias, I.; Alonso, M. A.; Marchesi,

[191085]V. T.; Benz, E. J., Jr.: Selective expression of an erythroid-

specific

[191086]isoform of protein 4.1. Proc. Nat. Acad. Sci. 85:
3713–3717, 1988.

[191087]

[191088]

[191089]20285.

[191090]Tchernia, G.; Mohandas, N.; Shohet, S. B.: Deficiency of
skeletal

[191091]membrane protein band 4.1 in homozygous hereditary el-
liptocytosis:

[191092]implications for erythrocyte membrane stability. J. Clin.
Invest. 68:

[191093]454–460, 1981.

[191094]

[191095]

[191096]20286.

[191097]Tomaselli, M. B.; John, K. M.; Lux, S. E.: Elliptical erythro-
cyte

[191098]membrane skeletons and heat-sensitive spectrin in
hereditary elliptocytosis. Proc.

[191099]Nat. Acad. Sci. 78: 1911–1915, 1981.

[191100]

[191101]

[191102]20287.

[191103]Brands, J. H. G. M.; Maassen, J. A.; Van Hemert, F. J.;
Amons,

[191104]R.; Moller, W.: The primary structure of the alpha subunit
of human

[191105]elongation factor 1: structural aspects of guanine-nu-
cleotide-binding

[191106]sites. Europ. J. Biochem. 155: 167-171, 1986.

[191107]

[191108]

[191109]20288.

[191110]Ditzel, H. J.; Masaki, Y.; Nielsen, H.; Farnaes, L.; Burton, D.

[191111]R.: Cloning and expression of a novel human antibody-
-antigen pair

[191112]associated with Felty's syndrome. Proc. Nat. Acad. Sci. 97:
9234-9239,

[191113]2000.

[191114]

[191115]

[191116]20289.

[191117]Lund, A.; Knudsen, S. M.; Vissing, H.; Clark, B.; Tommerup, N.

[191118]: Assignment of human elongation factor 1-alpha genes: EEF1A maps

[191119]to chromosome 6q14 and EEF1A2 to 20q13.3. Genomics 36: 359-361,

[191120]1996.

[191121]

[191122]

[191123]20290.

[191124]Opdenakker, G.; Cabeza-Arvelaiz, Y.; Fiten, P.; Dijkmans, R.; Van

[191125]Damme, J.; Volckaert, G.; Billiau, A.; Van Elsen, A.; Van der Schueren,

[191126]B.; Van den Berghe, H.; Cassiman, J.-J.: Human elongation factor

[191127]1-alpha: a polymorphic and conserved multigene family with multiple

[191128]chromosomal localizations. Hum. Genet. 75: 339-344,

1987.

[191129]

[191130]

[191131]20291.

[191132]Sparkes, R. S.; Sparkes, M. C.; Wilson, M. G.; Towner, J. W.;

[191133]Benedict, W.; Murphree, A. L.; Yunis, J. J.: Regional assign-
ment

[191134]of genes for human esterase D and retinoblastoma to
chromosome band

[191135]13q14. Science 208: 1042–1044, 1980.

[191136]

[191137]

[191138]20292.

[191139]Squire, J.; Dryja, T. P.; Dunn, J.; Goddard, A.; Hofmann, T.;

[191140]Musarella, M.; Willard, H. F.; Becker, A. J.; Gallie, B. L.;
Phillips,

[191141]R. A.: Cloning of the esterase D gene: a polymorphic gene
probe closely

[191142]linked to the retinoblastoma locus on chromosome 13.

Proc. Nat. Acad.

[191143]Sci. 83: 6573–6577, 1986.

[191144]

[191145]

[191146]20293.

[191147]Cohen, A. J.; Li, F. P.; Berg, S.; Marchetto, D. J.; Tsai, S.;

[191148]Jacobs, S. C.; Brown, R. S.: Hereditary renal–cell carcinoma
associated

[191149]with chromosomal translocation. New Eng. J. Med. 301:
592–595, 1979.

[191150]

[191151]

[191152]20294.

[191153]Gemmell, R. M.; West, J. D.; Boldog, F.; Tanaka, N.; Robin-
son,

[191154]L. J.; Smith, D. I.; Li, F.; Drabkin, H. A.: The hereditary re-
nal

[191155]cell carcinoma 3;8 translocation fuses FHIT to a patched-
related gene,

[191156]TRC8. Proc. Nat. Acad. Sci. 95: 9572–9577, 1998.

[191157]

[191158]

[191159]20295.

[191160]Nakano, A.; Pulkkinen, L.; Murrell, D.; Rico, J.; Lucky, A.
W.;

[191161]Garzon, M.; Stevens, C. A.; Robertson, S.; Pfendner, E.;
Uitto, J.

[191162]: Epidermolysis bullosa with congenital pyloric atresia:
novel mutations

[191163]in the beta-4 integrin gene (ITGB4) and genotype/phe-
notype correlations. Pediat.

[191164]Res. 49: 618-626, 2001.

[191165]

[191166]

[191167]20296.

[191168]Pulkkinen, L.; Bruckner-Tuderman, L.; August, C.; Uitto, J.:
Compound

[191169]heterozygosity for missense mutation (L156P) and non-
sense (R554X)

[191170]mutations in the beta-4 integrin gene (ITGB4) underlies
mild, nonlethal

[191171]phenotype of epidermolysis bullosa with pyloric atresia.

Am. J. Path. 152:

[191172]935–941, 1998.

[191173]

[191174]

[191175]20297.

[191176]Pulkkinen, L.; Rouan, F.; Bruckner–Tuderman, L.; Waller–
stein, R.;

[191177]Garzon, M.; Brown, T.; Smith, L.; Carter, W.; Uitto, J.: Novel
ITGB4

[191178]mutations in lethal and nonlethal variants of epidermoly–
sis bullosa

[191179]with pyloric atresia: missense versus nonsense. Am. J.

Hum. Genet. 63:

[191180]1376–1387, 1998.

[191181]

[191182]

[191183]20298.

[191184]Shaw, L. M.; Rabinovitz, I., Wang, H. H.–F.; Toker, A., Mer–
curio,

[191185]A. M.: Activation of phosphoinositide 3–OH kinase by the

alpha-6/beta-4

[191186]integrin promotes carcinoma invasion. Cell 91: 949-960, 1997.

[191187]

[191188]

[191189]20299.

[191190]Suzuki, S.; Naitoh, Y.: Amino acid sequence of a novel integrin

[191191]beta-4 subunit and primary expression of the mRNA in epithelial cells. EMBO

[191192]J. 9: 757-763, 1990.

[191193]

[191194]

[191195]20300.

[191196]Vidal, F.; Aberdam, D.; Miquel, C.; Christiano, A. M.; Pulkkinen,

[191197]L.; Uitto, J.; Ortonne, J.-P.; Meneguzzi, G.: Integrin beta-4 mutations

[191198]associated with junctional epidermolysis bullosa with pyloric atresia. Nature

[191199]Genet. 10: 229-234, 1995.

[191200]

[191201]

[191202]20301.

[191203]Krissansen, G. W.; Yuan, Q.; Jenkins, D.; Jiang, W.-M.;
Rooke,

[191204]L.; Spurr, N. K.; Eccles, M.; Leung, E.; Watson, J. D.: Chro-
mosomal

[191205]locations of the genes coding for the integrin beta-6 and
beta-7 subunits.

[191206]Immunogenetics 35: 58-61, 1992.

[191207]

[191208]

[191209]20302.

[191210]Baker, E.; Sutherland, G. R.; Jiang, W.-M.; Yuan, Q.; Leung,
E.;

[191211]Watson, J. D.; Krissansen, G. W.: Mapping of the human
integrin beta-7

[191212]gene (ITG-beta-7) to 12q13.13 by non-isotopic in situ
hybridization.

[191213]Mammalian Genome 2: 272-273, 1992.

[191214]

[191215]

[191216]20303.

[191217]Erle, D. J.; Ruegg, C.; Sheppard, D.; Pytela, R.: Complete
amino

[191218]acid sequence of an integrin beta subunit (beta-7) identified in leukocytes.

[191219]J. Biol. Chem. 266: 11009-11016, 1991.

[191220]

[191221]

[191222]20304.

[191223]Yuan, Q.; Kozak, C. A.; Jiang, W.; Hollander, D.; Watson, J.
D.;

[191224]Krissansen, G. W.: Genetic mapping of the gene coding for
the integrin

[191225]beta-7 subunit to the distal part of mouse chromosome
15. Immunogenetics 35:

[191226]403-407, 1992.

[191227]

[191228]

[191229]20305.

[191230]Chany, C.; Vignal, M.; Couillin, P.; Van Cong, N.; Boue, J.;
Boue,

[191231]A.: Chromosomal localization of human genes governing
the interferon-induced

[191232]antiviral state. Proc. Nat. Acad. Sci. 72: 3129–3133, 1975.

[191233]

[191234]

[191235]20306.

[191236]Dithmar, S.; Rusciano, D.; Lynn, M. J.; Lawson, D. H.; Arm-
strong,

[191237]C. A.; Grossniklaus, H. E.: Neoadjuvant interferon alfa-2b
treatment

[191238]in a murine model for metastatic ocular melanoma: a pre-
liminary study. Arch.

[191239]Ophthal. 118: 1085–1089, 2000.

[191240]

[191241]

[191242]20307.

[191243]Ezekowitz, R. A. B.; Mulliken, J. B.; Folkman, J.: Interferon

[191244]alfa-2a therapy for life-threatening hemangiomas of infancy. New

[191245]Eng. J. Med. 326: 1456-1463, 1992.

[191246]

[191247]

[191248]20308.

[191249]Hejny, C.; Sternberg, P., Jr.; Lawson, D. H.; Greiner, K.; Aaberg,

[191250]T. M., Jr.: Retinopathy associated with high-dose interferon alfa-2b

[191251]therapy. Am. J. Ophthal. 131: 782-787, 2001.

[191252]

[191253]

[191254]20309.

[191255]Aplin, H. M.; Hirst, K. L.; Crosby, A. H.; Dixon, M. J.: Mapping

[191256]of the human dentin matrix acidic phosphoprotein gene (DMP1) to the

[191257]dentinogenesis imperfecta type II critical region at chromosome 4q21. Genomics 30:

[191258]347–349, 1995.

[191259]

[191260]

[191261]20310.

[191262]Fisher, L. W.; McBride, O. W.; Termine, J. D.; Young, M. F.:

Human

[191263]bone sialoprotein: deduced protein sequence and chromosomal localization.

[191264]J. Biol. Chem. 265: 2347–2351, 1990.

[191265]

[191266]

[191267]20311.

[191268]Goold, R. D.; diSibio, G. L.; Xu, H.; Lang, D. B.; Dadgar, J.;

[191269]Magrane, G. G.; Dugaiczyk, A.; Smith, K. A.; Cox, D. R.;

Masters,

[191270]S. B.; Myers, R. M.: The development of sequence-tagged sites for

[191271]human chromosome 4. Hum. Molec. Genet. 2:

1271–1288, 1993.

[191272]

[191273]

[191274]20312.

[191275]Kerr, J. M.; Fisher, L. W.; Termine, J. D.; Wang, M. G.;
McBride,

[191276]O. W.; Young, M. F.: The human bone sialoprotein gene
(IBSP): genomic

[191277]localization and characterization. Genomics 17: 408–415,
1993.

[191278]

[191279]

[191280]20313.

[191281]Kim, R. H.; Shapiro, H. S.; Li, J. J.; Wrana, J. L.; Sodek, J.

[191282]: Characterization of the human bone sialoprotein (BSP)
gene and its

[191283]promoter sequence. Matrix Biol. 14: 31–40, 1994.

[191284]

[191285]

[191286]20314.

[191287]Young, M. F.; Ibaraki, K.; Kerr, J. M.; Lyu, M. S.; Kozak, C.

A.

[191288]: Murine bone sialoprotein (BSP): cDNA cloning, mRNA expression, and

[191289]genetic mapping. Mammalian Genome 5: 108–111, 1994.

[191290]

[191291]

[191292]20315.

[191293]Holbrook, S. T.; Ohls, R. K.; Schibler, K. R.; Yang, Y.-C.; Christensen,

[191294]R. D.: Effect of interleukin–9 on clonogenic maturation and cell–cycle

[191295]status of fetal and adult hematopoietic progenitors. Blood 77: 2129–2134,

[191296]1991.

[191297]

[191298]

[191299]20316.

[191300]Kelleher, K.; Bean, K.; Clark, S. C.; Leung, W.-Y.; Yang-Feng,

[191301]T. L.; Chen, J. W.; Lin, P.-F.; Luo, W.; Yang, Y.-C.: Human

interleukin-9:

[191302]genomic sequence, chromosomal location, and sequences
essential for

[191303]its expression in human T-cell leukemia virus
(HTLV)-I-transformed

[191304]human T cells. Blood 77: 1436-1441, 1991.

[191305]

[191306]

[191307]20317.

[191308]Merz, H.; Houssiau, F. A.; Orscheschek, K.; Renauld, J. C.;
Fliedner,

[191309]A.; Herin, M.; Noel, H.; Kadin, M.; Mueller-Hermelink, H.
K.; Van

[191310]Snick, J.; Feller, A. C.: Interleukin-9 expression in human
malignant

[191311]lymphomas: unique association with Hodgkin's disease
and large cell

[191312]anaplastic lymphoma. Blood 78: 1311-1317, 1991.

[191313]

[191314]

[191315]20318.

- [191316]Mock, B. A.; Krall, M.; Kozak, C. A.; Nesbitt, M. N.;
McBride,
[191317]O. W.; Renauld, J.-C.; Van Snick, J.: IL9 maps to mouse
chromosome
[191318]13 and human chromosome 5. Immunogenetics 31:
265–270, 1990.
- [191319]
- [191320]
- [191321]20319.
- [191322]Nicolaidis, N. C.; Holroyd, K. J.; Ewart, S. L.; Eleff, S. M.;
- [191323]Kiser, M. B.; Dragwa, C. R.; Sullivan, C. D.; Grasso, L.;
- Zhang, L.-Y.;
- [191324]Messler, C. J.; Zhou, T.; Kleeberger, S. R.; Beutow, K. H.;
- Levitt,
[191325]R. C.: Interleukin 9: a candidate gene for asthma. Proc.
Nat. Acad.
[191326]Sci. 94: 13175–13180, 1997.
- [191327]
- [191328]
- [191329]20320.

[191330]Postma, D. S.; Bleecker, E. R.; Amelung, P. J.; Holroyd, K.
J.;

[191331]Xu, J.; Panhuysen, C. I. M.; Meyers, D. A.; Levitt, R. C.: Ge-
netic

[191332]susceptibility to asthma: bronchial hyperresponsiveness
coinherited

[191333]with a major gene for atopy. New Eng. J. Med. 333:
894–900, 1995.

[191334]

[191335]

[191336]20321.

[191337]Van Snick, J.; Goethals, A.; Renauld, J.–C.; Van Roost, E.;
Uyttenhove,

[191338]C.; Rubira, M. R.; Moritz, R. L.; Simpson, R. J.: Cloning and
characterization

[191339]of a cDNA for a new mouse T cell growth factor (P40). J.
Exp. Med. 169:

[191340]363–368, 1989.

[191341]

[191342]

[191343]20322.

[191344]Yang, Y.-C.; Ricciardi, S.; Ciarletta, A.; Calvetti, J.; Keller,
her,

[191345]K.; Clark, S. C.: Expression cloning of a cDNA encoding a
novel human

[191346]hematopoietic growth factor: human homologue of
murine T-cell growth

[191347]factor P40. Blood 74: 1880-1884, 1989.

[191348]

[191349]

[191350]20323.

[191351]Ho, A. S. Y.; Liu, Y.; Khan, T. A.; Hsu, D.-H.; Bazan, J. F.;
Moore,

[191352]K. W.: A receptor for interleukin 10 is related to interferon
receptors. Proc.

[191353]Nat. Acad. Sci. 90: 11267-11271, 1993.

[191354]

[191355]

[191356]20324.

[191357]Liu, Y.; Wei, S. H.-Y.; Ho, A. S.-Y.; de Waal Malefyt, R.;
Moore,

[191358]K. W.: Expression cloning and characterization of a human
IL-10 receptor. J.

[191359]Immun. 152: 1821–1829, 1994.

[191360]

[191361]

[191362]20325.

[191363]Blixt, A.; Mahlapuu, M.; Aitola, M.; Peltö-Huikko, M.; Ener-
back,

[191364]S.; Carlsson, P.: A forkhead gene, FoxE3, is essential for
lens epithelial

[191365]proliferation and closure of the lens vesicle. Genes Dev.
14: 245–254,

[191366]2000.

[191367]

[191368]

[191369]20326.

[191370]Semina, E. V.; Brownell, I.; Mintz-Hittner, H. A.; Murray, J.
C.;

[191371]Jamrich, M.: Mutations in the human forkhead transcrip-
tion factor

[191372]FOXEO associated with anterior segment ocular dysgenesis and cataracts. Hum.

[191373]Molec. Genet. 10: 231–236, 2001.

[191374]

[191375]

[191376]20327.

[191377]Siegelmann–Danieli, N.; Buetow, K. H.: Constitutional genetic

[191378]variation at the human aromatase gene (Cyp19) and breast cancer risk. Brit.

[191379]J. Cancer 79: 456–463, 1999.

[191380]

[191381]

[191382]20328.

[191383]Simpson, E. R.; Michael, M. D.; Agarwal, V. R.; Hinshelwood, M.

[191384]M.; Bulun, S. E.; Zhao, Y.: Expression of the CYP19 (aromatase) gene:

[191385]an unusual case of alternative promoter usage. FASEB J. 11: 29–36,

[191386]1997.

[191387]

[191388]

[191389]20329.

[191390]Sparkes, R. S.; Mohandas, T.; Chen, S.; Besman, M. J.; Zollman,

[191391]S.; Shively, J. E.: Assignment of the aromatase gene to human chromosome

[191392]15q21. (Abstract) Cytogenet. Cell Genet. 46: 696–697, 1987.

[191393]

[191394]

[191395]20330.

[191396]Toda, K.; Merashima, M.; Kawamoto, T.; Sumimoto, H.; Yokoyama,

[191397]Y.; Kuribayashi, I.; Mitsuuchi, Y.; Maeda, T.; Yamamoto, Y.; Sagara,

[191398]Y.; Ikeda, H.; Shizuta, Y.: Structural and functional characterization

[191399]of human aromatase P-450 gene. Europ. J. Biochem. 193: 559–565,

[191400]1990.

[191401]

[191402]

[191403]20331.

[191404]Wang, Z. J.; Jeffs, B.; Ito, M.; Achermann, J. C.; Yu, R. N.;

[191405]Hales, D. B.; Jameson, J. L.: Aromatase (Cyp19) expression
is up-regulated

[191406]by targeted disruption of Dax1. Proc. Nat. Acad. Sci. 98:
7988–7993,

[191407]2001.

[191408]

[191409]

[191410]20332.

[191411]Whitlock, J. P., Jr.: The regulation of cytochrome P-450
gene

[191412]expression. Annu. Rev. Pharm. Toxicol. 26: 333–369,
1986.

[191413]

[191414]

[191415]20333.

[191416]Zhou, D.; Pompon, D.; Chen, S.: Structure–function studies
of

[191417]human aromatase by site–directed mutagenesis: kinetic
properties of

[191418]mutants pro308–to–phe, tyr361–to–phe, tyr361–to–leu,
and phe406–to–arg. Proc.

[191419]Nat. Acad. Sci. 88: 410–414, 1991.

[191420]

[191421]

[191422]20334.

[191423]Bruneau, G.; Gross, M.–S.; Krieger, M.; Bernheim, A.;
Thibault,

[191424]J.; Nguyen, V. C.: Preparation of a human DOPA decar–
boxylase cDNA

[191425]probe by PCR and its assignment to chromosome 7. Ann.
Genet. 33:

[191426]208–213, 1990.

[191427]

[191428]

[191429]20335.

[191430]Christenson, J. G.; Dairman, W.; Udenfriend, S.: On the
identity

[191431]of DOPA decarboxylase and 5-hydroxytryptophan decar-
boxylase (immunological

[191432]titration-aromatic L-amino acid decarboxylase-sero-
tonin-dopamine-norepinephrine).

[191433]Proc. Nat. Acad. Sci. 69: 343-347, 1972.

[191434]

[191435]

[191436]20336.

[191437]Craig, S. P.; Le Van Thai, A.; Weber, M.; Craig, I. W.: Locali-
sation

[191438]of the gene for human aromatic L-amino acid decarboxy-
lase (DDC) to

[191439]chromosome 7p13-p11 by in situ hybridisation. Cyto-
genet. Cell Genet. 61:

[191440]114-116, 1992.

[191441]

[191442]

[191443]20337.

[191444]Ichinose, H.; Kurosawa, Y.; Titani, K.; Fujita, K.; Nagatsu, T.

[191445]: Isolation and characterization of a cDNA clone encoding human aromatic

[191446]L-amino acid decarboxylase. Biochem. Biophys. Res. Commun. 164:

[191447]1024–1030, 1989.

[191448]

[191449]

[191450]20338.

[191451]Scherer, L. J.; McPherson, J. D.; Wasmuth, J. J.; Marsh, J. L.

[191452]: Human dopa decarboxylase: localization to human chromosome 7p11

[191453]and characterization of hepatic cDNAs. Genomics 13: 469–471, 1992.

[191454]

[191455]

[191456]20339.

[191457]Sumi-Ichinose, C.; Ichinose, H.; Takahashi, E.; Hori, T.;

Nagatsu,

[191458]T.: Molecular cloning of genomic DNA and chromosomal assignment of

[191459]the gene for human aromatic L-amino acid decarboxylase, the enzyme

[191460]for catecholamine and serotonin biosynthesis. Biochemistry 31:

[191461]2229–2238, 1992.

[191462]

[191463]

[191464]20340.

[191465]Luttrell, L. M.; Ferguson, S. S. G.; Daaka, Y.; Miller, W. E.;

[191466]Maudsley, S.; Della Rocca, G. J.; Lin, F.-T.; Kawakatsu, H.; Owada,

[191467]K.; Luttrell, D. K.; Caron, M. G.; Lefkowitz, R. J.: Beta-arrestin-dependent

[191468]formation of beta-2 adrenergic receptor–Src protein kinase complexes. Science 283:

[191469]655–661, 1999.

[191470]

[191471]

[191472]20341.

[191473]Attramadal, H.; Arriza, J. L.; Aoki, C.; Dawson, T. M.; Codina,

[191474]J.; Kwatra, M. M.; Snyder, S. H.; Caron, M. G.; Lefkowitz, R. J.:

[191475]Beta-arrestin-2, a novel member of the arrestin/
beta-arrestin gene

[191476]family. J. Biol. Chem. 267: 17882-17890, 1992.

[191477]

[191478]

[191479]20342.

[191480]Bohn, L. M.; Gainetdinov, R. R.; Lin, F.-T.; Lefkowitz, R. J.;

[191481]Caron, M. G.: Mu-opioid receptor desensitization by beta-arrestin-2

[191482]determines morphine tolerance but not dependence. Nature 408: 720-723,

[191483]2000.

[191484]

[191485]

[191486]20343.

[191487]Bohn, L. M.; Lefkowitz, R. J.; Gainetdinov, R. R.; Peppel, K.;

[191488]Caron, M. G.; Lin, F.-T.: Enhanced morphine analgesia in mice lacking

[191489]beta-arrestin 2. Science 286: 2495–2498, 1999.

[191490]

[191491]

[191492]20344.

[191493]Fong, F. M.; Premont, R. T.; Richardson, R. M.; Yu, Y.-R. A.; Lefkowitz,

[191494]R. J.; Patel, D. D.: Defective lymphocyte chemotaxis in beta-arrestin2–

[191495]and GRK6-deficient mice. Proc. Nat. Acad. Sci. 99: 7478–7483, 2002.

[191496]

[191497]

[191498]20345.

[191499]Kiselev, A.; Socolich, M.; Vinos, J.; Hardy, R. W.; Zuker, C. S.;

[191500]Ranganathan, R.: A molecular pathway for light-dependent photoreceptor

[191501]apoptosis in Drosophila. Neuron 28: 139–152, 2000.

[191502]

[191503]

[191504]20346.

[191505]McDonald, P. H.; Chow, C.–W.; Miller, W. E.; Laporte, S. A.;
Field,

[191506]M. E.; Lin, F.–T.; Davis, R. J.; Lefkowitz, R. J.: Beta–arrestin

[191507]2: a receptor–regulated MAPK scaffold for the activation of
JNK3. Science 290:

[191508]1574–1577, 2000.

[191509]

[191510]

[191511]20347.

[191512]Karathanasis, S. K.; Oettgen, P.; Haddad, I. A.; An–
tonarakis,

[191513]S. E.: Structure, evolution, and polymorphisms of the hu–
man apolipoprotein

[191514]A4 gene (APOA4). Proc. Nat. Acad. Sci. 83: 8457–8461,
1986.

[191515]

[191516]

[191517]20348.

[191518]Carrington, M.; Kissner, T.; Gerrard, B.; Ivanov, S.; O'Brien,

[191519]S. J.; Dean, M.: Novel alleles of the chemokine–receptor
gene CCR5. Am.

[191520]J. Hum. Genet. 61: 1261–1267, 1997.

[191521]

[191522]

[191523]20349.

[191524]Chaganti, R. S. K.: Personal Communication. New York, N.
Y.

[191525]10/22/1993.

[191526]

[191527]

[191528]20350.

[191529]Chaudhuri, A.; Polyakova, J.; Zbrzezna, V .; Pogo, A. O.:
The

[191530]coding sequence of Duffy blood group gene in humans
and simians: restriction

[191531]fragment length polymorphism, antibody and malarial
parasite specificities,
[191532]and expression in nonerythroid tissues in Duffy-negative
individuals. Blood 85:
[191533]615–621, 1995.

[191534]

[191535]

[191536]20351.

[191537]Chaudhuri, A.; Polyakova, J.; Zbrzezna, V.; Williams, K.;
Gulati,

[191538]S.; Pogo, A. O.: Cloning of glycoprotein D cDNA, which
encodes the

[191539]major subunit of the Duffy blood group system and the
receptor for

[191540]the Plasmodium vivax malaria parasite. Proc. Nat. Acad.
Sci. 90:

[191541]10793–10797, 1993.

[191542]

[191543]

[191544]20352.

[191545]Cook, P. J. L.; Page, B. M.; Johnston, A. W.; Stanford, W. K.;

[191546]Gavin, J.: Four further families informative for 1q and the
Duffy

[191547]blood group. Cytogenet. Cell Genet. 22: 378–380, 1978.

[191548]

[191549]

[191550]20353.

[191551]Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Jacobs, P. A.;
Polani,

[191552]P. E.: Segregation of genetic markers in families with
chromosome

[191553]polymorphisms and structural rearrangements involving
chromosome no.

[191554]1. Ann. Hum. Genet. 37: 261–274, 1974.

[191555]

[191556]

[191557]20354.

[191558]Crawford, M. N.; Punnett, H. H.; Carpenter, G. G.: Deletion
of

[191559]the long arm of chromosome 16 and an unexpected Duffy

blood group

[191560]phenotype reveal a possible autosomal linkage. Nature
215: 1075–1076,
[191561]1967.

[191562]

[191563]

[191564]20355.

[191565]Gelpi, A. P.; King, M. C.: Association of Duffy blood groups
[191566]with the sickle cell trait. Hum. Genet. 32: 65–68, 1976.

[191567]

[191568]

[191569]20356.

[191570]Hadley, T. J.; David, P. H.; McGinniss, M. H.; Miller, L. H.:

[191571]Identification of an erythrocyte component carrying the
Duffy blood

[191572]group Fy–a antigen. Science 223: 597–599, 1984.

[191573]

[191574]

[191575]20357.

[191576]Hadley, T. J.; Peiper, S. C.: From malaria to chemokine receptor:

[191577]the emerging physiologic role of the Duffy blood group antigen. Blood 89:

[191578]3077–3091, 1997.

[191579]

[191580]

[191581]20358.

[191582]Hamblin, M. T.; Di Rienzo, A.: Detection of the signature of

[191583]natural selection in humans: evidence from the Duffy blood group locus. Am.

[191584]J. Hum. Genet. 66: 1669–1679, 2000.

[191585]

[191586]

[191587]20359.

[191588]Hamblin, M. T.; Thompson, E. E.; Di Rienzo, A.: Complex signatures

[191589]of natural selection at the Duffy blood group locus. Am. J.

Hum.

[191590]Genet. 70: 369–383, 2002.

[191591]

[191592]

[191593]20360.

[191594]Hessner, M. J.; Pircon, R. A.; Johnson, S. T.; Luhm, R. A.:

Prenatal

[191595]genotyping of the Duffy blood group system by allele-specific polymerase

[191596]chain reaction. Prenatal Diag. 19: 41–45, 1999.

[191597]

[191598]

[191599]20361.

[191600]Horuk, R.; Chitnis, C. E.; Darbonne, W. C.; Colby, T. J.; Ry-

bicki,

[191601]A.; Hadley, T. J.; Miller, L. H.: A receptor for the malarial parasite

[191602]Plasmodium vivax: the erythrocyte chemokine receptor.

Science 261:

[191603]1182–1184, 1993.

[191604]

[191605]

[191606]20362.

[191607]Howard, P. N.; Stoddard, G. R.; Goddard, M. W.; Seely, J.
R.:

[191608]Giemsa banding of chromosome 1qh+ and linkage analysis. J. Med. Genet. 12:

[191609]44–48, 1975.

[191610]

[191611]

[191612]20363.

[191613]Iwamoto, S.; Li, J.; Sugimoto, N.; Okuda, H.; Kajii, E.: Characterization

[191614]of the Duffy gene promoter: evidence for tissue-specific abolishment

[191615]of expression in Fy(a–b–) of black individuals. Biochem. Biophys.

[191616]Res. Commun. 222: 852–859, 1996.

[191617]

[191618]

[191619]20364.

[191620]Lautenberger, J. A.; Stephens, J. C.; O'Brien, S. J.; Smith, M.

[191621]W.: Significant admixture linkage disequilibrium across 30 cM around

[191622]the FY locus in African Americans. *Am. J. Hum. Genet.* 66: 969–978,

[191623]2000.

[191624]

[191625]

[191626]20365.

[191627]Lee, C. S. N.; Ying, K. L.; Bowen, P.: Position of the Duffy

[191628]locus on chromosome 1 in relation to breakpoints for structural rearrangements. *Am.*

[191629]*J. Hum. Genet.* 26: 93–102, 1974.

[191630]

[191631]

[191632]20366.

[191633]Li, J.; Iwamoto, S.; Sugimoto, N.; Okuda, H.; Kajii, E.: Dinucleotide

[191634]repeat in the 3-prime flanking region provides a clue to the molecular

[191635]evolution of the Duffy gene. *Hum. Genet.* 99: 573–577,

1997.

[191636]

[191637]

[191638]20367.

[191639]Braun, A.; Kammerer, S.; Bohme, E.; Muller, B.; Roscher, A.

A.

[191640]: Identification of polymorphic sites of the human
bradykinin B(2)

[191641]receptor gene. Biochem. Biophys. Res. Commun. 211:
234–240, 1995.

[191642]

[191643]

[191644]20368.

[191645]Erdmann, J.; Hegemann, N.; Weidemann, A.; Kallisch, H.;
Hummel,

[191646]M.; Hetzer, R.; Fleck, E.; Regitz–Zagrosek, V.: Screening
the human

[191647]bradykinin B2 receptor gene in patients with cardiovascu-
lar diseases:

[191648]identification of a functional mutation in the promoter and
a new

[191649]coding variant (T21M). Am. J. Med. Genet. 80: 521–525,
1998.

[191650]

[191651]

[191652]20369.

[191653]Hess, J. F.; Borkowski, J. A.; Young, G. S.; Strader, C. D.;
Ransom,

[191654]R. W.: Cloning and pharmacological characterization of a
human bradykinin

[191655](BK–2) receptor. Biochem. Biophys. Res. Commun. 184:
260–268, 1992.

[191656]

[191657]

[191658]20370.

[191659]Kammerer, S.; Braun, A.; Arnold, N.; Roscher, A. A.: The
human

[191660]bradykinin B(2) receptor gene: full length cDNA, genomic
organization

[191661]and identification of the regulatory region. Biochem. Bio-
phys. Res.

[191662]Commun. 211: 226–233, 1995.

[191663]

[191664]

[191665]20371.

[191666]Ma, J.; Wang, D.; Ward, D. C.; Chen, L.; Dessai, T.; Chao, J.;

[191667]Chao, L.: Structure and chromosomal localization of the
gene (BDKRB2)

[191668]encoding human bradykinin B2 receptor. Genomics 23:
362–369, 1994.

[191669]

[191670]

[191671]20372.

[191672]Powell, S. J.; Slynn, G.; Thomas, C.; Hopkins, B.; Briggs, I.;

[191673]Graham, A.: Human bradykinin B2 receptor: nucleotide
sequence analysis

[191674]and assignment to chromosome 14. Genomics 15:
435–438, 1993.

[191675]

[191676]

[191677]20373.

[191678]Taketo, M.; Yokoyama, S.; Rochelle, J.; Kimura, S.; Higashida,

[191679]H.; Taketo, M.; Seldin, M. F.: Mouse B2 bradykinin receptor gene

[191680]maps to distal chromosome 12. Genomics 27: 222–223, 1995.

[191681]

[191682]

[191683]20374.

[191684]Chen, H.; Weber, A. J.: BDNF enhances retinal ganglion cell survival

[191685]in cats with optic nerve damage. Invest. Ophthalm. Vis. Sci. 42:

[191686]966–974, 2001.

[191687]

[191688]

[191689]20375.

[191690]Conover, J. C.; Erickson, J. T.; Katz, D. M.; Bianchi, L. M.; Poueymirou,

[191691]W. T.; McClain, J.; Pan, L.; Helgren, M.; Ip, N. Y.; Boland, P.; Friedman,

- [191692]B.; Wiegand, S.; Vejsada, R.; Kato, A. C.; DeChiara, T. M.;
Yancopoulos,
[191693]G. D.: Neuronal deficits, not involving motor neurons, in
mice lacking
[191694]BDNF and/or NT4. Nature 375: 235–238, 1995.
- [191695]
- [191696]
- [191697]20376.
- [191698]Hanson, I. M.; Seawright, A.; van Heyningen, V.: The hu-
man BDNF
[191699]gene maps between FSHB and HVBS1 at the boundary of
11p13–p14. Genomics 13:
[191700]1331–1333, 1992.
- [191701]
- [191702]
- [191703]20377.
- [191704]Hofer, M. M.; Barde, Y.–A.: Brain–derived neurotrophic
factor
[191705]prevents neuronal death in vivo. Nature 331: 261–262,
1988.

[191706]

[191707]

[191708]20378.

[191709]Huang, Z. J.; Kirkwood, A.; Pizzorusso, T.; Porciatti, V.;
Morales,

[191710]B.; Bear, M. F.; Maffei, L.; Tonegawa, S.: BDNF regulates
the maturation

[191711]of inhibition and the critical period of plasticity in mouse
visual

[191712]cortex. Cell 98: 739–755, 1999.

[191713]

[191714]

[191715]20379.

[191716]Kovalchuk, Y.; Hanse, E.; Kafitz, K. W.; Konnerth, A.: Post-
synaptic

[191717]induction of BDNF-mediated long-term potentiation. Sci-
ence 295:

[191718]1729–1734, 2002.

[191719]

[191720]

[191721]20380.

[191722]Liu, X.; Ernfors, P.; Wu, H.; Jaenisch, R.: Sensory but not
motor

[191723]neuron deficits in mice lacking NT4 and BDNF. Nature
375: 238–241,

[191724]1995.

[191725]

[191726]

[191727]20381.

[191728]Lu, Q.; Sun, E. E.; Klein, R. S.; Flanagan, J. G.: Ephrin–B re-
verse

[191729]signaling is mediated by a novel PDZ–RGS protein and se-
lectively inhibits

[191730]G protein–coupled chemoattraction. Cell 105: 69–79,
2001.

[191731]

[191732]

[191733]20382.

[191734]Masi, L.; Becherini, L.; Colli, E.; Gennari, L.; Mansani, R.;
Falchetti,

[191735]A.; Becorpi, A. M.; Cepollaro, C.; Gonnelli, S.; Tanini, A.;
Brandi,

[191736]M. L.: Polymorphisms of the calcitonin receptor gene are
associated

[191737]with bone mineral density in postmenopausal Italian
women. Biochem.

[191738]Biophys. Res. Commun. 248: 190–195, 1998.

[191739]

[191740]

[191741]20383.

[191742]Masi, L.; Becherini, L.; Gennari, L.; Colli, E.; Mansani, R.;
Falchetti,

[191743]A.; Cepollaro, C.; Gonnelli, S.; Tanini, A.; Brandi, M. L.: Al-
lelic

[191744]variants of human calcitonin receptor: distribution and as-
sociation

[191745]with bone mass in postmenopausal Italian women.
Biochem. Biophys.

[191746]Res. Commun. 245: 622–626, 1998.

[191747]

[191748]

[191749]20384.

[191750]Nakamura, M.; Zhang, Z. Q.; Shan, L.; Hisa, T.; Sasaki, M.;

Tsukino,

[191751]R.; Yokoi, T.; Kaname, A.; Kakudo, K.: Allelic variants of
human

[191752]calcitonin receptor in the Japanese population. Hum.

Genet. 99:

[191753]38–41, 1997.

[191754]

[191755]

[191756]20385.

[191757]Perez Jurado, L. A.; Li, X.; Francke, U.: The human calci-

tonin

[191758]receptor gene (CALCR) at 7q21.3 is outside the deletion
associated

[191759]with the Williams syndrome. Cytogenet. Cell Genet. 70:

246–249,

[191760]1995.

[191761]

[191762]

[191763]20386.

[191764]Taboulet, J.; Frenco, J. L.; Delage-Murroux, R.; Pichaud, F.;
de

[191765]Vernejoul, M. C.; Jullienne, A.: Evidence for 2 allelic forms
of

[191766]calcitonin receptor gene: distribution in normal and os-
teoporotic

[191767]women. (Abstract) J. Bone Miner. Res. 11 (suppl. 1): S204,
1996.

[191768]

[191769]

[191770]20387.

[191771]Taboulet, J.; Frenkian, M.; Frenco, J. L.; Feingold, N.; Julli-
enne,

[191772]A.; de Vernejoul, M. C.: Calcitonin receptor polymorphism
is associated

[191773]with a decreased fracture risk in post-menopausal
women. Hum. Molec.

[191774]Genet. 7: 2129-2133, 1998.

[191775]

[191776]

[191777]20388.

[191778]Amara, S. G.; Arriza, J. L.; Leff, S. E.; Swanson, L. W.;
Evans,

[191779]R. M.; Rosenfeld, M. G.: Expression in brain of a messenger RNA encoding

[191780]a novel neuropeptide homologous to calcitonin gene-related peptide.

[191781]Science 229: 1094–1097, 1985.

[191782]

[191783]

[191784]20389.

[191785]Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.; Geurts
van

[191786]Kessel, A. H. M.; Baylin, S. B.; Nelkin, B. D.; Jansz, H. S.;
Lips,

[191787]C. J. M.: The second human calcitonin/CGRP gene is located on chromosome

[191788]11. Hum. Genet. 70: 259–263, 1985.

[191789]

[191790]

[191791]20390.

[191792]Steenbergh, P. H.; Hoppener, J. W. M.; Zandberg, J.; Lips, C. J.

[191793]M.; Jansz, H. S.: A second human calcitonin/CGRP gene. FEBS Lett. 183:

[191794]403–407, 1985.

[191795]

[191796]

[191797]20391.

[191798]Arthur, J. S. C.; Elce, J. S.; Hegadorn, C.; Williams, K.; Greer,

[191799]P. A.: Disruption of the murine calpain small subunit gene, Capn4:

[191800]calpain is essential for embryonic development but not for cell growth

[191801]and division. Molec. Cell. Biol. 20: 4474–4481, 2000.

[191802]

[191803]

[191804]20392.

[191805]Ohno, S.; Emori, Y.; Suzuki, K.: Nucleotide sequence of a

cDNA

[191806]coding for the small subunit of human calcium-dependent protease. Nucleic

[191807]Acids Res. 14: 5559 only, 1986.

[191808]

[191809]

[191810]20393.

[191811]Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Ohmi-Imajoh,

[191812]S.; Suzuki, K.; Shimizu, Y.; Shimizu, N.: Four genes for the calpain

[191813]family locate on four distinct human chromosomes.

(Abstract) Cytogenet.

[191814]Cell Genet. 51: 1054-1055, 1989.

[191815]

[191816]

[191817]20394.

[191818]Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Shimizu, Y.;

[191819]Ohmi-Imajoh, S.; Shimizu, N.; Suzuki, K.: Four genes for

the calpain

[191820]family locate on four distinct human chromosomes. Cyto-
genet. Cell

[191821]Genet. 53: 225–229, 1990.

[191822]

[191823]

[191824]20395.

[191825]Sakihama, T.; Kakidani, H.; Zenita, K.; Yumoto, N.; Kikuchi,
T.;

[191826]Sasaki, T.; Kannagi, R.; Nakanishi, S.; Ohmori, M.; Takio,
K.; Titani,

[191827]K.; Murachi, T.: A putative Ca^{2+} –binding protein: struc-
ture of

[191828]the light subunit of porcine calpain elucidated by molecu-
lar cloning

[191829]and protein sequence analysis. Proc. Nat. Acad. Sci. 82:
6075–6079,

[191830]1985.

[191831]

[191832]

[191833]20396.

[191834]Berchtold, M. W.; Egli, R.; Rhyner, J. A.; Hameister, H.;
Strehler,

[191835]E. E.: Localization of the human bona fide calmodulin
genes CALM1,

[191836]CALM2, and CALM3 to chromosomes 14q24–q31,
2p21.1–p21.3, and 19q13.2–q13.3. Genomics 16:

[191837]461–465, 1993.

[191838]

[191839]

[191840]20397.

[191841]Chin, D.; Winkler, K. E.; Means, A. R.: Characterization of
substrate

[191842]phosphorylation and use of calmodulin mutants to ad-
dress implications

[191843]from the enzyme crystal structure of calmodulin–depen-
dent protein

[191844]kinase I. J. Biol. Chem. 272: 31235–31240, 1997.

[191845]

[191846]

[191847]20398.

[191848]Drum, C. L.; Yan, S.-Z.; Bard, J.; Shen, Y.-Q.; Lu, D.; Soe-laiman,

[191849]S.; Grabarek, Z.; Bohm, A.; Tang, W.-J.: Structural basis for the

[191850]activation of anthrax adenylyl cyclase exotoxin by calmodulin. Nature 415:

[191851]396–402, 2002.

[191852]

[191853]

[191854]20399.

[191855]Koller, M.; Schnyder, B.; Strehler, E. E.: Structural organization

[191856]of the human CaMIII calmodulin gene. Biochim. Biophys. Acta 1087:

[191857]180–189, 1990.

[191858]

[191859]

[191860]20400.

[191861]Kretsinger, R. H.; Rudnick, S. E.; Weissman, L. J.: Crystal structure

[191862]of calmodulin. J. Inorganic Biochem. 28: 289–302, 1986.

[191863]

[191864]

[191865]20401.

[191866]McPherson, J. D.; Hickie, R. A.; Wasmuth, J. J.; Meyskens,
F. L.;

[191867]Perham, R. N.; Strehler, E. E.; Graham, M. T.: Chromosomal
localization

[191868]of multiple genes encoding calmodulin. (Abstract) Cyto-
genet. Cell

[191869]Genet. 58: 1951 only, 1991.

[191870]

[191871]

[191872]20402.

[191873]Pegues, J. C.; Friedberg, F.: Multiple mRNAs encoding hu-
man calmodulin. Biochem.

[191874]Biophys. Res. Commun. 172: 1145–1149, 1990.

[191875]

[191876]

[191877]20403.

[191878]Rhyner, J. A.; Ottiger, M.; Wicki, R.; Greenwood, T. M.;
Strehler,

[191879]E. E.: Structure of the human CALM1 calmodulin gene and
identification

[191880]of two CALM1-related pseudogenes CALM1P1 and
CALM1P2. Europ. J. Biochem. 225:

[191881]71–82, 1994.

[191882]

[191883]

[191884]20404.

[191885]Scambler, P. J.; McPherson, M. A.; Bates, G.; Bradbury, N.
A.;

[191886]Dormer, R. L.; Williamson, R.: Biochemical and genetic ex-
clusion

[191887]of calmodulin as the site of the basic defect in cystic fi-
brosis. Hum.

[191888]Genet. 76: 278–282, 1987.

[191889]

[191890]

[191891]20405.

[191892]Schumacher, M. A.; Rivard, A. F.; Bachinger, H. P.; Adelman, J.

[191893]P.: Structure of the gating domain of a $\text{Ca}(2+)$ -activated K^+ channel

[191894]complexed with $\text{Ca}(2+)/$ calmodulin. Nature 410: 1120–1124, 2001.

[191895]

[191896]

[191897]20406.

[191898]Sen Gupta, B.; Detera-Wadleigh, S. D.; McBride, O. W.; Friedberg,

[191899]F.: A calmodulin pseudogene on human chromosome 17. Nucleic Acids

[191900]Res. 17: 2868 only, 1989.

[191901]

[191902]

[191903]20407.

[191904]Powers, P. A.; Liu, S.; Hogan, K.; Gregg, R. G.: Molecular characterization

[191905]of the gene encoding the gamma subunit of the human skeletal muscle

[191906]1,4-dihydropyridine-sensitive Ca(2+) channel (CACNLG), cDNA sequence,

[191907]gene structure, and chromosomal location. J. Biol. Chem. 268: 9275–9279,

[191908]1993.

[191909]

[191910]

[191911]20408.

[191912]Ambartsumian, N.; Tarabykina, S.; Grigorian, M.; Tulchinsky, E.;

[191913]Hulgaard, E.; Georgiev, G.; Lukanidin, E.: Characterization of two

[191914]splice variants of metastasis-associated human mts1 gene. Gene 159:

[191915]125–130, 1995.

[191916]

[191917]

[191918]20409.

[191919]Jackson-Grusby, L. L.; Swiergiel, J.; Linzer, D. I.: A growth-related

[191920]mRNA in cultured mouse cells encodes a placental calcium binding protein.

[191921]Nucleic Acids Res. 15: 6677-6690, 1987.

[191922]

[191923]

[191924]20410.

[191925]Iles, D. E.; Segers, B.; Weghuis, D. O.; Suikerbuijk, R.; Wieringa,

[191926]B.: Localization of the gamma-subunit of the skeletal muscle L-type

[191927]voltage-dependent calcium channel gene (CACNLG) to human chromosome

[191928]band 17q24 by in situ hybridization and identification of a polymorphic

[191929]repetitive DNA sequence at the gene locus. Cytogenet. Cell Genet. 64:

[191930]227-230, 1993.

[191931]

[191932]

[191933]20411.

[191934]Saris, C. J.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.;

Noonan,

[191935]D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and tissue
distribution

[191936]of the mRNA for bovine and murine p11, the S100-related
light chain

[191937]of the protein-tyrosine kinase substrate p36 (calpactin I).

J. Biol.

[191938]Chem. 262: 10663-10671, 1987.

[191939]

[191940]

[191941]20412.

[191942]Lefort, A.; Lecocq, R.; Libert, F.; Lamy, F.; Swillens, S.;

Vassart,

[191943]G.; Dumont, J. E.: Cloning and sequencing of a calcium-
binding protein

[191944]regulated by cyclic AMP in the thyroid. EMBO J. 8:

111-116, 1989.

[191945]

[191946]

[191947]20413.

[191948]Loughlin, J.; Irlen, C.; Sykes, B.: Exclusion of the cartilage

[191949]link protein and the cartilage matrix protein genes as the
mutant

[191950]loci in several heritable chondrodysplasias. Hum. Genet.

94: 698–700,

[191951]1994.

[191952]

[191953]

[191954]20414.

[191955]Osborne–Lawrence, S. L.; Sinclair, A. K.; Hicks, R. C.;

Lacey,

[191956]S. W.; Eddy, R. L., Jr.; Byers, M. G.; Shows, T. B.; Duby, A.

D.:

[191957]Complete amino acid sequence of human cartilage link
protein (CRTL1)

[191958]deduced from cDNA clones and chromosomal assignment
of the gene. Genomics 8:

[191959]562–567, 1990.

[191960]

[191961]

[191962]20415.

[191963]Watanabe, H.; Yamada, Y.: Mice lacking link protein develop dwarfism

[191964]and craniofacial abnormalities. Nature Genet. 21: 225–229, 1999.

[191965]

[191966]

[191967]20416.

[191968]Deak, F.; Piecha, D.; Bachrati, C.; Paulsson, M.; Kiss, I.: Primary

[191969]structure and expression of matrilin–2, the closest relative of cartilage

[191970]matrix protein within the von Willebrand factor type A–like module

[191971]superfamily. J. Biol. Chem. 272: 9268–9274, 1997.

[191972]

[191973]

[191974]20417.

[191975]Deak, F.; Wagener, R.; Kiss, I.; Paulsson, M.: The matrilins:

[191976]a novel family of oligomeric extracellular matrix proteins.

Matrix

[191977]Biol. 18: 55–64, 1999.

[191978]

[191979]

[191980]20418.

[191981]Hansson, A.–S.; Heinegard, D.; Holmdahl, R.: A new animal model

[191982]for relapsing polychondritis, induced by cartilage matrix protein

[191983](matrilin–1). J. Clin. Invest. 104: 589–598, 1999.

[191984]

[191985]

[191986]20419.

[191987]Jenkins, R. N.; Osborne–Lawrence, S. L.; Sinclair, A. K.; Eddy,

[191988]R. L., Jr.; Byers, M. G.; Shows, T. B.; Duby, A. D.: Structure and

[191989]chromosomal location of the human gene encoding cartilage matrix protein. J.

[191990]Biol. Chem. 265: 19624–19631, 1990.

[191991]

[191992]

[191993]20420.

[191994]Lang, B.; Rothenfusser, A.; Lanchbury, J. S.; Rauh, G.;
Breedveld,

[191995]F. C.; Urlacher, A.; Albert, E. D.; Peter, H.–H.; Melchers, I.:
Susceptibility

[191996]to relapsing polychondritis is associated with HLA–DR4.
Arthritis

[191997]Rheum. 36: 660–664, 1993.

[191998]

[191999]

[192000]20421.

[192001]Muratoglu, S.; Krysan, K.; Balazs, M.; Sheng, H.; Zakany,
R.; Modis,

[192002]L.; Kiss, I.; Deak, F.: Primary structure of human matrilin–
2, chromosome

[192003]location of the MATN2 gene and conservation of an AT–
AC intron in

[192004]matrilin genes. Cytogenet. Cell Genet. 90: 323–327, 2000.

[192005]

[192006]

[192007]20422.

[192008]Wagener, R.; Kobbe, B.; Paulsson, M.: Primary structure of
matrilin-3,

[192009]a new member of a family of extracellular matrix proteins
related

[192010]to cartilage matrix protein (matrilin-1) and von Willebrand
factor. FEBS

[192011]Lett. 413: 129–134, 1997.

[192012]

[192013]

[192014]20423.

[192015]Zeuner, M.; Straub, R. H.; Rauh, G.; Albert, E. D.;
Scholmerich,

[192016]J.; Lang, B.: Relapsing polychondritis: clinical and im-
munogenetic

[192017]analysis of 62 patients. J. Rheum. 24: 96–101, 1997.

[192018]

[192019]

[192020]20424.

[192021]Boldyreff, B.; Klett, C.; Gottert, E.; Geurts van Kessel, A.;
Hameister,

[192022]H.; Issinger, O.-G.: Assignment of casein kinase 2 alpha
sequences

[192023]to two different human chromosomes. Hum. Genet. 89:
79–82, 1992.

[192024]

[192025]

[192026]20425.

[192027]Doray, B.; Ghosh, P.; Griffith, J.; Geuze, H. J.; Kornfeld, S.

[192028]: Cooperation of GGAs and AP–1 in packaging MPRs at the
trans–Golgi

[192029]network. Science 297: 1700–1703, 2002.

[192030]

[192031]

[192032]20426.

[192033]Meisner, H.; Heller–Harrison, R.; Buxton, J.; Czech, M. P.:

Molecular

[192034]cloning of the human casein kinase II alpha subunit. Bio-chemistry 28:

[192035]4072–4076, 1989.

[192036]

[192037]

[192038]20427.

[192039]Wirkner, U.; Voss, H.; Lichter, P.; Ansorge, W.; Pyerin, W.:
The

[192040]human gene (CSNK2A1) coding for the casein kinase II
subunit alpha

[192041]is located on chromosome 20 and contains tandemly ar-
ranged Alu repeats. Genomics 19:

[192042]257–265, 1994.

[192043]

[192044]

[192045]20428.

[192046]Wirkner, U.; Voss, H.; Lichter, P.; Weitz, S.; Ansorge, W.;
Pyerin,

[192047]W.: Human casein kinase II subunit alpha: sequence of a

processed

[192048](pseudo)gene and its localization on chromosome 11.

Biochim. Biophys.

[192049]Acta 1131: 220–222, 1992.

[192050]

[192051]

[192052]20429.

[192053]Yang–Feng, T. L.; Zheng, K.; Kopatz, I.; Naiman, T.;

Canaani, D.

[192054]: Mapping of the human casein kinase II catalytic subunit

genes: two

[192055]loci carrying the homologous sequences for the alpha

subunit. Nucleic

[192056]Acids Res. 19: 7125–7129, 1991.

[192057]

[192058]

[192059]20430.

[192060]Lozeman, F. J.; Litchfield, D. W.; Piening, C.; Takio, K.;

Walsh,

[192061]K. A.; Krebs, E. G.: Isolation and characterization of human

cDNA

[192062]clones encoding the alpha and the alpha-prime subunits
of casein kinase

[192063]II. Biochemistry 29: 8436–8447, 1990.

[192064]

[192065]

[192066]20431.

[192067]Xu, X.; Toselli, P. A.; Russell, L. D.; Seldin, D. C.: Globo-
zoospermia

[192068]in mice lacking the casein kinase II alpha-prime catalytic
subunit. Nature

[192069]Genet. 23: 118–121, 1999.

[192070]

[192071]

[192072]20432.

[192073]Abrieu, A.; Kahana, J. A.; Wood, K. W.; Cleveland, D. W.:
CENP-E

[192074]as an essential component of the mitotic checkpoint in
vitro. Cell 102:

[192075]817–826, 2000.

[192076]

[192077]

[192078]20433.

[192079]Wood, K. W.; Sakowicz, R.; Goldstein, L. S. B.; Cleveland, D. W.

[192080]: CENP-E is a plus end-directed kinetochore motor required for metaphase

[192081]chromosome alignment. Cell 97: 357-366, 1997.

[192082]

[192083]

[192084]20434.

[192085]Yao, X.; Abrieu, A.; Zheng, Y.; Sullivan, K. F.; Cleveland, D.

[192086]W.: CENP-E forms a link between attachment of spindle microtubules

[192087]to kinetochores and the mitotic checkpoint. Nature Cell Biol. 2:

[192088]484-491, 2000.

[192089]

[192090]

[192091]20435.

[192092]Yen, T. J.; Compton, D. A.; Wise, D.; Zinkowski, R. P.;

Brinkley,

[192093]B. R.; Earnshaw, W. C.; Cleveland, D. W.: CENP-E, a novel
human centromere-associated

[192094]protein required for progression from metaphase to
anaphase. EMBO

[192095]J. 10: 1245-1254, 1991.

[192096]

[192097]

[192098]20436.

[192099]Yen, T. J.; Li, G.; Schaar, B. T.; Szilak, I.; Cleveland, D. W.

[192100]: CENP-E is a putative kinetochore motor that accumulates
just before

[192101]mitosis. Nature 359: 536-539, 1992.

[192102]

[192103]

[192104]20437.

[192105]Geissler, E. N.; Liao, M.; Brook, J. D.; Martin, F. H.; Zsebo,

[192106]K. M.; Housman, D. E.; Galli, S. J.: Stem cell factor (SCF), a novel

[192107]hematopoietic growth factor and ligand for c-kit tyrosine kinase receptor,

[192108]maps on human chromosome 12 between 12q14.3 and 12qter. Somat. Cell

[192109]Molec. Genet. 17: 207–214, 1991.

[192110]

[192111]

[192112]20438.

[192113]Heissig, B.; Hattori, K.; Dias, S.; Friedrich, M.; Ferris, B.;

[192114]Hackett, N. R.; Crystal, R. G.; Besmer, P.; Lyden, D.; Moore, M. A.

[192115]S.; Werb, Z.; Rafii, S.: Recruitment of stem and progenitor cells

[192116]from the bone marrow niche requires MMP–9 mediated release of Kit–ligand. Cell 109:

[192117]625–637, 2002.

[192118]

[192119]

[192120]20439.

[192121]Martin, F. H.; Suggs, S. V.; Langley, K. E.; Lu, H. S.; Ting, J.;

[192122]Okino, K. H.; Morris, C. F.; McNiece, I. K.; Jacobsen, F. W.;

Mendez,

[192123]E. A.; Birkett, N. C.; Smith, K. A.; and 15 others: Primary

structure

[192124]and functional expression of rat and human stem cell fac-

tor DNAs. Cell 63:

[192125]203-211, 1990.

[192126]

[192127]

[192128]20440.

[192129]Mathew, S.; Murty, V. V. V. S.; Hunziker, W.; Chaganti, R. S.

K.

[192130]: Subregional mapping of 13 single-copy genes on the

long arm of chromosome

[192131]12 by fluorescence in situ hybridization. Genomics 14:

775-779,

[192132]1992.

[192133]

[192134]

[192135]20441.

[192136]Li, X.-Y.; Mattei, M. G.; Zaleska-Rutczynska, Z.; Hooft van
Huijsduijnen,

[192137]R.; Figueroa, F.; Nadeau, J.; Benoist, C.; Mathis, D.: One
subunit

[192138]of the transcription factor NF- κ B maps close to the major
histocompatibility

[192139]complex in murine and human chromosomes. Genomics
11: 630-634, 1991.

[192140]

[192141]

[192142]20442.

[192143]Johnston, J.; Bollekens, J.; Allen, R. H.; Berliner, N.: Struc-
ture

[192144]of the cDNA encoding transcobalamin I, a neutrophil
granule protein. J.

[192145]Biol. Chem. 264: 15754-15757, 1989.

[192146]

[192147]

[192148]20443.

[192149]Johnston, J.; Yang-Feng, T.; Berliner, N.: Genomic structure and

[192150]mapping of the chromosomal gene for transcobalamin I (TCN1): comparison

[192151]to human intrinsic factor. Genomics 12: 459-464, 1992.

[192152]

[192153]

[192154]20444.

[192155]Yang-Feng, T. L.; Berliner, N.; Deverajan, P.; Johnston, J.: Assignment

[192156]of two human neutrophil secondary granule protein genes, transcobalamin

[192157]I and neutrophil collagenase to chromosome 11. (Abstract) Cytogenet.

[192158]Cell Genet. 58: 1974 only, 1991.

[192159]

[192160]

[192161]20445.

[192162]Courey, A. J.; Holtzman, D. A.; Jackson, S. P.; Tjian, R.:

Synergistic

[192163]activation by the glutamine-rich domains of human transcription factor

[192164]Sp1. Cell 59: 827–836, 1989.

[192165]

[192166]

[192167]20446.

[192168]Gaynor, R. B.; Shieh, B.-H.; Klisak, I.; Sparkes, R. S.; Lusi,

[192169]A. J.: Localization of the transcription factor SP1 gene to human

[192170]chromosome 12q12–q13.2. Cytogenet. Cell Genet. 64: 210–212, 1993.

[192171]

[192172]

[192173]20447.

[192174]Kadonaga, J. T.; Carner, K. R.; Masiarz, F. R.; Tjian, R.: Isolation

[192175]of a cDNA encoding transcription factor Sp1 and functional analysis

[192176]of the DNA binding domain. Cell 51: 1079–1090, 1987.

[192177]

[192178]

[192179]20448.

[192180]Matera, A. G.; Ward, D. C.: Localization of the human Sp1 transcription

[192181]factor gene to 12q13 by fluorescence in situ hybridization. Genomics 17:

[192182]793–794, 1993.

[192183]

[192184]

[192185]20449.

[192186]Szpirer, J.; Szpirer, C.; Riviere, M.; Levan, G.; Marynen, P.;

[192187]Cassiman, J.-J.; Wiese, R.; DeLuca, H. F.: The Sp1 transcription

[192188]factor gene (SP1) and the 1,25-dihydroxyvitamin D(3) receptor gene

[192189](VDR) are colocalized on human chromosome arm 12q and rat chromosome

[192190]

[192191]20450.

[192192]Genomics 11: 168–173, 1991.

[192193]

[192194]7. Wimmer, E. A.; Jackle, H.; Pfeifle, C.; Cohen, S. M.: A
Drosophila

[192195]homologue of human Sp1 is a head-specific segmentation
gene. Nature 366:

[192196]690–694, 1993.

[192197]

[192198]

[192199]20451.

[192200]Abbott, C.; Piaggio, G.; Ammendola, R.; Solomon, E.;
Povey, S.;

[192201]Gounari, F.; De Simone, V.; Cortese, R.: Mapping of the
gene TCF2

[192202]for the transcription factor LFB3 to human chromosome
17 by polymerase

[192203]chain reaction. Genomics 8: 165–167, 1990.

[192204]

[192205]

[192206]20452.

[192207]Bach, I.; Mattei, M.-G.; Cereghini, S.; Yaniv, M.: Two members

[192208]of an HNF1 homeoprotein family are expressed in human liver. Nucleic

[192209]Acids Res. 19: 3553–3559, 1991.

[192210]

[192211]

[192212]20453.

[192213]Bingham, C.; Ellard, S.; Allen, L.; Bulman, M.; Shepherd, M.; Frayling,

[192214]T.; Berry, P. J.; Clark, P. M.; Lindner, T.; Bell, G. I.; Ryffel,

[192215]G. U.; Nicholls, A. J.; Hattersley, A. T.: Abnormal nephron development

[192216]associated with a frameshift mutation in the transcription factor

[192217]hepatocyte nuclear factor–1–beta. Kidney Int. 57: 898–907, 2000.

[192218]

[192219]

[192220]20454.

[192221]Horikawa, Y.; Iwasaki, N.; Hara, M.; Furuta, H.; Hinokio, Y.;
Cockburn,

[192222]B. N.; Lindner, T.; Yamagata, K.; Ogata, M.; Tomonaga, O.;
Kuroki,

[192223]H.; Kasahara, T.; Iwamoto, Y.; Bell, G. I.: Mutation in hepa-
tocyte

[192224]nuclear factor-1-beta gene (TCF2) associated with MODY.
(Letter) Nature

[192225]Genet. 17: 384-385, 1997.

[192226]

[192227]

[192228]20455.

[192229]Menzel, R.; Kaisaki, P. J.; Rjasanowski, I.; Heinke, P.;
Kerner,

[192230]W.; Menzel, S.: A low renal threshold for glucose in dia-
betic patients

[192231]with a mutation in the hepatocyte nuclear factor-1-alpha
(HNF-1-alpha)

[192232]gene. Diabet. Med. 15: 816-820, 1998.

[192233]

[192234]

[192235]20456.

[192236]Nishigori, H.; Yamada, S.; Kohama, T.; Tomura, H.; Sho, K.; Horikawa,

[192237]Y.; Bell, G. I.; Takeuchi, T.; Takeda, J.: Frameshift mutation, A263fsinsGG,

[192238]in the hepatocyte nuclear factor-1-beta gene associated with diabetes

[192239]and renal dysfunction. Diabetes 47: 1354–1355, 1998.

[192240]

[192241]

[192242]20457.

[192243]Wild, W.; Pogge von Strandmann, E.; Nastos, A.; Senkel, S.; Lingott-Frieg,

[192244]A.; Bulman, M.; Bingham, C.; Ellard, S.; Hattersley, A. T.; Ryffel,

[192245]G. U.: The mutated human gene encoding hepatocyte nuclear factor

[192246]1-beta inhibits kidney formation in developing Xenopus embryos. Proc.

[192247]Nat. Acad. Sci. 97: 4695–4700, 2000.

[192248]

[192249]

[192250]20458.

[192251]Aplan, P. D.; Jones, C. A.; Chervinsky, D. S.; Zhao, X.;
Ellsworth,

[192252]M.; Wu, C.; McGuire, E. A.; Gross, K. W.: An scl gene prod-
uct lacking

[192253]the transactivation domain induces bony abnormalities
and cooperates

[192254]with LMO1 to generate T-cell malignancies in transgenic
mice. EMBO

[192255]J. 16: 2408–2419, 1997.

[192256]

[192257]

[192258]20459.

[192259]Aplan, P. D.; Lombardi, D. P.; Ginsberg, A. M.; Cossman, J.;
Bertness,

[192260]V. L.; Kirsch, I. L.: Disruption of the human SCL locus by
'illegitimate'

[192261]V-(D)-J recombinase activity. Science 250: 1426–1429,
1990.

[192262]

[192263]

[192264]20460.

[192265]Aplan, P. D.; Lombardi, D. P.; Reaman, G. H.; Sather, H. N.;
Hammond,

[192266]G. D.; Kirsch, I. R.: Involvement of the putative
hematopoietic transcription

[192267]factor SCL in T-cell acute lymphoblastic leukemia. Blood
79: 1327–1333,

[192268]1992.

[192269]

[192270]

[192271]20461.

[192272]Brown, L.; Cheng, J.-T.; Chen, Q.; Siciliano, M. J.; Crist, W.;

[192273]Buchanan, G.; Baer, R.: Site-specific recombination of the
tal-1

[192274]gene is a common occurrence in human T-cell leukemia.
EMBO J. 9:

[192275]3343–3351, 1990.

[192276]

[192277]

[192278]20462.

[192279]Collazo–Garcia, N.; Scherer, P.; Aplan, P. D.: Cloning and
characterization

[192280]of a murine SIL gene. Genomics 30: 506–513, 1995.

[192281]

[192282]

[192283]20463.

[192284]Izraeli, S.; Lowe, L. A.; Bertness, V. L.; Good, D. J.; Dor–
ward,

[192285]D. W.; Kirsch, I. R.; Kuehn, M. R.: The SIL gene is required
for

[192286]mouse embryonic axial development and left–right speci–
fication. Nature 399:

[192287]691–694, 1999.

[192288]

[192289]

[192290]20464.

[192291]Roy, A. L.; Du, H.; Gregor, P. D.; Novina, C. D.; Martinez,
E.;

[192292]Roeder, R. G.: Cloning of an Inr– and E–box binding pro–

tein, TFII-I,

[192293]that interacts physically and functionally with USF1. EMBO
J. 16:

[192294]7091–7104, 1997.

[192295]

[192296]

[192297]20465.

[192298]Steingrimsson, E.; Sawadogo, M.; Gilbert, D. J.; Zervos, A.
S.;

[192299]Brent, R.; Blonar, M. A.; Fisher, D. E.; Copeland, N. G.;
Jenkins,

[192300]N. A.: Murine chromosomal location of five bHLH-Zip
transcription

[192301]factor genes. Genomics 28: 179–183, 1995.

[192302]

[192303]

[192304]20466.

[192305]Zhong, G.; Fan, P.; Ji, H.; Dong, F.; Huang, Y.: Identifica-
tion

[192306]of a chlamydial protease-like activity factor responsible

for the

[192307]degradation of host transcription factors. J. Exp. Med.

193: 935–942,

[192308]2001.

[192309]

[192310]

[192311]20467.

[192312]Aasland, R.; Olsen, L. C.; Spurr, N. K.; Krokan, H. E.; Hel-
land,

[192313]D. E.: Chromosomal assignment of human uracil–DNA
glycosylase to

[192314]chromosome 12. Genomics 7: 139–141, 1990.

[192315]

[192316]

[192317]20468.

[192318]Dinner, A. R.; Blackburn, G. M.; Karplus, M.: Uracil–DNA
glycolase

[192319]acts by substrate autocatalysis. Nature 413: 752–755,
2001.

[192320]

[192321]

[192322]20469.

[192323]Haug, T.; Skorpen, F.; Kvaloy, K.; Eftedal, I.; Lund, H.;
Krokan,

[192324]H. E.: Human uracil–DNA glycosylase gene: sequence or-
ganization,

[192325]methylation pattern, and mapping to chromosome
12q23–q24.1. Genomics 36:

[192326]408–416, 1996.

[192327]

[192328]

[192329]20470.

[192330]Haug, T.; Skorpen, F.; Lund, H.; Krokan, H. E.: Structure of
the

[192331]gene for human uracil–DNA glycosylase and analysis of
the promoter

[192332]function. FEBS Lett. 353: 180–184, 1994.

[192333]

[192334]

[192335]20471.

- [192336]Meyer–Siegler, K.; Mauro, D. J.; Seal, G.; Wurzer, J.; deRiel,
[192337]J. K.; Sirover, M. A.: A human nuclear uracil DNA glycosy-
lase is
[192338]the 37–kDa subunit of glyceraldehyde–3–phosphate de-
hydrogenase. Proc.
[192339]Nat. Acad. Sci. 88: 8460–8464, 1991.
- [192340]
- [192341]
- [192342]20472.
- [192343]Nilsen, H.; Rosewell, I.; Robins, P.; Skjelbred, C. F.; Ander-
sen,
[192344]S.; Slupphaug, G.; Daly, G.; Krokan, H. E.; Lindahl, T.;
Barnes, D.
[192345]E.: Uracil–DNA glycosylase (UNG)–deficient mice reveal a
primary
[192346]role of the enzyme during DNA replication. Molec. Cell 5:
1059–1065,
[192347]2000.
- [192348]
- [192349]

[192350]20473.

[192351]Olsen, L. C.; Aasland, R.; Wittwer, C. U.; Krokan, H. E.;
Helland,

[192352]D. E.: Molecular cloning of human uracil–DNA glycosylase,
a highly

[192353]conserved DNA repair enzyme. EMBO J. 8: 3121–3125,
1989.

[192354]

[192355]

[192356]20474.

[192357]Vollberg, T. M.; Siegler, K. M.; Cool, B. L.; Sirover, M. A.:

[192358]Isolation and characterization of the human uracil DNA
glycosylase

[192359]gene. Proc. Nat. Acad. Sci. 86: 8693–8697, 1989.

[192360]

[192361]

[192362]20475.

[192363]Momeni, P.; Glockner, G.; Schmidt, O.; von Holtum, D.; Al-
brecht,

[192364]B.; Gillesen–Kaesbach, G.; Hennekam, R.; Meinecke, P.;

Zabel, B.;

[192365]Rosenthal, A.; Horsthemke, B.; Ludecke, H.-J.: Mutations
in a new

[192366]gene, encoding a zinc-finger protein, cause tricho-
rhino-phalangeal

[192367]syndrome type I. Nature Genet. 24: 71-74, 2000.

[192368]

[192369]

[192370]20476.

[192371]Komuro, I.; Wenninger, K. E.; Philipson, K. D.; Izumo, S.:
Molecular

[192372]cloning and characterization of the human cardiac
Na(+)/Ca(2+) exchanger

[192373]cDNA. Proc. Nat. Acad. Sci. 89: 4769-4773, 1992.

[192374]

[192375]

[192376]20477.

[192377]Kraev, A.; Chumakov, I.; Carafoli, E.: The organization of
the

[192378]human gene NCX1 encoding the sodium-calcium ex-

changer. Genomics 37:

[192379]105–112, 1996.

[192380]

[192381]

[192382]20478.

[192383]McDaniel, L. D.; Lederer, W. J.; Kofuji, P.; Schulze, D. H.;
Kieval,

[192384]R.; Schultz, R. A.: Mapping of the human cardiac
Na⁺/Ca²⁺ exchanger

[192385]gene (NCX1) by fluorescent in situ hybridization to chro-
mosome region

[192386]2p22–p23. Cytogenet. Cell Genet. 63: 192–193, 1993.

[192387]

[192388]

[192389]20479.

[192390]Shieh, B.–H.; Xia, Y.; Sparkes, R. S.; Klisak, I.; Lusk, A. J.;

[192391]Nicoll, D. A.; Philipson, K. D.: Mapping of the gene for the
cardiac

[192392]sarcolemmal Na⁽⁺⁾–Ca⁽²⁺⁾ exchanger to human chromo-
some 2p21–p23. Genomics 12:

[192393]616–617, 1992.

[192394]

[192395]

[192396]20480.

[192397]Wakimoto, K.; Kobayashi, K.; Kuro-o, M.; Yao, A.;
Iwamoto, T.;

[192398]Yanaka, N.; Kita, S.; Nishida, A.; Azuma, S.; Toyoda, Y.;
Omori, K.;

[192399]Imahie, H.; Oka, T.; Kudoh, S.; Kohmoto, O.; Yazaki, Y.;
Shigekawa,

[192400]M.; Imai, Y.; Nabeshima, Y.; Komura, I.: Targeted disruption of Na(+)/Ca(2+)

[192401]exchanger gene leads to cardiomyocyte apoptosis and defects in heartbeat. J.

[192402]Biol. Chem. 275: 36991–36998, 2000.

[192403]

[192404]

[192405]20481.

[192406]Chong, S. S.; Kristjansson, K.; Zoghbi, H. Y.; Hughes, M.
R.:

[192407]Molecular cloning of the cDNA encoding a human renal sodium phosphate

[192408]transport protein and its assignment to chromosome 6p21.3–p23. Genomics 18:

[192409]355–359, 1993.

[192410]

[192411]

[192412]20482.

[192413]Kos, C. H.; Tihy, F.; Murer, H.; Lemieux, N.; Tenenhouse, H. S.

[192414]: Comparative mapping of Na(+)-phosphate cotransporter genes, NPT1

[192415]and NPT2, in human and rabbit. Cytogenet. Cell Genet. 75: 22–24,

[192416]1996.

[192417]

[192418]

[192419]20483.

[192420]Zhang, X.-X.; Tenenhouse, H. S.; Hewson, A. S.; Eydoux, P.: Assignment

[192421]of renal specific Na(+)-phosphate cotransporter gene
Slc17a1 to mouse

[192422]chromosome bands 13A3-A4 by in situ hybridization. Cy-
togenet. Cell

[192423]Genet. 76: 180 only, 1997.

[192424]

[192425]

[192426]20484.

[192427]Ghishan, F. K.; Knobel, S.; Dasuki, M.; Butler, M.; Phillips,
J.

[192428]: Chromosomal localization of the human renal sodium
phosphate transporter

[192429]to chromosome 5: implications for X-linked hypophos-
phatemia. Pediat.

[192430]Res. 35: 510-513, 1994.

[192431]

[192432]

[192433]20485.

[192434]Hartmann, C. M.; Hewson, A. S.; Kos, C. H.; Hilfiker, H.;
Soumounou,

[192435]Y.; Murer, H.; Tenenhouse, H. S.: Structure of murine and human renal

[192436]type II Na(+)-phosphate cotransporter genes (Npt2 and NPT2). Proc.

[192437]Nat. Acad. Sci. 93: 7409–7414, 1996.

[192438]

[192439]

[192440]20486.

[192441]Kos, C. H.; Tihy, F.; Econs, M. J.; Murer, H.; Lemieux, N.; Tenenhouse,

[192442]H. S.: Localization of a renal sodium-phosphate cotransporter gene

[192443]to human chromosome 5q35. Genomics 19: 176–177, 1994.

[192444]

[192445]

[192446]20487.

[192447]Magagnin, S.; Werner, A.; Markovich, D.; Sorribas, V.; Stange,

[192448]G.; Biber, J.; Murer, H.: Expression cloning of human and rat renal

[192449]cortex Na/Pi cotransport. Proc. Nat. Acad. Sci. 90:
5979–5983, 1993.

[192450]

[192451]

[192452]20488.

[192453]McPherson, J. D.; Krane, M. C.; Wagner–McPherson, C. B.;
Kos, C.

[192454]H.; Tenenhouse, H. S.: High resolution mapping of the re-
nal sodium–phosphate

[192455]cotransporter gene (NPT2) confirms its localization to hu-
man chromosome

[192456]5q35. Pediat. Res. 41: 632–634, 1997.

[192457]

[192458]

[192459]20489.

[192460]Tenenhouse, H. S.; Werner, A.; Biber, J.; Ma, S.; Martel, J.;
Roy,

[192461]S.; Murer, H.: Renal Na(+)-phosphate cotransport in
murine X-linked

[192462]hypophosphatemic rickets: molecular characterization. J.
Clin. Invest. 93:

[192463]671–676, 1994.

[192464]

[192465]

[192466]20490.

[192467]Zhang, X.-X.; Tenenhouse, H. S.; Hewson, A. S.; Murer, H.;
Eydoux,

[192468]P.: Assignment of renal-specific Na(+)-phosphate co-
transporter gene

[192469]Slc17a2 to mouse chromosome band 13B by in situ hy-
bridization. Cytogenet.

[192470]Cell Genet. 77: 304–305, 1997.

[192471]

[192472]

[192473]20491.

[192474]Barnard, R.; Kelly, G.; Manzetti, S. O.; Harris, E. L.: Neither

[192475]the New Zealand genetically hypertensive strain nor Dahl
salt-sensitive

[192476]strain has an A1079T transversion in the alpha-1 isoform
of the Na(+),K(+)-ATPase

[192477]gene. Hypertension 38: 786–792, 2001.

[192478]

[192479]

[192480]20492.

[192481]Chehab, F. F.; Kan, Y. W.; Law, M. L.; Hartz, J.; Kao, F.-T.;
Blostein,

[192482]R.: Human placental Na⁺,K⁺-ATPase alpha subunit: cDNA
cloning, tissue

[192483]expression, DNA polymorphism, and chromosomal local-
ization. Proc.

[192484]Nat. Acad. Sci. 84: 7901-7905, 1987.

[192485]

[192486]

[192487]20493.

[192488]Dahl, L. K.; Heine, M.; Tassinari, L.: Role of genetic factors

[192489]in susceptibility to experimental hypertension due to
chronic excess

[192490]salt ingestion. Nature 194: 480-482, 1972.

[192491]

[192492]

[192493]20494.

[192494]Dahl, L. K.; Heine, M.; Thompson, K.: Genetic influence of the

[192495]kidneys on blood pressure: evidence from chronic renal homografts

[192496]in rats with opposite predispositions to hypertension.

Circ. Res. 40:

[192497]94–101, 1974.

[192498]

[192499]

[192500]20495.

[192501]Glorioso, N.; Filigheddu, F.; Troffa, C.; Soro, A.; Parpaglia,

[192502]P. P.; Tsikoudakis, A.; Myers, R. H.; Herrera, V. L. M.; Ruiz–Opazo,

[192503]N.: Interaction of alpha–1–Na,K–ATPase and Na,K,2Cl–cotransporter

[192504]genes in human essential hypertension. Hypertension 38: 204–209,

[192505]2001.

[192506]

[192507]

[192508]20496.

[192509]Herrera, V. L. M.; Ruiz-Opazo, N.: Alteration of alpha-1
Na⁺,K⁽⁺⁾-ATPase

[192510]86R-beta(+) influx by a single amino acid substitution.
Science 249:

[192511]1023-1026, 1990.

[192512]

[192513]

[192514]20497.

[192515]Andersson, S.; Russell, D. W.: Structural and biochemical
properties

[192516]of cloned and expressed human and rat steroid
5-alpha-reductases. Proc.

[192517]Nat. Acad. Sci. 87: 3640-3644, 1990.

[192518]

[192519]

[192520]20498.

[192521]Ellis, J. A.; Stebbing, M.; Harrap, S. B.: Genetic analysis of

[192522]male pattern baldness and the 5-alpha-reductase genes.

J. Invest.

[192523]Derm. 110: 849-853, 1998.

[192524]

[192525]

[192526]20499.

[192527]Harris, G.; Azzolina, B.; Baginsky, W.; Cimis, G.; Rasmus-
son, G.

[192528]H.; Tolman, R. L.; Raetz, C. R. H.; Ellsworth, K.: Identifica-
tion

[192529]and selective inhibition of an isozyme of steroid
5-alpha-reductase

[192530]in human scalp. Proc. Nat. Acad. Sci. 89: 10787-10791,
1992.

[192531]

[192532]

[192533]20500.

[192534]Hsieh, C.-L.; Milatovich, A.; Russell, D.; Francke, U.: Chro-
mosomal

[192535]mapping of human steroid 5 alpha-reductase gene
(SRD5A1) and pseudogene

[192536](SRD5AP1) in human and mouse. (Abstract) Cytogenet.

Cell Genet. 58:

[192537]1897 only, 1991.

[192538]

[192539]

[192540]20501.

[192541]Jenkins, E. P.; Andersson, S.; Imperato-McGinley, J.; Wilson, J.

[192542]D.; Russell, D. W.: Genetic and pharmacological evidence for more

[192543]than one human steroid 5-alpha-reductase. J. Clin. Invest. 89: 293-300,

[192544]1992.

[192545]

[192546]

[192547]20502.

[192548]Jenkins, E. P.; Hsieh, C.-L.; Milatovich, A.; Normington, K.; Berman,

[192549]D. M.; Francke, U.; Russell, D. W.: Characterization and chromosomal

[192550]mapping of a human steroid 5-alpha-reductase gene and
pseudogene and

[192551]mapping of the mouse homologue. Genomics 11:
1102-1112, 1991.

[192552]

[192553]

[192554]20503.

[192555]Thigpen, A. E.; Silver, R. I.; Guileyardo, J. M.; Casey, M. L.;

[192556]McConnell, J. D.; Russell, D. W.: Tissue distribution and
ontogeny

[192557]of steroid 5-alpha-reductase isozyme expression. J. Clin.
Invest. 92:

[192558]903-910, 1993.

[192559]

[192560]

[192561]20504.

[192562]He, Z.; Yamamoto, R.; Furth, E. E.; Schantz, L. J.; Naylor, S.

[192563]L.; George, H.; Billheimer, J. T.; Strauss, J. F., III: cDNAs
encoding

[192564]members of a family of proteins related to human sterol

carrier protein

[192565]2 and assignment of the gene to human chromosome

1p21-pter. DNA Cell

[192566]Biol. 10: 559-569, 1991.

[192567]

[192568]

[192569]20505.

[192570]Vesa, J.; Hellsten, E.; Barnoski, B. L.; Emanuel, B. S.; Billheimer,

[192571]J. T.; Mead, S.; Cowell, J. K.; Strauss, J. F., III; Peltonen, L.

[192572]: Assignment of sterol carrier protein X/sterol carrier protein 2

[192573]to 1p32 and its exclusion as the causative gene for infantile neuronal

[192574]ceroid lipofuscinosis. Hum. Molec. Genet. 3: 341-346, 1994.

[192575]

[192576]

[192577]20506.

[192578]Welch, C. L.; Xia, Y.-R.; Billheimer, J. T.; Strauss, J. F., III;

[192579]Lusis, A. J.: Assignment of the mouse sterol carrier protein gene

[192580](Scp2) to chromosome 4. Mammalian Genome 7: 624–625, 1996.

[192581]

[192582]

[192583]20507.

[192584]Yamamoto, R.; Kallen, C. B.; Babalola, G. O.; Rennert, H.; Billheimer,

[192585]J. T.; Strauss, J. F., III: Cloning and expression of a cDNA encoding

[192586]human sterol carrier protein 2. Proc. Nat. Acad. Sci. 88: 463–467,

[192587]1991.

[192588]

[192589]

[192590]20508.

[192591]Yamamoto, R.; Naylor, S. L.; George, H.; Billheimer, J. T.; Strauss,

[192592]J. F., III: Assignment of the gene encoding sterol carrier protein

[192593]2 to human chromosome 1pter-p21. (Abstract) Cytogenet.

Cell Genet. 58:

[192594]1866-1867, 1991.

[192595]

[192596]

[192597]20509.

[192598]DeBose-Boyd, R. A.; Brown, M. S.; Li, W.-P.; Nohturfft, A.;
Goldstein,

[192599]J. L.; Espenshade, P. J.: Transport-dependent proteolysis
of SREBP:

[192600]relocation of Site-1 protease from Golgi to ER obviates the
need for

[192601]SREBP transport to Golgi. Cell 99: 703-712, 1999.

[192602]

[192603]

[192604]20510.

[192605]Hua, X.; Wu, J.; Goldstein, J. L.; Brown, M. S.; Hobbs, H. H.:

[192606]Structure of the human gene encoding sterol regulatory
element binding

[192607]protein-1 (SREBF1) and localization of SREBF1 and SREBF2

to chromosomes

[192608]17p11.2 and 22q13. Genomics 25: 667–673, 1995.

[192609]

[192610]

[192611]20511.

[192612]Osborne, T. F.: CREating a SCAP-less liver keeps SREBPs
pinned

[192613]in the ER membrane and prevents increased lipid synthesis in response

[192614]to low cholesterol and high insulin. Genes Dev. 15:
1873–1878, 2001.

[192615]

[192616]

[192617]20512.

[192618]Shimano, H.; Shimomura, I.; Hammer, R. E.; Herz, J.; Goldstein,

[192619]J. L.; Brown, M. S.; Horton, J. D.: Elevated levels of SREBP-2
and

[192620]cholesterol synthesis in livers of mice homozygous for a
targeted

[192621]disruption of the SREBP-1 gene. J. Clin. Invest. 100:
2115–2124,

[192622]1997.

[192623]

[192624]

[192625]20513.

[192626]Shimomura, I.; Hammer, R. E.; Richardson, J. A.; Ikemoto, S.; Bashmakov,

[192627]Y.; Goldstein, J. L.; Brown, M. S.: Insulin resistance and diabetes
in adipose

[192628]mellitus in transgenic mice expressing nuclear SREBP-1c
in adipose

[192629]tissue: model for congenital generalized lipodystrophy.
Genes Dev. 12:

[192630]3182–3194, 1998.

[192631]

[192632]

[192633]20514.

[192634]Tobe, K.; Suzuki, R.; Aoyama, M.; Yamauchi, T.; Kamon, J.;
Kubota,

[192635]N.; Terauchi, Y.; Matsui, J.; Akanuma, Y.; Kimura, S.;
Tanaka, J.;

[192636]Abe, M.; Ohsumi, J.; Nagai, R.; Kadowaki, T.: Increased expression
of the sterol regulatory element-binding protein-1 gene

[192637]

in insulin

[192638]receptor substrate-2 -/- mouse liver. J. Biol. Chem. 276:
38337-38340,

[192639]2001.

[192640]

[192641]

[192642]20515.

[192643]Wang, X.; Sato, R.; Brown, M. S.; Hua, X.; Goldstein, J. L.:
SREBP-1,

[192644]a membrane-bound transcription factor released by
sterol-regulated

[192645]proteolysis. Cell 77: 53-62, 1994.

[192646]

[192647]

[192648]20516.

[192649]Yokoyama, C.; Wang, X.; Briggs, M. R.; Admon, A.; Wu, J.;
Hua,

[192650]X.; Goldstein, J. L.; Brown, M. S.: SREBP-1, a basic-he-
lix-loop-helix-leucine

[192651]zipper protein that controls transcription of the low den-

sity lipoprotein

[192652]receptor gene. Cell 75: 187–197, 1993.

[192653]

[192654]

[192655]20517.

[192656]Joosten, P. H. L. J.; Toepoel, M.; Mariman, E. C. M.; Van
Zoelen,

[192657]E. J. J.: Promoter haplotype combinations of the platelet-
derived

[192658]growth factor alpha-receptor gene predispose to human
neural tube

[192659]defects. Nature Genet. 27: 215–217, 2001.

[192660]

[192661]

[192662]20518.

[192663]Colombo, R.: Age and origin of the PRNP E200K mutation
causing

[192664]familial Creutzfeldt–Jacob disease in Libyan Jews. (Letter)
Am. J.

[192665]Hum. Genet. 67: 528–531, 2000.

[192666]

[192667]

[192668]20519.

[192669]Cousens, J.; Smith, P. G.; Ward, H.; Everington, D.; Knight, R.

[192670]S. G.; Zeidler, M.; Stewart, G.; Smith-Bathgate, E. A. B.; Macleod,

[192671]M.-A.; Mackenzie, J.; Will, R. G.: Geographical distribution of variant

[192672]Creutzfeldt-Jakob disease in Great Britain, 1994-2000. Lancet 357:

[192673]1002-1007, 2001.

[192674]

[192675]

[192676]20520.

[192677]de Silva, R.; Ironside, J. W.; McCardle, L.; Esmonde, T.; Bell,

[192678]J.; Will, R.; Windl, O.; Dempster, M.; Estibeiro, P.; Lathe, R.:

[192679]Neuropathological phenotype and 'prion protein' genotype correlation

[192680]in sporadic Creutzfeldt-Jakob disease. Neurosci. Lett. 179:

50–52,

[192681]1994.

[192682]

[192683]

[192684]20521.

[192685]Deslys, J.–P.; Jaegly, A.; d'Aignaux, J. H.; Mouthon, F.; de
Villemeur,

[192686]T. B.; Dormont, D.: Genotype at codon 129 and suscepti-
bility to Creutzfeldt–Jakob

[192687]disease. Lancet 351: 1251 only, 1998.

[192688]

[192689]

[192690]20522.

[192691]Dlouhy, S. R.; Hsiao, K.; Farlow, M. F.; Foroud, T.; Con-
neally,

[192692]P. M.; Johnson, P.; Prusiner, S. B.; Hodes, M. E.; Ghetti, B.:
Linkage

[192693]of the Indiana kindred of Gerstmann–Straussler–Scheinker
disease to

[192694]the prion protein gene. Nature Genet. 1: 64–67, 1992.

[192695]

[192696]

[192697]20523.

[192698]Doh-ura, K.; Kitamoto, T.; Sakaki, Y.; Tateishi, J.: CJD discrepancy.

[192699](Letter) Nature 353: 801–802, 1991.

[192700]

[192701]

[192702]20524.

[192703]Doh-ura, K.; Tateishi, J.; Sasaki, H.; Kitamoto, T.; Sakaki, Y.

[192704]: Pro-to-leu change at position 102 of prion protein is the most common

[192705]but not the sole mutation related to Gerstmann–Straussler syndrome. Biochem.

[192706]Biophys. Res. Commun. 163: 974–979, 1989.

[192707]

[192708]

[192709]20525.

[192710]Duchen, L. W.; Poulter, M.; Harding, A. E.: Dementia associated

[192711]with a 216 base pair insertion in the prion protein gene: clinical

[192712]and neuropathological features. Brain 116: 555–567, 1993.

[192713]

[192714]

[192715]20526.

[192716]Erginel-Unaltuna, N.; Peoc'h, K.; Komurcu, E.; Acuner, T. T.;

[192717]Issever, H.; Laplanche, J.–L.: Distribution of the M129V polymorphism

[192718]of the prion protein gene in a Turkish population suggests a high

[192719]risk for Creutzfeldt–Jacob disease. Europ. J. Hum. Genet. 9: 965–968,

[192720]2001.

[192721]

[192722]

[192723]20527.

[192724]Farlow, M. R.; Yee, R. D.; Dlouhy, S. R.; Conneally, P. M.;
Azzarelli,

[192725]B.; Ghetti, B.: Gerstmann–Straussler–Scheinker disease. I.
Extending

[192726]the clinical spectrum. *Neurology* 39: 1446–1452, 1989.

[192727]

[192728]

[192729]20528.

[192730]Flechsigg, E.; Shmerling, D.; Hegyi, I.; Raeber, A. J.; Fischer,

[192731]M.; Cozzio, A.; von Mering, C.; Aguzzi, A.; Weissmann, C.:
Prion

[192732]protein devoid of the octapeptide repeat region restores
susceptibility

[192733]to scrapie in PrP knockout mice. *Neuron* 27: 399–408,
2000.

[192734]

[192735]

[192736]20529.

[192737]Forloni, G.; Angeretti, N.; Chiesa, R.; Monzani, E.;
Salmona,

[192738]M.; Bugiani, O.: Neurotoxicity of a prion protein fragment.

(Letter) Nature 362:

[192739]543–546, 1993.

[192740]

[192741]

[192742]20530.

[192743]Gabizon, R.; Rosenmann, H.; Meiner, Z.; Kahana, I.; Kahana, E.;

[192744]Shugart, Y.; Ott, J.; Prusiner, S. B.: Mutation and polymorphism

[192745]of the prion protein gene in Libyan Jews with Creutzfeldt-Jakob disease

[192746](CJD). Am. J. Hum. Genet. 53: 828–835, 1993.

[192747]

[192748]

[192749]20531.

[192750]Gajdusek, D. C.: The transmissible amyloidoses: genetical control

[192751]of spontaneous generation of infectious amyloid proteins by nucleation

[192752]of configurational change in host precursors: kuru–
CJD–GSS–scrapie–BSE. Europ.

[192753]J. Epidemiol. 7: 567–577, 1991.

[192754]

[192755]

[192756]20532.

[192757]Gambetti, P.; Petersen, R.; Monari, L.; Tabaton, M.; Autilio–
Gambetti,

[192758]L.: Fatal familial insomnia and the widening spectrum of
prion diseases. Brit.

[192759]Med. Bull. 49: 980–994, 1993.

[192760]

[192761]

[192762]20533.

[192763]Ghetti, B.; Tagliavini, F.; Masters, C. L.; Beyreuther, K.; Gi–
accone,

[192764]G.; Verga, L.; Farlow, M. R.; Conneally, P. M.; Dlouhy, S. R.;
Azzarelli,

[192765]B.; Bugiani, O.: Gerstmann–Straussler–Scheinker disease.

II. Neurofibrillary

[192766]tangles and plaques with PrP-amyloid coexist in an affected family. Neurology 39:

[192767]1453-1461, 1989.

[192768]

[192769]

[192770]20534.

[192771]Giaccone, G.; Verga, L.; Bugiani, O.; Frangione, B.; Serban, D.;

[192772]Prusiner, S. B.; Farlow, M. R.; Ghetti, B.; Tagliavini, F.: Prion

[192773]protein preamyloid and amyloid deposits in Gerstmann-Straussler-Scheinker

[192774]disease, Indiana kindred. Proc. Nat. Acad. Sci. 89: 9349-9353, 1992.

[192775]

[192776]

[192777]20535.

[192778]Goldfarb, L. G.; Brown, P.; Cervenakova, L.; Gajdusek, D. C.:

[192779]Molecular genetic studies of Creutzfeldt-Jakob disease. Molec. Neurobiol. 8:

[192780]89–97, 1994.

[192781]

[192782]

[192783]20536.

[192784]Goldfarb, L. G.; Brown, P.; Haltia, M.; Cathala, F.; McCombie,
bie,

[192785]W. R.; Kovanen, J.; Cervenakova, L.; Goldin, L.; Nieto, A.;
Godec,

[192786]M. S.; Asher, D. M.; Gajdusek, D. C.: Creutzfeldt–Jakob
disease cosegregates

[192787]with the codon 178–asn PRNP mutation in families of European origin. Ann.

[192788]Neurol. 31: 274–281, 1992.

[192789]

[192790]

[192791]20537.

[192792]Goldfarb, L. G.; Brown, P.; McCombie, W. R.; Goldgaber,
D.; Swergold,

[192793]G. D.; Wills, P. R.; Cervenakova, L.; Baron, H.; Gibbs, C. J.,
Jr.;

[192794]Gajdusek, D. C.: Transmissible familial Creutzfeldt–Jakob disease

[192795]associated with five, seven, and eight extra octapeptide coding repeats

[192796]in the PRNP gene. Proc. Nat. Acad. Sci. 88: 10926–10930, 1991.

[192797]

[192798]

[192799]20538.

[192800]Goldfarb, L. G.; Brown, P.; Mitrova, E.; Cervenakova, L.; Goldin,

[192801]L.; Korczyn, A. D.; Chapman, J.; Galvez, S.; Cartier, L.; Rubenstein,

[192802]R.; Gajdusek, D. C.: Creutzfeldt–Jacob disease associated with the

[192803]PRNP codon 200–lys mutation: an analysis of 45 families. Europ. J.

[192804]Epidemiol. 7: 477–486, 1991.

[192805]

[192806]

[192807]20539.

[192808]Goldfarb, L. G.; Haltia, M.; Brown, P.; Nieto, A.; Kovanen, J.;

[192809]McCombie, W. R.; Trapp, S.; Gajdusek, D. C.: New mutation in scrapie

[192810]amyloid precursor gene (at codon 178) in Finnish Creutzfeldt–Jakob

[192811]kindred. (Letter) Lancet 337: 425, 1991.

[192812]

[192813]

[192814]20540.

[192815]Goldfarb, L. G.; Mitrova, E.; Brown, P.; Toh, B. H.; Gajdusek,

[192816]D. C.: Mutation in codon 200 of scrapie amyloid protein gene in two

[192817]clusters of Creutzfeldt–Jakob disease in Slovakia. (Letter) Lancet 336:

[192818]514–515, 1990.

[192819]

[192820]

[192821]20541.

[192822]Travis, G. H.; Sutcliffe, J. G.; Bok, D.: The retinal degeneration

[192823]slow (rds) gene product is a photoreceptor disc membrane-associated

[192824]glycoprotein. Neuron 6: 61–70, 1991.

[192825]

[192826]

[192827]20542.

[192828]Morris, C.; Heisterkamp, N.; Hao, Q. L.; Testa, J. R.; Groffen,

[192829]J.: The human tyrosine kinase gene (FER) maps to chromosome 5 and

[192830]is deleted in myeloid leukemias with a del(5q). Cytogenet. Cell Genet. 53:

[192831]196–200, 1990.

[192832]

[192833]

[192834]20543.

[192835]Herrera, V. L. M.; Xie, H. X.; Lopez, L. V.; Schork, N. J.; Ruiz-Opazo,

[192836]N.: The alpha-1 Na,K-ATPase gene is a susceptibility hypertension

[192837]gene in the Dahl salt-sensitive-HSD rat. J. Clin. Invest. 102: 1102-1111,

[192838]1998.

[192839]

[192840]

[192841]20544.

[192842]Williams, T. M.; Montoya, G.; Wu, Y.; Eddy, R. L.; Byers, M. G.;

[192843]Shows, T. B.: The TCF8 gene encoding a zinc finger protein (Nil-2-a)

[192844]resides on human chromosome 10p11.2. Genomics 14: 194-196, 1992.

[192845]

[192846]

[192847]20545.

[192848]van Nie, R.; Ivanyi, D.; Demant, P.: A new H-2-linked mutation,

[192849]rds, causing retinal degeneration in the mouse. Tissue Antigens 12:

[192850]106–108, 1978.

[192851]

[192852]

[192853]20546.

[192854]Vine, A. K.; Schatz, H.: Adult-onset foveomacular pigment
epithelial

[192855]dystrophy. Am. J. Ophthal. 89: 680–691, 1980.

[192856]

[192857]

[192858]20547.

[192859]Weleber, R. G.; Carr, R. E.; Murphey, W. H.; Sheffield, V. C.;

[192860]Stone, E. M.: Phenotypic variation including retinitis pig-
mentosa,

[192861]pattern dystrophy, and fundus flavimaculatus in a single
family with

[192862]a deletion of codon 153 or 154 of the peripherin/RDS
gene. Arch.

[192863]Ophthal. 111: 1531–1542, 1993.

[192864]

[192865]

[192866]20548.

[192867]Wells, J.; Wroblewski, J.; Keen, J.; Inglehearn, C.; Jubb, C.;

[192868]Eckstein, A.; Jay, M.; Arden, G.; Bhattacharya, S.; Fitzke, F.;
Bird,

[192869]A.: Mutations in the human retinal degeneration slow
(RDS) gene can

[192870]cause either retinitis pigmentosa or macular dystrophy.

Nature Genet. 3:

[192871]213–218, 1993.

[192872]

[192873]

[192874]20549.

[192875]Wroblewski, J. J.; Wells, J. A., III; Eckstein, A.; Fitzke, F.;

[192876]Jubb, C.; Keen, J.; Inglehearn, C.; Bhattacharya, S.; Arden,
G. B.;

[192877]Jay, M.; Bird, A. C.: Macular dystrophy associated with mu-
tations

[192878]at codon 172 in the human retinal degeneration slow
gene. Ophthalmology 101:

[192879]12–22, 1994.

[192880]

[192881]

[192882]20550.

[192883]Eph Nomenclature Committee: Unified nomenclature for
the Eph

[192884]family receptors and their ligands, the ephrins. Cell 90:
403–404,

[192885]1997.

[192886]

[192887]

[192888]20551.

[192889]Hattori, M.; Osterfield, M.; Flanagan, J. G.: Regulated
cleavage

[192890]of a contact–mediated axon repellent. Science 289:
1360–1365, 2000.

[192891]

[192892]

[192893]20552.

[192894]Maru, Y.; Hirai, H.; Yoshida, M. C.; Takaku, F.: Evolution,
expression,
[192895]and chromosomal location of a novel receptor tyrosine ki-
nase gene,
[192896]eph. Molec. Cell. Biol. 8: 3770–3776, 1988.

[192897]

[192898]

[192899]20553.

[192900]Yoshida, M. C.; Maru, H.; Hirai, H.; Takau, F.: Chromoso-
mal location
[192901]of a novel receptor tyrosine kinase gene, EPH, on chromo-
some 7. (Abstract) Cytogenet.
[192902]Cell Genet. 51: 1113 only, 1989.

[192903]

[192904]

[192905]20554.

[192906]Boyd, A. W.; Ward, L. D.; Wicks, I. P.; Simpson, R. J.; Sal-
varis,
[192907]E.; Wilks, A.; Welch, K.; Loudovaris, M.; Rockman, S.; Bus-
manis, I.

[192908]: Isolation and characterization of a novel receptor-type protein

[192909]tyrosine kinase (hek) from a human pre-B cell line. J. Biol. Chem. 267:

[192910]3262–3267, 1992.

[192911]

[192912]

[192913]20555.

[192914]Fox, G. M.; Holst, P. L.; Chute, H. T.; Lindberg, R. A.; Janssen,

[192915]A. M.; Basu, R.; Welcher, A. A.: cDNA cloning and tissue distribution

[192916]of five human EPH-like receptor protein-tyrosine kinases. Oncogene 10:

[192917]897–905, 1995.

[192918]

[192919]

[192920]20556.

[192921]Wicks, I. P.; Lapsys, N. M.; Baker, E.; Campbell, L. J.; Boyd,

[192922]A. W.; Sutherland, G. R.: Localization of a human receptor

tyrosine

[192923]kinase (ETK1) to chromosome region 3p11.2. Genomics
19: 38–41, 1994.

[192924]

[192925]

[192926]20557.

[192927]Wicks, I. P.; Wilkinson, D.; Salvaris, E.; Boyd, A. W.: Molec-
ular

[192928]cloning of HEK, the gene encoding a receptor tyrosine ki-
nase expressed

[192929]by human lymphoid tumor cell lines. Proc. Nat. Acad. Sci.
89: 1611–1615,

[192930]1992.

[192931]

[192932]

[192933]20558.

[192934]Patterson, D.; Sujansky, E.; Hart, I.; Bleskan, J.; Walton, K.;

[192935]Giang, J.; Shechter, I.: Recombinant 8 syndrome break-
point analysis.

[192936](Abstract) Am. J. Hum. Genet. 57: A91 only, 1995.

[192937]

[192938]

[192939]20559.

[192940]Elsea, S. H.; Mykytyn, K.; Ferrell, K.; Coulter, K. L.; Das, P.;

[192941]Dubiel, W.; Patel, P. I.; Metherall, J. E.: Hemizygosity for
the

[192942]COP9 signalosome subunit gene, SGN3, in the Smith–
Magenis syndrome. Am.

[192943]J. Med. Genet. 87: 342–348, 1999.

[192944]

[192945]

[192946]20560.

[192947]Koyama, K.; Fukushima, Y.; Inazawa, J.; Tomotsune, D.;
Takahashi,

[192948]N.; Nakamura, Y.: The human homologue of the murine
Llglh gene (LLGL)

[192949]maps within the Smith–Magenis syndrome region in
17p11.2. Cytogenet.

[192950]Cell Genet. 72: 78–82, 1996.

[192951]

[192952]

[192953]20561.

[192954]Forsberg, E.; Hirsch, E.; Frohlich, L.; Meyer, M.; Ekblom, P.;

[192955]Aszodi, A.; Werner, S.; Fassler, R.: Skin wounds and severed nerves

[192956]heal normally in mice lacking tenascin-C. Proc. Nat. Acad. Sci. 93:

[192957]6594-6599, 1996.

[192958]

[192959]

[192960]20562.

[192961]Gulcher, J. R.; Alexakos, M. J.; Le Beau, M. M.; Lemons, R. S.;

[192962]Stefansson, K.: Chromosomal localization of the human hexabrachion

[192963](tenascin) gene and evidence for recent reduplication within the gene. Genomics 6:

[192964]616-622, 1990.

[192965]

[192966]

[192967]20563.

[192968]Gulcher, J. R.; Nies, D. E.; Alexakos, M. J.; Ravikant, N. A.;

[192969]Sturgill, M. E.; Marton, L. S.; Stefansson, K.: Structure of
the

[192970]human hexabrachion (tenascin) gene. Proc. Nat. Acad. Sci.
88: 9438–9442,

[192971]1991.

[192972]

[192973]

[192974]20564.

[192975]Mitrovic, N.; Schachner, M.: Detection of tenascin–C in the
nervous

[192976]system of the tenascin–C mutant mouse. J. Neurosci. Res.
42: 710–717,

[192977]1995.

[192978]

[192979]

[192980]20565.

[192981]Nies, D. E.; Hemesath, T. J.; Kim, J.–H.; Gulcher, J. R.; Ste–

fansson,

[192982]K.: The complete cDNA sequence of human hexabrachion
(tenascin):

[192983]a multidomain protein containing unique epidermal
growth factor repeats. J.

[192984]Biol. Chem. 266: 2818–2823, 1991.

[192985]

[192986]

[192987]20566.

[192988]Williams, T. M.; Moolten, D.; Burlein, J.; Romano, J.; Bhaer-
man,

[192989]R.; Godillot, A.; Mellon, M.; Rauscher, F. J., III; Kant, J. A.:

[192990]Identification of a zinc finger protein that inhibits IL–2
gene expression. Science 254:

[192991]1791–1794, 1991.

[192992]

[192993]

[192994]20567.

[192995]Ludecke, H.–J.; Schaper, J.; Meinecke, P.; Momeni, P.;
Gross, S.;

- [192996] von Holtum, D.; Hirche, H.; Abramowicz, M. J.; Albrecht, B.; Apacik, C.; Christen, H.-J.; Claussen, U.; and 28 others: Genotypic and phenotypic spectrum in tricho-rhino-phalangeal syndrome types I and III. *Am. J. Hum. Genet.* 68: 81-91, 2001.
- [193000]
- [193001]
- [193002] 20568.
- [193003] Chiang, L.; Contreras, L.; Chiang, J.; Ward, P. H.: Human prostatic gastricsinogen: the precursor of seminal fluid acid proteinase. *Arch. Biochem. Biophys.* 210: 14-20, 1981.
- [193004]
- [193005]
- [193006]
- [193007]
- [193008] 20569.
- [193009] Hayano, T.; Sogawa, K.; Ichihara, Y.; Fujii-Kuriyama, Y.; Takahashi,

- [193010]K.: Primary structure of human pepsinogen C gene. J. Biol. Chem. 263: 1382–1385, 1988.
- [193012]
- [193013]
- [193014]20570.
- [193015]Pals, G.; Azuma, T.; Mohandas, T. K.; Bell, G. I.; Bacon, J.; Samloff, I. M.; Walz, D. A.; Barr, P. J.; Taggart, R. T.: Human pepsinogen C (progastricsin) polymorphism: evidence for a single locus located at 6p21.1–pter. Genomics 4: 137–145, 1989.
- [193019]
- [193020]
- [193021]20571.
- [193022]Randolph, L. M.; Azuma, T.; Petersen, G. M.; Sparkes, R. S.; Toyoda, M.; Wang, S. J.; Taggart, R. T.: Assignment of pepsinogen C (PGC)

[193024]to 6p21.3–p21.1. (Abstract) Cytogenet. Cell Genet. 51:
1063–1064,
[193025]1989.

[193026]

[193027]

[193028]20572.

[193029]Szymura, J. M.; Klein, J.: Linkage of a gene controlling uri-
nary
[193030]pepsinogen with the major histocompatibility complex of
the mouse.

[193031]Immunogenetics 13: 267–271, 1981.

[193032]

[193033]

[193034]20573.

[193035]Taggart, R. T.; Cass, L. G.; Mohandas, T. K.; Derby, P.;
Barr,

[193036]P. J.; Pals, G.; Bell, G. I.: Human pepsinogen C
(progastricsin):

[193037]isolation of cDNA clones, localization to chromosome 6,
and sequence

[193038]homology with pepsinogen A. J. Biol. Chem. 264:
375–379, 1989.

[193039]

[193040]

[193041]20574.

[193042]Taggart, R. T.; Mohandas, T. K.; Bell, G. I.: Assignment of
human

[193043]preprogastricsin (PGC) to chromosome 6 and regional lo-
calization of

[193044]PGC (6pter–p21.1), prolactin PRL (6pter–p21.1). (Abstract)
Cytogenet.

[193045]Cell Genet. 46: 701–702, 1987.

[193046]

[193047]

[193048]20575.

[193049]Takahara, K.; Fukushige, S.; Murotsu, T.; Ichihara, Y.;
Hayano,

[193050]T.; Ishihara, T.; Takahashi, K.: Assignment of human
pepsinogen C

[193051](PGC) gene to chromosome 6. Cytogenet. Cell Genet. 52:
100–101,

[193052]1989.

[193053]

[193054]

[193055]20576.

[193056]Weil, D.; Cottreau, D.; Cong, N. V.; Rebourcet, R.; Foubert, C.;

[193057]Gross, M.-S.; Dreyfus, J.-C.; Kahn, A.: Assignment of the gene for

[193058]F-type phosphofructokinase to human chromosome 10 by somatic cell

[193059]hybridization and specific immunoprecipitation. Ann. Hum. Genet. 44:

[193060]11-16, 1980.

[193061]

[193062]

[193063]20577.

[193064]Miwa, S.; Sato, T.; Murao, H.; Kozuru, M.; Ibayashi, H.: A new

[193065]type of phosphofructokinase deficiency: hereditary non-spherocytic

[193066]hemolytic anemia. Acta Haemat. Jpn. 35: 113–118, 1972.

[193067]

[193068]

[193069]20578.

[193070]Bernardi, G.: Personal Communication. Paris, France
5/13/1992.

[193071]

[193072]

[193073]20579.

[193074]Boulard, M. R.; Bois, M.; Reviron, M.; Najean, Y.: Red-cell
phosphofructokinase

[193075]deficiency. New Eng. J. Med. 291: 978–979, 1974.

[193076]

[193077]

[193078]20580.

[193079]Chadefaux, B.; Rethore, M. O.; Allard, D.: Regional map-
ping of

[193080]liver type 6-phosphofructokinase isoenzyme on chromo-
some 21. Hum.

[193081]Genet. 68: 136–137, 1984.

[193082]

[193083]

[193084]20581.

[193085]Cox, D. R.; Kawashima, H.; Vora, S.; Epstein, C. J.: Regional

[193086]mapping of SOD–1, PRGS, and PFK–L on human chromosome 21. (Abstract) Cytogenet.

[193087]Cell Genet. 37: 441–442, 1984.

[193088]

[193089]

[193090]20582.

[193091]Elson, A.; Levanon, D.; Brandeis, M.; Dafni, N.; Bernstein, Y.;

[193092]Danciger, E.; Groner, Y.: The structure of the human liver–type phosphofructokinase

[193093]gene. Genomics 7: 47–56, 1990.

[193094]

[193095]

[193096]20583.

[193097]Etiemble, J.; Kahn, A.; Boivin, P.; Bernard, J. F.; Goude-
mand,

[193098]M.: Hereditary hemolytic anemia with erythrocyte phos-
phofructokinase

[193099]deficiency. Hum. Genet. 31: 83–91, 1976.

[193100]

[193101]

[193102]20584.

[193103]Etiemble, J.; Picat, C.; Simeon, J.; Blatrix, C.; Boivin, P.:

[193104]Inherited erythrocyte phosphofructokinase deficiency:
molecular mechanism. Hum.

[193105]Genet. 55: 383–390, 1980.

[193106]

[193107]

[193108]20585.

[193109]Gehnrich, S. C.; Gekakis, N.; Sul, H. S.: Liver (B-type)
phosphofructokinase

[193110]mRNA: cloning, structure, and expression. J. Biol. Chem.
263: 11755–11759,

[193111]1988.

[193112]

[193113]

[193114]20586.

[193115]Kahn, A.; Etiemble, J.; Meienhofer, M. C.; Boivin, P.: Erythrocyte

[193116]phosphofructokinase deficiency associated with an unstable variant

[193117]of muscle phosphofructokinase. Clin. Chim. Acta 61: 415–419, 1975.

[193118]

[193119]

[193120]20587.

[193121]Levanon, D.; Danciger, E.; Dafni, N.; Groner, Y.: Genomic clones

[193122]of the human liver-type phosphofructokinase. Biochem. Biophys. Res.

[193123]Commun. 141: 374–380, 1986.

[193124]

[193125]

[193126]20588.

[193127]Pantelakis, S. N.; Karaklis, A. G.; Alexiou, D.; Vardas, E.;
Valaes,

[193128]T.: Red cell enzymes in trisomy 21. Am. J. Hum. Genet. 22:
184–193,

[193129]1970.

[193130]

[193131]

[193132]20589.

[193133]Van Keuren, M.; Drabkin, H.; Hart, I.; Harker, D.; Patter-
son,

[193134]D.; Vora, S.: Regional assignment of human liver-type
6–phosphofructokinase

[193135]to chromosome 21q22.3 by using somatic cell hybrids
and a monoclonal

[193136]anti-L antibody. Hum. Genet. 74: 34–40, 1986.

[193137]

[193138]

[193139]20590.

[193140]Vora, S.; Davidson, M.; Seaman, C.; Miranda, A. F.; Noble,
N.

[193141]A.; Tanaka, K. R.; Frenkel, E. P.; DiMauro, S.: Heterogeneity
of

[193142]the molecular lesions in inherited phosphofructokinase
deficiency. J.

[193143]Clin. Invest. 72: 1995–2006, 1983.

[193144]

[193145]

[193146]20591.

[193147]Vora, S.; Durham, S.; de Martinville, B.; Francke, U.: As-
signment

[193148]of the genes for liver type phosphofructokinase (PFKL) to
chromosome

[193149]21 and for muscle type (PFKM) to region p32–q32 of
chromosome 1. (Abstract) Cytogenet.

[193150]Cell Genet. 32: 324 only, 1982.

[193151]

[193152]

[193153]20592.

[193154]Vora, S.; Francke, U.: Assignment of the human gene for
liver

[193155]type phosphofructokinase isozyme to chromosome 21 using somatic cell

[193156]hybrids. (Abstract) Pediat. Res. 15: 570 only, 1981.

[193157]

[193158]

[193159]20593.

[193160]Vora, S.; Francke, U.: Assignment of the human gene for liver-type

[193161]6-phosphofructokinase isozyme (PFKL) to chromosome 21 by using somatic

[193162]cell hybrids and monoclonal anti-L antibody. Proc. Nat. Acad. Sci. 78:

[193163]3738-3742, 1981.

[193164]

[193165]

[193166]20594.

[193167]Wang, D.; Fang, H.; Cantor, C. R.; Smith, C. L.: A contiguous

[193168]NotI restriction map of band q22.3 of human chromosome 21. Proc.

[193169]Nat. Acad. Sci. 89: 3222–3226, 1992.

[193170]

[193171]

[193172]20595.

[193173]Vidal, R.; Revesz, T.; Rostagno, A.; Kim, E.; Holton, J. L.;
Bek,

[193174]T.; Bojsen-Moller, M.; Braendgaard, H.; Plant, G.; Ghiso, J.;
Frangione,

[193175]B.: A decamer duplication in the 3-prime region of the BRI
gene originates

[193176]an amyloid peptide that is associated with dementia in a
Danish kindred. Proc.

[193177]Nat. Acad. Sci. 97: 4920–4925, 2000.

[193178]

[193179]

[193180]20596.

[193181]Fathallah-Shaykh, H.; Wolf, S.; Wong, E.; Posner, J. B.:
Cloning

[193182]of a leucine-zipper protein recognized by the sera of pa-
tients with

[193183]antibody-associated paraneoplastic cerebellar degeneration. Proc.

[193184]Nat. Acad. Sci. 88: 3451–3454, 1991.

[193185]

[193186]

[193187]20597.

[193188]Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.; Posner, J. B.;

[193189]Siniscalco, M.: In situ mapping of the gene coding for a leucine

[193190]zipper DNA binding protein (CDR 2) to the region between two rare

[193191]fragile sites of autosome 16 (16p12–p13.1). (Abstract) Cytogenet.

[193192]Cell Genet. 58: 1999–2000, 1991.

[193193]

[193194]

[193195]20598.

[193196]Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.; Posner, J. B.;

[193197]Siniscalco, M.: In situ mapping of the gene coding for a leucine

[193198]zipper DNA binding protein (CDR62) to 16p12–16p13.1.

Genomics 13:

[193199]1340–1342, 1992.

[193200]

[193201]

[193202]20599.

[193203]Raimondi, E.; Rubboli, F.; Moralli, D.; Chini, B.; Fornasari, D.;

[193204]Tarroni, P.; De Carli, L.; Clementi, F.: Chromosomal localization

[193205]and physical linkage of the genes encoding the human alpha-3, alpha-5,

[193206]and beta-4 neuronal nicotinic receptor subunits. Genomics 12: 849–850,

[193207]1992.

[193208]

[193209]

[193210]20600.

[193211]Anand, R.; Lindstrom, J.: Nucleotide sequence of the human nicotinic

[193212]acetylcholine receptor beta-2 subunit gene. Nucleic Acids Res. 18:

[193213]4272, 1990.

[193214]

[193215]

[193216]20601.

[193217]De Fusco, M.; Becchetti, A.; Patrignani, A.; Annesi, G.; Gambardella,

[193218]A.; Quattrone, A.; Ballabio, A.; Wanke, E.; Casari, G.: The nicotinic

[193219]receptor beta-2 subunit is mutant in nocturnal frontal lobe epilepsy. Nature

[193220]Genet. 26: 275-276, 2000.

[193221]

[193222]

[193223]20602.

[193224]Lueders, K. K.; Elliott, R. W.; Marenholz, I.; Mischke, D.; DuPree,

[193225]M.; Hamer, D.: Genomic organization and mapping of the human and

[193226]mouse neuronal beta-1-nicotinic acetylcholine receptor genes. Mammalian

[193227]Genome 10: 900-905, 1999.

[193228]

[193229]

[193230]20603.

[193231]Phillips, H. A.; Favre, I.; Kirkpatrick, M.; Zuberi, S. M.; Goudie,

[193232]D.; Heron, S. E.; Scheffer, I. E.; Sutherland, G. R.; Berkovic, S.

[193233]F.; Bertrand, D.; Mulley, J. C.: CHRNA2 is the second acetylcholine

[193234]receptor subunit associated with autosomal dominant nocturnal frontal

[193235]lobe epilepsy. Am. J. Hum. Genet. 68: 225-231, 2001.

[193236]

[193237]

[193238]20604.

[193239]Picciotto, M. R.; Zoli, M.; Lena, C.; Bessis, A.; Lallemant, Y.;

[193240]LeNovere, N.; Vincent, P.; Pich, E. M.; Brulet, P.; Changeux, J.-P.

[193241]: Abnormal avoidance learning in mice lacking functional high-affinity

[193242]nicotine receptor in the brain. Nature 374: 65–67, 1995.

[193243]

[193244]

[193245]20605.

[193246]Zoli, M.; Picciotto, M. R.; Ferrari, R.; Cocchi, D.; Changeux,

[193247]J.-P.: Increased neurodegeneration during ageing in mice lacking

[193248]high-affinity nicotine receptors. EMBO J. 18: 1235–1244, 1999.

[193249]

[193250]

[193251]20606.

[193252]Koyama, K.; Sudo, K.; Nakamura, Y.: Mapping of the human nicotinic

[193253]acetylcholine receptor beta-3 gene (CHRNA3) within chromosome 12p12.3. Genomics 21:

[193254]460-461, 1994.

[193255]

[193256]

[193257]20607.

[193258]Willoughby, J. J.; Ninkina, N. N.; Beech, M. M.; Latchman, D. S.;

[193259]Wood, J. N.: Molecular cloning of a human neuronal nicotinic acetylcholine

[193260]receptor beta-3-like subunit. Neurosci. Lett. 155: 136-139, 1993.

[193261]

[193262]

[193263]20608.

[193264]Tarroni, P.; Rubboli, F.; Chini, B.; Zwart, R.; Oortgiesen, M.;

[193265]Sher, E.; Clementi, F.: Neuronal-type nicotinic receptors in human

[193266]neuroblastoma and small-cell lung carcinoma cell lines. FEBS Lett. 312:

[193267]66-70, 1992.

[193268]

[193269]

[193270]20609.

[193271]Bonner, T. I.; Buckley, N. J.; Young, A. C.; Brann, M. R.:

Identification

[193272]of a family of muscarinic acetylcholine receptor genes.

Science 237:

[193273]527-532, 1987.

[193274]

[193275]

[193276]20610.

[193277]Schwenk, W. F.; Rizza, R. A.; Mandarino, L. J.; Gerich, J. E.;

[193278]Hayles, A. B.; Haymond, M. W.: Familial insulin resistance

and acanthosis

[193279]nigricans: presence of a postbinding defect. Diabetes 35:

33-37,

[193280]1986.

[193281]

[193282]

[193283]20611.

[193284]Speckman, R. A.; Garg, A.; Du, F.; Bennett, L.; Veile, R.;
Arioglu,

[193285]E.; Taylor, S. I.; Lovett, M.; Bowcock, A. M.: Mutational and
haplotype

[193286]analyses of families with familial partial lipodystrophy
(Dunnigan

[193287]variety) reveal recurrent missense mutations in the globu-
lar C-terminal

[193288]domain of lamin A/C. Am. J. Hum. Genet. 66: 1192–1198,
2000. Note:

[193289]Erratum: Am. J. Hum. Genet. 67: 775 only, 2000.

[193290]

[193291]

[193292]20612.

[193293]Chan, J.; Watt, V. M.: Eek and erk, new members of the
eph subclass

[193294]of receptor protein-tyrosine kinases. Oncogene 6:
1057–1061, 1991.

[193295]

[193296]

[193297]20613.

[193298]Park, S.; Frisen, J.; Barbacid, M.: Aberrant axonal projections

[193299]in mice lacking EphA8 (Eek) tyrosine protein kinase receptors. EMBO

[193300]J. 16: 3106–3114, 1997.

[193301]

[193302]

[193303]20614.

[193304]Ganju, P.; Shigemoto, K.; Brennan, J.; Entwistle, A.; Reith, A.

[193305]D.: The Eck receptor tyrosine kinase is implicated in pattern formation

[193306]during gastrulation, hindbrain segmentation and limb development. Oncogene 9:

[193307]1613–1624, 1994.

[193308]

[193309]

[193310]20615.

[193311]Lindberg, R. A.; Hunter, T.: cDNA cloning and characterization

[193312]of eck, an epithelial cell receptor-tyrosine kinase in the eph/elk

[193313]family of protein kinases. Molec. Cell. Biol. 10: 6316-6324, 1990.

[193314]

[193315]

[193316]20616.

[193317]Sulman, E. P.; Tang, X. X.; Allen, C.; Biegel, J. A.; Pleasure,

[193318]D. E.; Brodeur, G. M.; Ikegaki, N.: ECK, a human EPH-related gene,

[193319]maps to 1p36.1, a common region of alteration in human cancers. Genomics 40:

[193320]371-374, 1997.

[193321]

[193322]

[193323]20617.

[193324]Colucci, F.; Schweighoffer, E.; Tomasello, E.; Turner, M.; Ortaldo,

[193325]J. R.; Vivier, E.; Tybulewicz, V. L. J.; Di Santo, J. P.: Natural

[193326]cytotoxicity uncoupled from the Syk and ZAP-70 intracellular kinases. Nature

[193327]Immun. 3: 288-294, 2002.

[193328]

[193329]

[193330]20618.

[193331]Ku, G.; Malissen, B.; Mattei, M.-G.: Chromosomal location of the

[193332]Syk and ZAP-70 tyrosine kinase genes in mice and humans. Immunogenetics 40:

[193333]300-302, 1994.

[193334]

[193335]

[193336]20619.

[193337]Toyabe, S.-I.; Watanabe, A.; Harada, W.; Karasawa, T.; Uchiyama,

[193338]M.: Specific immunoglobulin E responses in ZAP-70-deficient patients

[193339]are mediated by Syk-dependent T-cell receptor signalling.

Immunology 103:

[193340]164–171, 2001.

[193341]

[193342]

[193343]20620.

[193344]Sudbrak, R.; Golla, A.; Hogan, K.; Powers, P.; Gregg, R.; Du
Chesne,

[193345]I.; Lehmann–Horn, F.; Deufel, T.: Exclusion of malignant
hyperthermia

[193346]susceptibility (MHS) from a putative MHS2 locus on chro-
mosome 17q

[193347]and of the alpha–1, beta–1, and gamma subunits of the
dihydropyridine

[193348]receptor calcium channel as candidates for the molecular
defect. Hum.

[193349]Molec. Genet. 2: 857–862, 1993.

[193350]

[193351]

[193352]20621.

[193353]Naim, H. Y.; Sterchi, E. E.; Lentze, M. J.: Structure, biosyn–

thesis,

[193354]and glycosylation of human small intestinal maltase–
glucoamylase. J.

[193355]Biol. Chem. 263: 19709–19717, 1988.

[193356]

[193357]

[193358]20622.

[193359]Nichols, B. L.; Eldering, J.; Avery, S.; Hahn, D.; Quaroni, A.;

[193360]Sterchi, E.: Human small intestinal maltase–glucoamylase
cDNA cloning:

[193361]homology to sucrase–isomaltase. J. Biol. Chem. 273:
3076–3081, 1998.

[193362]

[193363]

[193364]20623.

[193365]Schuler, G. D.; Boguski, M. S.; Stewart, E. A.; Stein, L. D.;

Gyapay,

[193366]G.; Rice, K.; White, R. E.; Rodriguez–Tome, P.; Aggarwal,
A.; Bajorek,

[193367]E.; Bentolila, S.; Birren, B. B.; Butler, A.; Castle, A. B.; Chi–

annilkulchai,

[193368]N.; Chu, A.; Clee, C.; Cowles, S.; Day, P. J. R.; Dibling, T.;
and

[193369]84 others: A gene map of the human genome. Science
274: 540–546,

[193370]1996.

[193371]

[193372]

[193373]20624.

[193374]Dubiel, W.; Ferrell, K.; Rechsteiner, M.: Peptide sequencing
identifies

[193375]MSS1, a modulator of HIV Tat–mediated transactivation,
as subunit

[193376]7 of the 26 S protease. FEBS Lett. 323: 276–278, 1993.

[193377]

[193378]

[193379]20625.

[193380]Nelbock, P.; Dillon, P. J.; Perkins, A.; Rosen, C. A.: A cDNA

[193381]for a protein that interacts with the human immunodeficiency virus

[193382]Tat activator. Science 248: 1650–1653, 1990.

[193383]

[193384]

[193385]20626.

[193386]Shibuya, H.; Irie, K.; Ninomiya–Tsuji, J.; Goebel, M.;
Taniguchi,

[193387]T.; Matsumoto, K.: New human gene encoding a positive
modulator of

[193388]HIV Tat–mediated transactivation. Nature 357: 700–702,
1992.

[193389]

[193390]

[193391]20627.

[193392]Tanahashi, N.; Suzuki, M.; Fujiwara, T.; Takahashi, E.;
Shimbara,

[193393]N.; Chung, C. H.; Tanaka, K.: Chromosomal localization
and immunological

[193394]analysis of a family of human 26S proteasomal ATPases.
Biochem. Biophys.

[193395]Res. Commun. 243: 229–232, 1998.

[193396]

[193397]

[193398]20628.

[193399]Yanagi, S.; Shimbara, N.; Tamura, T.: Tissue and cell distribution

[193400]of a mammalian proteasomal ATPase, MSS1, and its complex formation

[193401]with the basal transcription factors. Biochem. Biophys. Res. Commun. 279:

[193402]568–573, 2000.

[193403]

[193404]

[193405]20629.

[193406]Levine, A.; Cantoni, G. L.; Razin, A.: Inhibition of promoter

[193407]activity by methylation: possible involvement of protein mediators. Proc.

[193408]Nat. Acad. Sci. 88: 6515–6518, 1991.

[193409]

[193410]

[193411]20630.

[193412]Nan, X.; Meehan, R. R.; Bird, A.: Dissection of the methyl-CpG

[193413]binding domain from the chromosomal protein MeCP2.
Nucleic Acids

[193414]Res. 21: 4886–4892, 1993.

[193415]

[193416]

[193417]20631.

[193418]Ohki, I.; Shimotake, N.; Fujita, N.; Jee, J.-G.; Ikegami, T.;
Nakao,

[193419]M.; Shirakawa, M.: Solution structure of the methyl-CpG
binding domain

[193420]of human MBD1 in complex with methylated DNA. Cell
105: 487–497,

[193421]2001.

[193422]

[193423]

[193424]20632.

[193425]Carrera, C. J.; Eddy, R. L.; Shows, T. B.; Carson, D. A.: As-

signment

[193426]of the gene for methylthioadenosine phosphorylase to human chromosome

[193427]9 by mouse-human somatic cell hybridization. Proc. Nat. Acad. Sci. 81:

[193428]2665-2668, 1984.

[193429]

[193430]

[193431]20633.

[193432]Chilcote, R. R.; Brown, E.; Rowley, J. D.: Lymphoblastic leukemia

[193433]with lymphomatous features associated with abnormalities of the short

[193434]arm of chromosome 9. New Eng. J. Med. 313: 286-291, 1985.

[193435]

[193436]

[193437]20634.

[193438]Nobori, T.; Takabayashi, K.; Tran, P.; Orvis, L.; Batova, A.; Yu,

[193439]A. L.; Carson, D. A.: Genomic cloning of methylthioadeno-

sine phosphorylase:

[193440]a purine metabolic enzyme deficient in multiple different cancers. Proc.

[193441]Nat. Acad. Sci. 93: 6203–6208, 1996.

[193442]

[193443]

[193444]20635.

[193445]Olopade, O. I.; Jenkins, R. B.; Ransom, D. T.; Malik, K.; Pomykala,

[193446]H.; Nobori, T.; Cowan, J. M.; Rowley, J. D.; Diaz, M. O.: Molecular

[193447]analysis of deletions of the short arm of chromosome 9 in human gliomas. Cancer

[193448]Res. 52: 2523–2529, 1992.

[193449]

[193450]

[193451]20636.

[193452]Olopade, O. I.; Pomykala, H. M.; Hagos, F.; Sveen, L. W.; Espinosa,

[193453]R., III; Dreyling, M. H.; Gursky, S.; Stadler, W. M.; Le Beau,

M.

[193454]M.; Bohlander, S. K.: Construction of a 2.8-megabase
yeast artificial

[193455]chromosome contig and cloning of the human methylthioadenosine phosphorylase

[193456]gene from the tumor suppressor region on 9p21. Proc.

Nat. Acad. Sci. 92:

[193457]6489–6493, 1995.

[193458]

[193459]

[193460]20637.

[193461]Ragione, F. D.; Takabayashi, K.; Mastropietro, S.; Mercurio,
C.;

[193462]Oliva, A.; Russo, G. L.; Pietra, V. D.; Borriello, A.; Nobori,
T.;

[193463]Carson, D. A.; Zappia, V.: Purification and characterization
of recombinant

[193464]human 5-prime-methylthioadenosine phosphorylase:
definite identification

[193465]of coding cDNA. Biochem. Biophys. Res. Commun. 223:
514–519, 1996.

[193466]

[193467]

[193468]20638.

[193469]Williams–Ashman, H. G.; Seidenfeld, J.; Galletti, P.: Trends
in

[193470]the biochemical pharmacology of

5–prime–deoxy–5–prime–methylthioadenosine. Biochem.

[193471]Pharm. 31: 277–288, 1982.

[193472]

[193473]

[193474]20639.

[193475]Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human
gene encoding

[193476]methionyl–tRNA synthetase to chromosome 12. (Abstract)
Cytogenet.

[193477]Cell Genet. 37: 437 only, 1984.

[193478]

[193479]

[193480]20640.

[193481]Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human

MARS gene,

[193482]encoding methioninyl-tRNA synthetase, to chromosome
12 using human

[193483]x Chinese hamster cell hybrids. Somat. Cell Molec. Genet.
10: 225-234,

[193484]1984.

[193485]

[193486]

[193487]20641.

[193488]Grewal, P. K.; Holzfeind, P. J.; Bittner, R. E.; Hewitt, J. E.

[193489]: Mutant glycosyltransferase and altered glycosylation of
alpha-dystroglycan

[193490]in the myodystrophy mouse. Nature Genet. 28: 151-154,
2001.

[193491]

[193492]

[193493]20642.

[193494]Zabel, B. U.; Eddy, R. L.; Scott, J.; Shows, T. B.: The human

[193495]nerve growth factor gene (NGF) is located on the short
arm of chromosome

[193496]1. (Abstract) Cytogenet. Cell Genet. 37: 614, 1984.

[193497]

[193498]

[193499]20643.

[193500]Adams, D. S.; Kiyokawa, M.; Getman, M. E.; Shashoua, V.
E.: Genes

[193501]encoding giant danio and golden shiner ependymin. Neu-
rochem. Res. 21:

[193502]377–384, 1996.

[193503]

[193504]

[193505]20644.

[193506]de Groen, P. C.; Eggen, B. J. L.; Gispen, W. H.; Schotman,
P.;

[193507]Schrama, L. H.: Cloning and promoter analysis of the hu-
man B-50/GAP-43

[193508]gene. J. Molec. Neurosci. 6: 109–119, 1995.

[193509]

[193510]

[193511]20645.

[193512]Kosik, K. S.; Orecchio, L. D.; Bruns, G. A. P.; Benowitz, L. I.;

[193513]MacDonald, G. P.; Cox, D. R.; Neve, R. L.: Human GAP-43:
its deduced

[193514]amino acid sequence and chromosomal localization in
mouse and human. Neuron 1:

[193515]127-132, 1988.

[193516]

[193517]

[193518]20646.

[193519]Reeves, R. H.; O'Hara, B. F.; Gearhart, J. D.: Localization on

[193520]mouse chromosome 16 of Smst and Gap43: a conserved
synteny with human

[193521]chromosome 3. (Abstract) Cytogenet. Cell Genet. 51:
1064, 1989.

[193522]

[193523]

[193524]20647.

[193525]Strittmatter, S. M.; Fankhauser, C.; Huang, P. L.; Mashimo,

H.;

[193526]Fishman, M. C.: Neuronal pathfinding is abnormal in mice lacking

[193527]the neuronal growth cone protein GAP-43. Cell 80: 445-452, 1995.

[193528]

[193529]

[193530]20648.

[193531]Bespalova, I. N.; Farjo, Q.; Mortlock, D. P.; Jackson, A. U.; Meisler,

[193532]M. H.; Swaroop, A.; Burmeister, M.: Mapping of the neural retina

[193533]leucine zipper gene, Nrl, to mouse chromosome 14. Mammalian Genome 4:

[193534]618-620, 1993.

[193535]

[193536]

[193537]20649.

[193538]Bessant, D. A. R.; Payne, A. M.; Mitton, K. P.; Wang, Q.-L.; Swain,

[193539]P. K.; Plant, C.; Bird, A. C.; Zack, D. J.; Swaroop, A.; Bhat-

tacharya,

[193540]S. S.: A mutation in NRL is associated with autosomal
dominant retinitis

[193541]pigmentosa. Nature Genet. 21: 355–356, 1999.

[193542]

[193543]

[193544]20650.

[193545]Bessant, D. A. R.; Payne, A. M.; Plant, C.; Bird, A. C.; Swa-
roop,

[193546]A.; Bhattacharya, S. S.: NRL S50T mutation and the impor-
tance of

[193547]'founder effects' in inherited retinal dystrophies. Europ. J.
Hum.

[193548]Genet. 8: 783–787, 2000.

[193549]

[193550]

[193551]20651.

[193552]Dahl, S. P.; Jackson, A.; Kimberling, W. J.; Blackwood, D.;
Swaroop,

[193553]A.: Genetic mapping of NRL, a human retina-specific gene

located

[193554]on chromosome 14. (Abstract) Am. J. Hum. Genet. 51
(suppl.): A185

[193555]only, 1992.

[193556]

[193557]

[193558]20652.

[193559]Farjo, Q.; Jackson, A.; Pieke-Dahl, S.; Scott, K.; Kimberling,

[193560]W. J.; Sieving, P. A.; Richards, J. E.; Swaroop, A.: Human
bZIP transcription

[193561]factor gene NRL: structure, genomic sequence, and fine
linkage mapping

[193562]at 14q11.2 and negative mutation analysis in patients
with retinal

[193563]degeneration. Genomics 45: 395-401, 1997.

[193564]

[193565]

[193566]20653.

[193567]Farjo, Q.; Jackson, A. U.; Xu, J.; Gryzenia, M.; Skolnick, C.;

[193568]Agarwal, N.; Swaroop, A.: Molecular characterization of
the murine

[193569]neural retina leucine zipper gene, Nrl. Genomics 18:
216–222, 1993.

[193570]

[193571]

[193572]20654.

[193573]Mears, A. J.; Kondo, M.; Swain, P. K.; Takada, Y.; Bush, R.
A.;

[193574]Saunders, T. L.; Sieving, P. A.; Swaroop, A.: Nrl is required
for

[193575]rod photoreceptor development. Nature Genet. 29:
447–452, 2001.

[193576]

[193577]

[193578]20655.

[193579]Swaroop, A.; Xu, J.; Agarwal, N.; Weissman, S. M.: A simple
and

[193580]efficient cDNA library subtraction procedure: isolation of
human retina-specific

[193581]cDNA clones. Nucleic Acids Res. 19: 1954 only, 1991.

[193582]

[193583]

[193584]20656.

[193585]Swaroop, A.; Xu, J.; Pawar, H.; Jackson, A.; Skolnick, C.;
Agarwal,

[193586]N.: A conserved retina-specific gene encodes a basic motif/leucine

[193587]zipper domain. Proc. Nat. Acad. Sci. 89: 266-270, 1992.

[193588]

[193589]

[193590]20657.

[193591]Wright, A. F.; Mansfield, D. C.; Bruford, E. A. Teague, P.
W.;

[193592]Thomson, K. L.; Riise, R.; Jay, M.; Patton, M. A.; Jeffery, S.;
Schinzel,

[193593]A.; Tommerup, N.: Fossarello, M.: Genetic studies in autosomal recessive

[193594]forms of retinitis pigmentosa. In: Anderson, R. E.; LaVail,
M. M.;

[193595]Hollyfield, J. G. (eds.): Degenerative diseases of the retina.

[193596]New York: Plenum Press (pub.) 1995. Pp. 293–302.

[193597]

[193598]

[193599]20658.

[193600]Yang–Feng, T. L.; Swaroop, A.: Neural retina–specific
leucine

[193601]zipper gene NRL (D14S46E) maps to human chromosome
14q11.1–q11.2. Genomics 14:

[193602]491–492, 1992.

[193603]

[193604]

[193605]20659.

[193606]Eddy, R. L.; Kretschmer, P. J.; Fairhurst, J. L.; Shows, T. B.;

[193607]Bohlen, P.; O'Hara, B.; Kovesdi, I.: A human gene family of
neurite

[193608]outgrowth–promoting proteins: the gene for a heparin
binding neurite

[193609]outgrowth–promoting factor maps to 7q22–qter.
(Abstract) Cytogenet.

[193610]Cell Genet. 58: 1920 only, 1991.

[193611]

[193612]

[193613]20660.

[193614]Lai, S.; Czubayko, F.; Riegel, A. T.; Wellstein, A.: Structure

[193615]of the human heparin-binding growth factor gene
pleiotrophin. Biochem.

[193616]Biophys. Res. Commun. 187: 1113-1122, 1992.

[193617]

[193618]

[193619]20661.

[193620]Li, Y.-S.; Hoffman, R. M.; Le Beau, M. M.; Espinosa, R., III;
Jenkins,

[193621]N. A.; Gilbert, D. J.; Copeland, N. G.; Deuel, T. F.: Charac-
terization

[193622]of the human pleiotrophin gene: promoter region and
chromosomal localization. J.

[193623]Biol. Chem. 267: 26011-26016, 1992.

[193624]

[193625]

[193626]20662.

[193627]Li, Y.-S.; Milner, P. G.; Chauhan, A. K.; Watson, M. A.;
Hoffman,

[193628]R. M.; Kodner, C. M.; Milbrandt, J.; Deuel, T. F.: Cloning
and expression

[193629]of a developmentally regulated protein that induces mitogenic and

[193630]neurite outgrowth activity. Science 250: 1690–1694, 1990.

[193631]

[193632]

[193633]20663.

[193634]Milner, P. G.; Shah, D.; Veile, R.; Donis–Keller, H.; Kumar, B.

[193635]V.: Cloning, nucleotide sequence, and chromosome localization of

[193636]the human pleiotrophin gene. Biochemistry 31: 12023–12028, 1992.

[193637]

[193638]

[193639]20664.

[193640]Brightbill, H. D.; Libraty, D. H.; Krutzik, S. R.; Yang, R.–B.;

[193641]Bellsie, J. T.; Bieharski, J. R.; Maitland, M.; Norgard, M. V.; Plevy,

[193642]S. E.; Smale, S. T.; Brennan, P. J.; Bloom, B. R.; Godowski,

P. J.;

[193643]Modlin, R. L.: Host defense mechanisms triggered by microbial lipoproteins

[193644]through toll-like receptors. Science 285: 732–736, 1999.

[193645]

[193646]

[193647]20665.

[193648]Jankovic, D.; Kullberg, M. C.; Hieny, S.; Caspar, P.; Collazo,

[193649]C. M.; Sher, A.: In the absence of IL-12, CD4+ T cell responses to

[193650]intracellular pathogens fail to default to a Th2 pattern and are host

[193651]protective in an IL-10-/- setting. Immunity 16: 429–439, 2002.

[193652]

[193653]

[193654]20666.

[193655]Jouanguy, E.; Lamhamedi-Cherradi, S.; Altare, F.; Fondaneche,

[193656]M.-C.; Tuerlinckx, D.; Blanche, S.; Emile, J.-F.; Gaillard,

J.-L.;

[193657]Schreiber, R.; Levin, M.; Fischer, A.; Hivroz, C.; Casanova,
J.-L.

[193658]: Partial interferon-gamma receptor 1 deficiency in a child
with tuberculoid

[193659]bacillus Calmette-Guerin infection and a sibling with clinical tuberculosis. J.

[193660]Clin. Invest. 100: 2658-2664, 1997.

[193661]

[193662]

[193663]20667.

[193664]Warrington, J. A.; Bengtsson, U.: High-resolution physical
mapping

[193665]of human 5q31-q33 using three methods: radiation hybrid
mapping, interphase

[193666]fluorescence in situ hybridization, and pulsed-field gel
electrophoresis. Genomics 24:

[193667]395-398, 1994.

[193668]

[193669]

[193670]20668.

[193671]Anderson, S. K.; Gallinger, S.; Roder, J.; Frey, J.; Young, H.

[193672]A.; Ortaldo, J. R.: A cyclophilin-related protein involved in
the

[193673]function of natural killer cells. Proc. Nat. Acad. Sci. 90:
542-546,

[193674]1993.

[193675]

[193676]

[193677]20669.

[193678]Frey, J. L.; Bino, T.; Kantor, R. R. S.; Segal, D. M.; Giardina,

[193679]S. L.; Roder, J.; Anderson, S.; Ortaldo, J. R.: Mechanism of
target

[193680]cell recognition by natural killer cells: characterization of a
novel

[193681]triggering molecule restricted to CD3- large granular lym-
phocytes. J.

[193682]Exp. Med. 174: 1527-1536, 1991.

[193683]

[193684]

[193685]20670.

[193686]Rinfret, A.; Anderson, S. K.: IL-2 regulates the expression
of

[193687]the NK-TR gene via an alternate RNA splicing mechanism.
Molec. Immun. 30:

[193688]1307-1313, 1993.

[193689]

[193690]

[193691]20671.

[193692]Simons-Evelyn, M.; Young, H. A.; Anderson, S. K.: Charac-
terization

[193693]of the mouse Nktr gene and promoter. Genomics 40:
94-100, 1997.

[193694]

[193695]

[193696]20672.

[193697]Young, H. A.; Jenkins, N. A.; Copeland, N. G.; Simek, S.;
Lerman,

[193698]M. I.; Zbar, B.; Glenn, G.; Ortaldo, J. R.; Anderson, S. K.:
Localization

[193699]of a novel natural killer triggering receptor locus to hu-
man chromosome

[193700]3p23–p21 and mouse chromosome 9. Genomics 16:
548–549, 1993.

[193701]

[193702]

[193703]20673.

[193704]Scott, I. C.; Halila, R.; Jenkins, J. M.; Mehan, S.; Apostolou,

[193705]S.; Winqvist, R.; Callen, D. F.; Prockop, D. J.; Peltonen, L.;
Kadler,

[193706]K. E.: Molecular cloning, expression and chromosomal lo-
calization

[193707]of a human gene encoding a 33 kDa putative metallopep-
tidase (PRSM1). Gene 174:

[193708]135–143, 1996.

[193709]

[193710]

[193711]20674.

[193712]Aliprantis, A. O.; Yang, R.–B.; Mark, M. R.; Suggett, S.; De-
vaux,

[193713]B.; Radolf, J. D.; Klimpel, G. R.; Godowski, P.; Zychlinsky,
A.:

[193714]Cell activation and apoptosis by bacterial lipoproteins
through toll-like

[193715]receptor-2. Science 285: 736-739, 1999.

[193716]

[193717]

[193718]20675.

[193719]Baldwin, A. S., Jr.: The NF-kappa-B and I-kappa-B pro-
teins: new

[193720]discoveries and insights. Annu. Rev. Immun. 14: 649-683,
1996.

[193721]

[193722]

[193723]20676.

[193724]Barnes, P. J.; Karin, M.: Nuclear factor-kappa-B -- A piv-
otal

[193725]transcription factor in chronic inflammatory diseases. New
Eng. J.

[193726]Med. 336: 1066-1071, 1997.

[193727]

[193728]

[193729]20677.

[193730]Chen, F.; Castranova, V.; Shi, X.; Demers, L. M.: New insights

[193731]into the role of nuclear factor- κ B, a ubiquitous transcription

[193732]factor in the initiation of diseases. Clin. Chem. 45: 7–17, 1999.

[193733]

[193734]

[193735]20678.

[193736]Heron, E.; Deloukas, P.; van Loon, A. P. G. M.: The complete

[193737]exon–intron structure of the 156–kb human gene NFKB1, which encodes

[193738]the p105 and p50 proteins of transcription factors NF- κ B and

[193739]I- κ B- γ : implications for NF-

κ B-mediated signal transduction. Genomics 30:

[193740]493–505, 1995.

[193741]

[193742]

[193743]20679.

[193744]Huxford, T.; Huang, D.-B.; Malek, S.; Ghosh, G.: The crystal
tal

[193745]structure of the I-kappa-B-alpha/NF-kappa-B complex
reveals mechanisms

[193746]of NF-kappa-B inactivation. Cell 95: 759-770, 1998.

[193747]

[193748]

[193749]20680.

[193750]Lin, L.; DeMartino, G. N.; Greene, W. C.: Cotranslational
biogenesis

[193751]of NF-kappaB p50 by the 26S proteasome. Cell 92:
819-828, 1998.

[193752]

[193753]

[193754]20681.

[193755]Liptay, S.; Schmid, R. M.; Perkins, N. D.; Meltzer, P.; Al-
therr,

[193756]M. R.; McPherson, J. D.; Wasmuth, J. J.; Nabel, G. J.: Related
subunits

[193757]of NF-kappa-B map to two distinct loci associated with translocations

[193758]in leukemia, NFKB1 and NFKB2. Genomics 13: 287-292, 1992.

[193759]

[193760]

[193761]20682.

[193762]Mathew, S.; Murty, V. V. V. S.; Dalla-Favera, R.; Chaganti, R.

[193763]S. K.: Chromosomal localization of genes encoding the transcription

[193764]factors, c-rel, NF-kappa-Bp50, NF-kappa-Bp65, and lyt-10 by fluorescence

[193765]in situ hybridization. Oncogene 8: 191-193, 1993.

[193766]

[193767]

[193768]20683.

[193769]Meyer, R.; Hatada, E. N.; Hohmann, H. P.; Haiker, M.; Bartsch,

[193770]C.; Rothlisberger, U.; Lahm, H. W.; Schlaeger, E. J.; van Loon, A.

[193771]P.; Scheidereit, C.: Cloning of the DNA-binding subunit of human

[193772]nuclear factor kappa B: the level of its mRNA is strongly regulated

[193773]by phorbol ester or tumor necrosis factor alpha. Proc. Nat. Acad.

[193774]Sci. 88: 966–970, 1991.

[193775]

[193776]

[193777]20684.

[193778]Ozes, O. N.; Mayo, L. D.; Gustin, J. A.; Pfeffer, S. R.; Pfeffer,

[193779]L. M.; Donner, D. B.: NF-kappa-B activation by tumour necrosis factor

[193780]requires the Akt serine-threonine kinase. Nature 401: 82–85, 1999.

[193781]

[193782]

[193783]20685.

[193784]Romashkova, J. A.; Makarov, S. S.: NF-kappa-B is a target of

[193785]AKT in anti-apoptotic PDGF signalling. Nature 401:
86–90, 1999.

[193786]

[193787]

[193788]20686.

[193789]Ryan, K. M.; Ernst, M. K.; Rice, N. R.; Vousden, K. H.: Role
[193790]of NF-kappa-B in p53-mediated programmed cell death.
Nature 404:

[193791]892–897, 2000.

[193792]

[193793]

[193794]20687.

[193795]Sen, R.; Baltimore, D.: Multiple nuclear factors interact
with

[193796]the immunoglobulin enhancer sequences. Cell 46:
705–716, 1986.

[193797]

[193798]

[193799]20688.

[193800]Sha, W. C.; Liou, H.-C.; Tuomanen, E. I.; Baltimore, D.:

Targeted

[193801]disruption of the p50 subunit of NF-kappa-B leads to
multifocal defects

[193802]in immune responses. Cell 80: 321-330, 1995.

[193803]

[193804]

[193805]20689.

[193806]Zhong, H.; May, M. J.; Jimi, E.; Ghosh, S.: The phosphory-
lation

[193807]status of nuclear NF-kappa-B determines its association
with CBP/p300

[193808]or HDAC-1. Molec. Cell 9: 625-636, 2002.

[193809]

[193810]

[193811]20690.

[193812]Claudio, E.; Brown, K.; Park, S.; Wang, H.; Siebenlist, U.:
BAFF-induced

[193813]NEMO-independent processing of NF-kappa-B2 in matur-
ing B cells. Nature

[193814]Immun. 3: 958–965, 2002.

[193815]

[193816]

[193817]20691.

[193818]Neri, A.; Chang, C.–C.; Lombardi, L.; Salina, M.; Corradini, P.;

[193819]Maiolo, A. T.; Chaganti, R. S. K.; Dalla–Favera, R.: B cell lymphoma–associated

[193820]chromosomal translocation involves candidate oncogene lyt–10, homologous

[193821]to NF–kappa–B p50. Cell 67: 1075–1087, 1991.

[193822]

[193823]

[193824]20692.

[193825]Taparowsky, E.; Shimizu, K.; Goldfarb, M.; Wigler, M.: Structure

[193826]and activation of the human N–ras gene. Cell 34: 581–586, 1983.

[193827]

[193828]

[193829]20693.

[193830]Yuasa, Y.; Gol, R. A.; Chang, A.; Chiu, I.-M.; Reddy, E. P.;
Tronick,

[193831]S. R.; Aaronson, S. A.: Mechanism of activation of an N-ras
oncogene

[193832]of SW-1271 human lung carcinoma cells. Proc. Nat. Acad.
Sci. 81:

[193833]3670-3674, 1984.

[193834]

[193835]

[193836]20694.

[193837]Ritzler, J. M.; Sawhney, R.; Geurts van Kessel, A. H. M.;
Grzeschik,

[193838]K.-H.; Schinzel, A.; Berchtold, M. W.: The genes for the
highly homologous

[193839]Ca(2+)-binding proteins oncomodulin and parvalbumin
are not linked

[193840]in the human genome. Genomics 12: 567-572, 1992.

[193841]

[193842]

[193843]20695.

[193844]Staubli, F.; Klein, A.; Rentsch, J. M.; Hameister, H.; Berchtold,

[193845]M. W.: Structure and chromosomal localization of the mouse oncomodulin

[193846]gene. Mammalian Genome 6: 769–777, 1995.

[193847]

[193848]

[193849]20696.

[193850]Bakin, A. V.; Curran, T.: Role of DNA 5–methylcytosine transferase

[193851]in cell transformation by fos. Science 283: 387–390, 1999.

[193852]

[193853]

[193854]20697.

[193855]Barker, P. E.; Rabin, M.; Watson, M.; Breg, W. R.; Ruddle, F. H.;

[193856]Verma, I. M.: Human c–fos oncogene mapped within chromosomal region

[193857]14q21–q31. Proc. Nat. Acad. Sci. 81: 5826–5830, 1984.

[193858]

[193859]

[193860]20698.

[193861]Dony, C.; Gruss, P.: Proto-oncogene c-fos expression in
growth

[193862]regions of fetal bone and mesodermal web tissue. Nature
328: 711–714,

[193863]1987.

[193864]

[193865]

[193866]20699.

[193867]Ekstrand, A. J.; Zech, L.: Human c-fos proto-oncogene
mapped to

[193868]chromosome 14, band q24.3–q31: possibilities for onco-
gene activation

[193869]by chromosomal rearrangements in human neoplasms.

Exp. Cell Res. 169:

[193870]262–266, 1987.

[193871]

[193872]

[193873]20700.

[193874]Glover, J. N. M.; Harrison, S. C.: Crystal structure of the
heterodimeric

[193875]bZIP transcription factor c-Fos--c-Jun bound to DNA. Na-
ture 373:

[193876]257-261, 1995.

[193877]

[193878]

[193879]20701.

[193880]Grigoriadis, A. E.; Wang, Z.-Q.; Cecchini, M. G.; Hofstetter,
W.;

[193881]Felix, R.; Fleisch, H. A.; Wagner, E. F.: c-Fos: a key regula-
tor

[193882]of osteoclast-macrophage lineage determination and
bone remodeling. Science 266:

[193883]443-448, 1994.

[193884]

[193885]

[193886]20702.

[193887]Grigoriadis, A. E.; Wang, Z.-Q.; Wagner, E. F.: Fos and bone cell

[193888]development: lessons from a nuclear oncogene. Trends Genet. 11:

[193889]:436–441, 1995.

[193890]

[193891]

[193892]20703.

[193893]Hayflick, L.: The limited in vitro lifetime of human diploid cell

[193894]strains. Exp. Cell Res. 37: 614–636, 1965.

[193895]

[193896]

[193897]20704.

[193898]Johnson, R. S.; Spiegelman, B. M.; Papaioannou, V.: Pleiotropic

[193899]effects of a null mutation in the c-fos proto-oncogene. Cell 71:

[193900]577–586, 1992.

[193901]

[193902]

[193903]20705.

[193904]Muller, R.; Tremblay, J. M.; Adamson, E. D.; Verma, I. M.:

Tissue

[193905]and cell type-specific expression of two human c-onc

genes. Nature 304:

[193906]454-456, 1983.

[193907]

[193908]

[193909]20706.

[193910]Rogaev, E. I.; Lukiw, W. J.; Vaula, G.; Haines, J. L.; Rogaeva,

[193911]E. A.; Tsuda, T.; Alexandrova, N.; Liang, Y.; Mortilla, M.;

Amaducci,

[193912]L.; Bergamini, L.; Bruni, A. C.; Foncin, J.-F.; Macciardi, F.;

Montesi,

[193913]M. P.; Sorbi, S.; Rainero, I.; Pinessi, L.; Polinsky, R. J.;

Frommelt,

[193914]P.; Duara, R.; Lopez, R.; Pollen, D.; Gusella, J. F.; Tanzi, R.;

Crapper

[193915]MacLachlan, D.; St. George-Hyslop, P. H.: Analysis of the

c-FOS gene

[193916]on chromosome 14 and the promoter of the amyloid precursor protein

[193917]gene in familial Alzheimer's disease. Neurology 43: 2275–2279, 1993.

[193918]

[193919]

[193920]20707.

[193921]Ruther, U.; Garber, C.; Komitowski, D.; Muller, R.; Wagner, E.

[193922]F.: Deregulated c-fos expression interferes with normal bone development

[193923]in transgenic mice. Nature 325: 412–416, 1987.

[193924]

[193925]

[193926]20708.

[193927]Saez, E.; Rutberg, S. E.; Mueller, E.; Oppenheim, H.; Smoluk,

[193928]J.; Yuspa, S. H.; Spiegelman, B. M.: c-fos is required for malignant

[193929]progression of skin tumors. Cell 82: 721–732, 1995.

[193930]

[193931]

[193932]20709.

[193933]Sanyal, S.; Sandstrom, D. J.; Hoeffler, C. A.; Ramaswami, M.:

[193934]AP-1 function upstream of CREB to control synaptic plasticity in Drosophila. Nature 416:

[193935]870–874, 2002.

[193936]

[193937]

[193938]20710.

[193939]Seshadri, T.; Campisi, J.: Repression of c-fos transcription

[193940]and an altered genetic program in senescent human fibroblasts. Science 247:

[193941]205–209, 1990.

[193942]

[193943]

[193944]20711.

[193945]van Straaten, F.; Muller, R.; Curran, T.; Van Beveren, C.;

Verma,

[193946]I. M.: Complete nucleotide sequence of a human c-onc gene: deduced

[193947]amino acid sequence of the human c-fos protein. Proc. Nat. Acad.

[193948]Sci. 80: 3183–3187, 1983.

[193949]

[193950]

[193951]20712.

[193952]Visvader, J.; Sassone-Corsi, P.; Verma, I. M.: Two adjacent promoter

[193953]elements mediate nerve growth factor activation of the c-fos gene

[193954]and bind distinct nuclear complexes. Proc. Nat. Acad. Sci. 85: 9474–9478,

[193955]1988.

[193956]

[193957]

[193958]20713.

[193959]Wang, Z.-Q.; Grigoriadis, A. E.; Mohle-Steinlein, U.; Wag-

ner,

[193960]E. F.: A novel target cell for c-fos-induced oncogenesis:
development

[193961]of chondrogenic tumours in embryonic stem cell
chimeras. EMBO J. 10:

[193962]2437–2450, 1991.

[193963]

[193964]

[193965]20714.

[193966]Wang, Z.-Q.; Ovitt, C.; Grigoriadis, A. E.; Mohle-Steinlein,
U.;

[193967]Ruther, U.; Wagner, E. F.: Bone and haematopoietic defects
in mice

[193968]lacking c-fos. Nature 360: 741–745, 1992.

[193969]

[193970]

[193971]20715.

[193972]Zhang, J.; Zhang, D.; McQuade, J. S.; Behbehani, M.; Tsien,
J.

[193973]Z.; Xu, M.: c-fos regulates neuronal excitability and sur-

vival. Nature

[193974]Genet. 30: 416–420, 2002.

[193975]

[193976]

[193977]20716.

[193978]Arheden, K.; Mandahl, N.; Strombeck, B.; Isaksson, M.;
Mitelman,

[193979]F.: Chromosome localization of the human oncogene INT1
to 12q13 by

[193980]in situ hybridization. Cytogenet. Cell Genet. 47: 86–87,
1988.

[193981]

[193982]

[193983]20717.

[193984]Gavin, B. J.; McMahon, J. A.; McMahon, A. P.: Expression of
multiple

[193985]novel Wnt–1/int–1–related genes during fetal and adult
mouse development. Genes

[193986]Dev. 4: 2319–2332, 1990.

[193987]

[193988]

[193989]20718.

[193990]Kirikoshi, H.; Sekihara, H.; Katoh, M.: WNT10A and WNT6,
clustered

[193991]in human chromosome 2q35 region with head-to-tail
manner, are strongly

[193992]coexpressed in SW480 cells. Biochem. Biophys. Res. Com-
mun. 283:

[193993]798-805, 2001.

[193994]

[193995]

[193996]20719.

[193997]McMahon, A. P.: The Wnt family of developmental regula-
tors. Trends

[193998]Genet. 8: 236-242, 1992.

[193999]

[194000]

[194001]20720.

[194002]Nusse, R.; Brown, A.; Papkoff, J.; Scambler, P.; Shackleford,

G.;

[194003]McMahon, A.; Moon, R.; Varmus, H.: A new nomenclature
for int-1 and

[194004]related genes: the Wnt gene family. (Letter) Cell 64:
231-232, 1991.

[194005]

[194006]

[194007]20721.

[194008]Nusse, R.; van't Veer, L.; Geurts van Kessel, A.; van Agth-
oven,

[194009]A.; Bootsma, D.; Varmus, H.: Chromosomal localization of
a human

[194010]homologue of a putative mammary tumor oncogene.
(Abstract) Cytogenet.

[194011]Cell Genet. 37: 556-557, 1984.

[194012]

[194013]

[194014]20722.

[194015]Pellegrino, J. E.; Lensch, M. W.; Muenke, M.; Chance, P. F.:
Clinical

[194016]and molecular analysis in Joubert syndrome. Am. J. Med.

Genet. 72:

[194017]59–62, 1997.

[194018]

[194019]

[194020]20723.

[194021]Thomas, K. R.; Capecchi, M. R.: Targeted disruption of the
murine

[194022]int-1 proto-oncogene resulting in severe abnormalities in
midbrain

[194023]and cerebellar development. Nature 346: 847–850, 1990.

[194024]

[194025]

[194026]20724.

[194027]Braaten, D.; Wellington, S.; Warburton, D.; Luban, J.: As-
signment

[194028]of cyclophilin A (PPIA) to human chromosome band 7p13
by in situ hybridization. Cytogenet.

[194029]Cell Genet. 74: 262 only, 1996.

[194030]

[194031]

[194032]20725.

[194033]Liu, J.; Albers, M. W.; Chen, C.-M.; Schreiber, S. L.; Walsh, C.

[194034]T.: Cloning, expression, and purification of human cyclophilin in

[194035]Escherichia coli and assessment of the catalytic role of cysteines

[194036]by site-directed mutagenesis. Proc. Nat. Acad. Sci. 87: 2304-2308,

[194037]1990.

[194038]

[194039]

[194040]20726.

[194041]Luban, J.; Bossolt, K. L.; Franke, E. K.; Kalpana, G. V.; Goff,

[194042]S. P.: Human immunodeficiency virus type 1 gag protein binds to cyclophilins

[194043]A and B. Cell 73: 1067-1078, 1993.

[194044]

[194045]

[194046]20727.

[194047]Takahashi, N.; Hayano, T.; Suzuki, M.: Peptidyl-prolyl cis-trans

[194048]isomerase is the cyclosporin A-binding protein cyclophilin. Nature 337:

[194049]473-475, 1989.

[194050]

[194051]

[194052]20728.

[194053]Willenbrink, W.; Halaschek, J.; Schuffenhauer, S.; Kunz, J.; Steinkasserer,

[194054]A.: Cyclophilin A, the major intracellular receptor for the immunosuppressant

[194055]cyclosporin A, maps to chromosome 7p11.2-p13: four pseudogenes map

[194056]to chromosomes 3, 10, 14, and 18. Genomics 28: 101-104, 1995.

[194057]

[194058]

[194059]20729.

[194060]Friedman, J.; Weissman, I.: Two cytoplasmic candidates for immunophilin

[194061]action are revealed by affinity for a new cyclophilin: one in the

[194062]presence and one in the absence of CsA. Cell 66: 799–806, 1991.

[194063]

[194064]

[194065]20730.

[194066]Liu, J.; Farmer, J. D., Jr.; Lane, W. S.; Friedman, J.; Weissman,

[194067]I.; Schreiber, S. L.: Calcineurin is a common target of cyclophilin–cyclosporin

[194068]A and FKBP–FK506 complexes. Cell 66: 807–815, 1991.

[194069]

[194070]

[194071]20731.

[194072]Dickinson, D. P.; Zhao, Y.; Thiesse, M.; Siciliano, M. J.: Direct

[194073]mapping of seven genes encoding human type 2 cystatins to a single

[194074]site located at 20p11.2. Genomics 24: 172–175, 1994.

[194075]

[194076]

[194077]20732.

[194078]Saitoh, E.; Isemura, S.; Sanada, K.; Ohnishi, K.: The human
cystatin

[194079]gene family: cloning of three members and evolutionary
relationship

[194080]between cystatins and Bowman–Birk type proteinase in-
hibitors. Biomed.

[194081]Biochim. Acta 50: 599–605, 1991.

[194082]

[194083]

[194084]20733.

[194085]Freije, J. P.; Pendas, A. M.; Velasco, G.; Roca, A.; Abraham-
son,

[194086]M.; Lopez–Otin, C.: Localization of the human cystatin D
gene (CST5)

[194087]to chromosome 20p11.21 by in situ hybridization. Cyto-
genet. Cell

[194088]Genet. 62: 29–31, 1993.

[194089]

[194090]

[194091]20734.

[194092]Takahashi, E.; Hori, T.; O'Connell, P.; Leppert, M.; White, R.

[194093]: R-banding and nonisotopic in situ hybridization: precise
localization

[194094]of the human type II collagen gene (COL2A1). Hum. Genet.
86: 14–16,

[194095]1990.

[194096]

[194097]

[194098]20735.

[194099]Takahashi, E.; Yamauchi, M.; Tsuji, H.; Hitomi, A.; Meuth,
M.;

[194100]Hori, T.: Chromosome mapping of the human cytidine-
5-prime-triphosphate

[194101]synthetase (CTPS) gene to band 1p34.1–p34.3 by fluores-
cence in situ

[194102]hybridization. Hum. Genet. 88: 119–121, 1991.

[194103]

[194104]

[194105]20736.

[194106]Takahashi, E.-I.; Yamauchi, M.; Ayusawa, D.; Kaneda, S.;
Seno,

[194107]T.; Meuth, M.; Hori, T.-A.: Chromosome mappings of the
human cytidine-5-prime-triphosphate

[194108]synthetase (CTPS) gene and the human ubiquitin-activat-
ing enzyme UBE1

[194109]gene by fluorescence in situ hybridization. (Abstract) Cy-
togenet.

[194110]Cell Genet. 58: 1864 only, 1991.

[194111]

[194112]

[194113]20737.

[194114]Thomas, P. E.; Sen, S.; Lamb, B. J.; Chu, E. H. Y.: Cloning
and

[194115]expression of mammalian CTP synthetase genes.
(Abstract) Am. J. Hum.

[194116]Genet. 45 (suppl.): A11 only, 1989.

[194117]

[194118]

[194119]20738.

[194120]Whelan, J.; Phear, G.; Yamauchi, M.; Meuth, M.: Clustered
base

[194121]substitutions in CTP synthetase conferring drug resistance
in Chinese

[194122]hamster ovary cells. Nature Genet. 3: 317–322, 1993.

[194123]

[194124]

[194125]20739.

[194126]Yamauchi, M.; Takahashi, E.; Whelan, J.; Phear, G.; Meuth,
M.:

[194127]Mapping and functional analysis of the cytidine triphos-
phate synthetase

[194128](CTPS) gene. (Abstract) Human Genome Mapping Work-
shop 93 1 only,

[194129]1993.

[194130]

[194131]

[194132]20740.

[194133]Nebert, D. W.; Gonzalez, F. J.: P450 genes: structure, evolution,

[194134]and regulation. Annu. Rev. Biochem. 56: 945–993, 1987.

[194135]

[194136]

[194137]20741.

[194138]Paolini, M.; Cantelli–Forti, G.; Perocco, P.; Pedulli, G. F.;

[194139]Abdel–Rahman, S. Z.; Legator, M. S.: Co–carcinogenic effect of beta–carotene.

[194140](Letter) Nature 398: 760–761, 1999.

[194141]

[194142]

[194143]20742.

[194144]Espinoza, H.; Cox, C. J.; Semina, E. V.; Amendt, B. A.: A molecular

[194145]basis for differential developmental anomalies in Axenfeld–Rieger

[194146]syndrome. Hum. Molec. Genet. 11: 743–753, 2002.

[194147]

[194148]

[194149]20743.

[194150]Letinic, K.; Zoncu, R.; Rakic, P.: Origin of GABAergic neurons
rons

[194151]in the human neocortex. Nature 417: 645–649, 2002.

[194152]

[194153]

[194154]20744.

[194155]McGuinness, T.; Porteus, M. H.; Smiga, S.; Bulfone, A.;
Kingsley,

[194156]C.; Qiu, M.; Liu, J. K.; Long, J. E.; Xu, D.; Rubenstein, J. L. R.

[194157]: Sequence, organization, and transcription of the Dlx–1
and the Dlx–2

[194158]locus. Genomics 35: 473–485, 1996.

[194159]

[194160]

[194161]20745.

[194162]Ozcelik, T.; Porteus, M. H.; Rubenstein, J. L. R.; Francke, U.

[194163]: DLX2 (TES1), a homeobox gene of the Distal-less family,
assigned

[194164]to conserved regions on human and mouse chromosomes

2. Genomics 13:

[194165]1157–1161, 1992.

[194166]

[194167]

[194168]20746.

[194169]Porteus, M. H.; Brice, A. E. J.; Bulfone, A.; Usdin, T. B.; Cia-
ranello,

[194170]R. D.; Rubenstein, J. L. R.: Isolation and characterization of
a

[194171]library of cDNA clones that are preferentially expressed in
the embryonic

[194172]telencephalon. Molec. Brain Res. 12: 7–22, 1992.

[194173]

[194174]

[194175]20747.

[194176]Qiu, M.; Bulfone, A.; Martinez, S.; Meneses, J. J.; Shima-
mura,

[194177]K.; Pedersen, R. A.; Rubenstein, J. L.: Null mutation of Dlx-2 results

[194178]in abnormal morphogenesis of proximal first and second branchial arch

[194179]derivatives and abnormal differentiation in the forebrain.

Genes

[194180]Dev. 9: 2523–2538, 1995.

[194181]

[194182]

[194183]20748.

[194184]Salvador, J. M.; Hollander, M. C.; Nguyen, A. T.; Kopp, J. B.;

[194185]Barisoni, L.; Moore, J. K.; Ashwell, J. D.; Fornace, A. J., Jr.:

[194186]Mice lacking the p53–effector gene Gadd45a develop a lupus–like syndrome. Immunity 16:

[194187]499–508, 2002.

[194188]

[194189]

[194190]20749.

[194191]Takekawa, M.; Saito, H.: A family of stress–inducible

GADD45-like

[194192]proteins mediate activation of the stress-responsive

MTK1/MEKK4 MAPKKK. Cell 95:

[194193]521–530, 1998.

[194194]

[194195]

[194196]20750.

[194197]Dhermy, D.; Garbarz, M.; Lecomte, M. C.; Feo, C.;

Bournier, O.;

[194198]Chaveroche, I.; Gautero, H.; Galand, C.; Boivin, P.: Heredi-

tary elliptocytosis:

[194199]clinical, morphological and biochemical studies of 38

cases. Nouv.

[194200]Rev. Franc. Hemat. 28: 129–140, 1986.

[194201]

[194202]

[194203]20751.

[194204]Fairbanks, G.; Steck, T. L.; Wallach, D. F. H.: Elec-

trophoretic

[194205]analysis of the major polypeptides of the human erythro-

cyte membrane. Biochemistry 10:

[194206]2606–2617, 1971.

[194207]

[194208]

[194209]20752.

[194210]Feo, C. J.; Fischer, S.; Piau, J. P.; Grange, M. J.; Tchernia,

[194211]G.: Premiere observation de l'absence d'une proteine de la
membrane

[194212]erythrocytaire (bande 4–1) dans un cas d'anemie ellipto-
cytaire familiale. Nouv.

[194213]Rev. Franc. Hemat. 22: 315–325, 1980.

[194214]

[194215]

[194216]20753.

[194217]Eddy, R. L.; Mahnke–Zizelman, D. K.; Bausch–Jurken, M.
T.; Sabina,

[194218]R. L.; Shows, T. B.: Distribution of the AMP deaminase
multigene

[194219]family within the human genome: assignment of the
AMPD2 to chromosome

[194220]1p21–p34 and AMPD3 to chromosome 11p13–pter.

(Abstract) Human Genome

[194221]Mapping Workshop 93 24 only, 1993.

[194222]

[194223]

[194224]20754.

[194225]Ogasawara, N.; Goto, H.; Yamada, Y.; Nishigaki, I.; Itoh, T.;
Hasegawa,

[194226]I.; Park, K. S.: Deficiency of AMP deaminase in erythro-
cytes. Hum.

[194227]Genet. 75: 15–18, 1987.

[194228]

[194229]

[194230]20755.

[194231]Sermsuvitayawong, K.; Wang, X.; Nagabukuro, A.; Mat-
suda, Y.; Morisaki,

[194232]H.; Toyama, K.; Mukai, T.; Morisaki, T.: Genomic organi-
zation of

[194233]Ampd3, heart-type AMPD gene, located in mouse chro-
mosome 7. Mammalian

[194234]Genome 8: 767–769, 1997.

[194235]

[194236]

[194237]20756.

[194238]Yamada, Y.; Goto, H.; Ogasawara, N.: Cloning and nucleotide sequence

[194239]of the cDNA encoding human erythrocyte-specific AMP deaminase. Biochim.

[194240]Biophys. Acta 1171: 125–128, 1992.

[194241]

[194242]

[194243]20757.

[194244]Yamada, Y.; Goto, H.; Ogasawara, N.: A point mutation responsible

[194245]for human erythrocyte AMP deaminase deficiency. Hum. Molec. Genet. 3:

[194246]331–334, 1994.

[194247]

[194248]

[194249]20758.

[194250]Allegrucci, C.; Liguori, L.; Mezzasoma, I.; Minelli, A.: A1
adenosine

[194251]receptor in human spermatozoa: its role in the fertiliza-
tion process. Molec.

[194252]Genet. Metab. 71: 381–386, 2000.

[194253]

[194254]

[194255]20759.

[194256]Boman, H.; Hermodson, M.; Hammond, C. A.; Motulsky, A.
G.: Analbuminemia

[194257]in an American Indian girl. Clin. Genet. 9: 513–526, 1976.

[194258]

[194259]